Title

Optimizing functional connectivity scanning conditions for predicting autistic traits

Author list

Corey Horien[1,2,3]*, Francesca Mandino[4], Abigail S. Greene[2,5], Xilin Shen[4], Kelly Powell[6],

Angelina Vernetti[6], David O'Connor[7], Brendan D. Adkinson[2], Link Tejavibulya[8],

James C. McPartland[6,9], Fred R. Volkmar[6,9], Marvin Chun[9,10], Katarzyna Chawarska[6,11,12],

Evelyn M.R. Lake[4,10,13], Monica D. Rosenberg[14,15], Theodore Satterthwaite[1,3,16], Dustin

Scheinost[4,6,10,11,17], Emily S. Finn[18], R. Todd Constable[2,4,13,17,19]*

Affiliations

[1]Department of Psychiatry, University of Pennsylvania, Philadelphia, PA, USA

[2]MD-PhD Program, Yale School of Medicine, New Haven, CT, USA

[3]Penn Lifespan Informatics and Neuroimaging Center (PennLINC), University of Pennsylvania,

Philadelphia, PA, USA

[4]Department of Radiology and Biomedical Imaging, Yale School of Medicine, New Haven, CT,

USA

[5]Department of Psychiatry, Brigham and Women's Hospital, Boston, MA, USA

[6]Child Study Center, Yale School of Medicine, New Haven, CT, USA

[7]BioMedical Engineering and Imaging Institute, Icahn School of Medicine at Mount Sinai, New

York, NY, USA

[8] Department of Radiology, Weill Cornell Medicine, New York, NY, USA
23
[9]Department of Psychology, Yale University, New Haven, CT, United States

[10]Wu Tsai Institute, Yale University, New Haven, CT, USA

[11]Department of Statistics and Data Science, Yale University, New Haven, CT, USA

[12]Department of Pediatrics, Yale School of Medicine, New Haven, CT, USA

[13]Department of Biomedical Engineering, Yale University, New Haven, CT, USA

[14]Department of Psychology, University of Chicago, Chicago, IL, USA

[15]Neuroscience Institute, University of Chicago, Chicago, IL, USA

[16]Penn-CHOP Lifespan Brain Institute, University of Pennsylvania, Philadelphia, PA, USA

[17]Interdepartmental Neuroscience Program, Yale University, New Haven, CT, USA

[18]Department of Psychological and Brain Sciences, Dartmouth College, Dartmouth, NH, USA

[19]Department of Neurosurgery, Yale School of Medicine, New Haven, CT, USA

*Corresponding authors:

Corey Horien

Psychiatry Office of Education

3535 Market Street, Suite 200

Philadelphia, PA 19104

corey.horien@pennmedicine.upenn.edu

R. Todd Constable

Magnetic Resonance Research Center

300 Cedar St

PO Box 208043

New Haven, CT 06520-8043

todd.constable@yale.edu

<u>Abstract</u>

Autism is a heterogeneous condition, and functional magnetic resonance imaging-based studies

have advanced understanding of neurobiological correlates of autistic features. Little work has

focused on the optimal brain states to reveal brain-phenotype relationships. Using connectome-

based predictive modelling, we interrogated four datasets to determine scanning conditions that

boost prediction of clinically relevant phenotypes and assess generalizability. In dataset one, a

sample of youth with autism and neurotypical participants (n = 63), we found that a sustained

attention task resulted in high prediction performance of autistic traits compared to a free-

viewing social attention task and a resting-state condition. In dataset two (n = 25), we observed

the predictive network model of autistic traits generated from the sustained attention task

generalized to predict measures of attention in neurotypical adults. In datasets three and four, we

determined the same predictive network model further generalized to predict measures of social

responsiveness in the Autism Brain Imaging Data Exchange (n = 229) and the Healthy Brain

Network (n = 643). Our data suggest an in-scanner sustained attention challenge can help

delineate robust markers of autistic traits.

Introduction

Autism spectrum disorder (referred to as "autism" hereafter) affects approximately 1% of children around the world[1] and is characterized by difficulties with social communication and interaction, restricted and repetitive behaviors, and sensory atypicalities[2]. There is a need to better appreciate the neurobiological correlates of autistic traits in youth, which will help improve understanding of the condition and might aid potential clinical utility. Furthermore, there is a growing movement to characterize conditions like autism along dimensions of function[3-7].

There are numerous approaches to characterize the brain-based correlates of autism traits using functional magnetic resonance imaging (fMRI) connectivity data, in which measures of similarity of the blood-oxygen-level-dependent (BOLD) signal are computed between different regions of interest[8]. In particular, prediction-based studies—using functional connectivity data to predict a phenotype—have proven promising. For instance, case-control studies have focused on classifying those with autism compared to neurotypical participants, showing that high prediction accuracy can be achieved on the basis of functional connectivity differences[9-18]. Another approach predicts continuous measures of a phenotype (a symptom scale or a behavioral test score)[18-21]. One method of dimensional prediction is connectome-based predictive modelling (CPM)[22,23], which seeks to identify the functional connections most strongly predictive of a given phenotype. Groups using CPM in autism samples have identified brain correlates of clinician-rated autism symptoms[24,25], and other traits, such as behavioral inhibition[26], social responsiveness[24,27], and attentional states[28]. Successful models tend to comprise distributed networks, spanning cortical, subcortical, and cerebellar regions. In particular, association cortices tend to be important in successful predictions, especially when the phenotypes relate to

attention[24,28]. In addition, the most successful CPM models, while complex, consistently tend to

comprise edges from about 5-10 percent of the connectome. This perhaps points to an optimal

number of edges that might be involved in mediating brain-behavior relationships in those with

autism.

Nevertheless, which conditions yield the best predictive modeling performance is still

largely understudied. Most studies have typically focused on resting-state fMRI, in which

participants rest quietly in the scanner. However, in neurotypical participants, the importance of

scanning condition (e.g., 'brain state') is being recognized[29-32] for prediction of various

phenotypes, including intelligence[33-35], attention[36,37], working memory[38,39], personality traits[40],

cognition and emotion scores[41], as well as for emphasizing individual differences in connectivity

patterns[42]. These studies suggest that predicting out-of-scanner phenotypes using connectivity

measured during task performance tend to increase prediction accuracy, particularly when the

task probes some aspect of the out-of-scanner item of interest (e.g., memory tasks in the scanner

tend to result in higher prediction of memory performance outside the scanner[38]).

In addition, there are a number of elegant studies showing that in-scanner attention tasks

can be used to inform the neurobiological organization of autism[43-47]. There are also other brain

imaging studies suggesting an overlap between the functional networks mediating ADHD and

autism[24,48]. At a behavioral level, the co-occurrence of autism and attention-deficit/hyperactivity

disorder (ADHD) symptoms has long been acknowledged[49-52].

Motivated by the importance of tasks in assessing phenotypes, as well as the importance

of attention in autism, here we consider brain state-associated improvements in prediction

performance in a sample of youth with autism and neurotypical participants. Using data from

three different scanning conditions—a task requiring sustained attention, a task requiring

selective social attention (SSA), and resting-state data—we applied CPM to probe brain-behavior

relationships. Specifically, the gradual onset continuous performance task (gradCPT)[36,53,54] tests

the ability to sustain attention to constantly changing stimuli. The SSA task captures the ability

to process dynamic, multimodal faces presented one-by-one within a complex visual scene. One

of the best replicated eye-tracking biomarkers in autism[55-59], the SSA is devoid of any narrative

content or interpersonal interaction and was designed such that speech (SP) and eye contact (EC)

were varied. The task design was constructed to determine the effect of each condition on

prediction performance.

We hypothesized that consistent with the social features of autism, prediction

performance of autistic traits would be highest in the SSA task and would increase with the

presence of increased social cues. Specifically, we expected that the condition containing both

eye contact and speech (EC+SP+) would yield the strongest prediction performance. We

hypothesized that the next highest prediction performance would result from the sustained

attention task, due to the restricted and repetitive behaviors observed in autism, and that both

tasks would outperform resting-state data. To determine if results were robust, we used three

other datasets to determine if successful models can generalize to external samples. One of the

datasets was used to assess the model's generalizability in predicting performance on an

attention task; the other two datasets were used to assess prediction of other autistic features.

Results

Overview

Four samples were used in this work (Figure 1). The first dataset comprised 63 subjects

from a sample described previously (mean age = 11.7 years, st. dev. = 2.8 years; 29 females;

mean IQ = 107.8, st. dev. = 15.1; Supplemental Table 1)[28,60]. Twenty participants had autism; 11

other participants had a neurodevelopmental condition (five had attention-deficit/hyperactivity

disorder (ADHD), two had anxiety disorder, and four were classified as belonging to the broader

autism phenotype)[61]. Five of the participants with autism diagnoses had a co-occurring ADHD

diagnosis; this was not significantly different compared to those without autism (X^2 (1, $N = 60$)

= 1.64 (Yates correction), $P = 0.2001$). Hereafter, we refer to this dataset as the "Yale youth

sample". It should be kept in mind that the sample comprised both a mixture of those with autism

and those without, and there are only 20 participants with an autism diagnosis. Autism symptoms

were scored using the Autism Diagnostic Observation Schedule-2 (ADOS-2)[62].

Figure 1. An overview of the datasets used in this study. The Yale youth sample was the first
dataset used. FMRI connectivity data from different scanning conditions (task and rest) were
used to generate connectome-based predictive models of Autism Diagnostic Observation
Schedule (ADOS) scores (red circular arrows denote a brain-behavior predictive model). A
summary predictive model was then generated and applied to the adult attention sample. The
goal of this step was to determine whether the model generalized to predict attention phenotypes
(d') in an external dataset. The summary model was also applied to ABIDE and to HBN to
determine if the model predicted SRS scores in external samples. ABIDE, autism brain imaging
data exchange; ADOS, Autism Diagnostic Observation Schedule; CPM, connectome-based
predictive model; d', d-prime (attention phenotype in the Adult Attention Sample); HBN,
Healthy Brain Network; SRS, social responsiveness scale.
23

Participants in the Yale youth sample completed three different scanning conditions

(Figure 2; see Methods for further description of each task). We note that the SSA clips were

counterbalanced across participants; the other scan conditions were not (Supplemental Materials;

Supplemental Figure 1A). A standard preprocessing approach was used to generate

connectomes[33,63-65] from the different scanning conditions using a 268-node atlas[66]. For each

subject, the mean time-course of each region of interest ("node") was computed, and the Pearson

correlation coefficient was calculated between each pair of nodes to achieve a symmetric 268 x

268 matrix of correlation values representing connections between nodes ("edges"). The Pearson

correlation coefficients were then transformed to z-scores via a Fisher transformation, and only

the upper triangle of the matrix was considered, yielding 35,778 unique edges.

Figure 2. A schematic showing the scanning conditions used in the Yale youth sample. Top panel: free-viewing selective social attention task. Four conditions were shown to participants: no eye contact, no speech (EC-SP-); no eye contact, with speech (EC-SP+); eye contact, with no speech (EC+SP-); and eye contact, with speech (EC+SP+). No responses were required of participants during the different conditions. Middle panel: the gradual onset continuous performance task (gradCPT) was used as a sustained attention task. Grayscale pictures of cities and mountains were presented with images gradually transitioning from one to the next; button presses were required for city scenes and withheld for mountain scenes. Bottom panel: resting-state condition, in which the participants viewed a fixation cross. Please see the Methods section for further details about each scanning condition.

Prediction performance is highest in the Yale youth sample using task data

We first assessed which scanning condition resulted in the highest prediction

performance of autistic traits in the Yale youth sample. To ensure consistent amounts of data

across scanning conditions, we discarded frames from the end of gradCPT and resting-state runs,

such that the total amount of data was the same as from the Selective Social Attention task runs

(four minutes of data). CPM[23] was then used to assess prediction performance of ADOS scores

(Supplemental Figure 1B) and was repeated 500 times. Head motion was controlled for during

CPM as before[28,67,68]. The median performing model is represented in the text below, as well

as prediction ranges where appropriate; significance was assessed via permutation testing

(Methods).

We found differential performance across the various task conditions (Figure 3;

Supplemental Table 2). For example, performance using the resting-state data was quite low

(rest 1, Spearman's rho = 0.093, P-value = 0.23; rest 2, Spearman's rho = 0.18, P-value =

0.1240), and prediction performance was noted to have substantial variance (i.e., using data

from resting-state run 1, the minimum Spearman's rho = -0.2017, maximum Spearman's rho =

0.337, with 15% of the prediction performance scores below zero). Performance was also

quite low in the SSA condition with no eye contact and no speech (EC-SP-; Spearman's rho =

-0.106, P-value = 0.822). Surprisingly, there was large variance in prediction performance

scores using the SSA condition with eye contact and speech (EC+SP+; minimum Spearman's

rho = -0.172; maximum Spearman's rho = 0.323, with 7.2% of the prediction performance

scores below zero). Prediction performance was higher in the other SSA conditions, but was

not statistically significant after correcting for multiple comparisons (EC+SP-: Spearman's

rho = 0.251, P-value = 0.052; EC-SP+: Spearman's rho = 0.266, P-value = 0.034). The only

condition that resulted in statistically significant brain-behavior predictions was gradCPT 1

(Spearman's rho = 0.441, P-value = 0.004, corrected). See Supplemental Table 2 for statistics

for all CPM analyses.

Prediction performance has been noted to increase with increasing amounts of data[69],

possibly due to an increase in reliability of functional connectivity estimates[70-72]. We tested

this possibility by combining data from gradCPT 1 and gradCPT 2, as well as resting-state

session 1 and resting-state session 2. More data led to a slight increase in prediction

performance using both gradCPT and rest (Figure 3), though only gradCPT prediction

performance was statistically significant after multiple comparisons correction (gradCPT

average: Spearman's rho = 0.445, *P*-value = 0.002, corrected; rest average: Spearman's rho =

0.296, *P*-value = 0.038).

To ensure results were internally consistent, we repeated the CPM analysis using a

multiverse approach, which assesses how results are affected by different analytical choices[73].

The point of this approach is not to determine what CPM pipeline results in the highest

prediction performance. Instead, the goal is to assess various analytical scenarios and

determine how arbitrary modelling choices impact CPM performance. In the Yale youth

sample, we first adjusted CPM models for age, sex, and IQ. Encouragingly, we found similar

results to above: gradCPT results in the highest prediction performance; the SSA task with no

eye contact and no speech results in the lowest (Supplemental Table 2). The other SSA task

conditions did not tend to result in high predictions; rest performance was also low. As an

additional internal control, we assessed if gradCPT task performance differed among those

with autism compared to those without; we observed no difference in performance on the

attention task (mean gradCPT score of participants with autism = 2.59 (st. dev. 0.96); mean score

of participants without autism = 2.66 (st. dev. 0.86; $t(55) = -0.27$, $P = 0.79$). Further, we

conducted additional controls in which we modelled additional autistic traits with less skewed

distributions than the ADOS scores used above; we again found successful prediction of ADOS

uncalibrated scores, SRS scores, and ADOS scores when balancing autistic and non-autistic

participants (Supplemental Figure 2; Supplemental Table 3).

We continued with the multiverse analysis and repeated CPM using the same pipeline

as above, except instead of predicting total ADOS scores, we attempted to predict the social

affect and the restricted and repetitive behaviors subscales of the ADOS. We observed the

same overall trend—gradCPT tends to result in the highest prediction performance, and the

resting-state and SSA task with no eye contact and no speech performed the poorest

(Supplemental Figure 1C; Supplemental Table 2). The SSA tasks resulted in increased

prediction performance of social affect scores, but predictions were not significant after

controlling for multiple comparisons. Prediction performance of restricted and repetitive

behaviors using the SSA data tended to be low.

To ensure results were robust to preprocessing pipeline, we reprocessed functional

data by altering the functional parcellation (using a 368-node instead of a 268-node atlas); we

also repeated analyses without performing global signal regression. In both cases, we

observed a similar pattern: gradCPT average had the best absolute prediction performance

across pipelines tested, while the SSA and rest conditions tended to vary, with many prediction

iterations below zero (Supplemental Figure 1C; Supplemental Table 4). Interestingly, we

observed, as have others[33,74,75], that GSR decreased the strength of brain-behavior relationships,

while the overall prediction patterns across scans remained consistent. In sum, these results

suggest that the CPM findings observed here were not being driven by a particular processing

choice.

Finally, given that we are using a dimensional approach to predict ADOS scores, and

ADOS scores are by definition higher in those with autism, we explored if case versus control

classification was possible. (Note there is no strict ADOS cut-off to classify autism—it is

simply one component of a diagnostic evaluation—but scores tend to be > 5 in those

diagnosed.) Modifying CPM and balancing cases and controls, we observed gradCPT data

resulted in the highest sensitivity, specificity, and accuracy in case-control classification,

particularly the gradCPT average data (Supplemental Figure 1D; Supplemental Table 4). The

fact that gradCPT afforded the highest predictions reiterates findings from the dimensional

CPM—that task brain states requiring sustained attention tend to outperform those from the

SSA and resting-state conditions.

Figure 3. CPM prediction performance across different scanning conditions for total ADOS
scores. The scan condition is shown on the x-axis; on the y-axis, Spearman's rho is shown for
the correlation of predicted and actual ADOS scores. For each condition, the median of the
500 iterations is shown as a solid black line in the violin plot; quartiles, as dotted lines. The
Selective Social Attention task conditions are shown in purple, gradCPT in turquoise, and
resting-state data in yellow. Statistical significance was obtained using permutation testing as

described in the Methods. Note that one-side significance testing was used, as we are interested
in statistically significant positive predictions only. Asterisk (*) indicates statistical significance
after correcting for multiple comparisons using the Benjamini–Hochberg method[135] (correcting
for ten tests: two for gradCPT, four for SSA, two for resting-state, one for gradCPT average, and
one for resting-state average). Exact Spearman's rho and P-values for statistically significant
conditions, gradCPT1: Spearman's rho = 0.441, P-value = 0.004; gradCPT average:
Spearman's rho = 0.445, P-value = 0.002. ADOS, autism diagnostic observation schedule;
Avg, average; EC-SP-, no eye contact, no speech; EC-SP+, no eye contact, with speech; EC+SP-
, eye contact, with no speech; EC+SP+, eye contact, with speech; mm, millimeters; SSA,
selective social attention task.
11

External validation of predictive models—attention prediction in the adult attention sample

Having determined that data derived from attention tasks are best for predicting

autistic traits, we next assessed generalizability of the attention predictive model. Previously,

we have shown it is possible to build predictive models of sustained attention in the Yale

youth sample and that such a model is related to autistic traits[28]. Therefore, we assessed the

extent to which predictive models of autistic traits are related to sustained attention. To ensure

generalizability was not driven by sample-specific noise, we tested the predictive model in an

external dataset of individuals performing the same gradCPT task (n=25 neurotypical adults,

13 females, mean age = 22.8 years, st. dev. = 3.5 years)[36]. Hereafter, we refer to this dataset as

the "adult attention sample" (Figure 1). The behavioral outcome of interest in this sample is

performance on the gradCPT, d' (sensitivity), the participant's hit rate minus false alarm rate

(mean d' = 2.11, st. dev. = 0.92).

We determined which edges tended to contribute consistently to successful prediction

of ADOS phenotypes in the Yale youth sample (see Methods, 'Testing generalizability of the

ADOS network') using the model generated from average gradCPT data in the prediction of

total ADOS scores. The resulting model (the 'ADOS consensus network') was used to

determine if there was a relationship between predicted ADOS scores and d' scores in the

adult attention sample. Specifically, we used the fMRI gradCPT task data from the adult

attention sample to generate a predicted ADOS score. Predicted ADOS scores were then

compared to actual d' scores across participants to assess accuracy. We point out this differs

from the Yale youth sample, where we were able to compare predicted ADOS scores with

observed ADOS scores. In the adult attention sample, the goal was to assess the relationship

between the model (trained to predict ADOS) and attention (d').

We observed a statistically significant relationship between predicted ADOS scores

and d' scores (Spearman's rho = -0.56, P = 0.0049, corrected; Figure 4A). Specifically,

higher predicted ADOS scores were associated with lower d' scores, indicating poorer

performance on the task and implying lower sustained attention. To ensure results were

robust, we repeated analyses controlling for potential confounds; predictions remained high

when adjusting for participant head motion (Spearman's rho $=$ -0.56, P $=$ 0.0043),

participant sex (Spearman's rho $=$ -0.49, P $=$ 0.0164), and participant age (Spearman's

rho $=$ -0.55, P $=$ 0.0066). In addition, we also assessed the relationship between predicted

ADOS scores and d' scores using only the ADOS positive network and then only the ADOS

negative network. We observed a statistically significant negative correlation in the ADOS

positive model (Spearman's rho = -0.59, P = 0.0021, corrected; Figure 4B) and in the ADOS

negative model (Spearman's rho = -0.46, P = 0.023, corrected; Figure 4C).

Lastly, we altered the stringency of how often an edge had to be included in the ADOS

consensus model (Methods). We observed consistent results across a range of thresholds

(Supplemental Table 5), increasing confidence that there is a relationship between ADOS

network strength and d' scores. In sum, these results suggest that the predictive model of

autistic traits captures variance related to sustained attention.

1
2

Figure 4. Generalization of the ADOS consensus network in the adult attention sample. A.
Results using the combined network model. B. Results using the positive-association network
model. C. Results using the negative-association network model. Predicted ADOS scores are
indicated on the *y*-axis; *d*' scores, on the *x*-axis. Higher predicted ADOS scores are associated
with lower *d*' scores, indicating poorer performance on the task and implying lower sustained
attention. Data are presented as a regression line (solid line) +/- 95% confidence intervals
(shaded areas). Spearman's rho was calculated and was used to assess statistical significance
(two-sided). The Benjamini–Hochberg method[135] was used to correct for three tests. ADOS,
autism diagnostic observation schedule; *P* = *P*-value.
12
13
14
External validation of predictive models—social responsiveness prediction in ABIDE and

HBN

After finding we could successfully predict attention scores, we set out to determine if

the predictive model from the Yale youth sample generalized to predict social responsiveness

in a large sample of participants from ABIDE (n=229, 65 females; mean age = 10.45 years, st.

dev. = 1.8 years; mean IQ = 113.7, st. dev. = 15.1; 77 individuals with autism)[76,77] described

elsewhere[24]. We used the same approach as in the adult attention sample to assess

generalizability. Specifically, we used the resting-state data from ABIDE and applied the

ADOS consensus model to predict social responsiveness scale (SRS) scores[78] across

participants (Methods). As with the other test of generalizability above, predicted ADOS scores

were then compared to actual SRS scores to assess accuracy.

We observed successful prediction of all SRS scales tested (Figure 5). In particular,

the model generalized to predict SRS total scores (Spearman's rho $=$ 0.17, $P =$ 0.008,

corrected, Figure 5A), as well as SRS subscales quantifying communication (Spearman's

rho $=$ 0.15, $P =$ 0.028, corrected, Figure 5B), mannerisms (Spearman's rho $=$ 0.21,

$P =$ 0.001, corrected, Figure 5C), and motivation (Spearman's rho $=$ 0.16, $P =$ 0.016,

corrected, Figure 5D). We also tested prediction of each SRS scale after adjusting for

participant age, sex, and head motion; predictions were essentially unchanged, further

supporting that the ADOS model is capturing variance related to the SRS scales

(Supplemental Table 6). As above, we altered how often an edge had to be included in the

ADOS CPM and retested predictions. In every case, we observed similar predictions across

various thresholds for all SRS scales (Supplemental Table 6).

We performed a similar test of generalization using data from the Healthy Brain

Network (HBN) (n=643, 264 females; mean age = 11.01 years, st. dev. = 2.63 years; mean IQ =

101.89, st. dev. = 16.7; 107 participants had a diagnosis of autism; 302 had a diagnosis of

ADHD). Using functional data obtained while participants watched a naturalistic movie clip

(scenes from Despicable Me; Methods), we applied the ADOS consensus model to predict

social responsiveness scale (SRS total) T-scores (mean = 56.36, st. dev. = 11.21)[78] across

participants. Predicted ADOS scores were then compared to actual SRS scores to assess

accuracy. Controlling for head motion, we again observed the model generalized to predict SRS

scores (Spearman's rho = 0.1002, P = 0.0111). Like before, we tested a variety of summary

models of differing sizes. We observed that results were stable when the consensus model had

both more and fewer edges, supporting that our results were robust to model size (Supplemental

Table 7). Taken together, these data indicate the ADOS model from the Yale youth sample

generalized to predict aspects of sociality in ABIDE and HBN.

Figure 5. Generalization of the ADOS consensus network to ABIDE. A. SRS total score
results. B. SRS communication score results. C. SRS mannerism score results. D. SRS
motivation score results. For all plots, actual SRS scores from each subscale are indicated on
the *x*-axis; predicted scores, on the *y*-axis. Data are presented as a regression line (solid line) +/-
95% confidence intervals (shaded areas). Spearman's rho was calculated and was used to
assess statistical significance (two-sided). The Benjamini–Hochberg method[135] was used to
correct for four tests. ADOS, autism diagnostic observation schedule; P = P-value.
12

Neuroanatomy of predictive edges

We next visualized brain connections in the ADOS consensus model. The network

comprised 2,014 total edges (1,001 edges in the positive-association and 1,013 edges in the

negative-association network), approximately 5.6% of the connectome, in line with other

models that have generalized to predict autism and attention symptoms[24,28]. Edges across the

brain were represented in the model, constituting a complex, distributed network (Figure 6A-

B). In particular, connections within and between heteromodal association networks were

found to contain the highest fraction of edges (Figure 6C-D; note that results have been

corrected for differing network size)[28]. For instance, the top three network pairs containing the

greatest proportion of edges in the positive-association network involved medial frontal,

frontoparietal, or default mode networks. In the negative-association network, the top three

network pairs involved connections within and between medial frontal, frontoparietal, or default

mode networks (e.g., in this case, the top network pair comprised connections within the medial

frontal network; the next highest network comprised connections between the medial frontal and

frontoparietal networks; and the third highest network connected the medial frontal and default

mode networks). In addition, 704/1001 of the edges in the positive-association network and

535/1013 of the edges connected to medial frontal, frontoparietal, or default mode networks.

We performed further visualizations using slightly different thresholding techniques; these

analyses again showed that association networks were important in the ADOS consensus

model (Supplemental Figure 1E).

Figure 6. Neuroanatomy of ADOS consensus network. A. The positive-association network.
B. The negative-association network. For both A and B: a circle plot is shown in the upper left.
The top of the circle represents anterior; the bottom, posterior. The left half of the circle plot
corresponds to the left hemisphere of the brain. A legend indicating the approximate anatomic
'lobe' is shown to the left. The same edges are plotted in the glass brains as lines connecting
different nodes (red circles); in these visualizations, nodes are sized according to degree, the
number of edges connected to that node. To aid in visualization, we have thresholded the
matrices to only show nodes with a degree threshold > 25. C. Matrix of the positive-association
network. D. Matrix of the negative-association network. For both C and D: The proportion of
edges in a given network pair is shown; data have been corrected to account for differing
network size. MF, medial frontal; FP, frontoparietal; DM, default mode; MT, motor; VI,
visual I; VII, visual II; VA, visual association; CO, cingulo-opercular; SB, subcortical; CB,
cerebellum.
15
16
17 *Testing for convergence of task versus rest CPM findings*
18
19   As a last test of generalizability, we repeated analyses, except we reversed the order of

20 which dataset was used to build the original model. As our primary aim was determining if

21 task data outperformed resting-state data in the prediction of autistic traits, and ABIDE does

not have task data, we used the Healthy Brain Network (HBN) sample. Specifically, we used

data from two conditions in which participants watched movies ('Despicable Me' and 'The

Present') in the scanner, along with a resting-state condition. Although HBN does not contain

the exact same tasks as the Yale youth sample, because the movie watching conditions

comprise a continuous task requiring sustained attention to an evolving narrative involving

multiple characters interacting in a complex scene, the sample still allows us to test broadly if

there is a relationship between autistic traits and attention. Note that the naturalistic movies

stand in distinction to the SSA tasks, which were designed to include only one actress

speaking at a time in short phrases (i.e., no narrative content or social interactions are

otherwise presented by the participant during the SSA tasks).

After ensuring all scan conditions had the same amount of data per scan (Methods), we

repeated CPM, controlling for participant head-motion as above, and attempted to predict SRS

scores. We observed successful prediction in the movie-watching conditions (median

performing model from Despicable Me: Spearman's rho = 0.14, P-value < 0.001, corrected;

median performing model from The Present: Spearman's rho = 0.11, P-value < 0.0001,

corrected). Prediction using resting-state data was non-significant (median performing model:

Spearman's rho = 0.05, P-value = 0.102, corrected (Supplemental Figure 1F).

To assess generalizability of the model, we used connections that tended to contribute

to successful predictions in the HBN DM condition to generate a consensus model (Methods).

The resulting DM consensus model was used to assess if there was a relationship between

predicted SRS scores and ADOS scores in the Yale youth sample. Specifically, we used the

gradCPT data in the Yale youth sample and applied the DM consensus model to predict social

responsiveness scale (SRS) scores[78] across participants (Methods), again controlling for head

motion. Predicted SRS scores were then compared to actual ADOS scores to assess accuracy.

Successful prediction was achieved (Spearman's rho = 0.26; *P*-value = 0.0422). To ensure

results were robust, we repeated analyses using different thresholds when generating the

consensus mask; we found results were stable across both looser and stricter thresholds

(Supplemental Table 8). In sum, the analyses in this section replicate our initial finding that task

data outperform resting state data in predicting autistic traits and that such a model can

generalize in external samples.

Discussion

We determined that using functional connectivity calculated from data acquired during

gradCPT resulted in the prediction of autistic traits. The resulting model generalized to

independent samples to predict attention and social phenotypes in neurotypical participants and

those with autism. Altogether, results highlight the potential of using in-scanner tasks,

particularly those demanding sustained attention, to more accurately determine brain-behavior

relationships in clinical samples.

There is a rich history of using tasks to probe the cognitive architecture of autism[79-84].

Nevertheless, most fMRI brain-behavior prediction studies in autism that use machine learning

techniques have typically relied on resting-state data (see[4] for a recent review). Our results

suggest that by optimizing the brain state under which data are acquired through task

engagement[85], more accurate brain-behavior relationships can be studied[29]. Improved brain-

behavior mapping increases the potential clinical utility of neuroimaging approaches[86] and might

help obtain a more accurate picture of brain circuits underlying the complex phenotypic

landscape of autism. Tasks also offer the advantage of improving the reliability of task-engaged

functional connections[87]. More generally, the results obtained here are in line with other work in neurotypical populations, indicating that predictions of phenotypes improve when using task as opposed to resting-state data[29,33,34,36,41,88,89]. We note that resting-state studies still retain utility, particularly in terms of ease of data collection and their ability to facilitate the collation of large datasets across centers.

Despite tasks outperforming rest-based models, our data indicate that not all tasks are equal. Our original hypothesis regarding prediction performance was that, reflective of social difficulties in autism, the SSA tasks would result in the highest prediction of ADOS scores. That an attention task—superficially unrelated to the core aspects of autism—outperformed a task designed to probe social abilities reinforces the importance of empirical work. The current results suggest that simply because a scanning condition is ostensibly related to a given phenotype does not mean that a brain-behavior relationship can be found in neurodevelopmental conditions.

It is perhaps puzzling that the SSA clips did not result in higher prediction performance. This could be due to the passive nature of the SSA clips, allowing the participants to let their minds wander[55,56]. As such, SSA clips could be interpreted as resembling resting-state[112]. The unconstrained nature of the resting-state might be suboptimal for probing certain aspects of brain-behavior relationships[29]. In parallel, it is perhaps also surprising that the gradCPT data resulted in the highest prediction performance. Beyond attention, the highly structured, rules-based design of gradCPT may effectively highlight variations in networks linked to autistic traits given the rules-based tendencies observed in autism[113].

Regardless of the exact mechanism, our finding that gradCPT led to the prediction of autistic and attention phenotypes adds to the growing literature suggesting an important link between autism and attention at the neurobiological level. Previous studies[24,47,93,94] have

indicated that complex models spanning numerous functional networks are important for

attention in autism. In addition, the size of the ADOS consensus model is of interest

(approximately 5% of the connectome) and is in line with the size of other models that have

generalized to predict autism and attention symptoms[24,28]. Notably, these models tend to share

a few characteristics with the ADOS consensus model. Edges tend to originate across the

entire brain, linking within and between most canonical network pairs. This finding

underscores that there is no one region or network preferentially involved in a given

phenotype, consistent with other psychiatric conditions[95]. The fact that distributed brain

connections mediate complex behaviors is a theme observed in multiple species, including the

global brain dynamics underlying homeostatic sleep behaviors in *Caenorhabditis elegans*[96], to

long-range connections underpinning social decisions in primates[97].

While the ADOS model is complex, post-hoc analyses revealed other neurobiological

trends. Similar to the ADOS model, previous models involved in predicting attention and

autism all tend to have particular representation in heteromodal association cortices[67],

including the default mode (recently reviewed in[98]). In work involving CPM, the default mode

network tends to be important, especially in the negative networks[24,26,28]. In CPM, edges

comprising the negative network are selected based on a negative correlation with the

behavior of interest. In the present work, it follows that higher connectivity in the default

mode, typically associated with more mind-wandering or internally-directed states[99], would be

associated with lower attention abilities, as well as potentially higher autistic traits. (A recent

mega-analysis has observed hyperconnectivity involving the DMN in both autistic and ADHD

participants as well.[100]) It is plausible that interactions among association networks—

including not only the default mode, but also the medial frontal and frontal parietal

networks—are needed to coordinate the complex cognitive demands associated with higher-

level brain functions[101], such as sustaining attention and picking up on social cues.

We contend that autistic traits and attentional abilities are hence multifaceted, interrelated

processes. Because the CPM also predicts attention in the adult sample, it indicates that the

attention task state is appropriately influencing brain circuits of interest involved in modulating

attention in the Yale autism sample. That the CPM goes on to predict a different phenotype

related to a core component of autism—namely, social difficulties—reiterates the shared

relationships between autistic features and attention. Indeed, autism and ADHD symptoms have

long been acknowledged to overlap[49-51,104]. In a more extreme example, it is possible the CPM is

more related to a broader pathology and could be applicable, to some degree, to many other

phenotypes, similar to the p-factor[105]. Instead of commenting on the validity of the p-factor as a

single unifying construct[106], we simply note that there are numerous examples of redundancy[107]

and degeneracy in biological circuits[108,109], especially in brain networks. Further, we do not

mean to argue that we have discovered a 'core autism/attention network.' Rather, it is logical that

these phenotypes, both defined to some extent by how complex information is perceived from

the outside world, are associated with overlapping circuitry.

A few additional items warrant discussion in the final paragraphs. Participant IQ in the

Yale youth sample and ABIDE was fairly high. More research should be conducted using

participants with a broad range of IQ scores to determine which scanning conditions are optimal

for prediction performance. The current work relied on a sample of 20/63 participants with

autism. Future work could aim to repeat analyses in more participants with autism. The autism

dataset contained mainly males; the importance of sex-based differences in brain circuitry[114-116]

and behavioral phenotypes[117] relevant for autism is increasingly well-described. It is possible

that a more active social task requiring participant engagement (unlike the SSA task) could

increase prediction performance. Likewise, given the success of the movie-watching conditions

of the HBN, perhaps more naturalistic movie clips (demanding the processing of narrative

content and complex, interpersonal relationships) are another promising path forward in autism

research[118,119]. Further, the role of arousal and its impact on predictive modelling could also be

studied. In addition, while we have taken numerous steps to lessen the effect of head motion, it

can still impact brain-behavior relationships; this must be kept in mind when interpreting the

current work. A potential limitation of the adult attention sample is that it comprises older

participants; another possible interpretation is that it strengthens the generalizability of the CPM.

That is, the fact the model generalized in a different age range eliminates the potential confound

that the model is confounded by age in the Yale youth sample (which in turn would drive

successful generalization in a test sample of similar ranges, but would fail to generalize in a

sample comprising different ages).

Also, the current work focused on prediction of traits in an adolescent dataset that is

relatively small; the adult attention sample was also quite small. Though replicable and

generalizable findings can still be determined by using robust methods[120-122] and we were well-

powered in the current study (see Methods for a power analysis), future studies in larger samples

are needed. Collecting small, unique samples also facilitates testing across diverse experimental

conditions, thereby enhancing generalizability[123]. We contend that in the age of big data, it is

essential to continue exploring brain-behavior associations in samples that might not contain

thousands of subjects, but comprise unique scanning conditions. Such an approach allows the

field to better determine which scans to include in big data endeavors and facilitates the

exploration of questions that may be difficult to address in large-scale studies. Balancing the

need for large samples with adequate scans per subject is a key question that has received much

attention in the neurotypical brain-wide association literature[124,125]. Related studies could be

pursued in the autism field. Including participants with moderate autism symptoms to tease apart

the brain correlates of more subtle phenotypes will be an important aspect of future work,

regardless of exact study design. More studies could also be conducted to assess task design and

prediction performance in much younger samples, such as toddlers and young children. Such

efforts aim to optimize the detection of brain-behavior relationships at earlier developmental

stages, ultimately providing better support for individuals with autism and their families.

Conclusions

We have shown in a preliminary study that sustained attention tasks, such as gradCPT,

can enhance the prediction of autistic traits. Such an approach leads to a robust marker that

generalizes to predict attention and social phenotypes in independent samples. Our findings

highlight the need to further investigate optimal brain states for modeling phenotypes in autism

and related conditions.

Methods

Description of datasets

We used four datasets in this work (Figure 1). The first dataset, the Yale youth sample,

comprised 63 subjects from a sample described elsewhere (mean age = 11.7 years, st. dev. = 2.8

years; 29 females; mean IQ = 107.8, st. dev. = 15.1)[28,60]. Twenty of the participants had autism

(five of whom had ADHD); 11 other participants had a neurodevelopmental condition (five with

ADHD, two with anxiety disorder, and four were classified as belonging to the broader autism

phenotype)[61]. Participants were scanned on a 3T Siemens Prisma System. See Supplementary

Material for full exclusion criteria and imaging parameters. Autism symptoms were scored using

the Autism Diagnostic Observation Schedule-2 (ADOS-2)[62] and were ascertained by trained

clinical psychologists; calibrated severity scores were used in the present work for the social

affect subscale (mean = 3.2, st. dev. = 2.9), the restricted and repetitive behavior subscale (mean

= 4.0, st. dev. 3.3), and the ADOS total score (mean = 3.1, st. dev. = 3.1). This sample was used

to conduct CPM, compare how scanning condition impacted performance, and generate a

consensus model. Note that although the sample is small, we were well-powered to detect

effects. Using the R package "pwr", with a sample size of 63 subjects, and assuming a

significance level of 0.05 (two-sided), a correlation between predicted and observed ADOS

scores in gradCPT average of 0.445 corresponds to a power of 0.96.

A second dataset of neurotypical adults, the adult attention sample (n=25, 13 females,

mean age = 22.8 years, st. dev. = 3.5 years) was used as a validation dataset and is described

elsewhere[36]. Participants were scanned on a 3T Siemens Trio TIM system. This sample was used

to determine if the consensus model generalized to predict attention.

A third dataset of individuals with and without autism (n=229, 65 females; mean age =

10.45 years, st. dev. = 1.8 years; mean IQ = 113.7, st. dev. = 15.1) comprising data from

ABIDE[76,77] was used as an additional validation dataset; processing of these data is described

elsewhere[24]. Note that 77/229 participants were diagnosed with autism. SRS[78] raw scores were

used from ABIDE and included the following scales: SRS total scores (mean = 42.4, st. dev. =

40.2); SRS communication (mean = 13.8, st. dev. = 14.2); SRS motivation (mean = 7.3, st. dev.

= 6.9); SRS mannerisms (mean = 7.1, st. dev. = 8.5). Seventy-seven of the participants had

autism. SRS was chosen due to the low numbers of subjects with ADOS scores (when using the

exclusion criteria described in[24]), along with the additional quality control exclusion criteria we

performed (i.e., there were 229 subjects with SRS data, compared to only 33 with ADOS; see

Supplemental Methods). This sample was used to determine if the consensus model generalized

to predict SRS scores.

A fourth dataset of individuals (n=643, 264 females; mean age = 11.01 years, st. dev. =

2.63 years; mean IQ = 101.89, st. dev. = 16.7) comprising data from the transdiagnostic HBN

sample[126] was used as an additional validation dataset. In the current study, 107 had a diagnosis

of autism, while 302 had a diagnosis of ADHD. Processing of these data is described

elsewhere[127]. SRS[78] total T-scores were used from HBN (mean = 56.36, st. dev. = 11.21). This

sample was used to determine if the consensus model generalized to predict SRS scores.

For the Yale youth sample and the adult attention sample, the Yale University

institutional review board oversaw and approved the ethics of the study. For HBN, the study was

approved by the Chesapeake Institutional Review Board. The data for ABIDE were approved by

each of the 16 sites contributing data (i.e. approval was required by each of the home institutions

prior to submission)[76]. Where appropriate, informed consent was obtained from the parents or

guardians of participants, and participants were paid for their involvement. Written assent was

obtained from children aged 13–17 years; verbal assent was obtained from participants under the

age of 13 years.

Preprocessing of functional imaging data

A standard preprocessing approach was used that has been described elsewhere[33,63-65].

Preprocessing steps were performed using BioImage Suite[128] unless otherwise noted, and

included: skull stripping the 3D magnetization prepared rapid gradient echo images using

optiBET[129] and performing linear and non-linear transformations to warp a 268-node functional

atlas[130] from Montreal Neurological Institute space to single subject space. Functional images

were motion-corrected using SPM8 (https://www.fil.ion.ucl.ac.uk/spm/software/spm8/).

Covariates of no interest were regressed from the data, including linear, quadratic, and cubic

drift, a 24-parameter model of motion[131], mean cerebrospinal fluid signal, mean white matter

signal, and the global signal. Data were temporally smoothed with a zero-mean unit-variance

low-pass Gaussian filter (approximate cutoff frequency of 0.12 Hz). Visual inspections were

performed after skull-stripping, non-linear, and linear registrations to ensure there were no errors

in processing. Head motion was computed as the mean frame-to-frame displacement (FFD) of

the participant's head[60] (see Supplemental Methods for additional motion control considerations,

as well as Supplemental Figure 1G). To ensure consistent amounts of data across scanning

conditions, we discarded frames from the end of the gradCPT and resting-state runs, such that

the total amount of data was the same as from the SSA task runs (four minutes of data).

Connectomes were generated using a 268-node atlas[22]. For each subject, the mean time-

course of each region of interest ("node") was computed, and the Pearson correlation coefficient

was computed between each pair of nodes to achieve a symmetric 268 x 268 matrix of

correlation values representing "edges" (connections between nodes). The Pearson correlation

coefficients were then transformed to *z*-scores via a Fisher transformation, and only the upper

triangle of the matrix was considered, yielding 35,778 unique edges.

The 268-node atlas was chosen because it is a full brain parcellation, covering cortical,

subcortical, and cerebellar regions, as well as striking a good balance in terms of number of

nodes and reducing computational resources while approximating biologically plausible brain

regions[66]. For example, the 268-node atlas results in 35,778 unique edges (because number of

edges = ((number of nodes x number of nodes) – number of nodes) / 2). Increasing the atlas size

by 100 to 368 nodes results in 67,528 edges, almost doubling the number of edges and thereby

doubling compute time. In addition, from a practical perspective, the 268-node parcellation

allows generalization testing in the adult attention, ABIDE samples, and HBN samples, which

had previously been processed using the 268-node atlas[24,28,36,127]. To ensure results were not

being driven by this specific parcellation, we repeated analyses using a 368-node atlas[132] (which

was generated using a different approach, namely, a spectral clustering algorithm in the cortex,

anatomical labelling of structures in subcortical regions, and by incorporating cerebellar regions

defined elsewhere[133]). We found CPM results did not differ, consistent with other brain-behavior

modelling studies showing that results tend to be stable across a range of parcellations[33,74].

Scanning conditions in the Yale Youth Sample

1) Scanning condition one: the free-viewing selective social attention task

14   Participants completed a novel version of a free-viewing selective social attention (SSA)

task[55,58] in which an actress is presented at the center of the screen and is surrounded by objects

in corners of the screen (Figure 2). Four types of clips were used in which the presence of speech

(SP) and eye contact (EC) were manipulated. The first condition included clips in which the

person smiled and made eye contact with the camera while speaking in full sentences (e.g.,

"Have you ever seen a monkey? Monkeys eat bananas, swing in trees, and chase each other.";

this was designated as the EC+SP+ condition). The second condition included a direct gaze

condition with no speech (EC+SP-), in which the person smiled directly at the viewer while

remaining silent. The third condition consisted of the person looking down at the table while

speaking in full sentences (EC-SP+). The fourth condition consisted of the person looking down

at the table and not speaking (EC-SP-). Each clip lasted two minutes and was shown twice over

four runs, such that eight clips were shown in total. To allow successful scene transitions in

between sentences, the direct gaze and speech condition lasted 2 minutes and 8 seconds. The

speech with no eye contact condition lasted 2 minutes and 6 seconds. In between clips during

each run, a white fixation cross on a black background was shown for 15 seconds. Clip order was

counterbalanced across participants (see Supplemental Materials for more about the

counterbalancing of clips, as well as study design considerations of the Yale youth sample). Clip

conditions were concatenated across runs, such that each resulting connectivity matrix comprised

four minutes of data from a single scanning condition. Both gradCPT and the SSA task were

presented using Psychtoolbox (version: 3.0.14; http://psychtoolbox.org/; MATLAB version

R2018a) on a Lenovo IdeaPad 720S computer, with Ubuntu 16.04 LTS installed.

2) Scanning condition two: the gradual onset continuous performance task (gradCPT)

The gradual onset continuous performance task (gradCPT; Figure 2) [36,53,54] was used in

datasets one and two. The gradCPT tests sustained attention and inhibition, producing a range of

performance scores across neurotypical[53,54] and neurodiverse populations[28]. Participants viewed

grayscale pictures of cities and mountains presented at the center of the screen, with images

gradually transitioning from one to the next every 1,000 ms. Subjects were instructed to respond

with a button press for city scenes and to withhold button presses for mountain scenes. City

scenes occurred randomly 90% of the time. Performance was calculated using d' (sensitivity),

the participant's hit rate minus false alarm rate. The task took 5 minutes to complete; participants

completed the task twice. Note that because of differences in task timing between gradCPT, the

selective social attention task, and resting-state, we trimmed the gradCPT and resting-state data

to include 4 minutes of data per scan (to match the selective social task time length of 4

minutes).

Subjects in the adult attention sample also performed gradCPT; the same parameters were

used as above, except scene transitions took 800 ms.

3) Scanning condition three: resting-state data

Resting-state data were also obtained. Subjects were instructed to keep their eyes open,

relax, and think of nothing in particular while they viewed a white fixation cross on a black

screen. Each scan lasted five minutes and was repeated twice per participant. Resting-state data

were also obtained in the ABIDE sample[76,77].

Scanning conditions in the HBN sample

We considered three conditions when performing CPM. Specifically, we used functional

data from two separate naturalistic movie conditions, in which participants watched *Despicable*

Me and *The Present*, as well as a single run of resting-state data. Because of differences in scan

time across conditions, we truncated time courses so they were consistent across all three

conditions, so that all matrices were generated using four minutes of data (the length of *The*

Present).

Connectome-based predictive modelling

CPM[23] (Supplemental Figure 1B) was used to predict ADOS scores from functional

connectivity data in the Yale youth sample. Briefly, using 10-fold cross-validation, connectivity

matrices from a given scan condition and ADOS scores were split into an independent training

set including subjects from 9 folds and a test set including the left-out fold. Linear regression

was used to relate edge strength to ADOS score in the training set. Edges most strongly

associated with ADOS scores were selected (feature selection threshold of $P = 0.05$) for both a

'positive network' (in which increased connectivity was associated with higher ADOS scores)

and a 'negative network' (in which increased connectivity was associated with lower ADOS

scores). We used partial correlation to control for mean participant head motion at the feature

selection step[28,67,68]. Note that this is the 'base model' we are considering throughout the results

(i.e., the results shown in Figure 3 used partial correlation to control for motion). Mean network

strength was computed in both the positive and negative networks, and the difference between

these network strengths was computed ('combined network strength'), as in previous work [33].

A linear model was then calculated relating combined network strength to ADOS scores

in the training set. In the last step, combined network strength was computed for the test set, and

the model was applied to generate ADOS predictions for these unseen participants.

Model performance was assessed[68] by comparing the similarity between predicted and

observed ADOS scores using both Spearman's correlation (to avoid distribution assumptions)[134].

Note that performance was assessed on the entire sample after cross-validation. That is, in a

given iteration, each of the ten folds resulted in a set of predictions for ADOS. After each

participant had a predicted score from being in the test set, Spearman's correlation was used to

assess the relationships between predicted and observed scores across the entire sample. We

performed 500 iterations of a given CPM analysis and selected the median-performing model;

we report this in the main text when discussing model performance. To calculate significance,

we randomly shuffled participant labels and attempted to predict ADOS scores. We repeated this

500 times and calculated the number of times a permuted predictive accuracy was greater than

the median of the unpermuted predictions to achieve a non-parametric *P*-value:

$P = (\#(\text{rho}_{\text{null}} \geq \text{rho}_{\text{median}})) / 500$

where $\#(\text{rho}_{\text{null}} \geq \text{rho}_{\text{median}})$ indicates the number of permuted predictions numerically

greater than or equal to the median of the unpermuted predictions[68]. We used the Benjamini–

Hochberg method[135] to correct for multiple comparisons, correcting for ten tests in the Yale

youth sample (two for gradCPT, four for SSA, two for resting-state, one for gradCPT average,

and one for resting-state average), three tests in the adult attention sample, and four tests in

ABIDE.

We note the same CPM approach was used in the HBN sample, except 5-fold cross-

validation was used, in line with the larger sample size[134]. All other parameters were the same as

above. We corrected for three tests in the HBN sample when correcting for multiple

comparisons.

Testing generalizability of the ADOS network

To determine if the ADOS networks from the Yale youth sample generalized to external

datasets (the adult attention sample, ABIDE, and HBN), we defined a consensus positive-

association network and a consensus negative-association network as edges that appear in at least

6/10 folds in 300/500 iterations of CPM. This process resulted in 1,001 edges in the positive-

association network and 1,013 edges in the negative-association network; hereafter, we refer to

the collection of edges in the positive and negative networks as the 'ADOS consensus network.'

We note the size of the ADOS consensus network is consistent with other CPM networks that

have generalized[36,136,137]. To ensure generalizability results were robust, we tested summary

networks of varying sizes (from liberal cases where an edge appeared in at least 1/10 folds and

50/500 iterations, to more stringent thresholds where an edge must appear in 10/10 folds and

500/500 iterations, moving in intervals of 1 fold and 50 iterations for each summary network).

To determine if the network predicted autistic traits, we then used the combined network

strength in the ADOS consensus network and computed model coefficients across the Yale youth

sample[36,67,74]. Model coefficients and the network masks were subsequently applied to the

ABIDE and HBN samples to predict SRS scores. Model performance was determined by

comparing the similarity between predicted and observed behavioral scores using Spearman's

correlation. We used the same approach to determine if the network predicted d' scores.

To further assess generalizability, we repeated testing if the ADOS network predicts d'

and SRS using a multiverse approach. A multiverse analysis assesses how results are affected

by different analytical choices[73]. Specifically, we tested if the ADOS positive and negative

networks generalized; we adjusted models for IQ, age, and sex; and, as mentioned above, we

tested a range of consensus network sizes. We point out the goal of a multiverse approach is

not to determine what pipeline results in the highest prediction performance. Instead, the point

is to assess various analytical scenarios and determine how different modelling choices impact

generalization. As such, we do not perform multiple comparisons correction when assessing

these results.

For completeness' sake, we include the additional multiverse analyses performed in

the Yale youth sample in this section. In this dataset, we adjusted CPM models for sex, age,

and IQ; we also used CPM to predict social affect and restricted and repetitive behavior

scores. Additionally, we assessed how altering the feature selection threshold impacted CPM.

Testing generalizability of the SRS network from HBN

After determining that *Despicable Me* was the highest performing condition for SRS

prediction, we reperformed CPM using the full 10 minutes of data from the functional scans

(recall that comparing how scan condition affected CPM performance forced us to truncate all

scans down to four minutes of data). After performing 500 iterations of CPM predicting SRS

scores in HBN as before (median Spearman's rho = 0.1398; $P = 0.006$), we generated a

consensus network similar to above. Specifically, we required an edge to appear in at least 3/5

folds in 300/500 iterations of CPM. This process resulted in 940 edges in the positive-association

network and 893 edges in the negative-association network. To ensure generalizability results

were robust, we again tested summary networks of varying sizes (from liberal cases where an

edge appeared in at least 1/5 folds and 50/500 iterations, to more stringent thresholds where an

edge must appear in 5/5 folds and 450/500 iterations, moving in intervals of 1 fold and 50

iterations for each summary network. The resulting summary networks were then applied to the

Yale youth sample to determine if the model generalized to predict ADOS scores using the

gradCPT average data.

Data availability

The functional parcellation is available here: (https://www.nitrc.org/frs/?group_id=51).

ABIDE data are available here: (https://fcon_1000.projects.nitrc.org/indi/abide/). HBN data are

available here: (https://fcon_1000.projects.nitrc.org/indi/cmi_healthy_brain_network/). To

protect the sensitive nature of data in the adult attention and Yale youth sample, please reach out

to the corresponding authors for questions about, or access to, these samples.

Code availability

Preprocessing was carried out using freely available software:

(https://medicine.yale.edu/bioimaging/suite/; this leads to the main page of BioImage Suite Web

1.2.0 (current build= 1.2.0, 2020/08/25, 11:43:02). Template scripts that were used for

preprocessing of the functional data, as well as CPM code, are available here:

(https://github.com/clhorien/tasks_versus_rest_in_autism_prediction/tree/main). CPM analyses

were conducted in MATLAB (R2021b).

Acknowledgements

This work was supported by the National Institutes of Health (P50MH115716, T32GM007205 to

C.H. and A.S.G., R25MH119043 to C.H., and TR001864 to A.S.G.). The authors thank Hedwig

Sarofin and Cheryl McMurray for assistance during the MRI sessions and Jitendra Bhawnani for

assistance with hardware and software. The authors also thank Scuddy Fontenelle IV, Emma

Brennan-Wydra, Chitra Banarjee, Rachel Foster, Veda Donthireddy, Kohrissa Joseph, Nicole

Powell, Chaela Nutor, Diogo Fortes, and Maureen Butler for assistance with data collection and

collation.

Author Contributions Statement

CH conceptualized the study, with guidance from DS and RTC. CH performed the analyses with

support from FM, ASG, XS, DO'C, EMRL, ESF, and RTC. CH, KC, and RTC designed the

Yale youth study with support from KP, AV, JCM, FRV, DS; CH collected and processed the

data for this study. MDR and MC designed the adult attention study; MDR collected and

processed the data for this study. EMRL processed the data for the ABIDE study. BDA and LT

processed the data for the HBN study. ESF and TS provided guidance on result interpretation.

CH wrote the manuscript, with contributions from FM, ASG, EMRL, TS, and RTC, and

comments from all authors.

Competing Interests Statement

J.C.M. consults or has consulted with Customer Value Partners, Bridgebio, Determined Health,

Apple, and BlackThorn Therapeutics, has received research funding from Janssen Research and

Development, serves on the Scientific Advisory Boards of Pastorus and Modern Clinics, and

receives royalties from Guilford Press, Lambert, Oxford, and Springer. B.D.A. is a founder of

Elevation Admissions.

Figure Legends

Figure 1. An overview of the datasets used in this study. The Yale youth sample was the first
dataset used. FMRI connectivity data from different scanning conditions (task and rest) were
used to generate connectome-based predictive models of Autism Diagnostic Observation
Schedule (ADOS) scores (red circular arrows denote a brain-behavior predictive model). A
summary predictive model was then generated and applied to the adult attention sample. The
goal of this step was to determine whether the model generalized to predict attention phenotypes
(d') in an external dataset. The summary model was also applied to ABIDE and to HBN to
determine if the model predicted SRS scores in external samples. ABIDE, autism brain imaging
data exchange; ADOS, Autism Diagnostic Observation Schedule; CPM, connectome-based
predictive model; d', d-prime (attention phenotype in the Adult Attention Sample); HBN,
Healthy Brain Network; SRS, social responsiveness scale.
27
Figure 2. A schematic showing the scanning conditions used in the Yale youth sample. Top
panel: free-viewing selective social attention task. Four conditions were shown to participants:
no eye contact, no speech (EC-SP-); no eye contact, with speech (EC-SP+); eye contact, with no
speech (EC+SP-); and eye contact, with speech (EC+SP+). No responses were required of

participants during the different conditions. Middle panel: the gradual onset continuous
performance task (gradCPT) was used as a sustained attention task. Grayscale pictures of cities
and mountains were presented with images gradually transitioning from one to the next; button
presses were required for city scenes and withheld for mountain scenes. Bottom panel: resting-
state condition, in which the participants viewed a fixation cross. Please see the Methods section
for further details about each scanning condition.
7
Figure 3. CPM prediction performance across different scanning conditions for total ADOS
scores. The scan condition is shown on the x-axis; on the y-axis, Spearman's rho is shown for
the correlation of predicted and actual ADOS scores. For each condition, the median of the
500 iterations is shown as a solid black line in the violin plot; quartiles, as dotted lines. The
Selective Social Attention task conditions are shown in purple, gradCPT in turquoise, and
resting-state data in yellow. Statistical significance was obtained using permutation testing as
described in the Methods. Note that one-side significance testing was used, as we are interested
in statistically significant positive predictions only. Asterisk (*) indicates statistical significance
after correcting for multiple comparisons using the Benjamini–Hochberg method[135] (correcting
for ten tests: two for gradCPT, four for SSA, two for resting-state, one for gradCPT average, and
one for resting-state average). Exact Spearman's rho and P-values for statistically significant
conditions, gradCPT1: Spearman's rho = 0.441, P-value = 0.004; gradCPT average:
Spearman's rho = 0.445, P-value = 0.002. ADOS, autism diagnostic observation schedule;
Avg, average; EC-SP-, no eye contact, no speech; EC-SP+, no eye contact, with speech; EC+SP-
, eye contact, with no speech; EC+SP+, eye contact, with speech; mm, millimeters; SSA,
selective social attention task.
24
Figure 4. Generalization of the ADOS consensus network in the adult attention sample. A.
Results using the combined network model. B. Results using the positive-association network
model. C. Results using the negative-association network model. Predicted ADOS scores are
indicated on the y-axis; d' scores, on the x-axis. Higher predicted ADOS scores are associated
with lower d' scores, indicating poorer performance on the task and implying lower sustained
attention. Data are presented as a regression line (solid line) +/- 95% confidence intervals
(shaded areas). Spearman's rho was calculated and was used to assess statistical significance
(two-sided). The Benjamini–Hochberg method[135] was used to correct for three tests. ADOS,
autism diagnostic observation schedule; $P\square=\square P$-value.
34
Figure 5. Generalization of the ADOS consensus network to ABIDE. A. SRS total score
results. B. SRS communication score results. C. SRS mannerism score results. D. SRS
motivation score results. For all plots, actual SRS scores from each subscale are indicated on
the x-axis; predicted scores, on the y-axis. Data are presented as a regression line (solid line) +/-
95% confidence intervals (shaded areas). Spearman's rho was calculated and was used to
assess statistical significance (two-sided). The Benjamini–Hochberg method[135] was used to
correct for four tests. ADOS, autism diagnostic observation schedule; $P\square=\square P$-value.
42
43 Figure 6. Neuroanatomy of ADOS consensus network. A. The positive-association network.
44 B. The negative-association network. For both A and B: a circle plot is shown in the upper left.
45 The top of the circle represents anterior; the bottom, posterior. The left half of the circle plot
46 corresponds to the left hemisphere of the brain. A legend indicating the approximate anatomic

'lobe' is shown to the left. The same edges are plotted in the glass brains as lines connecting different nodes (red circles); in these visualizations, nodes are sized according to degree, the number of edges connected to that node. To aid in visualization, we have thresholded the matrices to only show nodes with a degree threshold > 25. C. Matrix of the positive-association network. D. Matrix of the negative-association network. For both C and D: The proportion of edges in a given network pair is shown; data have been corrected to account for differing network size. MF, medial frontal; FP, frontoparietal; DM, default mode; MT, motor; VI, visual I; VII, visual II; VA, visual association; CO, cingulo-opercular; SB, subcortical; CB, cerebellum.

References

Zeidan, J. *et al.* Global prevalence of autism: A systematic review update. *Autism Res* **15**, 778-790, doi:10.1002/aur.2696 (2022).

American Psychiatric Assocaition. *Diagnostic and statistical manual of mental disorders: DSM-5.*, (American Psychiatric Association, 2013).

Feczko, E. & Fair, D. A. Methods and Challenges for Assessing Heterogeneity. *Biol Psychiatry* **88**, 9-17, doi:10.1016/j.biopsych.2020.02.015 (2020).

Horien, C. *et al.* Functional Connectome-Based Predictive Modeling in Autism. *Biol Psychiatry*, doi:10.1016/j.biopsych.2022.04.008 (2022).

Insel, T. *et al.* Research Domain Criteria (RDoC): Toward a New Classification Framework for Research on Mental Disorders. *Am J Psychiat* **167**, 748-751, doi:10.1176/appi.ajp.2010.09091379 (2010).

Duan, X., Shan, X., Uddin, L. Q. & Chen, H. The future of disentangling the heterogeneity of autism with neuroimaging studies. *Biol Psychiatry*, doi:10.1016/j.biopsych.2024.08.008 (2024).

Park, S. *et al.* Delineating a Pathway for the Discovery of Functional Connectome Biomarkers of Autism. *Adv Neurobiol* **40**, 511-544, doi:10.1007/978-3-031-69491-2_18 (2024).

Biswal, B., Yetkin, F. Z., Haughton, V. M. & Hyde, J. S. Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. *Magn Reson Med* **34**, 537-541, doi:10.1002/mrm.1910340409 (1995).

Anderson, J. S. *et al.* Functional connectivity magnetic resonance imaging classification of autism. *Brain* **134**, 3742-3754, doi:10.1093/brain/awr263 (2011).

Chen, C. P. *et al.* Diagnostic classification of intrinsic functional connectivity highlights somatosensory, default mode, and visual regions in autism. *Neuroimage Clin* **8**, 238-245, doi:10.1016/j.nicl.2015.04.002 (2015).

Guo, X. Y. *et al.* Diagnosing Autism Spectrum Disorder from Brain Resting-State Functional Connectivity Patterns Using a Deep Neural Network with a Novel Feature Selection Method. *Front Neurosci-Switz* **11**, doi:ARTN 460 10.3389/fnins.2017.00460 (2017).

Iidaka, T. Resting state functional magnetic resonance imaging and neural network classified autism and control. *Cortex* **63**, 55-67, doi:10.1016/j.cortex.2014.08.011 (2015).

Jahedi, A., Nasamran, C. A., Faires, B., Fan, J. & Muller, R. A. Distributed Intrinsic Functional Connectivity Patterns Predict Diagnostic Status in Large Autism Cohort. *Brain Connect* **7**, 515-525, doi:10.1089/brain.2017.0496 (2017).

Abraham, A. *et al.* Deriving reproducible biomarkers from multi-site resting-state data: An Autism-based example. *Neuroimage* **147**, 736-745, doi:10.1016/j.neuroimage.2016.10.045 (2017).

Emerson, R. W. *et al.* Functional neuroimaging of high-risk 6-month-old infants predicts a diagnosis of autism at 24 months of age. *Sci Transl Med* **9**, doi:ARTN eaag2882 10.1126/scitranslmed.aag2882 (2017).

Chen, H. *et al.* Multivariate classification of autism spectrum disorder using frequency-specific resting-state functional connectivity--A multi-center study. *Prog Neuropsychopharmacol Biol Psychiatry* **64**, 1-9, doi:10.1016/j.pnpbp.2015.06.014 (2016).

Uddin, L. Q. *et al.* Salience network-based classification and prediction of symptom severity in children with autism. *Jama Psychiat* **70**, 869-879, doi:10.1001/jamapsychiatry.2013.104 (2013).

Yahata, N. *et al.* A small number of abnormal brain connections predicts adult autism spectrum disorder. *Nat Commun* **7**, 11254, doi:10.1038/ncomms11254 (2016).

Hong, S. J. *et al.* Atypical functional connectome hierarchy in autism. *Nat Commun* **10**, 1022, doi:10.1038/s41467-019-08944-1 (2019).

Ilioska, I. *et al.* Connectome-wide Mega-analysis Reveals Robust Patterns of Atypical Functional Connectivity in Autism. *Biol Psychiatry* **94**, 29-39, doi:10.1016/j.biopsych.2022.12.018 (2023).

Xiao, J. *et al.* Linked Social-Communication Dimensions and Connectivity in Functional Brain Networks in Autism Spectrum Disorder. *Cereb Cortex* **31**, 3899-3910, doi:10.1093/cercor/bhab057 (2021).

Finn, E. S. *et al.* Functional connectome fingerprinting: identifying individuals using patterns of brain connectivity. *Nat Neurosci* **18**, 1664-1671, doi:10.1038/nn.4135 (2015).

Shen, X. L. *et al.* Using connectome-based predictive modeling to predict individual behavior from brain connectivity. *Nat Protoc* **12**, 506-518, doi:10.1038/nprot.2016.178 (2017).

Lake, E. M. R. *et al.* The Functional Brain Organization of an Individual Allows Prediction of Measures of Social Abilities Transdiagnostically in Autism and Attention-Deficit/Hyperactivity Disorder. *Biol Psychiatry* **86**, 315-326, doi:10.1016/j.biopsych.2019.02.019 (2019).

Ma, X. *et al.* Connectome-based prediction of the severity of autism spectrum disorder. *Psychoradiology* **3**, kkad027, doi:10.1093/psyrad/kkad027 (2023).

Rohr, C. S., Kamal, S. & Bray, S. Building functional connectivity neuromarkers of behavioral self-regulation across children with and without Autism Spectrum Disorder. *Dev Cogn Neuros-Neth* **41**, doi:ARTN 100747 10.1016/j.dcn.2019.100747 (2020).

Dufford, A., Kimble, V., Tejavibulya, L., Dadashkarimi, J. & Scheinost, D. Predicting Transdiagnostic Social Impairments in Childhood Using Connectome-Based Predictive Modeling. *Biol Psychiat* **91**, S87, doi:10.1016/j.biopsych.2022.02.234 (2022).

Horien, C. *et al.* A generalizable connectome-based marker of in-scan sustained attention in neurodiverse youth. *Cereb Cortex* **33**, 6320-6334, doi:10.1093/cercor/bhac506 (2023).

Finn, E. S. Is it time to put rest to rest? *Trends Cogn Sci* **25**, 1021-1032, doi:10.1016/j.tics.2021.09.005 (2021).

O'Connor, D., Horien, C, Mandino, F, Constable, RT. Identifying dynamic reproducible brain states using a predictive modelling approach. *bioRxiv*, doi:https://doi.org/10.1101/2022.10.14.512147 (2022).

Zhang, X., Hulvershorn, LA, Constable, RT, Zhao, Y, Wang, S. Cost efficiency of fMRI studies using resting-state vs task-based functional connectivity. *arXiv*, doi: https://doi.org/10.48550/arXiv.2411.01092 (2024).

Ramduny, J. & Kelly, C. Connectome-based fingerprinting: reproducibility, precision, and behavioral prediction. *Neuropsychopharmacology* **50**, 114-123, doi:10.1038/s41386-024-01962-8 (2024).

Greene, A. S., Gao, S. Y., Scheinost, D. & Constable, R. T. Task-induced brain state manipulation improves prediction of individual traits. *Nat Commun* **9**, doi:ARTN 2807 10.1038/s41467-018-04920-3 (2018).

Jiang, R. T. *et al.* Task-induced brain connectivity promotes the detection of individual differences in brain-behavior relationships. *Neuroimage* **207**, doi:ARTN 116370 10.1016/j.neuroimage.2019.116370 (2020).

Greene, A. S., Gao, S., Noble, S., Scheinost, D. & Constable, R. T. How Tasks Change Whole-Brain Functional Organization to Reveal Brain-Phenotype Relationships. *Cell Rep* **32**, 108066, doi:10.1016/j.celrep.2020.108066 (2020).

Rosenberg, M. D. *et al.* A neuromarker of sustained attention from whole-brain functional connectivity. *Nat Neurosci* **19**, 165-+, doi:10.1038/nn.4179 (2016).

Yoo, K. *et al.* A brain-based general measure of attention. *Nat Hum Behav* **6**, 782-795, doi:10.1038/s41562-022-01301-1 (2022).

Ju, S. *et al.* Connectome-based predictive modeling shows sex differences in brain-based predictors of memory performance. *Front Dement* **2**, 1126016, doi:10.3389/frdem.2023.1126016 (2023).

Avery, E. W. *et al.* Distributed Patterns of Functional Connectivity Predict Working Memory Performance in Novel Healthy and Memory-impaired Individuals. *J Cognitive Neurosci* **32**, 241-255, doi:10.1162/jocn_a_01487 (2020).

Hardikar, S. *et al.* Personality traits vary in their association with brain activity across situations. *Commun Biol* **7**, 1498, doi:10.1038/s42003-024-07061-0 (2024).

Finn, E. S. & Bandettini, P. A. Movie-watching outperforms rest for functional connectivity-based prediction of behavior. *Neuroimage* **235**, 117963, doi:10.1016/j.neuroimage.2021.117963 (2021).

Finn, E. S. *et al.* Can brain state be manipulated to emphasize individual differences in functional connectivity? *Neuroimage* **160**, 140-151, doi:10.1016/j.neuroimage.2017.03.064 (2017).

Keehn, B., Nair, A., Lincoln, A. J., Townsend, J. & Muller, R. A. Under-reactive but easily distracted: An fMRI investigation of attentional capture in autism spectrum disorder. *Dev Cogn Neurosci* **17**, 46-56, doi:10.1016/j.dcn.2015.12.002 (2016).

Keehn, B., Shih, P., Brenner, L. A., Townsend, J. & Muller, R. A. Functional connectivity for an "Island of sparing" in autism spectrum disorder: An fMRI study of visual search. *Hum Brain Mapp* **34**, 2524-2537, doi:10.1002/hbm.22084 (2013).

Vaidya, C. J. *et al.* Controlling attention to gaze and arrows in childhood: an fMRI study of typical development and Autism Spectrum Disorders. *Dev Sci* **14**, 911-924, doi:10.1111/j.1467-7687.2011.01041.x (2011).

Rahko, J. S. *et al.* Attention and Working Memory in Adolescents with Autism Spectrum Disorder: A Functional MRI Study. *Child Psychiatry Hum Dev* **47**, 503-517, doi:10.1007/s10578-015-0583-6 (2016).

Fitzgerald, J. *et al.* Disrupted functional connectivity in dorsal and ventral attention networks during attention orienting in autism spectrum disorders. *Autism Res* **8**, 136-152, doi:10.1002/aur.1430 (2015).

Kernbach, J. M. *et al.* Shared endo-phenotypes of default mode dsfunction in attention deficit/hyperactivity disorder and autism spectrum disorder. *Transl Psychiatry* **8**, 133, doi:10.1038/s41398-018-0179-6 (2018).

Stevens T, P. L., Barnard-Brak L. The comorbidity of ADHD in children diagnosed with autism spectrum disorder. *Research in Autism Spectrum Disorders* **31**, 11-18 (2016).

Reiersen, A. M. & Todd, R. D. Co-occurrence of ADHD and autism spectrum disorders: phenomenology and treatment. *Expert Rev Neurother* **8**, 657-669, doi:10.1586/14737175.8.4.657 (2008).

Antshel, K. M. & Russo, N. Autism Spectrum Disorders and ADHD: Overlapping Phenomenology, Diagnostic Issues, and Treatment Considerations. *Curr Psychiatry Rep* **21**, 34, doi:10.1007/s11920-019-1020-5 (2019).

Alvarez-Fernandez, S. *et al.* Perceived social support in adults with autism spectrum disorder and attention-deficit/hyperactivity disorder. *Autism Res* **10**, 866-877, doi:10.1002/aur.1735 (2017).

Esterman, M., Noonan, S. K., Rosenberg, M. & DeGutis, J. In the Zone or Zoning Out? Tracking Behavioral and Neural Fluctuations During Sustained Attention. *Cereb Cortex* **23**, 2712-2723, doi:10.1093/cercor/bhs261 (2013).

Rosenberg, M., Noonan, S., DeGutis, J. & Esterman, M. Sustaining visual attention in the face of distraction: a novel gradual-onset continuous performance task. *Atten Percept Psycho* **75**, 426-439, doi:10.3758/s13414-012-0413-x (2013).

Chawarska, K., Macari, S. & Shic, F. Context modulates attention to social scenes in toddlers with autism. *J Child Psychol Psychiatry* **53**, 903-913, doi:10.1111/j.1469-7610.2012.02538.x (2012).

Chawarska, K., Macari, S. & Shic, F. Decreased spontaneous attention to social scenes in 6-month-old infants later diagnosed with autism spectrum disorders. *Biol Psychiatry* **74**, 195-203, doi:10.1016/j.biopsych.2012.11.022 (2013).

Shic, F., Macari, S. & Chawarska, K. Speech disturbs face scanning in 6-month-old infants who develop autism spectrum disorder. *Biol Psychiatry* **75**, 231-237, doi:10.1016/j.biopsych.2013.07.009 (2014).

Shic, F., Wang, Q., Macari, S. L. & Chawarska, K. The role of limited salience of speech in selective attention to faces in toddlers with autism spectrum disorders. *J Child Psychol Psychiatry* **61**, 459-469, doi:10.1111/jcpp.13118 (2020).

Campbell, D. J., Shic, F., Macari, S. & Chawarska, K. Gaze response to dyadic bids at 2 years related to outcomes at 3 years in autism spectrum disorders: a subtyping analysis. *J Autism Dev Disord* **44**, 431-442, doi:10.1007/s10803-013-1885-9 (2014).

Horien, C. *et al.* Low-motion fMRI data can be obtained in pediatric participants undergoing a 60-minute scan protocol. *Sci Rep-Uk* **10**, doi:ARTN 21855

10.1038/s41598-020-78885-z (2020).

Ingersoll, B. Broader autism phenotype and nonverbal sensitivity: evidence for an association in the general population. *J Autism Dev Disord* **40**, 590-598, doi:10.1007/s10803-009-0907-0 (2010).

Lord C, R. M., DiLavore PC, Risi S, Gotham K, Bishop S. *Autism Diagnostic Observation Schedule, Second Edition.*, (Western Psychological Services, 2012).

Greene, A. S. *et al.* Brain-phenotype models fail for individuals who defy sample stereotypes. *Nature* **609**, 109-118, doi:10.1038/s41586-022-05118-w (2022).

Horien, C. *et al.* Considering factors affecting the connectome-based identification process: Comment on Waller et al. *Neuroimage* **169**, 172-175, doi:10.1016/j.neuroimage.2017.12.045 (2018).

Rapuano, K. M. *et al.* Behavioral and brain signatures of substance use vulnerability in childhood. *Dev Cogn Neurosci* **46**, 100878, doi:10.1016/j.dcn.2020.100878 (2020).

Shen, X., Tokoglu, F., Papademetris, X. & Constable, R. T. Groupwise whole-brain parcellation from resting-state fMRI data for network node identification. *Neuroimage* **82**, 403-415, doi:10.1016/j.neuroimage.2013.05.081 (2013).

Dufford, A. J. *et al.* Predicting Transdiagnostic Social Impairments in Childhood using Connectome-based Predictive Modeling. *medRxiv*, 2022.2004.2007.22273518, doi:10.1101/2022.04.07.22273518 (2022).

Scheinost, D. *et al.* Functional connectivity during frustration: a preliminary study of predictive modeling of irritability in youth. *Neuropsychopharmacology* **46**, 1300-1306, doi:10.1038/s41386-020-00954-8 (2021).

Taxali, A., Angstadt, M., Rutherford, S. & Sripada, C. Boost in Test-Retest Reliability in Resting State fMRI with Predictive Modeling. *Cereb Cortex* **31**, 2822-2833, doi:10.1093/cercor/bhaa390 (2021).

Birn, R. M. *et al.* The effect of scan length on the reliability of resting-state fMRI connectivity estimates. *Neuroimage* **83**, 550-558, doi:10.1016/j.neuroimage.2013.05.099 (2013).

Laumann, T. O. *et al.* Functional System and Areal Organization of a Highly Sampled Individual Human Brain. *Neuron* **87**, 657-670, doi:10.1016/j.neuron.2015.06.037 (2015).

Noble, S. *et al.* Influences on the Test-Retest Reliability of Functional Connectivity MRI and its Relationship with Behavioral Utility. *Cereb Cortex* **27**, 5415-5429, doi:10.1093/cercor/bhx230 (2017).

Steegen, S., Tuerlinckx, F., Gelman, A. & Vanpaemel, W. Increasing Transparency Through a Multiverse Analysis. *Perspect Psychol Sci* **11**, 702-712, doi:10.1177/1745691616658637 (2016).

Ju, Y. *et al.* Connectome-based models can predict early symptom improvement in major depressive disorder. *J Affect Disord* **273**, 442-452, doi:10.1016/j.jad.2020.04.028 (2020).

Li, J. *et al.* Global signal regression strengthens association between resting-state functional connectivity and behavior. *Neuroimage* **196**, 126-141, doi:10.1016/j.neuroimage.2019.04.016 (2019).

Di Martino, A. *et al.* Data Descriptor: Enhancing studies of the connectome in autism using the autism brain imaging data exchange II. *Scientific Data* **4**, doi:ARTN 170010 10.1038/sdata.2017.10 (2017).

Di Martino, A. *et al.* The autism brain imaging data exchange: towards a large-scale evaluation of the intrinsic brain architecture in autism. *Molecular Psychiatry* **19**, 659-667, doi:10.1038/mp.2013.78 (2014).

Constantino, J. N. *et al.* Validation of a brief quantitative measure of autistic traits: comparison of the social responsiveness scale with the autism diagnostic interview-revised. *J Autism Dev Disord* **33**, 427-433, doi:10.1023/a:1025014929212 (2003).

Koshino, H. *et al.* Functional connectivity in an fMRI working memory task in high-functioning autism. *Neuroimage* **24**, 810-821, doi:10.1016/j.neuroimage.2004.09.028 (2005).

Keehn, B., Shih, P., Brenner, L. A., Townsend, J. & Muller, R. A. Functional connectivity for an "island of sparing" in autism spectrum disorder: an fMRI study of visual search. *Hum Brain Mapp* **34**, 2524-2537, doi:10.1002/hbm.22084 (2013).

McGrath, J. *et al.* Atypical visuospatial processing in autism: insights from functional connectivity analysis. *Autism Res* **5**, 314-330, doi:10.1002/aur.1245 (2012).

Gilbert, S. J., Bird, G., Brindley, R., Frith, C. D. & Burgess, P. W. Atypical recruitment of medial prefrontal cortex in autism spectrum disorders: an fMRI study of two executive function tasks. *Neuropsychologia* **46**, 2281-2291, doi:10.1016/j.neuropsychologia.2008.03.025 (2008).

Just, M. A., Cherkassky, V. L., Keller, T. A., Kana, R. K. & Minshew, N. J. Functional and anatomical cortical underconnectivity in autism: evidence from an FMRI study of an executive function task and corpus callosum morphometry. *Cereb Cortex* **17**, 951-961, doi:10.1093/cercor/bhl006 (2007).

Knaus, T. A., Silver, A. M., Lindgren, K. A., Hadjikhani, N. & Tager-Flusberg, H. fMRI activation during a language task in adolescents with ASD. *J Int Neuropsychol Soc* **14**, 967-979, doi:10.1017/S1355617708081216 (2008).

Greene, A. S., Horien, C., Barson, D., Scheinost, D. & Constable, R. T. Why is everyone talking about brain state? *Trends Neurosci* **46**, 508-524, doi:10.1016/j.tins.2023.04.001 (2023).

Finn, E. S. & Rosenberg, M. D. Beyond fingerprinting: Choosing predictive connectomes over reliable connectomes. *Neuroimage* **239**, 118254, doi:10.1016/j.neuroimage.2021.118254 (2021).

Rai, S., Graff, K., Tansey, R. & Bray, S. How do tasks impact the reliability of fMRI functional connectivity? *Hum Brain Mapp* **45**, e26535, doi:10.1002/hbm.26535 (2024).

Greene, A. S., Gao, S. Y., Noble, S., Scheinost, D. & Constable, R. T. How Tasks Change Whole-Brain Functional Organization to Reveal Brain-Phenotype Relationships. *Cell Rep* **32**, doi:ARTN 108066 10.1016/j.celrep.2020.108066 (2020).

Sripada, C., Angstadt, M., Rutherford, S., Taxali, A. & Shedden, K. Toward a "treadmill test" for cognition: Improved prediction of general cognitive ability from the task activated brain. *Hum Brain Mapp* **41**, 3186-3197, doi:10.1002/hbm.25007 (2020).

Durston, S. *et al.* Differential patterns of striatal activation in young children with and without ADHD. *Biol Psychiatry* **53**, 871-878, doi:10.1016/s0006-3223(02)01904-2 (2003).

Yerys, B. E. *et al.* The fMRI success rate of children and adolescents: typical development, epilepsy, attention deficit/hyperactivity disorder, and autism spectrum disorders. *Hum Brain Mapp* **30**, 3426-3435, doi:10.1002/hbm.20767 (2009).

Pecci-Terroba, C. *et al.* Subgrouping autism and ADHD based on structural MRI population modelling centiles. *Mol Autism* **16**, 33, doi:10.1186/s13229-025-00667-z (2025).

Di Martino, A. *et al.* Shared and distinct intrinsic functional network centrality in autism and attention-deficit/hyperactivity disorder. *Biol Psychiatry* **74**, 623-632, doi:10.1016/j.biopsych.2013.02.011 (2013).

Itahashi, T. *et al.* Neural correlates of shared sensory symptoms in autism and attention-deficit/hyperactivity disorder. *Brain Commun* **2**, fcaa186, doi:10.1093/braincomms/fcaa186 (2020).

Greene, A. S. & Constable, R. T. Clinical Promise of Brain-Phenotype Modeling: A Review. *Jama Psychiat* **80**, 848-854, doi:10.1001/jamapsychiatry.2023.1419 (2023).

Nichols, A. L. A., Eichler, T., Latham, R. & Zimmer, M. A global brain state underlies C. elegans sleep behavior. *Science* **356**, doi:10.1126/science.aam6851 (2017).

Dal Monte, O., Chu, C. C. J., Fagan, N. A. & Chang, S. W. C. Specialized medial prefrontal-amygdala coordination in other-regarding decision preference. *Nat Neurosci* **23**, 565-574, doi:10.1038/s41593-020-0593-y (2020).

Harikumar, A., Evans, D. W., Dougherty, C. C., Carpenter, K. L. H. & Michael, A. M. A Review of the Default Mode Network in Autism Spectrum Disorders and Attention Deficit Hyperactivity Disorder. *Brain Connect* **11**, 253-263, doi:10.1089/brain.2020.0865 (2021).

Menon, V. 20 years of the default mode network: A review and synthesis. *Neuron* **111**, 2469-2487, doi:10.1016/j.neuron.2023.04.023 (2023).

Norman, L. J. *et al.* Cross-sectional mega-analysis of resting-state alterations associated with autism and attention-deficit/hyperactivity disorder in children and adolescents. *Nature Mental Health* **3**, 709-723, doi:10.1038/s44220-025-00431-5 (2025).

Chopra, S. *et al.* Generalizable and replicable brain-based predictions of cognitive functioning across common psychiatric illness. *Sci Adv* **10**, eadn1862, doi:10.1126/sciadv.adn1862 (2024).

Poldrack, R. A. Inferring mental states from neuroimaging data: from reverse inference to large-scale decoding. *Neuron* **72**, 692-697, doi:10.1016/j.neuron.2011.11.001 (2011).

Mundy, P. A review of joint attention and social-cognitive brain systems in typical development and autism spectrum disorder. *Eur J Neurosci* **47**, 497-514, doi:10.1111/ejn.13720 (2018).

Segura, P. *et al.* Connectome-based symptom mapping and in silico related gene expression in children with autism and/or attention-deficit/hyperactivity disorder. *medRxiv*, 2024.2012.2009.24318621, doi:10.1101/2024.12.09.24318621 (2024).

Caspi, A. *et al.* The p Factor: One General Psychopathology Factor in the Structure of Psychiatric Disorders? *Clin Psychol Sci* **2**, 119-137, doi:10.1177/2167702613497473 (2014).

Watts, A. L., Greene, A. L., Bonifay, W. & Fried, E. I. A critical evaluation of the -factor literature. *Nat Rev Psychol* **3**, 73-90, doi:10.1038/s44159-023-00260-2 (2024).

Kafri, R., Levy, M. & Pilpel, Y. The regulatory utilization of genetic redundancy through responsive backup circuits. *Proc Natl Acad Sci U S A* **103**, 11653-11658, doi:10.1073/pnas.0604883103 (2006).

Tononi, G., Sporns, O. & Edelman, G. M. Measures of degeneracy and redundancy in biological networks. *Proc Natl Acad Sci U S A* **96**, 3257-3262, doi:10.1073/pnas.96.6.3257 (1999).

Edelman, G. M. & Gally, J. A. Degeneracy and complexity in biological systems. *Proc Natl Acad Sci U S A* **98**, 13763-13768, doi:10.1073/pnas.231499798 (2001).

Emerson, R. W. *et al.* Functional neuroimaging of high-risk 6-month-old infants predicts a diagnosis of autism at 24 months of age. *Sci Transl Med* **9**, doi:10.1126/scitranslmed.aag2882 (2017).

Tang, S. *et al.* Reconciling Dimensional and Categorical Models of Autism Heterogeneity: A Brain Connectomics and Behavioral Study. *Biol Psychiatry* **87**, 1071-1082, doi:10.1016/j.biopsych.2019.11.009 (2020).

Chou, Y. H. *et al.* Maintenance and Representation of Mind Wandering during Resting-State fMRI. *Sci Rep* **7**, 40722, doi:10.1038/srep40722 (2017).

Bettoni, R. *et al.* Learning and generalization of repetition-based rules in autism. *Psychol Res* **87**, 1429-1438, doi:10.1007/s00426-022-01761-0 (2023).

Bedford, S. A. *et al.* Large-scale analyses of the relationship between sex, age and intelligence quotient heterogeneity and cortical morphometry in autism spectrum disorder. *Mol Psychiatry* **25**, 614-628, doi:10.1038/s41380-019-0420-6 (2020).

Lai, M. C. *et al.* Imaging sex/gender and autism in the brain: Etiological implications. *J Neurosci Res* **95**, 380-397, doi:10.1002/jnr.23948 (2017).

Floris, D. L. *et al.* Towards robust and replicable sex differences in the intrinsic brain function of autism. *Mol Autism* **12**, 19, doi:10.1186/s13229-021-00415-z (2021).

Chouinard, B., Gallagher, L. & Kelly, C. He said, she said: Autism spectrum diagnosis and gender differentially affect relationships between executive functions and social communication. *Autism* **23**, 1793-1804, doi:10.1177/1362361318815639 (2019).

Vanderwal, T., Eilbott, J. & Castellanos, F. X. Movies in the magnet: Naturalistic paradigms in developmental functional neuroimaging. *Dev Cogn Neurosci* **36**, 100600, doi:10.1016/j.dcn.2018.10.004 (2019).

Manoli, A., Van Overwalle, F., Grosse Wiesmann, C. & Valk, S. L. Functional recruitment and connectivity of the cerebellum is associated with the emergence of Theory of Mind in early childhood. *Nat Commun* **16**, 5273, doi:10.1038/s41467-025-60523-9 (2025).

Rosenberg, M. D. & Finn, E. S. How to establish robust brain-behavior relationships without thousands of individuals. *Nat Neurosci* **25**, 835-837, doi:10.1038/s41593-022-01110-9 (2022).

Spisak, T., Bingel, U. & Wager, T. D. Multivariate BWAS can be replicable with moderate sample sizes. *Nature* **615**, E4-E7, doi:10.1038/s41586-023-05745-x (2023).

Kang, K. *et al.* Study design features increase replicability in brain-wide association studies. *Nature*, doi:10.1038/s41586-024-08260-9 (2024).

Kiar, G. *et al.* Why experimental variation in neuroimaging should be embraced. *Nat Commun* **15**, 9411, doi:10.1038/s41467-024-53743-y (2024).

Ooi, L. Q. R. *et al.* Longer scans boost prediction and cut costs in brain-wide association studies. *Nature*, doi:10.1038/s41586-025-09250-1 (2025).

Marek, S. *et al.* Reproducible brain-wide association studies require thousands of individuals. *Nature* **603**, 654-660, doi:10.1038/s41586-022-04492-9 (2022).

Alexander, L. M. *et al.* An open resource for transdiagnostic research in pediatric mental health and learning disorders. *Sci Data* **4**, 170181, doi:10.1038/sdata.2017.181 (2017).

Adkinson, B. D. *et al.* Brain-phenotype predictions of language and executive function can survive across diverse real-world data: Dataset shifts in developmental populations. *Dev Cogn Neurosci* **70**, 101464, doi:10.1016/j.dcn.2024.101464 (2024).

Joshi, A. *et al.* Unified framework for development, deployment and robust testing of neuroimaging algorithms. *Neuroinformatics* **9**, 69-84, doi:10.1007/s12021-010-9092-8 (2011).

Lutkenhoff, E. S. *et al.* Optimized brain extraction for pathological brains (optiBET). *Plos One* **9**, e115551, doi:10.1371/journal.pone.0115551 (2014).

Finn, E. S. *et al.* Functional connectome fingerprinting: identifying individuals using patterns of brain connectivity. *Nat Neurosci* **18**, 1664-1671, doi:10.1038/nn.4135 (2015).

Satterthwaite, T. D. *et al.* An improved framework for confound regression and filtering for control of motion artifact in the preprocessing of resting-state functional connectivity data. *Neuroimage* **64**, 240-256, doi:10.1016/j.neuroimage.2012.08.052 (2013).

Horien, C., Shen, X. L., Scheinost, D. & Constable, R. T. The individual functional connectome is unique and stable over months to years. *Neuroimage* **189**, 676-687, doi:10.1016/j.neuroimage.2019.02.002 (2019).

Buckner, R. L., Krienen, F. M., Castellanos, A., Diaz, J. C. & Yeo, B. T. The organization of the human cerebellum estimated by intrinsic functional connectivity. *J Neurophysiol* **106**, 2322-2345, doi:10.1152/jn.00339.2011 (2011).

Scheinost, D. *et al.* Ten simple rules for predictive modeling of individual differences in neuroimaging. *Neuroimage* **193**, 35-45, doi:10.1016/j.neuroimage.2019.02.057 (2019).

Benjamini, Y., Hochberg, Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society, Series B* **57**, 289-300 (1995).

Yip, S. W., Scheinost, D., Potenza, M. N. & Carroll, K. M. Connectome-Based Prediction of Cocaine Abstinence. *Am J Psychiatry* **176**, 156-164, doi:10.1176/appi.ajp.2018.17101147 (2019).

Rosenberg, M. D. *et al.* Methylphenidate Modulates Functional Network Connectivity to Enhance Attention. *J Neurosci* **36**, 9547-9557, doi:10.1523/Jneurosci.1746-16.2016 (2016).