High Prevalence and Genotypic Diversity of Persistent *Chlamydia trachomatis* Infections Among South African Adolescent Girls and Young Women: A Tale of Two Cities

Smritee Dabee,[1,2] Shaun Barnabas,[1,3] Bart Versteeg,[4] Brian Kullin,[1,2] Shameem Z. Jaumdally,[1] Hoyam Gamieldien,[1,2] Nonhlanhla Mkhize,[5] Etienne Muller,[5] Venessa Maseko,[5] Katherine Gill,[1,3] Darren P. Martin,[1] Glenda Gray,[6,7] Linda-Gail Bekker[1,3] David A. Lewis,[8,9] Heather B. Jaspan,[1,10] Sylvia M. Bruisten,[11,12] Jo-Ann S. Passmore[1,2,13#]

[1]Institute of Infectious Disease and Molecular Medicine (IDM), Division of Medical Virology, Dept. Pathology, University of Cape Town, Cape Town, South Africa; [2]DST-NRF CAPRISA Centre of Excellence in HIV Prevention, University of Cape Town, Cape Town, South Africa; [3]Desmond Tutu HIV Centre, University of Cape Town, South Africa; [4]Knowledge institute of the Dutch association of Medical Specialists, Utrecht, Netherlands; [5]Centre for HIV and STIs, National Institute for Communicable Disease, National Health Laboratory Service, South Africa; [6]Perinatal HIV Research Unit, Faculty of Health Sciences, University of the Witwatersrand, Diepkloof, Johannesburg, South Africa; [7]South African Medical Research Council, Cape Town, South Africa; [8]Western Sydney Sexual Health Centre, Western Sydney Local Health District, Parramatta, Australia; [9]Sydney Infectious Diseases Institute & Sydney Medical School-Westmead, University of Sydney, Sydney, Australia; [10]Seattle Children's Research Institute, University of Washington, Seattle, Washington, USA; [11]Department of Infectious Diseases, Public Health Service Amsterdam, Amsterdam, the Netherlands; [12]Amsterdam Infection and Immunity Institute, Academic Medical Center, University of Amsterdam, Amsterdam, the Netherlands; [13]National Health Laboratory Service, Groote Schuur Hospital, Cape Town, South Africa

#Address correspondence to Professor Jo-Ann S. Passmore, Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Medical School, Anzio Road, Observatory, 7925 Cape Town, South Africa. E-mail: Jo-ann.Passmore@uct.ac.za, Tel: +21-21-6507963

Word count: 1999, Abstract (95), Figures: 2, Supplementary Tables: 1

Running title: Chlamydia Genotypes in South Africa

Summary: *Chlamydia trachomatis* infection is highly prevalent and asymptomatic among South African adolescent girls and young women, with marked regional and genotypic diversity. Frequent persistence or reinfection highlights the need for improved screening, surveillance, and targeted prevention strategies.

Abstract

Chlamydia trachomatis (CT) is the most common sexually transmitted bacterial infection globally, significantly affecting adolescent girls and young women (AGYW) in South Africa. This study investigated CT prevalence and genotypic diversity among 298 AGYW from Cape Town and Johannesburg, revealing an overall prevalence of 29.5%. Prevalence was higher in Cape Town (41.6%) compared to Johannesburg (17.4%; $p<0.0001$). Genetic analysis identified 34 sequence types, including 15 novel variants. Longitudinal data highlighted frequent reinfections or persistent infections despite treatment. These findings underscore the importance of addressing CT's genetic diversity for improved reproductive health strategies and vaccine development.

Words: 95

Keywords: Chlamydia trachomatis, adolescent female, South Africa, molecular epidemiology, multilocus sequence typing, persistence

Introduction

Adolescent girls and young women (AGYW) in sub-Saharan Africa are at heightened risk for

sexually transmitted infections (STIs), including *Chlamydia trachomatis* (CT), which

exacerbates HIV acquisition risks and reproductive complications (1). The asymptomatic

nature of most CT infections in women complicates diagnosis and management (2),

necessitating a deeper understanding of its epidemiology and genotypic diversity. South

Africa shows marked regional variation in CT prevalence and associated complications,

potentially reflecting differences in circulating genotypes linked to inflammation, persistence,

or increased risk (3–5). Characterising CT genotypes can therefore inform regional

epidemiology and support global surveillance efforts.

Here we characterise CT genotypes among AGYW from two urban South African

settings (Cape Town and Johannesburg) using high resolution multi-locus sequence typing

(MLST) to further explore regional differences in the prevalence of different CT genetic

variants (6). These data align with South Africa's genomic surveillance goals, which could be

extremely valuable for monitoring the emergence and spread of pathogenic or treatment-

resistant CT strains (7).

MLST typing of CT variants is also useful in that it can (i) indicate whether

epidemiologically linked patients experiencing recurrent infections are acquiring them from

independent sources, and (ii) differentiate between unrelated new and persistent infections.

Inadequately treated CT infections can persist or recur, perpetuating inflammation and

increasing the risk of long-term complications.

Evaluating CT genotypes associated with AGYW from two geographically distinct

locations in South Africa addresses a gap in our understanding of the interplay between CT

genetic diversity and regional epidemiological patterns of CT infections. Closing this gap

will inform both general STI and HIV prevention strategies, and the design of CT vaccines

tailored to the needs of South African AGYW.

Methods

Cohort Description: This study recruited sexually active AGYW aged 16–22 years from

two urban centres in South Africa: the Desmond Tutu Health Centre (DTHC) in Cape Town

and the Perinatal HIV Research Unit (PHRU) in Johannesburg, using a targeted enrolment

strategy with predefined sample sizes (4). These two major urban sites in South Africa, with

distinct metropolitan geographies, were selected to allow exploration of shared versus site-

specific risk factors. Participants were excluded if they were living with HIV, pregnant, not

sexually active, or if they had douched, used spermicides within the past two days, or taken

antibiotics within the last two weeks. Ethical approval was obtained from the Universities of

Cape Town (UCT HREC #267/2013) and Witwatersrand Research Ethics Committees

(WITS HREC #M130745). Written informed consent was provided by participants aged 18

years or older, while those under 18 years provided written assent alongside parental or

guardian consent. At enrolment, participants completed a questionnaire capturing socio-

demographic characteristics, reproductive health, and sexual behaviour.

Sample Collection: During each clinical visit, a speculum examination was conducted. A

Dacron vulvovaginal swab was collected for STI testing and a lateral vaginal wall

FLOQSwab™ sample for preparing a Gram-stained slide for Nugent scoring to diagnose

bacterial vaginosis (BV) (4). Vulvovaginal swabs were analyzed for discharge-causing STIs,

including *Chlamydia trachomatis* (CT), using real-time multiplex polymerase chain reaction

(PCR) assays (4). BV was categorized based on Nugent scoring: scores of 7–10 indicated BV

positivity, 4–6 were intermediate, and 0–3 were BV-negative. Participants with positive

results received treatment according to South African national guidelines, with a contact slip

being given as part of the care protocol (4).

Chlamydia MLST: The genotypic diversity of *C. trachomatis* was analyzed using MLST

and *ompA* genotyping (6)(7). Sequence data from five non-housekeeping gene loci (Uppsala

typing scheme) were analysed using the CT database hosted on PubMLST. Allelic

combinations that were not present in the CT MLST database were assigned preliminary

sequence type (ST) numbers (ST551-ST560). The *ompA* genovars were assigned by

comparison to an offline reference database described previously (6). BioEdit was used for

sequence editing, with sequences assembled and aligned using the ClustalW tool in MEGA4.

Minimum-spanning trees were generated using BioNumerics (Applied Maths, Sint-Martens-

Latem, Belgium) to explore genetic relationships between strains. To generate a global CT

phylogeny based on the MLST scheme proposed by Klint et al. (8), loci sequences were

downloaded for all currently recognised MLST subtypes from the PubMLST database (9),

and used to generate in-frame concatenated alignments for each ST using mafft (10), used to

construct a maximum likelihood phylogenetic tree using IQ-TREE2 (11) and best fit gene-

based partitioned models selected using ModelFinder (12).

Statistical tests: Differences between sites and age groups were assessed using chi-squared

or Fisher's exact tests for categorical variables and the Kruskal–Wallis test for non-

parametric comparisons. All tests were two-sided with p<0.05 considered significant, and

analyses were performed using Stata v14.2 (StataCorp, College Station, TX, USA).

Results

A total of 298 sexually active AGYW from two communities — Cape Town (n=149) and

Johannesburg (n=149) — were included in this study. Participants from Cape Town were

followed longitudinally over three visits spanning 6–8 months, whereas those from

Johannesburg were enrolled for a single visit (Supplementary Table 1). Overall contraceptive

use varied significantly between the sites (Kruskal Wallis p<0.0001). While all AGYW in

Cape Town used contraception (predominantlyNet-EN; 105/149; 70.5%), in Johannesburg,

90.6% used contraception, with condoms being the most common type (77/149; 63.8%). The

prevalence of treatable STIs in the cohort was notably high, affecting 74.2% of participants

(221/298). Among these, 10.1% (30/298) had multiple concurrent STIs, and 37.2% (111/298)

had both an STI and BV (4).

CT was the most prevalent STI in the study, affecting 88 of 298 participants (29.5%)

at enrolment, with prevalence being higher in Cape Town (62/149; 41.6%) than

Johannesburg (26/149; 17.4%; p<0.0001). All CT cases were asymptomatic (4). Younger

adolescents were disproportionately affected, with a prevalence of 33.8% (69/204) among

those aged 16–19 years, compared to 20.2% (19/94) in those aged 20–22 years (Chi-squared

p=0.02). Among participants infected with CT, 71.6% (63/88) were co-infected with another

STI or had BV (Nugent score 7–10), highlighting the high burden of co-occurring

reproductive tract conditions in this population.

OmpA typing and MLST were performed on 128 samples with available DNA,

including 20/26 cases from Johannesburg and 108/121 from Cape Town. Of these, 18

samples had either insufficient DNA or low bacterial loads. From these CT cases, 34 STs

were identified by MLST, with 25/34 circulating in Cape Town and 12/34 in Johannesburg

(Figure 1A). Of these, 15 were novel STs not previously reported. The three most common

were ST12d (13.3%), ST530b (12%), and ST3 (9.3%). ST12d was the most common ST in

Cape Town but was absent in Johannesburg. ST530b was the most common ST in

Johannesburg. STs found in Johannesburg but not Cape Town included ST062a, ST086, and

ST560.

Despite demographic differences between Cape Town and Johannesburg, a minimum

spanning tree analysis revealed no clear clustering of genotypes by location (Figure 1B).

Based on sequences from *ompA*, CT genovars from Cape Town and Johannesburg largely

overlapped, with genovar E and D being common in both regions (Figure 1C). However,

certain notable differences were evident, with genovar F only being found in Cape Town and

genovar G being more common in Johannesburg than Cape Town. Mapping STs detected in

the current study to the global MLST-based phylogeny (Figure 1D), revealed that there was

no clustering by site and that the overall diversity of STs was high.

In Cape Town, where adolescents were followed longitudinally, 33.9% who were

infected at visit 1 were also infected with CT at visit 2, and 18.2% were infected at visit 3

(Figure 2A). Among these, 10 individuals tested positive for CT at all three study visits, 10

were infected at the first two visits, and 5 were infected at visits 1 and 3 only. For those

infected at >1 study visit, the majority harboured the same ST across visits, suggesting

persistent infections or reinfection from the same source (Figure 2B). Limited sample size,

particularly after stratification by visit and ST, precludes formal inference about whether

prior infection increased the likelihood of infection at follow-up.

Discussion

This study highlights the high prevalence, persistence, and genotypic diversity of CT infections among AGYW in South Africa, a group disproportionately affected by STIs and their associated adverse reproductive health outcomes. The findings emphasize the urgent need for targeted interventions to address South Africa's significant chlamydial burden, particularly its implications for HIV acquisition and complications like pelvic inflammatory disease and infertility. CT prevalence in this cohort was 29.5%, with significant geographic variation: 41.6% in Cape Town compared to 17.4% in Johannesburg. The higher prevalence in Cape Town likely reflects differences in recruitment settings (reproductive health clinics versus general populations) and contraceptive use patterns. However, a limitation of this study is that participants were recruited using a targeted enrolment strategy, and a formal response rate could therefore not be calculated, which may limit the generalisability of the findings. All CT infections were asymptomatic, and younger AGYW (aged 16–19 years) were disproportionately affected, underscoring the need for early sexual health education and routine screening. Co-infections were common, with 71.6% of CT-positive participants also diagnosed with another STI or BV, which may exacerbate genital inflammation and increase HIV susceptibility (13).

Genotypic analysis using MLST revealed significant genetic diversity with 34 STs identified, including 15 novel STs. Regional differences were observed, with ST12d predominant in Cape Town and ST530b in Johannesburg. The lack of geographical clustering indicates widespread circulation of certain genotypes. Notably, whereas globally prevalent *ompA* genovars such as D and E (1) were dominant in both the South African cities studied, another globally dominant genotype, genovar F, was entirely absent in Johannesburg. Genovar G, a less globally common but potentially more pathogenic genotype (14), was

detected at a higher frequency in Johannesburg. Given that the infectiousness of CT is

significantly influenced by the *ompA* genotype of the infecting strain (15), these findings

underscore the importance of regional surveillance to identify locally relevant

epidemiological patterns with respect to informing tailored interventions (5). Future studies

should investigate the functional implications of these genotypic variations to better

understand their role in disease transmission and clinical outcomes.

Persistent or recurrent CT infections were common in the Cape Town cohort, where

participants were followed longitudinally. Among those with repeated infections, the same

ST was frequently identified across visits, indicating either reinfection from untreated

partners or incomplete clearance of infections. This calls attention to the importance of

partner notification, treatment, and test of cure to ensure resolution (2) in that persistent

infections likely contribute to sustained inflammation, which may exacerbate reproductive

health risks, including HIV susceptibility. However, limited sample sizes within STs

precluded assessment of type-specific CT persistence or recurrence.

From a public health perspective, addressing the high burden of CT among young

women in South Africa requires a multifaceted approach. Although there has been a shift in

policy in some countries (like the Netherlands) around the need for ongoing STI screening

and treatment, these data from South Africa strongly argue that expanding access to effective

and affordable STI POC testing in regions with high HIV burden, integrating partner

notification systems, and promoting consistent condom usage are needed for controlling CT

infections.

These findings stress the need for vaccines targeting CT, particularly those tailored to

the most prevalent genotypes in high-burden regions (8). Collar and colleagues (1) present

compelling evidence emphasizing the necessity for CT vaccine strategies to account for

regional variations in CT genotypes. This approach is critical to ensuring that interventions

are not only effective but also equitable, addressing the unique epidemiological and genetic

diversity of CT across diverse communities. Genomic surveillance is set to play a pivotal role

in addressing CT burden (5). Incorporating genetic diversity data into vaccine development

pipelines should also ensure broader protection against locally prevalent rather than just

globally dominant CT genotypes (5).

In conclusion, this study provides an analysis of CT prevalence, persistence, and

genotypic diversity among AGYW in South Africa. The findings emphasize the need for

targeted interventions to mitigate the health risks associated with CT. By integrating genomic

surveillance, public health initiatives, and community-based strategies, we can address the

multifaceted burden of CT and improve reproductive health outcomes for AGYW in South

Africa.

Acknowledgements

We extend our gratitude to the WISH study team, with special thanks to Ms. Penelope

Ngcobo for her efforts in participant recruitment and Sr. Janine Nixon. We are deeply

appreciative of the young women who participated in the study, without whom this research

would not have been possible. The data underlying this article are available in the PubMLST

Chlamydia trachomatis database at https://pubmlst.org/organisms/chlamydia-trachomatis, and

can be accessed using the assigned multilocus ST identifiers, including novel STs reported in

this study.

Funding sources

This work was supported by the EDCTP Strategic Primer grant (SP.2011.41304.038; PI: J.

Passmore, UCT) and a self-initiated grant from the South African Medical Research Council

(PI: J. Passmore, UCT).

Conflict of Interest

All authors declare no conflicts of interest related to this work.

References

1.   Delany-Moretlwe S, Mgodi N, Bekker LG, Baeten JM, Li C, Donnell D, et al. High

prevalence and incidence of gonorrhoea and chlamydia in young women eligible for HIV

pre-exposure prophylaxis in South Africa and Zimbabwe: results from the HPTN 082 trial.

Sex Transm Infect. 2023;99(7):433-9. Epub 20230308. doi: 10.1136/sextrans-2022-055696.

PubMed PMID: 36889914; PubMed Central PMCID: PMC10555488.

2.   Fortas C, Delarocque-Astagneau E, Randremanana RV, Crucitti T, Huynh BT.

Asymptomatic infections with Chlamydia trachomatis, Neisseria gonorrhoeae, and

Trichomonas vaginalis among women in low- and middle-income countries: A systematic

review and meta-analysis. PLOS Glob Public Health. 2024;4(5):e0003226. Epub 20240523.

doi: 10.1371/journal.pgph.0003226. PubMed PMID: 38781286; PubMed Central PMCID:

PMC11115196.

3.   Mkhize P, Mehou-Loko C, Maphumulo N, Radzey N, Abrahams AG, Sibeko S, et al.

Differences in HIV risk factors between South African adolescents and adult women and

their association with sexually transmitted infections. Sex Transm Infect. 2024. Epub

20241204. doi: 10.1136/sextrans-2024-056260. PubMed PMID: 39632075.

4.   Barnabas SL, Dabee S, Passmore JS, Jaspan HB, Lewis DA, Jaumdally SZ, et al.

Converging epidemics of sexually transmitted infections and bacterial vaginosis in southern

African female adolescents at risk of HIV. Int J STD AIDS. 2018;29(6):531-9. Epub

20171204. doi: 10.1177/0956462417740487. PubMed PMID: 29198180.

5.   Collar AL, Clarke TN, Jamus AN, Frietze KM. Ensuring equity with pre-clinical

planning for chlamydia vaccines. NPJ Vaccines. 2023;8(1):131. Epub 20230906. doi:

10.1038/s41541-023-00726-7. PubMed PMID: 37673890; PubMed Central PMCID:

PMC10482967.

6.   Herrmann B, Isaksson J, Ryberg M, Tangrot J, Saleh I, Versteeg B, et al. Global

Multilocus Sequence Type Analysis of Chlamydia trachomatis Strains from 16 Countries. J

Clin Microbiol. 2015;53(7):2172-9. Epub 20150429. doi: 10.1128/JCM.00249-15. PubMed

PMID: 25926497; PubMed Central PMCID: PMC4473235.

7.   Bom RJ, Christerson L, Schim van der Loeff MF, Coutinho RA, Herrmann B,

Bruisten SM. Evaluation of high-resolution typing methods for Chlamydia trachomatis in

samples from heterosexual couples. J Clin Microbiol. 2011;49(8):2844-53. Epub 20110608.

doi: 10.1128/JCM.00128-11. PubMed PMID: 21653758; PubMed Central PMCID:

PMC3147734.

8.   Klint M, Fuxelius HH, Goldkuhl RR, Skarin H, Rutemark C, Andersson SGE,

Persson K, Herrmann B. High-resolution genotyping of Chlamydia trachomatis strains by

multilocus   sequence   analysis.   J   Clin   Microbiol.   2007;45:1410–1414.

doi:10.1128/JCM.02301-06.

9.   Jolley KA, Bray JE, Maiden MCJ. Open-access bacterial population genomics:

BIGSdb software, the PubMLST.org website and their applications. Wellcome Open Res.

2018;3:124. doi:10.12688/wellcomeopenres.14826.1.

10.   Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:

improvements   in   performance   and   usability.   Mol   Biol   Evol.   2013;30:772–780.

doi:10.1093/molbev/mst010.

11.   Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler

A, Lanfear R. IQ-TREE 2: new models and efficient methods for phylogenetic inference in

the genomic era. Mol Biol Evol. 2020;37:1530–1534. doi:10.1093/molbev/msaa015.

12.   Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS.

ModelFinder: fast model selection for accurate phylogenetic estimates. Nat Methods.

2017;14:587–589. doi:10.1038/nmeth.4285.

13.   Masson L, Passmore JA, Liebenberg LJ, Werner L, Baxter C, Arnold KB, et al.

Genital inflammation and the risk of HIV acquisition in women. Clin Infect Dis.

2015;61(2):260-9. Epub 20150421. doi: 10.1093/cid/civ298. PubMed PMID: 25900168;

PubMed Central PMCID: PMC4565995.

14.   Anttila T, Saikku P, Koskela P, Bloigu A, Dillner J, Ikaheimo I, et al. Serotypes of

Chlamydia trachomatis and risk for development of cervical squamous cell carcinoma.

JAMA. 2001;285(1):47-51. doi: 10.1001/jama.285.1.47. PubMed PMID: 11150108.

15.   Ardizzone CM, Albritton HL, Lillis RA, Bagnetto CEL, Shen L, Cavacini LA, et al.

Human genital antibody-mediated inhibition of Chlamydia trachomatis infection and

evidence for ompA genotype-specific neutralization. PLoS One. 2021;16(10):e0258759.

Epub 20211018. doi: 10.1371/journal.pone.0258759. PubMed PMID: 34662351; PubMed

Central PMCID: PMC8523062.

Figure 1. MLST sequence types (ST) and *ompA* genovars in CT isolates from AGYW from

Cape Town and Johannesburg. (A) Percentage distribution of MLST STs by region. Each

stacked bar represents the proportion of detected STs in Johannesburg and Cape Town.

*indicate STs that were unique to each region. (B) Minimum-spanning tree analysis of the

identified MLST STs, showing the genetic relationships between STs across regions. The

size of the nodes represents the frequency of the sequence type, while connecting lines

indicate isolates differing at a single locus. Colours correspond to geographic locations (Cape

Town shaded in green and Johannesburg in red). (C) Distribution of *ompA* genovars,

stratified by geographic location. (D) Maximum likelihood phylogenetic tree generated from

a concatenated gene alignment for all currently recognised CT MLST STs. Coloured circles

represent STs detected in the current study.

Figure 2. Transition in CT infection status over time in AGYW from Cape Town. (A)

Longitudinal changes in CT infection status across three study visits. Participants were

categorized based on their infection status at each visit (CT negative [green]; incident [blue],

cleared [visit 2: lilac or visit 3: purple], persisted [dark peach] or recurred [light orange]. (B)

Comparison of MLST STs identified at multiple visits in AGYW with recurrent infections.

The data illustrate whether recurrent infections were associated with the same ST (suggesting

persistence or reinfection by untreated partners) or different STs (indicating new infections).

A

Transition in *C. trachomatis* status over time

B

V1	V2	V3
169b	135b*	
559		559
62a	62a	
62a	62a	
	530b	530b
268a	268a	268a
3	3	3
551	551	551
530b	530b	530b
530b	530b	
553	530c*	554a*
528b	528b	188b
188b	188b	188b
12d	12d	
12d	12d	12d
12d		12d
188b		188b
12d		12d
319		137
85	85	85
12d	12d	
3	3	
555	555	
87	87	
556	556	
12d	3	
3	3	
100b	100b	
56a	56a	
530b	530b	