

1 **Factors associated with unintended pregnancies among unmarried adolescents and the**
2 **potential of using mobile money shops: A Cross-sectional study in Eastern Uganda**

3 **Authors:**

- 4 1. Makiko Komasa, Ogata Sadako Research Institute for Peace and Development, Japan
5 International Cooperation Agency, Tokyo, Japan and Department of Global Health
6 Research, Juntendo University, Tokyo, Japan. mkomasa@juntendo.ac.jp.
- 7 2. Miho Sato, School of Tropical Medicine and Global Health, Nagasaki University, Nagasaki,
8 Japan. mihos@nagasaki-u.ac.jp.
- 9 3. Kiyoko Saito, Ogata Sadako Research Institute for Peace and Development, Japan
10 International Cooperation Agency, Tokyo, Japan. Saito.Kiyoko@jica.go.jp.
- 11 4. Sumihisa Honda, Graduate School of Biomedical Sciences, Nagasaki University, Nagasaki,
12 Japan. honda@nagasaki-u.ac.jp.
- 13 5. Robert Ssekitoleko, School of Biomedical Sciences, Makerere University, Kampala,
14 Uganda. rsseki@gmail.com.
- 15 6. Peter Waiswa, School of Public Health, Makerere University, Kampala, Uganda.
16 pwaiswa@musph.ac.ug.
- 17 7. Kharim Mwebaza Muluya, Iganga Health District Office, Iganga, Uganda.
18 muluyak@gmail.com.
- 19 8. Sheba Gitta, Busoga Health Forum, Jinja, Uganda, Uganda. shebagitta@gmail.com.
- 20 9. Myo Nyein Aung, Department of Global Health Research, Juntendo University, Tokyo,
21 Japan. myo@juntendo.ac.jp.

22

23 **Institutional address:**

24 Ogata Sadako Research Institute for Peace and Development, Japan International Cooperation
25 Agency (JICA), 10-5, Ichigaya-honmuracho, Shinjuku-ku, 162-8433, Tokyo, Japan, TEL: +81-
26 3-3269-2911, FAX: +81-3-3269-2054

27 **Corresponding author:**

28 Makiko Komasa, Ogata Sadako Research Institute for Peace and Development, Japan
29 International Cooperation Agency (JICA), 10-5, Ichigaya-honmuracho, Shinjuku-ku, 162-8433,

NOTE: This preprint reports new research that has not been certified by peer review and should not be used to guide clinical practice.

30 Tokyo, Japan, mkomasa@juntendo.ac.jp. 0000-0002-0864-0657.

31

32 **Abstract**

33 **Background:** Adolescent sexual and reproductive health and rights (SRHR) remains a critical
34 public health issue in low- and middle-income countries, with adverse health, educational, and
35 economic consequences. Uganda faces significant challenges with high rates of unintended
36 pregnancies among adolescents. This study investigated the factors associated with unintended
37 pregnancies among unmarried adolescents in Eastern Uganda and explored the potential of
38 using mobile money vendors to provide SRHR services.

39 **Methods:** A cross-sectional study was conducted among 1267 unmarried girls/boys, aged 15-
40 19 years, in the Busoga region. Data were collected through face-to-face interviews, using a
41 structured questionnaire. Sociodemographic characteristics, sexual behaviors, contraceptive
42 knowledge, and environmental factors were analyzed. Multivariate logistic regression analysis
43 identified factors associated with unintended pregnancies.

44 **Results:** Of the adolescents who had sexual intercourse (n=876), 22.5% (n=197) experienced
45 unintended pregnancies. Fundamental factors associated with lower prevalence of pregnancy
46 included being a current student (adjusted odds ratio [AOR]: 0.36, 95% confidence interval
47 [CI]: 0.24-0.54), living with parents (AOR: 0.48, 95% CI: 0.33-0.69), and teacher engagement
48 (AOR: 0.56, 95% CI: 0.37-0.84). Factors associated with higher prevalence of pregnancy

49 included knowledge regarding contraceptive use (AOR: 2.28, 95% CI: 1.16-4.49), parental
50 communication (AOR: 1.91, 95% CI: 1.32-2.75), parental contraception support (AOR: 1.64,
51 95% CI: 1.03-2.59), and mobile phone possession (AOR: 2.31, 95% CI: 1.09-4.90). Public
52 health facilities and community-based distributors, including mobile money vendors, were
53 considered comfortable channels for obtaining contraceptives for adolescents who had
54 experienced pregnancy.

55 **Conclusions:** Unintended pregnancies among unmarried adolescents in Eastern Uganda were
56 influenced by educational, parental, and environmental factors. Enhancing SRHR education
57 from earlier ages, parental involvement, and leveraging community resources such as mobile
58 money vendors could improve adolescent SRHR outcomes. Innovative approaches outside
59 conventional health and education sectors are necessary for sustainable and effective adolescent
60 SRHR programs.

61

62 **Trial registration:** This study was registered at Japan's University Hospital Medical
63 Information Network (UMIN000053332) on 12 January 2024.

64

65 **Keywords:** sexual and reproductive health and rights, unintended pregnancy, unmarried
66 adolescent, mobile money vendor, sexual behavior, teenagers, Uganda.

67

68 **Plain English summary**

69 Adolescent sexual and reproductive health and rights (SRHR) is a pressing public health issue
70 globally. Uganda has a high rate of unintended pregnancies among adolescents. This study
71 investigated the factors associated with unintended pregnancies among unmarried adolescents
72 in Eastern Uganda and explored the potential of using mobile money vendors to provide SRHR
73 services.

74 This cross-sectional study was conducted using a structured questionnaire at 60 vendors. It
75 included 1267 unmarried girls/boys aged 15-19 years. Of 876 adolescents who ever had sexual
76 intercourse, 22.5% experienced unintended pregnancies. Factors associated with a lower
77 prevalence of pregnancy were: a current student (adjusted odds ratio [AOR] = 0.36), living with
78 parents (AOR = 0.48), and teacher engagement (AOR = 0.56). Contrarily, factors associated
79 with a higher prevalence of pregnancy were: knowledge regarding contraceptive use (AOR =
80 2.28), parental communication (AOR = 1.91), parental contraception support (AOR = 1.64), and
81 mobile phone possession (AOR = 2.30), which may imply that teenagers who had experienced
82 pregnancy tend to become more serious about contraception and engagement of parents and
83 teachers enhanced. In addition, community-based SRHR service distributors, including mobile
84 money vendors, were considered comfortable channels.

85 In conclusion, unintended pregnancies among unmarried adolescents in Eastern Uganda
86 were influenced by educational, parental, and environmental factors. Enhancing SRHR
87 education before starting sexually active and parental and school involvement may reduce
88 unintended pregnancies among unmarried adolescents in Eastern Uganda. In addition,
89 leveraging accessible community-based distributors, such as mobile money vendors, may be a
90 potential channel for delivering SRHR information and contraceptives.

91

92 **Background**

93 Adolescents sexual and reproductive health and rights (SRHR) is a pressing public health issue
94 globally. In low- and middle-income countries, approximately 21 million girls aged 15-19 years
95 become pregnant and almost half of these pregnancies are unintended [1]. These early
96 pregnancies not only contribute to maternal and child mortality but also lead to unsafe abortions
97 [1, 2]. The impact of early pregnancy extends beyond health, affecting a girl's education and
98 income-earning potential and perpetuating a cycle of poverty that can last for generations [3,
99 4]. This situation is a result of unmet needs for SRHR services among young populations [5, 6].

100 Sub-Saharan Africa (SSA) has high rates of adolescent pregnancies, and this trend is
101 expected to increase [7]. Among the 29 SSA countries, Uganda ranks the fifth with a high
102 prevalence of unintended pregnancies [8]. Many girls engage in sexual activity before marriage.
103 The median age for the first sexual relationship among women aged 25-49 years was 16.9 years
104 in 2021 [9]. The maternal mortality rate remains high at 189 per 100,000 live births, with 17%
105 of these involving girls aged 15-19 years [2, 9]. The rate of any contraceptive use among all
106 women aged 15-49 year remains at 44.0% and for modern methods at 40.1% [9]. Ninety-seven
107 percent of pregnant female students drop out of school, despite the education policy
108 encouraging students who have given birth to return to school [10]. Additionally, there has been
109 an increase in the number of unintended pregnancies resulting from "transactional sex," where

110 girls economically affected by coronavirus disease 2019 crisis engage in sex in exchange for
111 small amounts of financial aid or transportation [11]. There remains a need to improve the
112 quality and quantity of public SRHR services in Uganda, including youth-friendly family
113 planning services [12, 13]. The Ugandan government has implemented various policies and
114 programs over the past 15 years, such as expanding the compulsory education period, enacting
115 defilement laws, prohibiting marriage for girls under 18 years of age, prohibiting sexual
116 intercourse with girls under 18 years of age, and promoting adolescent contraception [11, 14].
117 However, adolescent pregnancy rates have remained stagnant over the past decade [14].

118 Since the International Conference on Population and Development in 1994, numerous
119 interventions have been undertaken to improve adolescent SRHR [15]. Most of these
120 interventions have been conducted in schools, primary healthcare facilities, and communities
121 by peer-educators [16-18]. However, some studies highlighted that many programs have not
122 adequately reached unmarried teenagers who need SRHR services [16, 19]. Additionally, the
123 number of SRHR services has been limited, and their adoption has been insufficient, making
124 many of these programs unsustainable and ineffective despite substantial investments [19].
125 Consequently, youth SRHR programs have stagnated for almost two decades [6, 20]. In light
126 of these challenges, several studies have stressed the need for innovative approaches to ensure
127 sustainability and scalability [17, 18, 21, 22].

128 In our exploration of effective approaches in Uganda, we focused on utilizing mobile

129 money shops (vendors) as a channel to connect with adolescents and provide them with SRHR
130 services [23]. These vendors are available throughout the country; as in 2023, 63% of Uganda's
131 population was using mobile phones [24]. The number of registered mobile phone customers
132 increased by 1.3 times from 2015 to 2019 [25]. Our preliminary study also discovered that 85%
133 of adolescents accessed vendors more than weekly to use various services such as money
134 transfers, purchasing airtime/data, and making cash deposits [23].

135 Given the situation in Uganda, we initiated a hybrid effectiveness-implementation trial to
136 reach teenagers using mobile money shops to reduce unintended pregnancies [23]. Based on
137 baseline data collected as part of the trial, this study aimed to explore the current situation of
138 sexual behavior and determine factors associated with unintended pregnancy among
139 adolescents aged 15-19 years. More importantly, this study assessed the potential of an
140 innovative approach based on vendors in Uganda.

141

142 **Methods**

143 **Study design and setting**

144 The detailed study design has been published elsewhere [23] and was registered at J アパ根セ
145 ‘the University Hospital Medical Information Network (UMIN000053332) in Japan. Here,
146 we provide the fundamental parts of the study design for interpreting the study results. This was
147 a cross-sectional study of unmarried adolescents aged 15-19 years in the Busoga region of

148 Eastern Uganda. The center of the Busoga region is approximately 80 km from Kampala, the
149 capital city. Two municipalities in the Busoga region with similar characteristics were selected:
150 Iganga and Bugiri. In 2022, the population of Iganga municipality was 55,263 and that of Bugiri
151 was 28,747, whose teenage population (10-17 years old) accounted for 20.9% and 19.3%,
152 respectively . We selected 30 vendors from each municipality according to predefined study
153 criteria. Owners of mobile money businesses (vendors) aged 18-30 years, who have operated
154 for more than one year, and are willing to participate in the study were selected. Stand-alone
155 shops were selected to secure the privacy of customers and commodities. The geographical
156 distribution, diversity, population density, availabilities of health facilities and schools of each
157 municipality were also considered. We recruited teenage customers from selected vendor sites.
158 Recruitment ceased after reaching the desired sample size of approximately 20 per vendor.

159

160 **Participants**

161 The eligible study participants were unmarried girls/boys aged 15-19 years who used the
162 selected vendors in the target municipalities and had no plans to move out of the municipalities
163 during the study period. Adolescents aged 13-14 years were excluded because they tended to
164 be less sexually active and used fewer vendors.

165 We applied a non-randomized prospective controlled design because vendor customers
166 cannot be assigned to specific vendors, and they choose several vendors at their own

167 convenience. We decided on a sample size of 1200 for the intervention study in the two
168 municipalities using a formula based on the available data and previous studies, using a
169 calculation method explained elsewhere [23]. Adolescents under 18 years of age who agreed to
170 contact their parents/guardians during the screening process were recruited.

171

172 **Data collection**

173 A structured questionnaire was developed using research instruments based mainly on the 2016
174 Uganda Demographic and Health Survey [26] and other studies conducted in SSA [27-33]. We
175 have added questions regarding social and physical opportunities, and those related to vendors'
176 usage [23]. We developed a questionnaire in English and translated it into the local language.
177 Before finalizing the questionnaire, we conducted a series of pre-tests with 26 adolescents (10
178 boys and 16 girls) and modified them for easy answering. Although most adolescents used
179 English daily, some participants were comfortable using their local language to discuss sensitive
180 topics in SRHR. We employed data collectors who could manage the local language and had
181 academic backgrounds in the social welfare or health fields. The questionnaire included five
182 categories: (1) sociodemographic characteristics, (2) sexual behavior, (3) contraception
183 knowledge and capability, (4) environmental factors preventing pregnancy, and (5) preference
184 for contraceptive practices.

185 The questionnaire was administered face-to-face by trained data collectors using

186 preprogrammed tablet computers [23]. Interviews were conducted in a private location near
187 vendors' or teenagers' homes between February and March 2024. Data collection lasted
188 between 30 and 40 min per interview.

189

190 **Data analysis**

191 The primary outcome was the experience of pregnancy experience of girls or impregnation of
192 boys, regardless of sex. First, we described the sociodemographic characteristics of the
193 participants and their sexual behaviors with the comparison of the two groups
194 (pregnancy/impregnation [pregnant] versus never-pregnant) using the chi-square test. Second,
195 we described the variables in each category using two-group comparisons. Third, multivariate
196 logistic regression analysis was performed to identify the factors associated with the primary
197 outcome. Variables that showed a statistically significant difference in bilateral regression were
198 considered for multivariate logistic regression. In addition, we removed correlated variables
199 and added the basic variables identified in a previous study, which accounted for 21 variables.
200 Statistical significance was set at $p < 0.05$. Data analysis was conducted using STATA v.18.

201

202 **Ethical consideration**

203 Ethical approval was obtained from the Ethics Review Committee of Uganda Christian
204 University (UCUREC-2023-710), the Uganda National Council for Science and Technology

205 (SS2284ES), and the Ethics Review Committee of the JICA Ogata Sadako Research Institute
206 for Peace and Development in Japan (JICA Ogata RI, reference: JICA [DI] 202312220002).
207 Written informed consent was obtained from all adolescent participants and the
208 parents/guardians of adolescents aged 15-17 years. Adolescents may feel uncomfortable
209 participating in the study as the content includes their sexual behaviors. To mitigate their anxiety,
210 we informed them that their participation was truly voluntary without any harm and that they
211 could withdraw their participation at any time, even after the survey and collected information
212 were strictly confidentially managed. To obtain consent from parents/guardians of adolescents
213 under 18 years of age, highly experienced and skilled researchers visited the homes of expected
214 participants and explained the study to the parents/guardians.

215

216 **Results**

217 We recruited 1281 eligible teenagers for the intervention [Additional file_01]. We excluded 14
218 participants who did not responded their pregnancy experiences, and 1267 participants were
219 included in the study. Table 1 shows the sociodemographic characteristics of the participants in
220 two groups: pregnant versus never-pregnant. Out of the analyzed participants, 69.1% had sexual
221 intercourse (876/1267). Among the respondents who had sexual intercourse, 22.5% (197/876)
222 had gotten pregnant. Of the respondents, boys were 52.5% and those aged below 18 years were
223 30.8%. More than half of them (57.7%) completed lower secondary school (senior 4). The

224 highest proportion of religion was Islam (54.3%). We recruited both in- and out-of-school
 225 adolescents; 51.2% were currently in-school students and 39.8% were working. Only sex and
 226 religion were not statistically different between the two groups. Table 2 presents respondents'
 227 sexual behaviors. Overall, 84.7% (1073/1267) ever had a girlfriend(s)/boyfriend(s) (i.e.,
 228 intimate partner[s]). Among them, 81.6% (876/1073) had ever experienced sexual intercourse.
 229 The mean age at first intercourse was 15.39 years (standard deviation [SD]: 1.98) and was
 230 significantly different between the pregnant (mean: 15.04 years, SD: 2.01) and never-pregnant
 231 (mean: 15.48 years, SD: 1.96) groups (difference: 0.44 years, $p < 0.0053$). In the last four
 232 months, 88.7% (777/876) had intimate partners, and 84.7% (660/777) had sexual intercourse.
 233 Out of these, 81.5% (538/660) had taken contraceptives during intercourse in the last four
 234 months. Out of those who got pregnant, 51.0% (97/190, 7 did not responded about outcome of
 235 pregnancy, out of 197) had a live birth(s).

236 **Table 1 Basic characteristics of participants(n = 1267)**

Variables	Total (n=1267)		Pregnant (n=197)		Never pregnant (n=1070)		p-value
	n	%	n	%	n	%	
1) Gender							0.321
Boys	665	52.5	97	49.2	568	53.1	
Girls	602	47.5	100	50.8	502	46.9	
2) Age							$p < 0.001$
15-17 years old	390	30.8	24	12.2	366	34.2	
18-19	887	69.2	173	87.8	704	65.8	
3) Education level completed							0.017
Primary school or less	429	33.9	83	42.1	346	32.3	
Lower secondary school	731	57.7	103	52.3	628	58.7	
Secondary school or higher	107	8.5	11	5.6	96	9.0	
4) Religion							0.204
Christianity	327	25.8	54	27.4	273	25.5	
Islam	688	54.3	113	57.4	575	53.7	

Others	252	19.9	30	15.2	222	20.8	
5) Currently in education							p
Yes	649	51.2	45	22.8	604	56.5	p < 0.001
No	618	48.8	152	77.2	466	43.6	
6) Currently in working							p < 0.001
Yes	504	39.8	114	57.9	390	36.5	
No	763	60.2	83	42.1	680	63.6	

237

238 **Table 2 Sexual behavior**

Variables	Total		Pregnant		Never pregnant		p-value
	n	%	n	%	n	%	
1) Ever had girl/boyfriends (n=1267)							
Yes	1073	84.7	197	100.0	876	81.9	p < 0.001
No	194	15.3	0	0.0	194	18.1	
2) Ever had sexual intercourse (n=1073)							
Yes	876	81.6	197	100.0	679	77.5	p < 0.001
No	190	17.7	0	0.0	190	21.9	
Declined to answer	7	0.7	0	0.0	7	0.8	
3) Age at the first sexual intercourse (n=866)							
Age (SD)	15.39	(1.98)	15.04	(2.01)	15.48	(1.96)	0.005
4) Having girl/boy friends in the last four months (n=876)							
Yes	777	88.7	180	91.4	597	87.9	0.394
No	89	10.2	15	7.6	74	10.9	
Declined to answer	10	1.1	2	1.0	8	1.2	
5) Sexual intercourse in the last four months (n = 777)							
Yes	660	84.9	163	90.6	497	83.3	0.051
No	107	13.8	15	8.3	92	15.4	
Declined to answer	10	1.3	2	1.1	8	1.3	
6) Use of contraceptives in the last four months (n = 660)							
Yes	538	81.5	127	77.9	411	82.7	0.384
No	116	17.6	34	20.9	82	16.5	
Declined to answer	6	0.9	2	1.2	4	0.8	

239

240 Table 3 shows factors related to contraception knowledge and capability. A total of 96.1%
 241 had heard any contraceptives, while the knowledge regarding how to use any contraceptives
 242 was lower (79.9%). Knowledge regarding the menstrual cycle and timing of pregnancy was low
 243 (overall 9.1%). Friends were the most frequent source of contraceptive information, followed

244 by school, television/radio, and public health facilities. Partner agreement on contraceptive use
245 was 80.8%, and differences between the two groups. Knowledge regarding how to use
246 contraceptives and sources of contraceptive information (public health facility, private facility,
247 school, community-based distributor, and parent/relatives) were significantly different between
248 the pregnant and never-pregnant groups. Table 4 describes the environmental factors preventing
249 pregnancy. Nearly 70% of the respondents lived with parents, showing a higher percentage of
250 never-pregnant than the pregnant group ($p < 0.001$). Communication with parents/guardians
251 about intimate partners was low among all respondents (27.6%). Perception of support by
252 healthcare workers, teachers, and the community was relatively high among all respondents
253 (82.7%, 68.4%, and 69.1%, respectively), with statistically significant differences between the
254 two groups, except for community support. Regarding mobile phone possession, 81.2% of
255 teenagers had it, with the pregnant group having significantly higher ownership than the never-
256 pregnant group ($p < 0.001$). Regarding vendor service usage, 64.2% used vendor services at
257 least once a day. The most frequent use of service was airtime/data for all, while the pregnant
258 group showed a higher percentage of withdraws than the never-pregnant group. Table 5 shows
259 the results of preferences on contraception practice. Desire to acquire knowledge on
260 contraception and develop intimate relationships were high in both the pregnant and never-
261 pregnant groups (92.9% and 83.2%, respectively). The most preferred source of SRHR
262 information was the public health facility, and others were varied. Regarding comfortable

263 channels of obtaining contraceptives, public health facilities were the most preferred source),
 264 followed by pharmacies) and private health facilities. Statistically significant differences
 265 between the two groups were: desire to develop intimate relationship, preferred source of SRHR
 266 information, some of channel of obtaining contraceptives. Nearly 70% expected vendors to be
 267 suitable channels for receiving SRHR information and contraceptives before the trial
 268 intervention started.

269 **Table 3 Knowledge and capability of contraception (n = 1267)**

Variables	Total (n=1267)		Pregnant (n=197)		Never pregnant (n=1070)		p-value
	n	%	n	%	n	%	
1) Knowledge on contraceptives							0.146
Yes	1218	96.1	193	98.0	1,025	95.8	
No	49	3.9	4	2.0	45	4.2	
2) Knowledge on how to use contraceptives							p < 0.001
Yes	1012	79.9	184	93.4	828	77.4	
No	255	20.1	13	6.6	242	22.6	
3) Knowledge on the fertility period							0.567
Yes	115	9.1	20	10.2	95	8.9	
No	1,152	90.9	177	89.9	975	91.1	
4) Source of contraceptive information							p < 0.001
1. Public health facility							
Yes	640	50.5	141	71.6	499	46.6	
No	627	49.5	56	28.4	571	53.4	
2. Private facility							0.005
Yes	198	15.6	44	22.3	154	14.4	
No	1069	84.4	153	77.7	916	85.6	
3. School							0.007
Yes	790	62.4	106	53.8	684	63.9	
No	477	37.7	91	46.2	386	36.1	
4. Community-based distributor							p < 0.001
Yes	240	18.9	56	28.4	184	17.2	
No	1027	81.1	141	71.6	886	82.8	
5. TV/radio							0.511
Yes	732	57.8	118	59.9	614	57.4	
No	535	42.2	79	40.1	456	42.6	
6. SNS							

Yes	357	28.2	64	32.5	293	27.4	0.143
No	910	71.8	133	67.5	777	72.6	
7. Parents/relatives							
Yes	480	37.9	88	44.7	392	36.6	0.033
No	787	62.1	109	55.3	678	63.4	
8. Friends							
Yes	866	68.4	133	67.5	733	68.5	0.783
No	401	31.7	64	32.5	337	31.5	
5) Agreement on contraceptive use by partner (n=876)							
Yes	708	80.8	162	82.2	546	80.4	0.585
No	80	9.1	19	9.6	61	9.0	
I don't know	88	10.1	16	8.1	72	10.6	

270

271 **Table 4 Environment related to prevent pregnancy (n = 1267)**

Variables	Total (n=1267)		Pregnant (n=197)		Never pregnant (n=1070)		p-value
	n	%	n	%	n	%	
1) Living with parent (s)							
Yes	828	65.4	98	50.3	730	68.2	p < 0.001
No	439	34.7	99	49.8	340	31.8	
2) Communicating with parents/guardians on intimate partner							
Yes	349	27.6	96	48.7	253	23.6	p < 0.001
No	918	72.4	101	51.3	817	76.4	
3) Supporting contraceptive use by parents/guardians							
Yes	618	48.8	135	68.5	483	45.1	p < 0.001
No	451	35.6	41	20.8	410	38.3	
I don't know	198	15.6	21	10.7	177	16.5	
4) Supporting contraceptive use by healthcare workers							
Yes	1048	82.7	178	90.4	870	81.3	0.008
No	100	7.9	9	4.6	91	8.5	
I don't know	119	9.4	10	5.1	109	10.2	
5) Teachers' awareness of student's life/worries							
Yes	866	68.4	110	55.8	756	70.7	p < 0.001
No	302	23.8	64	32.5	238	22.2	
I don't know	99	7.8	23	11.7	76	7.1	
6) Community supports on teenagers							
Yes	875	69.1	142	72.1	733	68.5	0.523
No	332	26.2	48	24.4	284	26.5	
I don't know	60	4.7	7	3.6	53	5.0	
7) Mobile phone possession							
Yes	1029	81.2	186	94.4	843	78.8	p < 0.001
No	238	18.8	11	5.6	227	21.2	
8) Frequency of access to vendors							

Once a day or more	813	64.2	125	63.5	688	64.3	0.770
Weekly	340	26.8	53	26.9	287	26.8	
Less than weekly	93	7.3	14	7.1	79	7.4	
Others	21	1.7	11	5.0	2.54	16.0	
9) Most frequent use of vendors							
Withdraws	194	15.3	42	21.3	152	14.2	0.081
Deposit	401	31.7	58	29.4	343	32.1	
Airtime/data	661	52.2	96	48.7	565	52.8	
Others	11	0.9	1	0.5	10	0.9	

272

273 **Table 5 Preferences for contraceptive practice (n = 1267)**

Variables	Total (n=1267)		Pregnant (n=197)		Never pregnant (n=1070)		p-value
	n	%	n	%	n	%	
1) Desire for knowledge of contraception							
Yes	1177	92.9	189	95.9	988	92.3	0.192
No	53	4.2	5	2.5	48	4.5	
I don't know	37	2.9	3	1.5	34	3.2	
2) Desire for developing intimate relationship							
Yes	1054	83.2	184	93.4	870	81.3	p < 0.001
No	105	8.3	5	2.5	100	9.4	
I don't know	108	8.5	8	4.1	100	9.4	
3) Preferred source of SRHR information							
Public health facility	505	39.9	106	53.8	399	37.3	p < 0.001
Others	672	53.0	83	42.1	589	55.1	
I don't want to know	90	7.1	8	4.1	82	7.7	
4) Comfortable channel to obtain contraceptives							
1. Public health facility							
Yes	952	75.1	160	81.2	792	74.0	0.032
No	315	24.9	37	18.8	278	26.0	
2. Private facility							
Yes	366	28.9	58	29.4	308	28.8	0.852
No	901	71.1	139	70.6	762	71.2	
3. Pharmacy/drug shop							
Yes	445	35.1	61	31.0	384	35.9	0.183
No	822	64.9	136	69.0	686	64.1	
4. Community-based distributor							
Yes	53	4.2	23	11.7	30	2.8	p < 0.001
No	1214	95.8	174	88.3	1,040	97.2	
5) Agree on vendor as suitable channel for SRHR services (n=1264)							
Yes	880	69.6	151	76.7	729	68.3	0.013
No	295	23.3	30	15.2	265	24.8	
Don't to know	89	7.0	16	8.1	73	6.8	

274

275

276 Table 6 presents factors associated with experiencing pregnancy/impregnation. Girls'
277 pregnancy rate was higher than the boys' impregnation rate (adjusted odds ratio [AOR]: 2.29,
278 95% confidence interval [CI]: 1.52-3.44). Factors associated with the lower prevalence of
279 pregnancy were currently student status (AOR: 0.36, 95% CI: 0.24-0.54), living with parents
280 (AOR: 0.48, 95% CI: 0.33-0.69), and teachers' engagement (AOR: 0.56, 95% CI: 0.37-0.84).
281 Contrarily, factors associated with the higher pregnancy were knowledge regarding how to use
282 contraceptives (AOR: 2.28, 95% CI: 1.16-4.49), communication with parents/guardians (AOR:
283 1.91, 95% CI: 1.32-2.75), supporting contraceptive use by parents (AOR: 1.64, 95% CI: 1.03-
284 2.59), and mobile phone possession (AOR: 2.31, 95% CI: 1.09-4.90). In addition, adolescents
285 who had experienced pregnancy were more likely to use public health facilities (AOR: 1.66,
286 95% CI: 1.10-2.49) and community-based distributors (AOR: 1.72, 95% CI: 1.11-2.66).
287 Healthcare workers and community support were not correlated with the pregnancy experiences.

288 **Table 6 Factors associated with experiencing pregnancy (n = 1267)**

289 (Insert attached file)

290

291 **Discussion**

292 This study illustrates the current sexual behaviors of adolescents who used the services of
293 vendors in Eastern Uganda. Nearly 70% of the analyzed participants had sexual intercourse.
294 The approximate age at sexual debut was approximately 15 years for both sexes—more than

295 20% of those who had sexual experience had ever been pregnant. Parental engagement was a
296 key factor associated with pregnancy. Teachers' engagement with students' lives reflected a
297 reduction in pregnancy risks. In contrast, mobile phone possession and desire to get
298 contraceptives from community-based distributors were positively correlated with pregnancy
299 experiences.

300

301 **Active sexual behavior**

302 Adolescents in SSA are sexually active [34, 35]. The Ugandan national average of percentage
303 of those who had sexual intercourse among unmarried population aged 15-49 years were 59.6%
304 for male and 45.6% of female in 2021 [9]. Our respondents showed more active sexual
305 behaviors than the national trend, even limiting the younger population. The Ugandan national
306 average age at first intercourse among adults aged 20-49 years was 18.3 years for males and
307 16.6 years for females [9]. Compared with the national average, our study participants had an
308 early sexual initiation (15.4 years for both sexes; 5.0 years for males; 15.9 years for females).
309 These trends were much lower than those in other SSA countries, for example, 17.0 years for
310 females in Nigeria [27]. Furthermore, the pregnant group experienced the first sexual
311 intercourse 0.44 years earlier than the other group in this study. This was in line with previous
312 studies that revealed that earlier first intercourse was associated with higher unintended
313 pregnancy rates in SSA [14, 34-36]. Our study found that the experience of pregnancy was 0.36

314 times lower among school students than among non-students, implying that pregnancy led to
315 drop-out from school or out-school teens are more likely to face pregnancy risks. This supports
316 that longer school years may reduce pregnancy risks, and extension of compulsory education
317 years by the governmental strategy may partially effect to its purpose [14]. In addition, the
318 Uganda government had issues several laws to reduce unintended girls pregnancy risk since
319 2000, such as the Unholy Law and the prohibition of marriage for girls under the age of 18
320 years [14]. However the data are limited to prove the effectiveness of these governmental
321 compulsory measures [14]. Therefore, further research is needed to capture explicit adolescent
322 needs to lower sexual activities nationwide and strengthen effective, reality-based policies [12,
323 14].

324

325 **Contraceptive knowledge and practice**

326 Non-practice of contraception is a critical determinant of unintended adolescent pregnancies in
327 SSA [8, 36]. Most respondents in this study had a basic knowledge of contraceptive methods,
328 which is similar to the national trend among sexually active unmarried both-sex teenagers
329 (97.8% for boys and 99.4% for girls aged 15-19 years) [26]. However, the participants' practical
330 knowledge on how to use the methods was 16% points lower than their basic knowledge. In
331 addition, only 9.1% of the adolescents had correct knowledge of the fertile period of the
332 menstrual cycle, which was lower than the national average among girls aged 15-19 years

333 (13.8%) in 2016 [26]. Our descriptive analysis revealed that adolescents with pregnancy
334 experience tended to have more knowledge and access to contraception and support from
335 parents/guardians and healthcare workers. Although this cross-sectional study could not
336 identify causality, the results may be interpreted that teens those who experienced pregnancy
337 became more serious about searching for information and mobilizing services after pregnancy.
338 It can be said that Uganda's SRHR education either at school, community, nor home does not
339 provide preventative information of pregnancy and meet their real needs, although the Ministry
340 of Education and Sports encourages the improvement of SRHR education in schools. In our
341 study, the most common source of contraceptive information was schools; however, this was
342 not the preferred source of SRHR information. A previous study concluded that insisting on
343 abstinence in school education was ineffective in preventing teenage pregnancies [37].
344 Providing practical knowledge on sex, pregnancy, contraception, and sexually transmitted
345 infections is crucial among adolescents at the scale, through home, school, and community [37-
346 39]. Some studies emphasized that younger people were exposed to more unsafe sexual
347 intercourse and were flexible in absorbing concepts related to gender, suggesting that sex
348 education from an early age before sexual initiation was effective [36, 40-42]. A study on
349 children aged 10-14 years in Uganda supported this suggestion [43]. Education content should
350 include sex, pregnancy, delivery, sexually transmitted infections, other health risks, the efficacy
351 of vulnerable populations, and future income potential [3, 7, 11, 32, 37, 38, 44]. The content

352 should be designed according to the needs of different adolescent age groups [41, 45]. About
353 half of those who had ever been pregnant were most likely to terminate their pregnancies. Since
354 abortion for non-medical reasons is illegal in Uganda, statistics on abortion data were collected
355 only through post-abortion care at health facilities and the actual situation is underreported. The
356 previous qualitative studies captured that many girls have used traditional medicines or unsafe
357 means to terminate unwanted pregnancies and received no after-abortion care [11, 12, 14].
358 Ignoring high-risk, unsafe self-abortion or necessary after-abortion care is likely to lead to
359 maternal death and adverse maternal outcomes [12]. Sexual education emphasizing the risks of
360 abortion and infertility is needed [11, 37].

361

362 **Parental involvement**

363 Parental involvement was an essential factor influencing adolescent unintended pregnancy in
364 several studies in SSA countries [31, 32, 38]. In this study, "living with parents" was indicated
365 to be a deterrent factor to sexual behavior. Studies in Ghana have reported a similar situation in
366 which parental involvement reduced pregnancy occurrence [31, 38]. Conversely, parental
367 interaction and parental contraceptive support were positively correlated with pregnancy
368 experience, which may imply that parents' substantive care and support increased after their
369 children's pregnancy experience, similar to the results of a Ghanaian study [31]. Parents' neglect
370 of their children and hesitancy to teach sex education because they did not know how to interact

371 with them was also noted in a previous study in Uganda [11]. Parents' motivation for sex
372 education for their children, their own acquisition of correct SRHR knowledge, and their ability
373 to interact with the children need to be improved [27, 32, 38]. Urgent development of
374 educational program guidance for families (parents/guardians) and communities, including
375 practical sex education content and skills to teach children in Uganda [11] is required.

376

377 **Contraceptive accessibility in the community**

378 The prevalence of unintended pregnancies is closely related to access to contraceptives [44].
379 Paid services at private clinics or pharmacies were hurdles for teenagers in Uganda and they
380 tend to avoid public health facilities because they follow formal procedures, ask detailed
381 questions about personal information and the purpose of contraceptive use, and, in some cases,
382 may be known to their parents [11, 12]. This study found that motivation for contraceptive use
383 was higher among those who had experienced pregnancy, and in fact, they mobilized a variety
384 of contraceptive sources. These facts imply that teenagers who experienced pregnancies have
385 strengthened their contraceptive capabilities. A Ghanaian study also pointed out that many girls
386 who had experienced pregnancy developed competencies to deal with pregnancy prevention
387 [31]. Our findings also indicated that easily accessible community-based distributors showed a
388 preference for and a higher desire for future use among teenagers who had experienced
389 pregnancy.

390 This study also found Approximately 70% of adolescents thought that the vendor was a
391 suitable channel for receiving SRHR services. Furthermore, adolescents who have experienced
392 pregnancy are more likely to prefer accessing vendor-based SRHR service. These facts suggest
393 that vendors may be effective channels for providing SRHR services, especially for high risk
394 population. Chandra-Mouli pointed out that interventions implemented in many programs such
395 as facility-based interventions, including school education, health center activities, and youth
396 centers with peer-educators in the community, have not been cost-effective or sustainable [19].
397 The Ugandan government has called for innovative approaches that utilize local mechanisms
398 outside the health and education sectors [46, 47]. In response to the global community calling
399 for innovative platforms to reach vulnerable populations nearly a decade ago [4, 17, 21], our
400 results suggest that vendors in Uganda could be a cost-effective platform to reach adolescents
401 and strengthen their competencies in accessing contraceptives and practice.

402

403 **Limitation**

404 This study has several limitations. First, we collected data from adolescents who used the
405 vendor services, meaning that our findings could not represent the general population. Second,
406 since the survey was dependent on self-reporting, the study is susceptible to recall and social
407 desirability biases. Third, the nature of the quantitative study and cross-sectional design could
408 not describe causality of pregnancy, and limited further interpretation of underlying social,

409 cultural, and historical factors. Fourth, our study did not explore the economic status of the
410 adolescents because of difficulties in accessing accurate data and comparing adolescents living
411 with their parents and others using a structured questionnaire. In addition, we did not include
412 patients with human immunodeficiency virus (HIV) acquired immunodeficiency syndrome
413 (AIDS) in this study. As there is still high stigma attached to HIV/AIDS status in Uganda, we
414 determined that obtaining such information through self-reports would not provide reliable
415 information. To overcome these limitations, a qualitative study with a diverse population of
416 children, parents, and social decision-makers is required. Despite these limitations, to the best
417 of our knowledge, this is the first community-based quantitative study of sexual behaviors and
418 contraception preferences among unmarried girls and boys aged 15-19 years in Uganda and
419 identified the unmet needs of sexual and reproductive health services .

420

421 **Conclusions**

422 This study revealed a picture of early sexual debut and pregnancy risk among adolescents aged
423 15-19 years in Eastern Uganda. This study found that the important factors underlying such
424 situations were a lack of knowledge of pregnancy risk and correct knowledge of contraception
425 as well as a lack of practical sexual education by parents and schools. However, there was light
426 on the existence of a contemporary community resource with a high affinity for adolescents,
427 known as mobile money vendors. Young vendors who are in the community, geographically

428 accessible, and have a close relationship with adolescents could be a potential channel for
429 delivering necessary information on pregnancy risk and promoting safe contraceptive use
430 among adolescents. Further studies are needed to develop a model for promoting modern
431 contraceptive use among vendors, test its scalability, and share the results with policymakers in
432 Uganda.

433

434 **List of abbreviations**

435 AIDS, acquired immunodeficiency syndrome

436 AOR, adjusted odds ratio

437 CI, confidence interval

438 HIV, human immunodeficiency virus

439 SRHR, sexual and reproductive health and rights

440 SSA, Sub-Saharan Africa

441

442 **Declarations**

443 **Ethics approval and consent to participate:** Ethical approval was obtained from the Ethics
444 Review Committee of Uganda Christian University (UCUREC-2023-710), Uganda National
445 Council for Science and Technology (SS2284ES), and Ethics Review Committee of the JICA
446 Ogata Sadako Research Institute for Peace and Development in Japan (JICA Ogata RI,

447 reference: JICA [DI] 202312220002). Written informed consent was obtained from all
448 adolescent participants and the parents/guardians of adolescents aged 15-17 years.

449

450 **Consent for publication:**

451 Not applicable.

452

453 **Availability of data and materials:**

454 The dataset supporting the conclusions of this article is available in the repository of the
455 University Hospital Medical Information Network (UMIN000053332) dataset in
456 <https://rctportal.niph.go.jp/en>.

457

458 **Competing interests:**

459 The authors declare that they have no competing interests.

460

461 **Funding:**

462 This work was supported by JSPS KAKANHI grant number JP 23K16346 from Grants-in-Aid
463 for Scientific Research from the Ministry of Education, Culture, Sports, Science and
464 Technology of Japan.

465

466 **Authors' contributions:**

467 MK, MS, RS, KS, SG and AMN were involved in the development of the conception and study
468 design. MK, MS, and RS were responsible for field research implementation. SH provided

469 technical advice on the overall methodology, including the sampling. MK was involved in
470 drafting the manuscript. PW supervised the study and was involved in critical revision of the
471 manuscript. MS, RS, MKM and AMN provided technical comments on the manuscript draft.
472 MK was responsible for the fund. All authors were involved in the final approval of the
473 manuscript and the decision to submit the manuscript for publication.

474

475 **Acknowledgements:**

476 We would like to express our gratitude to all study participants, the Ministry of Health for their
477 invaluable support. Our heartfelt thanks also go to all members of the research team of the
478 Busoga Health Forum, whose efforts facilitated the implementation of this study, including the
479 development of a network involving various stakeholders and healthcare facilities.

480

481 **Authors' information:**

482 ¹Ogata Sadako Research Institute for Peace and Development, Japan International Cooperation
483 Agency, Tokyo, Japan. ²Department of Global Health Research, Juntendo University, Tokyo,
484 Japan. ³School of Tropical Medicine and Global Health, Nagasaki University, Nagasaki, Japan.
485 ⁴Graduate School of Biomedical Sciences, Nagasaki University, Nagasaki, Japan. ⁵School of
486 Biomedical Sciences, Makerere University, Kampala, Uganda. ⁶School of Public Health,
487 Makerere University, Kampala, Uganda. ⁷Iganga Health District office, Iganga, Uganda.
488 ⁸Busoga Health Forum, Jinja, Uganda.

489

490 References

- 491 1. WHO. Adolescent pregnancy. 2023. [https://www.who.int/news-room/fact-](https://www.who.int/news-room/fact-sheets/detail/adolescent-pregnancy)
492 [sheets/detail/adolescent-pregnancy](https://www.who.int/news-room/fact-sheets/detail/adolescent-pregnancy). Accessed 8 Jul 2024.
- 493 2. UNFPA. State of world population 2022_ Seeing the unseen 2022.
- 494 3. Mann L BD, Black KI. Teenage Pregnancy. *Australian Journal of General Practice*
495 2020, 49, 310-316.
- 496 4. Baird S, Ezeh A, Azzopardi P, Choonara S, Kleinert S, Sawyer S, Patton G, Viner R.
497 Realising transformative change in adolescent health and wellbeing: a second Lancet
498 Commission. *Lancet* 2022, 400(10352):545-547.
- 499 5. Vaivada T, Lassi ZS, Irfan O, Salam RA, Das JK, Oh C, Carducci B, Jain RP, Als D,
500 Sharma N *et al*. What can work and how? An overview of evidence-based interventions
501 and delivery strategies to support health and human development from before
502 conception to 20 years. *The Lancet* 2022, 399(10337):1810-1829.
- 503 6. Kruk ME, Lewis TP, Arsenault C, Bhutta ZA, Irimu G, Jeong J, Lassi ZS, Sawyer SM,
504 Vaivada T, Waiswa P *et al*. Improving health and social systems for all children in
505 LMICs: structural innovations to deliver high-quality services. *The Lancet* 2022,
506 399(10337):1830-1844.
- 507 7. Ahinkorah BO, Kang M, Perry L, Brooks F, Hayen A. Prevalence of first adolescent
508 pregnancy and its associated factors in sub-Saharan Africa: A multi-country analysis.
509 *PLoS One* 2021, 16(2):e0246308.
- 510 8. Ameyaw EK, Budu E, Sambah F, Baatiema L, Appiah F, Seidu AA, Ahinkorah BO.
511 Prevalence and determinants of unintended pregnancy in sub-Saharan Africa: A multi-
512 country analysis of demographic and health surveys. *PLoS One* 2019, 14(8):e0220970.
- 513 9. Uganda Bureau of Statistics. Demographic and Health Survey 2022. 2022.
- 514 10. Ministry of Education and Sports. National sexuality education framework 2018. 2018.
- 515 11. Nabugoomu J, Seruwagi GK, Hanning R. What can be done to reduce the prevalence of
516 teen pregnancy in rural Eastern Uganda?: multi-stakeholder perceptions. *Reprod Health*
517 2020, 17(1):134.
- 518 12. Atuyambe LM, Kibira SP, Bukenya J, Muhumuza C, Apolot RR, Mulogo E.
519 Understanding sexual and reproductive health needs of adolescents: evidence from a
520 formative evaluation in Wakiso district, Uganda. *Reprod Health* 2015, 12:35.
- 521 13. Mambo SB, Sikakulya FK, Ssebuufu R, Mulumba Y, Wasswa H, Mbina SA, Rusatira
522 JC, Bhondoekhan F, Kamyuka LK, Akib SO *et al*. Challenges in Access and Utilization
523 of Sexual and Reproductive Health Services Among Youth During the COVID-19
524 Pandemic Lockdown in Uganda: An Online Cross-Sectional Survey. *Frontiers in*
525 *Reproductive Health* 2022, 3.
- 526 14. Amongin D, Benova L, Nakimuli A, Nakafeero M, Kaharuzza F, Atuyambe L, Hanson
527 C. Trends and determinants of adolescent childbirth in Uganda- analysis of rural and

- 528 urban women using six demographic and health surveys, 1988-2016. *Reprod Health*
529 2020, 17(1):74.
- 530 15. Starrs AM, Ezeh AC, Barker G, Basu A, Bertrand JT, Blum R, Coll-Seck AM, Grover
531 A, Laski L, Roa M *et al.* Accelerate progress-sexual and reproductive health and rights
532 for all: report of the Guttmacher-Lancet Commission. *Lancet* 2018, 391(10140):2642-
533 2692.
- 534 16. Denno DM, Chandra-Mouli V, Osman M. Reaching youth with out-of-facility HIV and
535 reproductive health services: a systematic review. *J Adolesc Health* 2012, 51(2):106-
536 121.
- 537 17. Denno DM, Hoopes AJ, Chandra-Mouli V. Effective strategies to provide adolescent
538 sexual and reproductive health services and to increase demand and community support.
539 *J Adolesc Health* 2015, 56(1 Suppl):S22-41.
- 540 18. Belaid L, Dumont A, Chaillet N, Zertal A, De Brouwere V, Hounton S, Ridde V.
541 Effectiveness of demand generation interventions on use of modern contraceptives in
542 low- and middle-income countries. *Trop Med Int Health* 2016, 21(10):1240-1254.
- 543 19. Chandra-Mouli V, Lane C, Wong S. What Does Not Work in Adolescent Sexual and
544 Reproductive Health: A Review of Evidence on Interventions Commonly Accepted as
545 Best Practices. *Glob Health Sci Pract* 2015, 3(3):333-340.
- 546 20. Bhutta ZA, Boerma T, Black MM, Victora CG, Kruk ME, Black RE. Optimising child
547 and adolescent health and development in the post-pandemic world. *The Lancet* 2022,
548 399(10337):1759-1761.
- 549 21. Salam RA, Das JK, Lassi ZS, Bhutta ZA. Adolescent Health Interventions: Conclusions,
550 Evidence Gaps, and Research Priorities. *J Adolesc Health* 2016, 59(4S):S88-S92.
- 551 22. Igras S, Plesons M, Chandra-Mouli V. Building evidence on what works (and what does
552 not): practical guidance from the World Health Organization on post-project evaluation
553 of adolescent sexual and reproductive health projects. *Health Policy Plan* 2021,
554 36(5):594-605.
- 555 23. Komasawa M, Sato M, Ssekitoleko R, Waiswa P, Gitta S, Nabugoomu J, Honda S, Saito
556 K, Aung MN. Study protocol for a type-II hybrid effectiveness-implementation trial to
557 reach teenagers using mobile money shops to reduce unintended pregnancies in Uganda.
558 *BMJ Open* 2024, 14(4):e084539.
- 559 24. Uganda Bureau of StatisticsStatistics. 2020 Statistical Abstract. 2020.
- 560 25. Kepios: Digital 2023 uganda. 2023. [https://datareportal.com/reports/digital-2023-](https://datareportal.com/reports/digital-2023-uganda)
561 [uganda](https://datareportal.com/reports/digital-2023-uganda). Accessed 8 Jul 2024.
- 562 26. Uganda Bureau of StatisticsStatistics. Demographic and Health Survey 2016. 2018.
- 563 27. Crawford EE, Atchison CJ, Ajayi YP, Doyle AM. Modern contraceptive use among
564 unmarried girls aged 15-19 years in South Western Nigeria: results from a cross-
565 sectional baseline survey for the Adolescent 360 (A360) impact evaluation. *Reprod*
566 *Health* 2021, 18(1):6.

- 567 28. Orach CG, Otim G, Aporomon JF, Amone R, Okello SA, Odongkara B, Komakech H.
568 Perceptions, attitude and use of family planning services in post conflict Gulu district,
569 northern Uganda. *Confl Health* 2015, 9:24.
- 570 29. Okawa S, Mwanza-Kabaghe S, Mwiya M, Kikuchi K, Jimba M, Kankasa C, Ishikawa
571 N. Sexual and reproductive health behavior and unmet needs among a sample of
572 adolescents living with HIV in Zambia: a cross-sectional study. *Reprod Health* 2018,
573 15(1):55.
- 574 30. Bakesiima R, Beyeza-Kashesya J, Tumwine JK, Chalo RN, Gemzell-Danielsson K,
575 Cleve A, Larsson EC. Effect of peer counselling on acceptance of modern
576 contraceptives among female refugee adolescents in northern Uganda: A randomised
577 controlled trial. *PLoS One* 2021, 16(9):e0256479.
- 578 31. Ahorlu CK, Pfeiffer C, Obrist B. Socio-cultural and economic factors influencing
579 adolescents' resilience against the threat of teenage pregnancy: a cross-sectional survey
580 in Accra, Ghana. *Reprod Health* 2015, 12:117.
- 581 32. Yakubu I, Salisu WJ. Determinants of adolescent pregnancy in sub-Saharan Africa: a
582 systematic review. *Reprod Health* 2018, 15(1):15.
- 583 33. Zard M, Lau LS, Bowser DM, Fouad FM, Lucumi DI, Samari G, Harker A, Shepard
584 DS, Zeng W, Moresky RT *et al.* Leave no one behind: ensuring access to COVID-19
585 vaccines for refugee and displaced populations. *Nat Med* 2021, 27(5):747-749.
- 586 34. Sedgh G, Finer LB, Bankole A, Eilers MA, Singh S. Adolescent pregnancy, birth, and
587 abortion rates across countries: levels and recent trends. *J Adolesc Health* 2015,
588 56(2):223-230.
- 589 35. Ferede TA, Muluneh AG, Wagnaw A, Walle AD. Prevalence and associated factors of
590 early sexual initiation among youth female in sub-Saharan Africa: a multilevel analysis
591 of recent demographic and health surveys. *BMC Womens Health* 2023, 23(1):147.
- 592 36. Ahinkorah BO. Individual and contextual factors associated with mistimed and
593 unwanted pregnancies among adolescent girls and young women in selected high
594 fertility countries in sub-Saharan Africa: A multilevel mixed effects analysis. *PLoS One*
595 2020, 15(10):e0241050.
- 596 37. Stanger-Hall KF, Hall DW. Abstinence-only education and teen pregnancy rates: why
597 we need comprehensive sex education in the U.S. *PLoS One* 2011, 6(10):e24658.
- 598 38. Krugu JK, Mevissen FE, Prinsen A, Ruiters RA. Who's that girl? A qualitative analysis
599 of adolescent girls' views on factors associated with teenage pregnancies in Bolgatanga,
600 Ghana. *Reprod Health* 2016, 13:39.
- 601 39. Zulaika G, Nyothach E, van Eijk AM, Obor D, Mason L, Wang D, Chen T, Kerubo E,
602 Opollo V, Ngere I *et al.* Factors associated with the prevalence of HIV, HSV-2,
603 pregnancy, and reported sexual activity among adolescent girls in rural western Kenya:
604 A cross-sectional analysis of baseline data in a cluster randomized controlled trial. *PLoS*
605 *Med* 2021, 18(9):e1003756.

- 606 40. Cohn LD, Macfarlane S, Yanez C, Imai WK. Risk-perception: Differences between
607 adolescents and adults. *Health Psychol* 1995, 14(3):217-222.
- 608 41. Goldfarb ES, Lieberman LD. Three Decades of Research: The Case for Comprehensive
609 Sex Education. *J Adolesc Health* 2021, 68(1):13-27.
- 610 42. Yah CS, Ndlovu S, Kutuywayo A, Naidoo N, Mahuma T, Mullick S. The prevalence of
611 pregnancy among adolescent girls and young women across the Southern African
612 development community economic hub: A systematic review and meta-analysis. *Health*
613 *Promot Perspect* 2020, 10(4):325-337.
- 614 43. Kemigisha E, Bruce K, Ivanova O, Leye E, Coene G, Ruzaaza GN, Ninsiima AB,
615 Mlahagwa W, Nyakato VN, Michielsen K. Evaluation of a school based comprehensive
616 sexuality education program among very young adolescents in rural Uganda. *BMC*
617 *Public Health* 2019, 19(1):1393.
- 618 44. WHO. Every newborn: An action plan to end preventable deaths. 2014. www.who.int.
619 Accessed 8 Jul 2024.
- 620 45. Osadolor UE, Amoo EO, Azuh DE, Mfonido-Abasi I, Washington CP, Ugbenu O.
621 Exposure to Sex Education and Its Effects on Adolescent Sexual Behavior in Nigeria. *J*
622 *Environ Public Health* 2022, 2022:3962011.
- 623 46. Ministr of Health. The second national family planning costed implementation plan
624 2020/21- 2024/25 (FP-CIP II). 2021.
- 625 47. Ministry of Health. National guidelines for conducting family planning outreaches in
626 Uganda. 2017.
- 627