Generalized Linear Mixed Model Approach for Analyzing Water, Sanitation, and Hygiene Facilities in Bangladesh: Insights from BDHS 2022 Data

Mahmila Sanjana Mim[1*], Md. Lutfor Rahaman[2¶] and Anamul Haque Sajib[2¶]

[1]Department of Statistics, University of Dhaka, Dhaka—1000, Bangladesh

[2]Department of Statistics, University of Dhaka, Dhaka—1000, Bangladesh

*Corresponding Author's Email: mahmilasanjana@gmail.com

Authors details:

Mahmila Sanjana Mim, Email: mahmilasanjana@gmail.com; ORCID ID: 0009-0002-4419-1542

Md. Lutfor Rahaman, Email: lutfor3021@du.ac.bd.com

Anamul Haque Sajib, Email: sajibstat@du.ac.bd; ORCID ID: 0009-0000-0583-4045

Abstract

This study aims to identify key determinants and barriers to effective WASH facilities in Bangladesh, crucial for preventing health issues and ensuring fair access. For identifying the vital factors associated with access to WASH facilities, data obtained from Bangladesh Demographic and Health Survey (BDHS) 2022 have been analyzed. Since the survey employed a two-stage stratified clustered sampling approach, the responses from each cluster are likely to be correlated and exhibit variation between clusters, reflecting the inherent heterogeneity. To overcome this problem and incorporate cluster effect along with fixed effects, Generalized Linear Mixed Model (GLMM) for binary response has been employed rather than the conventional fixed effect logistic regression model for analyzing individual and combined basic WASH facilities in Bangladesh. The superiority of GLMM over fixed effect logistic regression model for binary response have been observed based on Akaike Information Criterion (AIC) and likelihood ratio test (LRT). The study shows that, age of household head, partner's education level, media exposure, women empowerment, wealth index, mother's age and mother's education of children under 5 had significant association with basic handwashing facility. Basic sanitation facility was associated with division, place of residence, sex and age of the household head, household size, partner's education level, working status, wealth index, mother's age, and mother's education of children under 5. Wealth index significantly affected the access to basic drinking water facility. Moreover, combined WASH facilities were influenced by the sex and age of the household head, household size, partner's education level, working status, media exposure, wealth index, mother's age, and mother's education of children under 5. Findings of this study can be used to enhance accessibility to basic WASH facilities, contributing to the achievement of 6th Sustainable Development Goal (SDG).

Keywords: Water, Sanitation, and Hygiene (WASH), Generalized Linear Mixed Model (GLMM), intra-cluster correlation, Sustainable Development Goal (SDG).

Introduction

The United Nations adopted the Sustainable Development Goals (SDGs) in 2015 to urge everyone to work together to eradicate poverty, safeguard the environment, and achieve peace and prosperity for all by 2030. There are 17 SDGs and they are interlinked. They acknowledge that the impact of one goal depends on the progress of others, and that development requires a harmonious approach among social, economic and environmental dimensions. Among 17 SDGs the 6th goal is related to the clean water and sanitation which aims to ensure the availability and sustainable management of water and sanitation for all [1]. Evidence demonstrates that water pollution and poor sanitation can cause many transmitted diseases and conditions such as bacillary diarrhoea, cholera, typhoid, polio, viral hepatitis A, acute respiratory infections etc. [2-8]. The World Health Organization (WHO) says that almost 2 million people die every year around the world as they do not have enough WASH facilities, and most of them are children [8]. Although, an impressive progress on improving access to safe water, sanitation and hygiene (WASH) has been witnessed globally, the year 2020 saw 3.85 billion people without basic hygiene service, 1.7 billion people without basic water service, and 780 million people without improved toilets at their health care facility [9]. This indicates that many people's lives are affected by unmet needs of WASH facilities. Hence, lack of adequate access to WASH services can exert widespread influence on one's health, food availability, income and education.

Like many other developing countries, Bangladesh is also facing problems due to lack of proper WASH facilities. According to the latest data from the Joint Monitoring Programme (JMP) of

UNICEF and WHO, only 59% of the population in Bangladesh had safely managed drinking water in 2020, and only 39% had safely managed sanitation. Moreover, basic hygiene facilities had been found only among 58% of the population [10]. Poor WASH facilities in Bangladesh lead to various negative consequences. Therefore, it is necessary to improve the WASH facilities. The government should take proper steps to ensure the availability of basic WASH services. For this purpose, the factors which are hindering the availability of basic WASH facilities should be identified properly.

Considering the harmful effects of inadequate WASH facilities, this study aims to identify the factors contributing to insufficient WASH services in Bangladesh. For doing this, the Bangladesh Demographic and Health Survey (BDHS), 2022 data have been utilized. The BDHS delivers nationally representative data on numerous of population health and demographics in Bangladesh. Since BDHS employed a two-stage stratified cluster sampling approach, data from respondents within the same cluster are likely correlated. To ensure consistent and efficient parameter estimates, this correlation must be accounted for during estimation. A generalized linear mixed model (GLMM) can be utilized to model such correlated data [11].

To the best of our knowledge, no existing literature has identified the key determinants and barriers to effective individual and combined basic WASH facilities in Bangladesh using the latest BDHS 2022 data. Motivated by this lack, our study has aimed to bridge the existing research gap.

Data and methodology

Data sources

In this study, the secondary data extracted from the BDHS 2022 have been used which can be freely downloaded from the DHS website upon request. The BDHS 2022 was conducted under the authority of the National Institute of Population Research and Training (NIPORT), Ministry of

Health and Family Welfare (MOHFW), Medical Education and Family Welfare Division. Mitra and Associates which is a private research agency, implemented this survey from June 2022 to December 2022. The BDHS 2022 sample underwent a two-stage stratified selection. Initially, 675 enumeration areas (EAs) were chosen based on their size, with independent selection within each stratum. In the second stage, 45 households per cluster were systematically selected from the new household listings. The survey targeted all ever-married women aged 15-49 who were either usual residents or stayed the night before the survey in the selected households [12]. After excluding missing values, our final dataset included 15,610 observations.

Study variables

Dependent variables

The study has focused on household access to individual and combined basic WASH services. According to the WHO/UNICEF Joint Monitoring Programme (JMP) guidelines, these services were categorized into different levels: basic, limited, unimproved and no service (Table 1 and Table 2) [13].

Table 1. WHO/UNICEF JMP ladder for WASH services.

Service level	Water	Sanitation	Hygiene
Basic	Drinking water from an improved source, provided collection time is not more than 30 min for a round trip, including queuing	Use of improved facilities that are not shared with other households	Availability of a handwashing facility on premises with soap and water
Limited	Drinking water from an improved source for which collection time exceeds 30 min for a round trip, including queuing	Use of improved facilities shared between two or more households	Availability of a handwashing facility on premises without soap and water

Unimproved	Drinking water from an unprotected dug well or unprotected spring	Use of pit latrines without a slab or platform, hanging latrines or bucket latrines	Not applicable
No service	Surface water	Open defecation	No handwashing facility on premise

Table 2. JMP classification of improved/unimproved water and sanitation facility type

Facility types	Water	Sanitation
Improved facilities	Piped supplies • Tap water in the dwelling, yard or plot • Public standposts Non-piped supplies • Boreholes/tubewells • Protected wells and springs • Rainwater • Packaged water, including bottled water and sachet water • Delivered water, including tanker trucks and small carts	Network sanitation • Flush and pour flush toilets connected to sewers On-site sanitation • Flush and pour flush toilets or latrines connected to septic tanks or pit • Ventilated improved pit latrines • Pit latrines with slabs • Composting toilets, including twin pit latrines and container-based system
Unimproved facilities	Non-piped supplies • Unprotected wells and springs	On-site sanitation • Pit latrines without slabs • Hanging latrines • Bucket latrines

In order to create dependent variables (individual basic WASH facilities — basic handwashing facility, basic sanitation facility and basic drinking water facility) by following the guidelines of WHO/UNICEF Joint Monitoring Programme (JMP), a dichotomization has been performed:

$$\text{Individual Basic WASH Facilities} = \begin{cases} \text{Yes} = 1, & \text{when the service level is basic} \\ \text{No} = 0, & \text{otherwise} \end{cases}$$

While constructing the basic drinking water variable, the collection time for drinking water from an improved source has not been considered because of extensive missing data. Finally, the last binary dependent variable has been generated for households that combined all three individual basic WASH facilities, termed as combined basic WASH facilities. A household has been considered to have access to the combined basic WASH facilities only if all of these three basic facilities are available. That is,

$$\text{Combined Basic WASH Facilities} = \begin{cases} \text{Yes} = 1, & \text{all of three basic services are available} \\ \text{No} = 0, & \text{otherwise} \end{cases}$$

Covariates

Based on literature reviews [14-28], the following covariates have been considered in this study: division (Barisal, Chittagong, Dhaka, Khulna, Mymensingh, Rajshahi, Rangpur and Sylhet), place of residence (urban and rural), sex of household head (male and female), age of household head (< 30, $30 - 39$, $40 - 49$, $50 - 59$ and ≥ 60), household size (≤ 5 and > 5), partner's education level (no education, primary, secondary and higher), working status (yes and no), media exposure (exposed and non-exposed), migration (migrant and non-migrant), women empowerment (yes and no), wealth index (poor, middle and rich), children aged 5 and under in household (yes and no), mother's age of under 5 children (≤ 30 and > 30), mother's education of under 5 children (no education, primary, secondary and higher). Here, the variable media exposure has been defined by categorizing the respondents who read magazines or newspapers, listen to radio, or watch television once a week. The migration status has been determined based on whether individuals have resided in their current homes for less than two years. The variable women empowerment has been generated on the basis of active participation of women in four types of decision: about

their own health care, major household purchases, visiting family or relatives and decisive actions on money earned by their husbands. If a woman is involved any of these four decisions, she has been placed into category: yes, otherwise she has been placed into category: no. Though the dataset originally divides the variable wealth index into five categories: poorest, poorer, middle, richer and richest, for the purposes of this study, these have been consolidated into three groups: the poor category has been formed by combining the poorest and poorer groups, the middle category has remained unchanged and the rich category has been created by merging the richer and richest groups.

Statistical analyses

The study variables taken into consideration in this paper have been summarized using univariate analysis, such as frequency distribution. The Chi-Square test has been used as a bivariate analysis to measure the connection between the response variable and the categorical explanatory variables. Finding the explanatory variables that are significantly associated with the response variable is one of the primary goals of bivariate analysis. Significant explanatory variables identified in bivariate analysis have been regarded as such in multivariate analysis.

Generalized Linear Mixed Model (GLMM)

To take into account the correlation between the observations inside each cluster, the Generalized Linear Mixed Model (GLMM) has been used in this investigation. In contrast to the Generalized Linear Model (GLM), random effects have been incorporated alongside fixed effects to get estimates of the regression parameters that are more accurate. GLMMs extend GLMs to accommodate correlated responses within the same group, which contravenes the independence assumption. GLMM is regarded as a superior method for analyzing data that demonstrates clustering effect [29].

Let, y_{ip} be p^{th} individual from i^{th} cluster where $i = 1, 2,..., q$ and, $p = 1, 2,..., n_i$. Let, x_{ip} be a vector of covariates for p^{th} individual from i^{th} cluster related with fixed effect parameter denoted by β as variation may exist in number of subjects per cluster. Also, let, u be a $(q \times 1)$ vector of random effects associated with q clusters and z_{ip} be a unique vector of dimensions $(q \times 1)$, which is composed entirely of zeros except for a single entry of 1 at the i^{th} position where, $i = 1, 2,..., q$. One may write,

$$\mu_{ip} = E(y_{ip} \mid u_i),$$

where, u_i be the random effect of the i^{th} cluster. The linear predictor in GLMM has the following form [11],

$$g(\mu_{ip}) = x_{ip}'\beta + z_{ip}'u.$$

For continuous response variable,

$$y_{ip} = x_{ip}'\beta + z_{ip}'u + e_{ip},$$

which can be written in matrix notation as,

$$Y = X\beta + Zu + e,$$

here, Y denotes $n \times 1$ column vector, X denotes $n \times k$ matrix of covariates, β is $k \times 1$ vector of coefficients of fixed effects, Z is $n \times q$ design matrix of q random effects, u denotes $q \times 1$ vector of random effects associated with q clusters and the $n \times 1$ vector of residuals is denoted by e where, $n = \sum_{i=1}^{q} n_i$. u follows normal distribution with mean 0 and covariance matrix, Σ [16].

Because of introducing a random slope in the model, the numbers of columns in design matrix, Z will be double although the number of rows will remain same. Hence, the covariance matrix, Σ, will become a (2×2) matrix as follows,

$$\Sigma = \begin{pmatrix} \sigma^2_{intercept} & \sigma^2_{intercept,slope} \\ \sigma^2_{slope,intercept} & \sigma^2_{slope} \end{pmatrix}$$

In this study, only random intercept term is being considered which turns Σ into a scaler. In this case, the variance-covariance matrix of residuals is,

$$Var(e_{ip}) = \sigma^2_e.$$

In this structure, it is assumed that the variance of residual is homogeneous for all observations and also they are independent of each other. The between group (or cluster) variance is denoted by σ^2_u and the within group variance is denoted by σ^2_e. The likelihood function for the individuals associated with i^{th} cluster is,

$$L_i\big((\beta,\sigma^2_u)\,|\,x_{ip},u_i\big) = f(y_i\,|\,x_{ip},u_i) = \prod_{p=1}^{n_i} f(y_i\,|\,x_{ip},u_i)$$

The marginal likelihood function is [30],

$$L\big((\beta,\sigma^2_u)\,|\,x_{ip}\big) = \int_{-\infty}^{\infty} \left[\prod_{i=1}^{q} L_i\big((\beta,\sigma^2_u)\,|\,u_i\big)\right] g(u_i)du_i.$$

It is quite impossible to have explicit solution from this expression. For this reason, some techniques have been utilized for approximation in order to obtain the maximum likelihood estimates. Some of these approximation techniques are: Gauss-Hermite quadrature, Laplace approximation, Penalized quasi-likelihood method [31-33].

GLMM for Binary Response: Mixed Effect Logistic Regression Model

When the response variable is binary, employing a mixed effect logistic regression within the Generalized Linear Mixed Models (GLMM) framework is suitable for analyzing such data. The link function under this model is of the following form [34],

$$g(\mu_{ip}) = log\left(\frac{\mu_{ip}}{1 - \mu_{ip}}\right),$$

and,

$$log\left(\frac{\mu_{ip}}{1 - \mu_{ip}}\right) = x_{ip}{}'\beta + z_{ip}{}'u,$$

where,

$$u_i \sim N(0,\Sigma)$$

Because of considering only random intercept term in this study, Σ can be replaced by σ_u^2,

$$u_i \sim N\left(0,\sigma_u^2\right)$$

The conditional probability of p^{th} observation from i^{th} cluster given the value of covariate of that observation and the random cluster effect is given by,

$$\mu_{ip} = E\left[Y_{ip} \mid x_{ip}, u_i\right] = Pr\left[Y_{ip} = 1 \mid x_{ip}, u_i\right] = \frac{e^{\beta' x_{ip} + u_i}}{1 + e^{\beta' x_{ip} + u_i}}$$

Intra-Cluster Correlation (ICC)

Intra-cluster correlation is a statistical measure which is used to measure how similar the outcomes are that belong to the same cluster [35]. For binary response variable, the model can be written as,

$$Y_{ip} = \mu + u_i + e_{ip}; \qquad u_i = z_{ip}{}'u,$$

where, μ is fixed, u_i follows normal distribution with zero mean and variance σ^2, e_{ip} has standard logistic distribution with zero mean and variance σ_e^2. If the response variable is binary type, the within cluster variance σ_e^2 is replaced by $\frac{\pi^2}{3}$ for convenience [36]. Then the intra-cluster correlation (ICC) for binary outcome denoted by ρ can be defined as,

$$\rho = \frac{\sigma_u^2}{\sigma_u^2 + \frac{\pi^2}{3}}$$

Since, the response variables of our study (individual and combined basic WASH facilities) are binary, mixed effect logistic regression model within GLMM framework has been employed for analyzing data. Based on likelihood ratio test (LRT) and AIC values, we have compared the performance of the fixed effect logistic regression model and the mixed effect logistic regression model. AIC has the following mathematical form:

$$AIC = -2ln(L) + 2\gamma,$$

where, n is the number of observations and γ is the number of parameters to be estimated. The model with the lowest AIC is typically preferred since it offers a better fit. All the analyses of our study have been performed by using Statistical Package for Social Science (SPSS) software version 20 and R version 4.1.1.

Results

Univariate analysis

Table 3 presents the frequency, valid percentage, and cumulative percentage of each study variable. It shows that 59.7% of households had access to basic handwashing facility, 62.7% to

basic sanitation, 98.7% to basic drinking water, and 45.0% to combined WASH facilities. The highest number of respondents were from Dhaka (15.7%) and the lowest from Sylhet (10.2%). A larger proportion resided in rural areas (64.6%) compared to urban areas (35.4%). Most household heads were male (88.6%), with 28.9% aged 40-49 years and 13.5% aged under 30. The majority had a household size of 5 or fewer (74.7%). Among respondents' partners, 23.0% were illiterate, 31.0% had secondary education (highest), and 16.6% had higher education (lowest). Additionally, 33.2% of respondents were currently working, 58.4% had media exposure, and 90.6% were non-migrants. Women empowerment stood at 89.1%. Of the 15,610 respondents, 37.9% were from poor families, 19.8% from middle-class, and 42.3% from rich families. Nearly half (47.9%) had children aged 5 or under. Among these mothers, 40.0% were aged 30 or younger, and 60.0% were older than 30. Educational levels among these mothers were: 14.3% illiterate, 28.4% primary, 34.1% secondary, and 23.2% higher education.

Table 3. Univariate analysis of dependent and independent variables.

Variable	Frequency	Valid Percent	Cumulative Percent
Basic Handwashing Facility			
Yes	9324	59.7%	59.7%
No	6286	40.3%	100.0%
Basic Sanitation Facility			
Yes	9784	62.7%	62.7%
No	5826	37.3%	100.0%
Basic Drinking Water Facility			
Yes	15410	98.7%	98.7%
No	200	1.3%	100.0%
Combined WASH Facilities			
Yes	7018	45.0%	45.0%
No	8592	55.0%	100.0%
Division			
Barishal	1662	10.6%	10.6%
Chattogram	2239	14.3%	25.0%
Dhaka	2450	15.7%	40.7%
Khulna	2060	13.2%	53.9%
Mymensingh	1690	10.8%	64.7%
Rajshahi	2056	13.2%	77.9%
Rangpur	1860	11.9%	89.8%
Sylhet	1593	10.2%	100.0%
Place of Residence			

Urban	5528	35.4%	35.4%
Rural	10082	64.6%	100.0%
Sex of Household Head			
Male	13827	88.6%	88.6%
Female	1783	11.4%	100.0%
Age of Household Head			
<30	2115	13.5%	13.5%
30-39	3710	23.8%	37.3%
40-49	4506	28.9%	66.2%
50-59	3002	19.2%	85.4%
≥60	2277	14.6%	100.0%
Household Size			
≤5	11653	74.7%	74.7%
>5	3957	25.3%	100.0%
Partner's Education Level			
No Education	3594	23.0%	23.0%
Primary	4583	29.4%	52.4%
Secondary	4836	31.0%	83.4%
Higher	2597	16.6%	100.0%
Working Status			
Yes	5177	33.2%	33.2%
No	10433	66.8%	100.0%
Media Exposure			
Exposed	9112	58.4%	58.4%
Non-exposed	6498	41.6%	100.0%
Migration			
Migrant	1464	9.4%	9.4%
Non-migrant	14146	90.6%	100.0%
Woman Empowerment			
Yes	13912	89.1%	89.1%
No	1698	10.9%	100.0%
Wealth Index			
Poor	5912	37.9%	37.9%
Middle	3089	19.8%	57.7%
Rich	6609	42.3%	100.0%
Children Aged 5 and Under in Household			
Yes	7483	47.9%	47.9%
No	8127	52.1%	100.0%
Mother's Age of Under 5 Children			
≤30	6250	40.0%	40.0%
>30	9360	60.0%	100.0%
Mother's Education of Under 5 Children			
No Education	2233	14.3%	14.3%
Primary	4430	28.4%	42.7%
Secondary	5327	34.1%	76.8%
Higher	3620	23.2%	100.0%

Bivariate analysis

Bivariate analysis represents the most fundamental form of quantitative analysis used to uncover the empirical correlation between two distinct variables. Essentially, it seeks to ascertain the presence of a link between the two variables. In the realm of bivariate analysis, the technique of cross tabulation is employed to examine the connection between two variables through their arrangement in a tabular format. Results obtained from bivariate analysis for individual and combined basic WASH facilities are presented in Table 4

Table 4. Bivariate frequency distribution of individual and combined basic WASH facilities among the different categories of selected covariates along with p-value.

Explanatory Variables	Handwashing Facility			Sanitation Facility			Drinking Water Facility			Combined WASH Facilities		
	Yes	No	p-value	Yes	No	p-value	Yes	No	p-value	Yes	No	p-value
Division			0.153			< 0.001			0.496			0.151
Barisal	861 (51.8%)	801 (48.2%)		1182 (71.1%)	480 (28.9%)		1643 (98.9%)	19 (1.1%)		698 (42.0%)	964 (58.0%)	
Chittagong	1417 (63.3%)	822 (36.7%)		1472 (65.7%)	767 (34.3%)		2209 (98.7%)	30 (1.3%)		1112 (49.7%)	1127 (50.3%)	
Dhaka	1536 (62.7%)	914 (37.3%)		1438 (58.7%)	1012 (41.3%)		2429 (99.1%)	21 (0.9%)		1116 (45.6%)	1334 (54.4%)	
Khulna	1275 (61.9%)	785 (38.1%)		1348 (65.4%)	712 (34.6%)		1991 (96.7%)	69 (3.3%)		978 (47.5%)	1082 (52.5%)	
Mymensingh	862 (51.0%)	828 (49.0%)		937 (55.4%)	753 (44.6%)		1690 (100.0%)	0 (0.0%)		605 (35.8%)	1085 (64.2%)	
Rajshahi	1258 (61.2%)	798 (38.8%)		1333 (64.8%)	723 (35.2%)		2051 (99.8%)	5 (0.2%)		969 (47.1%)	1087 (52.9%)	
Rangpur	1203 (64.7%)	657 (35.3%)		1085 (58.3%)	775 (41.7%)		1860 (100.0%)	0 (0.0%)		841 (45.2%)	1019 (54.8%)	
Sylhet	912 (57.3%)	681 (42.7%)		989 (62.1%)	604 (37.9%)		1537 (96.5%)	56 (3.5%)		699 (43.9%)	894 (56.1%)	
Place of Residence			< 0.001			< 0.001			0.473			< 0.001
Urban	3934 (71.2%)	1594 (28.8%)		3725 (67.4%)	1803 (32.6%)		5462 (98.8%)	66 (1.2%)		3155 (57.1%)	2373 (42.9%)	
Rural	5390 (53.5%)	4692 (46.5%)		6059 (60.1%)	4023 (39.9%)		9948 (98.7%)	134 (1.3%)		3863 (38.3%)	6219 (61.7%)	
Sex of Household Head			0.005			< 0.001			0.630			< 0.001
Male	8204 (59.3%)	5623 (40.7%)		8585 (62.1%)	5242 (37.9%)		13652 (98.7%)	175 (1.3%)		6139 (44.4%)	7688 (55.6%)	
Female	1120 (62.8%)	663 (37.2%)		1199 (67.2%)	584 (32.8%)		1758 (98.6%)	25 (1.4%)		879 (49.3%)	904 (50.7%)	
Age of Household Head			< 0.001			< 0.001			0.735			< 0.001
<30	1085 (51.3%)	1030 (48.7%)		888 (42.0%)	1227 (58.0%)		2076 (98.2%)	39 (1.8%)		619 (70.7%)	1496 (29.3%)	
30-39	2179 (58.7%)	1531 (41.3%)		2126 (57.3%)	1584 (42.7%)		3678 (99.1%)	32 (0.9%)		1581 (57.4%)	2129 (42.6%)	
40-49	2744 (60.9%)	1762 (39.1%)		2968 (65.9%)	1538 (34.1%)		4451 (98.8%)	55 (1.2%)		2119 (53.0%)	2387 (47.0%)	
50-59	1905 (63.5%)	1097 (36.5%)		2141 (71.3%)	861 (28.7%)		2965 (98.8%)	37 (1.2%)		1537 (48.8%)	1465 (51.2%)	
≥60	1411 (62.0%)	866 (38.0%)		1661 (72.9%)	616 (27.1%)		2240 (98.4%)	37 (1.6%)		1162 (51.0%)	1115 (49.0%)	
Household Size			0.021			< 0.001			0.482			< 0.001

≤5	6899 (59.2%)	4754 (40.8%)		7003 (60.1%)	4650 (39.9%)		11508 (98.8%)	145 (1.2%)		5059 (43.4%)	6594 (56.6%)	
>5	2425 (61.3%)	1532 (38.7%)		2781 (70.3%)	1176 (29.7%)		3902 (98.6%)	55 (1.4%)		1959 (49.5%)	1998 (50.5%)	
Partner's Education Level			< 0.001			< 0.001			< 0.001			< 0.001
No Education	1696 (47.2%)	1898 (52.8%)		1839 (51.2%)	1755 (48.8%)		3532 (98.3%)	62 (1.7%)		1068 (29.7%)	2526 (70.3%)	
Primary	2331 (50.9%)	2252 (49.1%)		2497 (54.5%)	2086 (45.5%)		4516 (98.5%)	67 (1.5%)		1577 (34.4%)	3006 (65.6%)	
Secondary	3100 (64.1%)	1736 (35.9%)		3216 (66.5%)	1620 (33.5%)		4777 (98.8%)	59 (1.2%)		2383 (49.3%)	2453 (50.7%)	
Higher	2197 (84.6%)	400 (15.4%)		2232 (85.9%)	365 (14.1%)		2585 (99.5%)	12 (0.5%)		1990 (76.6%)	607 (23.4%)	
Working Status			< 0.001			< 0.001			0.039			< 0.001
Yes	2916 (56.3%)	2261 (43.7%)		3040 (58.7%)	2137 (41.3%)		5097 (98.5%)	80 (1.5%)		2077 (40.1%)	3100 (59.9%)	
No	6408 (61.4%)	4025 (38.6%)		6744 (64.6%)	3689 (35.4%)		10313 (98.8%)	120 (1.2%)		4941 (47.4%)	5492 (52.6%)	
Media Exposure			< 0.001			< 0.001			< 0.001			< 0.001
Exposed	6096 (66.9%)	3016 (33.1%)		6165 (67.7%)	2947 (32.3%)		9040 (99.2%)	72 (0.8%)		4826 (53.0%)	4286 (47.0%)	
Non-exposed	3228 (49.7%)	3270 (50.3%)		3619 (55.7%)	2879 (44.3%)		6370 (98.0%)	128 (2.0%)		2192 (33.7%)	4306 (66.3%)	
Migration			0.001			0.376			0.359			0.001
Migrant	932 (63.7%)	532 (36.3%)		902 (61.6%)	562 (38.4%)		1449 (99.0%)	15 (1.0%)		718 (49.0%)	746 (51.0%)	
Non-migrant	8392 (59.3%)	5754 (40.7%)		8882 (62.8%)	5264 (37.2%)		13961 (98.7%)	185 (1.3%)		6300 (44.5%)	7846 (55.5%)	
Woman Empowerment			< 0.001			0.779			0.019			0.002
Yes	8398 (60.4%)	5514 (39.6%)		8725 (62.7%)	5187 (37.3%)		13744 (98.8%)	168 (1.2%)		6316 (45.4%)	7596 (54.6%)	
No	926 (54.5%)	772 (45.5%)		1059 (62.4%)	639 (37.6%)		1666 (98.1%)	32 (1.9%)		702 (41.3%)	996 (58.7%)	
Wealth Index			< 0.001			< 0.001			< 0.001			< 0.001
Poor	2171 (36.7%)	3741 (63.3%)		2523 (42.7%)	3389 (57.3%)		5795 (98.0%)	117 (2.0%)		1101 (18.6%)	4811 (81.4%)	
Middle	1722 (55.7%)	1367 (44.3%)		1846 (59.8%)	1243 (40.2%)		3061 (99.1%)	28 (0.9%)		1191 (38.6%)	1898 (61.4%)	
Rich	5431 (82.2%)	1178 (17.8%)		5415 (81.9%)	1194 (18.1%)		6554 (99.2%)	55 (0.8%)		4726 (71.5%)	1883 (28.5%)	
Children Aged 5 and Under in Household			< 0.001			< 0.001			0.682			< 0.001

Yes	4399 (58.8%)	3084 (41.2%)		4465 (59.7%)	3018 (40.3%)		7390 (98.8%)	93 (1.2%)		3212 (42.9%)	4271 (57.1%)	
No	4925 (60.6%)	3202 (39.4%)		5319 (65.4%)	2808 (34.6%)		8020 (98.7%)	107 (1.3%)		3806 (46.8%)	4321 (53.2%)	
Mother's Age of Under 5 Children			< 0.001			< 0.001			0.893			< 0.001
≤30	3551 (56.8%)	2699 (43.2%)		3411 (54.6%)	2839 (45.4%)		6169 (98.7%)	81 (1.3%)		2433 (38.9%)	3817 (61.1%)	
>30	5773 (61.7%)	3587 (38.3%)		6373 (68.1%)	2987 (31.9%)		9241 (98.7%)	119 (1.3%)		4585 (49.0%)	4775 (51.0%)	
Mother's Education of Under 5 Children			< 0.001			< 0.001			< 0.001			< 0.001
No Education	1045 (46.8%)	1188 (53.2%)		1153 (51.6%)	1080 (48.4%)		2189 (97.6%)	53 (2.4%)		682 (30.5%)	1551 (69.5%)	
Primary	2172 (49.0%)	2258 (51.0%)		2392 (54.0%)	2038 (46.0%)		4369 (98.6%)	61 (1.4%)		1439 (32.5%)	2991 (67.5%)	
Secondary	3192 (59.9%)	2135 (40.1%)		3331 (62.5%)	1996 (37.5%)		5267 (98.9%)	60 (1.1%)		2374 (44.6%)	2953 (55.4%)	
Higher	2915 (80.5%)	705 (19.5%)		2908 (80.3%)	712 (19.7%)		3594 (99.3%)	26 (0.7%)		2523 (69.7%)	1097 (30.3%)	

Table 4 reveals that access to basic handwashing facility was significantly associated with the place of residence, household head's sex and age, household size, partner's education level, working status, media exposure, migration status, women empowerment, wealth index, presence of children aged 5 and under, and the mother's age and education of these children. Division, place of residence, sex and age of the household head, household size, partner's education level, working status, media exposure, wealth index, presence of children aged 5 and under, mother's age, and education of these children were all significantly correlated with access to basic sanitation facilities, as shown in Table 4. Moreover, Table 4 shows a significant relationship between access to basic drinking water facility and the partner's education level, employment position, media exposure, women's empowerment, wealth index, and mother's education of children under 5. It is also found from Table 4 that access to combined WASH facilities were significantly associated with the place of residence, sex and age of the household head, household size, partner's education level, working status, media exposure, migration status, women empowerment, wealth index, presence of children aged 5 and under, and the mother's age and education of these children. All of these associations have been found significant as the corresponding p-values are less than $\alpha = 0.05$.

Generalized Linear Mixed Model: Analyzing WASH Facilities

To capture the clustering effect adequately, mixed-effect logistic regression model under GLMM framework has been utilized. In this model, a random effect component has been incorporated for each cluster. This assumes that the baseline odds of the event occurring are consistent for all respondents within the same cluster, while the odds vary from one cluster to another. Four separate

set of GLMMs for binary responses have been considered for basic handwashing facility, sanitation facility, drinking water facility and combined WASH facilities. Significant explanatory variables found in the bivariate analysis have been considered as explanatory variables in GLMMs. The odds ratios (OR), 95% confidence intervals (CI) and p-values obtained from mixed effect logistic regression for analyzing individual and combined basic WASH facilities are presented in Table 5

Table 5. Odds ratios (OR), 95% confidence intervals (CI) and p-values obtained from mixed effect logistic regression for analyzing individual and combined basic WASH facilities.

Covariates	Handwashing Facility			Sanitation Facility			Drinking Water Facility			Combined WASH Facilities		
	OR	p-value	95% CI	OR	p-value	95% CI	OR	p-value	95% CI	OR	p-value	95% CI
Intercept	0.324	<0.001	(0.256, 0.409)	0.261	<0.001	(0.195, 0.349)	18142.3	<0.001	(4813.020, 68385.767)	0.068	<0.001	(0.053, 0.087)
Division												
Barisal	-	-	-	-	-	-	-	-	-	-	-	-
Chittagong	-	-	-	0.643	0.001	(0.490, 0.842)	-	-	-	-	-	-
Dhaka	-	-	-	0.339	<0.001	(0.260, 0.443)	-	-	-	-	-	-
Khulna	-	-	-	0.522	0.011	(0.396, 0.688)	-	-	-	-	-	-
Mymensingh	-	-	-	0.505	0.006	(0.380, 0.671)	-	-	-	-	-	-
Rajshahi	-	-	-	0.582	<0.001	(0.442, 0.767)	-	-	-	-	-	-
Rangpur	-	-	-	0.592	<0.001	(0.449, 0.781)	-	-	-	-	-	-
Sylhet	-	-	-	0.520	0.007	(0.388, 0.696)	-	-	-	-	-	-
Place of Residence												
Urban	-	-	-	-	-	-	-	-	-	-	-	-
Rural	0.892	0.111	(0.776, 1.025)	1.578	<0.001	(1.359, 1.831)	-	-	-	1.040	0.588	(0.903, 1.197)
Sex of Household Head												
Male	-	-	-	-	-	-	-	-	-	-	-	-
Female	1.050	0.450	(0.925, 1.193)	1.220	0.003	(1.070, 1.391)	-	-	-	1.143	0.042	(1.005, 1.301)
Age of Household Head												
<30	-	-	-	-	-	-	-	-	-	-	-	-
30-39	1.108	0.133	(0.972, 1.264)	1.448	0.003	(1.265, 1.657)	-	-	-	1.405	0.013	(1.213, 1.627)
40-49	1.230	0.008	(1.056, 1.433)	1.919	<0.001	(1.644, 2.241)	-	-	-	1.595	<0.001	(1.356, 1.877)
50-59	1.452	0.020	(1.232, 1.712)	2.555	<0.001	(2.159, 3.024)	-	-	-	2.048	<0.001	(1.724, 2.434)
≥60	1.169	0.054	(0.997, 1.370)	2.413	<0.001	(2.051, 2.840)	-	-	-	1.747	<0.001	(1.479, 2.064)
Household Size												
≤5	-	-	-	-	-	-	-	-	-	-	-	-
>5	1.080	0.115	(0.981, 1.189)	1.480	<0.001	(1.339, 1.636)	-	-	-	1.262	0.008	(1.145, 1.392)
Partner's Education Level												
No Education	-	-	-	-	-	-	-	-	-	-	-	-
Primary	0.969	0.573	(0.870, 1.080)	1.027	0.630	(0.921, 1.147)	0.916	0.749	(0.533, 1.573)	1.031	0.607	(0.917, 1.160)
Secondary	1.077	0.212	(0.957, 1.211)	1.116	0.074	(0.990, 1.258)	0.750	0.368	(0.400, 1.404)	1.153	0.026	(1.019, 1.304)
Higher	1.670	<0.001	(1.406, 1.985)	2.036	<0.001	(1.700, 2.438)	1.939	0.169	(0.755, 4.977)	2.034	<0.001	(1.718, 2.407)
Working Status												
No	-	-	-	-	-	-	-	-	-	-	-	-
Yes	0.967	0.311	(0.878, 1.043)	0.873	0.002	(0.799, 0.953)	0.768	0.237	(0.496, 1.189)	0.873	0.003	(0.799, 0.953)
Media Exposure												

Non-exposed	-	-	-	-	-	-	-	-	-	-	-	-
Exposed	1.147	0.001	(1.054, 1.248)	1.066	0.150	(0.976, 1.164)	1.226	0.402	(0.760, 1.978)	1.176	<0.001	(1.077, 1.284)
Migration												
Non-migrant	-	-	-	-	-	-	-	-	-	-	-	-
Migrant	0.901	0.151	(0.783, 1.038)	-	-	-	-	-	-	0.998	0.974	(0.863, 1.154)
Woman Empowerment												
No	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1.165	0.014	(1.032, 1.316)	-	-	-	1.156	0.589	(0.682, 1.959)	1.165	0.335	(0.936, 1.212)
Wealth Index												
Poor	-	-	-	-	-	-	-	-	-	-	-	-
Middle	1.952	<0.001	(1.763, 2.162)	2.203	<0.001	(1.982, 2.449)	2.210	0.006	(1.252, 3.902)	2.522	<0.001	(2.255, 2.820)
Rich	5.888	<0.001	(5.276, 6.572)	7.308	<0.001	(6.472, 8.253)	2.111	0.011	(1.191, 3.741)	8.559	<0.001	(7.639, 9.590)
Children Aged 5 and Under in Household												
No	-	-	-	-	-	-	-	-	-	-	-	-
Yes	1.061	0.184	(0.973, 1.156)	0.972	0.533	(0.889, 1.064)	-	-	-	1.034	0.471	(0.944, 1.131)
Mother's Age of Under 5 Children												
≤30	-	-	-	-	-	-	-	-	-	-	-	-
>30	1.182	0.002	(1.061, 1.316)	1.565	<0.001	(1.405, 1.743)	-	-	-	1.502	<0.001	(1.346, 1.677)
Mother's Education of Under 5 Children												
No Education	-	-	-	-	-	-	-	-	-	-	-	-
Primary	1.051	0.427	(0.929, 1.189)	1.119	0.082	(0.987, 1.268)	1.202	0.543	(0.665, 2.173)	1.010	0.881	(0.882, 1.156)
Secondary	1.288	<0.001	(1.127, 1.471)	1.338	0.019	(1.166, 1.534)	1.791	0.074	(0.946, 3.394)	1.332	0.054	(1.155, 1.537)
Higher	1.990	<0.001	(1.681, 2.355)	1.809	<0.001	(1.523, 2.150)	1.721	0.193	(0.760, 3.897)	1.998	<0.001	(1.681, 2.374)
Variance Component	**0.4357**			**0.4906**			**52.17**			**0.4512**		
ICC	**0.117**			**0.130**			**0. 941**			**0.121**		

ICC=0.117 for basic handwashing facility indicates a moderate level of similarity within clusters. This means that about 11.7% of the total variation in handwashing facility availability can be attributed to differences between clusters, while the remaining 88.3% is due to individual differences within clusters. This suggests that there is some degree of clustering in the data which should be accounted for in the analysis to obtain accurate estimates. Age of household head, partner's education level, media exposure, women empowerment, wealth index, mother's age and mother's education of under 5 children have been found to have significant association with basic handwashing facility. The households with household head aged between 40-49 years and 50-59 years had respectively 23.0% and 45.2% higher odds of having basic handwashing facility compared to the households with household head aged less than 30 years. In case of partner's education level, the study shows that, those who had received higher level of education had 67.0% higher odds of having basic handwashing facility compared to those with no education. The study has also found that, those who were exposed to media had 14.7% higher odds of having basic handwashing facility compared to those who were not exposed to media. Women who were empowered had 16.5% higher odds of having basic handwashing facility compared to the women who were not empowered. Those who belonged to the middle class, had 95.2% higher odds of having basic handwashing facility compared to the poor while the rich had 488.8% higher odds of having basic handwashing facility compared to the poor. The mothers of under 5 children aged more than 30 years had 18.2% higher odds of having basic handwashing facility compared to the mothers who were aged 30 years or less. The mothers who had received secondary and higher level of education had respectively 28.8% and 99.0% higher odds of having basic handwashing facility compared to the mothers with no education.

For basic sanitation facility, ICC=0.130 which indicates about 13.0% of the total variation in sanitation facility availability can be attributed to differences between clusters, while the remaining 87.0% is due to individual differences within clusters. Table 5 demonstrates that, division, place of residence, sex and age of household head, household size, partner's education level, working status, wealth index, mother's age and mother's education of under 5 children had significant association with basic sanitation facility. Chittagong, Dhaka, Khulna, Mymensingh, Rajshahi, Rangpur and Sylhet had respectively 35.7%, 66.1%, 47.8%, 49.5, 41.8%, 40.8% and 48.0% lower odds of having basic sanitation facility compared to Barisal. Those living in rural areas had 57.8% higher odds of having basic sanitation facility compared to those living in urban areas. The households with female household head had 22.0% higher odds of having basic sanitation facility compared to those with male household head. The households with household head aged between 30-39, 40-49 years, 50-59 years, 60 years and above had respectively 44.8%, 91.9%, 155.5% and 141.3% higher odds of having basic sanitation facility compared to the households with household head aged less than 30 years. Table 9 clarifies that, the households with more than five members had 48.0% higher odds of having basic sanitation facility compared to the households with five or less members. In case of partner's education level, the study shows that, those who had received higher level of education had 103.6% higher odds of having basic sanitation facility compared to those with no education. The working respondents had 12.7% lower odds of having basic sanitation facility compared to the non-working respondents. Moreover, those who belonged to the middle class, had 120.3% higher odds of having basic sanitation facility compared to the poor while the rich had 630.8% higher odds of having basic sanitation facility compared to the poor. The mothers of under 5 children aged more than 30 years had 56.5% higher odds of having basic sanitation facility compared to the mothers who were aged 30 years or less. The mothers who had

received secondary and higher level of education had respectively 33.8% and 80.9% higher odds of having basic sanitation facility compared to the mothers with no education.

Table 5 reveals that, ICC=0.941 for basic drinking water facility which suggests a very high level of similarity within clusters. This shows that while individual differences within clusters account for only 5.9% of the variance in the availability of basic drinking water facilities, cluster differences account for 94.1% of the total variance. It has been found that, only wealth index had significant association with having basic drinking water facility. Table 5 shows that, those who belonged to the middle class, had 121.0% higher odds of having basic drinking water facility compared to the poor and the rich had 111.1% higher odds of having basic drinking water facility compared to the poor.

Table 5 suggests ICC=0.121 for combined WASH facilities which indicates a moderate level of similarity within clusters. This means that about 12.1% of the total variation in combined WASH facilities availability can be attributed to differences between clusters, while the remaining 87.9% is due to individual differences within clusters. It can be seen from Table 5 that, sex and age of household head, household size, partner's education level, working status, media exposure, wealth index, mother's age and mother's education of under 5 children had significant effect on combined WASH facilities. The households with female household head had 14.3% higher odds of having combined WASH facilities compared to those with male household head. The households with household head aged between 30-39, 40-49 years, 50-59 years, 60 years and above had respectively 40.5%, 59.5%, 104.8% and 74.7% higher odds of having combined WASH facilities compared to the households with household head aged less than 30 years. The households with more than five members had 26.2% higher odds of having combined WASH facilities compared to the households with five or less members. The study shows that, those whose partners had

received secondary and higher level of education had respectively 15.3% and 103.4% higher odds of having combined WASH facilities compared to those whose partners were illiterate. The working respondents had 12.7% lower odds of having combined WASH facilities compared to the non-working respondents. The study has also found that, those who were exposed to media had 17.6% higher odds of having combined WASH facilities compared to those who were not exposed to media. Those who belonged to the middle class, had 152.2% higher odds of having combined WASH facilities compared to the poor while the rich had 755.9% higher odds of having combined WASH facilities compared to the poor. The mothers of under 5 children aged more than 30 years had 50.2% higher odds of having combined WASH facilities compared to the mothers who were aged 30 years or less. The mothers who had received higher level of education had 99.9% higher odds of having combined WASH facilities compared to the mothers with no education.

Mixed Effect vs. Fixed Effect Logistic Regression Model

For analyzing individual and combined WASH facilities, the performances of mixed effect logistic regression model (model 1) and fixed effect logistic regression model (model 2) have been compared on the basis of AIC values and by performing likelihood ratio test (LRT).

Performance Comparison Based on LRT

The log-likelihood values for mixed effect logistic regression model (model 1) and fixed effect logistic regression model (model 2) along with χ^2-value and p-value are presented in Table 6.

Table 6. Performance Comparison of Mixed effect logistic regression model (Model 1) and Fixed effect logistic regression model (Model 2) for individual and combined WASH facilities Based on LRT.

Facility	Model	Log-likelihood	χ^2	p-value
Basic Handwashing Facility	Model 1	-8663.7		
	Model 2	-8900.0	473.15	<0.001

Facility	Model			
Basic Sanitation Facility	Model 1	-8290.4		
	Model 2	-8557.5	534.33	<0.001
Basic Drinking Water Facility	Model 1	-570.2		
	Model 2	-1029.9	861.97	<0.001
Combined WASH Facilities	Model 1	-8218.7		
	Model 2	-8435.5	433.65	<0.001

Here, p-value has been obtained for null hypothesis that the fixed effect logistic regression model performs equally well as mixed effect logistic regression model. Since, p-value has been found to be less than 0.001, the null hypothesis can be rejected which reveals that the mixed effect model outperforms the fixed effect model significantly. This indicates that the mixed effect logistic regression model offers a substantially better fit to the data than the fixed effect logistic regression model for analyzing individual and combined WASH facilities. Hence, the use of the mixed effect logistic regression model is appropriate in this case.

Performance Comparison Based on AIC Values

The AIC values for mixed effect logistic regression model (model 1) and fixed effect logistic regression model (model 2) are presented in table 7. The model with a lower AIC value suggests a better fit for the data.

Table 7. Performance Comparison of Mixed effect logistic regression model (Model 1) and Fixed effect logistic regression model (Model 2) for individual and combined WASH facilities Based on AIC values.

Facility	Model	AIC
Basic Handwashing Facility	Model 1	17373.3
	Model 2	17844.0
Basic Sanitation Facility	Model 1	16636.8
	Model 2	17169.0
Basic Drinking Water Facility	Model 1	1166.4
	Model 2	2083.8
Combined WASH Facilities	Model 1	16483.4
	Model 2	16915.0

From Table 7, it has been found that, the mixed effect logistic regression model (model 1) has lower AIC value than the usual logistic regression model with fixed effects (model 2). Therefore, mixed effect logistic regression model seems to be a better fit for analyzing individual and combined WASH facilities.

Discussion

Traditional logistic regression models are not suitable for analyzing clustered data, as responses within the same cluster are not independent, despite clusters being independent of each other. To account for this dependency, a GLMM is necessary for obtaining consistent and efficient estimators for the regression parameters of interest. This paper reveals several key determinants and barriers to WASH facilities in Bangladesh using clustered data from BDHS 2022, employing a GLMM for binary response which is also known as mixed effect logistic regression model. Based on AIC values and LRT, this study has found that the mixed effect logistic regression model outperforms the fixed effect logistic regression model. This justifies the adoption of the mixed effect logistic regression model. Also, the existence of correlation within the same cluster has been justified by calculating ICC while analyzing individual and combined basic WASH facilities.

Among all the divisions, Barisal has been found to have higher likelihood of having basic sanitation facility which remains consistent with the previous study [18]. This research confirms the results of past studies by showing that wealth index had a positive association with both individual and combined basic WASH facilities [15-17,19,22,25]. Moreover, it has been discovered that media exposure was positively associated with basic handwashing facility and combined WASH facilities, whereas working status was negatively associated with both basic sanitation and combined WASH facilities. Additionally, the study has discovered that basic

handwashing facility, sanitation facility, and combined WASH facilities were positively impacted by the age of the household head. The female-headed households and the households with more than 5 members have been found to have higher likelihood of having basic sanitation and combined WASH facilities. Furthermore, the study shows that the rural residents were more likely to have basic sanitation facility. These findings also align with the findings of previous researches [15-18,24-25].

Partner's education level, under 5 children's mothers' age and education played important role in having adequate basic handwashing facility, sanitation facility and combined WASH facilities. The respondents having partners with low education level or no education, have been found to be less likely to have these facilities. Also, the younger mothers and the mothers with low education level or no education have been found to have lower likelihood of having these facilities. These results are in line with earlier investigations [27]. The study further highlights a positive correlation between basic handwashing facility and women's empowerment, reinforcing findings from previous research [28].

Conclusion

This study identifies key determinants and barriers to effective WASH practices in Bangladesh, essential for preventing waterborne diseases and ensuring fair access to services. The findings show that improving handwashing facility requires prioritizing partner and mother education, media exposure, and women empowerment. Sanitation facilities should address regional differences, household sizes, and economic support. Drinking water facility access must focus on lower wealth indices. Combined WASH facilities benefit from targeting working respondents, male-headed and smaller households, and enhancing education and media exposure.

Limitations and Further Scopes

In this study, cross-sectional data have been utilized, which may introduce selection bias and information bias. Consequently, the assessment of changes in WASH facilities over time—critical for understanding trends and causal relationships—may be limited. To enhance the depth of findings, researchers can consider longitudinal studies. Additionally, exploring informal social accountability mechanisms (such as rewards, sanctions, voice, and responsiveness), identifying additional factors influencing access to basic WASH services, and evaluating behavior change interventions would enrich the research.

Acknowledgments

The authors are grateful to the Demographic and Health Survey (DHS) for executing a nationwide survey and made its data publicly available.

References

1. United Nations General Assembly. *Transforming our world: the 2030 Agenda for Sustainable Development*. 2015. Available from: https://sdgs.un.org/2030agenda

2. Wolf J, Hunter PR, Freeman MC, Cumming O, Clasen T, Bartram J, et al. "Impact of drinking water, sanitation and handwashing with soap on childhood diarrhoeal disease: updated meta-analysis and meta-regression". In: *Tropical medicine & international health* 23.5 (2018), pp. 508–525. doi: 10.1111/tmi.13051

3. Luby SP, Rahman M, Arnold BF, Unicomb L, Ashraf S, Winch PJ, et al. "Effects of water quality, sanitation, handwashing, and nutritional interventions on diarrhoea and child growth in rural Bangladesh: a cluster randomised controlled trial". In: *The Lancet Global Health* 6.3 (2018), e302–e315. doi: 10.1016/s2214-109x(17)30490-4

4. Kathryn H Jacobsen and James S Koopman. "The effects of socioeconomic development on worldwide hepatitis A virus seroprevalence patterns". In: *International journal of epidemiology* 34.3 (2005), pp. 600–609. doi: 10.1093/ije/dyi062

5. Im J, Islam MT, Ahmmed F, Kim DR, Islam Khan A, Zaman K, et al. "Can existing improvements of water, sanitation, and hygiene (WASH) in urban slums reduce the burden of typhoid fever in these settings?" In: *Clinical Infectious Diseases* 72.11 (2021), e720–e726. doi: 10.1093/cid/ciaa1429

6. Man Mohan Mehndiratta, Prachi Mehndiratta, and Renuka Pande. "Poliomyelitis: historical facts, epidemiology, and current challenges in eradication". In: *The Neurohospitalist* 4.4 (2014), pp. 223–229. doi: 10.1177/1941874414533352

7. Ashraf S, Islam M, Unicomb L, Rahman M, Winch PJ, Arnold BF, et al. "Effect of improved water quality, sanitation, hygiene and nutrition interventions on respiratory illness in young

children in rural Bangladesh: a multi-arm cluster-randomized controlled trial". In: *The American journal of tropical medicine and hygiene* 102.5 (2020), p. 1124. doi: [10.4269/ajtmh.19-0769](#)

8. Nicholas John Ashbolt. "Microbial contamination of drinking water and disease outcomes in developing regions". In: *Toxicology* 198.1-3 (2004), pp. 229– 238. doi: [10.1016/j.tox.2004.01.030](#)

9. World Health Organization and United Nations Children's Fund. *Progress on WASH in health care facilities 2000–2021: special focus on WASH and infection prevention and control*. World Health Organization, 2023.

   Available from: https://www.who.int/publications/i/item/9789240058699

10. World Health Organization and United Nations Children's Fund. *Progress on household drinking water, sanitation and hygiene 2000-2020: five years into the SDGs*. World Health Organization, 2021. Available from: https://www.who.int/publications/i/item/9789240030848

11. Walter W Stroup. *Generalized linear mixed models: modern concepts, methods and applications*. CRC press, 2012. doi: [10.1201/b13151](#)

12. National Institute of Population Research & Training (NIPORT) and ICF. *Bangladesh Demographic and Health Survey 2022: Final Report*. Dhaka, Bangladesh. 2024. Available from: https://www.dhsprogram.com/pubs/pdf/FR386/FR386.pdf

13. World Health Organization and United Nations Children's Fund (UNICEF). *Progress on drinking water, sanitation and hygiene 2017 update and SDG baselines*. 2017. Available from: https://data.unicef.org/resources/progress-drinking-water-sanitation-hygiene-2017-update-sdg-baselines/

14. Girmay AM, Weldegebriel MG, Mengesha SD, Serte MG, Weldetinsae A, Alemu ZA, et al. "Factors influencing access to basic water, sanitation, and hygiene (WASH) services in schools of Bishoftu Town, Ethiopia: a cross-sectional study". In: *Discover Sustainability* 4.1 (2023), p. 5. doi: 10.1007/s43621-023-00122-0

15. Akter J, Islam MR, Akter S, Rahman MM, Hossain F, Anam MR, et al. "Equity in access to safely managed sanitation and prevalence of diarrheal diseases in Bangladesh: A national and sub-national analysis". In: *BMC Infectious Diseases* 22.1 (2022), p. 871. doi: 10.1186/s12879-022-07884-4

16. Gaffan N, Kpozèhouen A, Dégbey C, Glèlè Ahanhanzo Y, Glèlè Kakaï R, Salamon R. "Household access to basic drinking water, sanitation and hygiene facilities: secondary analysis of data from the demographic and health survey V, 2017–2018". In: *BMC Public Health* 22.1 (2022), p. 1345. doi: 10.1186/s12889-022-13665-0

17. Andualem Z, Dagne H, Azene ZN, Taddese AA, Dagnew B, Fisseha R, et al. "Households access to improved drinking water sources and toilet facilities in Ethiopia: a multilevel analysis based on 2016 Ethiopian Demographic and Health Survey". In: *BMJ open* 11.3 (2021), e042071. doi: 10.1136/bmjopen-2020-042071

18. Ahmed MS, Islam MI, Das MC, Khan A, Yunus FM. "Mapping and situation analysis of basic WASH facilities at households in Bangladesh: evidence from a nationally representative survey". In: *PloS One* 16.11 (2021), e0259635. doi: 10.1371/journal.pone.0259635

19. Simelane MS, Shongwe MC, Vermaak K, Zwane E. "Determinants of households' access to improved drinking water sources: a secondary analysis of eswatini 2010 and 2014 multiple indicator cluster surveys". In: *Advances in Public Health* 2020.1 (2020), p. 6758513. doi: 10.1155/2020/6758513

20. Akpakli DE, Manyeh AK, Akpakli JK, Kukula V, Gyapong M. "Determinants of access to improved sanitation facilities in rural districts of southern Ghana: evidence from Dodowa Health and Demographic Surveillance Site". In: *BMC Research Notes* 11 (2018), pp. 1–7. doi: 10.1186/s13104-018-3572-6

21. Tuyet-Hanh TT, Lee JK, Oh J, Van Minh H, Ou Lee C, Hoan LT, et al. "Household trends in access to improved water sources and sanitation facilities in Vietnam and associated factors: findings from the Multiple Indicator Cluster Surveys, 2000–2011". In: *Global Health Action* 9.1 (2016), p. 29434. doi: 10.3402/gha.v9.29434

22. Decio Ribeiro Sarmento MPH. "An analysis of access to improved drinking water and sanitation and distance to the water source in a newly independent country, Timor-Leste: assessing geographical and socioeconomic disparities". In: (2015). doi: 10.57709/7000548

23. Gurung R, Tirkey C, Takri KK, Diyali N, Choubey M, Rai R. "Determinants of access to improved drinking water and sanitation in India: evidence from India Human Development Survey-II (IHDS)". In: *Water Policy* 25.10 (2023), pp. 980–995. doi: 10.2166/wp.2023.083

24. Gebremichael SG, Yismaw E, Tsegaw BD, Shibeshi AD. "Determinants of water source use, quality of water, sanitation and hygiene perceptions among urban households in North-West Ethiopia: A cross-sectional study". In: *Plos one* 16.4 (2021), e0239502. doi: 10.1371/journal.pone.0239502

25. Azanaw J, Abera E, Malede A, Endalew M. "A multilevel analysis of improved drinking water sources and sanitation facilities in Ethiopia: Using 2019 Ethiopia mini demographic and health survey". In: *Frontiers in Public Health* 11 (2023), p. 1063052. doi: 10.3389/fpubh.2023.1063052

26. Ayisha Matuamo Mahama, Kwabena Asomanin Anaman, and Isaac Osei Akoto. "Factors influencing householders' access to improved water in low-income urban areas of Accra, Ghana". In: *Journal of water and health* 12.2 (2014), pp. 318–331. doi: 10.2166/wh.2014.149

27. Shalik Ram Dhital, Catherine Chojenta, and Deborah Loxton. "Multi-level factors associated with utilization of water, sanitation and hygiene services by mothers in Nepal". In: *Plos one* 19.3 (2024), e0283379. doi: 10.1371/journal.pone.0283379

28. Daniel Bogale. "Association of Women's Empowerment With Water, Sanitation, and Hygiene (Wash) Services in Ethiopia: A Cross-sectional Analysis" [Preprint]. 13 January 2021 [version 1]. doi: 10.21203/rs.3.rs-142619/v1

29. Göran Broström and Henrik Holmberg. "Generalized linear models with clustered data: Fixed and random effects models". In: *Computational Statistics & Data Analysis* 55.12 (2011), pp. 3123–3134. doi: 10.1016/j.csda.2011.06.011

30. Tuerlinckx F, Rijmen F, Verbeke G, De Boeck P. "Statistical inference in generalized linear mixed models: A review". In: *British Journal of Mathematical and Statistical Psychology* 59.2 (2006), pp. 225–255. doi: 10.1348/000711005X79857

31. Jianxin Pan and Robin Thompson. "Gauss-Hermite quadrature approximation for estimation in generalised linear mixed models". In: *Computational statistics* 18 (2003), pp. 57–78. doi: 10.1007/s001800300132

32. Dian Handayani, Khairil Anwar Notodiputro, Kusman Sadik, Anang Kurnia. "A comparative study of approximation methods for maximum likelihood estimation in generalized linear mixed models (GLMM)". In: *AIP Conference Proceedings*. Vol. 1827. 1. AIP Publishing. 2017. doi: 10.1063/1.4979449

33. Xu Ning, Francis Hui, and Alan Welsh. "Asymptotic Results for Penalized Quasi-Likelihood Estimation in Generalized Linear Mixed Models". In: *arXiv preprint arXiv:2405.01026* (2024). doi: 10.48550/arXiv.2405.01026

34. D Hedeker and RD Gibbons. "Mixed-effects regression models for binary outcomes". In: *Longitudinal data analysis*. Hoboken: Wiley-Interscience (2006), pp. 149–86.

35. Shersten Killip, Ziyad Mahfoud, and Kevin Pearce. "What is an intracluster correlation coefficient? Crucial concepts for primary care researchers". In: *The Annals of Family Medicine* 2.3 (2004), pp. 204–208. doi: 10.1370/afm.141

36. Sandra M. Eldridge, Obioha C. Ukoumunne, and John B. Carlin. "The intra-cluster correlation coefficient in cluster randomized trials: a review of definitions". In: *International Statistical Review* 77.3 (2009), pp. 378-394. doi: 10.1111/j.1751-5823.2009.00092.x