Title: Partner Military Deployment During Wartime Is Associated with Maternal Depression and Impaired Bonding: A Matched-Control Study from the Israel-Hamas War

Hadas Allouche-Kam[1,2], Sabrina J Chan[1], Isha H Arora[1], Christina T Pham[1], Inbal Reuveni[3], Eyal Sheiner[4], Sharon Dekel*[1,5]

[1]Massachusetts General Hospital, Department of Psychiatry, Boston, MA

[2]Department of Obstetrics and Gynecology, Hadassah Medical Center, Hebrew University of Jerusalem, Jerusalem, Israel.

[3]Department of Psychiatry, Tel Aviv Sourasky Medical Center, Tel Aviv University, Tel Aviv

[4]Department of Obstetrics and Gynecology, Soroka University Medical Center, Ben Gurion University of the Negev, Beer-Sheva, Israel.

[5]Harvard Medical School, Department of Psychiatry, Boston, MA

*Address correspondence and reprint requests to Dr. Sharon Dekel, Department of Psychiatry, Massachusetts General Hospital. B62 113st, Boston MA. sdekel@mgh.harvard.edu

Abstract

Purpose. The pregnancy and postpartum periods represent a time of heightened psychological vulnerability with implications for the offspring. Knowledge of the mental health of perinatal women exposed to armed conflict when their partner is in military deployment is scarce. Methods. This matched-control, survey-based study included a sample of 429 women recruited during the first months of the Israel-Hamas War who were pregnant or within six months postpartum. Women who had a partner in military deployment were matched primarily on demographics, prior mental health, and trauma exposure to women whose partner was no longer deployed. Results. We found that nearly 44% of pregnant women with a partner deployed endorsed probable depression. This group was more than twice as likely to endorse probable depression than matched pregnant controls. Likewise, postpartum women with a partner deployed reported significantly more maternal-infant attachment problems than the matched postpartum group of partners not deployed. Importantly, analysis showed that partner's active deployment was related to maternal depression and attachment problems via reduced perceived social support. Conclusions. Partner military deployment during conditions of war can serve as a major psychological stressor for pregnant and postpartum women. It can heighten psychiatric morbidity and interfere with attachment to the infant in part by diminished social support. Implementation of community-based services for the peripartum population is crucial during times of war and other large-scale traumas.

Keywords.

Postpartum Depression; Maternal-Infant Attachment; Social Support; Partner Deployment; October 7th; Israel-Hamas War.

58   **Article Highlights**

59   Partner military deployment increases risk for antepartum depression and maternal-infant bonding problems.

60   Reduced social support explains these maternal outcomes.

61   Clinical attention to the wellbeing of the peripartum population is warranted during times of collective trauma.

Introduction

On October 7, 2023, the Israel-Hamas conflict escalated when the Hamas organization infiltrated southern Israel through coordinated land, sea, and air attacks. Israeli civilians residing in communities near the border, Nova festival participants, and soldiers were murdered, raped, and taken hostage, without distinction between children and elderly or Jewish and non-Jewish victims (Secretary-General of the United Nations 2024). This event, classified as a war crime by the UN Human Rights Council (Office of the United Nations High Commissioner for Human Rights 2024), resulted in over 1,250 fatalities and 251 hostages (U.S. Department of Defense; Levany et al. 2023; Secretary-General of the United Nations 2024). In response, Israel declared war, and the Israeli Defense Forces (IDF) invaded Gaza, striking multiple Hamas targets. There was a rapid and massive deployment of reserve duty by the IDF, with many soldiers serving on active duty for months.

Mental health effects of man-made trauma, such as war and terrorism, on individuals directly exposed, those indirectly exposed, and the community at large have been extensively studied (Silver 2002; Yehuda and Hyman 2005; Perlman et al. 2011; Morina et al. 2018; Castro-Vale et al. 2019; Brown 2020; Hoppen et al. 2021). Man-made trauma shatters people's core cognitive schemas that the world is a predictable and safe place (Chrisman and Dougherty 2014). The amplitude of studies report ensuing psychiatric conditions among direct and indirect trauma-exposed individuals (Punamaki et al. 2018; Druetz et al. 2020; Klapper-Goldstein et al. 2024; Krupelnytska and Morozova-Larina 2025) — most notably post-traumatic stress disorder (PTSD) and depression, which may critically and persistently undermine the person's physical health (VanItallie 2002; Glaesmer et al. 2011; Nemeroff and Goldschmidt-Clermont 2012; Roberts et al. 2015; Rosenbaum et al. 2015; Perkins et al. 2021). In the general community, an estimated 1 in 4 individuals residing in a war-conflict zone will be affected by PTSD (Ahmed et al. 2024) with potential intergenerational effects on the offspring, underscoring the magnitude of psychiatric morbidity in exposed communities (Devakumar et al. 2014; Weinstein et al. 2018; Yehuda and Lehrner 2018; Castro-Vale et al. 2019; Brown 2020; Hoppen et al. 2021). Not surprisingly, a national survey revealed elevated mental health

complications among the Israeli community following the October 7th, 2023, attack (Levi-Belz et al. 2024; Hasson-Ohayon and Horesh 2024).

There is a critical gap in knowledge concerning the effects of war exposure on the perinatal population. Pregnancy and the postpartum period have been repeatedly documented as times of potential psychological vulnerability (Louise M. Howard 2020; Wisner et al. 2024). Perinatal depression is the leading pregnancy complication, affecting 1 out of 7 women residing in non-war-conflict zones (Gavin et al. 2005; Curry et al. 2019). It increases risk for maternal (Ackerman-Banks et al. 2023; Meaidi 2024), fetus (Alder et al. 2007; Grote et al. 2010), and infant health complications (Jarde et al. 2016; Smith et al. 2020; Rogers et al. 2020; Simonovich et al. 2021). In its extreme form, it can result in maternal death and infanticide (Jones et al. 2014). Although maternal stress has been linked to poor childbirth outcomes, studies on the adversity of war and terrorism on perinatal mental health are largely lacking (Yehuda et al. 2005; Engel et al. 2005; Klapper-Goldstein et al. 2024; Krupelnytska and Morozova-Larina 2025).

An important factor to consider in understanding the unique impact of war on the perinatal population is partner military deployment. Partners often serve as the primary source of emotional reassurance and practical aid (Houts et al. 2008). Lack of partner support increases risk for perinatal depression and maternal-infant attachment problems (Bilszta et al. 2008; Biaggi et al. 2016). Although the literature is in its infancy, existing studies on partner deployment reveal substantial depression in women residing in non-war zones (Razurel et al. 2013; Tarney et al. 2015; Godier-McBard et al. 2019; Pretorius et al. 2024).

A large body of research underscores the adaptive role of social support in coping following a traumatic event (Wickramaratne et al. 2022; Calhoun et al. 2022). Post-trauma recovery has been understood within an interpersonal framework, whereby relational partners facilitate coping through the provision of social support (Calhoun et al. 2022). Partner deployment may act as a stressor that heightens feelings of helplessness and loneliness while depleting psychological resources needed to cope with trauma exposure during the demanding perinatal period

(Robrecht et al. 2008; Klaman and Turner 2016; Godier-McBard et al. 2019; Pretorius et al. 2024). To our knowledge, no study has examined the impact of partners' deployment on mental health and maternal-infant attachment among pregnant and postpartum women who have been exposed, both directly and indirectly, to war-related stressors.

We studied a sample of 165 pregnant and 264 postpartum women in the first months following the events of October 7th and the outbreak of the war, including some who had partners actively deployed. We asked whether partner deployment is associated with maternal mental health and maternal-infant attachment, and whether social support mediates the relationship between partner deployment and maternal outcomes.

Materials and Methods

The study commenced on January 19th, 2024, during the first months of the Israel-Hamas ("Iron Swords") War to understand the impact of terrorism and war on maternal health (n = 1,085). In the months leading up to study recruitment (i.e., October 2023-January 2024), over 10,600 rockets and mortar rounds were launched toward Israel and during the recruitment, there were 425 missile alerts across the country (The Meir Amit Intelligence and Terrorism Information Center 2024; Israel's National Emergency Portal 2024) and 10 terrorist attacks (The Meir Amit Intelligence and Terrorism Information Center 2024).

Women residing in Israel during the war who were pregnant or gave birth within the past six months were recruited via social media and professional organizations. They were asked to complete a survey about mental health, maternal-infant attachment, trauma exposure including partner deployment, and obstetrics and childbirth-related factors. The study was approved by Mass General Brigham Human Research Committee. Participants provided implied consent by providing responses in study questionnaires.

A total of 466 (42.9%) of participants reported partner deployed since October 7th, 2023, either currently or no longer. 7.9% were missing information on important study outcomes, resulting in a sample of 429 participants.

This includes 165 (38.5%) pregnant women and 264 (61.5%) postpartum women. 99.3% were Jewish. Information

regarding the sample characteristics is provided in Table 1.

Geographic exposure zones were classified based on IDF Home Front Commands: Red, Orange, Yellow and

Green, defined by time available to reach shelter for threat of rocket and missile attacks from the time of the siren

(i.e., immediate to 15 seconds, i.e., highest level of proximity to combat zone, 30, 45, and 60-90 seconds). The

majority of the sample resided during October 7th in the Green zone (41.7%), 24.9% and 24.5% in the Orange and

Yellow zones, respectively, and 8.9% in the Red zone with 21.05% displaced. 74.1% of participants knew people

who were murdered, killed, injured, and/or taken hostage as a result of October 7th and the war.

Insert Table 1 here

Measures

Depression symptoms were assessed using the Edinburgh Postnatal Depression Scale (EPDS), a 10-item

questionnaire assessing depression symptoms over the past seven days (Cox et al. 1987). Higher scores indicate a

greater level of symptoms. The EPDS is the recommended screening tool in perinatal settings, as determined by

American College of Obstetricians and Gynecologists (ACOG) (American College of Obstetricians and

Gynecologists 2023) and the US Preventive Services Task Force (Curry et al. 2019). We used the cutoff score of 14

to indicate probable depression (Cox et al. 1987). Reliability was high ($\alpha = 0.86$).

Posttraumatic stress disorder (PTSD) symptoms were measured with the well-validated PTSD Checklist for

DSM-5 (PCL-5)(Weathers et al. 2013). This self-report tool is designed to assess the severity of post-traumatic stress

disorder (PTSD) symptoms in the past month in regard to a specified trauma (here, October 7th events) and

recommended by the Veterans of Affairs (VA) for provisional PTSD (Weathers 2013; Bovin et al. 2016). It lists the

20 DSM-5 PTSD symptoms. Higher scores indicate higher symptom levels. We used the cutoff score of 33 to

determine probable PTSD (Bovin et al. 2016) ($\alpha = 0.91$).

Maternal-fetal attachment in pregnancy was measured using the Maternal-Fetal Attachment Scale (MFAS)

(Cranley 1981). This is a 24-item questionnaire designed to evaluate the emotional connection between a mother

and her fetus. Higher scores indicate greater maternal attachment. It shows strong psychometric properties (Lindgren

2001; Wittkowski et al. 2020) ($\alpha = 0.88$).

Maternal-infant attachment during the postpartum period was assessed using the Maternal Attachment

Inventory (MAI) (Müller 1994). This 26-item self-report scale evaluates a mother's perceptions and emotions

regarding her infant during the first year postpartum. It accords with observational data of bond behavior

(Wittkowski et al. 2020). Higher scores indicate attachment problems ($\alpha = 0.94$).

The Multidimensional Scale of Perceived Social Support (MSPSS) is a widely used 12-item questionnaire

to assess perceived social support with higher scores indicating greater support. The scale demonstrates strong

reliability and validity across trauma-exposed and peripartum populations (Dambi et al. 2018; Bedaso et al. 2021)

($\alpha = 0.93$).

Socio-demographics, prior mental health problems, and obstetric information (e.g. gestational week, giving

birth before or after the war, delivery mode, neonatal intensive care unit (NICU) admission) were measured by single

items. Trauma exposure included information about participants' place of residence on October 7th, defined by

Home Front Command zones, whether participants knew people murdered, taken hostage, or injured, and degree of

exposure to October 7th events/war via social media (i.e., exposure to uncensored movies).

Data Analysis

Demographic data for maternal age and Home Front Commands exposure zone were noted to be missing at random,

and multiple imputations using the *mice* R package (Buuren and Groothuis-Oudshoorn 2011) were performed. We

conducted a full propensity score procedure using *MatchIt* R package (Ho et al. 2011) for pregnant and postpartum

participants, separately, and matched participants whose partner was in active military deployment (study group)

and participants whose partner was no longer in active deployment (control group). Propensity scores were estimated using logistic regression, modeling the probability of being in the study group as a function of selected baseline characteristics. Full matching was applied, which retained all participants by constructing matched sets where each set contained at least one individual from the study group and at least one from the control group. Participants were assigned weights based on the structure of matched sets, and all statistical analyses incorporated these weights. Groups were matched on demographics (i.e., maternal age, education, income), primiparity, prior mental health, exposure to October 7th and the war (i.e., residing in the Red zone, knowing civilians murdered and/or civilians/soldiers taken hostage, and watching related uncensored videos). Pregnant women were also matched on gestational week period and postpartum women on whether they gave birth before or after October 7th.

Following the matching, we compared groups on depression and PTSD, maternal-infant attachment, and social support. Kolmogorov-Smirnov and Shapiro-Wilk tests revealed that all factors in consideration were significantly positively skewed ($p < 0.001$). Analyses were conducted separately for pregnant and postpartum women. To estimate differences between women with a partner currently in active deployment or not, Mann-Whitney U tests were performed for quantitative measures and logistic regressions for estimating differences in categorical measures. Confidence Intervals (CIs) for the tests were based on bootstrapping of 1000 resampling cycles. All analyses were conducted in R (R Core Team 2023).

To examine whether social support mediated the effects of study group (women with versus without a partner who is deployed) on maternal outcomes, we conducted a mediation model using observed measures within a Structural Equation Modeling (SEM) framework, estimating a saturated model in *lavaan* R package(Rosseel 2012). Significance was estimated by bias-corrected bootstrap analysis with 2000 sampling cycles. Missing data were handled with the Full Information Maximum Likelihood (FIML) procedure. The group (0 = partner in non-active deployment, 1 = partner in active deployment) served as the predictor, social support as the mediator, and peripartum depression and attachment problems as the outcomes in separate analyses.

Results

The rate of clinically significant depression symptoms (≥ 14, EPDS), indicative of probable depression, in pregnant women with partners in active deployment was 43.96% in comparison to 25.49% in the matched group of partners not deployed and represented a two-fold increase ($\chi^2 = 4.01$, Odds Ratio, OR = 2.29, 95% CI = 1.10 – 5.00, p = 0.05). No differences were found in depression rates among postpartum women with and without a partner deployed and in probable PTSD rates for pregnant and postpartum women by deployment status (Table 2). The depression rate in the pregnant active partner deployed group was also higher than in postpartum women with a partner deployed ($\chi^2 = 8.51$, OR = 2.46, 95% CI = 1.38 – 4.44, p = 0.00).

Insert Table 2 here

There were differences in maternal-infant attachment related to partner deployment. The active deployment group endorsed more attachment problems than controls (Z = 2.36, r = 0.15, CI = 0.03 – 0.26, p = 0.02) (Table 2), although there were no differences between the groups in delivery mode ($\chi^2 = 4.83$, p = 0.19) and NICU admission ($\chi^2 = 7.5e^{-30}$, p = 1) (Table 1). No group differences were noted in maternal-fetal attachment (Table 2).

Both pregnant and postpartum women with a partner in active deployment reported significantly lower levels of social support than matched controls (Pregnant group: $Z = 3.16$, r = 0.25, CI = 0.1 – 0.38, p = 0.00; Postpartum group: $Z = 3.07$, r = 0.19, CI = 0.07 – 0.30, p = 0.00) (Table 2). For pregnant women, social support significantly mediated the path between study group and maternal depression (Figure 1). Partner in active deployment was related to lower social support ($\beta = -0.30$, p < 0.001), which in turn was associated with higher depression ($\beta = 0.39$, p < .001) (Figure 1). For postpartum women, social support mediated the path between study group and maternal-infant attachment, although the results were near significance. Partner in active deployment was related to lower social support ($\beta = -0.26$, p < 0.001), which in turn was associated with more attachment problems ($\beta = -0.16$, p < 0.1) (Figure 2).

Insert Mediation Analysis here

Discussion and Conclusions

This study examined the psychological adjustment of pregnant and postpartum women exposed to war-related

stressors during a time that their partner was in military deployment. Among pregnant women who were similar on

demographic background, prior mental health, war-related exposure, and gestational week, those with a partner

actively deployed were twice as likely to endorse probable depression compared to those with a partner no longer

in active deployment. Similarly, postpartum women with a deployed partner reported greater difficulties bonding

with their infant compared to those whose partner was not actively deployed, despite being comparable on other

factors. As importantly, maternal adversity associated with having a partner deployed was mediated via reduced

social support, underscoring the clinically significant impact of partner absence on maternal psychological health

during times of collective trauma.

The findings reveal considerable psychological morbidity among perinatal women during wartime, with

elevated rates of both depressive and posttraumatic symptoms. Under these conditions, partner deployment was

associated with a markedly increased risk for probable depression among pregnant women, with prevalence

approaching 44%. This accords with findings of elevated antepartum depression among American women living

outside active war zones during their partner's deployment (Haas et al. 2005; Smith et al. 2010; Levine et al. 2015;

Tarney et al. 2015; Pretorius et al. 2024). A body of studies demonstrate the adverse impact of antepartum depression

on maternal, fetus, and infant health (Bauer et al. 2016; Jarde et al. 2016; Smith et al. 2020; Simonovich et al. 2021;

Ackerman-Banks et al. 2023; Meaidi 2024). Depression increases risk for preterm labor, low birth weight and

decreased head circumference (Engel et al. 2005; Ohlsson Arne 2011). It is linked with infant neurodevelopmental

risks (Field 2011; Rogers et al. 2020; Fan et al. 2024; Miller et al. 2024) potentially via fetal programming, in which

maternal distress contributes to the maternal-fetal biological milieu and, in turn, increases offspring's susceptibility

to disease states (Yehuda et al. 2005; Tegethoff et al. 2011; Yehuda and Lehrner 2018). Depression in pregnancy

strongly associates with postpartum symptoms and depression in subsequent pregnancies (Milgrom et al. 2008;

O'Hara and McCabe 2013).

To our knowledge, this study is the first to examine how partner deployment may undermine the formation

of maternal-infant attachment. This bond is derived from daily, positive mother-infant interactions that are

instrumental to the child's emotional and social development (Moehler et al. 2006; O'Dea et al. 2023). While various

mechanisms have been put forth for explaining the transmission of trauma across generations (Schore 2001; Yehuda

et al. 2005; Lev-Wiesel 2007; Yehuda and Lehrner 2018; Bhattacharya et al. 2019), maternal-infant attachment may

mediate the effect of maternal traumatic distress on the offspring's welfare (Schore 2001; Benoit et al. 2010; Enlow

et al. 2014; Ahlfs-Dunn et al. 2022). Stress and trauma may undermine maternal emotional regulation and maternal

sensitive care (Enlow et al. 2014; Kumar et al. 2020; Dekel et al. 2020). This study further suggests that partner

deployment may add to the impact of maternal trauma exposure on maternal-infant bonding.

The findings underscore the critical role of social support for maternal wellbeing during times of war. Partner

deployment may substantially diminish women's overall perceived social support, which in turn can adversely affect

maternal mental health and attachment with the infant. A body of research has documented the positive role of social

support in post-trauma psychological adaptation (Holt-Lunstad and Uchino 2015; Gariépy et al. 2016;

Wickramaratne et al. 2022; Calhoun et al. 2022). In particular, emotional and instrumental support from partners

has been identified as critical for maternal wellbeing in non-conflict contexts (Bilszta et al. 2008; Leahy-Warren et

al. 2012; Tanner Stapleton et al. 2012; Pilkington et al. 2015). These findings highlight the need for a more nuanced

understanding of the pathways through which partner deployment during a vulnerable period influences maternal

mental health, including direct effects via reduction of specific types of social support. Scalable interventions, such

as telehealth-based mental health support, online maternal support groups, virtual partner involvement and

community-based services, may provide alternative sources of support and help mitigate the adverse effects of partner absence (Kees and Rosenblum 2015; Nolan et al. 2019; Seagle et al. 2021; Hendrikx and Murphy 2021). Strengths of this study include the assessment of both pregnant and postpartum women exposed to twofold levels of adversity: from partner military deployment and from residing in war zones themselves. We also employed a match-controlled study design to understand the unique contribution of partner deployment to maternal outcomes. Shortcomings include the reliance on an internet-based, convenience sample. This approach enabled rapid recruitment of women under conditions of armed conflict, a population that would otherwise be difficult to enroll. Although the sample largely represents the Jewish Israeli population on demographics, there could be a selection bias, and the findings may reflect cultural-specific factors. We did not include clinical assessments; the self-reported mental health instruments accord strongly with diagnostic tools; however, there could be reporting bias. The use of a cross-sectional design precludes causal interpretations. Future longitudinal studies of perinatal women and their partners in various war zones are warranted to fully elucidate the impact of partner deployment on maternal mental health and their temporal relationships, as well as risk and protective factors.

In summary, among women with deployed partners, we find higher levels of probable depression during pregnancy, as well as lower maternal-infant attachment in the postpartum. Partner deployment adds a significant amount of stress by reducing social support and undermining maternal mental health during times of heightened stress, such as living under conditions of war. Because maternal wellbeing is a bedrock of a healthy society (Dekel et al. 2024), there is a critical need to ensure that resources for social support are in place and available. On a community level, attention to the welfare of perinatal women exposed to large-scale traumas is warranted.

Author Contribution

HAK contributed to project management activities and wrote parts of the manuscript.

SC created the online survey.

IHA conducted the statistical analysis and wrote the statistical approach and results.

CP contributed to the manuscript writing and editing.

IR and ES served as study consultants.

SD is the principal investigator of the larger project. She designed the conceptual study paradigm, collected the data, supervised the manuscript preparation, and wrote the manuscript.

All authors read and approved the final manuscript.

Funding Source. SD was supported by grants from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (R01HD108619; R21HD100817; and R21HD109546). The sponsor was not involved in study design; in the collection, analysis or interpretation of data; in the writing of the report; or in the decision to submit this article for publication. HAK is a recipient of a fellowship grant from the American Physicians Fellowship for Medicine in Israel.

Conflict of Interest. All authors have no competing interests.

Data availability. The datasets generated and/or analyzed during the current study are available from the corresponding author on reasonable request.

References

Ackerman-Banks CM, Lipkind HS, Palmsten K, et al (2023) Association of prenatal depression with new cardiovascular disease within 24 months postpartum. J Am Heart Assoc 12(9): https://doi.org/10.1161/JAHA.122.028133

Ahlfs-Dunn SM, Benoit D, Huth-Bocks AC (2022) Intergenerational transmission of trauma from mother to infant: the mediating role of disrupted prenatal maternal representations of the child. Attach Hum Dev 24:229–251. https://doi.org/10.1080/14616734.2021.1933769

Ahmed SH, Zakai A, Zahid M, et al (2024) Prevalence of post-traumatic stress disorder and depressive symptoms among civilians residing in armed conflict-affected regions: A systematic review and meta-analysis. Gen Psychiatr 37(3): https://doi.org/10.1136/gpsych-2023-101438

Alder J, Fink N, Bitzer J, et al (2007) Depression and anxiety during pregnancy: A risk factor for obstetric, fetal and neonatal outcome? A critical review of the literature. J Matern Fetal Neonatal Med 20:189–209. https://doi.org/10.1080/14767050701209560

American College of Obstetricians and Gynecologists (2023) Screening and diagnosis of mental health conditions during pregnancy and postpartum. Clinical Practice Guideline No. 4. Obstetrics & Gynecology 141:1232–1261. https://doi.org/10.1097/AOG.0000000000005200

Bauer A, Knapp M, Parsonage M (2016) Lifetime costs of perinatal anxiety and depression. J Affect Disord 192:83–90. https://doi.org/10.1016/j.jad.2015.12.005

Bedaso A, Adams J, Peng W, Sibbritt D (2021) The relationship between social support and mental health problems during pregnancy: a systematic review and meta-analysis. Reprod Health 18(162): https://doi.org/10.1186/s12978-021-01209-5

Benoit M, Bouthillier D, Moss E, et al (2010) Emotion regulation strategies as mediators of the association between level of attachment security and PTSD symptoms following trauma in adulthood. Anxiety Stress Coping 23:101–118. https://doi.org/10.1080/10615800802638279

Bhattacharya S, Fontaine A, MacCallum PE, et al (2019) Stress across generations: DNA methylation as a potential mechanism underlying intergenerational effects of stress in both post-traumatic stress disorder and pre-clinical predator stress rodent models. Front Behav Neurosci 13(113): https://doi.org/10.3389/fnbeh.2019.00113

Biaggi A, Conroy S, Pawlby S, Pariante CM (2016) Identifying the women at risk of antenatal anxiety and depression: A systematic review. J Affect Disord 191:62–77. https://doi.org/10.1016/j.jad.2015.11.014

Bilszta J, Tang M, Meyer D, et al (2008) Single motherhood versus poor partner relationship: Outcomes for antenatal mental health. Aust N Z J Psychiatry 42(1):56–65. https://doi.org/10.1080/00048670701732731

Bovin MJ, Marx BP, Weathers FW, et al (2016) Psychometric properties of the PTSD checklist for diagnostic and statistical manual of mental disorders-fifth edition (PCL-5) in veterans. Psychol Assess 28:1379–1391. https://doi.org/10.1037/pas0000254

Brown R (2020) The intergenerational impact of terror: Did the 9/11 tragedy impact the initial human capital of the next generation? Demography 57(4):1459–1481. https://doi.org/10.1007/s13524-020-00876-6

Buuren S van, Groothuis-Oudshoorn K (2011) MICE: Multivariate Imputation by Chained Equations in R. J Stat Softw 45(3): https://doi.org/10.18637/jss.v045.i03

Calhoun CD, Stone KJ, Cobb AR, et al (2022) The role of social support in coping with psychological trauma: An integrated biopsychosocial model for posttraumatic stress recovery. Psychiatric Quarterly 93(4):949–970. https://doi.org/10.1007/s11126-022-10003-w

Castro-Vale I, Severo M, Carvalho D, Mota-Cardoso R (2019) Intergenerational transmission of war-related trauma assessed 40 years after exposure. Ann Gen Psychiatry 18(14): https://doi.org/10.1186/s12991-019-0238-2

Chrisman AK, Dougherty JG (2014) Mass trauma: Disasters, terrorism, and war. Child Adolesc Psychiatric Clin N
Am 23(2):257–279. https://doi.org/10.1016/j.chc.2013.12.004
Cox JL, Holden JM, Sagovsky R (1987) Detection of postnatal depression: Development of the 10-item
Edinburgh Postnatal Depression Scale. Br J Psychiatry 150:782–786. https://doi.org/10.1192/bjp.150.6.782
Cranley MS (1981) Development of a tool for the measurement of maternal attachment during pregnancy. Nurs
Res 30:281–4. https://doi.org/PMID:6912989.
Curry SJ, Krist AH, Owens DK, et al (2019) Interventions to prevent perinatal depression: US preventive services
task force recommendation statement. JAMA 321(6):580–587
Dambi JM, Corten L, Chiwaridzo M, et al (2018) A systematic review of the psychometric properties of the cross-
cultural translations and adaptations of the Multidimensional Perceived Social Support Scale (MSPSS).
Health Qual Life Outcomes 16:80. https://doi.org/10.1186/s12955-018-0912-0
Dekel S, Ein-Dor T, Dishy GA, Mayopoulos PA (2020) Beyond postpartum depression: posttraumatic stress-
depressive response following childbirth. Arch Womens Ment Health 23:557–564.
https://doi.org/10.1007/s00737-019-01006-x
Dekel S, Papadakis JE, Quagliarini B, et al (2024) Preventing posttraumatic stress disorder following childbirth: a
systematic review and meta-analysis. Am J Obstet Gynecol 230:610-641.e14.
https://doi.org/10.1016/j.ajog.2023.12.013
Devakumar D, Birch M, Osrin D, et al (2014) The intergenerational effects of war on the health of children. BMC
Med 12(57): https://doi.org/10.1186/1741-7015-12-57
Druetz T, Browne L, Bicaba F, et al (2020) Effects of terrorist attacks on access to maternal healthcare services: A
national longitudinal study in Burkina Faso. BMJ Glob Health 5(9): https://doi.org/10.1136/bmjgh-2020-
002879
Engel SM, Berkowitz GS, Wolff MS, Yehuda R (2005) Psychological trauma associated with the World Trade
Center attacks and its effect on pregnancy outcome. Paediatr Perinat Epidemiol 19(5):334–341.
https://doi.org/10.1111/j.1365-3016.2005.00676.x
Enlow MB, Egeland B, Carlson E, et al (2014) Mother-infant attachment and the intergenerational transmission of
posttraumatic stress disorder. Dev Psychopathol 26:41–65. https://doi.org/10.1017/S0954579413000515
Fan X, Wu N, Tu Y, et al (2024) Perinatal depression and infant and toddler neurodevelopment: A systematic
review and meta-analysis. Neurosci Biobehav Rev 159:105579.
https://doi.org/10.1016/j.neubiorev.2024.105579
Field T (2011) Prenatal depression effects on early development: A review. Infant Behav Dev 34:1–14.
https://doi.org/10.1016/j.infbeh.2010.09.008
Gariépy G, Honkaniemi H, Quesnel-Vallée A (2016) Social support and protection from depression: Systematic
review of current findings in Western countries. British Journal of Psychiatry 209(4):284–293.
https://doi.org/10.1192/bjp.bp.115.169094
Gavin NI, Gaynes BN, Lohr KN, et al (2005) Perinatal depression. Obstet Gynecol 106(5):1071–1083.
https://doi.org/10.1097/01.AOG.0000183597.31630.db
Glaesmer H, Brähler E, Gündel H, Riedel-Heller SG (2011) The association of traumatic experiences and
posttraumatic stress disorder with physical morbidity in old age: A German population-based study.
Psychosom Med 73:401–406. https://doi.org/10.1097/PSY.0b013e31821b47e8
Godier-McBard LR, Ibbitson L, Hooks C, Fossey M (2019) Military spouses with deployed partners are at greater
risk of poor perinatal mental health: a scoping review. BMJ Mil Health 165:363–370.
https://doi.org/10.1136/jramc-2018-001069
Grote NK, Bridge JA, Gavin AR, et al (2010) A meta-analysis of depression during pregnancy and the risk of
preterm birth, low birth weight, and intrauterine growth restriction. Arch Gen Psychiatry 67:1012–1024.
https://doi.org/10.1001/archgenpsychiatry.2010.111

Haas DM, Pazdernik LA, Olsen CH (2005) A cross-sectional survey of the relationship between partner
deployment and stress in pregnancy during wartime. Women's Health Issues 15:48–54.
https://doi.org/10.1016/j.whi.2004.12.002
Hasson-Ohayon I, Horesh D (2024) A unique combination of horror and longing: Traumatic grief in post–October
7, 2023, Israel. J Trauma Stress 37:348–351
Hendrikx LJ, Murphy D (2021) Supporting the mental health needs of military partners through the together
webinar program: Pilot Randomized Controlled Trial. JMIR Ment Health 8(10):e25622.
https://doi.org/10.2196/25622
Ho DE, Imai K, King G, Stuart EA (2011) MatchIt: Nonparametric preprocessing for parametric causal inference.
J Stat Softw 42(8):1–28. https://doi.org/10.18637/jss.v042.i08
Holt-Lunstad J, Uchino BN (2015) Social support and health. In: Health behavior: Theory, research, and practice,
5th ed. Jossey-Bass/Wiley, Hoboken, NJ, US, pp 183–204
Hoppen TH, Priebe S, Vetter I, Morina N (2021) Global burden of post-traumatic stress disorder and major
depression in countries affected by war between 1989 and 2019: a systematic review and meta-analysis. BMJ
Glob Health 6:e006303. https://doi.org/10.1136/bmjgh-2021-006303
Houts Renate, Barnett-Walker KC, Paley B, Cox MJ (2008) Patterns of couple interaction during the transition to
parenthood. Pers Relatsh 15:103–122. https://doi.org/10.1111/j.1475-6811.2007.00187.x
Israel's National Emergency Portal (2024) Alerts History- Israel's National Emergency Portal | Iron Swords War.
https://alerts-history.oref.org.il/12481-he/Pakar.aspx. Accessed 21 Jul 2025
Jarde A, Morais M, Kingston D, et al (2016) Neonatal outcomes in women with untreated antenatal depression
compared with women without depression: A systematic review and meta-analysis. JAMA Psychiatry
73:826–837. https://doi.org/10.1001/jamapsychiatry.2016.0934
Jones I, Chandra PS, Dazzan P, Howard LM (2014) Bipolar disorder, affective psychosis, and schizophrenia in
pregnancy and the post-partum period. The Lancet 384:1789–1799. https://doi.org/10.1016/S0140-
6736(14)61278-2
Kees M, Rosenblum K (2015) Evaluation of a psychological health and resilience intervention for military
spouses: A pilot study. Psychol Serv 12:222–230. https://doi.org/10.1037/ser0000035
Klaman SL, Turner K (2016) Prevalence of perinatal depression in the military: A systematic review of the
literature. Matern Child Health J 20:52–65. https://doi.org/10.1007/s10995-016-2172-0
Klapper-Goldstein H, Pariente G, Wainstock T, et al (2024) The association of delivery during a war with the risk
for postpartum depression, anxiety and impaired maternal-infant bonding, a prospective cohort study. Arch
Gynecol Obstet 310:2863–2871. https://doi.org/10.1007/s00404-024-07715-8
Krupelnytska L, Morozova-Larina O (2025) Perinatal experiences of Ukrainian women at the beginning of the
war. J Reprod Infant Psychol 43:532–549. https://doi.org/10.1080/02646838.2023.2240827
Kumar SA, Franz MR, Brock RL, DiLillo D (2020) Posttraumatic stress and parenting behaviors: The mediating
role of emotion regulation. J Fam Violence 35:417–426. https://doi.org/10.1007/s10896-019-00124-5
Leahy-Warren P, McCarthy G, Corcoran P (2012) First-time mothers: social support, maternal parental self-
efficacy and postnatal depression. J Clin Nurs 21:388–397. https://doi.org/10.1111/j.1365-2702.2011.03701.x
Levany S, Shahar G, Greenberg D (2023) Calling for an immediate release of captive children in Gaza. The
Lancet 402:2070–2071. https://doi.org/10.1016/S0140-6736(23)02519-9
Levi-Belz Y, Groweiss Y, Blank C, Neria Y (2024) PTSD, depression, and anxiety after the October 7, 2023 attack
in Israel: a nationwide prospective study. EClinicalMedicine 68:102418.
https://doi.org/10.1016/j.eclinm.2023.102418
Levine JA, Bukowinski AT, Sevick CJ, et al (2015) Postpartum depression and timing of spousal military
deployment relative to pregnancy and delivery. Arch Gynecol Obstet 292:549–558.
https://doi.org/10.1007/s00404-015-3672-7

Lev-Wiesel R (2007) Intergenerational transmission of trauma across three generations: A preliminary study. Qualitative Social Work 6:75–94. https://doi.org/10.1177/1473325007074167

Lindgren K (2001) Relationships among maternal–fetal attachment, prenatal depression, and health practices in pregnancy. Res Nurs Health 24:203–217. https://doi.org/10.1002/nur.1023

Louise M. Howard HK (2020) Perinatal mental health: A review of progress and challenges. 19(3):313–327. https://doi.org/10.1002/wps.20769

Meaidi A (2024) Perinatal depression and incident maternal cardiovascular disease: a neglected association. Eur Heart J 45:2876–2877. https://doi.org/10.1093/eurheartj/ehae340

Milgrom J, Gemmill AW, Bilszta JL, et al (2008) Antenatal risk factors for postnatal depression: A large prospective study. J Affect Disord 108:147–157. https://doi.org/10.1016/j.jad.2007.10.014

Miller ES, Costantine MM, Mele L, et al (2024) The association between perinatal depressive symptoms and child neurodevelopment. Am J Obstet Gynecol MFM 6(11): https://doi.org/10.1016/j.ajogmf.2024.101488

Moehler E, Brunner R, Wiebel A, et al (2006) Maternal depressive symptoms in the postnatal period are associated with long-term impairment of mother-child bonding. Arch Womens Ment Health 9:273–278. https://doi.org/10.1007/s00737-006-0149-5

Morina N, Stam K, Pollet T V., Priebe S (2018) Prevalence of depression and posttraumatic stress disorder in adult civilian survivors of war who stay in war-afflicted regions. A systematic review and meta-analysis of epidemiological studies. J Affect Disord 239:328–338. https://doi.org/10.1016/j.jad.2018.07.027

Müller ME (1994) A questionnaire to measure mother-to-infant attachment. J Nurs Meas 2(2):129–141. https://doi.org/10.1891/1061-3749.2.2.129

Nemeroff CB, Goldschmidt-Clermont PJ (2012) Heartache and heartbreak—the link between depression and cardiovascular disease. Nat Rev Cardiol 9:526–539. https://doi.org/10.1038/nrcardio.2012.91

Nolan J, Lindeman S, Varghese FP (2019) Mobile app interventions for military and veteran families: Before, during, and after deployment. Psychol Serv 16:208–212. https://doi.org/10.1037/ser0000272

O'Dea GA, Youssef GJ, Hagg LJ, et al (2023) Associations between maternal psychological distress and mother-infant bonding: a systematic review and meta-analysis. Arch Womens Ment Health 26:441–452. https://doi.org/10.1007/s00737-023-01332-1

Office of the United Nations High Commissioner for Human Rights (2024) 7 October: UN experts call for end of violence and accountability after year of human loss and suffering and blatant disregard for international law

O'Hara MW, McCabe JE (2013) Postpartum depression: Current status and future directions. Annu Rev Clin Psychol 9(1):379–407. https://doi.org/10.1146/annurev-clinpsy-050212-185612

Ohlsson Arne SPS (2011) Effects of the September 11, 2001 disaster on pregnancy outcomes: a systematic review. Acta Obstet Gynecol Scand 90:6–18. https://doi.org/10.1111/j.1600-0412.2010.01020.x

Perkins JD, Wilkins SS, Kamran S, Shuaib A (2021) Post-traumatic stress disorder and its association with stroke and stroke risk factors: A literature review. Neurobiol Stress 14:100332. https://doi.org/10.1016/j.ynstr.2021.100332

Perlman SE, Friedman S, Galea S, et al (2011) Short-term and medium-term health effects of 9/11. The Lancet 378:925–934. https://doi.org/10.1016/S0140-6736(11)60967-7

Pilkington PD, Milne LC, Cairns KE, et al (2015) Modifiable partner factors associated with perinatal depression and anxiety: A systematic review and meta-analysis. J Affect Disord 178:165–180. https://doi.org/10.1016/j.jad.2015.02.023

Pretorius K, Sposato MF, Trueblood-Miller W (2024) Perinatal mental health and active-duty military spouses: a scoping review. BMC Pregnancy Childbirth 24(1): https://doi.org/10.1186/s12884-024-06727-1

Punamaki RL, Diab SY, Isosavi S, et al (2018) Maternal pre- and postnatal mental health and infant development in war conditions: The Gaza infant study. Psychol Trauma 10(2):144–153. https://doi.org/10.1037/tra0000275.supp

R Core Team (2023) R: A language and environment for statistical computing

Razurel C, Kaiser B, Sellenet C, Epiney M (2013) Relation Between Perceived Stress, Social Support, and Coping Strategies and Maternal Well-Being: A Review of the Literature. Women Health 53(1):74–99. https://doi.org/10.1080/03630242.2012.732681

Roberts AL, Agnew-Blais JC, Spiegelman D, et al (2015) Posttraumatic stress disorder and incidence of type 2 diabetes mellitus in a sample of women: A 22-year longitudinal study. JAMA Psychiatry 72:203–210. https://doi.org/10.1001/jamapsychiatry.2014.2632

Robrecht DT, Millegan J, Leventis LL, et al (2008) Spousal military deployment as a risk factor for postpartum depression. J Reprod Med 53:860–4

Rogers A, Obst S, Teague SJ, et al (2020) Association between maternal perinatal depression and anxiety and child and adolescent development: A meta-analysis. JAMA Pediatr 174(11):1082–1092. https://doi.org/10.1001/jamapediatrics.2020.2910

Rosenbaum S, Stubbs B, Ward PB, et al (2015) The prevalence and risk of metabolic syndrome and its components among people with posttraumatic stress disorder: a systematic review and meta-analysis. Metabolism 64:926–933. https://doi.org/10.1016/j.metabol.2015.04.009

Rosseel Y (2012) lavaan: An R package for structural equation modeling. J Stat Softw 48(2): https://doi.org/10.18637/jss.v048.i02

Schore AN (2001) Effects of a secure attachment relationship on right brain development, affect regulation, and infant mental health. Infant Ment Health J 22:7–66. https://doi.org/10.1002/1097-0355(200101/04)22:1<7::AID-IMHJ2>3.0.CO;2-N

Seagle ES, Xu J, Edwards N, McComb SA (2021) Social networking, social support, and well-being for the military spouse. The Journal for Nurse Practitioners 17(5):600–604. https://doi.org/10.1016/j.nurpra.2020.12.013

Secretary-General of the United Nations (2024) Message to mark one year since the attacks of 7 October 2023

Silver RC (2002) Nationwide longitudinal study of psychological responses to September 11. JAMA 288(10):1235. https://doi.org/10.1001/jama.288.10.1235

Simonovich SD, Nidey NL, Gavin AR, et al (2021) Meta-analysis of antenatal depression and adverse birth outcomes in US populations, 2010–20. Health Aff 40(10):1560–1565. https://doi.org/10.1377/hlthaff.2021.00801

Smith A, Twynstra J, Seabrook JA (2020) Antenatal depression and offspring health outcomes. Obstet Med 13:55–61. https://doi.org/10.1177/1753495X19843015

Smith DC, Munroe ML, Foglia LM, et al (2010) Effects of deployment on depression screening scores in pregnancy at an army military treatment facility. Obstetrics & Gynecology 116:679–684. https://doi.org/10.1097/AOG.0b013e3181eb6c84

Tanner Stapleton LR, Schetter CD, Westling E, et al (2012) Perceived partner support in pregnancy predicts lower maternal and infant distress. Journal of Family Psychology 26:453–463. https://doi.org/10.1037/a0028332

Tarney CM, Berry-Caban C, Jain RB, et al (2015) Association of spouse deployment on pregnancy outcomes in a U.S. military population. Obstetrics & Gynecology 126(3):569–574. https://doi.org/10.1097/AOG.0000000000001003

Tegethoff M, Greene N, Olsen J, et al (2011) Stress during pregnancy and offspring pediatric disease: A national cohort study. Environ Health Perspect 119:1647–1652. https://doi.org/10.1289/ehp.1003253

The Meir Amit Intelligence and Terrorism Information Center (2024) Operation Iron Swords. https://www.terrorism-info.org.il/en/operation-iron-swords-updated-to-1-p-m-february-21-2024/. Accessed 21 Jul 2025

U.S. Department of Defense Statement by Secretary of Defense Lloyd J. Austin III Marking one year since Hamas's October 7th terrorist assault on the state of Israel. In: Oct. 7, 2024

VanItallie TB (2002) Stress: A risk factor for serious illness. Metabolism 51:40–45.
https://doi.org/10.1053/meta.2002.33191
Weathers FW, LBT, KTM, PPA, MBP, & SPP (2013) The PTSD Checklist for DSM-5 (PCL-5). Scale available
from the National Center for PTSD at www.ptsd.va.gov.
Weinstein Y, Levav I, Gelkopf M, et al (2018) Association of maternal exposure to terror attacks during pregnancy
and the risk of schizophrenia in the offspring: A population-based study. Schizophr Res 199:163–167.
https://doi.org/10.1016/j.schres.2018.04.024
Wickramaratne PJ, Yangchen T, Lepow L, et al (2022) Social connectedness as a determinant of mental health: A
scoping review. PLoS One 17:e0275004. https://doi.org/10.1371/journal.pone.0275004
Wisner KL, Murphy C, Thomas MM (2024) Prioritizing maternal mental health in addressing morbidity and
mortality. JAMA Psychiatry 81(5):521–526. https://doi.org/10.1001/jamapsychiatry.2023.5648
Wittkowski A, Vatter S, Muhinyi A, et al (2020) Measuring bonding or attachment in the parent-infant-
relationship: A systematic review of parent-report assessment measures, their psychometric properties and
clinical utility. Clin Psychol Rev 82:101906. https://doi.org/10.1016/j.cpr.2020.101906
Yehuda R, Engel SM, Brand SR, et al (2005) Transgenerational effects of posttraumatic stress disorder in babies of
mothers exposed to the World Trade Center attacks during pregnancy. Journal of Clinical Endocrinology and
Metabolism 90:4115–4118. https://doi.org/10.1210/jc.2005-0550
Yehuda R, Hyman SE (2005) The impact of terrorism on brain, and behavior: What we know and what we need to
know. Neuropsychopharmacology 30:1773–1780. https://doi.org/10.1038/sj.npp.1300817
Yehuda R, Lehrner A (2018) Intergenerational transmission of trauma effects: putative role of epigenetic
mechanisms. World Psychiatry 17:243–257. https://doi.org/10.1002/wps.20568
575
576

Table 1. Socio-demographics, trauma exposure, and obstetrical information for pregnant and postpartum women by partner deployment status

	Pregnant Women (N = 165)		Postpartum Women (N = 264)	
	Active Partner Deployment	Non-Active Partner Deployment	Active Partner Deployment	Non-Active Partner Deployment
	n (%) / M (SD)	n (%) / M (SD)	n (%) / M (SD)	n (%) / M (SD)
Demographic Factors				
Maternal Age	31.86 ± 4.06	31.31 ± 4.43	31.79 ± 4.55	32.22 ± 4.05
Mental Illness before October 7th	9 (8.41)	2 (3.45)	10 (6.99)	14 (11.57)
Marital Status				
Married or domestic partnership	106 (99.07)	57 (98.28)	142 (99.3)	121(100.00)
Single, in a relationship	1 (0.93)	1 (1.72)	1 (0.70)	0
Employment				
Employed	75 (70.09)	45 (78.95)	97 (68.31)	91 (75.83)
Home-employed	10 (9.35)	4 (7.02)	10 (7.04)	7 (5.83)
Self-employed	10 (9.35)	4 (7.02)	15 (10.56)	13 (10.83)
Student	11 (10.28)	4 (7.02)	12 (8.45)	5 (4.17)
Unemployed	1 (0.93)	0	8 (5.63)	4 (3.33)
Education				
High School or equivalent	5 (4.67)	3 (5.17)	5 (3.50)	2 (1.65)
Some college	5 (4.67)	1 (1.72)	7 (4.90)	8 (6.61)
Bachelor's Degree	57 (53.27)	36 (62.07)	78 (54.55)	56 (46.28)
Master's Degree	34 (31.76)	15 (25.86)	43 (30.07)	46 (38.02)

Doctoral or Medical Degree	6 (5.61)	3 (5.17)	10 (6.99)	9 (7.44)
Household Income (Monthly)				
Up to 7106 NIS	7 (6.54)	4 (6.90)	7 (4.90)	8 (6.61)
Up to 12862 NIS	20 (18.69)	11 (18.97)	36 (25.17)	29 (23.97)
Up to 22847 NIS	46 (42.99)	26 (44.83)	57 (39.86)	33 (27.27)
Over 22847 NIS	34 (31.78)	17 (29.31)	43 (30.07)	51 (42.25)
Trauma Exposure Factors				
Home Zones				
Red	9 (8.41)	5 (8.62)	15 (10.49)	9 (7.44)
Being Displaced	2 (22.22)	1 (20.00)	4 (26.67)	1 (11.11)
Orange	27 (25.23)	13 (22.41)	35 (24.48)	32 (26.45)
Yellow	30 (28.04)	10 (17.24)	35 (24.48)	30 (24.79)
Green	41 (38.32)	30 (51.72)	58 (40.56)	50 (41.32)
Watch uncensored videos (past week)	7 (6.54)	0	12 (8.39)	12 (9.92)
Know civilians murdered	42 (39.25)	20 (34.48)	61 (42.66)	39 (32.23)
Know civilians and/or soldiers taken hostage	16 (14.95)	7 (12.07)	25 (17.48)	21 (17.36)
Obstetrics Factors				
Primiparity	25 (23.36)	17 (29.31)	36 (25.17)	43 (35.54)
Gestational age				
1st Trimester	23 (21.49)	5 (8.62)	-	-
2nd Trimester	39 (36.45)	25 (43.10)	-	-
3rd Trimester	45 (42.06)	28 (48.28)	-	-
Birth after October 7th	-	-	86 (60.14)	65 (53.72)
Mode of Delivery				

Vaginal delivery	-	-	106 (79.70)	85 (71.43)
Assisted vaginal delivery	-	-	11 (8.27)	8 (6.72)
Scheduled caesarean delivery	-	-	6 (4.51)	11 (9.24)
Emergent caesarean delivery	-	-	10 (7.52)	15 (12.61)
NICU admittance	-	-	9 (6.82)	8 (6.72)

Note. NIS, New Israeli Shekel. All exposure factors related to the October 7th events and the war. Home Zone refers to geographic zones as defined by the Israeli Defense Forces (IDF) Home Front Command to indicate the available time to reach shelter during missile attacks: Red (≤15 seconds), Orange (30 seconds), Yellow (45 seconds), and Green (60-90 seconds). Watched uncensored videos in the past week; Values pertain to high frequency of exposure (score 8 or above on a 1-10 scale regarding amount (hours) of exposure daily). Gestational age was reported according to trimesters (weeks) (0-13, first trimester; 14-27, second trimester; 28-42, third trimester). NICU = Neonatal Intensive Care Unit.

Table 2. Maternal depression, mother-infant bonding, and social support by partner deployment status

	Pregnant Women (N = 165)					Postpartum Women (N = 264)					OR / Effect Size (95% CI)
	Active Partner Deployment		Non-Active Partner Deployment		χ^2 / Z	Active Partner Deployment		Non-Active Partner Deployment		χ^2 / Z	
	%/M	n/SD	%/M	n/SD		%/M	n/SD	%/M	n/SD		
Probable Depression	43.96	40	25.49	13	4.01*	23.89	27	23.08	24	0	2.29 (1.10 – 5.00)[1]
											1.05 (0.56 – 1.97)[2]
Probable PTSD	20.24	17	19.15	9	1.8e^{-30}	21	21	20	18	0	1.07 (0.44 – 2.73)[1]
											1.06 (0.52 – 2.17)[2]
Social Support	45.76	9.60	50.84	6.11	3.16**	47.53	9.65	52.12	6.98	3.07**	0.25 (0.10 – 0.38)[1]
											0.19 (0.07 – 0.30)[2]
Social Support (Significant Other)	15.73	3.36	17.86	1.93	3.97***	16.51	3.29	18.30	2.37	3.90***	0.31 (0.16 – 0.43)[1]
											0.24 (0.12 – 0.35)[2]
	15.16	3.96	16.82	2.50	2.25*	15.48	4.04	17.03	3.36	2.62**	0.17 (0.02 – 0.30)[1]

	M	SD	M	SD	t	M	SD	M	SD	t	OR (CI)
Social Support (Family)											$0.16 (0.04 - 0.28)^2$
Social Support (Friends)	14.86	3.86	16.16	3.13	2.02*	15.54	3.59	16.79	3.13	2.22*	$0.16 (0.02 - 0.30)^1$ $0.14 (0.02 - 0.26)^2$
Maternal Attachment	61.65	14.00	63.02	15.12	0.61	32.44	7.28	30.27	5.50	2.36*	$0.05 (0.00 - 0.21)^1$ $0.15 (0.03 - 0.26)^2$

Note. Probable depression was defined based on a score of 14 or higher on the Edinburgh Postnatal Depression Scale (EPDS $\geq$ 14). Probable posttraumatic stress disorder (PTSD) regarding October 7th was defined based on a score of 33 or higher on the Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5 $\geq$ 33). Maternal attachment refers to bonding with the fetus (higher scores better bonding) and bonding with the infant (higher scores worse attachment). M = Mean. SD = Standard deviation. OR = odd ratios, CI = confidence interval.

[1] = Active partner deployment v/s non-active partner deployment for pregnant women

[2] = Active partner deployment v/s non-active partner deployment for postpartum women

***p<0.001, **p<0.01, *p<0.05

Figure 1. **Associations between partner deployment, social support and antepartum depression.** Mediation analysis linking study group (partner actively deployed v/s not) to symptoms of depression during pregnancy via perceived social support. Solid lines represent significant paths; Gray dashed lines represent non-significant paths. Values are standardized scores. ***p<0.001

Figure 2. **Associations between partner deployment, social support and mother-infant attachment problems**. Mediation analysis linking study group (partner actively deployed v/s not) to maternal-infant attachment during the postpartum period via perceived social support. Solid lines represent significant or near significance paths. Values are standardized scores. ***p<0.001, +<0.10