Functional Heterogeneity in Non-Suicidal Self-Injury across Psychiatric Disorders: Neural and Psychosocial Correlates

Mingzhu Li, MMed[1,2 †], Yang Xiao, MSc[1,2 †], Yuqi Ge, MMed[1,2], Huiru Yan, MSn[5], Xueni Li, MD [1,2], Weihua Yue, MD[1,2,3,4], Hao Yan, PhD[1,2*]

[1]Institute of Mental Health, Peking University Sixth Hospital, Beijing 100191, China

[2]NHC Key Laboratory of Mental Health (Peking University), National Clinical Research Center for Mental Disorders (Peking University Sixth Hospital), Beijing 100191, China

[3]PKU-IDG/McGovern Institute for Brain Research, Peking University, Beijing, 100871, China

[4]Research Unit of Diagnosis and Treatment of Mood Cognitive Disorder (2018RU006), Chinese Academy of Medical Sciences, Beijing 100191, China

[5]Peking University School of Nursing, Beijing 100191, China.

*Correspondence: Hao Yan

Email: hao_y@bjmu.edu.cn

Address: 51 Huayuanbei Road, Haidian District, Beijing 100191, China.

Tel: 010-82805307

† These authors contributed equally to this work.

Abstract

Non-suicidal self-injury (NSSI) is a common behavior among adolescents,

particularly within psychiatric populations. While neurobiological and psychosocial

risk factors have been extensively studied, the mechanisms underlying NSSI's

heterogeneity remain unclear. This study investigated 304 hospitalized

adolescents/young adults (16–25 years) with NSSI and comorbid psychiatric

diagnoses (major depressive disorder [MDD], bipolar disorder [BD], eating disorders

[ED]) using psychological assessments and resting-state fMRI data from 163

participants. Orthogonal projection non-negative matrix factorization of Ottawa Self-

Injury Inventory responses identified two core motivational dimensions: self-related

(emotional regulation) and social-related (interpersonal influence). The self-related

factor correlated with amygdala-centered cortico-limbic emotional regulation

networks and predominated in affective disorders (MDD/BD), while the social-related

factor linked to frontoparietal cognitive control and frontotemporal social cognition

networks, particularly in ED. Fuzzy C-means clustering revealed three NSSI

functional subtypes, independent of diagnostic categories : self-subtype primarily

driven by self-related functions, social-subtype influenced by both self- and social-

related functions with greater exposure to psychosocial risks, and non-specific

subtype with undifferentiated motivations. No subtype was exclusively driven by

social-related functions. The "self-social" dual-dimensional neural model

demonstrated distinct subtype-specific profiles in functional connectivity,

psychosocial risk exposure, and clinical features. Self-related mechanisms primarily

engaged emotional regulation circuits, whereas social-related mechanisms emphasize

the role of psychosocial risk factors and cognitive-emotional circuits. These findings

provide neural evidence for the functional heterogeneity of NSSI and highlight the

need for personalized interventions. Treatments targeting emotion regulation may

benefit all subtypes, individuals with prominent social-related motivations may

additionally require interventions aimed at improving interpersonal functioning.

Introduction

Non-suicidal self-injury (NSSI), defined as the deliberate self-inflicted damage to

body tissues without suicidal intent (e.g., cutting, burning, or hitting) [1], is a prevalent

behavior among adolescents and young adults, particularly in psychiatric populations

[2]. Lifetime prevalent rates of NSSI range from 13.4–17.2% in community samples [3] to

as high as 67% in psychiatric inpatient settings [4]. NSSI frequently co-occurs with

major depressive disorders (MDD), bipolar disorders (BD), and eating disorders (ED)

[5], underscoring its transdiagnostic nature. Although not intended to be suicidal, NSSI

is associated with severe adverse outcomes, including increased suicide risk [6] and

impaired social functioning [7], underscoring its significant public health impact.

NSSI is a complex behavior shaped by the interplay of social, psychological, and

biological factors [8]. Psychosocial risk variables, such as childhood abuse, adverse life

events, limited family and social support, depression, anxiety and deficits in emotion

regulation are well-documented contributors to NSSI [9–11]. These variables can be

broadly categorized into interpersonal and intrapersonal domains, both of which play

critical roles in the onset and maintenance of NSSI [12]. Although advances in

psychosocial research have successfully informed emotion regulation-focused

psychological interventions [13], the lack of mechanistic elucidation at the biological

level continues to hinder treatment innovation. To date, no targeted somatic treatments,

such as psychopharmacology or neuromodulation, are available for NSSI [14],

highlighting the urgent need to elucidate its neurobiological mechanisms [15].

From a neurodevelopmental perspective, distal risk factors (e.g., childhood trauma) during critical periods of neural maturation can influence proximal biological alterations, including structural and functional changes in key brain regions [16]. Neuroimaging studies investigating the neural correlates of NSSI have identified abnormalities in the frontal-limbic system, encompassing the amygdala, anterior cingulate cortex (ACC), medial prefrontal cortex (mPFC), dorsolateral prefrontal cortex (dlPFC), hippocampus, and insula [17,18]. These regions are integral to emotion expression and regulation [19], self-referential processing/interpersonal interactions [20], and reward processing [21]. For instance, hyperactivity in the amygdala and insula has been linked to heightened emotional reactivity and salience detection [19], while dysregulation in the mPFC and ACC is associated with impaired self-regulation and social cognition [20]. These findings suggest that NSSI may arise from dysregulated bottom-up and top-down neural circuits involved in emotion and cognitive control [17].

NSSI is a highly heterogeneous behavior, varying in frequency, methods, severity, and underlying motivations [22]. The functional model of NSSI, which distinguishes between intrapersonal (e.g., emotion regulation) and interpersonal (e.g., emotion expression and social influence) functions, provides a valuable framework for understanding this heterogeneity [23,24]. Studies indicate that 66–81% of individuals engage in NSSI primarily for intrapersonal functions, while 33–56% report interpersonal motivations [25]. These functional dimensions are associated with differing clinical outcomes, with intrapersonal functions linked to higher suicide risk

[26,27]. Moreover, inter- and intrapersonal functions appear to be differentially associated

with abovementioned psychosocial factors and neurobiological mechanisms,

respectively.

In this study, we aim to bridge the gap between self-injurious behaviors and their

underlying neurobiological and psychosocial correlates. First, we identify

conceptually meaningful and compact factor structures for NSSI functions in

hospitalized adolescents with psychiatric disorders and derived functional subtypes.

Next, we examined the neural correlates of different factors of NSSI functions. Finally,

we explored the factor related neural characteristics and psychosocial risk variables

associated with the different functional subtypes. By integrating functional, neural,

and psychosocial dimensions, this study seeks to advance our understanding of NSSI

mechanisms and inform the development of targeted psychological and

neuromodulation interventions.

Methods

2.1 Participants

The present study included inpatients at Peking University Sixth Hospital (Beijing,

Chian) from July 2021 to September 2022. A total of 304 patients were enrolled and

completed the full set of questionnaires. Among them, 163 patients voluntarily

underwent magnetic resonance imaging (MRI) data acquisition. This study was

approved by the Ethics Committee of Peking University Sixth Hospital.

Inclusion criteria were: (1) aged 16-25 years, of any gender; (2) inpatient

diagnosis meeting the International Classification of Diseases, 10th Revision (ICD-10)

criteria for depressive episode or recurrent depressive disorder, bipolar disorder, and

eating disorders, as assessed by two attending or senior psychiatrists; (3) at least one

episode of NSSI; (4) provision of written informed consent by the patient or their

guardian.

Exclusion criteria were: (1) individuals exhibiting only suicidal behavior without

NSSI; (2) those with mental developmental delay; (3) individuals with severe,

unstable physical illnesses; (4) history of epilepsy or febrile seizures; (5) pregnant or

lactating women, or those planning pregnancy.

Among the participants, 167 were diagnosed with major depressive disorders, 89

with bipolar disorders, and 48 with eating disorders. This study included patients with

varying severities of NSSI behaviors: 178 participants had repeated NSSI behaviors

(NSSI behaviors occurred for at least 5 days in the past year, meeting the DSM-5

diagnostic criteria for NSSI disorder), while 126 patients had occasional NSSI

behaviors (NSSI behaviors occurred on at least one day but fewer than 5 days). Given

that the focus of this study is on the functions of NSSI behaviors, the functional

classifications were applied to both frequent and occasional NSSI behaviors to

maintain consistency across varying severities.

2.2 Measurements and Assessments

Basic demographic characteristics, medical history, comorbidities, and clinical

information regarding current psychiatric disorders were collected using a self-

designed Case Report Form (CRF). Suicidal ideation and suicidal behavior were also

recorded in the CRF based on documentation in the patients' hospitalization records

regarding the presence of suicidal ideation and suicidal behavior prior to the current

admission.

The Ottawa Self-Injury Inventory (OSI) was used to assess the frequency and

functions of NSSI behaviors among participants. This scale has demonstrated good

reliability and validity in Chinese adolescent samples [28]. The study specifically

focused on the NSSI functions assessed by the 29 items of the OSI function subscale

(OSI-F) [28], which captures various motivations and intentions underlying NSSI

behaviors. Responses were rated from 0 (never) to 4 (always), with higher scores

indicating stronger functional tendencies.

The Self-Rating Anxiety Scale (SAS) [29] and the Self-Rating Depression Scale

(SDS) [29] were used to evaluate anxiety and depressive symptoms, respectively.

Additionally, several other scales were utilized to assess individual characteristics and

environmental risk factors, including the NEO Five-Factor Inventory (NEOFFI) [30],

the revised Chinese Internet Addiction Scale (CIAS-R) [31], the Adolescent Self-Rating

Life Events Check List (ASLEC) [29], the Childhood Trauma Questionnaire (CTQ) [32],

the Family Assessment Device (FAD) [29], and the Social Support Rate Scale (SSRS) [29].

Suicidal ideation and suicide attempt within the past year were also recorded.

2.3 Factor Analysis of NSSI Functions

Orthonormal projective non-negative factorization (OPNMF) [33,34] was employed to

identify latent factors underlying the NSSI functional items from the OSI-F. OPNMF

is a variant of NMF, which is widely used in recent biomedical studies[35]. This method

introduces two constraints: a projective constraint, which ensures stable and

deterministic item-to-factor assignments, and an orthonormality constraint, which

promotes compact and non-redundant factor definitions. Here, the OPNMF achieved

factorization for given data (OSI-F) by yielding a probabilistic parcellation to assign

latent factor loadings for each item of NSSI functions (a dictionary matrix containing

factors), and these loadings quantified the extent of which each item belongs to a

factor. The advantages of OPNMF enabled us to acquire a sparse, stable and

interpretable structure of latent factor of NSSI functions.

To determine the optimal number of factors, we employed a set of sophisticated

evaluation strategies based on cross-validation with 10,000 split-half analyses.

Specifically, the entire sample used in the current study was divided into two halves,

and the OPNMF was conducted separately on each half to derive the dictionary

matrices. We subsequently calculated the congruency between item-to-factor

assignments to examine the stability of the factor solutions at various evaluation

strategies (describe in the Supplement), including the adjusted Rand index (RI),

variation of information (VI), and concordance index (CI) between the dictionaries [35].

Additionally, we also assessed the generalizability of the factor model by using the

reconstruction error (RE) calculating the absolute differences between the projection

of the dictionary from one split-sample data on the other split-sample [35]. The internal

consistency of the optimal OPNMF model was assessed using the Pearson's

correlations with an additional bootstrapping validation procedure in which we

conducted 10,000 times resampling and re-calculate the correlation coefficient. The

between-items relationships under optimal factor structure were also calculated using

the Pearson's correlation.

2.4 Clustering of NSSI Functional Subtypes

Factor loadings from the optimal OPNMF model were used to cluster patients into

different NSSI functional subtypes. Considering the extent of differentiation of

vulnerability factors related to disorder or symptoms exhibited a certain of

heterogeneity in different patients, we adopted a soft clustering approach (Fuzzy C-

means) to allow probabilistic assignment to multiple subtypes. This analytic

combination has been infrequently explored in NSSI populations.

Fuzzy C-means clustering is a widely used soft clustering algorithm that estimates the

membership likelihood of each individual to each cluster, rather than forcing

exclusive assignment. This method allows us to identify those patients with an explicit

inclination to specific NSSI functions and ambiguous subgroups that have not yet

been differentiated into specific categories, based on a cutoff threshold.

In this study, the fuzzy C-means clustering [36] was implemented to NSSI factor

loadings of all patients to encode the membership likelihood of each patient in any

given subtyping clusters, after adjusting for age, gender, years of education, suicidal

ideation, and suicide attempts. The fuzzy silhouette index (SI), Xie and Beni index

(XB), and partition entropy (PE) were calculated at range of 2 to 6 as internal validity

indices to determine the optimal number of clusters (details in Supplement). Higher SI

and lower XB and PE indicated better clustering quality. To choose the cutoff to

define the optimal membership discrimination, the elbow method was used at range of

70% to 90% by identifying the elbow point, where the rate of decrease in explained

variance slows significantly. The elbow point represented a balance between

maximizing cluster compactness and minimizing the number of clusters.

2.5 Resting-State Functional Connectivity (rsFC) Network Construction

Resting-state functional MRI data were acquired from 163 participants using the

Siemens Prisma 3.0 Tesla MRI scanner. Data preprocessing was conducted using the

DPABI (Data Processing Assistant for Resting State fMRI Advanced Edition) [37]

software based on the MATLAB (The MathWorks, Natick, MA, USA). Standard

processing procedures were followed. The scanning parameters and detailed

information for each preprocessing step are available in the Supplement.

This study quantified resting-state functional connectivity (rsFC) based on the

spontaneous fluctuations of resting-state bold oxygenation level dependent (BOLD)

signals, generating a whole-brain rsFC matrix for each participant. We referred to a

previous brain parcellation scheme comprising a total of 210 cortical regions [38] and 14

subcortical regions [39]. The average time series signals of each brain region were

extracted, and the Pearson's correlation coefficients were computed between the time

series signals of each pair of brain regions to determine functional connectivity

strength. The Fischer's *r-to-z* transformation was utilized to convert the correlation

coefficients into normalized *z*-scores. A 224×224 functional connectivity matrix was

constructed for each participant.

2.6 Modeling Neural Correlates of NSSI Functions Using rsFC

To better understand the neural correlates of NSSI functions, we trained a machine-

learning algorithm based on regularized canonical correlation (RCC) with a hyper-

parameterized feature selection space following a previous study [40] (details in

Supplement).

Briefly, RCC based on L2 penalty (λ) was used to predict each factor loadings of

NSSI functions for all patients. Out-of-sample predictions were obtained using a

nested cross-validation scheme using a 90% random selection strategy (10-fold),

repeated 100 times. During the training of the model in each fold, we ran a random

hyperparameter search consisting of 10 iterations in a 10-fold cross-validation using

the grid search and chose the best model for prediction. The optimized number of

rsFC features was defined as the top-ranked features based on a bootstrapping scheme

with 95% subsampling and 100 replications. The hyperparameter space was defined

as follows: N_{features} ranged from 100 to 400 top-ranked rsFC features, and λ ranged

from 1 to 10. We evaluated the performance of each model on the entire predicted set

using canonical correlation coefficients. Statistical significance of the coefficients was

evaluated using a non-parametric permutation test with 10,000 iterations.

The correlations between the FC scores and each functional factor were using the

Pearson correlation analysis. The significant threshold for all statistical analysis was

set as $p < 0.001$ adjusted by multiple comparison correction using false discovery rate

(FDR) method.

2.7 Statistical Analyses

Pearson's correlation analysis was used to examine the associations between factors

loadings and SAS/SDS scores. One-way analysis of variance (ANOVA) with a post-

hoc Turkey test was employed to analyze differences in scale scores (e.g., CTQ,

ALSE) across different subtypes or diagnostic groups (MDD, BD and ED). For

ordinal and categorical variables, such as NSSI frequency and gender, chi-square tests

were conducted to assess group differences. The statistical analyses above were

conducted using SPSS, version 27 (IBM, Armonk, NY). OPNMF and fuzzy C-means

clustering were performed using MATLAB (version 2012a, MathWorks, Natick, MA,

USA). All analyses controlled for age, gender, years of education, suicidal ideation,

and suicide attempts. A $p < 0.05$ was considered the threshold for statistical

significance, multiple comparisons were corrected by FDR method.

Results

3.1 Demographic and Clinical Characteristics

A total of 304 participants were included and completed the full assessment, with 163

of them also undergoing MRI for subsequent analysis of neural correlates. Statistical

comparisons indicated that the MRI subset did not significantly differ from the overall

sample in terms of age, gender distribution, level of depression and anxiety,

personality traits, and various assessments for the environmental risk factors (Table 1).

Despite a high proportion of participants in both groups reporting suicidal ideation

and attempts, no significant differences were observed (Table 1). Regarding the

severity of NSSI, no significant differences were found between the two samples in

terms of NSSI frequency, age at first episode, or functional scores (Table 1and Table

S1).

------------------------------- Insert Table 1 here -------------------------------

3.2 Two Factors of NSSI Functions

A robust two-factor model was obtained using OPNMF based on 304 patients (Figure

2A). The first factor comprised 14 NSSI functional items in OSI-F, it mainly related to

self-perception and emotional regulation and was named as "self-related factor". The

second factor comprised 15 items mainly related to interpersonal functioning and was

named as "social-related factor". Model evaluation metrics indicated that a two-factor

solution provided the highest stability (with the highest adjusted RI and CI values, and

the lowest VI value, Figure S1A) and the strongest generalizability (with the lowest

out-of-sample increase in RE value, Figure S1A). These results were validated

through 1,000 times split-half cross-validation.

The items within the two NSSI functional factors decomposed by OPNMF

exhibited higher correlations internally (Figure 2B). A strong negative correlation was

observed between the two factors ($r = -0.766$, Figure S1B). Correlation analysis

(Figure 2C) revealed influences of depressive and anxious emotions on the two

functional factor-loadings. Although SDS and SAS scores were highly correlated among participants ($r = 0.776$, $p < 0.001$), the two factors effectively captured distinct internal characteristics of the patients. Specifically, the self-related factor-loading showed a significant positive correlation with SDS scores ($r = 0.167$, $p = 0.004$), but no significant correlation with SAS scores ($r = 0.075$, $p = 0.194$). However, the social-related factor loading was significantly correlated with SAS scores ($r = 0.229$, $p < 0.001$) but not with SDS scores ($r = 0.075$, $p = 0.191$). Further analysis based on disease diagnosis revealed that the self-related factor-loading was higher in mood disorders (MDD = BD > ED), whereas the social-related factor showed higher loadings in eating disorders (ED > MDD = BD, Figure 2C).

-------------------------------- Insert Figure 2 here --------------------------------

3.3 Subtypes of NSSI Function

Fuzzy C-means clustering applied to the two-factor loadings revealed the optimal solution was to divide patients into two clusters (Figure S2). The core members of each cluster were selected based on their likelihoods derived from the fuzzy C-means algorithm, with a cutoff threshold of 0.80. Consequently, two core subtypes of NSSI functions were defined, with 52 ambiguous patients located near the intersection of the clusters (Figure 3A). 意味这目前不以任何功能为主导，定义为…

The largest group included 154 patients (50.7%) and was characterized by higher loadings on the self-related factor and lower loadings on the social-related factor, leading to its designation as the "self-subtype" (Figure 3A). The "social-subtype"

comprised 98 patients (32.2%) and was characterized by higher loadings on the

social-related factor and lower loadings on the self-related factor. A total of 52

ambiguous patients (17.1%), dispersed between the other two subtypes, were

categorized as "non-specific subtype". No significant differences were found in

gender, age, or diagnostic distribution across the three subtypes. Additionally, the

subtypes showed similar profiles regarding suicidal ideation, suicide attempts, age of

onset, and NSSI frequency/ severity (Table 2 and Table S2).

Regarding NSSI function, social-subtype exhibited a significantly higher total

function score compared to the other two subtypes (Cohen's d = 0.97, $p < 0.001$).

Interestingly, in contrast to the factor loadings, social-subtype showed higher scores

on both self-related and social-related OSI-F items (Table 2), with multiple functions

being reported (Cohen's d = 0.96, $p < 0.001$). These differences were not significant

between self-subtype and non-specific subtype.

Levels of anxiety and depression were significantly different across three

subtypes (Figure 3B and Table 2; SAS: Cohen's d = 0.84, $p = 0.008$; SDS: Cohen's d

= 0.78, $p = 0.049$). Specifically, SAS scores in social-subtype were significantly

higher than those in non-specific subtype (Cohen's d = 0.54, $p = 0.02$); SDS scores in

self-subtype and social-subtype were significantly higher compared to non-specific

subtype (self vs non-specific: Cohen's d = 0.37, $p = 0.024$, social vs non-specific:

Cohen's d = 0.39, $p = 0.023$).

Regarding potential NSSI risk factors, no significant differences were found in

the total scores for the FAD, CTQ, and SSRS, which assess family functioning,

childhood trauma, and social support, respectively, among the three subtypes.

Differences were primarily observed in ASLEC, NEOFFI, and CIAS (Figure 3D and

Table 2). Specifically, social-subtype reported significantly higher scores on several

dimensions of the ASLEC (Cohen's d = 0.95, $p < 0.001$), including "catch hell,"

"loss," "interpersonal pressure," "learning pressure," and "healthy adaptation" ($p <$

0.05, FDR corrected), suggested a higher level of exposure to life events. Similarly,

social-subtype exhibited significantly higher levels of internet addiction, as measured

by the CIAS (Cohen's d = 0.85, $p = 0.006$). In terms of personality traits assessed with

NEOFFI, social-subtype scored significantly higher in neuroticism compared to the

other subtypes (Cohen's d = 0.87, $p = 0.002$) and significantly lower in agreeableness

(Cohen's d = 0.90, $p < 0.001$).

------------------------------ Insert Figure 3 and Table 2 here ------------------------------

3.4 rsFC Features Related to Functional Factors and Subtypes

We conducted RCCA to investigate the rsFC features that best related to the two-

factors of NSSI function using MRI data from 163 participants. To minimize

overfitting, we performed a subsampling and cross-validated procedure (see

Supplement). This procedure identified two dimensions of rsFC features related to

NSSI functions.

The first dimension was significantly positively correlated with loading of self-

related factor (Figure 4A, $r = 0.656$, $p = 0.005$), while the second dimension was

associated with the loading of social-related factor ($r = 0.602$, $p = 0.012$). We then

measured the regional contributions of each dimension (Figure 4A). The top five

representative brain regions for the first dimension included the right amygdala

(13.5%), left amygdala (6.5%), left caudal cuneus gyrus (4.0%), left ventromedial

parietooccipital sulcus (2.6%) and right putamen (2.6%). For the social-related

dimension, the top representative brain regions are in the right dorsolateral frontal

cortex (Brodmann area, [BA] 6, 5.4%), left ventrolateral frontal cortex (BA8, 4.6%),

left medial parietal area (BA7, 4.4%), left dorsal frontal area (BA9/46, 3.9%) and

right amygdala (3.4%).

We then examined the rsFC correlates of each functional factors and the

differences between subtypes. The self-related functions were associated with

extensive cortico-subcortical functional connectivity (Figure S6). Specifically, the 23

significant connections (surviving a threshold of $p < 0.001$) were observed between

the amygdala and various cortical regions, including the superior frontal gyrus (SFG),

precentral gyrus, superior temporal gyrus (STG), parahippocampal gyrus, postcentral

gyrus, insular gyrus, and occipital lobe. Additionally, rsFC between the putamen and

the occipital lobe, as well as subcortical connections between the putamen and the

amygdala, were observed (Figure 4A, Table S6). Regarding these self-related

connections, self-subtype showed a significantly higher rsFC compared to social-

subtype and non-specific subtype, with no differences between the latter two subtypes

(Figure 4B).

In contrast, the social-related factor demonstrated a more localized set of rsFC,

with six significant connections identified. These connections involved rsFC between

the fronto-parietal and fronto-temporal network, as well as connections between the

inferior temporal gyrus and hippocampus, dorsolateral superior frontal gyrus and

insular gyrus (Figure 4A, Table S6). Social-subtype exhibited elevated rsFC compared

to self-subtype, while no significant differences were observed between social-

subtype/self-subtype and non-specific subtype (Figure 4B).

------------------------------ Insert Figure 4 here ------------------------------

3.5 Sensitivity Analysis

We conducted an OPNMF on the revised OSI-F (with 7 items deleted) [41] for

sensitivity analysis. The results indicated that both the self-related and social-related

factors comprised 11 items each, confirming a stable and generalizable two-factor

structure (Figure S3A-B). The model maintained good inter-factor independence

(Figure S3C, $r = -0.715$, $p < 0.001$) and high internal factor consistency ($r = 0.968$, p

< 0.001). Further imaging verification found that the two factors were still

significantly correlated with the previously obtained rsFC characteristics (Figure S3D;

self-related: $r = 0.689$, $p < 0.001$; social-related: $r = 0.774$, $p < 0.001$). Additionally,

whether for patients with repeated NSSI behaviors or those with occasional NSSI

behaviors, the two-factor structure was found to be optimal. Despite the presence of

some potentially confusing individual items, the model consistently upheld strong

inter-factor independence and internal factor consistency in both groups (Figures S4-

S5). The correlation with rsFC characteristics were also reliably replicated across

these patient groups (Figures S4-S5).

Discussion

This study identified two conceptually meaningful and compact factors underlying the

motivations and reasons of NSSI behaviors: self-related functions, associated with

emotional regulation and self-perception, and social-related functions, linked to

interpersonal influence and social interactions. These findings align with the

previously proposed intrapersonal and interpersonal functional models of NSSI [24,42],

providing further empirical support for the heterogeneity of NSSI behaviors. The self-

related factor emphasizes on reinforcement from the self, including affect regulation,

self-punishment, anti-dissociation, anti-suicide, and marking distress. In contrast, the

social-related factor focuses on reinforcement from others or the environment,

involving interpersonal influence, establishing boundaries and seeking peer bonding.

Importantly, the identified subtypes revealed that self-related and social-related

functions are not mutually exclusive but instead represent overlapping motivations,

with varying emphases across individuals. While both subtypes exhibited self-related

motivations, the social-subtype also demonstrated significant interpersonal concerns,

suggesting that self-related motivations are the dominant driver of NSSI behaviors.

Psychosocial factors further differentiate these subtypes. Individuals in the

social-subtype reported greater exposure to adverse life events, higher tendency

toward problematic internet use, higher neuroticism and lower agreeableness. These

findings align with studies showing that social distress can exacerbate emotional

dysregulation, reinforcing both self- and social-related motivations for NSSI [43,44].

Additionally, higher neuroticism was linked to emotional dysregulation and

vulnerability to mental health disorders [45], while lower agreeableness is associated

with difficulties in interpersonal relationships [46]. These findings suggest offer a

streamlined, diagnosis-independent intervention framework for NSSI：emotion

regulation strategies may benefit all NSSI subtypes, while those in the social-subtype

may additionally require to improve interpersonal effectiveness. Such tailored

approaches could address the distinct psychosocial profiles associated with each

subtype, potentially enhancing treatment outcomes.

Although the diagnostic distribution of mental disorders did not show significant

differences across three subtypes, we observed a notable pattern in the factor loadings.

Specifically, the self-related factor loading was higher in mood disorders (MDD and

BD), while the social-related factor loading was higher in eating disorders. This

suggests that patients with eating disorders may be more likely to engage in NSSI for

social-related reasons, such as managing interpersonal relationships or coping with

social pressures [47,48]. Given the relatively small sample size of eating disorder patients

in this study, it is important to replicate these findings in larger cohorts to further

explore the potential distinctiveness of social-related NSSI motivations in this

population.

Neuroimaging analyses revealed distinct neural correlates for self-related and

social-related functions. Key findings revealed the amygdala as the central hub for

self-related motivations, consistent with its well-documented roles in emotion

processing and regulation [49,50]. Three observed amygdala-centered network (frontal,

temporal and insula lobe) constitutes a core substrate for self-related functions. The

fronto-amygdala circuitry forms a top-down neural circuit crucial for emotion

regulation [49,51,52], where enhanced connectivity may paradoxically increase

rumination and worry [53,54]. The temporal-amygdala connectivity supports emotional

memory [55–57] and processing of affectively charged information [58], potentially

underlying self-punitive and distress-marking functions. Connections with temporal

and insular lobes are involved in self-referential processing [59] and may facilitate the

transmission of negative emotions, contributing to the emotional dysregulation

characteristic of NSSI populations[60,61]. The insula is central to interoception,

emotional self-awareness, decision-making, and cognitive control [62–64]. It has been

linked to attention modulation and pain perception in NSSI patients [65], as well as

deficits in interoceptive awareness and suicidal ideation [66]. These networks are

integral to emotion regulation, self-referential processing, and interoceptive awareness,

supporting the emotional and self-punitive motivations underlying self-subtype of

NSSI.

In contrast, social-related functions were linked to fronto-parietal, fronto-insular ,

and fronto-temporal networks, which are involved in social cognition, interpersonal

behavior, and the regulation of social-emotional information. The dorsal fronto-

parietal network supports top-down attentional orienting processes [67,68] and is

recruited during the relational integration of social information [69]. The medial

prefrontal cortex and precuneus (the medial portion of BA7) within this network play

pivotal roles in adaptive interpersonal behavior [70] and self-related mental

representations [71,72]. The insula exhibits consistent activation under social exclusion [73].

The fronto-insula hyperconnectivity appears linked to maladaptive introspection [74]

and negative emotional information [75]. Concurrently, the fronto-temporal network also

participates in the "social brain", and within this network, fusiform gyrus has been

implicated in socioemotional processes [76] interpersonal influence [71].These neural

circuits reflect the higher rejection sensitivity combined with ineffective attempts to

regulate feelings of rejection in individuals with NSSI, underscoring their difficulties

in emotion regulation within social contexts. Moreover, these findings support the

potential translational utility of neuromodulation strategies targeting subtype-specific

neural circuits in adolescents with NSSI.

The application of fuzzy clustering allowed us to capture the heterogeneity of

NSSI behaviors more effectively [35]. The identification of an undifferentiated subtype

(non-specific subtype), characterized by lower scores on both self- and social-related

motivations, suggests the existence of individuals who have yet to develop a dominant

function for NSSI. This subgroup highlights the potential for shifts in motivational

bases over time, warranting further longitudinal investigation to track changes in

NSSI functions and their associated neurobiological profiles.

Despite these advances, several limitations should be acknowledged. Firstly, the

cross-sectional design limits our ability to infer causality or assess the temporal

stability of the identified functional subtypes. Longitudinal studies are needed to

explore the evolution of NSSI motivations and their neural correlates over time,

especially for non-specific subtype. Secondly, these results need to be replicated and

validated using independent samples. Finally, the relatively small sample size of

certain subgroups, such as individuals with eating disorders, may limit the

generalizability of our results. Meanwhile, as a cross-culture behavior, future studies

with larger, culturally diverse cohorts are needed to further explore the distinctiveness

of social-related NSSI expression and motivations.

In conclusion, this study bridges the gap between behavioral functions and

neurobiological phenotypes in NSSI, an area that has received limited attention in

prior research. By identifying distinct NSSI subtypes with corresponding neural and

psychosocial profiles, our findings provide a foundation for developing more precise

and effective interventions. Tailoring treatments to the specific functional and

neurobiological characteristics of individuals with NSSI may enhance therapeutic

outcomes and ultimately reduce the burden of this debilitating behavior.

Acknowledgements

We extend our gratitude to all participants in this study. This study was supported by

the National Natural Science Foundation of China (82071505, 81771443,

81361120395). HY was partly supported by Clinical Medicine Plus X – Young

Scholars Project, Peking University the Fundamental Research Funds for the Central

Universities (PKU2024LCXQ046) and Capital's Funds for Health Improvement and

Research (2024-2-4115). We also acknowledge the development of a preliminary

MATLAB-based toolkit for classifying NSSI subtypes, available at

https://github.com/XiaoYangLab/NSSI_factor .

Declaration of Interest

The authors report on biomedical financial interests or potential conflicts of interest.

Reference

Nock MK. Self-Injury. *Annu Rev Clin Psychol* 2010; 6: 339–363.

Singhal A, Ross J, Seminog O, Hawton K, Goldacre MJ. Risk of self-harm and suicide in people with specific psychiatric and physical disorders: comparisons between disorders using English national record linkage. *J R Soc Med* 2014; 107: 194–204.

Swannell SV, Martin GE, Page A, Hasking P, St John NJ. Prevalence of Nonsuicidal Self-Injury in Nonclinical Samples: Systematic Review, Meta-Analysis and Meta-Regression. *Suicide Life Threat Behav* 2014; 44: 273–303.

Wolff J, Frazier EA, Esposito-Smythers C, Burke T, Sloan E, Spirito A. Cognitive and Social Factors Associated with NSSI and Suicide Attempts in Psychiatrically Hospitalized Adolescents. *Journal of Abnormal Child Psychology* 2013; 41: 1005–1013.

Wang L, Liu J, Yang Y, Zou H. Prevalence and risk factors for non-suicidal self-injury among patients with depression or bipolar disorder in China. *BMC Psychiatry* 2021; 21: 389.

Ribeiro JD, Franklin JC, Fox KR, Bentley KH, Kleiman EM, Chang BP *et al.* Self-injurious thoughts and behaviors as risk factors for future suicide ideation, attempts, and death: a meta-analysis of longitudinal studies. *Psychol Med* 2016; 46: 225–236.

Tan AC, Rehfuss MC, Suarez EC, Parks-Savage A. Nonsuicidal self-injury in an adolescent population in Singapore. *Clin Child Psychol Psychiatry* 2014; 19: 58–76.

Qu D, Wen X, Liu B, Zhang X, He Y, Chen D *et al.* Non-suicidal self-injury in Chinese population: a scoping review of prevalence, method, risk factors and preventive interventions. *The Lancet Regional Health–Western Pacific* 2023; 37.

Qu D, Wen X, Liu B, Zhang X, He Y, Chen D *et al.* Non-suicidal self-injury in Chinese population: a scoping review of prevalence, method, risk factors and preventive interventions. *Lancet Reg Health West Pac* 2023; 37: 100794.

Wang Y-J, Li X, Ng CH, Xu D-W, Hu S, Yuan T-F. Risk factors for non-suicidal self-injury (NSSI) in adolescents: A meta-analysis. *EClinicalMedicine* 2022; 46: 101350.

Wolff JC, Thompson E, Thomas SA, Nesi J, Bettis AH, Ransford B *et al.* Emotion dysregulation and non-suicidal self-injury: A systematic review and meta-analysis. *Eur Psychiatry* 2019; 59: 25–36.

Tatnell R, Kelada L, Hasking P, Martin G. Longitudinal Analysis of Adolescent NSSI: The Role of Intrapersonal and Interpersonal Factors. *Journal of Abnormal Child Psychology* 2014; 42: 885–896.

Calvo N, García-González S, Perez-Galbarro C, Regales-Peco C, Lugo-Marin J, Ramos-Quiroga J-A *et al.* Psychotherapeutic interventions specifically developed for NSSI in adolescence: A systematic review. *Eur Neuropsychopharmacol* 2022; 58: 86–98.

Jiang Z, Wang Z, Diao Q, Chen J, Tian G, Cheng X *et al.* Efficacy of interventions for suicide and self-injury in children and adolescents: a meta-analysis. *Sci Rep* 2023; 12: 12313.

Kaess M, Hooley JM, Klimes-Dougan B, Koenig J, Plener PL, Reichl C *et al.* Advancing a temporal framework for understanding the biology of nonsuicidal self-injury: An expert review. *Neuroscience & Biobehavioral Reviews* 2021; 130: 228–239.

Whittle S, Zhang L, Rakesh D. Environmental and neurodevelopmental contributors to youth mental illness. *Neuropsychopharmacology* 2024; 50: 201–210.

Auerbach RP, Pagliaccio D, Allison GO, Alqueza KL, Alonso MF. Neural Correlates Associated With Suicide and Nonsuicidal Self-injury in Youth. *Biological Psychiatry* 2021; 89: 119–133.

Brañas MJAA, Croci MS, Ravagnani Salto AB, Doretto VF, Martinho E, Macedo M *et al.* Neuroimaging Studies of Nonsuicidal Self-Injury in Youth: A Systematic Review. *Life* 2021; 11: 729.

Morawetz C, Riedel MC, Salo T, Berboth S, Eickhoff SB, Laird AR *et al.* Multiple large-scale neural networks underlying emotion regulation. *Neuroscience & Biobehavioral Reviews* 2020; 116: 382–395.

Ho TC, Walker JC, Teresi GI, Kulla A, Kirshenbaum JS, Gifuni AJ *et al.* Default mode and salience network alterations in suicidal and non-suicidal self-injurious thoughts and behaviors in adolescents with depression. *Transl Psychiatry* 2021; 11: 38.

Liu X, Hairston J, Schrier M, Fan J. Common and distinct networks underlying reward valence and processing stages: a meta-analysis of functional neuroimaging studies. *Neurosci Biobehav Rev* 2011; 35: 1219–1236.

Wang Z, Li D, Chen Y, Tao Z, Jiang L, He X *et al.* Understanding the subtypes of non-suicidal self-injury: A new conceptual framework based on a systematic review. *Psychiatry Research* 2024; 334: 115816.

Nock MK, Prinstein MJ, Nock MK, Prinstein MJ. A Functional Approach to the

Assessment of Self-Mutilative Behavior. *Journal of Consulting and Clinical Psychology* 2004; 72: 885–890.

Klonsky ED, Glenn CR, Styer DM, Olino TM, Washburn JJ. The functions of nonsuicidal self-injury: converging evidence for a two-factor structure. *Child Adolesc Psychiatry Ment Health* 2015; 9: 44.

Taylor PJ, Jomar K, Dhingra K, Forrester R, Shahmalak U, Dickson JM. A meta-analysis of the prevalence of different functions of non-suicidal self-injury. *Journal of Affective Disorders* 2018; 227: 759–769.

Nock MK, Prinstein MJ. Contextual features and behavioral functions of self-mutilation among adolescents. *Journal of abnormal psychology* 2005; 114: 140.

Park Y, Qu W, Ammerman BA. Characteristics and Functions of Non-Suicidal Self-Injury That Inform Suicide Risk. *Archives of Suicide Research* 2024; : 1–14.

[CHENG Wen-hong ZF et al, XIAO Ze-ping. Study on reliability and validity of Chinese version of Ottawa self-injury inventory. *Journal of Shanghai Jiao Tong University (Medical Science)* 2015; 35: 460-.

Xiang Dong Wang, Wang XL, Ma H. *Rating scales for mental health*. Beijing: Chinese Mental Health Journal Press, 1999.

Huang XT. Handbook of personality: theory and research. *Shanghai: East China Normal University Press, doi* 2003; 10: 2093310.

Bai Y, Fan F. A Study on the Internet Dependence of College Students: the Revising and Applying of a Measurement. *Psychological Development and Education* 2005; 21: 99–104.

Zhao X, Zhang Y, Li L, Zhou Y. Evaluation on reliability and validity of Chinese version of childhood trauma questionnaire. *Chinese Journal of Tissue Engineering Research* 2005; : 209–211.

Sotiras A, Resnick SM, Davatzikos C. Finding imaging patterns of structural covariance via Non-Negative Matrix Factorization. *NeuroImage* 2015; 108: 1–16.

Zhirong Yang, Oja E. Linear and Nonlinear Projective Nonnegative Matrix Factorization. *IEEE Trans Neural Netw* 2010; 21: 734–749.

Chen J, Patil KR, Weis S, Sim K, Nickl-Jockschat T, Zhou J *et al.* Neurobiological Divergence of the Positive and Negative Schizophrenia Subtypes Identified on a New Factor Structure of Psychopathology Using Non-negative Factorization: An International Machine Learning Study. *Biological Psychiatry* 2020; 87: 282–293.

Bezdek JC. *Pattern Recognition With Fuzzy Objective Function Algorithms*. Pattern Recognition with Fuzzy Objective Function Algorithms, 1981.

Yan CG, Wang XD, Zuo XN, Zang YF. DPABI: Data Processing & Analysis for (Resting-State) Brain Imaging. *Neuroinformatics* 2016; 14: 339–351.

Fan L, Li H, Zhuo J, Zhang Y, Wang J, Chen L *et al.* The Human Brainnetome Atlas: A New Brain Atlas Based on Connectional Architecture. *Cereb Cortex* 2016; 26: 3508–3526.

Desikan RS, Ségonne F, Fischl B, Quinn BT, Dickerson BC, Blacker D *et al.* An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage* 2006; 31: 968–980.

Buch AM, Vértes PE, Seidlitz J, Kim SH, Grosenick L, Liston C. Molecular and network-level mechanisms explaining individual differences in autism spectrum disorder. *Nat Neurosci* 2023; 26: 650–663.

Chen H, Pan B, Zhang C, Guo Y, Zhou J, Wang X. Revision of the non-suicidal self-injury behavior scale for adolescents with mental disorder. *Zhong Nan Da Xue Xue Bao Yi Xue Ban* 2022; 47: 301–308.

Klonsky ED, Glenn CR. Assessing the Functions of Non-suicidal Self-injury: Psychometric Properties of the Inventory of Statements About Self-injury (ISAS). *J Psychopathol Behav Assess* 2009; 31: 215–219.

Tang J, Ma Y, Lewis SP, Chen R, Clifford A, Ammerman BA *et al.* Association of Internet Addiction With Nonsuicidal Self-injury Among Adolescents in China. *JAMA Network Open* 2020; 3: e206863–e206863.

Yen J-Y, Yen C-F, Chen C-S, Wang P-W, Chang Y-H, Ko C-H. Social anxiety in online and real-life interaction and their associated factors. *Cyberpsychol Behav Soc Netw* 2012; 15: 7–12.

Ormel J, Jeronimus BF, Kotov R, Riese H, Bos EH, Hankin B *et al.* Neuroticism and common mental disorders: meaning and utility of a complex relationship. *Clin Psychol Rev* 2013; 33: 686–697.

Graziano WG, Jensen-Campbell LA, Hair EC. Perceiving interpersonal conflict and reacting to it: the case for agreeableness. *J Pers Soc Psychol* 1996; 70: 820–835.

Muehlenkamp JJ, Takakuni S, Brausch AM, Peyerl N. Behavioral functions underlying NSSI and eating disorder behaviors. *J Clin Psychol* 2019; 75: 1219–1232.

Kiekens G, Claes L. Non-Suicidal Self-Injury and Eating Disordered Behaviors: An Update on What We Do and Do Not Know. *Curr Psychiatry Rep* 2020; 22: 68.

Morawetz C, Riedel MC, Salo T, Berboth S, Eickhoff SB, Laird AR *et al.* Multiple large-scale neural networks underlying emotion regulation. *Neuroscience & Biobehavioral Reviews* 2020; 116: 382–395.

Ochsner KN, Silvers JA, Buhle JT. Functional imaging studies of emotion regulation: a synthetic review and evolving model of the cognitive control of emotion. *Ann N Y Acad Sci* 2012; 1251: E1-24.

Ochsner KN, Bunge SA, Gross JJ, Gabrieli JD. Rethinking feelings: an FMRI study of the cognitive regulation of emotion. *Journal of cognitive neuroscience* 2002; 14: 1215–1229.

Refojo D, Schweizer M, Kuehne C, Ehrenberg S, Thoeringer C, Vogl AM *et al.* Glutamatergic and dopaminergic neurons mediate anxiogenic and anxiolytic effects of CRHR1. *Science* 2011; 333: 1903–1907.

Murphy ER, Barch DM, Pagliaccio D, Luby JL, Belden AC. Functional connectivity of the amygdala and subgenual cingulate during cognitive reappraisal of emotions in children with MDD history is associated with rumination. *Dev Cogn Neurosci* 2016; 18: 89–100.

Sokołowski A, Kowalski J, Dragan M. Neural functional connectivity during rumination in individuals with adverse childhood experiences. *Eur J Psychotraumatol* 2022; 13: 2057700.

LaBar KS, Cabeza R. Cognitive neuroscience of emotional memory. *Nat Rev Neurosci* 2006; 7: 54–64.

Smith APR, Stephan KE, Rugg MD, Dolan RJ. Task and content modulate amygdala-hippocampal connectivity in emotional retrieval. *Neuron* 2006; 49: 631–638.

Kark SM, Slotnick SD, Kensinger EA. Repetition Enhancement of Amygdala and Visual Cortex Functional Connectivity Reflects Nonconscious Memory for Negative Visual Stimuli. *J Cogn Neurosci* 2016; 28: 1933–1946.

LeDoux JE. Emotion Circuits in the Brain. .

van der Meer L, Costafreda S, Aleman A, David AS. Self-reflection and the brain: a theoretical review and meta-analysis of neuroimaging studies with implications for schizophrenia. *Neurosci Biobehav Rev* 2010; 34: 935–946.

Nam G, Moon H, Lee J-H, Hur J-W. Self-referential processing in individuals with nonsuicidal self-injury: An fMRI study. *Neuroimage Clin* 2022; 35: 103058.

Liao K, Yu R, Chen Y, Chen X, Wu X, Huang X *et al.* Alterations of regional

brain activity and corresponding brain circuits in drug-naïve adolescents with nonsuicidal self-injury. *Sci Rep* 2024; 14: 24997.

Farb NAS, Segal ZV, Anderson AK. Attentional modulation of primary interoceptive and exteroceptive cortices. *Cereb Cortex* 2013; 23: 114–126.

Craig ADB. How do you feel--now? The anterior insula and human awareness. *Nat Rev Neurosci* 2009; 10: 59–70.

Critchley HD. Neural mechanisms of autonomic, affective, and cognitive integration. *J Comp Neurol* 2005; 493: 154–166.

Krause-Utz A, Frost R, Winter D, Elzinga BM. Dissociation and Alterations in Brain Function and Structure: Implications for Borderline Personality Disorder. *Curr Psychiatry Rep* 2017; 19: 1–22.

Ho TC, Walker JC, Teresi GI, Kulla A, Kirshenbaum JS, Gifuni AJ *et al.* Default mode and salience network alterations in suicidal and non-suicidal self-injurious thoughts and behaviors in adolescents with depression. *Transl Psychiatry* 2021; 11: 38.

Petersen SE, Posner MI. The attention system of the human brain: 20 years after. *Annu Rev Neurosci* 2012; 35: 73–89.

Corbetta M, Shulman GL. Control of goal-directed and stimulus-driven attention in the brain. *Nat Rev Neurosci* 2002; 3: 201–215.

Magis-Weinberg L, Blakemore S-J, Dumontheil I. Social and Nonsocial Relational Reasoning in Adolescence and Adulthood. *J Cogn Neurosci* 2017; 29: 1739–1754.

Bickart KC, Dickerson BC, Feldman Barrett L. The amygdala as a hub in brain networks that support social life. *Neuropsychologia* 2014; 63: 235–248.

Quevedo K, Martin J, Scott H, Smyda G, Pfeifer JH. The neurobiology of self-knowledge in depressed and self-injurious youth. *Psychiatry Res Neuroimaging* 2016; 254: 145–155.

Cavanna AE, Trimble MR. The precuneus: a review of its functional anatomy and behavioural correlates. *Brain* 2006; 129: 564–583.

Cacioppo S, Frum C, Asp E, Weiss RM, Lewis JW, Cacioppo JT. A Quantitative Meta-Analysis of Functional Imaging Studies of Social Rejection. *Scientific Reports* 2013; 3: 2027.

Kaiser RH, Whitfield-Gabrieli S, Dillon DG, Goer F, Beltzer M, Minkel J *et al.* Dynamic Resting-State Functional Connectivity in Major Depression.

Neuropsychopharmacology 2016; 41: 1822–1830.

75 Kaiser RH, Snyder HR, Goer F, Clegg R, Ironside M, Pizzagalli DA. Attention Bias in Rumination and Depression: Cognitive Mechanisms and Brain Networks. *Clin Psychol Sci* 2018; 6: 765–782.

76 Suarez GL, Burt SA, Gard AM, Burton J, Clark DA, Klump KL *et al.* The impact of neighborhood disadvantage on amygdala reactivity: Pathways through neighborhood social processes. *Dev Cogn Neurosci* 2022; 54: 101061.

Tables and Figure legends

Table1. Demographic and clinical characteristics

R/O, Repeated/Occasional; Y/N, Yes/No; M/F, Male/Female; MDD, major depressive disorder; BD, bipolar disorder; ED, eating disorders; SAS, Self-Rating Anxiety Scale; SDS, Self-Rating Depression Scale; ASLEC, Adolescent Self-Rating Life Events Checklist; FAD, Family Assessment Device; CTQ, Childhood Trauma Questionnaire; SSRS, Social Support Rating Scale; CIAS, Chinese Internet Addiction Scale; NEOFFI, NEO Five-Factor Inventory.
[a] The χ^2 was obtained by a chi-square test.

Table2. Characteristics and functions of NSSI between three functional subtypes

	Self-subtype (N = 154, 50.7%)	Social-subtype (N = 98, 32.2%)	Non-specific subtype (N = 52, 17.1%)	F/χ^2	p
Sex (F/M)	31/123	18/80	12/40	0.471[a]	0.790
Age (year)	18.70 ± 2.46	18.36 ± 2.25	18.69 ± 2.49	0.004	0.996
Diagnosis (MDD/BD/ED)	88/47/19	53/26/19	26/16/10	3.109[a]	0.540
Suicidal Ideation (Y/N)	122/32	78/20	44/8	0.755[a]	0.685
Suicidal Attempt (Y/N)	90/64	62/36	27/25	1.83[a]	0.400
NSSI Frequency (R/O)	92/62	57/41	29/23	0.262[a]	0.877
Age of onset	15.31 ± 3.33	15.30 ± 3.39	14.48 ± 2.36	1.329	0.266
Self-related factor scores	29.58 ± 11.73	29.22 ± 12.32	23.75 ± 15.06	4.614	0.010*
Self-related factor loading	0.07 ± 0.02	0.01 ± 0.01	0.03 ± 0.02	329.324	<0.001***
Social-related factor scores	7.88 ± 5.82	23.40 ± 9.58	13.40 ± 7.44	207.829	<0.001***
Social-related factor loading	0.01 ± 0.01	0.08 ± 0.03	0.04 ± 0.02	557.919	<0.001***
Functional scores at onset	37.51 ± 16.26	52.92 ± 20.88	37.25 ± 21.72	35.553	<0.001***
SAS	59.81 ± 14.70	63.43 ± 14.78	55.75 ± 12.99	4.952	0.008**
SDS	71.27 ± 13.41	71.68 ± 13.93	66.40 ± 12.59	3.047	0.049*
ASLEC	46.28 ± 20.02	61.51 ± 23.55	44.09 ± 21.28	18.227	<0.001***
FAD	148.60 ± 23.23	147.89 ± 22.19	145.09 ± 19.18	0.486	0.616
CTQ	47.92 ± 15.13	49.84 ± 14.22	46.19 ± 14.59	1.112	0.330
SSRS	28.84 ± 6.44	29.84 ± 6.77	31.33 ± 6.52	2.886	0.057
CIAS	45.64 ± 14.03	41.50 ± 9.05	40.71 ± 10.15	5.212	0.006**
NEOFFI					
Neuroticism	36.81 ± 6.12	39.04 ± 5.91	35.63 ± 6.55	6.355	0.002**
Extraversion	21.44 ± 6.69	22.27 ± 6.99	23.26 ± 6.27	1.517	0.221
Openness to Experience	33.56 ± 6.91	34.64 ± 6.01	32.87 ± 4.96	1.524	0.219

Agreeableness	33.52 ± 5.34	30.81 ± 5.00	32.73 ± 4.47	8.532	<0.001[***]
Conscientiousness	29.44 ± 7.29	28.99 ± 7.18	29.02 ± 7.22	0.138	0.871

Note: R/O, Repeated/Occasional; Y/N, Yes/No; M/F, Male/Female; MDD, major depressive disorder; BD, bipolar disorder; ED, eating disorders; SAS, Self-Rating Anxiety Scale; SDS, Self-Rating Depression Scale; ASLEC, Adolescent Self-Rating Life Events Checklist; FAD, Family Assessment Device; CTQ, Childhood Trauma Questionnaire; SSRS, Social Support Rating Scale; CIAS, Chinese Internet Addiction Scale; NEOFFI, NEO Five-Factor Inventory.
*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.
[a] The χ^2 was obtained by a chi-square test.

FIGURE 1

Figure 1. Neurobiological vulnerabilities and psychosocial risks in psychiatric disorders with NSSI. A sample of 304 inpatients diagnosed with major depressive disorder (MDD: N = 167), bipolar disorder (BD: N = 89), and eating disorders (ED: N = 48) reported the functions underlying their NSSI behavior. The self-reported functions were analyzed using orthogonal projection non-negative matrix factorization (OPNMF), which identified two primary functional factors: social-related and self-related functions. Functional connectivity (FC) features and corresponding subtypes were determined based on these factors. Subsequent analyses examined differences in FC characteristics and associated risk factors across the subtypes, offering valuable insights into the neurobiological vulnerabilities and psychosocial risks linked to NSSI behavior. MDD, major depressive disorder; BD, bipolar disorder; ED, eating disorders; OPNMF, orthogonal projection non-negative matrix factorization; OSI-F, functional items in the Ottawa Self-Injury Inventory.

FIGURE 2

Figure 2. Self–Social dimensional model of NSSI functions. **(A)** The most stable and generalizable two-factor structure of NSSI functions identified by OPNMF (N = 304), consisting of a self-related factor (14 items) and a social-related factor (15 items) derived from the 29 OSI-F items. **(B)** A heatmap illustrating the correlations between and within the two derived factors. The strength of correlations is represented by color: red indicates positive correlations, blue indicates negative correlations, with darker shades denoting stronger correlations. **(C)** Scatter plots display Pearson's correlations between levels of anxiety and depression and the two factor loadings, adjusted for age, sex, education, SI, and SA. **(D)** One-way ANOVA results comparing the two factor loadings across the three diagnostic groups, controlling for covariates. For each box plot, the box illustrates the Standard Error of the Mean (SEM, centered on the mean), whiskers denote the 5% and 95% values, and the horizontal line signifies the median. SAS, Self-Rating Anxiety Scale; SDS, Self-Rating Depression Scale; MDD, major depressive disorder; BD, bipolar disorder; ED, eating disorders; $*p < 0.05$; $**p < 0.01$; $***p < 0.001$.

FIGURE 3

Figure 3. Three NSSI subtypes identified via clustering based on the self–social motivational structure. **(A)** the scatter plot presents a two-dimensional visualization of subtype clustering results. Red points represent social-subtype (social-related subtype, N = 98), gray points represent self-subtype (self-related subtype, N = 154), and blue points represent non-specific subtype (N = 52). **(B)** Violin plots depict the comparison results of the two-factor loadings, depression, and anxiety levels across the three subtypes. **(C)** The bubble chart shows a comparison of risk factors for NSSI among the three subtypes, incorporating six questionnaires: the NEO Five-Factor Inventory (NEOFFI), the revised Chinese Internet Addiction Scale (CIAS), the Adolescent Self-Rating Life Events Check List (ASLEC), the Childhood Trauma Questionnaire (CTQ), the Family Assessment Device (FAD), and the Social Support Rate Scale (SSRS). The horizontal dotted line denotes the significance threshold ($p < 0.05$, FDR corrected), with larger bubbles indicating more significant differences between the groups. **(D)** Risk factors differences between three functional subtypes (post-hoc Turkey test, FDR corrected). Violin plots represent means (dot) and 95% confidence intervals (line). *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$.

FIGURE 4

Figure 4. Functional connectivity features correlating with functional factors and subtypes. **(A)** RCC revealed two dimensions related to two functional factors (top: self-related factor; bottom: social-related factor). Scatterplots illustrate the association between connectivity scores and the loading of functional factor scores for each RCC dimension across participants. The schematic diagram of the brain depicts the spatial distribution of regional proportions for each dimension. For each dimension linked to one functional factor, the rsFCs significantly contributing to it were extracted, and the regional proportions were calculated, with darker colors indicating greater contributions. The circular diagram on the far right illustrates the rsFCs significantly associated with the raw scores of the two factors ($p < 0.001$, FDR corrected). Blue lines indicate negative correlations, while red lines represent positive correlations. Node colors differentiate the macroscopic brain regions: purple represents the frontal lobe, blue-purple the temporal lobe, yellow the parietal lobe, orange the insular lobe, dark red the limbic lobe, pink the occipital lobe, and magenta the subcortical nuclei. **(B)** Between-subtype differences in factor-related functional connectivity (top: rsFC correlated with the self-related factor; bottom: rsFC correlated with the social-related factor). From left to right, self-subtype is compared with social-subtype and non-specific subtype, followed by comparisons between the latter two. Red lines indicate that the t-value of comparison is greater than 0, while blue lines indicate that the *T*-value is less than 0.

Self-related Function

Social-related Function

NSSI   OSI-F   OPNMF

Functional Subtypes

Neural Correlates

Psychosocial Risks

A Factor structure of NSSI functions

Item 17: To help me escape from uncomfortable feelings or moods
Item 1: To release unbearable tension
Item 21: To relieve feelings of sadness or feeling "down"
Item 19: To experience physical pain in one area, when the other pain I feel is unbearable
Item 6: To punish myself
Item 25: To produce a sense of being real when I feel numb and ``unreal''
Item 8: To relieve nervousness/fearfulness
Item 10: To distract me from unpleasant memories
Item 2: To experience a ``high'' like a drug high
Item 28: For no reason that I know about - it just happens sometimes
Item 7: To provide a sense of excitement that feels exhilarating
Item 4: To stop feeling alone and empty
Item 23: To stop me from thinking about ideas of killing myself
Item 24: To stop me from acting out ideas to kill myself
Item 18: To satisfy voices inside or outside of me telling me to do it

Self-related factor

Item 26: To release frustration
Item 13: To release anger
Item 15: To show others how hurt or damaged I am
Item 5: To get care and attention from other people
Item 27: To get out of doing something that I don't want to do
Item 22: To have control in a situation where no one can influence me
Item 20: To stop people from expecting so much from me
Item 9: To avoid getting in trouble for something I did
Item 3: To stop my parents from being angry at me
Item 29: To prove to myself how much I can take
Item 14: To stop my friends/boyfriend/girlfriend from being anger at me
Item 11: To change my body image and/or appearance
Item 12: To belong to a group
Item 16: To show others how strong or though I am

Social-related factor

Coefficients of the loading matrix
1 — 10

B Correlations between factor items

Item: 1 2 4 6 7 8 10 17 18 19 21 23 24 25 28 | 3 5 9 11 12 13 14 15 16 20 22 26 27 29

Self-related factor

Social-related factor

Correlations (*r*)
-0.8 — 0.8

C Correlations between factor-loading and clinical features

r = 0.167**
p = 0.004

r = 0.075
p = 0.194

r = 0.075
p = 0.191

r = 0.229***
p < 0.001

Loading of self-related factor

Loading of social-related factor

SDS scores

SAS scores

Diagnosis labels

MDD   BD   ED

***
**

***
*

A Dimensional representation of connectivity features related with NSSI-factor

B Between-subtypes difference in factor-related functional connectivity

A Factor structure space for clustering subtypes

B Between-subtypes comparison results

C Between-subtypes comparison results of risk factors

D Significant between-subtypes comparison results of risk factors