Explainable advanced electrocardiography at rest can rule out myocardial ischemia on stress echocardiography

Kevin X Yang[1,2], Johan von Scheele[3], Maren Maanja[3], Daniel E Loewenstein[3],

Todd T Schlegel[3,4], Martin Ugander[1,2,3*], Rebecca Kozor[1*]

[1] Kolling Institute, Royal North Shore Hospital, and Faculty of Medicine and Health,

University of Sydney, Sydney, Australia

[2] St Vincent's Clinical School, University of New South Wales, Sydney, Australia

[3] Department of Clinical Physiology, Karolinska University Hospital, and Karolinska

Institutet, Stockholm, Sweden

[4] Nicollier-Schlegel SARL, Trélex, Switzerland

* equal contributions as senior author

Key words: Advanced ECG; Stress echocardiography

Corresponding author:

Dr Rebecca Kozor, BSc (Med) MBBS PhD FRACP FCSANZ

Associate Professor, Faculty of Medicine and Health, University of Sydney

Cardiologist, Royal North Shore Hospital | North Shore Private Hospital

Address: Kolling Building, Level 12 | Royal North Shore Hospital | St Leonards NSW 2065 |

Australia

Email: rebecca.kozor@sydney.edu.au

Abstract

Background: Stress echocardiography (SE) is relatively resource intensive and has a low incidence of abnormal tests for detecting coronary artery disease (CAD) in low-to-intermediate risk patients. This study aimed to derive and determine the diagnostic performance of a resting advanced electrocardiography (A-ECG) score for detecting inducible myocardial ischemia on SE in patients with low-to-intermediate risk stable chest pain. **Methods:** Patients were included if they presented with low-to-intermediate risk stable chest pain to the emergency department, had acute coronary syndrome ruled out by electrocardiography (ECG) and high-sensitivity troponin, and subsequently underwent outpatient SE. Patients were excluded if they had known CAD or confounders on resting ECG. A-ECG was retrospectively applied to a standard resting 12-lead ECG and a multivariable logistic regression score was derived to predict myocardial ischemia on SE. **Results:** Among 292 patients (51% male, age 58±14 years), 24 (8%) exhibited inducible myocardial ischemia on SE. A 3-parameter A-ECG score had an area under the receiver-operating characteristic curve (AUC [bootstrapped 95% confidence interval]) of 0.85 [0.75–0.93], sensitivity 92 [67–100]%, specificity 67 [64–94]%, positive predictive value 22 [20–55]%, negative predictive value 99 [96–100]%, positive likelihood ratio 2.8 [2.5–12.0] and inverse negative likelihood ratio 8.1 [2.5-18.0]. **Conclusions:** An A-ECG score had a good overall diagnostic performance and excellent performance for ruling out inducible myocardial ischemia on SE. This supports the use of an A-ECG score to triage and improve the selection of patients with low-intermediate risk stable chest pain that should undergo further testing with SE.

Introduction

Stress echocardiography (SE) is well-validated for diagnostic and prognostic use in coronary artery disease (CAD)[1,2]. It is routinely performed to investigate stable chest pain in patients with a low-intermediate risk of CAD. However, SE is relatively time and resource-consuming, requiring considerable expertise for reliable interpretation. The incidence of inducible myocardial ischemia on SE and other stress imaging modalities has also declined over time, now as low as 4% in contemporary populations[3,4]. This incurs unnecessary costs to patients and the healthcare system without substantial clinical utility. As such, there is an unmet need for a triage tool that more accurately predicts which patients will exhibit inducible myocardial ischemia.

The standard resting 12-lead electrocardiogram (ECG) is simple, low-cost, quickly performed, and widely available. It is ubiquitously conducted for patients presenting to the emergency department with chest pain. However, visual assessment of the resting ECG has poor diagnostic accuracy for detecting CAD[5]. Advanced electrocardiography (A-ECG) involves the measurement of various advanced and conventional ECG measures and subsequently using multivariable statistics to detect heart disease[6]. A-ECG incorporates signal averaging, derived vectorcardiography, and measurement of QRS and T waveform complexity by singular value decomposition[7,8]. A-ECG analysis can be applied to a standard resting 12-lead digital ECG, providing diagnostic information that cannot be discerned by visual assessment. In one study, A-ECG was shown to rule out significant CAD based on cardiac computed tomography angiography (CCTA) with high specificity[9]. This supports its use as a powerful triaging tool for chest pain by identifying patients who do not require further investigations. However, A-ECG has not yet been validated using SE.

Therefore, the aim of the study was to derive and determine the diagnostic performance of an

A-ECG score for detecting inducible myocardial ischemia on SE in patients with low-to-

intermediate risk stable chest pain. We hypothesised that an A-ECG score from the resting

12-lead ECG would have a clinically useful diagnostic performance for detecting inducible

myocardial ischemia on SE in these patients.

Methodology

Study participants

The study population consisted of adult patients retrospectively recruited between February

2017 and March 2022 who presented with cardiac-suspected chest pain to the emergency

department at Royal North Shore Hospital, had myocardial infarction ruled out by resting

ECG and blood troponin levels, were deemed low-to-intermediate risk by HEART score[10],

and were subsequently referred to the rapid access chest pain clinic (RACC) for SE. Patients

were excluded if they had a pacemaker or known CAD defined as having a history of

myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting or

documented coronary artery stenosis >50% on invasive angiography or CCTA. Patient data

was retrospectively obtained from electronic medical records. The study was approved by the

Northern Sydney Local Health District Human Research Ethics Committee

(2019/STE10454). All patients provided written informed consent, or a waiver of consent

was provided by the ethics committee.

Stress echocardiography

All SE tests were performed using an exercise treadmill following the Bruce protocol[11].

Inducible myocardial ischemia on SE was defined as a stress-induced left ventricular regional

wall motion abnormality as determined by the reporting cardiologist in the RACC. Patients

were excluded if they had an uninterpretable test or if their peak heart rate was less than 85%

of their maximum predicted heart rate (220 – age) without ischemia.

Exercise electrocardiography

Treadmill exercise ECG (ExECG) was performed simultaneously alongside SE in all

patients. Inducible myocardial ischemia on ExECG was defined as $\geq$ 1 millimetre of

horizontal or down-sloping ST depression in 2 contiguous leads during exercise or recovery.

All ST segment measurements were performed 80 milliseconds after the J-point. Patients

who tested negative on SE but had equivocal findings on ExECG (horizontal/down-sloping

ST segment depression < 1 millimetre) were excluded.

Resting ECG Acquisition and Analysis

A 10-second resting digital ECG was performed (Philips, Best, the Netherlands) during

RACC visit prior to SE. Patients were excluded if their digital ECG could not be

retrospectively retrieved. A large proportion of digital ECGs were not stored due to technical

issues in the early establishment of the RACC. Available ECG files were anonymized,

reviewed, and excluded if they contained atrial fibrillation or atrial flutter, QRS duration >

120 milliseconds, missing leads, or excessive noise. Additionally, a large proportion of files

contained 50 Hz noise from electrical interference which was not detectable at the time of

recording due to a noise filter, and needed to be excluded. Included ECGs were analysed

using in-house software to produce conventional and advanced ECG parameters.

Conventional parameters included all major intervals, axes, voltages and ST segment levels.

Advanced parameters included derived vectorcardiographic measures and waveform

complexity measures of the QRS and T wave quantified by singular value decomposition.

Statistical Analysis

All statistical analyses were performed using the programming language R 4.1.2 (R Core Team, R Foundation for Statistical Computing, Vienna, Austria). Categorical variables are expressed as frequencies and percentages. Continuous variables are expressed as mean± standard deviation. Baseline characteristics between groups were compared using Fisher's exact test for categorical variables and one-way analysis of variance for continuous variables. A multivariable logistic regression model was trained for inducible myocardial ischemia on SE. Feature selection was performed by first identifying univariately significant parameters using the Mann-Whitney U test. Exhaustive search was then performed on these parameters to identify the best performing model, with the number of parameters included in the model limited to 3 (~1 parameter per 10 events). Performance was measured using the area under the receiver operating characteristic curve (AUC), sensitivity, specificity, predictive values, and likelihood ratios. The multivariate statistical significance of model parameters were evaluated using the likelihood-ratio chi-square test. Multicollinearity between model parameters was evaluated using variance inflation index. The optimal probability threshold was determined as the value that maximised Youden's index. The intercept for the logistic model was adjusted so that the probability threshold for a positive result was set to 50%. Non-parametric percentile bootstrap was performed with 2000 samples to assess the stability of the coefficients and calculate 95% confidence intervals for all performance metrics[12]. A two-sided p-value < 0.05 was considered statistically significant.

Results

Study population

The patient inclusion flowchart is shown in Figure 1. From 1,670 patients with chest pain who were managed through the RACC during the time period, 589 underwent SE plus had an available digital clinic ECG, and after exclusions, 292 patients were included as the final study population. The baseline characteristics are presented in Table 1. The mean age was 58±14 years and 150 (51%) were males. The majority of patients had a left ventricular ejection fraction (LVEF) > 50%, 6 (2%) had a mildly reduced LVEF (40–50%), and none had an LVEF <40%. The mean HEART score was 3±1, reflecting patients had a low-to-intermediate 6-week risk of MACE[10]. Both SE and ExECG were negative for inducible myocardial ischemia in 240 patients (82%), forming the negative SE group. Inducible myocardial ischemia was present on SE in 24 patients (8%), forming the positive SE group. The final subgroup were those with a negative SE but a positive ExECG positive (29 patients (10%), forming the discordant ExECG group. Patients in the positive SE and discordant ExECG groups were older than those in the negative group (p=0.01), otherwise the three groups were similar in baseline characteristics.

Development of the A-ECG Score

The best 3-parameter A-ECG score for predicting inducible myocardial ischemia by SE consisted of one conventional ECG measure and two advanced ECG measures, as follows: (1) the natural log of the R wave duration in lead V6 (R Dur V6), (2) the normalised 4th singular value of the T wave (4th T Eigen), and (3) the azimuth direction at the final eighth of the spatial ST-T loop (ST-T Azimuth). All included measures were minimally correlated by variance inflation factor (< 1.1) and had stable coefficients across bootstrapping. These measures are compared between the negative and positive groups in Table 2. The regression

coefficients are reported in *Table 3*. After re-arranging the logit function, the A-ECG score

for inducible ischemia on SE was calculated as:

$$AECG\ Score = \frac{1}{1 + e^{-[-15 + R\ Dur\ V6\ *\ 3.5 + 4th\ T\ Eigen\ *\ 0.16 + ST-T\ Azimuth\ *\ (-0.013)]}} * 100$$

A-ECG Score Performance

The diagnostic performance of the optimal 3-parameter score for detecting inducible

myocardial ischemia on SE is presented in Table 4 and Figure 2. The probability cut-off with

the highest Youden's index was 0.06, and this was scaled to 50% by adjusting the intercept of

the model. Of the 24 patients in the positive SE group, 2 (8%) had scores below the adjusted

threshold of 50%, representing false negatives. 78 out of the 239 patients (33%) in the

negative group had scores above 50%, representing false positives. As an example, Figure 3

compares the resting ECGs between two representative patients with high and low A-ECG

scores for inducible myocardial ischemia on SE.

Discussion

This study demonstrates that a 3-parameter A-ECG score applied to a standard resting 12-

lead ECG has a good AUC and excellent negative predictive value and inverse negative

likelihood for ruling out inducible myocardial ischemia on SE. This is the first study to

investigate the ability of A-ECG to detect inducible myocardial ischemia on SE, and thus

could be used as a simple test to help triage low-to-intermediate risk patients into needing a

SE or not.

SE has been validated as an excellent non-invasive test for the diagnosis of significant CAD

and is widely performed to investigate stable chest pain[1,2]. Although invasive angiography is

the gold standard for the diagnosis of CAD, SE provides independent prognostic

information[13,14]. For these reasons, SE was chosen as the primary outcome to train the logistic

model.

Explainable A-ECG Measures

Measures of waveform complexity are obtained through singular value decomposition, a

method of dimensionality reduction. In healthy patients, most of the variance can be

explained from a single eigenvalue (the 1st eigenvalue). In patients with cardiac disease,

higher numbered eigenvectors account for greater voltage as the signal becomes more

complex[15]. The normalised 4th singular value of the T wave was included in our model, being

elevated in patients with positive SE. This indicates elevated T wave complexity and

abnormal repolarisation. Large studies have shown measures of waveform complexity to

predict incident heart failure and cardiovascular mortality[8,16].

The QRS complex reflects depolarisation of the ventricular myocardium. The natural log of

the R wave duration in lead V6 was included in our model, being elevated in patients with

positive SE. This may reflect conduction delay caused by ischemia to the conduction system

and myocardium. In general populations, QRS duration is independently associated with

CAD, myocardial infarction and cardiovascular mortality[17,18]. Also, in patients referred to

stress echocardiography for suspected CAD, QRS duration is an independent predictor of

cardiac death and non-fatal infarction[19].

Vectorcardiography (VCG) is the representation of the heart's electrical activity in three

spatial planes[20]. In A-ECG analysis, the VCG is derived from the standard 12-lead ECG

using validated methods[21], providing ready access to hundreds of additional VCG parameters.

Our model contained one VCG-derived measure: the azimuth direction at the final eighth of

the spatial ST-T loop, which was more positively directed in patients with negative SE. The

mean values for this measure in our negative SE cohort are similar to those previously

reported for healthy men[22], whereas those in our positive SE cohort likely diverge from

normal[23]. However, the mechanisms behind these observed directional differences in T wave

orientation require further elucidation.

A-ECG Model Performance

The high NPV and low PPV observed in our model is inherent to the low prevalence of

outcome events. If used in a similar population to ours, patients with chest pain that receive

an A-ECG score < 50% are highly unlikely to exhibit inducible myocardial ischemia on SE

and thus we suggest would not require a SE or further testing for CAD. The poorer PPV

indicates that a positive A-ECG score does not guarantee inducible ischemia on SE. At face

value, this would greatly reduce its clinical utility. However, the utility of the A-ECG score is

not to replace SE, but to optimise its testing population. In our study cohort, if the 163 out of

263 patients with an A-ECG score < 50% were discharged without referral to SE, only 2 of

these (1.2%) would have gone on to test positive. Doing so would greatly reduce the level of

unnecessary testing and associated costs while missing very few patients that would

otherwise test positive.

Comparison to Clinical Risk Scores

Current triage of patients to SE depends on risk of major adverse cardiovascular events and

pre-test probability for CAD based on clinical assessment. Clinical risk scores such as the

Diamond-Forrester and Duke clinical cores were developed to estimate pre-test probability

for CAD based on easily obtained clinical findings[24,25]. However, these scores have been

shown to overestimate pre-test probability for CAD in contemporary populations[26,27]. This

may contribute to the decrease in incidence of positive SE results over the last decades[3,4].

There is a growing sentiment that the level of cardiac testing is inappropriately high,

incurring unnecessary costs to patients and the healthcare system[28]. One study aimed to

identify the clinical factors that specifically predicted inducible myocardial ischemia on

nuclear stress testing[29]. Known CAD, typical angina and reduced LVEF were identified as

the strongest independent predictors[29]. Our study offers an alternative method of estimating

risk of CAD, by using advanced and conventional ECG parameters. We did not incorporate

any clinical variables into our model in order to specifically assess the predictive power of A-

ECG variables alone. While clinical predictors are robust due to their extensive real-world

usage and validation[26], A-ECG measures have a theoretical basis alongside proven clinical

utility in many large studies[7,8,16,30].

Comparison to Conventional ECG

The resting conventional ECG is readily available and forms a component of many clinical

risk scores for CAD. Previous studies have attempted to quantify the diagnostic potential of

the resting conventional ECG for CAD outside of STEMI. One cross-sectional study

demonstrated that patients with resting ECG abnormalities have a higher prevalence of

subclinical atherosclerosis by coronary calcium score[31]. Another study determined that

resting ECG abnormalities defined by the Minnesota code could predict myocardial ischemia

on stress perfusion magnetic resonance imaging with a hazard ratio of 2.5 (95% CI: 1.4-

4.4)[32]. Our study did not use pre-determined ECG abnormalities to predict myocardial

ischemia. Rather, features selection methods were used to select the optimal parameters from

a large pool of conventional and advanced ECG parameters. As a result, our A-ECG score

can predict myocardial ischemia on SE even on resting ECGs that demonstrate no remarkable

findings to the naked eye. An example of this is demonstrated in Figure 3.

Comparison to Other A-ECG Scores

The excellent performance of A-ECG for predicting inducible myocardial ischemia on SE is

in alignment with previous studies which have demonstrated its ability to distinguish various

cardiac conditions[6,9,33-35]. A-ECG has been shown to outperform optimized pooled criteria

from the conventional ECG. A 7-parameter A-ECG score has been shown to have a

sensitivity of 89% and specificity of 94% for predicting pooled CAD, left ventricular

hypertrophy, and left ventricular systolic dysfunction[6]. Also, within our same cohort of chest

pain patients, A-ECG has been validated with a high specificity of 86% for ruling out

significant stenosis on CCTA[9]. A recent study showed that A-ECG can accurately diagnose

apical hypertrophic cardiomyopathy while also distinguishing from other cardiac disease

using linear discriminant analysis[35]. This was able to accurately identify CAD with an AUC

of 0.99 and specificity of 98%[35].

Comparison to Artificial Intelligence

There is an emerging body of research addressing the medical applications of machine

learning, with the ECG being a prime target due to its raw digital data that is available in

large numbers for training[36]. Multivariable logistic regression itself is a form of supervised

machine learning used for regression and classification. Artificial intelligence (AI) usually

refers to unsupervised deep learning, which is beneficial in that it does not require human

selection of parameters that is limited by current scientific understanding[36]. Studies have

shown that AI can accurately predict various cardiac conditions from a resting ECG[37,38].

However, the prevailing issue with unsupervised AI is that its decision process cannot be

interpreted by humans – a so-called "black box". Consequently, there exists a sentiment that

AI may not be appropriate for guiding medical decisions[39]. By comparison, the A-ECG score

from the current study is comprised of known A-ECG measures and can therefore be

considered transparent or explainable, when compared to unsupervised AI methods.

Clinical Implications

The current study demonstrates the utility of a resting 12-lead A-ECG score to screen low-to-

intermediate risk chest pain patients for those that would exhibit inducible myocardial

ischemia on SE. The conventional 12-lead ECG is routinely performed in the emergency

department and outpatient clinics as a quick, cost-effective, and simple test for chest pain. An

A-ECG score inherits these advantages plus provides diagnostic information that cannot be

discerned by the human eye. Automatic A-ECG analysis of a resting 12-lead ECG can

quickly identify the probability of testing positive for inducible myocardial ischemia on SE.

This analysis can be applied to ECGs acquired from any vendor that stores a digital raw data

file that is accessible to the user. Hence, the addition of an A-ECG score may significantly

improve triage to SE and reduce unnecessary testing in this patient population.

Limitations

The study was performed retrospectively from a single centre and may not be applicable to

other patient populations. It also contains a limited sample size, particularly in the outcome

group. There exists the possibility of overfitting, whereby the model trains on noise specific

to our patient sample. Consequently, the diagnostic performance of the A-ECG score may

decrease when applied to another sample. Due to the limited data, it was not feasible to split

patients into a training and test group. Instead, the model was trained on all available data and

resampled using non-parametric bootstrapping to obtain 95% confidence intervals. Also, the

number of parameters incorporated into the model was limited to 3 to prevent overfitting.

We did not correlate the SE results with any further tests such as invasive or CT coronary

angiography.

Future studies should aim to test the reproducibility of the A-ECG score in an independent

validation cohort, the clinical implementation in an emergency department setting, and

assessing short and long term clinical and economical outcomes.

Conclusions

A 3-parameter resting A-ECG score has a good AUC and excellent inverse negative

likelihood ratio for detecting inducible myocardial ischemia on SE. This supports the use of

A-ECG assessment on a resting 12-lead ECG to effectively identify patients with chest pain

at low-to-intermediate risk that do or do not require further investigation for CAD. However,

further larger prospective validation in independent cohorts is warranted.

Disclosures

Todd T Schlegel is owner and founder of Nicollier-Schlegel SARL, which performs ECG interpretation consultancy using software used in the current study. Todd T Schlegel and Martin Ugander are founders of Advanced ECG Systems, a company that is developing commercial applications of advanced ECG technology used in the current study. Rebecca Kozor has a financial interest in Advanced ECG Systems through being married to Martin Ugander.

Funding

Rebecca Kozor was funded in part by a grant from New South Wales Health.

Authorship

Kevin X Yang: Software, Validation, Formal Analysis, Investigation, Data curation, Writing - Original Draft, Writing – Review & Editing, Visualisation. **Johan von Scheele:** Software, Validation, Formal analysis, Investigation, Data curation, Writing – Original Draft, Writing – Review & Editing, Visualisation. **Maren Maanja:** Software, Formal analysis, Writing – Review & Editing. **Daniel E Loewenstein:** Software, Formal analysis, Resources, Writing – Review & Editing. **Todd T Schlegel:** Software, Formal analysis, Resources, Writing – Review & Editing. **Martin Ugander:** Conceptualisation, Methodology, Resources, Writing - Review & Editing, Supervision, Project Administration, **Rebecca Kozor:** Conceptualisation, Methodology, Resources, Data Curation, Writing - Review & Editing, Supervision, Project Administration.

References

Pontone, G. *et al.* Diagnostic performance of non-invasive imaging for stable coronary artery disease: A meta-analysis. *Int. J. Cardiol.* **300**, 276-281 (2020).

Fleischmann, K. E., Hunink, M. G., Kuntz, K. M. & Douglas, P. S. Exercise echocardiography or exercise SPECT imaging? A meta-analysis of diagnostic test performance. *JAMA* **280**, 913-920 (1998).

Cortigiani, L., Ramirez, P., Coltelli, M., Bovenzi, F. & Picano, E. Drop-off in positivity rate of stress echocardiography based on regional wall motion abnormalities over the last three decades. *Int. J. Cardiovasc. Imaging* **35**, 627-632 (2019).

Bouzas-Mosquera, A. *et al.* Temporal changes in the use and results of exercise echocardiography. *Eur. Heart J. Cardiovasc. Imaging* **16**, 1207-1212 (2015).

Mahmoodzadeh, S., Moazenzadeh, M., Rashidinejad, H. & Sheikhvatan, M. Diagnostic performance of electrocardiography in the assessment of significant coronary artery disease and its anatomical size in comparison with coronary angiography. *J. Res. Med. Sci.* **16**, 750-755 (2011).

Schlegel, T. T. *et al.* Accuracy of advanced versus strictly conventional 12-lead ECG for detection and screening of coronary artery disease, left ventricular hypertrophy and left ventricular systolic dysfunction. *BMC Cardiovasc. Disord.* **10**, 28 (2010).

Kardys, I. *et al.* Spatial QRS-T angle predicts cardiac death in a general population. *Eur. Heart J.* **24**, 1357-1364 (2003).

Rautaharju, P. M., Kooperberg, C., Larson, J. C. & LaCroix, A. Electrocardiographic predictors of incident congestive heart failure and all-cause mortality in postmenopausal women: the Women's Health Initiative. *Circulation* **113**, 481-489 (2006).

Rajamohan, M. *et al.* Advanced Electrocardiography Has A High Negative Predictive Value For Ruling Out Significant Coronary Artery Disease By Cardiovascular CT. *J. Cardiovasc. Comput. Tomogr.* **15**, S1 (2021).

Six, A. J., Backus, B. E. & Kelder, J. C. Chest pain in the emergency room: value of the HEART score. *Neth. Heart J.* **16**, 191-196 (2008).

Bruce, R. A. Exercise testing of patients with coronary heart disease. Principles and normal standards for evaluation. *Ann. Clin. Res.* **3**, 323-332 (1971).

Carpenter, J. & Bithell, J. Bootstrap confidence intervals: when, which, what? A practical guide for medical statisticians. *Stat. Med.* **19**, 1141-1164 (2000).

Gilchrist, I., Foy, A. & Davidson, W. A FALSE-POSITIVE STRESS ECHOCARDIOGRAM IS NOT THE SAME AS A NEGATIVE RESULT. *J. Am. Coll. Cardiol.* **69**, 1426-1426 (2017).

Sicari, R., Palinkas, A., Pasanisi, E. G., Venneri, L. & Picano, E. Long-term survival of patients with chest pain syndrome and angiographically normal or near-normal coronary arteries: the additional prognostic value of dipyridamole echocardiography test (DET). *Eur. Heart J.* **26**, 2136-2141 (2005).

Priori, S. G. *et al.* Evaluation of the Spatial Aspects of T-Wave Complexity in the Long-QT Syndrome. *Circulation* **96**, 3006-3012 (1997).

Okin, P. M. *et al.* Principal component analysis of the T wave and prediction of cardiovascular mortality in American Indians: the Strong Heart Study. *Circulation* **105**, 714-719 (2002).

Deen, J. F. *et al.* Comparison of QRS Duration and Associated Cardiovascular Events in American Indian Men Versus Women (The Strong Heart Study). *Am. J. Cardiol.* **119**, 1757-1762 (2017).

Badheka, A. O. *et al.* QRS Duration on Electrocardiography and Cardiovascular Mortality (from the National Health and Nutrition Examination Survey—III). *Am J Cardiol* **112**, 671-677 (2013).

Schinkel, A. F. *et al.* Prognostic significance of QRS duration in patients with suspected coronary artery disease referred for noninvasive evaluation of myocardial ischemia. *Am. J. Cardiol.* **104**, 1490-1493 (2009).

Pérez Riera, A. R. *et al.* Significance of vectorcardiogram in the cardiological diagnosis of the 21st century. *Clin. Cardiol.* **30**, 319-323 (2007).

Kors, J. A., van Herpen, G., Sittig, A. C. & van Bemmel, J. H. Reconstruction of the Frank vectorcardiogram from standard electrocardiographic leads: diagnostic comparison of different methods. *Eur. Heart J.* **11**, 1083-1092 (1990).

Draper, H. W., Peffer, C. J., Stallmann, F. W., Littmann, D. & Pipberger, H. V. THE CORRECTED ORTHOGONAL ELECTROCARDIOGRAM AND VECTORCARDIOGRAM IN 510 NORMAL MEN (FRANK LEAD SYSTEM). *Circulation* **30**, 853-864 (1964).

Pipberger, H. V., Arms, R. J. & Stallmann, F. W. Automatic screening of normal and abnormal electrocardiograms by means of digital electronic computer. *Proc Soc Exp Biol Med* **106**, 130-132 (1961).

Diamond, G. A. & Forrester, J. S. Analysis of probability as an aid in the clinical diagnosis of coronary-artery disease. *N. Engl. J. Med.* **300**, 1350-1358 (1979).

Pryor, D. B., Harrell, F. E., Jr., Lee, K. L., Califf, R. M. & Rosati, R. A. Estimating the likelihood of significant coronary artery disease. *Am. J. Med.* **75**, 771-780 (1983).

Genders, T. S. *et al.* A clinical prediction rule for the diagnosis of coronary artery disease: validation, updating, and extension. *Eur. Heart J.* **32**, 1316-1330 (2011).

Wasfy, M. M. *et al.* Comparison of the Diamond-Forrester method and Duke Clinical Score to predict obstructive coronary artery disease by computed tomographic angiography. *Am. J. Cardiol.* **109**, 998-1004 (2012).

Ladapo, J. A., Blecker, S. & Douglas, P. S. Physician decision making and trends in the use of cardiac stress testing in the United States: an analysis of repeated cross-sectional data. *Ann. Intern. Med.* **161**, 482-490 (2014).

Rozanski, A. *et al.* The prevalence and predictors of inducible myocardial ischemia among patients referred for radionuclide stress testing. *J. Nucl. Cardiol.* (2021).

Yamazaki, T., Froelicher, V. F., Myers, J., Chun, S. & Wang, P. Spatial QRS-T angle predicts cardiac death in a clinical population. *Heart Rhythm* **2**, 73-78 (2005).

Kang, M., Chang, Y., Kang, J., Kim, Y. & Ryu, S. Electrocardiogram Risk Score and Prevalence of Subclinical Atherosclerosis: A Cross-Sectional Study. *J. Pers. Med.* **12** (2022).

Kaolawanich, Y., Thongsongsang, R., Songsangjinda, T. & Boonyasirinant, T. Clinical values of resting electrocardiography in patients with known or suspected chronic coronary artery disease: a stress perfusion cardiac MRI study. *BMC Cardiovasc. Disord.* **21**, 621 (2021).

Johnson, K. *et al.* Advanced Electrocardiography Identifies Left Ventricular Systolic Dysfunction in Non-Ischemic Cardiomyopathy and Tracks Serial Change over Time. *J. Cardiovasc. Dev. Dis.* **2**, 93-107 (2015).

Fernlund, E., Liuba, P., Carlson, J., Platonov, P. G. & Schlegel, T. T. MYBPC3 hypertrophic cardiomyopathy can be detected by using advanced ECG in children and young adults. *J. Electrocardiol.* **49**, 392-400 (2016).

Hughes, R. K. *et al.* Accurate diagnosis of apical hypertrophic cardiomyopathy using explainable advanced electrocardiogram analysis. *Europace* **26** (2024).

Siontis, K. C., Noseworthy, P. A., Attia, Z. I. & Friedman, P. A. Artificial intelligence-enhanced electrocardiography in cardiovascular disease management. *Nat. Rev. Cardiol.* **18**, 465-478 (2021).

Huang, P. S. *et al.* An Artificial Intelligence-Enabled ECG Algorithm for the Prediction and Localization of Angiography-Proven Coronary Artery Disease. *Biomedicines* **10**, 394 (2022).

Attia, Z. I. *et al.* An artificial intelligence-enabled ECG algorithm for the identification of patients with atrial fibrillation during sinus rhythm: a retrospective analysis of outcome prediction. *Lancet* **394**, 861-867 (2019).

Rudin, C. Stop explaining black box machine learning models for high stakes decisions and use interpretable models instead. *Nat. Mach. Intell.* **1**, 206-215 (2019).

Table 1. Patient demographics stratified by stress testing result.

	Total (n = 292)	Negative (n = 239)	Discordant ExECG (n = 29)	Positive SE (n = 24)	p-value
Demographics					
Age, years	58±14	56±14	63±11	61±14	0.01
Male sex, n (%)	150 (51)	122 (51)	15 (52)	13 (54)	0.97
BMI, kg/m²	27±5	27±5	26±5	28±6	0.27
Resting SBP, mmHg	130±15	129±15	130±16	134±17	0.26
Resting DBP, mmHg	76±10	76±10	76±9	74±11	0.67
HEART Score	3±1	3±1	3±1	4±1	0.17
Peak HR, % of MPHR	100±8	101±8	100±9	100±10	0.80
Baseline LVEF					0.049
>50%, n (%)	286 (98)	236 (99)	28 (97)	22 (92)	
40-50%, n (%)	6 (2)	3 (1)	1 (3)	2 (8)	
Comorbidities, n (%)					
Hypertension	124 (42)	99 (41)	13 (45)	12 (50)	0.71
Diabetes	34 (12)	24 (10)	7 (24)	3 (12)	0.08
Dyslipidemia	142 (49)	113 (47)	17 (59)	12 (50)	0.51
Obesity	76 (26)	64 (27)	4 (14)	8 (33)	0.24
Current Smoker	34 (12)	27 (11)	3 (10)	4 (17)	0.68
Ex-Smoker	81 (28)	66 (28)	6 (21)	9 (38)	0.39
Stroke/TIA	7 (2)	5 (2)	2 (7)	0 (0)	0.24
AF/AFL	12 (4)	8 (3)	2 (7)	2 (8)	0.21
COPD	4 (1)	3 (1)	1 (3)	0 (0)	0.55
Asthma	19 (6)	16 (7)	3 (10)	0 (0)	0.30
GERD	41 (14)	33 (14)	6 (21)	2 (8)	0.46
Medications, n (%)					
Aspirin	23 (8)	16 (7)	4 (14)	3 (12)	0.19
Statin	72 (25)	60 (25)	6 (21)	6 (25)	0.93
ACE/ARB	83 (28)	66 (28)	8 (28)	9 (38)	0.58
CCB	34 (12)	26 (11)	3 (10)	5 (21)	0.33
Beta-blocker	14 (5)	9 (4)	4 (14)	1 (4)	0.07
Insulin	8 (3)	4 (2)	3 (10)	1 (4)	0.03
Oral hypoglycaemic	32 (11)	25 (10)	6 (21)	1 (4)	0.15

Continuous variables are expressed as mean± standard deviation and compared using one way analysis of variance. Categorical variables are expressed as frequencies (percentage) and compared using Fisher's exact test. Abbreviations: SE = stress echocardiography; ExECG = exercise electrocardiography, Discordant ExECG = positive ExECG with negative SE; n = frequency; BMI = body mass index; mmHg = millimetres of mercury; kg/m² = kilograms/metres²; SBP = systolic blood pressure; DBP = diastolic blood pressure; MPHR = maximum predicted heart rate (220 – age); HR = heart rate; LVEF = left ventricular ejection fraction; TIA = transient ischemic attack; AF = atrial fibrillation; AFL = atrial flutter; COPD = chronic obstructive pulmonary disease; GERD = gastroesophageal reflux disease; ACE = angiotensin-converting enzyme inhibitor; ARB = angiotensin II receptor blocker; CCB = calcium channel blockers

Table 2. Parameter Means with Mann Whitney U test.

Parameter	Total (n = 263)	Negative SE (n = 239)	Positive SE (n = 24)	p-value
Natural log of the R wave duration in lead V6 (log ms)	3.9±0.3	3.9±0.3	4.2±0.3	<0.001
Normalised 4^{th} singular value of the T wave (µV)	8.1±4.7	7.7±3.8	13±8.8	0.001
Azimuth direction at the final eighth of the ST-T loop (degrees)	17±53	20±50	-13±67	0.009

Continuous variables are expressed as mean± standard deviation and compared using Mann Whitney U test. Abbreviations: SE = stress echocardiography.

Table 3. Regression coefficients for the logistic model.

	Coefficient	Chi-square	p-value
Intercept	-15		-
Natural log of the R wave duration in lead V6 (log ms)	3.5	17	<0.001
Normalised 4th singular value of the T wave (μV)	0.16	17	<0.001
Azimuth direction at the final eighth of the ST-T loop (degrees)	-0.013	9.3	0.002

Coefficients are assessed using the likelihood-ratio chi-square test.

Table 4. Diagnostic performance of 3-parameter score for inducible myocardial ischemia on stress echocardiography.

Diagnostic performance measure	Estimate [95% confidence interval]
Area under the receiver operating characteristics curve	0.85 [0.75–0.93]
Sensitivity, %	92 [67–100]
Specificity, %	67 [64–94]
Positive predictive value, %	22 [20–55]
Negative predictive value, %	99 [96–100]
Positive likelihood ratio, %	2.8 [2.5–12.0]
Negative likelihood ratio, %	0.12 [0.00–0.39]
Inverse negative likelihood ratio, %	8.1 [2.5-18.0]

Estimates are from the unchanged training data. 95% confidence intervals are from bootstrapped percentiles.

Figure 1. Patient inclusion flow chart and stress testing results.

Abbreviations: ECG = electrocardiogram; SE = stress echocardiography; ExECG = exercise

electrocardiography; A-ECG = advanced electrocardiography; CAD = coronary artery

disease; MI = myocardial infarction; PCI = percutaneous coronary intervention; CABG =

coronary artery bypass grafting; MPHR = maximum predicted heart rate (220-age);

Equivocal ExECG changes = horizontal/down-sloping ST segment depression < 1 millimetre

Figure 2. Empirical receiver-operating characteristic (ROC) curve and its 95% confidence

interval for the 3-parameter score trained for inducible myocardial ischemia on stress

echocardiography. The random chance line is given by**:** true positive rate (sensitivity) = false

positive rate (1- specificity), representing an ROC curve with no discrimination.

Abbreviations: ROC = receiver-operating characteristic

Figure 3a. Patient A

Figure 3b. Patient B

Figure 3. Comparison of A-ECG results of two representative patients.

Both Patient A and Patient B had no signs of ischemia on resting ECGs and normal resting echocardiography. On stress echocardiography, Patient A showed no evidence of inducible myocardial ischemia, with a low corresponding A-ECG score for inducible myocardial ischemia (2%). Patient B developed hypokinesis of the basal-to-mid left ventricular inferior wall, with a high corresponding A-ECG score for inducible myocardial ischemia (95%).

Table 5 – Clinical and ECG information comparison between Patient A and Patient B

	Patient A	Patient B
Myocardial ischemia on SE		
True result	Negative	Positive
A-ECG score	95%	2%
A-ECG measures		
Natural log of the R wave duration in lead V6 (log ms)	3.6	3.8
Normalised 4th singular value of the T wave (µV)	5.5	17.7
Azimuth direction at the final eighth of the ST-T loop (degrees)	150.0	-153.78
Clinical Information		
Age, years	66-70	81-85
Sex	Female	Female
BMI, kg/m²	22.3	22.6
Resting SBP, mmHg	140	142
Resting DBP, mmHg	70	70
HEART Score	3	3
Peak HR, % of MPHR	106	85

Abbreviations: SE = stress echocardiography; A-ECG = advanced electrocardiography; BMI = body mass index; SBP = systolic blood pressure; DBP = diastolic blood pressure; HR = heart rate; MPHR = maximum predicted heart rate (220 – age)