Gene-environment interaction analysis in atopic eczema: evidence from large population

datasets and modelling *in vitro*.

Marie Standl[1,2]*, Ashley Budu-Aggrey[3]*, Luke J Johnston[4], Martina S Elias[4], S Hasan Arshad[5,6,7], Peter

Bager[8], Veronique Bataille[9], Helena Blakeway[3], Klaus Bonnelykke[10], Dorret Boomsma[11], Ben M

Brumpton[12], Mariona Bustamante Pineda[13,14,15], Archie Campbell[4], John A Curtin[16], Anders Eliasen[10],

João PS Fadista[8], Bjarke Feenstra[8], Trine Gerner[17], Carolina Medina Gomez[18], Sarah Grosche[19],

Kristine B. Gutzkow[20], Anne-Sofie Halling[21], Caroline Hayward[4], John Henderson[3], Esther Herrera-

Luis[22], John W Holloway[6,23], Joukejan Hottenga[11], Jonathan O'B Hourihane[24,25,26], Chen Hu[27,28],

Kristian Hveem[12,29], Amaia Irizar[15,30,31], Benedicte Jacquemi[32], Leon Jessen[10], Sara Kress[33], Ramesh J

Kurukulaaratchy[5,6,7], Susanne Lau[34], Sabrina Llop[15,35], Mari Løset[12,36], Ingo Marenholtz[19,37], Dan

Mason[38], Daniel L McCartney[4], Mads Melbye[39,40,41], Erik Melén[42], Camelia Minica[11], Clare S Murray[16],

Tamar Nijsten[28], Luba M Pardo[28], Suzanne Pasmans[28], Craig E Pennell[43,44], Maria R Rinnov[17,45], Gillian

Santorelli[38], Tamara Schikowski[33], Darina Sheehan[26,46,47], Angela Simpson[16], Cilla Söderhäll[48], Laurent

F Thomas[12,49,50], Jacob P Thyssen[21], Maties Torrent[51], Toos van Beijsterveldt[11], Alessia Visconti [9,52],

Judith M. Vonk[53,54], Carol A Wang[43,44], Cheng-Jian Xu[55,56], Ali H Ziyab[57], UK-Translational Research

Network in Dermatology, BIOMAP consortium, Adnan Custovic[58], Paola Di Meglio[59,60], Liesbeth

Duijts[61,62], Carsten Flohr[63], Alan D Irvine[64], Gerard H Koppelman[54,65], Young-Ae Lee[19,34,37], Nick J

Reynolds[66,67], Catherine Smith[60], Sinéad M Langan[68,69]*, Lavinia Paternoster[3,70]* and Sara J

Brown[4,71]*§.

*authors contributed equally

§corresponding author

Affiliations

1. Institute of Epidemiology, Helmholtz Zentrum München - German Research Center for Environmental

Health, Neuherberg, Germany.

2. German Center for Child and Adolescent Health (DZKJ), partner site Munich, Munich, Germany.

3. MRC Integrative Epidemiology Unit, Bristol Medical School, University of Bristol, UK.

4. Centre for Genomic and Experimental Medicine, Institute of Genetics and Cancer, Crewe Road South,

University of Edinburgh, Edinburgh, UK.

5. Clinical and Experimental Sciences, Faculty of Medicine, University of Southampton, Southampton, UK.

6. NIHR Southampton Biomedical Research Centre, University Hospitals Southampton NHS Foundation Trust,

Southampton, UK.

7. Asthma and Allergy Research Centre, Isle of Wight, UK.

8. Department of Epidemiology Research, Statens Serum Institute, Copenhagen, Denmark.

9. Department of Twin Research and Genetic Epidemiology, King's College London, London, UK.

10. COPSAC, Copenhagen Prospective Studies on Asthma in Childhood, Herlev and Gentofte Hospital,

University of Copenhagen, Copenhagen, Denmark.

11. Vrije Universiteit, Department of Biological Psychology Van der Boechorststraat, Amsterdam, The

Netherlands.

12. HUNT Center for Molecular and Clinical Epidemiology, Department of Public Health and Nursing,

Norwegian University of Science and Technology, Trondheim, Norway.

13. ISGlobal, Institute for Global Health, Barcelona, Spain.

14. Universitat Pompeu Fabra (UPF), Barcelona, Spain.

15. Spanish Consortium for Research on Epidemiology and Public Health (CIBERESP), Madrid, Spain.

16. Division of Immunology, Immunity to Infection and Respiratory Medicine, School of Biological Sciences, The

University of Manchester, Manchester Academic Health Science Centre, and Manchester University NHS

Foundation Trust, Manchester, United Kingdom.

17. Department of Dermatology and Allergy, Herlev and Gentofte Hospital, University of Copenhagen,

Hellerup, Denmark.

18. Department of Internal Medicine, Erasmus MC, University Medical Center Rotterdam, 3000 CA Rotterdam,

The Netherlands.

19. Max-Delbrück-Center (MDC) for Molecular Medicine, 13125, Berlin, Germany.

20. Department of Air Quality and Noise, Division of Climate and Environmental Health, Norwegian Institute of

Public Health (NIPH), Oslo, Norway.

21. Department of Dermatology and Venereology, Bispebjerg Hospital, University of Copenhagen, Bispebjerg

Bakke 23, build. 4, 2400 Copenhagen N, Denmark.

22. Department of Epidemiology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, *MD*

21205, USA.

23. Human Development and Health, Faculty of Medicine, University of Southampton, Southampton, UK.

24. Royal College of Surgeons in Ireland, Dublin, Ireland.

25. Children's Health Ireland, Dublin, Ireland.

26. INFANT Centre, University College Cork, Cork, Ireland.

27. The Generation R Study Group, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The

Netherlands.

28. Department of Dermatology, Erasmus MC, University Medical Center Rotterdam, Rotterdam, The

Netherlands.

29. Department for Research, St. Olavs hospital, Trondheim University Hospital, Trondheim, Norway.

30. Department of Preventive Medicine and Public health, UPV/EHU, Spain.

31. Biogipuzkoa HRI, Donostia, Spain.

32. Univ Rennes, Inserm, EHESP, Irset (Institut de recherche en santé, environnement et travail) - UMR_S 1085,

F-35000 Rennes, France.

33. IUF-Leibniz Research Institute for Environmental Medicine, Düsseldorf, Germany.

34. Department of Pediatric Respiratory Medicine, Immunology, and Intensive Care Medicine, Charité-

Universitätsmedizin Berlin, Berlin, Germany.

35. Epidemiology and Environmental Health Joint Research Unit, FISABIO-Universitat Jaume I–Universitat de

València, Valencia, Spain.

36. Department of Dermatology, Clinic of Orthopedy, Rheumatology and Dermatology, St. Olavs hospital,

Trondheim University Hospital, Trondheim, Norway.

37. Clinic for Pediatric Allergy, Experimental and Clinical Research Center, Charité University Medical Center,

Berlin, Germany.

38. Bradford Institute for Health Research, Bradford Teaching Hospitals NHS Foundation Trust, Bradford,

England, BD9 6RJ, UK.

39. Danish Cancer Institute, Copenhagen, Denmark.

40. Department of Clinical Medicine, University of Copenhagen, Copenhagen, Denmark.

41. Department of Pediatrics, Stanford University School of Medicine, Stanford, USA.

42. Department of Clinical Science and Education, Södersjukhuset, Karolinska Institutet, Stockholm, Sweden.

43. School of Medicine and Public Health, The University of Newcastle, New South Wales, NSW 2308,

Australia.

44. Mothers and Babies Research Program, Hunter Medical Research Institute, New South Wales, NSW 2305,

Australia.

45. Department of Neonatology, Rigshospitalet, University of Copenhagen, Copenhagen, Denmark.

46. University College Cork, College Road, Cork, Ireland.

47. Cork University Hospital, Wilton, Cork, Ireland.

48. Department of Women's and Children's Health, Karolinska Institutet, Stockholm, Sweden.

49. Department of Clinical and Molecular Medicine, Norwegian University of Science and Technology,

Trondheim, Norway.

50. BioCore - Bioinformatics Core Facility, Norwegian University of Science and Technology, Trondheim,

Norway.

51. Area de Salut de Menorca, ib-salut, Balearic Islands, Spain.

52. Center for Biostatistics, Epidemiology and Public Health, Department of Clinical and Biological Sciences,

University of Turin, Turin, Italy.

53. University of Groningen, University Medical Center Groningen, Department of Epidemiology, Groningen,

The Netherlands.

54. University of Groningen, University Medical Center Groningen, Groningen Research Institute for Asthma

and COPD (GRIAC), Groningen, The Netherlands.

55. Centre for Individualised Infection Medicine (CiiM), a joint venture between the Helmholtz Centre for

Infection Research (HZI) and Hannover Medical School (MHH). Hannover, Germany.

56. TWINCORE, a joint venture between the Helmholtz-Centre for Infection Research (HZI) and the Hannover

Medical School (MHH), Hannover, Germany.

57. Department of Community Medicine and Behavioral Sciences, College of Medicine, Kuwait University,

Kuwait.

58. National Heart and Lung Institute, Imperial College London, UK.

59. The Francis Crick Institute, London, UK.

60. St John's Institute of Dermatology, School of Basic & Medical Biosciences, King's College London.

61. Department of Pediatrics, divisions of Respiratory Medicine and Allergology, and Neonatology, Erasmus

MC, University Medical Center Rotterdam, Rotterdam, The Netherlands.

62. Department of Neonatal and Intensive Care, division of Neonatology, Erasmus MC, University Medical

Center Rotterdam, Rotterdam, The Netherlands.

63. Department of Paediatric Dermatology, St John's Institute of Dermatology, Guy's and St Thomas' NHS

Foundation Trust and King's College London, London, UK.

64. Department of Clinical Medicine, Trinity College Dublin, Dublin, Ireland.

65. University of Groningen, University Medical Center Groningen, Beatrix Children's Hospital, Department of

Pediatric Pulmonology and Pediatric Allergology, Groningen, The Netherlands.

66. Translational and Clinical Research Institute, Faculty of Medical Sciences, Newcastle University, Newcastle

upon Tyne, UK.

67. Department of Dermatology and NIHR Newcastle Biomedical Research Centre, Newcastle upon Tyne

Hospitals NHS Foundation Trust, Newcastle upon Tyne, UK.

68. Faculty of Epidemiology and Population Health, London School of Hygiene & Tropical Medicine, London,

UK.

69. St John's Institute of Dermatology, Guy's and St Thomas' NHS Foundation Trust, London, UK.

70. NIHR Bristol Biomedical Research Centre, University Hospital Bristol and Weston NHS Foundation Trust and

University of Bristol, Bristol, UK.

71. Department of Dermatology, NHS Lothian, Lauriston Building, Edinburgh, Scotland, UK.

Abstract

Background

Environmental factors play a role in the pathogenesis of complex traits including atopic eczema (AE)

and a greater understanding of gene-environment interactions (G*E) is needed to define

pathomechanisms for disease prevention. We analysed data from 16 European studies to test for

interaction between the 24 most significant AE-associated loci identified from genome-wide

association studies and 18 early-life environmental factors. We tested for replication using a further

10 studies and *in vitro* modelling to independently assess findings.

Results

The discovery analysis showed suggestive evidence for interaction (p<0.05) between 7 environmental

factors (antibiotic use, cat ownership, dog ownership, breastfeeding, elder sibling, smoking and

washing practices) and at least one established variant for AE, 14 interactions in total

(maxN=25,339). In replication analysis (maxN=252,040) dog exposure*rs10214237 (on chromosome

5p13.2 near *IL7R*) was nominally significant ($OR_{interaction}$=0.91 [0.83-0.99] P=0.025), with a risk effect

of the T allele observed only in those not exposed to dogs. A similar interaction with rs10214237 was

observed for siblings in the discovery analysis ($OR_{interaction}$=0.84[0.75-0.94] P=0.003), but replication

analysis was under-powered $OR_{interaction}$=1.09[0.82-1.46]). Rs10214237 homozygous risk genotype is

associated with lower IL-7R expression in human keratinocytes, and dog exposure modelled *in vitro*

showed a differential response according to rs10214237 genotype.

Conclusions

Interaction analysis and functional assessment provide evidence that early-life dog exposure may

modify the genetic effect of rs10214237 on AE via *IL7R*, supporting observational epidemiology

showing a protective effect for dog ownership. The lack of evidence for other G*E studied here

implies that only weak effects are likely to occur.

Background

Atopic eczema (AE, synonymous with atopic dermatitis or eczema [1]) is a chronic inflammatory skin

and systemic condition affecting approximately 20% of children and 10% of adults in high-income

countries. Eczema is the dermatosis which contributes the greatest number of disability-adjusted life

years worldwide [2] and co-morbid conditions, including asthma and allergies, obesity,

cardiovascular disease, anxiety and depression add substantially to the social, academic,

occupational, and financial impact [3]. Atopic eczema is a heritable trait [4] but the rapid rise in

prevalence in industrialised areas over the past 30 years [3, 5] illustrates the importance of

environmental factors in aetiology. A greater understanding of environmental effects in driving

pathology could facilitate disease prevention.

The European Academy of Allergy and Clinical Immunology published an umbrella review of

systematic reviews in allergy epidemiology and identified a relative lack of research in eczema

genetic epidemiology and environmental effects [6]. The investigation of environmental factors using

observational epidemiology is inherently challenging in the context of AE because there are multiple

confounding factors and possible reverse causation [7]. Genetic studies, however, have made

substantial progress in defining mechanisms in eczema predisposition and pathogenesis, including

skin barrier dysfunction and aberrant immune response [8]. The evidence of individual variation in

susceptibility to environmental allergens and irritants supports the concept of gene-environment

interaction (G*E) [9] playing a role in AE and loss-of-function variants in *FLG* encoding the skin barrier

protein filaggrin have been implicated [10]. Knowledge of genetic risk may therefore provide an

opportunity to identify key environmental effects and clarify important disease biology.

We aimed to investigate evidence of interaction between the most highly significant eczema risk loci

defined by genome-wide association studies [11] and environmental risk factors selected based on

previous literature [7, 10] and importance to patients and carers [12]. We used early-life

environmental exposures (*in utero* and up to the first 12 months of life) to minimise reverse

causation and focus on disease pathogenesis. G*E was tested in cohorts and data from European

populations, in discovery and replication phases as a pragmatic approach to maximise sample size.

Mechanistic assessment was carried out *in vitro* in a skin keratinocyte model to validate the observed

interactions.

Results

Analysis was conducted to assess observational association (of environmental effects) followed by

interaction effect (of environmental and genetic risk factors) in the discovery cohorts; next the

nominally significant findings and those with *a priori* evidence were tested for replication in available

larger cohorts.

Discovery analysis

In meta-analyses of between 1,084 and 22,263 participants (dependent on exposure, **Additional file**

1) we found strong evidence for antibiotic use increasing risk of AE (*in utero* p=0.004, at 6 months

p=0.001 and at 12 months $p=6x10^{-4}$); weaker evidence was found for a protective effect of dog

ownership (p=0.03), protective effect of childhood smoke exposure (p=0.038) and risk effect of NO_2

levels (p=0.035) (M1 models, **Additional file 2**). Little evidence (p>0.05) was found for main effects of

caesarean delivery, cat ownership, breastfeeding, elder siblings, *in utero* smoke exposure, washing

practices at 6 months and 2 years, PM10 exposure and house dust mite exposure at birth or 1 year

(M1 models, **Additional file 2**).

Of the 432 interactions tested (between 24 genetic variants and 18 environmental exposures), we

found no significant interactions that passed multiple testing correction, yet 14 nominally significant

(p_{int}<0.05) interactions (**Figure 1, Table 1**). Of these, 8 interactions indicated a higher genetic risk in

the presence of the exposure (OR > 1) and 6 indicated a higher genetic risk in the unexposed stratum

(OR < 1). Of the 18 environmental exposures tested, the two with the strongest evidence for

interaction with *FLG* null variants were exposure to tobacco smoke between 0 and 2 years

(p_{int}=0.018) and washing practices during the same period (p_{int}=0.045). There was little evidence

(p>0.05) for interactions between *FLG* null variants and other tested exposures, though confidence

intervals for some interaction estimates were wide (**Additional file 2**). Notably, there was little

evidence for interaction between *FLG* null variants and cat exposure (p=0.36), with strong effects of

FLG in both the unexposed and exposed strata.

Sensitivity analyses, additionally adjusting for family history of AE and socioeconomic status, broadly

supported the results of the main analyses (**Additional file 2**), but many of the sensitivity analyses

are based on much smaller sample sizes because of the requirement for data on additional

covariates. There was little evidence of heterogeneity between cohorts (smallest $p_{het=}$0.01) amongst

the 14 reported interactions.

Replication analysis

We took the 14 interactions with nominal evidence forward to replication, but also the exposures

that had prior literature suggesting an interaction with *FLG* null variants (cat, siblings and breast-

feeding [10]). In total, 19 interactions based on 8 different exposures and 10 genetic variants were

included in the replication.

In replication analysis dog exposure and rs1041237 showed evidence for interaction (p=0.025), **Table**

1. In an analysis stratified by dog exposure the T allele increases risk of atopic eczema (OR=1.14, 95%

CI 1.08 to 1.22), but only amongst those who are not exposed to a dog in the family home. In

individuals who are exposed to dog in early life, this variant appears to have little effect (OR=0.98,

95% CI 0.68-1.11 **Figure 2**).

Availability of environmental data for replication varied, with many of our attempted replications of

interactions being insufficiently powered to be conclusive. Washing practices (0-2y) and antibiotic

use *in utero* interactions had only 3 and 4% power respectively (given the interaction effects

observed in the discovery phase, **Additional file 3**). The tobacco exposure in utero interaction only

reached 11% power and the four sibling interactions had between 8 and 37% power (dependent on

variant). The breast-feeding duration interaction only had 4% power in the replication phase and so

we extended the replication analysis to 'ever breastfed' to increase the power to 56%. The

interactions with dog, cat and tobacco smoke exposure 0-2 years were all sufficiently powered (88%,

72-88% and 99%, respectively, **Additional file 3**). The previously reported interactions between *FLG*

null mutations and cat, siblings and ever breastfed had 99% power given their reported interaction

effects (**Additional file 3**).

***In silico* follow-up of rs10214237*dog interaction**

Rs10214237 is an intergenic variant (T>C) on chromosome 5p13.2; this was identified in association

with eczema by a genome-wide association study (GWAS) [11] in which *IL7R* was prioritised as the

likely causal gene based on evidence including eQTL colocalisation in macrophages and monocytes

[13, 14]. The top single nucleotide variant (SNV) at this locus in more recent meta-analysis [13] is

rs10214273, but this variant is in complete linkage disequilibrium with rs10214237 in European

populations (R^2=1, LDLink version 5.6.6, LDPair tool). Global population data from gnomAD shows

ancestral difference in allele frequency, with rs10214237 being more frequent in European and South

Asian populations (MAF 0.28 and 0.20 respectively) compared to African people (MAF 0.07) (1KG

data accessed 10 Jan 2025).

Rs10214237 is within a region of open chromatin in keratinocytes and fibroblasts, but not the

lymphoblastoid cell line GM12878 (UCSC Genome Browser 06 Feb and 27 Nov 2024). Open Targets

V2G analyses confirm *IL7R* as most likely gene affected by this SNV based on pQTL, sQTL and eQTL (06 Feb and 27 Nov 2024). GTEx data show that expression of *IL7R* is higher with T:T genotype in whole blood and cultured fibroblasts and in newly generated data we show that individuals with the T:T genotype have slightly higher *IL7R* mRNA expression in primary human keratinocytes than those with the C:C genotype (**Additional file 4**). Single cell data from the Human Protein Atlas [15, 16] confirms that IL-7R is expressed at protein level in human keratinocytes, in addition to circulating immune cells.

In vitro testing of the effects of dog allergen on human keratinocytes

Human keratinocytes comprise the outermost layer of skin and can therefore represent the first line of interaction in an allergen encounter *in utero* or early life. To further investigate the effect of dog exposure in early life, primary normal human keratinocytes were exposed to clinical-grade dog epithelial extract, a standardised reagent used for allergy testing in the clinic [17]. Dog allergen exposure stimulated an up-regulation in *CXCL8* (IL-8), *CSF2*, *CCL2* and *TNF* mRNA but the atopy-related cytokines *IL33* and *TSLP* mRNA were down-regulated (**Figure 3A**). Network analysis of the proteins encoded by the upregulated transcripts showed significant enrichment for IL-10 signalling (**Figure 3B**, Reactome pathway FDR 7.71e-08) which plays a suppressive role in contact dermatitis and atopic eczema [18]. To test the keratinocyte response more broadly, we used an ELISA panel of 64 cytokine, chemokines and receptors (**Additional file 5**). This confirmed the signature of increased IL-10 signalling (**Additional file 5**).

Next, using primary human keratinocytes of known rs10214237 genotype and focusing on CXCL8 (IL-8), CCL2 and IL-6 as molecules of relevance to IL-7R signalling in epithelial cells, we investigated the effect of dog allergen exposure, with and without IL-7 stimulation (**Figure 3C-E**). There was no difference in expression levels after IL-7 stimulation, but on stimulation with dog extract (or IL-7 plus dog extract), keratinocytes of T:T genotype (homozygous for the eczema-risk allele) showed a greater response than the C:C genotype.

Together these observations provide a possible mechanistic explanation for the finding that the T

allele at rs10214237 increases risk for atopic eczema; the T:T genotype shows greater IL-7R mRNA

expression, but in the context of dog exposure the risk effect is overshadowed by an increase in

cytokines and chemokines in the IL-10 pathway which suppresses eczema to a greater extent in T:T

than C:C individuals.

Discussion

Our collaborative work represents the largest and most comprehensive analysis to date investigating

G*E in atopic eczema, using a systematic approach focussed on the most significant genetic loci and

selected environmental factors. We first meta-analysed data from available observational studies to

test for association and then applied interaction analysis to investigate G*E. Statistical power remains

a limiting factor and the nominal significance level (p<0.05 without correction for multiple testing)

means cautious interpretation is needed. We have identified important negative results as well as

one interaction with functional validation *in vitro* and others that warrant further follow up.

A variety of sources provide evidence that G*E plays a role in the aetiology of atopic eczema. These

include rapidly rising prevalence [5], clinical observation [4], epidemiological studies [10], and *in vitro*

analyses demonstrating molecular effects that include aryl hydrocarbon receptor signalling [19].

Some authors have even stated that 'atopic eczema is an environmental disease' [20]. Our meta-

analysis of observational associations provides evidence that early-life exposure to antibiotics and

NO_2 levels associate with increased risk of AE, whilst early-life exposure to dog or tobacco smoke is

associated with a lower risk of AE in the populations studied. However, these associations may be

affected by bias through confounding and reverse causation.

Statistical interaction analysis indicates that early-life dog exposure may modify the genetic effect of

rs10214237. Functional genetic analyses show an effect mediated via the gene *IL7R* which encodes

the alpha-subunit of the IL-7 receptor. Rs10214237 T:T genotype was associated with an increased

risk of atopic eczema in population as a whole and in the sub-population without dog exposure

(**Figure 2**) consistent with the T:T genotype showing greater IL7R mRNA expression (**Additional file**

4). The IL-7 receptor is a heterodimer composed of IL7R-alpha and IL2R-gamma. It is expressed is

multiple cell-types and tissues, including T-cells, NK-cells, glandular and stratified epithelial cells (data

from Human Protein Atlas [15, 16]). IL7R-alpha also contributes to a heteromeric complex with the

thymic stromal lymphopoietin (TSLP) receptor but our experimental work to test TSLP as an

alternative ligand in keratinocytes was not informative (data not shown) likely, in part, because the

TSLPR is only very lowly expressed in this cell type [21].

Our detailed *in vitro* work focussed on human epidermal keratinocytes as the earliest tissue to

encounter dog allergen in the initiation of atopic disease, *in utero* or early infancy. We have shown

that keratinocytes display a direct response to dog allergen exposure, with down-regulation of IL-33

and TSLP mRNA (both inducers of type 2 immune responses in atopy [22, 23]) and upregulation of a

network of genes encoding chemokines and cytokines of IL-10 signalling (Reactome pathway HAS-

6783783), contributing to the suppression of atopic inflammation [18]. This is consistent with

observational epidemiology showing an apparent protective effect of dog exposure early in life [24]

[25]. Gene ontology analysis of the same network indicates a role in cellular response to

lipopolysaccharide (GO:0071222), likely to reflect a response to gram negative components of the

canine microbiome.

The proposed interaction with genotype was investigated using keratinocytes of known rs10214237

status. Here the T:T genotype showed a greater increase in IL-10 signalling in response to dog

allergen exposure than the C:C genotype, which is consistent with the suppression of atopic eczema

risk on a population level in the dog-exposed T:T individuals, whilst non-dog-exposed T:T individuals

remain at risk of disease. The interaction is analogous to a 17q21*dog interaction demonstrated in

asthma [26] in which the risk of persistent wheeze is attenuated by dog ownership [26]. There is an

interesting parallel in the interaction of rs10214237 with exposure to older siblings, in which the

older sibling abrogates risk effect for rs10214237. We speculate that this may be related to the

increased microbial exposure experienced by an infant with older siblings (or a dog) in the

household, and there is evidence of shared skin and gut microbiome between humans and their

pets[27], but it could also reflect lifestyle choices of dog-owning families and these hypotheses

require further testing.

There are some limitations to this work. The discovery analysis used selected SNVs to represent

known eczema risk loci, rather than conducting a genome-wide interaction analysis. This restricted

approach has been shown to be effective in other traits [28]; it is needed because of power

constraints, even in large population datasets. A post-hoc estimation of statistical power (**Additional**

file 3) showed that our replication sample sizes were insufficient for some interactions. Therefore,

where replication results do not meet our pre-specified significance threshold it is not possible to

definitively *exclude* an interaction, but we report the interaction effect sizes for which we had good

statistical power, to demonstrate the magnitude of interactions which are unlikely to exist, given our

null results (**Additional file 3**). Furthermore, by focusing on selected SNVs within the known AE risk

loci, we acknowledge that there may be loci in which an effect is only apparent in the context of

interaction with an environmental exposure. These would not be detected by our analysis strategy

and genome-wide interaction analysis should be considered in future work if far larger sample sizes

than used here become available. An important limitation to this work is the use of European cohort

data including people of predominantly white ancestry; this reflects the current sparsity of diverse

ancestries in population genetic studies of sufficient size to carry out these analyses. The observed

differences in allele frequency of rs10214237 in African compared to European and South Asian

populations illustrates the limited transferability of this variant effect across population, although

other population-specific variants in the same locus may contribute to similar mechanistic effects.

International efforts are on-going to address this limitation [29], and future G*E studies are needed

to investigate population-specific environmental effects. More detailed sub-phenotyping of AE may,

in the future, reveal that more specific genetic and environmental drivers exist in distinct ancestral or

sub-phenotype groups.

In our previous systematic review focusing on gene-environment interactions with *FLG* null mutations

[10] we found some published evidence for *FLG**environment interactions with exposures including

early-life cat ownership, older siblings, water hardness, phthalate exposure, and prolonged

breastfeeding from the small number of previous studies. The lack of replication of *FLG**cat

ownership interaction in the large well-powered study reported here, and another recent meta-

analysis [30] represents an interesting null finding, contrasting with two small birth cohort studies

([31, 32] n=379 and n=503) which reported p values for interaction <0.01 with evidence for increased

risks of atopic eczema in those with *FLG* null mutations exposed to cat in early life. Evidence for these

G*E interactions came from small numbers of individuals with *FLG* mutation, cat exposure and

development of atopic eczema (five people in one study [31]). We had very good power (99%) for

the interaction magnitude previously reported (OR_{int}=11[31]) and 80% for an interaction as small as

OR_{int}=1.26, suggesting very little evidence in our data for this interaction. We found little evidence for

FLG*breastfeeding, consistent with our systematic review [10], where studies reported no evidence

for interactions with breastfeeding, although an *FLG**breastfeeding duration interaction was reported

from the Isle of Wight cohort [33]. Here, our post-hoc power calculation (**Additional file 3**) showed

adequate power (99%) for the *FLG**breastfed-ever interaction, but low power (<1%) for

FLG*breastfeeding duration analyses, which may explain the discrepancy.

Conclusions

We report observational evidence for an association of atopic eczema with exposure to antibiotics,

NO_2, and tobacco smoke in early life, but the precise nature and mechanisms of action of these

environmental factors on atopic skin inflammation remain unclear. We also detected an

observational association between early life dog exposure and reduction in prevalence of atopic

eczema. Further interaction analysis and functional assessments have provided evidence that dog

exposure reduces the genetic risk effect of rs10214237 in a pathway via *IL7R* and possibly IL-10, to

suppress skin inflammation. There may be an equivalent interaction effect with siblings, but this is

not possible to model *in vitro*. The lack of statistical evidence for other G*E explored in this analysis

suggests that only weak interactions are likely to exist, indicating that on a population level the

interactions tested and found to be null are unlikely to have important contributions to AE

pathogenesis. Therefore further, larger longitudinal studies should focus on alternative mechanistic

questions.

Methods

Aim

This work aimed to investigate evidence of interaction between 24 genetic risk loci for atopic eczema

and 18 early-life environmental effects.

Study design and setting

Genetic risk loci were defined by the 24 top hits at each locus from genome-wide association analysis

[11, 13] and coded for the risk-increasing allele as effect allele (**Additional file 8**). *FLG* null genotype

was coded as presence/absence (0/1) of any of the loss-of-function variants prevalent in the white

European population (R501X, 2282del4, R2447X, S3247X as previously reported [11, 34]).

Environmental exposures were selected on the basis of our recently published literature review [10],

interest from representative of a national eczema support group [12] and refined for pragmatic

reasons, based on data availability.

Genetic epidemiology and interaction analysis was used for discovery and replication. *In vitro*

modelling was performed to independently assess the one G*E effect that showed a nominally

significant interaction in the discovery and replication analyses.

Characteristics of participants

Cohort descriptions

The discovery analysis included 16 population-based cohorts from people of European ancestry (N =

25,339) and a further 10 European population-based cohorts were included in the replication stage

(N = 254,532), giving a maximum total of 279,871 (maxN) in the final meta-analysis (**Additional file**

1A and 1B). Disease status was determined by either parental report or doctor diagnosis for those

who had "ever had eczema". Further details on the phenotype definitions for the included studies

can be found in **Additional file 6**.

Keratinocyte culture and gene expression

Primary human keratinocytes were isolated from normal human skin samples excised during routine

surgical procedures, with patient consent, under governance of the Lothian Bioresource (reference

SR1665). Samples were genotyped for rs10214237 using KASP™ (LGC Genomics, Teddington,

England). IL7R mRNA expression was quantified in 34 keratinocyte samples (3 of C:C genotype, 15 T:C

and 16 T:T) using RT-qPCR. RNA was isolated with TRIzol (15596026, Invitrogen, Carlsbad, USA) and

spin filtration columns using Direct-zol (R2072, Zymo, Irvine, USA). cDNA was prepared using

200ng/ml random primers (48190011, Invitrogen, Carlsbad, USA) with reverse transcriptase using

SuperScript IV (18090050, Invitrogen, Carlsbad, USA). qPCR was carried out using exon-spanning

probes (*IL7R*: HS00902334_m1, Thermo, Waltham, USA) and (*EF1A*: HS.PT.58.24345862, Integrated

DNA Technologies ,San Diego, USA) with TaqMan Universal Master Mix II (4440040, Thermo,

Waltham, USA) and run on a CFX384 PCR Detection System (Bio-Rad, Hercules, USA) using cycling

conditions: 95°C for 10 mins, 40 cycles of [95°C for 15 secs, 60°C for 1 min]. Fold changes in gene

expression were derived via the 2(-Delta Delta C[T]) method, using *EF1A* as the reference gene.

***In vitro* analysis for rs10214237*dog interaction**

To investigate the effect of dog allergen on human keratinocytes, monolayers were treated for 8h

with 10ug/ml dog allergen (Can f 1, catalogue E802, Immunotek, Madrid, Spain). RNA isolation and

RT-qPCR were carried out as above (*CXCL8*: Hs.PT.58.38869678.g, *CSF2*: Hs.PT.58.20138984, *CCL2*:

Hs.PT.58.45467977, *TNF*: Hs.PT.58.45380900, *IL33*: Hs.PT.58.21416460, *EF1A*: HS.PT.58.24345862,

Integrated DNA Technologies, San Diego, USA) and (*TSLP*: Hs00263639_m1, Thermo, Waltham, USA).

Experiments were replicated using keratinocytes from 5-12 independent donors.

To investigate a genotype-specific effect of IL-7 and/or dog allergen stimulation, keratinocytes were

grown to confluency and treated with 100ng/ml recombinant human IL-7 (rhIL-7) (BioTechne,

Minneapolis, USA, catalogue: 207-IL) and 500ng/ml Dog dander (Lofarma, Milan, Italy) for 8 hours.

The carrier solution for dog dander (Lofarma) or 0.002% BSA, used as a carrier protein for rhIL-7 were

negative controls and experiments were conducted in duplicate for each condition.

One-way analysis of variance (ANOVA) with Dunnett's post-hoc test for multiple comparisons was

used to compare samples' means and results displayed showing standard error of the mean (SEM).

Gene ontology, network and pathway analyses were conducted using STRING v12.0.

Statistical genetic analysis

The early-life environmental exposures for investigation included pet ownership for cat and dog

separately, house dust mite exposure, washing practices (to represent environmental irritants),

cigarette tobacco smoking within the household, antibiotic use, environmental pollution, breast

feeding mode of delivery and presence of older siblings. These are listed in **Additional file 7**, with

details of their definition and coding. The exposures were tested for interaction effects with 24 SNPs

previously reported for eczema risk [11, 35] (**Additional file 8**). This involved fitting a statistical model

to include the main effect of the SNP upon eczema (G) (extracted from Paternoster et al, 2015[11]),

the main effect of the environmental factors upon eczema (E), and the product of the SNP effect and

the environmental effect (G*E). Logistic regression models were applied to identify the main effect of

each environmental factor (models M1 to M4, **Additional file 9**), and to test for interaction between

the exposure and each SNP while adjusting for sex (models I1 to I3, **Additional file 9**). Sensitivity

analyses were also performed while adjusting for family history of atopic disease (asthma, eczema or

hay fever) and parental education as a proxy for socioeconomic status (models S1 to S3, **Additional**

file 9).

Analyses were performed separately within each cohort and then combined by performing fixed-

effects meta-analyses. Genetic data was imputed separately for each cohort. Further imputation

details can be found in the **Additional file 6**.

 Power calculation

Posthoc estimates of statistical power were calculated in Quanto (version 1.2.4). These were

informed by effect size estimates from the discovery analyses or previously published studies,

assuming case-to-control ratio of 1:3, and alpha=0.004 in replication analyses (0.05/14 for multiple

testing of 14 gene-environment pairs) (**Additional file 3**).

Abbreviations

ANOVA   Analysis of variance

FLG   Gene encoding filaggrin

G*E   Gene-environment interaction

GWAS   Genome-wide association study

maxN   Maximum number of individuals in the analysis

OR   Odds ratio

SEM   Standard error of mean

SNV   Single nucleotide variant

TSLP   Thymic stromal lymphopoietin

Declarations

Ethics approval and consent to participate

Each contributing cohort has ethical approval for the sharing of anonymised data from study

participants, with their written informed consent.

Consent for publication

The named authors provide consent for publication of this work.

Availability of data and materials

All results supporting the conclusions of this article are included within manuscript and Additional

files. Specifically, full results for all the models tested are given in **Additional files 2 and 10**. Each

cohort contributing to the analysis has their own controlled-access procedures and should be

contacted directly to obtain access to individual level data.

Competing interests

SJB has received research funding (but no personal financial benefits) from the Wellcome Trust

(220875/Z/20/Z), UKRI, Medical Research Council, Rosetrees Trust, Stoneygates Trust, British Skin

Foundation, Charles Wolfson Charitable Trust, anonymous donations from people with eczema,

Unilever, Pfizer, Abbvie, Sosei-Heptares, Janssen, European Lead Factory (which includes multiple

industry partners). SJB, AB-A, KB, ADI, GHK, CS, SML and LP have received funding from the BIOMAP-

IMI consortium (EU H2020 project ref No 821511) which receives support from several

pharmaceutical industry partners. LP has received honorarium payment for a scientific talk on

eczema genetics from LEO Pharma.

GHK reports grants from the Netherlands Lung Foundation, ZON-MW, Ubbo Emmius Foundation,

TEVA the Netherlands, GSK, Vertex, outside the submitted work (money to institution). His institution

received compensation for consultancy or lectures from Astra Zeneca, Boehringer Ingelheim and

Sanofi.

Funding

This project has received funding from the Innovative Medicines Initiative 2 Joint Undertaking (JU)

under grant agreement No 821511(BIOMAP). The JU receives support from the European Union's

Horizon 2020 research and innovation programme and EFPIA. This publication/dissemination reflects

only the author's view and the JU is not responsible for any use that may be made of the information

it contains. M.S. has received funding from the European Research Council (ERC) under the European

Union's Horizon 2020 research and innovation programme (Grant Agreement No. 949906). NJR's

research/laboratory is funded in part by the Newcastle NIHR Biomedical Research Centre (BRC), the

Newcastle NIHR Medtech and In vitro diagnostics Co-operative and the NIHR Newcastle Patient

Safety Research Collaboration and NJR is a NIHR Senior Investigator. LP has received funding from a

MRC Population Health Scientist Fellowship (MR/J012165/1) and receives support from the UK

Medical Research Council Integrative Epidemiology Unit at the University of Bristol

(MC_UU_00011/1, MC_UU_00032/01). SJB holds a Wellcome Trust Senior Research Fellowship

(220875/Z/20/Z). The MAS study up to adolescence was funded by several grants from the German

Federal Ministry of Education and Research (07015633, 07 ALE 27, 01EE9405/5, and 01EE9406). EHL

was supported by a fellowship awarded by MCIN/AEI/10.13039/501100011033 and by "ESF Investing

in your future" [PRE2018-083837]. MAAS was supported by the Asthma UK Grants No 301 (1995-

1998), No 362 (1998-2001), No 01/012 (2001-2004), No 04/014 (2004-2007), BMA James Trust

(2005) and The JP Moulton Charitable Foundation (2004-current), The North west Lung Centre

Charity (1997-current) and the Medical Research Council (MRC) G0601361 (2007-2012),

MR/K002449/1 (2013-2014) and MR/L012693/1 (2014-2018), and MR/S025340/1 UNICORN (Unified

Cohorts Research Network): Disaggregating asthma (2020-2024). This study was delivered through

the National Institute for Health and Care Research (NIHR) Manchester Biomedical Research Centre

(BRC) (NIHR203308). The views expressed are those of the author(s) and not necessarily those of the

NIHR or the Department of Health and Social Care. The PIAMA study has been supported by project

grants from the Netherlands Organization for Health Research and Development; the Netherlands

Authors' contributions

AB-A, PB, KB, DB, SJB, MBP, ACu, CF, JHe, JWH, JO'BH, ADI, GK, PdM, SML, Y-AL, DM, EM, CSM, DMM,

SP, LP, CP, NJR, AS, CS, MS, JPT and CW made substantial contributions to the conception or design of

the work; all authors contributed to the acquisition, analysis, or interpretation of data; AB-A, SJB,

ACu, LD, CF, ADI, GK, SML, Y-AL, LP, NJR, CS and MS drafted the work or substantively revised it.

All authors have approved the submitted version and have agreed both to be personally accountable

for the author's own contributions and to ensure that questions related to the accuracy or integrity

of any part of the work, even ones in which the author was not personally involved, are

appropriately investigated, resolved, and the resolution documented in the literature.

Acknowledgements

We are grateful to all patients and participants in the contributing cohort studies and to the skin

donors and surgical team at the Western General Hospital, Edinburgh, for providing samples of

normal skin tissue for this research; to our UK-TREND collaborators including Mike Cork for

informative discussion; to Heather J Cordell for advice on statistical genetics approaches; to Lisa

Maier for preparing **Figure 1;** and to Dr Lauren Kelly and Ms Lucy Glass for administrative support.

Authors' information

MS, AB-A, LP are experts in statistical genetic analysis of complex traits including AE; SML is a clinical

academic dermatologist with epidemiology expertise; SJB is a clinical academic dermatologist with

expertise in genetic epidemiology and functional genetics. NJR is a clinical academic dermatologist

with expertise in immune mediated inflammatory skin disorders and precision medicine.

References

1.   Johansson S, Bieber T, Dahl R, Friedmann P, Lanier B, Lockey R, Motala C, Ortega Martell J, Platts-Mills

T, Ring J, et al: **Revised nomenclature for allergy for global use: Report of the Nomenclature Review**

Committee of the World Allergy Organization, October 2003. *J Allergy Clin Immunol* 2004, **113:**832-

836.

2.   Mehrmal S, Uppal P, Giesey RL, Delost GR: **Identifying the prevalence and disability-adjusted life**

years of the most common dermatoses worldwide. *J Am Acad Dermatol* 2020, **82:**258-259.

3.   Urban K, Chu S, Giesey RL, Mehrmal S, Uppal P, Nedley N, Delost GR: **The global, regional, and**

national burden of atopic dermatitis in 195 countries and territories: An ecological study from the

Global Burden of Disease Study 2017. *JAAD Int* 2021, **2:**12-18.

4.   Langan SM, Irvine AD, Weidinger S: **Atopic dermatitis.** *Lancet* 2020, **396:**345-360.

5.   Odhiambo JA, Williams HC, Clayton TO, Robertson CF, Asher MI, Group IPTS: **Global variations in**

prevalence of eczema symptoms in children from ISAAC Phase Three. *J Allergy Clin Immunol* 2009,

124:1251-1258 e1223.

6.   Genuneit J, Seibold AM, Apfelbacher CJ, Konstantinou GN, Koplin JJ, La Grutta S, Logan K, Perkin MR,

Flohr C, Task Force 'Overview of Systematic Reviews in Allergy Epidemiology ' of the EIGoE: **Overview**

of systematic reviews in allergy epidemiology. *Allergy* 2017, **72:**849-856.

7.   Rutter CE, Silverwood RJ, Williams HC, Ellwood P, Asher I, Garcia-Marcos L, Strachan DP, Pearce N,

Langan SM, Group IPTS: **Are Environmental Factors for Atopic Eczema in ISAAC Phase Three due to**

Reverse Causation? *J Invest Dermatol* 2018.

8.   Brown SJ: **What Have We Learned from GWAS for Atopic Dermatitis?** *J Invest Dermatol* 2021, **141:**19-

22.

9.   Ottman R: **Gene-environment interaction: definitions and study designs.** *Prev Med* 1996, **25:**764-770.

10.   Blakeway H, Van-de-Velde V, Allen VB, Kravvas G, Palla L, Page MJ, Flohr C, Weller RB, Irvine AD,

570   McPherson T, et al: **What is the evidence for interactions between filaggrin null mutations and**

571   **environmental exposures in the aetiology of atopic dermatitis? - A systematic review.** *Br J Dermatol*

572   2019.

573   11.   Paternoster L, Standl M, Waage J, Baurecht H, Hotze M, Strachan DP, Curtin JA, Bonnelykke K, Tian C,

574   Takahashi A, et al: **Multi-ancestry genome-wide association study of 21,000 cases and 95,000**

575   **controls identifies new risk loci for atopic dermatitis.** *Nat Genet* 2015, **47:**1449-1456.

576   12.   Brown SJ: **Discussion with Eczema Outreach Scotand** 2018.

577   13.   Budu-Aggrey A, Kilanowski A: **European and multi-ancestry genome-wide association meta-analysis**

578   **of atopic dermatitis highlights importance of systemic immune regulation.** *Nat Commun* 2023, **in**

579   **press**.

580   14.   Sobczyk MK, Richardson TG, Zuber V, Min JL, Gaunt TR, Paternoster L, eQtlgen Consortium BCG:

581   **Triangulating Molecular Evidence to Prioritize Candidate Causal Genes at Established Atopic**

582   **Dermatitis Loci.** *J Invest Dermatol* 2021, **141:**2620-2629.

583   15.   Thul PJ, Akesson L, Wiking M, Mahdessian D, Geladaki A, Ait Blal H, Alm T, Asplund A, Bjork L, Breckels

584   LM, et al: **A subcellular map of the human proteome.** *Science* 2017, **356**.

585   16.   **Human Protein Atlas available from** http://www.proteinatlas.org

586   17.   Patel G, Saltoun C: **Skin testing in allergy.** *Allergy Asthma Proc* 2019, **40:**366-368.

587   18.   Boyman O, Werfel T, Akdis CA: **The suppressive role of IL-10 in contact and atopic dermatitis.** *J Allergy*

588   *Clin Immunol* 2012, **129:**160-161.

589   19.   Hidaka T, Ogawa E, Kobayashi EH, Suzuki T, Funayama R, Nagashima T, Fujimura T, Aiba S, Nakayama K,

590   Okuyama R, Yamamoto M: **The aryl hydrocarbon receptor AhR links atopic dermatitis and air**

591   **pollution via induction of the neurotrophic factor artemin.** *Nat Immunol* 2017, **18:**64-73.

20. Luschkova D, Zeiser K, Ludwig A, Traidl-Hoffmann C: **Atopic eczema is an environmental disease.**

Allergol Select 2021, **5:**244-250.

21. Zhong W, Wu X, Zhang W, Zhang J, Chen X, Chen S, Huang H, Yang Y, Yu B, Dou X: **Aberrant Expression**

of Histamine-independent Pruritogenic Mediators in Keratinocytes may be Involved in the

Pathogenesis of Prurigo Nodularis. *Acta Derm Venereol* 2019, **99:**579-586.

22. Ebina-Shibuya R, Leonard WJ: **Role of thymic stromal lymphopoietin in allergy and beyond.** *Nat Rev*

Immunol 2023, **23:**24-37.

23. Liew FY, Girard JP, Turnquist HR: **Interleukin-33 in health and disease.** *Nat Rev Immunol* 2016, **16:**676-

689.

24. Thorsteinsdottir S, Thyssen JP, Stokholm J, Vissing NH, Waage J, Bisgaard H: **Domestic dog exposure at**

birth reduces the incidence of atopic dermatitis. *Allergy* 2016, **71:**1736-1744.

25. Eapen AA, Sitarik AR, Cheema G, Kim H, Ownby D, Johnson CC, Zoratti E: **Effect of prenatal dog**

exposure on eczema development in early and late childhood. *J Allergy Clin Immunol Pract* 2022,

10:3312-3314 e3311.

26. Tutino M, Granell R, Curtin JA, Haider S, Fontanella S, Murray CS, Roberts G, Arshad SH, Turner S,

Morris AP, et al: **Dog ownership in infancy is protective for persistent wheeze in 17q21 asthma-risk**

carriers. *J Allergy Clin Immunol* 2023, **151:**423-430.

27. Lehtimaki J, Sinkko H, Hielm-Bjorkman A, Laatikainen T, Ruokolainen L, Lohi H: **Simultaneous allergic**

traits in dogs and their owners are associated with living environment, lifestyle and microbial

exposures. *Sci Rep* 2020, **10:**21954.

28. Shungin D, Deng WQ, Varga TV, Luan J, Mihailov E, Metspalu A, Consortium G, Morris AP, Forouhi NG,

Lindgren C, et al: **Ranking and characterization of established BMI and lipid associated loci as**

candidates for gene-environment interactions. *PLoS Genet* 2017, **13:**e1006812.

29.   Borrell LN, Elhawary JR, Fuentes-Afflick E, Witonsky J, Bhakta N, Wu AHB, Bibbins-Domingo K,

Rodriguez-Santana JR, Lenoir MA, Gavin JR, 3rd, et al: **Race and Genetic Ancestry in Medicine - A**

Time for Reckoning with Racism. *N Engl J Med* 2021, **384:**474-480.

30.   Thyssen JP, Ahluwalia TS, Paternoster L, Ballardini N, Bergstrom A, Melen E, Chawes BL, Stokholm J,

Hourihane JO, O'Sullivan DM, et al: **Interaction between filaggrin mutations and neonatal cat**

exposure in atopic dermatitis. *Allergy* 2020, **75:**1481-1485.

31.   Bisgaard H, Simpson A, Palmer CN, Bonnelykke K, McLean I, Mukhopadhyay S, Pipper CB, Halkjaer LB,

Lipworth B, Hankinson J, et al: **Gene-environment interaction in the onset of eczema in infancy:**

filaggrin loss-of-function mutations enhanced by neonatal cat exposure. *PLoS Med* 2008, **5:**e131.

32.   Schuttelaar ML, Kerkhof M, Jonkman MF, Koppelman GH, Brunekreef B, de Jongste JC, Wijga A,

McLean WH, Postma DS: **Filaggrin mutations in the onset of eczema, sensitization, asthma, hay fever**

and the interaction with cat exposure. *Allergy* 2009, **64:**1758-1765.

33.   Ziyab AH, Mukherjee N, Ewart S, Arshad SH, Karmaus W, Turati F, Bertuccio P, Galeone C, Pelucchi C,

Naldi L, et al: **Filaggrin gene loss-of-function variants modify the effect of breast-feeding on eczema**

risk in early childhood. *Allergy* 2016, **71:**1371-1373.

34.   Sandilands A, Terron-Kwiatkowski A, Hull P, O'Regan G, Clayton T, Watson R, Carrick T, Evans A, Liao H,

Zhao Y, et al: **Comprehensive analysis of the gene encoding filaggrin uncovers prevalent and rare**

mutations in ichthyosis vulgaris and atopic eczema. *Nat Genet* 2007, **39:**650-654.

35.   Budu-Aggrey A, Kilanowski A, Sobczyk MK, andMe Research T, Shringarpure SS, Mitchell R, Reis K,

Reigo A, Estonian Biobank Research T, Magi R, et al: **European and multi-ancestry genome-wide**

association meta-analysis of atopic dermatitis highlights importance of systemic immune regulation.

Nat Commun 2023, **14:**6172.

Figures and Table

Figure 1. Heatmap to summarise results of interaction analyses

Strength of colour indicates beta in which blue is positive and red is a negative direction of effect; diameter of

circle indicates sample size; 14 nominally significant interactions ($p_{int}<0.05$) are highlighted with black outline;

one association was reported in only one cohort, so was not pursued further.

Figure 2a: Unexposed stratum

Figure 2b: Exposed stratum

Figure 2. Forest plot showing interaction of dog exposure with rs1041237 in exposed and

unexposed strata

Interaction analysis for discovery (N=18,045), replication (N=47,185) and combined meta-analysis (total N=

65,230) show the T allele of rs1041237 increases risk of atopic eczema only amongst those who are not

exposed to a dog in the family home. Full names and study cohort descriptions are given in **Additional file 6**.

Figure 3. *In vitro* testing of the effects of dog allergen on primary human keratinocytes.

(3A) Dog allergen exposure stimulated a reduction in *IL33* and *TSLP* mRNA but upregulation of *CXCL8* (IL-8),

CSF2, *CCL2* and *TNF*; negative indicates keratinocyte media with dog allergen carrier solution; 5-12 donor

isolates shown, bars represent SEM one-way ANOVA, Dunnet post hoc test compared to negative control,

$p < 0.01$, ***$p < 0.001$ ****$p < 0.0001$. **(3B) IL-10 signalling was the most significantly enriched Reactome

pathway (4 out of 45 genes/proteins, FDR 7.71e-08). **(3C-3E)** Effects of IL-7 and dog allergen stimulation on

primary human keratinocytes with different rs10214237 genotypes in which T is eczema risk allele; graphs

represent the mean fold change in cytokine mRNA expression relative to the housekeeping gene *EF1A*, from 4

keratinocyte isolates with T:T genotype and 2 keratinocyte isolates from donors of C:C genotype; untreated

indicates keratinocyte media only and negative is keratinocyte media with dog allergen carrier solution; BSA as

0.0002% included for as carrier protein for recombinant Il-7; two-way ANOVA with Dunnett's post-hoc test,

compared to the negative control, bars represent SEM, *$p < 0.05$, **$p < 0.01$.

Exposure	SNV or gene	Discovery							Replication					p-value combined
		N	N studies	OR	95% CI	p-value	p-value random	p-value heterogeneity	N	N studies	OR	95% CI	p-value	
Antibiotic use in utero	rs10995251	11575	7	1.19	[1.00-1.42]	0.045	0.059	0.34	2666	1	0.88	[0.55-1.41]	0.59	0.09
Cat ownership	rs16948048	14063	10	0.86	[0.76-0.97]	0.012	0.29	0.054	49212	3	1	[0.92-1.08]	0.98	0.17
Cat ownership	rs2227483	14644	11	1.13	[1.01-1.27]	0.037	0.18	0.014	49212	3	1.05	[0.97-1.13]	0.25	0.036
Cat ownership	rs2228145	12994	9	1.14	[1.01-1.29]	0.037	0.037	0.61	49212	3	1	[0.92-1.08]	0.93	0.3
Cat ownership	rs2897442	12702	9	0.87	[0.75-1.00]	0.044	0.11	0.35	49212	3	1.02	[0.94-1.11]	0.59	0.58
Dog ownership	rs10214237	14656	11	0.83	[0.72-0.96]	0.011	0.014	0.37	47185	2	0.91	[0.83-0.99]	0.025	0.0013
Duration of any breastfeeding	rs13015714	14474	9	1.02	[1.00-1.03]	0.049	0.068	0.32	4252	5	0.99	[0.95-1.02]	0.47	0.14
Elder siblings	rs10214237	19155	11	0.84	[0.75-0.94]	0.003	0.003	0.99	5049	4	1.09	[0.82-1.46]	0.55	0.011
Elder siblings	rs10995251	18608	10	0.89	[0.80-1.00]	0.042	0.042	0.44	7529	6	1.03	[0.85-1.25]	0.74	0.11
Elder siblings	rs7146581	19176	11	1.25	[1.10-1.42]	0.00042	0.0031	0.32	7529	6	1.04	[0.84-1.29]	0.73	0.0012
Elder siblings	rs7927894	18263	10	1.13	[1.01-1.26]	0.031	0.031	0.53	7529	6	1	[0.83-1.21]	0.97	0.058
Smoking in household up to 2 years	FLG*	15618	12	1.33	[1.05-1.68]	0.018	0.022	0.36	147880	7	1.01	[0.81-1.25]	0.93	0.096
Smoking in household in utero	rs16948048	16062	11	1.15	[1.02-1.31]	0.028	0.028	0.53	8078	5	1.04	[0.85-1.27]	0.72	0.039
Washing practices up to 2 years	FLG*	6962	3	0.71	[0.51-0.99]	0.045	0.12	0.31	1061	1	1.93	[0.29-12.99]	0.5	0.063

Table 1. Nominally significant interaction results from discovery and replication analyses

Results of testing for interaction between 24 genetic variants and 18 environmental exposures; *combined null genotype for 2 or more loss-of-function mutations in *FLG* as detailed in cohort descriptions (**Additional file 6**); nominal significance defined as unadjusted $p_{int}<0.05$. N, number; OR, odds ratio; p-value indicates significance from fixed effects meta-analysis; p-value random indicates significance from random effects meta-analysis; p-value combined indicates significance from combined fixed effects meta-analysis of discovery and replication data.

List of additional Files

Additional file 1. List of cohorts and available exposure data

1A. Included cohorts and exposure availability at discovery stage.

1B. Included cohorts and exposure availability at replication stage.

Additional file 2. Full results of the discovery analysis

Exposure, environmental exposure; N, number of individuals; N_studies, number of studies;

OR_fixed, odds ratio from fixed effect.

Additional file 3. Estimation of statistical power

Posthoc power calculations performed to facilitate interpretation of negative findings.

Additional file 4. *IL7R* mRNA expression in cells of different rs10214237 genotype

Rs10214237 T:T genotype is associated with higher expression level of *IL7R* mRNA than C:C

genotype; **4A** and **4B**, Screenshots from GTEx Portal

(https://www.gtexportal.org/home/snp/rs10214237 accessed 14/04/2024) showing T:T genotype is

associated with higher *IL7R* mRNA expression in cultured fibroblasts and whole blood; **4C**, shows

higher mRNA expression levels in primary human keratinocytes of T:T than C:C genotype.

Additional file 5. Results of cytokine, chemokine and receptor expression on human primary

keratinocytes following dog allergen exposure

5A. Protein detection by ELISA.

5B. STRING network analysis of upregulated genes.

5C. Cytokine expression from primary human keratinocytes.

5D. STRING network analysis of genes showing no significant change in expression.

Additional file 6. Supplementary methods

Additional file 7. Definition and coding of environmental exposures

Additional file 8. Table of selected SNVs, risk alleles and risk allele frequencies

Additional file 9. Logistic regression models on ever having atopic eczema

Additional file 10. Full results of the replication analysis

Exposure, environmental exposure; N, number of individuals; N_studies, number of studies;

OR_fixed, odds ratio from fixed effect meta-analysis; CI_fixed, 95%-confidence interval from fixed

effect meta-analysis; p_fixed, p-value from fixed effect meta-analysis; OR_random, odds ratio from

random effects meta-analysis; CI_random, 95%-confidence interval from random effects meta-

analysis; p_random, p-value from random effects meta-analysis; p_heterogeneity, p-value from Q-

statistic.