1 **Risk assessment of *Escherichia coli* O157:H7 along the farm-to-fork fresh-cut romaine lettuce**
2 **supply chain**
3

Names and e-mail addresses for all authors
Ece Bulut[1*]   eb643@cornell.edu
Sarah I. Murphy[1]   sim39@cornell.edu
Laura K. Strawn[2]   laurakstrawn@vt.edu
Michelle D. Danyluk[3] mddanyluk@ufl.edu
Martin Wiedmann[4]   martin.wiedmann@cornell.edu
Renata Ivanek[1]   ri25@cornell.edu
11
Author affiliations
[1]Department of Population Medicine and Diagnostic Sciences, Cornell University, Ithaca, NY
14853, USA
[2]Department of Food Science and Technology, Virginia Tech, Blacksburg, VA 24061, USA
[3]Food Science and Human Nutrition Department, University of Florida, Lake Alfred, FL 33850,
USA
[4]Department of Food Science, Cornell University, Ithaca, NY 14853, USA
19
20 **Contact information for corresponding author**
21 eb643@cornell.edu
22 602 Tower Rd, Ithaca, NY 14853

Abstract

Outbreaks of *Escherichia coli* O157:H7 (ECO157) in romaine lettuce remain an ongoing public health concern. Quantitative microbial risk assessment (QMRA) models are key tools for identifying control measures to mitigate foodborne diseases. Here, we introduce a comprehensive QMRA framework along the farm-to-fork fresh-cut romaine lettuce chain, including a novel preharvest difference equation model, to predict annual ECO157 illness cases in the United States and evaluate control strategies. We demonstrated the importance of managing irrigation-related contamination at preharvest to control illness cases. Wildlife intrusions pose lower health risk, followed by runoffs and biological soil amendments of animal origin. When preharvest contamination persists and combines with time-temperature abuses at postharvest, the predicted ECO157 illness cases rise considerably. We showed a broad range of interventions targeting both preharvest and postharvest stages can effectively improve the microbial safety of fresh-cut romaine. The comprehensive practices and interventions explored in this study will aid decision-makers in establishing/enhancing food safety best management practices.

Keywords: Quantitative microbial risk assessment, Escherichia coli O157:H7, lettuce, farm to fork

Main

The number of detected foodborne illness outbreaks linked to leafy greens has increased in the United States (USA) between 1996 and 2016, with *Escherichia coli* O157:H7 (ECO157)-contaminated romaine lettuce emerging as the leading source, followed by other common serotypes such as ECO26 and ECO111 [1]. Between 2015 and 2021, ECO157-contaminated romaine has been implicated in 7 outbreaks in the USA, 6 of which were multistate incidents [2,3]. These outbreaks resulted in 4,274 laboratory-confirmed illnesses, leading to 766 hospitalizations and 11 deaths.

Contamination of leafy greens, including romaine, can occur at any point during preharvest, harvest and postharvest, involving various risk factors [4]. Preharvest risk factors may involve domestic animals in the fields [5,6], wild animals [4,7-9], biological soil amendments of animal origin (BSAAO) amended soil [10], irrigation water [11] and runoff from animal operations [12,13]. Harvest and postharvest risk factors may include human behavior [14], harvesting/production equipment [15,16] and production water [17]. In addition, the lack of processing steps that ensure effective removal or inactivation of ECO157 before consumption [14], coupled with the recent increase in romaine consumption [18], further elevates the risk.

In the USA, 90% of the leafy greens are grown by the members of the California or Arizona Leafy Green Products Handler Marketing Agreements (LGMA) [19,20]. These agreements have established guidelines to minimize contamination during the production and harvesting of leafy greens. The guidelines cover best practices in areas such as personnel training, agricultural

water management, soil amendments, harvest equipment cleaning, packaging, transportation,

produce testing, and prevention of animal intrusion. For example, both LGMAs prohibit the use

of Type B agricultural waters—potentially contaminated with fecal indicators—for overhead

spray irrigation of leafy greens within 21 days of harvest. If Type B water is used, it must be

treated to meet Type A standards (agricultural water unlikely to contain fecal contamination)

and must pass microbiological testing criteria. Additionally, the use of untreated, un-

composted, incompletely composted or non-thermally treated BSAAOs is prohibited. Leafy

greens cannot be produced from fields where such BSAAOs have been applied until one year

has passed. If wildlife intrusion is detected, affected crops must be isolated within a three-foot

radius. Furthermore, the position of animal feeding operations near leafy green farms must be

assessed for any issues that might impact safety. The FDA Food Safety Modernization Act

(FSMA) Produce Safety Rule (PSR) mandates similar standards for agricultural water, soil

amendments, animal management, personnel training and equipment cleaning [21]. While

FSMA PSR does not require microbiological testing of agricultural water [22], it mandates

routine inspections to ensure agricultural water safety and proper maintenance [22]. The

application of raw BSAAOs is regulated by the '90-120-day' rule, which requires that raw

BSAAOs be incorporated into the soil at least 120 days before harvest [23]. Stabilized compost

must meet a ECO157 count of < 0.3 MPN per gram of BSAAO [24]. Produce likely to be

contaminated by animal intrusion should not be harvested [25]. The impact of nearby animal

feeding operations should be evaluated based on their effects on agricultural water, such as

drainage or discharge of untreated water [26].

Quantitative microbial risk assessment (QMRA) models are commonly used to quantify the risk

of microbial contamination or illness associated with exposure to microbial hazards. The

outcome of these models provides valuable information for decision makers for setting

regulatory standards, designing interventions to reduce microbial risks, or assessing the

effectiveness of control measures [27]. Over the years, numerous QMRA models have been

developed for leafy greens contaminated with E. coli strains, including ECO157 [28-41]. These

risk assessments have investigated numerous pathways of E. coli contamination along the farm-

to-fork leafy greens supply chain. However, none had comprehensively evaluated a broader

range of factors related to ECO157 exposure and transmission, particularly for the preharvest

lettuce supply chain. Furthermore, there is no standard QMRA model currently in use by

governmental agencies or industry for ECO157 on lettuce. While some tools, such as FDA-iRISK

[42] and microbial risk assessments [43,44] or guidelines [45] from the USDA (developed over

20 years ago) exist, they do not specifically account for the health risks associated with ECO157

contamination of lettuce or leafy greens. Thus, an updated and comprehensive risk assessment

tool for ECO157 on lettuce is needed to thoroughly evaluate the potential health implications of

a wider range of exposure and transmission factors for ECO157 on lettuce.

We developed a comprehensive QMRA aiming to model ECO157 contamination along the

fresh-cut romaine lettuce production and consumption chain in the USA. Our model

incorporates a broad set of factors, utilizes a novel difference equation modeling approach at

the preharvest stage, and accounts for ECO157 exposure and spread throughout the farm-to-

fork chain, as informed by existing literature and expert elicitation. This QMRA was designed as

a decision support tool for managing various risk factors of ECO157 contamination in the USA

and is publicly available via a GitHub repository

(https://github.com/IvanekLab/Romaine_ECO157_QMRA/).

Methods

Overview of the model

In this study, we adapted elements from QMRAs developed for leafy greens in Spain *[27]*,

Belgium *[39]* and USA *[32,36,38]*. We integrated these QMRAs with a modeling method for

partitioning and mixing processes for microbial organisms in the food supply chain *[46]*, as well

as the concept of non-persister and persister cell state dynamics *[47]* for modeling biphasic

decay of ECO157 on romaine and in soil. We also used the FSMA PSR, California and Arizona

LGMA guidelines, and an expert opinions *[48]* to capture practices specific to the USA. This

resulted in a comprehensive risk assessment model that simulates various field and

processing/handling conditions. We predicted the number of annual illness cases in the USA

attributed to consuming ECO157-contaminated fresh-cut romaine, which is our main model

output. We integrated two models into an exposure assessment: the preharvest model

(including preharvest and harvest sub-models) and postharvest model (including processing-

retail, and consumer sub-models). In the preharvest model, we used a difference equation

model to track the population dynamics of ECO157 in soil and romaine prior to harvest. Figure 1

provides an overview of the QMRA model's structure and Figure 2 describes the parameters

used in the model. The unit of analysis in the preharvest model was a romaine 'batch', but it

becomes a 'package' and then a 'serving' in the postharvest model. We defined a 'lot' as all

romaine packages produced in a day between two cleaning and sanitation shifts. We assumed

that a lot was harvested from a single 1-acre field and processed in a day. Based on this

assumption, each lot contained 12 romaine batches or 26,000 romaine plants. To quantify the

variability and uncertainty of the model outcomes, we evaluated the variability and uncertainty

separately in a one-dimensional Monte Carlo simulation using Latin Hypercube Sampling with

100,000 iterations. In the variability analysis, variable parameters changed while uncertain

parameters remained fixed to their medians—and vice versa for uncertainty analysis. This

approach maintained model interpretability while revealing the distinct impacts of variability

and uncertainty on predictions. This was executed in Microsoft Excel 2016 (Microsoft

Corporation, WA USA), utilizing the @RISK add-on software (version 8.8.1, Lumivero, CO USA).

Model validation was carried out by comparing the predicted annual illness cases with the data

from existing surveillance studies [49,50]. Expert elicitation, along with sensitivity and

uncertainty analyses were conducted to guide the selection of promising risk management

strategies and inform parameter choices in scenario analyses. Regression trees were used in the

preharvest and postharvest models to identify interaction of factors that lead to ECO157

contamination in romaine and illness cases. The following sections detailed the baseline model

structure.

Preharvest model

The preharvest model included growing and harvesting romaine in an open field. We

considered a square field measuring approximately 63 m by 63 m (equivalent to 1-acre) for the

cultivation of romaine. A total of 26,000 romaine plants were harvested from the modeled 1-

acre field in a day. Each plant was measured 25cm in width, length and height, and weighed

300g *[51]*. The allocated area for each individual plant within the field was 30cm by 50cm *[36]*.

We assumed continuous romaine production throughout the year, with daily harvests,

occurring in one of three distinct geographical locations: (i) the Coastal Region and Central

Valley in CA (CC), (ii) the Desert Region (Imperial) in CA (DR), and (iii) the Yuma Region in AZ (Y).

These locations were selected since they are the top leafy green producers in the USA (CA

accounts for 70%, and AZ for 30% of US lettuce production) *[52-54]*. The choice of location

affected several model parameters, such as the time from seed planting to harvest, and the

type of irrigation systems used in the fields (Figure 2, Supplementary Tables S1, S2 and S3).

Additionally, the month of harvest affected the parameters related to weather conditions and

survival of ECO157 (Figure 2, Supplementary Table S3).

Contamination of romaine in the field before harvest was attributed to four sources based on

the available information in the literature and the expert elicitation approach: (i) runoff from a

neighboring dairy farm, (ii) intrusion by feral swine, (iii) irrigation water, and (iv) the application

of BSAAO (i.e., treated/composted/processed dairy manure, which may include incompletely

treated/composted/processed dairy manure). All calculations were conducted for a randomly

selected 1/12 portion of the modeled field (i.e., 1/12 lot), which we refer to as the 'batch area',

containing 2,125 romaine plants that weigh 637.5kg at harvest. The plants were grown and

harvested as a batch unit and population dynamics of ECO157 were tracked for the batch. The

size of the batch area enabled the development of a more realistic model by accounting for the

risk factors associated with the exposure and spread of ECO157 throughout the farm-to-fork

chain. For example, we assumed that only the batch area closest to the edge of the field (and

thus to a hypothetical neighboring dairy farm) would be impacted by dairy farm runoff. Due to

the rapid decay of ECO157 on field-grown romaine, we adopted a 14-day 'countdown' period,

representing the final 14 days before harvest, during which we tracked the introduction of

ECO157 to the romaine in the field [39]. The duration of this period was chosen assuming that

furrow irrigation happened every 10 days on average and that an average 'waiting time' of 4

days was used before harvest (as explained in the later text, overhead spray and furrow

irrigation cease during the waiting time). Throughout this countdown period, we tracked

ECO157 counts on day t on both the romaine (LN_{t+1}; CFU/romaine batch) and in the top 1cm

layer of soil of the batch area (referred to as 'soil batch'; SN_{t+1}; CFU/soil batch) using a system

of coupled difference equations (1) - (6) coded within the QMRA in @RISK (notations were

explained in the following paragraphs and also in Supplementary Tables S4 and S5):

$LN_{t+1} = LN_t + Irr_{L_t} + Pig_{L_t} + Runoff_{L_t} + \left(R_{splash_t} + I_{splash_t}\right) \times (SN_t + SP_t) -$

$LNtoLP_t - LNdecay_t$ \hfill (1)

$LP_{t+1} = LP_t + LNtoLP_t - LPdecay_t$ \hfill (2)

$C_{L_{t+1}} = LN_{t+1} + LP_{t+1}$ \hfill (3)

$SN_{t+1} = SN_t + Irr_{S_t} + Pig_{S_t} + Runoff_{S_t} - \left(R_{splash_t} + I_{splash_t}\right) \times SN_t - SNtoSP_t -$

$SNdecay_t$ \hfill (4)

$$SP_{t+1} = SP_t + SNtoSP_t - \left(R_{splash_t} + I_{splash_t}\right) \times SP_t - SPdecay_t \qquad (5)$$

$$C_{S_{t+1}} = SN_{t+1} + SP_{t+1} \qquad (6)$$

where $t = 0,1,2,3,\dots,13,14$ were days of the countdown period (i.e., at t=0 there were 14

days until harvest). If in a time step, the calculated LN_{t+1}, LP_{t+1}, SN_{t+1} or $SP_{t+1} < 0$, we set

the corresponding LN_{t+1}, LP_{t+1}, SN_{t+1} or SP_{t+1} to 0.

We adopted the concept of non-persister and persister cell state dynamics to provide a more

accurate die-off mechanism for ECO157 [55], reflecting the well-known biphasic decay

phenomenon (i.e., fast initial decay followed by slower decay) of the ECO157 cells on leafy

green and in soil environments [56-59]. In Equation (1), LN_{t+1} represents non-persister ECO157

count in romaine batches on day t+1. It was determined by the previous day's count in romaine

batches (LN_t); accumulation through irrigation (Irr_{L_t}), runoff ($Runoff_{L_t}$), intrusion by feral

swine (Pig_{L_t}), rain splash ($R_{splash_t} \times (SN_t + SP_t)$) and irrigation splash ($I_{splash_t} \times (SN_t +$

$SP_t)$); removal of cells that switched from non-persister to persister state ($LNtoLP_t$) and daily

reduction on romaine ($LNdecay_t$). Equation (2) describes the persister ECO157 count in

romaine batches on day t+1 (LP_{t+1}). Total ECO157 count (non-persister and persister cells) in

romaine batches on day t+1 ($C_{L_{t+1}}$) was calculated in Equation (3). Equations (4), (5) and (6)

estimated the total ECO157 count (non-persister and persister) in the soil batch on day t+1

($C_{S_{t+1}}$) and were similar to the calculations for the corresponding romaine batch. It was

assumed that the cross-contamination of ECO157 cells from soil to romaine via splash events

triggers an adaptation process, resulting in a transformation of any transferred cells from a

persister state to a non-persister state when they land on romaine leaves *[47]*.

The ECO157 counts in the romaine batch ($C_{L_{14}}$) and soil batch ($C_{S_{14}}$) at *t*=14 were used in a

harvest sub-model (explained below) to calculate the preharvest outcome: the ECO157 count in

the harvested romaine batch (C_{L_h}). From there, the ECO157 prevalence (P_{Lh_f}) and the ECO157

count in harvested romaine batches (C_{Lh_f}) were calculated in one simulation (one simulation

consists of 100,000 iterations) and used as inputs in the postharvest model. Specifically, P_{Lh_f}

was derived by counting the number of iterations out of 100,000 where the ECO157 count on

the harvested romaine batch was at least one (where $C_{L_h} \geq 1$ CFU/romaine batch). The P_{Lh_f}

was established as a Triangular distribution, where the mean of the distribution corresponded

to the predicted preharvest prevalence in the simulation with 100,000 iterations mentioned

above. The distribution's minimum and maximum were set at half and twice of the predicted

prevalence, respectively. The C_{Lh_f} was derived by fitting a distribution around the predicted

ECO157 counts in contaminated harvested romaine batches (where $C_{L_h} \geq 1$ CFU/romaine

batch) using @RISK software. The distribution that best fits the data was determined using

Akaike Information Criterion, Bayesian Information Criterion and Average Log-Likelihood

methods (Supplementary Table S2).

Accumulation and loss of ECO157 contamination in soil batch (C_S)

We assumed that irrigation water, BSAAO, swine, and cattle feces were the only sources of

ECO157 contamination for the modeled field, explained in the following sections.

BSAAO Application (accumulation in soil): We modeled ECO157 prevalence (P_{man}) and count

(C_{man}) in BSAAO by accounting for the reports of ECO157 in stacked/piled manure from dairy

farms in the USA *[60]*. The maximum ECO157 count in BSAAO was the maximum ECO157 count

found on raw dairy feces *[36]*. This represented an unintentional application of an incompletely

treated BSAAO, though the application probability of such BSAAO was low. The timing of

BSAAO application (D_{man}) was set on the same day of planting romaine seeds. The number of

days from planting romaine seeds to harvest ($Daytogrow$) was set between 65 and 130 days

depending on the field location (CC, DR or Y regions) *[48]*. We considered a constant daily decay

of ECO157 among both non-persister and persister cells in BSAAO (R_{dman} and R_{dman_P}), from

the day of BSAAO application to the time of harvest *[61,62]*. A total of 10% of the simulated

fields received BSAAO application (Pr_{man}) (BSAAO may be incompletely treated), assuming that

those were organic operations that used BSAAO as nitrogen source. The BSAAO application

(M_{man}) ranged from 5 to 15 tons per acre *[63]*. When applying BSAAO, we incorporated a

dilution factor (DiF_{man}) assuming a uniform distribution of BSAAO across the tilled column of

the soil batch. This approach considered that a gram of BSAAO has a volume (Rh_{con}) of 2.5cm^3

[64] and was assumed to be uniformly distributed across the topsoil (Dh_{con}) up to 15 to 20cm

depth *[65]*. In this and other instances of using a dilution factor in the soil column in our model,

the assumption of a uniform distribution of ECO157 within the top layer of soil batch was

justified since the unit of analysis of romaine contamination was the whole batch. For example,

transfer of 100 ECO157 cells from one contaminated spot of soil onto a single romaine plant

(which becomes contaminated with 100 ECO157) and transfer of 1 ECO157 cell from each of

100 different contaminated soil spots resulting in 100 plants contaminated with 1 ECO157 each,

both resulted in the same contamination at the batch level (i.e., 100 ECO157 in a given romaine

batch). The estimation of the initial ECO157 count in soil batch (C_{S_0}) is provided in

Supplementary Table S4.

Runoff from a neighboring dairy farm (accumulation in soil) ($Runoff_{S_t}$): We adapted the

runoff contamination component from Mishra et al.'s (2016) QMRA model [36]. Runoff events

were considered to occur on rainy days only (at a probability $1 - Bs_t$; a day could either be

sunny or rainy). They lead to contamination of batch area at the edge of the field, determined

randomly by the '$edge$' parameter. We assumed that 5 to 15kg of dairy cattle feces (M_{cow})

could enter the batch area during a runoff event, where the quantity of fecal material entering

the batch area was proportional to the monthly average precipitation of the field location

($Prec_{norm_m}$). We incorporated ECO157 count (C_{cow}) and month-specific ECO157 prevalence

(P_{cow_m}, Supplementary Table S3) in dairy cattle feces from Cooley et al. (2013) [8]. The spread

of runoff across the $edge$ batch area affected both romaine batch and soil batch, determined

by the 'Pr_{Ltos}' parameter. This parameter was a proportion describing the area a romaine plant

occupies relative to the soil area allocated for plant growth. The runoff in soil was uniformly

distributed and spread into the soil. The ECO157 in runoff was spread to a soil depth of 1cm by

a dilution factor (DiF_{cow}). The estimation of the term $Runoff_{S_t}$ in the model (1) - (6); i.e., the

ECO157 count in soil batch added via runoff on day t, is provided in Supplementary Table S5.

Intrusion by feral swine (accumulation in soil) (Pig_{S_t}): Wildlife intrusion in our model was

integrated using the risk model developed by Mishra et al. (2016) *[36]*. A contamination event

was dependent on the probability of feral swine intruding the field ($Pr_{pig_{count_t}}$), derived from

reported feral swine population density in Salinas Valley, CA (~ 5 swine/km^2) *[6]*. In the event of

an intrusion, we assumed that between 1-5 adult swine would enter the field and defecate, as

they could be found individually and in family groups in CA *[66]*. In $Pr_{pig_{count_t}}$, the probability of

intrusion by one animal was the highest, and the probability of intrusion by more animals

decreased progressively. The fecal quantity found on the field after the defecations (M_{pig}) was

estimated by the daily fecal production reported per adult swine ($Mdef_{pig}$) *[67]*. In addition, we

factored in a defecation frequency of 4 times/swine/day ($Tdef_{pig}$) *[68]* and assumed at least

one of those defecation events occurred in the modeled field ($Ndef_{pig}$). Feral swine could

defecate on romaine plants or field soil, determined by the 'Pr_{Ltos}' parameter. When feces fall

on soil, we assumed the ECO157 in fecal material was spread evenly within the top 1cm of the

soil batch, which we modeled using a dilution factor (DiF_{pig}). The ECO157 count (C_{pig}) and the

ECO157 prevalence (P_{pig_m}) in swine feces were modeled based on the data provided by Cooley

et al. (2013) *[8]*. The term Pig_{S_t} in the model (1) - (6); i.e., ECO157 count in soil batch added via

feral swine on day t, was estimated as shown in Supplementary Table S5.

Irrigation (accumulation in soil) (Irr_{s_t}): Contamination of soil by irrigation occurred through

contaminated water that was delivered to the field and not captured by the romaine plants on

the field. The ECO157 prevalence ($P_{irr} = P_{irr_W}$ or P_{irr_S}) and count ($C_{irr_t} = C_{irr_W}$ or C_{irr_S}) in

the irrigation water depended on the water source: surface or well (ground) waters. Water

source was selected based on its respective frequency of use in a field location [69]. We

assumed the value for the ECO157 prevalence in well water (P_{irr_W}) as 0.1%, given the limited

literature on well waters, and considering that well waters are generally free of ECO157. Our

scenario analysis in Supplementary Table S6 showed that decreasing this prevalence tenfold—

to 0.01%—resulted in a 20.7% decrease in predicted illness risk compared to the baseline

model. This finding suggests that ECO157 levels in well waters could have a modest yet

noteworthy impact on lettuce safety. ECO157 prevalence in surface water (P_{irr_S}) was based on

the ECO157 survey on surface waters in Monterey County, CA [8]. The ECO157 counts in

surface water included two parts: (i) the generic *E. coli* count and (ii) a factor that converts the

generic *E. coli* count to that of ECO157 [34]. The ECO157 count in well water (C_{irr_W}) was the

LGMA testing limit of generic *E. coli* for Type A agricultural waters, set at 10MPN/100ml [19,20].

For consistency across model calculations, we assumed that the most probable number (MPN)

was equivalent to colony forming units (CFU) [70]. The generic *E. coli* count part of the surface

water (C_{irr_S}) was represented as a PERT distribution, with the minimum and most likely values

taken from the standard testing limits set by the LGMAs for Type B agricultural waters used in

drip and furrow irrigation systems [19,20]. The maximum value of the PERT distribution was

based on the highest generic *E. coli* count observed in surface waters from CA region [71].

Starting in 2023, the LGMAs implemented more stringent testing limits for overhead spray

irrigated surface waters, setting the threshold at 235MPN/100ml generic *E. coli* for any single

sample. However, we applied the less stringent testing limit historically used by LGMA

(576MPN/100ml) in the baseline model to account for agricultural waters in poorer conditions

and to reflect the practices more relevant to the period for which the model validation data

were available. However, our scenario analysis in Supplementary Table S6 predicted that

adopting the more stringent testing limit led to 31.1% reduction in predicted illness risk

compared to our baseline model, suggesting that the 2023 California and Arizona LGMAs may

have a positive impact on leafy green safety. Our model included different irrigation systems

(f_{irr} = overhead spray (O), furrow (F), or drip (D)) based on the field location (Figure 2,

Supplementary Table S1). Each system delivered varying volumes of irrigation water per

irrigation event (g_{irr}), considering that most of the water (i.e., 51% of the total volume

required for romaine growth) was applied in the last 30 days before harvest (W_O, W_F, W_D) [48].

The volume of water captured by a romaine batch (v_{irr}) also differed for each system [27,72].

Overhead spray and furrow irrigation only occurred on sunny days (Bs_t) [73], and differed in

frequency: Overhead spray, applied once every 2-3 days, was reflected in probability Pr_O

($fO_t = Binomial(1, Pr_O)$) and furrow was applied about once every 10 days, reflected in

probability Pr_F ($fF_t = Binomial(1, Pr_F)$). Drip irrigation occurred every day ($fD_t = Pr_{D_t} = 1$)

regardless of sunny or rainy weather. We used a dilution factor (DiF_{irr}) to simulate an even

distribution of ECO157 introduced by irrigation across the field and spread of cells to a depth of

1cm. To align with typical harvest practices in the USA [74], we introduced a waiting period

ranging from 2 to 8 days before harvest ($BwaitP_t$) and terminated overhead spray and furrow

irrigation during the selected waiting time. Shallowly buried drip lines were assumed for drip-

irrigated fields, making them unaffected by the waiting period. The term Irr_{s_t} in the model (1) -

(6), the ECO157 count in soil batch added via irrigation water on day t, is provided in

Supplementary Table S5.

Soil splash via rain or irrigation (i.e., loss from soil) $(R_{splash_t}, I_{splash_t})$: The effect of irrigation

and rain splash in the field was based on the QMRA model developed by Allende et al. (2018)

[27]. Accordingly, splash of soil (and ECO157) to romaine would be facilitated either via

irrigation water on a sunny day or rainwater on a rainy day. The maximum quantity of soil

splashed by rain was assumed to be equal to that splashed by overhead spray irrigation.

However, the quantity of soil splashed by rain was also proportional to the standardized

monthly precipitation level ($Prec_{norm_m}$), reflecting the intensity of rain. We assumed that the

quantity of soil splashed on sunny days varied by different irrigation systems, with overhead

spray irrigation had the potential of causing the most amount soil to splash and drip irrigation

causing the least (M_{irr}). The amount of soil splashed during irrigation was also affected by the

intensity of irrigation systems, where each system assumed to mimic an intermediate rainfall,

equal to the median of the standardized precipitation $\left(Prec_{norm_m}\right)$. In case of a splash, a

fraction of ECO157 in soil would transfer onto the romaine (LsR during rain or LsI during

irrigation, Supplementary Table S4) via transfer of contaminated soil *[75]*. The terms for

modeling rain (R_{splash_t}) and irrigation splash (I_{splash_t}) in the model (1) - (6) were derived as

shown in Supplementary Table S5.

Daily decay of ECO157 (loss from soil) $(SNdecay_t, SPdecay_t)$: Daily inactivation of persister

and non-persister cells in soil was constant. We used four different decay rates to represent

non-persister *[36,61]* and persisters' *[62]* decay in BSAAO-amended soil and soil with no BSAAO

application. We used slower inactivation in BSAAO amended soil (R_{dman} for non-persisters and

R_{dman_P} for persisters) and faster inactivation in soil with no BSAAO application (R_{ds} for non-

persisters and R_{ds_P} for persisters). Persister cells decayed at a rate 10 times slower than that of

their non-persister counterparts, based on the decay curve provided in Chen (2020) for ECO157

persisters in soil *[62]*. Decay parameters in soil ($SNdecay_t$, $SPdecay_t$) in the model (1) - (6) are

explained in Supplementary Table S5.

Switch to persister cells (no change in the cell count in soil) ($SNtoSP_t$): A small fraction of

non-persister ECO157 cells were transitioned to persister cells on each day in soil *[47,76,77]*.

Persisters coexisted with non-persisters and comprised up to 0.01% of the population in soil

environments *[76,77]*. The cells could also switch between the persister and non-persister

states *[78]*. Although the characteristics and the formation mechanisms of persisters are

complex *[78]*, our model focused only on their enhanced ability to survive, which facilitated

modeling biphasic decay. While the direct estimates for the switch rates in BSAAO-amended

soil were unavailable, we used the study by Hofstrenge et al. (2013) *[77]*, which indicated a

strong one-to-one correlation between switch rates ($NtoP_{man}$, $NtoP_S$) and the fraction of

persisters ($Pmax_{man}$, $Pmax_S$). We assumed that the persister state was more prevalent in

BSAAO-amended soil batches ($NtoP_{man}$). This assumption was based on the observation that

pathogens regrew and recovered to a culturable state months after being introduced into soils

through dairy BSAAO application *[79]*. This delayed recovery might suggest the initial presence

of persister cells within the BSAAO that was applied on the field *[80]*. Details on the switch

terms ($SNtoSP_t$) in the model (1) - (6) are provided in Supplementary Table S5.

Accumulation and loss of ECO157 in romaine batches

Given the rapid decay of ECO157 on field romaine [27,39,81], we assumed that the initial

ECO157 count on the field romaine was negligible one day prior to the start of the 14-day

countdown period ($C_{L_0} = 0$). However, the sensitivity of model predictions to this assumption

was tested in scenario analysis (see Supplementary Table S7).

Irrigation (accumulation on romaine): Contamination of romaine through irrigation followed a

similar mechanism to that described for soil contamination. The volume of irrigation water

captured by the romaine batches (v_{irr}) transferred a fraction of ECO157 cells from water to

the romaine plants (Tr_{ecirr}). The total volume of water delivered to the batch area (g_{irr}) was

either captured by romaine or spread into the soil. Thus, we modelled the fraction of irrigation

water delivered to the soil $\left(1 - \frac{v_{irr}}{g_{irr}}\right)$ and romaine ($\frac{v_{irr}}{g_{irr}}$). The term (Irr_{L_t}) describing the

irrigation contamination in the model (1) - (6) is explained in Supplementary Table S5.

Runoff from a neighboring dairy farm and intrusion by feral swine (accumulation on romaine)

(Pig_{L_t} and $Runoff_{L_t}$): The effect of wildlife and runoff contamination on romaine was

modelled mechanistically similar to their effect on soil detailed previously. The difference was

the proportion of feces/runoff that was spread in either the soil or on romaine, as determined

by the parameter 'Pr_{Ltos}' (for soil: $1 - Pr_{pigLtos}$, and for romaine: Pr_{Ltos}). The feces/runoff

was modelled as spreading uniformly through the romaine batches. The terms Pig_{L_t} and

$Runoff_{L_t}$ describing the ECO157 count in romaine batches via feral swine intrusion and runoff

events in the model (1) - (6) are explained in Supplementary Table S5.

Soil splash via rain or irrigation (accumulation on romaine)(R_{splash_t}, I_{splash_t}): The daily loss

of ECO157 from the soil due to rain or irrigation splash was applied to a given romaine batch,

resulting in ECO157 accumulation on romaine batches. The transferred ECO157 cells were

distributed uniformly in a given romaine batch (Supplementary Table S5).

Daily decay of ECO157 (loss from romaine) ($LNdecay_t$, $LPdecay_t$): For non-persister ECO157

decay on romaine ($LNdecay_t$, Supplementary Table S5), the daily decay rate (R_{dL}) followed the

month-specific solar decay pattern (R_{sun_m}) throughout the year [27]. Specifically, a slower

decay rate, reflecting winter conditions, was applied from December to April (for production in

Y and DR regions), while a faster decay rate, reflecting summer conditions, was implemented

from May to October (for production in Y region). Additionally, the parameter 'h_{sun_m}', which

indicated the average daily sunlight hours in a given month, was determined based on weather

data specific to both field location and the time of romaine harvest. The decay for persisters

(R_{dL_P}) was assumed 10 times slower than the non-persisters ($LPdecay_t$, Supplementary Table

S5) [47,62].

Switch to persister cells (no change in the cell count on romaine) ($LNtoLP$): The switch rate

from non-persister to persister cells on romaine ($LNtoLP_t$, Supplementary Table S5) was

constant and established based on the data from the study by Munther et al. (2020) [47].

Harvest (CC_{blade}): After modeling the population dynamics of ECO157 in soil and on romaine

plants during the 14-day countdown period (equations (1) - (6)), we proceeded with modeling

the harvesting process, which accounted for cross-contamination of romaine from soil caused

by contaminated coring knife blades *[15,38]*.

Postharvest model

The bacterial population dynamics outlined by Pang et al. (2017), Danyluk and Schaffner (2011)

and Nauta (2005) served as the baseline for the postharvest practices (i.e., cold storage prior to

processing, chlorine wash, retail, transportation and consumer stages) *[32,38,46]*.

Cold storage before chlorine wash (CC_{coldst}): The harvested romaine was transported to cold

storage, where it was kept for a period ranging from 1 to 72 hours (t_{coldst}). In this stage, we

used 5°C as the minimum temperature for ECO157 growth. Time spent (t_{coldst}) above 5°C

increased ECO157 counts (growth rate, G_{coldst}) and time spent ≤ 5°C decreased ECO157 counts

(decay rate, D_{coldst}) *[38,82]*. Cold storage temperature was assumed to follow a triangular

distribution centered around the standard refrigeration temperature, allowing inadequate

storage ($T_{endcoldst}$). We assumed that the temperature of romaine at the time of harvest

($T_{harvest}$) was equivalent to the air temperature during the harvesting process, under the

assumption that harvesting occurs early in the morning *[83]*. Accordingly, we used field

location- and month-specific weather data for determining the harvesting temperature.

CC_{coldst} in Supplementary Table S2 represented the change in ECO157 count on a given

romaine batch during cold storage.

Chlorine wash ($CC_{w_{agent}}$): The washing step was assumed to reduce ECO157 count by 0.52

$Log_{10}CFU/g$ romaine on average, with a range of reduction between 0.01 $Log_{10}CFU/g$ romaine

and 1 $Log_{10}CFU/g$ romaine (Supplementary Table S2) [84].

Partitioning of the romaine batches into packages (Sc_{clstrb}, $Cluster_{parb}$): The partitioning of

a chlorine-washed romaine batches (C_{wash}) into individual packages included potential

clustering of ECO157 cells (or aggregation of cells) [46,85]. To estimate the ECO157 count

within a contaminated romaine package (C_{pack}), we used the following equations:

$$C_{pack} = Log_{10}\big(Binomial\big(C_{wash},\ P_{contpack}\big) + 1\big) \qquad (7)$$

where

$$P_{contpack} = \frac{1 - Sc_{clstrb}}{A_{pack}} + Sc_{clstrb} \times Beta(Cluster_{parb}, Cluster_{parb} \times (A_{pack} - 1)))$$

$$P0_{pack} = EXP\left(\frac{\Gamma\big(Cluster_{parb} \times A_{pack}\big) \times \Gamma\big(Cluster_{parb} \times (A_{pack}-1) + C_{wash}\big)}{\Gamma\big(Cluster_{parb} \times (A_{pack}-1)\big) \times \Gamma\big(Cluster_{parb} \times A_{pack} + C_{wash}\big)}\right) \qquad (8)$$

Here, $P_{contpack}$ is an intermediary parameter used to determine the ECO157 count within a

given package, accounting for the presence of clustering (Sc_{clstrb}; set to 0 when clustering was

absent and 1 when present) and the degree of clustering when it occurred (with maximum

clustering when $Cluster_{parb} = 0$, decreasing as $Cluster_{parb}$ approaches positive infinity). In

the baseline model, we assumed no clustering ($Sc_{clstrb} = 0$), and as a result, the ECO157 cells

from the romaine batches were evenly distributed to each package ($\frac{1}{A_{pack}}$). The '$P0_{pack}$'

parameter in Supplementary Table S2 (equation (8)) determined the probability of an

uncontaminated package [46]. We derived the probability of a package being contaminated as

$(1 - P0_{pack})$.

Retail storage, transport from retail to home, and consumer (CC_{retail}, CC_{trspt}, CC_{home}): We

used the same growth and decay approach implemented in cold storage process to model retail

storage, transport from retail to home, and consumer steps (i.e., home storage) albeit with

stage specific time and temperature parameters [38,81]. The values of the time and

temperature parameters were previously parameterized by Pang et al. (2017) [38].

Partitioning of the romaine package into servings (Sc_{clstrp}, $Cluster_{parp}$): We adopted a

serving size of 85g for romaine (Ser) based on reference serving size provided by USDA (2015)

[86]. To distribute ECO157 cells from romaine packages to servings, we followed the same

approach used for partitioning romaine batches to packages (no partitioning at the baseline

model, Equations (7) - (8)) [46].

Dose-response

Following the partitioning process, the ECO157 count (C_{serW}) and prevalence (P_{ser}) in romaine

servings were estimated. The C_{serW} was used in a Beta-Poisson dose-response model ($Pill$)

(Supplementary Table S2) [32,87]. We estimated our main outcome, the predicted number of

illness cases per year in the USA from consuming ECO157-contaminated romaine servings

($Ncase_{US}$), by using the dose-response model along with the number of contaminated servings

consumed annually in the USA ($Ncsc_{US}$).

Expert elicitation: Several parameters across the farm-to-fork chain, as well as specific

interventions for the scenario analysis, were determined by using the input from study authors

and industry experts who served on an advisory council for the USDA-NIFA-SCRI grant #2019-

51181-30016. Three informal interviews were conducted with the advisory council members,

who were from produce production and processing, distribution and retail, and sanitation

industries. Additionally, an online survey was created in Qualtrics (Provo, UT, USA) and

administered to the advisory council (12 respondents) to identify the intervention strategies of

greatest interest to industry. Based on the feedback, (i) cattle vaccination, (ii) transition from

overhead spray and furrow to drip irrigation, and (iii) transition from overhead spray to furrow

irrigation, were the most requested intervention strategies and were included in the scenario

analysis. Further, in another online survey with the council conducted via Qualtrics, we

presented the scope of the risk assessment model, including contamination sources and

interventions.

Model validation: The main model outcome was validated with the information from the CDC

for the year 2011 (most recent data). Since the most recent data was from 2011, we used the

2011 USA population data ($USpop$) *[88]* along with 2011 USA fresh romaine and leaf lettuce

consumption data (consumer weight without accounting for the food loss at consumer level)

[18]. The validation estimated the USA foodborne diseases associated with leafy greens

contaminated with ECO157 *[49]*. Accordingly, the estimated number of foodborne diseases due

to ECO157 contamination in the USA averages 63,153 cases annually, with 90% credible interval

from 17,587 to 149,631. Of these, 19.3% - 31.5% are attributed to leafy greens *[50]*, resulting in

an average of 12,189 – 19,893 cases each year in the USA linked to consuming contaminated

leafy greens. Considering the 90% credible interval, the true range of annual illness cases

attributable to consumption of romaine could be between 3,394 and 47,134.

Sensitivity and uncertainty analysis: We evaluated the effect of variable and uncertain

parameters in the baseline model to determine the key factors influencing the main outcome

(the predicted number of illnesses in the USA). We used @RISK and R (*dplyr* package) for the

sensitivity and uncertainty analyses (for preharvest and postharvest models) and generated

tornado plots of Spearman rank-order correlation coefficients (SRCCs) *[89]*. The SRCC analysis

for the baseline preharvest model was based on a set of core and derived parameters, where

derived parameters were calculated from core parameters. However, no parameter was

included in the analysis more than once. This approach was taken to ensure that the results

were more interpretable and informative. More information about the selection of parameters

is provided in Supplementary Table S8. Parameters with absolute value of the SRCC $|\rho| < 0.05$

were considered negligible and were excluded from the tornado plots. Likewise, only

parameters with statistically significant SRCC were presented in graphs, adhering to a

Bonferroni-corrected alpha threshold of 0.05 divided by the number of evaluated parameters

(Sensitivity analysis: 21 for preharvest, 18 for postharvest; uncertainty analysis: 11 for

preharvest). The absolute value of the SRCC, or $|\rho|$, was interpreted according to its common

usage in the literature (ρ=0 to <0.1, none; ρ=0.1 to <0.3, poor; ρ=0.3 to <0.6, fair; ρ=0.6 to <0.8,

moderate; ρ=0.8 to 1.0 very strong correlation) *[90,91]*. In the sensitivity analysis of variable

parameters, all uncertain parameters were set to their fixed deterministic values. A parameter

was classified as an uncertainty parameter when no information was available in the literature,

requiring assumptions. In the uncertainty analysis of the preharvest model, we fixed the

variable parameters to the median values of their respective distributions and created a

distribution for each of the uncertain parameters by establishing a triangular distribution. The

minimum of each triangular distribution was set as a 50% reduction from the baseline, and the

maximum was set as a 50% increase from the baseline for all uncertain parameters, with the

original baseline value as the mean/mode. In the preharvest model, this definition applied to

the following parameters: (i) volume of irrigation water captured by romaine through furrow or

drip irrigation per irrigation event (V_F, V_D); (ii) fraction of soil batch transferred to romaine

batches via splash caused by furrow or drip irrigation (Tr_{man_F}, Tr_{man_D}); (iii) dilution factors

used for the distribution of irrigation water, feces and runoff in soil batch

(DiF_{irr}, DiF_{pig}, DiF_{cow}); (iv) switch rate from non-persister to persister cells in BSAAO-

amended soil ($NtoP_{man}$); (v) ECO157 prevalence (P_{irr_w}) in well water; and (vi) transfer of

ECO157 cells from irrigation water to the romaine batch in an irrigation event (Tr_{ecirr}).

Regression trees: Decision trees are developed for both preharvest and postharvest models

using the *rpart* package in R *[92]*. All analyses were implemented at default settings in

rpart.control function. To prevent tree overfitting, we used 10-fold cross-validation *[81]*. Similar

to the SRCC analysis, the regression tree for the preharvest model was based on the set of the

core and derived parameters previously determined (Supplementary Table S8).

Scenario analysis of risk management strategies: Based on the sensitivity and uncertainty

analyses, as well as the results of expert elicitation, we identified nine risk management

strategies for evaluation in the scenario analysis (Table 1, Supplementary Table S9). These

scenarios are (i) cattle vaccination, (ii - iv) treatment of surface water, used for overhead spray

irrigation, by either chlorine, peracetic acid or ultraviolet radiation, (v) transition from overhead

spray to furrow irrigation system, (vi) transition from overhead spray and furrow to drip

irrigation, (vii) chlorine wash alternatives during postharvest processing, (viii) temperature

reduction during retail, and (ix) consumer wash (the modifications to the parameters to apply

to these scenarios were provided in Table 1). In addition, we evaluated the effect of parameters

related to clustering of ECO157 cells during the partitioning process (i.e., romaine batch to

package and package to serving) as an additional scenario (Table 1). The effect of each risk

management strategy or clustering was assessed based on their impact on the main outcome.

578
Results

Preharvest model: Under the baseline model conditions, our model predicted a 1.00%

prevalence of ECO157 in romaine batches across all three regions. The median ECO157 count in

contaminated romaine batches (637.5kg romaine/batch or 2,125 plants/batch) after harvest

(C_{Lh_f}) was 3.30 Log_{10}CFU/romaine batch ($5^{th} - 95^{th}$ percentile: $0.59 - 5.41$).

The sensitivity analysis of the variable parameters in the preharvest model (Figure 3),

determined by SRCC, indicated that the ECO157 count in irrigation water (C_{irr_P}) has a fair-

strength (0.29) positive correlation with the ECO157 count in romaine batches (outcome of

preharvest). The rest of the parameters were either poorly correlated or showed no

correlation.

To assess the impact of preharvest contamination sources (irrigation, wildlife, runoff, and

BSAAO) on the ECO157 counts in romaine batches, we ran the baseline preharvest model

separately for each source and visualized their effects using violin plots (i.e., we evaluated the

effect of each source individually, excluding the effect of the other three sources). Figure 4 (a)

shows the distribution of ECO157 counts in contaminated romaine batches (i.e., with ECO157

counts ≥1 CFU/romaine batch), for each preharvest source, including the median, interquartile

range, and 5^{th} to 95^{th} percentile across 100,000 iterations. Irrigation was identified as the

primary contributor to ECO157 counts in romaine batches (Figure 4 (b)) with a median ECO157

count of 3.91 $Log_{10}CFU$/romaine batch ($5^{th} - 95^{th}$ percentile: $0.00 - 5.29$) and an ECO157

prevalence of 0.53%. This was followed by wildlife, runoff and BSAAO contamination sources

(Figure 4 (b)). While wildlife (median ECO157 count: 2.27 $Log_{10}CFU$/romaine batch, $5^{th} - 95^{th}$

percentile: $0.47 - 4.47$) and runoff (median ECO157 count: 2.28 $Log_{10}CFU$/romaine batch, $5^{th} -$

95^{th} percentile: $0.48 - 5.28$) both led to similar median ECO157 counts when contamination

occurred, contamination by wildlife intrusion events was more frequent (0.26%) than runoff

contamination (0.21%). The predicted ECO157 count in romaine batches contributed by BSAAO

contamination was low and occurred in 2 out of 100,000 batches (0.002% occurrence), with

ECO157 counts of 8 and 16 CFU/romaine batch. This result on BSAAO contamination was not

negligible given the volume of romaine production in the USA: Between 2018 and 2022, an

average of 3,578 million kilograms of lettuce were produced annually, with romaine and leaf

lettuce together accounting for 53% of the total production *[101]*.

We also evaluated the impact of three irrigation systems—overhead spray, furrow, and drip—

on ECO157 counts in romaine batches (Figure 4 (c) and (d)). We found that overhead spray

irrigation was the primary contributor to ECO157 counts in romaine batches (median ECO157

count: 4.00 Log_{10}CFU/romaine batch, $5^{th} - 95^{th}$ percentile: $2.53 - 5.30$), leading to ECO157

counts in romaine batches most frequently (0.47%). In contrast, we predicted that furrow and

drip irrigation rarely caused ECO157 counts in romaine batches (0.04% and 0.02%, respectively)

and resulted in substantially lower ECO157 counts when contamination occurred (Furrow

median ECO157 count: 0.00 Log_{10}CFU/romaine batch, $5^{th} - 95^{th}$ percentile: $0.00 - 1.40$; drip

median ECO157 count: 0.30 Log_{10}CFU/romaine batch, $5^{th} - 95^{th}$ percentile: $0.00 - 1.54$).

We ran a sensitivity analysis (using SRCC) for the variable parameters in the baseline irrigation

model (considering only the three irrigation systems with no other preharvest contamination

sources), as well as for each individual irrigation source contamination separately — i.e., (i)

baseline irrigation model with contamination from only overhead spray irrigation, (ii) baseline

irrigation model with contamination from only furrow irrigation, and (iii) baseline irrigation

model with contamination from only drip irrigation. Similar to what was observed in the

baseline model, we found a poor to fair correlation between the ECO157 counts in a given

romaine batch and the ECO157 count in irrigation water (C_{irr_P}). Other variables were also

either poorly correlated or had no correlation with ECO157 counts in romaine batches,

regardless of the irrigation system (Supplementary Table S10).

The preharvest regression tree captured the importance of irrigation and wildlife-related

parameters in predicting ECO157 count in a romaine batch (Figure 5). The highest ECO157

counts in romaine batches were predicted when the ECO157 count in irrigation water was ≥ 0.3

CFU/100ml and applied via an overhead spray system. On the other hand, irrigation from furrow

or drip systems led to substantially lower ECO157 counts in romaine batches. Also

demonstrated was the importance of wildlife intrusions. When wildlife intrusion occurred for

one or more days during the countdown period, it resulted in a mean of 1,000 CFU ECO157 per

romaine batch, given that the fecal material from the feral swine contained 3 Log_{10}CFU ECO157

or more per gram of feces.

Postharvest model: The postharvest model predicted that the median number of cases

resulting from the consumption of romaine was 19,040 annually ($5^{th} - 95^{th}$ percentile: 42 −

3.6×10^6). Although our predicted median number of illness cases aligned with the reported

range for the annual median illness cases (12,189 − 19,893), the predicted distribution of illness

cases was wider than the reported validation data (reported 90% CI: 3,394 − 47,134 cases

annually). The validation data fell between 32^{nd} and 64^{th} percentiles of the predicted illness

cases. Further predictions on hospitalizations, cases of haemolytic uraemic syndrome (HUS) and

mortality were provided in Supplementary Table S11, based on CDC surveys on ECO157

outbreaks *[102]*. Considering the number of romaine servings consumed per year in the USA,

the annual median risk of ECO157 due to consumption of ECO157-contaminated fresh-cut

romaine was predicted as 0.000132% (or 1 illness in approximately 759,641 consumed

servings). In addition, the predicted median prevalence of ECO157 contaminated romaine

packages was 0.14% following the retail storage, but in 51% of the iterations the contaminated

package (225g) had only 1 CFU per package.

The sensitivity analysis of variable parameters in postharvest model (Figure 6), determined by

SRCC, showed that the median number of illness cases was very strongly correlated with the

ECO157 count in romaine batches entering processing (C_{Lh_f}). The remaining parameters were

either poorly correlated or showed no correlation with the number of illness cases.

The postharvest regression trees demonstrated how ECO157 counts, and consequently the

number of illness cases, substantially increases due to both preharvest and postharvest

conditions (Figure 7). Specifically, the ECO157 prevalence and counts in preharvest romaine

batches, as well as the temperatures and durations of postharvest retail and home storage, play

a critical role in this increase. In Figure 7 (a), the regression tree shows a high number of illness

cases, although occurring with a small probability, when $C_{Lh_f} \geq 3.9$ Log$_{10}$CFU/romaine batch,

$T_{retail} \geq 5°C$ and $T_{home} \geq 5°C$. Illness cases were lowest when $C_{Lh_f} < 1.6$ Log$_{10}$CFU/romaine

batch and $T_{home} < 5 °C$. Figure 7 (b) indicated that the ECO157 counts in contaminated

servings were increased by a combination of time-temperature abuses during retail and home

storage: ≥ 4 day storage at $\geq 8.9°C$ in retail, combined with home storage at $\geq 5.1°C$ resulted

in the highest ECO157 counts in romaine servings. On the other hand, the ECO157 counts were

lowest when $C_{Lh_f} < 5.2$ Log$_{10}$CFU/romaine batch, $T_{retail} < 9°C$ and $T_{home} < 5.1°C$. Figure 7 (c)

indicated that the highest prevalence of ECO157 contaminated romaine servings occurred

when the preharvest ECO157 counts and prevalence in romaine batches was higher, along with

retail storage at $\geq 5°C$.

In the preharvest model, none of the uncertainty variables were correlated with the predicted

ECO157 count in romaine batches. Additionally, only approximately 0.3% of illness cases were

attributed to the presence of persister cells in the baseline model, as detailed in Supplementary

Table S12.

Scenario analysis

In the 9 scenarios we tested (excluding clustering effect), the median number of illness cases

decreased by 17.1% to 99.7% (Table 1). Out of the nine scenarios tested independently, six

targeted reducing ECO157 counts at the preharvest stage. Among scenarios tested, the most

effective was a postharvest "best case" wash alternative with 3.0 to 5.0 Log$_{10}$CFU reduction

[97], which achieved a 99.7% reduction in median illness cases. The reduction rate in this

scenario was a general estimate based on experimental results from various investigated wash

methods—such as electrolyzed water combined with ultrasound or UV exposure, and

alternative sanitizers like bacteriophages, phage lysins, or nanoparticles *[97]*. Because these

alternative washes are still experimental and (i) likely cannot be implemented at scale or

outside of laboratory settings and (ii) would not have been applied in the production of all

lettuce produced nationwide, we also evaluated more moderate and realistic reductions in

ECO157 levels. Specifically, we modeled additional scenarios with median reductions of 1.0, 1.5,

2.0 and 2.5 $Log_{10}CFU$. These corresponded to reductions in the median number of illness cases

by 61.9%, 87.3%, 95.7% and 98.4%, respectively. Surface water treatments and transitioning to

furrow or drip irrigation resulted in slightly lower but highly effective reductions in ECO157

counts at the preharvest stage, achieving between 90.5% and 96.8% reductions in median

illness cases. In contrast, consumer washing (28.0%), cattle vaccination (25.9%) and

temperature reduction at retail (17.1%) were considerably less effective at reducing the

number of illness cases.

The scenario analyses also showed that the degree of clustering assumed during the

partitioning process affects the risk of illness. Increased clustering (in a batch and package)

increased the dose of exposure and thus the probability of illness given exposure, but it also

reduced the prevalence of ECO157 positive servings, which led to reduced number of cases

(Table 1).

Discussion

This risk assessment model simulated ECO157 counts during the production and consumption

of fresh-cut romaine, providing a comprehensive view of how various preharvest and

postharvest factors influence food safety risk. Our findings indicated that the model's

predictions of ECO157 counts, and associated illness cases aligned with surveillance studies in

the USA. Irrigation emerged as the most significant source of ECO157 counts in fresh-cut

romaine, particularly when considering the impact of other contamination sources:

inadequately treated BSAAO, runoff, and wildlife intrusion. However, the public health impact

could be mitigated through effective interventions. Our predictions also indicated a high

number of illness cases due to the combined effects of preharvest and postharvest factors.

Additionally, we identified key knowledge gaps in assessing human ECO157 exposure through

contaminated leafy greens.

The developed risk assessment tracks the dynamics of ECO157 cross-contamination between

romaine and soil prior to harvest.

We developed a novel difference equation model, which was incorporated into our preharvest

model, quantifying ECO157 counts in both soil and romaine in the field. Similar approaches

have been used in other risk assessment models, such as those describing the transmission of

Listeria monocytogenes contamination in a smoked salmon facility *[103]*, *Salmonella* in pig

slaughterhouses *[104]*, and *Salmonella* Typhimurium among farm swine *[105]*. This approach

enabled us to track daily dynamics of ECO157 counts on romaine during the days prior to

harvest. It allowed us to track the specific impacts of irrigation, wildlife, runoff and BSAAO; as

well as to perform calculations related to ECO157 growth, biphasic decay and cross-

contamination. Each of these factors has distinct characteristics that would be impossible to

accurately capture with a static equation.

Model predictions for preharvest and retail level ECO157 counts are consistent with previous reports.

Our findings generally aligned with existing literature and foodborne illness surveillance data for the USA. The predicted annual median illness cases were in agreement with the average observed foodborne illnesses linked to ECO157-contaminated romaine in the USA. However, the distribution of the predicted illness cases was wider than the reported illness range. One potential reason for this discrepancy was that that we did not account for romaine loss/waste at the consumer stage, assuming instead that all retail romaine would be consumed. This assumption is likely inaccurate, given that it has been estimated that 39.7% lettuce loss occurred at the consumer level in the USA in 2011 [18]. Another reason for the right skewness may be explained by the interactions between various parameters in the model, which will be discussed in the following sections. The predicted ECO157 prevalence at preharvest (1.00%) falls within the range of previously reported values for leafy greens from CA's central coast region [106], which ranged from <0.01% in 2007 to 2.5% in 2013. While a previous study by Zhang et al. (2018) reported a 0.02% average prevalence of ECO157 on romaine at the retail level in the USA (95% Confidence Interval by Clopper-Pearson method: <0.01 - 0.10, based on 5,548 samples of 25g each) [11], our predicted median ECO157 prevalence of contaminated romaine packages was over 7 times higher (0.14%, predicted for 225g packages). One potential explanation for the discrepancy between the reported and predicted prevalence could be the limitation of microbial tests in detecting contamination when smaller sample sizes are used. For instance, in Zhang et al.'s (2018) study, the detection limit was 1 CFU in a 25g sample (BAM method for diarrheagenic *E. coli*), equivalent to 0.04 CFU/g. For a 225g sample—such as the

packages in our model—the detection limit would be 1 CFU per 225g, or 0.0044 CFU/g. Under

this scenario, testing the entire 225g package could detect ECO157 levels 9 times lower than a

25g sample. Assuming a 225g sample size, estimating ECO157 counts could yield a prevalence 9

times higher than that observed in retail samples, potentially leading to 0.18% (0.02% × 9)

ECO157 prevalence. In our risk assessment model, we considered any package (and serving)

contaminated if it contained even a single ECO157 cell. Nevertheless, the previously observed

ECO157 prevalence (0.02%) *[11]* falls within the predicted distribution of ECO157 prevalence

simulated in our study (P_{pack}, Supplementary Table S2).

Irrigation is predicted to be the most important source of ECO157 contamination of fresh-cut

romaine with appropriate interventions reducing public health impact.

Our findings suggested that mitigation strategies at preharvest should prioritize reducing

contamination of irrigation water, particularly in cases of overhead spray irrigation supplied via

surface water sources. Aligned with our findings, contaminated irrigation water has been

defined as a major risk factor in the contamination of leafy greens *[11]*. Notably, contaminated

surface water irrigation has either been directly linked to or strongly associated with several

multistate ECO157 outbreaks of leafy greens in the USA *[107,108,109]*, including spinach *[107]*

and iceberg lettuce outbreaks in 2006 *[109]*, and a romaine outbreak in 2018 *[108]*. For the

romaine outbreak in 2018, ECO157 introduction into the irrigation water through aerial and

land-based spray applications of crop protection chemicals was proposed as a plausible route of

introduction *[110]*. On the other hand, the spinach and iceberg lettuce outbreaks appeared to

have been facilitated by cross-contamination, where contaminated surface water and

wastewater flowed into groundwater or surface water used for irrigating the lettuce plants

[107,109].

Our model also suggests that treatment of surface irrigation water—an important intervention

strategy already required by the California and Arizona LGMAs since the 2023 update—may

achieve a substantial reduction of human illness cases. This outcome was predicted for various

surface water treatments, including chlorination, PAA or UV. We consistently observed a strong

effectiveness across all treatment methods in reducing illness cases, based on the assumption

that the different methods achieved ECO157 reduction in surface irrigation water, ranging from

a minimum of 1.00 Log_{10} CFU/ml (chlorination) to a maximum of 3.09 Log_{10} CFU/ml (PAA).

Previous research has also reported that these methods reduce *E. coli* counts by up to 6

Log_{10}CFU/ml in agricultural water sources, with PAA typically having the strongest impact

[96,111-116]. However, the effectiveness of these methods can be highly variable, even when

implemented correctly *[115]*. Their site-specific applicability is influenced by a range of factors

including water quality (such as total suspended solids, turbidity, pH, oxygen demand and

temperature), applied dose, contact time, and contaminant characteristics *[117-119]*. In an

open leafy green production field, these factors can be affected by rainfall, runoff or sediment

activity upstream of the water sources *[116]*. On the other hand, both California and Arizona

LGMAs prohibit the use of surface irrigation waters that fail to meet the performance

requirements. To use such waters, treatment and testing are required to demonstrate

treatment efficacy and compliance with microbial standards. Therefore, future models should

(i) better account for the variability in treatment effectiveness and (ii) assess the impact of

reducing treatment variability: A less variable treatment with lower $Log_{10}CFU/ml$ reduction may

be more effective than one with a higher reduction but greater variability. Addressing these

challenges can help leafy green growers, who must navigate multiple treatment options

influenced by various factors while meeting LGMAs' requirements, make more informed

decisions.

Our data also suggested that transitioning from overhead spray irrigation to furrow or, ideally,

drip irrigation systems could provide a viable solution for reducing illness cases (Figure 4, Table

1). Despite introducing numerous parameters specific to each irrigation system that reasonably

differentiate between the three irrigation systems, our model suggested that the primary driver

behind the higher number of illness cases associated with overhead spray irrigation was its

capacity for increased direct contact between irrigation water and romaine (Supplementary

Table S10). Conversely, drip irrigation, delivering irrigation water to the plant roots, minimized

water contact, and consequently resulted in the lowest number of illness cases among the

three irrigation systems. Numerous field studies *[120-122]* and risk assessments *[40,41]* have

consistently demonstrated that overhead spray irrigation system leads to higher counts of

generic *E. coli* and ECO157 on romaine compared to furrow or drip irrigation systems, and

argued that the type of irrigation system plays a critical role in mitigating the risk of ECO157

contamination on leafy greens *[111,123,124]*. Aligned with our model's results, research also

supports that furrow irrigation leads to greater produce contamination than drip irrigation,

highlighting drip irrigation as the most effective irrigation method for reducing health risks from

contaminated irrigation water *[40,72,125]*. In our study, transitioning fully from overhead spray

and furrow irrigation to drip irrigation was predicted to be more effective than chlorination and

UV treatments of surface irrigation water, though less effective than PAA treatment.

The combined effect of preharvest and postharvest factors defines the public health risk.

The developed model illustrated that interactions of preharvest and postharvest factors affect

the number of predicted illness cases (Figure 7). This aligns with findings from observational

studies, which support the multifactorial nature of food safety *[126]*. Our model highlighted

how certain co-occurring factors could, though rarely, lead to a very large number of illnesses.

Specific factors, such as higher ECO157 counts in irrigation water, the use of overhead spray

irrigation systems instead of furrow or drip irrigation, or increased number of wildlife intrusion,

led to elevated ECO157 counts on preharvest romaine (Figure 5), which were further increased

by improper handling and storage conditions during postharvest, including time-temperature

abuse from harvest until consumption (Figure 7). This highlights the importance of simulation

tools like the one developed here that can be used to study impacts and prevention of rare

events (occurring individually or in combination) which would be challenging to study

empirically.

Several aspects of farm-to-fork continuum of fresh-cut romaine production and consumption

were included in the model but did not emerge as significant risk factors. These included

harvesting months, weather conditions (i.e., precipitation intensity, number of sunny days,

decay related to sun exposure) and several others. Weather conditions, including temperature,

seasonality and precipitation events, have previously shown to influence pathogen counts on

leafy greens *[10,35,127]*. In addition, historically, more leafy greens outbreaks have occurred in

summer and fall seasons than spring and winter seasons in the USA *[1]*. The likely reason why

our model did not capture the significance of these factors, including weather conditions, was

that weather conditions alone was insufficient to exacerbate the risk of contamination unless

cooccurring with other factors that introduce ECO157 contamination into the field. For

example, heavy rain will introduce contamination via runoff from a neighboring dairy farm only

if there was a nearby dairy farm and cattle on the farm that shed ECO157. Similarly, heavy rain

would contaminate romaine via splash but only if soil was already contaminated. In addition, it

was possible that certain factors that may increase ECO157 contamination during certain

months or under certain weather conditions were not or not fully included in our model (e.g.,

increased leaf damage or blistering induced by certain weather conditions) *[128]*.

Our study shows that a "best case" highly effective produce washing (3.0 to 5.0 $Log_{10}CFU/g$

reduction) is predicted to reduce the illness cases by over 99%, even with high preharvest

contamination levels (Table 1). This best case scenario was based on research studies on

alternative sanitizers that reported reductions around 5.0 $Log_{10}CFU$; however, most of these

studies have been limited to small-scale experiments and have not been adopted by industry

[97]. This high level of reduction however appears to typically be unachievable, at least with

current approaches and at realistic production scales, as indicated by studies that suggest

substantially lower reduction rates *[97]*. In the baseline model, the impact of postharvest

washing on illness risk was limited due to the use of a conservative and likely more realistic

ECO157 reduction — a maximum of 1.03 $Log_{10}CFU/g$ reduction (Figure 6). There are however

some reports that suggest that chlorine-based produce washes can achieve higher ECO157

reductions by preventing cross-contamination of ECO157 in the romaine batch during washing

[17,129]. Importantly, our scenario analysis (Table 1) showed that even a modest

improvement—such as a median reduction of 1.5 Log_{10}CFU/g rather than the baseline 1.03

Log_{10}CFU/g reduction, or just ~0.5 Log_{10}CFU/g improvement—could result in 61.9% decrease in

predicted median illness cases. This supports the importance of continuous improvement of

produce wash treatments as even relatively small improvements in effectiveness may lead to

tangible public health improvements. Our findings also suggest that it is important for industry

to verify produce wash effectiveness under realistic commercial scale conditions to correctly

assess risk reductions that can be expected by a given produce wash system. The reduction is

influenced by various factors, including the concentration and state (free or bound) of chlorine,

pH of the water, the duration of contact time, and the characteristics of the organism

[84,130,131]. Addressing these parameters has been the focus of multiple studies, which aim to

close knowledge gaps on produce wash efficacy *[130,132-134]*.

Model limitations and key knowledge gaps

Our model here reflected a synthesis of the current knowledge on ECO157 contamination

throughout the farm-to-fork supply chain of fresh-cut romaine lettuce in the USA. However, we

acknowledge limitations related to the model structure and parameterization. The most

consequential assumption in our model was about dilution factors in preharvest, which

facilitated modeling the spread of ECO157 from BSAAO, runoff, irrigation and fecal material on

the field. For ECO157 spread via BSAAO, we assumed a typical tiling depth of 15cm to 20cm. In

contrast, for irrigation, swine feces introduction caused by feral swine, and runoff, we assumed

a uniform spread of ECO157 to a depth of 1cm. As shown in Supplementary Table S13, the

depth significantly impacts the predicted ECO157 counts. Therefore, our analysis indicates that

using a 1cm depth was a conservative approach because the risk of romaine contamination

from contaminated soil decreases as the depth of bacterial spread increases. Despite this, our

assumptions about spread were considered reasonable based on the complexity of acquiring an

accurate depth of ECO157 spread in these various media (runoff, swine feces and BSAAO)

within 24 hours of their introduction to the field. Another limitation was that, while BSAAOs of

poultry origin are the most common in fields across the USA, our study focused on those of

dairy origin. Another assumption in the preharvest model was a constant and low-level ECO157

transfer from water and soil to romaine through furrow and drip irrigation. However, since

these irrigation methods involve less direct contact with the romaine compared to overhead

spray irrigation, we do not expect this assumption to have a major impact on the study

conclusions —a finding also supported by the preharvest uncertainty analysis. Additionally, we

did not account for the exterior leaves removed in the field, as we do not know the fraction of

cells on these outer vs inner leaves. However, based on our assumptions, the impact of this

practice was minimal as a 33.3% reduction in the ECO157 count on romaine resulted in 13.5%

reduction in the median predicted illness cases (as detailed in Supplementary Table S14).

Preharvest contamination was modeled at batch level, ignoring the spatial distribution of

contamination among plants and soil in different locations of the field. However, we consider

this to be a reasonable simplification since romaine was washed in batches, and the process of

washing may further distribute ECO157 within a romaine batch. The preharvest model focused

on open-field romaine production, excluding other potential production methods, such as

greenhouses or vertical farming. This decision is expected to have minimal impact, as the

majority of lettuce in the USA is grown in open fields *[101]*.

During postharvest, presence of cell clustering during partitioning — a poorly understood

phenomenon in romaine packaging and servings — was predicted to moderately reduce illness

cases (up to 48% reduction when high clustering). This reduction was due to the prediction of a

lower ECO157 prevalence in batches, as clustering decreased the probability of consumers

being exposed to servings with high ECO157 counts, as also predicted by Zoellner et al. (2019)

[85]. Clustering acted as a form of containment, minimizing the spread of contamination. For

the industry, this insight may encourage mechanical processing that minimizes disturbance of

clustered cells, preventing cells from spreading across the produce. For example, preventing

cross-contamination during postharvest wash process (e.g., by adding chlorine), helps contain

contamination within a single romaine batch, thereby reducing the risk of human illness (Table

1). We assumed that 100% of annual romaine consumption in the US is produced domestically,

even though USDA estimates suggest that only about 85% is US-grown *[138]*, based on reported

import and export data *[139]*. We acknowledge this as a limitation of our study. The dose-

response model used here describes the distribution of susceptibility to ECO157 and accounts

for population variability. Based on the assumptions of the Beta-Poisson model, (i) each cell is

capable of causing illness, and (ii) each cell is equally infective *[87]*. However, ECO157 illness is

a complex phenomenon influenced by factors such as the genetic diversity of ECO157 strains,

variations in virulence factors, and host susceptibility. As a result, some strains may have a

higher or lower infectious dose (ID_{50}), which is estimated to be between 10 and 100 bacteria,

according to the National Advisory Committee on Microbiological Criteria for Foods (NACMFS)

[140]. Therefore, our results should be interpreted with caution, as these factors are not fully

captured by the model.

Conclusion

The risk assessment model presented here provides a comprehensive analysis of the impact of

various preharvest and postharvest factors and their interactions. Our results suggest that

illnesses occur mainly (52%) due to ECO157 contamination originating from untreated irrigation

water applied through overhead spray irrigation. Our data further suggests that preharvest

ECO157 risk, including that from irrigation, can be reduced either through water treatments or

by switching to furrow or drip irrigation. In postharvest, maintaining the cold chain is crucial to

prevent high numbers of illness cases. While our data indicates that an effective postharvest

wash can help mitigate the public heath impact of both preharvest and postharvest

contamination, future work on the impact of postharvest wash variability on public health risk

is important. We identify knowledge gaps regarding the microbial quality of irrigation waters,

as well as the effectiveness of preharvest surface water treatments and postharvest wash

processes. In conclusion, the developed QMRA is expected to inform development of

sustainable food safety strategies along the supply chain and guide further research to address

key knowledge gaps.

References

1. Turner, K., Moua, C. N., Hajmeer, M., Barnes, A. & Needham, M. Overview of leafy
greens–related food safety incidents with a California link: 1996 to 2016. *J. Food Prot.*
82, 405–414 (2019).
2. CDC. NORS Dashboard. https://wwwn.cdc.gov/norsdashboard/ (2022).
3. CDC. Active investigations in foodborne outbreaks. https://www.cdc.gov/foodborne-
outbreaks/active-investigations/all-foodborne-outbreak-notices.html (2024).
4. Coulombe, G., Catford, A., Martinez-Perez, A. & Buenaventura, E. Outbreaks of
Escherichia coli O157:H7 infections linked to romaine lettuce in Canada from 2008 to
2018: An analysis of food safety context. *J. Food Prot.* **83**, 1444–1462 (2020).
https://doi.org/10.4315/JFP-20-029
5. Benjamin, L. A. et al. Risk factors for *Escherichia coli* O157 on beef cattle ranches located
near a major produce production region. *Epidemiol. Infect.* **143**, 81–93 (2015).
6. Jay, M. T. et al. *Escherichia coli* O157:H7 in feral swine near spinach fields and cattle,
central California coast. *Emerg. Infect. Dis.* **13**, 1908 (2007).
7. California Food Emergency Response Team (CalFERT). Investigation of an *Escherichia coli*
O157:H7 outbreak associated with Dole pre-packaged spinach. California Department of
Health Services, Sacramento, CA (2007).
8. Cooley, M. B. et al. Development of a robust method for isolation of Shiga toxin-positive
Escherichia coli (STEC) from fecal, plant, soil, and water samples from a leafy greens
production region in California. *PLOS ONE* **8**, e65716 (2013).
https://doi.org/10.1371/journal.pone.0065716
9. Atwill, E. R. et al. Transfer of *Escherichia coli* O157:H7 from simulated wildlife scat onto
romaine lettuce during foliar irrigation. *J. Food Prot.* **78**, 240–247 (2015).
https://doi.org/10.4315/0362-028X.JFP-14-277
10. Oliveira, M., Viñas, I., Usall, J., Anguera, M. & Abadias, M. Presence and survival of
Escherichia coli O157:H7 on lettuce leaves and in soil treated with contaminated
compost and irrigation water. *Int. J. Food Microbiol.* **156**, 133–140 (2012).
https://doi.org/10.1016/j.ijfoodmicro.2012.03.014
11. Zhang, G. et al. Survey of foodborne pathogens, aerobic plate counts, total coliform
counts, and *Escherichia coli* counts in leafy greens, sprouts, and melons marketed in the
United States. *J. Food Prot.* **81**, 400–411 (2018).
12. Ackers, M. L. et al. An outbreak of *Escherichia coli* O157:H7 infections associated with
leaf lettuce consumption. *J. Infect. Dis.* **177**, 1588–1593 (1998).
13. Hillborn, E. D. et al. A multistate outbreak of *Escherichia coli* O157:H7 infections
associated with consumption of mesclun lettuce. *Arch. Intern. Med.* **159**, 1758–1764
(1999).
14. FAO. Prevention and control of microbiological hazards in fresh fruits and vegetables,
Part 4: Specific commodities.
https://openknowledge.fao.org/server/api/core/bitstreams/151c9dda-508b-4b70-972a-
85725eb12145/content (2023).
15. Yang, Y., Luo, Y., Millner, P., Turner, E. & Feng, H. Assessment of *Escherichia coli*
O157:H7 transference from soil to iceberg lettuce via a contaminated field coring
harvesting knife. *Int. J. Food Microbiol.* **153**, 345–350 (2012).

16. Buchholz, A. L., Davidson, G. R., Marks, B. P., Todd, E. C., Ryser, E. T. Transfer of *Escherichia coli* O157:H7 from equipment surfaces to fresh-cut leafy greens during processing in a model pilot-plant production line with sanitizer-free water. *J. Food Prot.* **75**, 1920–1929 (2012). https://doi.org/10.4315/0362-028X.JFP-11-558

17. López-Gálvez, F., Allende, A., Selma, M. V. & Gil, M. I. Prevention of *Escherichia coli* cross-contamination by different commercial sanitizers during washing of fresh-cut lettuce. *Int. J. Food Microbiol.* **133**, 167–171 (2009). https://doi.org/10.1016/j.ijfoodmicro.2009.05.017

18. USDA ERS. Food availability per capita data system. https://www.ers.usda.gov/data-products/food-availability-per-capita-data-system/ (2020).

19. California LGMA. About us. https://lgma.ca.gov/about-us (2024).

20. Arizona LGMA. About us. https://www.arizonaleafygreens.org/about-us (2023).

21. FDA. FSMA Final Rule on Produce Safety: Standards for the Growing, Harvesting, Packing, and Holding of Produce for Human Consumption. https://www.fda.gov/food/food-safety-modernization-act-fsma/fsma-final-rule-produce-safety (2024).

22. FSMA PSR § 112.41, § 112.42. https://www.ecfr.gov/current/title-21/chapter-I/subchapter-B/part-112 (2024).

23. CFR § 205.203. https://www.ecfr.gov/current/title-7/subtitle-B/chapter-I/subchapter-M/part-205/subpart-C/section-205.203 (2024).

24. CFR § 112.55. https://www.ecfr.gov/current/title-21/chapter-I/subchapter-B/part-112/subpart-F/section-112.55 (2024).

25. CFR §112.83 and §112.112. https://www.regulations.gov/document/FDA-2011-N-0921-18558 (2024).

26. FSMA PSR. https://www.regulations.gov/document/FDA-2011-N-0921-18558 (2015).

27. Allende, A., Truchado, P., Lindqvist, R. & Jacxsens, L. Quantitative microbial exposure modelling as a tool to evaluate the impact of contamination level of surface irrigation water and seasonality on fecal hygiene indicator *E. coli* in leafy green production. *Food Microbiol.* **75**, 82–89 (2018).

28. Koseki, S. & Isobe, S. Prediction of pathogen growth on iceberg lettuce under real temperature history during distribution from farm to table. *Int. J. Food Microbiol.* **104**, 239–248 (2005).

29. Franz, E., Semenov, A. V. & van Bruggen, A. H. Modelling the contamination of lettuce with *Escherichia coli* O157:H7 from manure-amended soil and the effect of intervention strategies. *J. Appl. Microbiol.* **105**, 1569–1584 (2008).

30. Franz, E., Tromp, S. O., Rijgersberg, H. & Van Der Fels-Klerx, H. J. Quantitative microbial risk assessment for *Escherichia coli* O157:H7, *Salmonella*, and *Listeria monocytogenes* in leafy green vegetables consumed at salad bars. *J. Food Prot.* **73**, 274–285 (2010).

31. Tromp, S. O., Rijgersberg, H. & Franz, E. Quantitative microbial risk assessment for *Escherichia coli* O157:H7, *Salmonella enterica*, and *Listeria monocytogenes* in leafy green vegetables consumed at salad bars, based on modeling supply chain logistics. *J. Food Prot.* **73**, 1830–1840 (2010).

32. Danyluk, M. D. & Schaffner, D. W. Quantitative assessment of the microbial risk of leafy greens from farm to consumption: preliminary framework, data, and risk estimates. *J. Food Prot.* **74**, 700–708 (2011).

33. Rodríguez, F. P. et al. A mathematical risk model for *Escherichia coli* O157:H7 cross-contamination of lettuce during processing. *Food Microbiol.* **28**, 694–701 (2011).

34. Ottoson, J. R., Nyberg, K., Lindqvist, R. & Albihn, A. Quantitative microbial risk assessment for *Escherichia coli* O157 on lettuce, based on survival data from controlled studies in a climate chamber. *J. Food Prot.* **74**, 2000–2007 (2011).

35. Castro-Ibáñez, I., Gil, M. I., Tudela, J. A., Ivanek, R. & Allende, A. Assessment of microbial risk factors and impact of meteorological conditions during production of baby spinach in the Southeast of Spain. *Food Microbiol.* **49**, 173–181 (2015).

36. Mishra, A., Pang, H., Buchanan, R. L., Schaffner, D. W. & Pradhan, A. K. A system model for understanding the role of animal feces as a route of contamination of leafy greens before harvest. *Appl. Environ. Microbiol.* **83**, e02775-16 (2017).

37. Kundu, A., Wuertz, S. & Smith, W. A. Quantitative microbial risk assessment to estimate the risk of diarrheal diseases from fresh produce consumption in India. *Food Microbiol.* **75**, 95–102 (2018).

38. Pang, H, Lambertini, E., Buchanan, R. L., Schaffner, D. W. & Pradhan, A. K. Quantitative microbial risk assessment for *Escherichia coli* O157:H7 in fresh-cut lettuce. *J. Food Prot.* **80**, 302–311 (2017).

39. O'Flaherty, E., Solimini, A. G., Pantanella, F., De Giusti, M. & Cummins, E. Human exposure to antibiotic-resistant *Escherichia coli* through irrigated lettuce. *Environ. Int.* **122**, 270–280 (2019).

40. Rock, C. M. et al. Review of water quality criteria for water reuse and risk-based implications for irrigated produce under the FDA Food Safety Modernization Act, produce safety rule. *Environ. Res.* **172**, 616–629 (2019).

41. Bozkurt, H., Bell, T., van Ogtrop, F., Phan-Thien, K. Y. & McConchie, R. Assessment of microbial risk during Australian industrial practices for *Escherichia coli* O157:H7 in fresh-cut cos lettuce: A stochastic quantitative approach. *Food Microbiol.* **95**, 103691 (2021).

42. FDA. FDA-iRISK. https://irisk.foodrisk.org/ (2025).

43. USDA. Risk assessment of *E. coli* O157:H7 in ground beef. https://www.fsis.usda.gov/node/2003 (2001).

44. Ebel, E. et al. Draft risk assessment of the public health impact of *Escherichia coli* O157:H7 in ground beef. *J. Food Prot.* **67**, 1991–1999 (2004).

45. USDA. Microbial risk assessment guideline. Pathogenic microorganisms with focus on food and water. https://www.fsis.usda.gov/sites/default/files/media_file/2020-07/Microbial_Risk_Assessment_Guideline_2012-001.pdf (2012).

46. Nauta, M. J. Microbiological risk assessment models for partitioning and mixing during food handling. *Int. J. Food Microbiol.* **100**, 311–322 (2005).

47. Munther, D. S., Carter, M. Q., Aldric, C. V., Ivanek, R. & Brandl, M. T. Formation of *Escherichia coli* O157:H7 persister cells in the lettuce phyllosphere and application of differential equation models to predict their prevalence on lettuce plants in the field. *Appl. Environ. Microbiol.* **86**, e01602-19 (2020).

48. Turini, T. et al. Iceberg lettuce production in California.
https://escholarship.org/uc/item/7w47j6zv (2011).

49. Scallan, E. et al. Foodborne illness acquired in the United States—major pathogens.
Emerg. Infect. Dis. **17**, 7–15 (2011).

50. Painter, J. A. et al. Attribution of foodborne illnesses, hospitalizations, and deaths to
food commodities by using outbreak data, United States, 1998–2008. *Emerg. Infect. Dis.*
19, 407–415 (2013).

51. Longfellow's Greenhouses.
http://plants.longfellowsgreenhouses.com/12100007/Plant/Print?id=29076.

52. Rural Migration News (RMN). https://migration.ucdavis.edu/rmn/blog/post/?id=2699
(2022).

53. Agriculture Marketing Resource Center. Lettuce. https://www.agmrc.org/commodities-
products/vegetables/lettuce (2021).

54. Geisseler, D. & Horwath, W. R. Lettuce production in California.
https://www.cdfa.ca.gov/is/ffldrs/frep/FertilizationGuidelines/pdf/Lettuce_Production_
CA.pdf (2016).

55. Brouwer, A. F. et al. Modeling biphasic environmental decay of pathogens and
implications for risk analysis. *Environ. Sci. Technol.* **51**, 2186–2196 (2017).

56. Bezanson, G. et al. Comparative examination of *Escherichia coli* O157:H7 survival on
romaine lettuce and in soil at two independent experimental sites. *J. Food Prot.* **75**, 480–
487 (2012).

57. Erickson, M. C. et al. Surface and internalized *Escherichia coli* O157:H7 on field-grown
spinach and lettuce treated with spray-contaminated irrigation water. *J. Food Prot.* **73**,
1023–1029 (2010).

58. Moyne, A. L. et al. Fate of *Escherichia coli* O157:H7 in field-inoculated lettuce. *Food*
Microbiol. **28**, 1417–1425 (2011).

59. Williams, T. R., Moyne, A. L., Harris, L. J. & Marco, M. L. Season, irrigation, leaf age, and
Escherichia coli inoculation influence the bacterial diversity in the lettuce phyllosphere.
PLoS One **8**, e68642 (2013).

60. Chen, Z. et al. Prevalence of *Escherichia coli* O157 and *Salmonella* spp. in solid bovine
manure in California using real-time quantitative PCR. *Lett. Appl. Microbiol.* **69**, 23–29
(2019).

61. Rogers, S. W. et al. Decay of bacterial pathogens, fecal indicators, and real-time
quantitative PCR genetic markers in manure-amended soils. *Appl. Environ. Microbiol.* **77**,
4839–4848 (2011).

62. Chen, H. Contribution of the persister state to the tolerance of *Escherichia coli* O157:H7
to stresses encountered within the fresh produce chain. Master's Thesis, University of
Guelph (2020).

63. Gravuer, K. Compost application rates for California croplands and rangelands for a
CDFA Healthy Soils Incentives Program.
https://www.cdfa.ca.gov/oefi/healthysoils/docs/CompostApplicationRate_WhitePaper.
pdf (2016).

64. Aqua-Calc. https://www.aqua-calc.com/ (2023).

65. Valenzuela, H. R., Kratky, B. & Cho, J. Lettuce production guidelines for Hawaii. https://www.ctahr.hawaii.edu/oc/freepubs/pdf/res-164.pdf (1995).

66. California Department of Fish and Wildlife. Wild pig. https://wildlife.ca.gov/Hunting//Wild-Pig (2001).

67. Hamilton, D. W., Luce, W. G. & Heald, A. D. Production and characteristics of swine manure. Oklahoma State University Extension Facts Publication F-1735 (1997).

68. Ferretti, F., Storer, K., Coats, J. & Massei, G. Temporal and spatial patterns of defecation in wild boar. *Wildl. Soc. Bull.* **39**, 65–69 (2015).

69. USGS. Freshwater withdrawals in the United States. https://www.usgs.gov/special-topics/water-science-school/science/freshwater-withdrawals-united-states (2018).

70. Smith, B. A., Fazil, A. & Lammerding, A. M. A risk assessment model for *Escherichia coli* O157:H7 in ground beef and beef cuts in Canada: Evaluating the effects of interventions. *Food Control* **29**, 2 (2013).

71. Benjamin, L. et al. Occurrence of generic *Escherichia coli*, *E. coli* O157, and *Salmonella* spp. in water and sediment from leafy green produce farms and streams on the Central California coast. *Int. J. Food Microbiol.* **165**, 65–76 (2013).

72. Stine, S. W. et al. Application of microbial risk assessment to the development of standards for enteric pathogens in water used to irrigate fresh produce. *J. Food Prot.* **68**, 913–918 (2005).

73. Allende, A. et al. Quantitative contamination assessment of *Escherichia coli* in baby spinach primary production in Spain: Effects of weather conditions and agricultural practices. *Int. J. Food Microbiol.* **257**, 238–246 (2017).

74. FDA. Standards for the Growing, Harvesting, Packing, and Holding of Produce for Human Consumption. Federal Register. https://www.federalregister.gov/documents/2015/11/27/2015-28159/standards-for-the-growing-harvesting-packing-and-holding-of-produce-for-human-consumption (2015).

75. Girardin, L. et al. Behaviour of the pathogen surrogates Listeria innocua and Clostridium sporogenes during production of parsley in fields fertilized with contaminated amendments. *FEMS Microbiol. Ecol.* **54**, 287-295 (2005).

76. Alzahrani, A. Persister formation and revival in Shiga Toxin-Producing *Escherichia coli* subjected to environmental stress conditions. *PhD Thesis, University of Guelph.* *https://atrium.lib.uoguelph.ca/server/api/core/bitstreams/e5a584c9-3e95-41a1-aa4a-7e0abf5e0c42/content* (2020).

77. Hofsteenge, N., van Nimwegen, E. & Silander, O.K. Quantitative analysis of persister fractions suggests different mechanisms of formation among environmental isolates of *E. coli. BMC Microbiol* **13**, 25 (2013).

78. Niu, H., Gu, J. & Zhang, Y. Bacterial persisters: molecular mechanisms and therapeutic development. *Sig. Transduct. Target. Ther.* **9**, 174 (2024).

79. Wang, D. et al. Comparative persistence of *Salmonella* and *Escherichia coli* O157:H7 in loam or sandy loam soil amended with bovine or swine manure. *Can. J. Microbiol.* **64**, 979–991 (2018).

1170 80. Tran, D. T. Q. et al. Environmental drivers for persistence of *Escherichia coli* and
1171   *Salmonella* in manure-amended soils: A meta-analysis. *J. Food Prot.* **83**, 1268–1277
1172   (2020).
1173 81. McKellar, R. C., LeBlanc, D. I., Rodríguez, F. P. & Delaquis, P. Comparative simulation of
1174   *Escherichia coli* O157:H7 behaviour in packaged fresh-cut lettuce distributed in a typical
1175   Canadian supply chain in the summer and winter. *Food Control* **35**, 192–199 (2014).
1176 82. McKellar, R. C. LeBlanc, D. I., Lu, J. & Delaquis, P. Simulation of *Escherichia coli* O157:H7
1177   behavior in fresh-cut lettuce under dynamic temperature conditions during distribution
1178   from processing to retail. *Foodborne Pathog. Dis.* **9**, 239–244 (2012).
1179 83. When to harvest vegetables. University of Georgia Extension.
1180   https://extension.uga.edu/publications/detail.html?number=C935&title=when-to-
1181   harvest-vegetables (2022).
1182 84. Keskinen, L. A., Burke, A. & Annous, B. A. Efficacy of chlorine, acidic electrolyzed water
1183   and aqueous chlorine dioxide solutions to decontaminate *Escherichia coli* O157:H7 from
1184   lettuce leaves. *Int. J. Food Microbiol.* **132**, 134–140 (2009).
1185 85. Zoellner, C., Wiedmann, M. & Ivanek, R. An assessment of Listeriosis risk associated with
1186   a contaminated production lot of frozen vegetables consumed under alternative
1187   consumer handling scenarios. *J. Food Prot.* **82**, 2174–2193 (2019).
1188 86. USDA. *Lettuce, Cos or Romaine, Raw; Portion*. https://fdc.nal.usda.gov/food-
1189   details/746769/nutrients/ (2015).
1190 87. Cassin, M. H., Lammerding, A. M., Todd, E. C., Ross, W., & McColl, R. S. Quantitative risk
1191   assessment for *Escherichia coli* O157:H7 in ground beef hamburgers. *Int. J. Food*
1192   *Microbiol.* **41**, 21–44 (1998).
1193 88. U.S. Census Bureau. Census Bureau projects U.S. population of 312.8 million on New
1194   Year's Day. https://www.census.gov/newsroom/releases/archives/population/cb11-
1195   219.html (2011).
1196 89. Wickham, H., François, R., Henry, L., Müller, K. & Vaughan, D. dplyr: A Grammar of Data
1197   Manipulation. R package version 1.1.4. https://dplyr.tidyverse.org. (2023).
1198 90. Akoglu, H. User's guide to correlation coefficients. *Turk. J. Emerg. Med.* **18**, 91–93
1199   (2018).
1200 91. Mukaka, M. M. Statistics corner: A guide to appropriate use of correlation coefficient in
1201   medical research. *Malawi Med. J.* **24**, 69–71 (2012).
1202 92. Therneau, T., Atkinson, B. & Ripley, B. rpart: Recursive Partitioning and Regression
1203   Trees. https://cran.r-project.org/package=rpart (2018).
1204 93. Mir, R. et al. Cattle intestinal microbiota shifts following *Escherichia coli* O157:H7
1205   vaccination and colonization. *PLOS ONE* **14**, e0226099 (2019).
1206 94. Chang, T. Evaluation of multiple disinfection methods to mitigate contaminated
1207   irrigation water. Master's Thesis, University of Tennessee
1208   https://trace.tennessee.edu/utk_gradthes/3352 (2015).
1209 95. Dery, J. L., Gerrity, D. & Rock, C. Minimizing Risks: Use of surface water in pre-harvest
1210   agricultural irrigation; part ii: sodium and calcium hypochlorite (chlorine) treatment
1211   methods. https://repository.arizona.edu/bitstream/handle/10150/669941/AZ1831-
1212   2020.pdf?sequence=1&isAllowed=y (2011).

96. Beauvais, W. et al. The effectiveness of treating irrigation water using ultraviolet radiation or sulphuric acid fertilizer for reducing generic *Escherichia coli* on fresh produce – a controlled intervention trial. *J. Appl. Microbiol.* **131**, 1360–1377 (2021).

97. FAO/WHO. Joint FAO/WHO expert meeting on microbiological risk assessment on the prevention and control of microbiological hazards in fresh fruits and vegetables - part 4: commodity-specific interventions. https://www.who.int/publications/m/item/joint-fao-who-expert-meeting-on-microbiological-risk-assessment-on-the-prevention-and-control-of-microbiological-hazards-in-fresh-fruits-and-vegetables-part4 (2022).

98. FDA, University of California Davis. *Head Lettuce.* https://www.wifss.ucdavis.edu/wp-content/uploads/2016/10/HeadLettuce_PDF.pdf

99. EcoSure 2007 Cold Temperature Database. https://www.foodrisk.org/resources/display/21

100. Verrill, L. Consumer vegetable and fruit washing practices in the United States, 2006 and 2010. *Food Prot. Trends* **32**, 164-172 (2012).

101. USDA ERS. *Vegetables and Pulses Outlook* (2023). Available at: https://downloads.usda.library.cornell.edu/usda-esmis/files/cj82k729x/pn89fk86d/3f463k190/VGS-370.pdf

102. Tack, D. M. et al. Shiga toxin-producing *Escherichia coli* outbreaks in the United States, 2010-2017. *Microorganisms*, **9**, 1529 (2021).

103. Ivanek, R., Gröhn, Y. T., Wiedmann, M. & Wells, M. T. Mathematical model of *Listeria monocytogenes* cross-contamination in a fish processing plant. *J. Food Prot.* **67**, 2688–2697 (2004).

104. de Freitas Costa, E., Corbellini, L. G., da Silva, A. P. S. P. & Nauta, M. A stochastic model to assess the effect of meat inspection practices on the contamination of pig carcasses. *Risk Anal.* **37**, 1849–1864 (2017).

105. Ivanek, R., Snary, E. L., Cook, A. J., & Gröhn, Y. T. A mathematical model for the transmission of *Salmonella Typhimurium* within a grower-finisher pig herd in Great Britain. *J. Food Prot.* **67**, 2403–2409 (2004).

106. Karp, D. S. et al. Co-managing fresh produce for nature conservation and food safety. *Proc. Natl. Acad. Sci. U. S. A.* **112**, 11126–11131 (2015).

107. Gelting, R. J. et al. Irrigation water issues potentially related to the 2006 multistate *E. coli* O157:H7 outbreak associated with spinach. *Agric. Water Manag.* **98**, 1395–1402 (2011).

108. Bottichio, L. et al. Shiga toxin-producing *Escherichia coli* infections associated with romaine lettuce—United States, 2018. *Clin. Infect. Dis.* **71**, e323–e330 (2020).

109. Gelting, R. J. et al. A systems analysis of irrigation water quality in an environmental assessment of an *E. coli* O157:H7 outbreak in the United States linked to iceberg lettuce. *Agric. Water Manag.* **150**, 11–118 (2015).

110. FDA. *Environmental Assessment Factors Potentially Contributing to Contamination of Romaine Lettuce Implicated in Outbreaks.* (2019). Available at: https://www.fda.gov/food/outbreaks-foodborne-illness/environmental-assessment-factors-potentially-contributing-contamination-romaine-lettuce-implicated#summary

111. Allende, A. & Monaghan, J. Irrigation water quality for leafy crops: a perspective of risks and potential solutions. *Int. J. Environ. Res. Public Health* **12**, 7457–7477 (2015).

112. Dery, J. L., Choppakatla, V., Sughroue, J. & Rock, C. Minimizing Risks: Use of Surface Water in Pre-Harvest Agricultural Irrigation; Part III: Peroxyacetic Acid (PAA) Treatment Methods. Available at: https://extension.arizona.edu/sites/extension.arizona.edu/files/pubs/az1884-2021.pdf

113. Banach, J. L. et al. Application of water disinfection technologies for agricultural waters. *Agric. Water Manag.* **244**, 106527 (2021).

114. Collvignarelli, M. et al. Overview of the main disinfection processes for wastewater and drinking water treatment plants. *Sustainability* **10**, 86 (2018).

115. Rock, C. CPS Rapid Response – Yuma Valley, Final Report. *Center for Produce Safety* (2019).

116. Rock, C., Cooper, K. & Yemmireddy, V. K. Agriculture Water Treatment – Southwest Region Final Project Report. Center for Produce Safety (2021).

117. USEPA. Peracetic Acid Fact Sheet https://www.epa.gov/sites/default/files/2019-08/documents/disinfection_-_paa_fact_sheet_-_2012.pdf (2012).

118. *USEPA*. Chlorination Fact Sheet. https://www3.epa.gov/npdes/pubs/chlo.pdf (1999).

119. *USEPA*. UV Disinfection Fact Sheet. https://www3.epa.gov/npdes/pubs/uv.pdf (1999).

120. Fonseca, J. M. et al. *Escherichia coli* survival in lettuce fields following its introduction through different irrigation systems. *J. Appl. Microbiol.* **110**, 893–902 (2011).

121. Moyne, A. L. et al. Fate of *Escherichia coli* O157:H7 in field-inoculated lettuce. *Food Microbiol.* **28**, 1417–1425 (2011).

122. Solomon, E. B., Potenski, C. J. & Matthews, K. R. Effect of irrigation method on transmission to and persistence of *Escherichia coli* O157:H7 on lettuce. *J. Food Prot.* **65**, 673–676 (2002).

123. Gil, M. I., Tudela, J. A., Luna, M. C. & Allende, A. Water management and its effect on the postharvest quality of fresh-cut vegetables. *Stewart Postharvest Rev.* **9**, 1–8 (2013).

124. Gurtler, J. B. & Gibson, K. E. Irrigation water and contamination of fresh produce with bacterial foodborne pathogens. *Curr. Opin. Food Sci.* **47**, 100889 (2022).

125. Song, I., Stine, S., Choi, C. & Gerba, C. Comparison of crop contamination by microorganisms during subsurface drip and furrow irrigation. *J. Environ. Eng.* **132**, 1243–1248 (2006).

126. Park, S. et al. Multifactorial effects of ambient temperature, precipitation, farm management, and environmental factors determine the level of generic *Escherichia coli* contamination on preharvested spinach. *Appl. Environ. Microbiol.* **81**, 2635–2650 (2015).

127. Holvoet, K., Sampers, I., Seynnaeve, M., & Uyttendaele, M. Relationships among hygiene indicators and enteric pathogens in irrigation water, soil and lettuce and the impact of climatic conditions on contamination in the lettuce primary production. *Int. J. Food Microbiol.* **171**, 21–31 (2014).

128. Erickson, M. C. et al. Preharvest internalization of *Escherichia coli* O157:H7 into lettuce leaves, as affected by insect and physical damage. *J. Food Prot.* **73**, 1809–1816 (2010).

129. Petri, E., Rodríguez, M. & García, S. Evaluation of combined disinfection methods for reducing *Escherichia coli* O157:H7 population on fresh-cut vegetables. *Int. J. Environ. Res. Public Health* **12**, 8678–8690 (2015).

130. Mokhtari, A. et al. Evaluation of chlorine dioxide and peracetic acid on *Escherichia coli*
O157:H7 reduction on lettuce. *Food Control* **62**, 36–40 (2016).

131. Fu, T.-J., Li, Y., Awad, D., Zhou, T.-Y. & Liu, L. Factors affecting the performance and
monitoring of a chlorine wash in preventing *Escherichia coli* O157:H7 cross-
contamination during postharvest washing of cut lettuce. *Food Control* **94**, 212–221
(2018).

132. Gombas, D. et al. Guidelines to validate control of cross-contamination during washing
of fresh-cut leafy vegetables. *J. Food Prot.* **80**, 312–330 (2017).

133. Luo, Y. et al. Determination of free chlorine concentrations needed to prevent
Escherichia coli O157:H7 cross-contamination during fresh-cut produce wash. *J. Food*
Prot. **74**, 352–358 (2011).

134. Gómez-López, V. M., Lannoo, A. S., Gil, M. I. & Allende, A. Minimum free chlorine
residual level required for the inactivation of *Escherichia coli* O157:H7 and
trihalomethane generation during dynamic washing of fresh-cut spinach. *Food Control*
42, 132–138 (2014).

135.   Latorre, A. et al. Quantitative risk assessment of listeriosis due to consumption of
raw milk. *J. Food Prot.* **74**, 1268–1281 (2011).

136.   Pouillot, R., Lubran, M. B., Cates, S. C. & Dennis, S. Estimating parametric
distributions of storage time and temperature of ready-to-eat foods for U.S. households.
J. Food Prot. **73**, 312–321 (2010).

137. Climate Data Online. https://www.ncei.noaa.gov/cdo-web/

138. USDA. U.S. lettuce production shifts regionally by season.
https://www.ers.usda.gov/data-products/charts-of-note/chart-detail?chartId=106516
(2023).

139. USDA ERS. Vegetables and pulses data - data by commodity - imports and exports.
https://data.ers.usda.gov/reports.aspx?ID=4027&programArea=veg&top=5&HardCopy
=True&RowsPerPage=25&groupName=Vegetables&commodityName=Lettuce#P91068
9dff5c747409da423e3f8f44e38_4_292 (2025).

140. National Advisory Committee on Microbiological Criteria for Foods (NACMSF).
Response to questions posed by the food and drug administration regarding virulence
factors and attributes that define foodborne shiga toxin-producing *Escherichia coli*
(STEC) as severe human pathogens. *J. Food Prot*, **82**, 724–767 (2019).

1332

Acknowledgements

1334 This work was supported through a grant (award 2019-51181-30016) from the U.S. Department

1335 of Agriculture (USDA), National Institute of Food and Agriculture (NIFA). Partial support

1336 received from a grant (award 2020-67021-32855) from the USDA NIFA.

1337

1338

Author contributions

E.B.: conceptualization, data curation, formal analysis, investigation, methodology, software,

validation, visualization, writing- original draft, writing – review and editing; S.I.M.:

conceptualization, data curation, methodology, software, writing – review and editing; L.K.S.:

conceptualization, funding acquisition, writing – review and editing; M.D.D.: conceptualization,

funding acquisition, writing – review and editing; M.W.: conceptualization, funding acquisition,

writing – review and editing; R.I.: conceptualization, funding acquisition, investigation,

methodology, project administration, software, supervision, validation, visualization, writing –

review and editing. All authors have seen and approved the manuscript.

1348

Data availability statement

1350 The code associated with this project is stored in the GitHub repository linked here:

1351 https://github.com/IvanekLab/Romaine_ECO157_QMRA/ and is publicly accessible.

1352

Competing interests

1354 The authors declare no competing interests.

1355

1356 **Figure 1. (a)** Conceptual framework of the quantitative microbial risk assessment (QMRA)

1357 model for predicting illnesses due to *Escherichia coli* O157 (ECO157) contamination in the farm-

1358 to-fork fresh-cut romaine supply chain. **(b)** Conceptual framework of the difference equation

1359   model for population dynamics on the romaine batch and in soil batch during the 14-day

1360   preharvest countdown period.

1361

1362   **Figure 2.** Parameter IDs and definitions used in **(a)** Preharvest model **(b)** Postharvest model;

1363   and **(c)** Parameters varying by the harvest month of romaine production

1364

Figure 3. Tornado plot showing the impact of variable parameters on the preharvest output:

the predicted *E. coli* O157:H7 (ECO157) count on romaine batches after harvest, ready for

processing. The Spearman rank-order correlation coefficients (SRCC) are shown next to each

parameter bar. Only variables statistically significant after the Bonferroni correction and with a

correlation coefficient > 0.05 are shown. All values are positive, indicating positive correlation

(i.e., an increase in each parameter value increases ECO157 count in romaine batches).

1371

Figure 4. *Escherichia coli* O157:H7 (ECO157) counts in contaminated romaine batches in the

baseline model (with all four sources of preharvest contamination accounted for), as well as in

models considering each individual source of preharvest contamination separately **(a)**. The

ECO157 prevalence in romaine batches contributed by each individual preharvest

contamination source included in the model, one at a time **(b)**. ECO157 counts in contaminated

romaine batches in the baseline irrigation model (considering only the three irrigation systems

with no other preharvest sources), as well as in irrigation models considering each individual

irrigation source contamination separately **(c)**. The ECO157 prevalence in romaine batches

contributed by each individual irrigation contamination source included in the irrigation model, one at a time **(d)**.

Figure 5. Preharvest model regression tree predicting the mean *Escherichia coli* O157:H7 (ECO157) counts in romaine batches under various preharvest conditions. Branch labels indicate the rules used for splitting each node, and the circled numbers at the end of each branch indicate the predicted mean ECO157 counts, expressed in CFU/romaine batch. The intensity of the circle colors was proportional to the value of predicted ECO157 counts for a given branch (blue: low; red: high). Underneath each circle (e.g., 0), the percentage of model iterations falling in that branch (e.g., 89%) are displayed (e.g., 89% is shown under 0, indicating that a mean ECO157 count of 0 CFU/romaine batch was observed for this branch, which appeared in 89% of 100,000 iterations).

Figure 6. Tornado plot showing the impact of variable parameters on the postharvest output, i.e., the median predicted annual number of *E. coli* O157:H7 (ECO157) illness cases. The spearman rank-order correlation coefficients (SRCC) are shown next to each parameter bar. Only variables statistically significant after the Bonferroni correction and with a correlation coefficient > 0.05 are shown. Positive values indicate positive correlation (i.e., an increase in a given parameter increases the number of illness cases), while negative values indicate negative correlation (i.e., an increase in a given parameter decreases the number of illness cases).

Figure 7. Regression trees for postharvest model, predicting illness outcome: **(a)** the mean

number of illness cases from *Escherichia coli* O157:H7 (ECO157), **(b)** ECO157 counts in

contaminated servings in CFU, and **(c)** ECO157 prevalence in romaine servings (unitless). Branch

labels indicated the rules used for splitting each node, and the circled numbers at the end of

each branch indicated the predicted mean value for the outcomes. The intensity of the circle

colors was proportional to the value of predictions at the branch (blue: low; red: high).

Underneath each circle (e.g., 63 cases), the percentage of model iterations falling in that branch

(e.g., 12%) were displayed (e.g., 12% was shown under 63, indicating that a mean illness case

number of 63 was observed for this branch, which appeared in 12% of 100,000 iterations).

1410

1411 **Table 1. Description of the scenarios and changes in the illness outcome for potential risk**

1412 **management strategies**

Scenario	Definition (Reference)	ID of parameters affected (unit)	Parameter value in the scenario	Predicted illness cases resulted from the tested scenario: 5th; 50th; 95th percentile
Baseline	No changes made in the model.	None	All parameters were at their baseline conditions.	42; 19,040; 3.6×10^6
Cattle vaccination	ECO157 counts were reduced by 1-Log_{10}CFU/g in BSAAO and cattle feces [93].	C_{man} (Log_{10}CFU/g BSAAO) and C_{cow} (Log_{10}CFU/g feces)	$C_{man} = Uniform(3.1, 8.4) - 1$ $C_{cow} = Pert(0.89, 3.08, 8.4) - 1$	43; 14,113; 2.6×10^6 Median number of cases reduced by 25.9%.
Surface water treatment for overhead spray irrigation by chlorine	ECO157 counts in surface water were reduced when chlorine was applied [94].	C_{irr_s} (CFU/ml)	$\dfrac{C_{irr_s}}{10^{\text{Uniform}(1.00, 2.78)}}$	27; 1,803; 4.7×10^5 Median number of cases reduced by 90.5%.
Surface water treatment for overhead spray irrigation by peracetic acid (PAA)	ECO157 counts in surface water were reduced when PAA was applied [94,95].	C_{irr_s} (CFU/ml)	$\dfrac{C_{irr_s}}{10^{\text{Uniform}(2.48, 3.09)}}$	20; 608; 2.7×10^5 Median number of cases reduced by 96.8%.
Surface water treatment for overhead spray irrigation by Ultraviolet radiation (UV)	ECO157 counts in surface water were reduced when UV was applied [94,96].	C_{irr_s} (CFU/ml)	$\dfrac{C_{irr_s}}{10^{\text{Uniform}(1.30, 2.91)}}$	27; 1,705; 4.4×10^5 Median number of cases reduced by 91.1%.
Transition from overhead spray to furrow irrigation (i.e., all irrigation was through either furrow or drip)	All overhead spray irrigated fields were transitioned to furrow irrigated fields.	Prp_O and Prp_F (unitless)	$Prp_O =$ $CC: 0.0; \ DR: 0.0; \ Y: 0.0$ $Prp_F =$ $CC: 0.4; \ DR: 0.7; \ Y: 1.0$	12; 715; 6.3×10^5 Median number of cases reduced by 96.3%.

Transition from overhead spray and furrow to drip irrigation (all irrigation was through drip)	All overhead spray and furrow irrigated fields were transitioned to drip irrigated fields.	Prp_O, Prp_F and $Prpr_D$ (unitless)	Prp_O and Prp_F= $CC: 0.0;\ DR: 0.0;\ Y: 0.0$ Prp_D= $CC: 1.0;\ DR: 1.0;\ Y: 1.0$	14; 706; 5.3×10^5 Median number of cases reduced by 96.3%.
Produce wash alternatives with varying Log_{10}CFU reductions	Alternative produce washes at postharvest with varying reductions were used instead of the baseline chlorine wash (all scenarios, except CC_{ch5} [97] are based on assumptions).	CC_{ch20} (Log_{10}CFU)	CC_{ch20}= $Pert(0.01,0.52,1.03)$ was changed to $CC_{ch1}= Pert(0.5,1.0,1.5)$	31; 7,251; 1.2×10^6 Median number of cases reduced by 61.9%.
			$CC_{ch2}= Pert(1.0,1.5,2.0)$	25; 2,426; 4.1×10^5 Median number of cases reduced by 87.3%.
			$CC_{ch3}= Pert(1.5,2.0,2.5)$	22; 829; 1.4×10^5 Median number of cases reduced by 95.7%.
			$CC_{ch4}= Pert(2.0,2.5,3.0)$	21; 307; 5.3×10^4 Median number of cases reduced by 98.4%.
			$CC_{ch5}=Uniform(3.0,5.0)$	18; 56; 5.3×10^3 Median number of cases reduced by 99.7%.
Temperature reduction during retail	Temperature distribution was changed [98,99].	T_{retail} (°C)	T_{retail}= $Normal(4.4441,2.9642,$ $Truncate(0,20.56))$ was changed to $Uniform(0,4.4)$	38; 15,792; 5.5×10^5 Median number of cases reduced by 17.1%.
Consumer wash	62% of consumers washed the product (yielding a median of 0.99 Log_{10} CFU ECO157 reduction before consumption) [100].	CC_{conw} (unitless)	CC_{conw}=0 was changed to $Triang(0.65,0.99,0.99)\times$ 0.62	32; 13,717; 1.2×10^6 Median number of cases reduced by 28.0%.
Clustering effect	Clustering was allowed during partitioning of a romaine batch to packages and package to servings. Smaller $Clstr_{parb}$ or $Clstr_{parp}$ values indicated higher clustering of cells [46,85].	$Clstr_{parb}$ (unitless), $Clstr_{parp}$ (unitless)	$Clstr_{parb}$=0.1 $Clstr_{parp}$=0.1	38; 9,862; 2.8×10^5 Median number of cases reduced by 48.2%.
			$Clstr_{parb}$=1 $Clstr_{parp}$=1	44; 20,103; 2.9×10^6 Median number of cases increased by 5.6%.

1413

a

Preharvest

CFU/romaine batch
CFU/soil batch

| **Growing** |
| *CC & Growth* |

◦ Runoff
◦ Wildlife
◦ BSAAO
◦ Irrigation

■ Cattle vac
■ Tr from O to F irr
■ Tr to D irr
■■■ Chlorine, PAA or UV irr water treatments

CFU/romaine batch
CFU/soil batch

| **Waiting time** |
| *Growth* |

◦ Runoff
◦ Wildlife
◦ BSAAO

CFU/romaine batch
CFU/soil batch

| **Harvest** |
| *CC* |

▪ By blades

Postharvest

CFU/romaine batch

| **Cold storage** |
| *Growth* |

▪ At facility, before wash

CFU/romaine package

| **Processing** |
| *Inactiv & Partitioning* |

▪ Wash, cut, pack
■ Chlorine wash alternative

CFU/romaine package

| **Retail** |
| *Growth* |

▪ Storage & display
■ Temperature reduction

CFU/romaine package

| **Transportation** |
| *Growth* |

▪ Retail to home

CFU/romaine serving

| **Consumer** |
| *Growth & Partitioning* |

▪ Storage & consumption
■ Consumer wash

| **Number of ECO157 per contaminated fresh-cut romaine lettuce serving at point-of-consumption** |

| **Predicted number of annual illnesses in the USA** |

❖ Beta-Poisson dose-response
❖ Per-capita annual fresh-cut romaine lettuce consumption in the US

KEY
■ Risk intervention
◦ Contamination source
❖ Parameter
▪ Modeling detail
Process (in each box)
Unit (left column)

BSAAO: Biological soil amendment of animal origin**; Cattle vac:** Cattle vaccination; **Tr from O to F irr:** Transition from overhead spray to furrow irrigation; **Tr to D irr**: Transition from overhead spray and furrow to drip irrigation; **Chlorine, PAA or UV irr water treatments**: Surface irrigation treatments either with chlorine, peracetic acid or ultraviolet radiation; *CC*: Cross-contamination; *Inactiv*: Inactivation; *ECO157*: *Escherichia coli* O157:H7.

BSAAO: Biological soil amendment of animal origin; **ECO157**: *Escherichia coli* O157:H7.

Parameter	Description (units)	Parameter	Description (units)
t	Day of the countdown period in the difference equation model (1)-(6) (day)	Tr_{manO}	Fraction of the top 1cm layer of soil batch transferred to a romaine batch via splash caused by overhead spray irrigation per irrigation event (unitless)
A_{field}	Area of the field (approximately 1-acre field) (cm^2)	Tr_{manF}	Fraction of the top 1cm layer of soil batch transferred to a romaine batch via splash caused by furrow irrigation per irrigation event (unitless)
A_{batch}	Area of a field that grows a romaine batch ("batch area", 1/12 of the field area) (cm^2)	Tr_{manD}	Fraction of the top 1cm layer of soil batch transferred to a romaine batch via splash caused by drip irrigation per irrigation event (unitless)
N_{plant}	Number of plants harvested from the field (romaine plants)	M_{irr}	Mass of soil transferred to a romaine batch via splash per irrigation event (g/day)
$Plant_{width}$	The area that a single mature romaine plant occupied on the field, length x width of one plant (cm^2)	Tr_{ecirr}	Fraction of ECO157 cells transferred from irrigation water to romaine (unitless)
$Plant_{space}$	The area allocated for one romaine plant in the field (cm^2)	DiF_{irr}	Dilution factor for ECO157 cells in irrigation water in the top 1cm layer of soil batch (unitless)
$Plant_{barea}$	Number of plants grown in a batch area (romaine plants)	LN_o	ECO157 count in a romaine batch at the start of the 14-day countdown period (t=0) (CFU)
$Plant_{wght}$	Weight of one mature romaine plant (g)	CC_{UV}	Log$_{10}$ ECO157 count reduction of surface water contamination by UV treatment before overhead spray irrigation (Log$_{10}$CFU)
f_{loc}	Select geographic location where the romaine batches were grown (unitless). Notations: CC: Coastal Region and Central Valley, CA DR: Desert (Imperial) Region, CA Y: Yuma Region, AZ	CC_{PAA}	Log$_{10}$ ECO157 count reduction of surface water contamination by PAA treatment before overhead spray irrigation (Log$_{10}$CFU)
f_{month}	Select the month of harvest for the given geographic location (unitless). Notations: January: 1, February: 2, March: 3, April: 4, May: 5, June: 6, July: 7, August: 8, September: 9, October: 10, November: 11, December: 12	CC_{Ch}	Log$_{10}$ ECO157 count reduction of surface water contamination by Chlorine treatment before overhead spray irrigation (Log$_{10}$CFU)
f_w	Select the source of irrigation water (unitless). Notations: S: Surface water W: Well water	$C_{cowprep}$	Log$_{10}$ ECO157 count on fresh cattle feces (Log$_{10}$CFU/g)
f_{irr}	Select the irrigation type for the given geographic location (unitless). Notations: O: Overhead spray irrigation F: Furrow irrigation D: Drip irrigation	C_{cowtot}	Log$_{10}$ ECO157 count on fresh cattle feces, accounting for the effect of ECO157 vaccine intervention, if implemented (Log$_{10}$CFU/g)
$Modeldays$	Countdown period before harvest, modeled via a set of difference equations (day)	C_{cow}	ECO157 count on fresh cattle feces (CFU/g)
$edge$	Batch area was at the edge of the field and thus could be contaminated by runoff (unitless): no=0 (baseline), yes=1	C_{cowP}	ECO157 count on any cattle feces, which returns 0 CFU for non-contaminated feces and C_{cow} for contaminated feces (CFU/g)
C_{man}	ECO157 count per g of BSAAO (CFU/g)	C_{pig}	Log$_{10}$ ECO157 count on fresh swine feces (Log$_{10}$CFU/g)
Sc_{vac}	Cattle vaccination intervention against ECO157 (unitless): no=0 (baseline), yes=1	C_{pigP}	ECO157 count on any swine feces, which returns 0 CFU for non-contaminated feces and C_{pig} for contaminated feces (CFU/g)
Rh_{con}	Volume of 1g BSAAO/swine feces/cattle feces (cm^3/g)	Pr_{pig}	Probability of feral swine entering the field on day t (unitless)
Dh_{con}	Depth of tiling on the field (cm)	$Pr_{pigCountt}$	Wildlife intrusion occurred on day t (unitless): no=0, yes=1
M_{man}	Mass of BSAAO applied on the batch area (g)	$Pr_{pigCountSum}$	Total number of days during the 14-day countdown when wildlife intrusion occurred (day)
DiF_{man}	Dilution of ECO157 cells in BSAAO in the volume of the tilled soil batch layer (unitless)	$Mdef_{pig}$	Mass of feces excreted by a feral swine per day (g swine feces/day)
Pr_{man}	BSAAO was used (unitless): no=0, yes=1	$Tdef_{pig}$	Number of times a feral swine defecated per day (times/day)
P_{man}	BSAAO applied on a field was contaminated with ECO157 (unitless): no=0, yes=1	$Ndef_{pig}$	Number of times feral swine (between 1-5 swine) defecated on the field (times/day)
C_{manP}	ECO157 count in any BSAAO applied soil, which returns 0 CFU for a non-contaminated soil and C_{man} for a contaminated soil (CFU/g)	M_{pig}	Mass of feces excreted by feral swine intruding the batch area (g)
P_{irrS}	ECO157 prevalence in surface water (unitless)	DiF_{pig}	Dilution factor for ECO157 cells in feral swine feces in soil batch associated with plants near which the swine defecated (unitless)
P_{irrW}	ECO157 prevalence in well water (unitless)	M_{cow}	Mass of cattle runoff (carrying fresh cattle feces only) introduced into the edge batch area soil per runoff event (g/day)
P_{irr}	Irrigation water contaminated with ECO157 applied during the countdown period (unitless): no=0, yes=1	DiF_{cow}	Dilution factor for ECO157 cells in cattle runoff introduced into the top 1cm layer of soil batch (unitless)
Sc_{wt}	Water treatment intervention applied prior to the overhead spray irrigation (O) using surface water (S) (unitless): None=0 (baseline), Ultra-Violet (UV)=1, Peracetic Acid (PAA)=2, Chlorine=3	Pr_{LtoS}	Proportion of fecal matter (swine and runoff) landing on romaine rather than in soil (unitless)
C_{irrS}	ECO157 count per ml of contaminated surface water post treatment (CFU/ml)	Tr_{ecsoil}	Proportion of ECO157 cells in soil transferring to romaine during a rain or irrigation event (unitless)
C_{irrW}	ECO157 count per ml of contaminated well water (CFU/ml)	R_{dman}	ECO157 decay rate for non-persisters in BSAAO amended soil (Log$_{10}$CFU/day)[a]
C_{irr}	ECO157 count per ml of contaminated irrigation water applied on day t (CFU/ml)	R_{dL}	ECO157 decay rate for non-persisters on romaine in field (1/day) [a,b]
C_{irrP}	ECO157 count on any irrigation water, which returns 0 CFU for a non-contaminated water and C_{irr} for a contaminated water (CFU/ml)	R_{ds}	ECO157 decay rate for non-persisters in soil (1/day) [a]
$type_{irr}$	Probability the selected irrigation type was occurring on a given day (unitless)	R_{dmanP}	ECO157 decay rate for persisters in BSAAO amended soil (Log$_{10}$CFU/day) [a]
Pr_O	Probability of overhead spray irrigation on day t (unitless)	R_{dLP}	ECO157 decay rate for persisters on romaine in field (1/day) [a]
Pr_F	Probability of furrow irrigation on day t (unitless)	R_{dsP}	ECO157 decay rate in soil for persisters (1/day) [a]
Pr_{D_t}	Probability of drip irrigation on day t (unitless)	$NtoP_{man}$	Switch rate from normal to persister in BSAAO amended soil (1/day) [a]
W_O	Volume of water applied through overhead spray irrigation to a romaine batch per an irrigation event (ml/day)	$NtoP_L$	Switch rate from normal to persister on romaine (1/day) [a]
W_F	Volume of water applied through furrow irrigation to a romaine batch per an irrigation event (ml/day)	$NtoP_S$	Switch rate from normal to persister in soil (1/day) [a]
W_D	Volume of water applied through drip irrigation to a romaine batch per an irrigation event (ml/day)	$Pmax_{man}$	Maximum fraction of persisters in BSAAO amended soil (unitless)
$Daytogrow$	Days from seeding to harvest (day)	$Pmax_S$	Maximum fraction of persisters in soil (unitless)
D_{man}	Number of days before BSAAO application (day)	$Pmax_L$	Maximum fraction of persisters on romaine (unitless)
g_{irr}	Volume of water applied per an irrigation event in a batch area (ml/day)	$BwaitP$	Waiting time: The period before harvest during which overhead spray and furrow irrigation was stopped (day)
V_O	Volume of irrigation water captured per g of romaine per an overhead spray irrigation event (ml/g/day)	Tr_{blade}	Average amount of soil attached to a harvesting blade after contact with soil in one cross contamination event (g/blade)
V_F	Volume of irrigation water captured per romaine batch per a furrow irrigation event (ml/day)	$Tr_{ecblade}$	Proportion of ECO157 cells transferred from a harvesting blade to a romaine plant in one cross contamination event (unitless)
V_D	Volume of irrigation water captured per romaine batch per a drip irrigation event (ml/day)	N_{blade}	ECO157 count in soil attached on blades (CFU/soil batch x soil batch/blade=CFU/blade)
v_{irr}	Volume of irrigation water captured by romaine batch per an irrigation event (ml/day)	CC_{blade}	ECO157 cells transferred from harvesting blades to a romaine batch (CFU/romaine batch)
		C_{L_h}	Preharvest outcome: ECO157 count on harvested romaine batch (CFU)

b

Parameter	Description (units)	Parameter	Description (units)
$C_{L_{hf}}$	*E. coli* O157:H7 (ECO157) count in a contaminated harvested romaine batch (Log$_{10}$CFU/romaine batch)	D_{trnspt}	Transportation die-off rate (Log$_{10}$CFU/h)
$P_{L_{hf}}$	ECO157 prevalence in a harvested romaine batch (unitless)	$GorD_{trnspt}$	Growth (G) or die off (D) based on transport temperature
t_{coldst}	Cold storage time (h)	CC_{trspt}	Change in ECO157 count during transportation (Log$_{10}$CFU)
$T_{endcoldst}$	Temperature after cold storage (°C)	C_{trnspt}	ECO157 count in a contaminated package after transportation (Log$_{10}$CFU/romaine package)
T_{coldst}	Temperature during cold storage (°C)	t_{fcons}	Time to first consumption (h)
G_{coldst}	Cold storage growth rate (Log$_{10}$CFU/h)	t_{lcons}	Time to last consumption (h)
D_{coldst}	Cold storage die-off rate (Log$_{10}$CFU/h)	t_{home}	Home storage time (h)
$GorD_{coldst}$	Growth (G) or die off (D) based on cold storage temperature (Log$_{10}$CFU/h)	T_{home}	Home storage temperature (°C)
CC_{coldst}	Change in ECO157 count during cold storage (Log$_{10}$CFU)	G_{home}	Home growth rate (Log$_{10}$CFU/h)
$C_{endcoldst}$	ECO157 count on a contaminated romaine batch after cold storage, going to processing (CFU/romaine batch)	D_{home}	Home die-off rate (Log$_{10}$CFU/h)
B	Romaine batch size (g)	$GorD_{home}$	Growth (G) or die off (D) based on home storage temperature
Sc_{wash}	1 or 2 based on the scenarios below, (baseline=1)	CC_{home}	Change in ECO157 count during home storage (Log$_{10}$CFU)
CC_{ch20}	Log$_{10}$CFU ECO157 reduction with chlorine washing (Log$_{10}$CFU/g romaine)	C_{home}	ECO157 count in a contaminated package after home storage (Log$_{10}$CFU/romaine package)
CC_{ch30}	Chlorine wash alternative with a median of 3.22 Log$_{10}$CFU reduction instead of 0.52 in the baseline chlorine wash (Log$_{10}$CFU)	$P0_{ser}$	Probability a non-contaminated serving was partitioned from a contaminated romaine package
$CC_{w_{agent}}$	ECO157 reduction associated with the scenario selected for the washing step (Log$_{10}$CFU)	P_{ser}	ECO157 prevalence on a romaine serving
C_{wash}	ECO157 count on a contaminated romaine batch after washing (CFU/romaine batch)	Sc_{clstrp}	Clustering in package: no=0 (baseline), yes=1
$P0_{pack}$	Probability a non-contaminated package is partitioned from a contaminated romaine batch (unitless)	$Clstr_{parp}$	Clustering parameter
P_{pack}	ECO157 prevalence on a romaine package (unitless)	Ser	Serving size (g romaine/ romaine serving)
Sc_{clstrb}	Clustering in romaine batch: no=0 (baseline), yes=1	Ser_{pack}	Number of servings in a package (romaine servings)
$Clstr_{parb}$	Clustering parameter	$P_{contserp_{prep}}$	Distribute ECO157 cells to a serving when clustering (unitless)
$W_{romainepa}$	Weight of shredded romaine per package (8 oz) (g)	$P_{contser}$	Proportion of ECO157 cells from contaminated package in a serving (unitless)
A_{pack}	Number of packages in a romaine batch (packages)	C_{serH}	ECO157 count in a contaminated serving after home storage (Log$_{10}$CFU/serving romaine)
$P_{contpack_{dist}}$	Distribute ECO157 cells to a package when clustering	Sc_{conw}	Consumer wash romaine before consumption: no=0 (baseline), yes=1
$P_{contpack}$	Proportion of ECO157 cells from a contaminated romaine batch in a package (unitless)	P_{conw}	Probability a USA consumer washes the prepacked romaine (unitless)
C_{pack}	ECO157 count in a contaminated package (Log$_{10}$CFU/pack romaine)	$CC_{conw_{prep}}$	ECO157 reduction by consumer wash with water prep cell (Log$_{10}$CFU)
t_{retail}	Retail storage time (h)	CC_{conw}	Log$_{10}$CFU ECO157 reduction by consumer wash with water (Log$_{10}$CFU)
T_{retail}	Retail storage temperature (°C)	C_{serW}	ECO157 count in a contaminated serving after consumer wash (CFU/romaine serving)
G_{retail}	Retail growth rate (Log$_{10}$CFU/g romaine/h)	$alpha$	Dose-response parameter alpha (unitless)
D_{retail}	Retail die-off rate (Log$_{10}$CFU/g romaine/h)	$beta$	Dose-response parameter beta (unitless)
$GorD_{retail}$	Growth (G) or die off (D) based on retail temperature (Log$_{10}$CFU/h)	$Pill$	Probability of illness as a result of consumption of a contaminated romaine serving (unitless)
CC_{retail}	Change in ECO157 count during retail storage (Log$_{10}$CFU)	$USpop$	USA population (person)
C_{retail}	ECO157 count in a contaminated package after retail storage (Log$_{10}$CFU/package romaine)	A	Annual romaine consumption in the USA (g/consumer/year)
$t_{transport}$	Transportation time (h)	Np	Number of romaine servings consumed per person per year (romaine serving/consumer/year)
T_{trhom}	Package temperature before putting into home refrigerator (°C)	Nsc_{US}	Total number of romaine servings consumed per year in the USA (romaine serving/year)
T_{trnspt}	Temperature during transportation (°C)	$Ncsc_{US}$	Number of contaminated romaine servings consumed per year in the USA (romaine serving/year)
G_{trnspt}	Transportation growth rate (Log$_{10}$CFU/)	$Ncase_{US}$	Number of illness cases per year in the USA (person)

c

Parameter	Description (units)	Parameter	Description (units)
P_{sun_m}	Probability of a sunny day (unitless)	R_{sun_m}	Solar decay rate (1/day)
P_{cow_m}	ECO157 prevalence in fresh cattle feces (unitless)	$Prec_{norm_m}$	Precipitation level normalized (unitless)
P_{pig_m}	ECO157 prevalence in fresh feral swine feces (unitless)	$Prec_{actual_m}$	Actual (average) level of precipitation during the year (mm)
h_{sun_m}	Average sunny hours per day (h)	$T_{harvest_m}$	Temperature at harvest (°C)

C_{irr_P}: ECO157 count per 100 ml of contaminated irrigation water applied on day t; C_{pig_p}: ECO157 count on any swine feces, which returns 0 CFU for non-contaminated feces and C_{pig} for contaminated feces; C_{cow_P}: ECO157 count on any cattle feces, which returns 0 CFU/g for non-contaminated feces and C_{cow} for contaminated feces; $type_{irr}$: Probability the selected irrigation type is occurring on a given day; **O**: Overhead spray irrigation; **F**: Furrow irrigation; **D**: Drip irrigation; $Pr_{pig_{CountSum}}$: Total number of days during the 14-day countdown when wildlife intrusion occurred; $edge$: Batch area was at the edge of the field and thus could be contaminated by runoff.

a

$C_{Lh_f} < \dfrac{2.7\ Log_{10}CFU}{romaine\ batch}$ $C_{Lh_f} \geq \dfrac{2.7\ Log_{10}CFU}{romaine\ batch}$

$C_{Lh_f} < \dfrac{1.6\ Log_{10}CFU}{romaine\ batch}$ $C_{Lh_f} \geq \dfrac{1.6\ Log_{10}CFU}{romaine\ batch}$ $C_{Lh_f} < \dfrac{3.9\ Log_{10}CFU}{romaine\ batch}$ $C_{Lh_f} \geq \dfrac{3.9\ Log_{10}CFU}{romaine\ batch}$

$T_{home} < 5°C$ $T_{home} \geq 5°C$ $T_{home} < 5°C$ $T_{home} \geq 5°C$ $T_{home} < 5.1°C$ $T_{home} \geq 5.1°C$ $T_{home} < 5°C$ $T_{home} \geq 5°C$

$T_{retail} < 7°C$ $T_{retail} \geq 7°C$ $T_{retail} < 5°C$ $T_{retail} \geq 5°C$

63	500	1,300	8,000	16,000	126,000	100,000	1,000,000	500,000	6,300,000
12%	4%	15%	5%	23%	8%	19%	5%	5%	4%

c

C_{Lh_f}: ECO157 count on a contaminated romaine batch after harvest; T_{home}: Home storage temperature; T_{retail}: Retail storage temperature; t_{lcons}: Time to last consumption; P_{L_hf}: ECO157 prevalence in a romaine batch after harvest; t_{retail}: Retail storage time.