Full Title:

Long COVID affects working memory: Assessment using a single rapid online test

Short title:

Long COVID affects working memory

Aziz UR Asghar[1,2], Hong Kiu Yuen[1,3], Murat Aksoy[1], Abayomi Salawu[1,3], Heidi A Baseler[2,4,5]*

[1] Hull York Medical School, University of Hull, Hull, United Kingdom, [2] York Biomedical Research Institute, University of York, York, United Kingdom, [3] Hull University Teaching Hospitals National Health Service (NHS) Trust, Hull, United Kingdom, [4] Hull York Medical School, University of York, York, United Kingdom, [5] Department of Psychology, University of York, York, United Kingdom

***** Corresponding author

Email: heidi.baseler@hyms.ac.uk

Abstract

Long COVID, or post-COVID-19 syndrome, is characterized by persistent symptoms following SARS-CoV-2 infection, including cognitive impairments such as brain fog that adversely affects quality of life. The aim of this study was to evaluate the impact of long COVID on working memory using a single rapid online anonymous survey and working memory quiz. We analyzed working memory scores in relation to long COVID status (clinically diagnosed, self-reported, and non-long COVID), age, number of COVID-19 infections and long COVID duration, and subjective ratings of brain fog severity and overall life impact of symptoms. The study recruited 1064 participants aged 16 to over 85 years (83% from the United Kingdom) with 39% reporting long COVID. The results demonstrate that long COVID significantly impairs working memory, with the greatest deficits observed in clinically diagnosed cases. In addition, we found that working memory declined with age and multiple COVID-19 infections particularly in the diagnosed long COVID group. Moreover, individuals in the diagnosed long COVID group reported experiencing the most severe brain fog and the greatest impact of long COVID symptoms on their lives, and both of these factors were strongly correlated with lower memory scores. Participants who had been living with long COVID for a longer duration had lower memory scores. However, this relationship was likely influenced by the SARS-CoV-2 variant, as individuals with prolonged long COVID symptoms were more likely to have been infected with earlier variants (alpha and wild-type) which are linked to more severe disease outcomes. Our results suggest that long COVID is associated with significant working memory impairments, emphasizing the need for targeted interventions and support strategies for those with long COVID to address memory problems.

Introduction

Long COVID, also known as post COVID-19 syndrome/condition, broadly refers to the signs and symptoms that develop during or after an infection with the SARS-CoV-2 virus that persists over a number of months and has no alternative explanation [1, 2]. A systematic review of long COVID prevalence, which included 194 studies with 735,006 participants, found that 45% of those who survived COVID-19 continued to experience symptoms at approximately four months [3].

Approximately 1.9 million people in the United Kingdom self-reported long COVID symptoms with 41% experiencing symptoms for at least 2 years and 79% reported that their symptoms negatively impacted their daily activities [4]; over 200 long COVID symptoms have been documented, and the most commonly reported were fatigue (72%), difficulty concentrating (51%), muscle aches (49%) and shortness of breath (48%).

Substantial cognitive performance deficits in tasks requiring memory, attention and reasoning have been reported for people with COVID-19 symptoms that persisted for up to two years [5-11]. The cognitive impact of long COVID, often generally described as 'brain fog' (including problems with memory and performing daily tasks), is a major contributor to poor functioning and reduced quality of life experienced by patients with long COVID [12-15]. A recent systematic review and meta-analysis of studies evaluating brain fog, cognitive function and mental health symptoms in people with long COVID found that the average overall prevalence of memory loss was 21% and as high as 61% [16]

Working memory is the process of temporarily storing and retrieving information relevant to performing the current task [17, 18]. Working memory is integral in everyday tasks including problem solving, holding a conversation, reading comprehension and decision making. Working memory deficits have been consistently reported when testing cognitive function in long COVID [19-22].

Studies investigating working memory deficits related to COVID-19 or long COVID are often measured within the context of a battery of multiple cognitive tests [5, 19-25]. This is potentially problematic for two reasons. Firstly, when embedded within a more complex cognitive task (for example, spatial mental rotation), it is difficult to interpret performance on the working memory component alone. Moreover, it is possible that a participant's performance on a working memory task could be affected by previous different cognitive tasks. Secondly, undertaking numerous cognitive tests within a given study requires statistical correction for multiple comparisons, which reduces the statistical power and increases the likelihood of type II errors (false negatives). Conversely, if statistical correction is not applied to correct for multiple cognitive tests, there is the risk for type I errors (false positives). Since the majority of people with long COVID report fatigue and difficulty in concentrating as major symptoms, completing a lengthy survey with multiple cognitive tests would be burdensome and may decrease completion rates or yield unreliable data. We have previously shown, using a single cognitive task requiring minimal time and effort, that working memory was negatively impacted in people who had been infected with COVID-19 compared to those who reported not having had COVID-19 [26].

The main aim of our study was to investigate the impact of long COVID on working memory function. To capture a broad demographic sample, we distributed an online survey with a single working memory quiz that could be completed relatively rapidly. We utilized the same memory quiz as we have used previously to explore working memory function in COVID-19, which was anonymous and incorporated elements of gamification [26]. In our analysis we evaluate the impact of several variables on objective memory scores including long COVID status (diagnosed long COVID, self-reported long COVID and non-long COVID), age, gender, number of COVID-19 infections and long COVID duration. In addition, we examined the relationship between objective memory scores and participant subjective ratings of brain fog, and the overall impact of long COVID symptoms on their life.

Materials and methods

Ethics

The study was undertaken in accordance with the Declaration of Helsinki [27]. Ethical approval was given by the Hull York Medical School Ethics Committee (Reference 22-23 41). The Long COVID Online Rapid Objective Neuro-Memory Assessment (L-CORONA) was made publicly available, and data collected from the 16th of March 2023 to the 31st of December 2023. The link to the online survey and memory quiz was disseminated mainly via direct communication through personal and professional contacts as well as through local intranets, social media (including long COVID support groups) and advertisements posted in public places. Participants aged 16 years old and over were invited to complete the survey

and memory quiz regardless of whether they had long COVID. After providing information at the beginning of the survey about the study, written informed consent was obtained by asking participants, '*Do you consent to take part?*', and only those who answered '*Yes*' were allowed to proceed. The L-CORONA survey and memory quiz complied with General Data Protection Regulations (GDPR), was fully anonymous, and no personal contact details were acquired. The Qualtrics platform (University of York license) collected and stored the encrypted data, and access was restricted to the study investigators only. No compensation was provided to participants for completing the assessment.

Long COVID definitions

Some studies reporting the impact of COVID-19 on cognitive function do not provide an explicit definition of long COVID but instead have inferred long COVID status based upon ongoing symptomology [21]. The NHS England's definition of post COVID-19 syndrome, which is based on the National Institute for Health and Care Excellence definition [1], is '*…signs and symptoms that develop during or after COVID-19 and continue for more than 12 weeks and are not explained by an alternative diagnosis*' [28]. This definition is broadly consistent with that of the World Health Organization which defines post COVID-19 condition (long COVID) as, '*…the continuation or development of new symptoms 3 months after the initial SARS-CoV-2 infection, with these symptoms lasting for at least 2 months with no other explanation*'[2]. In our current study, we adopted the NHS England definition of long COVID with a questionnaire designed to exclude people who had symptoms less than 12 weeks after being infected with COVID-19. In addition, we designed our study to

compare people diagnosed with long COVID by a medical professional with those who self-reported having long COVID.

Long COVID survey and memory quiz structure

The long COVID assessment was similar in structure to our previous COVID-19 online survey and memory quiz [26]. To encourage participant recruitment, the long COVID assessment was designed to be engaging and relatively quick to complete. The Qualtrics platform (University of York license) was used to develop the online survey and memory quiz. Participants accessed the long COVID assessment using a Uniform Resource Locator (URL) or a quick response (QR) code, which made it available on smartphones, tablets, laptops and PCs with internet access. No time limit was imposed upon participants to complete the long COVID assessment. Screenshots of some survey questions and a sample of the memory quiz are illustrated in Fig 1.

Fig 1. The 'Long COVID Survey and Memory Quiz' presented on a smartphone. a) Example questions from the survey, and examples of the animal, number, fruit and object categories shown from the memory quiz. **b)** A series of screenshots representing a single trial from the working memory quiz.

The survey component consisted of a series of '*Yes/No*' questions asking participants about their long COVID status. First, we asked, '*Have you been diagnosed with long COVID (post-COVID-19 syndrome) by a medical professional?*'. If a participant answered '*Yes*', they were designated into the diagnosed long COVID group. Those answering '*No*' were then asked, '*Do you think that you may have long COVID?*', and we included the NHS England definition of long COVID (post COVID-19 syndrome): '*Signs and symptoms that develop during or after COVID-19 and continue for more than 12 weeks and are not explained by an alternative diagnosis.*' [28]. Those that answered '*Yes*' to this question were designated as the self-reported long COVID group. Those who answered '*No*' to both questions were designated as the non-long COVID group. If they were diagnosed with long COVID or self-reported having long COVID, they were asked how many months it had been since their long COVID symptoms started (using a drop-down menu, from less than 3 months to 60 months). We next asked long COVID participants to subjectively rate the overall life impact of their long COVID symptoms (using *'None', 'Mild', 'Moderate' or 'Severe'*). All participants (regardless of long COVID status) were asked whether they had been infected with COVID-19 in the last 12 weeks ('*Yes/No*'), and how many times they had been infected with COVID-19 ('*Not had COVID-19*', '*One*', '*Two*', '*Three*', or '*Four or more times*'). All participants were also asked to subjectively rate their level of '*brain fog*' (using *'None', 'Mild', 'Moderate' or 'Severe'*). The following definition of '*brain fog*' was provided to aid participants: '*for example, difficulty in thinking clearly or quickly, concentrating, having a conversation, remembering things, or getting tired easily after mental effort'.* This definition of brain fog was based on our previous findings showing that these COVID-19 cognitive symptoms were highly correlated with one another [26]. Finally, participants were asked to select their age range (16-17, 18-24, 25-34, 35-44, 45-54, 55-64, 65-74, 75-84 and 85+), gender, and country of current

residence. Following completion of these survey questions, participants next completed the objective working memory quiz. To increase levels of concentration and compliance, the participants were instructed to '*Please complete the quiz in a QUIET place WITHOUT DISTRACTIONS.*' The memory quiz consisted of the same visual working memory/recall task that we utilized previously to investigate the impact of COVID-19 on working memory [26]. In brief, the working memory quiz employed aspects of gamification such that the task was relatively easy to understand with familiar and engaging images. Participants progressed through the quiz with increasing difficulty levels and motivational feedback was given for responses. At the end of the quiz, participants were notified of their overall score along with the scores for the sub-categories. Four image categories were displayed to participants consisting of animals, numbers, fruits, and objects (Fig 1a). The stimulus set comprised of 55 unique images: animals, fruits, objects (15 unique images in each of these categories), and ten single-digit whole numbers (0 to 9). For each image category, two, three, four or five different random images were shown consecutively. Each image was presented for 500ms and the interstimulus interval was randomly varied between 500-600ms (Fig 1b). A grid of six images was presented and participants were instructed to select the images (two, three, four or five) viewed prior. Participant responses were not time restricted.

Data Analysis

Based on our inclusion criteria, participants under the age of 16 were excluded (N=3). Participants that had COVID-19 infections within the last 12 weeks were also excluded from the analysis (N = 146), as a recent COVID-19 infection is known to affect working memory [26]. In addition, we excluded one participant who reported having long COVID for 49

months before completing the survey/memory quiz on 21st March 2023. This meant that their long-COVID symptoms would have started on the 21st of February 2019, a time before COVID-19 was identified.

A score of 1 was given for each correct trial in the memory quiz, yielding a possible maximum objective memory score of 16. Statistical Package for the Social Sciences (SPSS Version 29.0.1.0, IBM Corporation, Armonk, N.Y., USA) was utilized to perform statistical analyses. To determine the relative contribution of each independent variable to the objective memory scores we used categorical regression (Optimal scaling-CATREG, SPSS). First, we assessed the contributions of long COVID status (three groups: non-long COVID, diagnosed long COVID and self-reported long COVID), age (nine groups: 16-17, 18-24, 25-34, 35-44, 45-54, 55-64, 65-74, 75-84 and 85+), number of COVID-19 infections (five groups: none, 1, 2, 3, and 4 or more) and gender (three groups: female, male, other/prefer not to say). A second categorical regression was performed using the long COVID groups only with the following variables: long COVID status (two groups: diagnosed long COVID and self-reported long COVID), age, number of COVID-19 infections, gender, and long COVID duration in months (from less than 3 months up to 60 months). Next, we investigated the effect of the significant individual variables arising from the categorical regression analysis on objective memory scores, using non-parametric statistics: Mann-Whitney U test and Kruskal-Wallis test with Dunn's pairwise comparisons (Bonferroni corrected). We evaluated relationships between pairs of variables using the Kendall's tau-b correlation coefficients. All testing was two-tailed, and we set statistical significance levels to p values less than 0.05.

Results

Demographics of participants

We included 1064 participants that fully completed the L-CORONA survey and memory quiz in the period between 17th March 2023 and 31st December 2023. Participant demographics are summarized in Table 1. In terms of long COVID status, 39% of participants had long COVID (N=415), and 61% did not have long COVID ('Non-Long COVID group'). Participants ranged from 16 to over 85 years of age, with the highest number of participants in the age range 45 to 54 years old. The majority of the participants resided in the United Kingdom (83%), with the remainder residing in 25 other countries around the globe. Over 94% of participants (N=1001) completed the survey and memory quiz within 15 minutes (median = 7.2 minutes): long COVID group (median = 7.3 mins, n=393); non-long COVID group (median = 6.9 minutes, n=608).

Table 1. Participant demographics.

Long COVID status	N
Non-Long COVID	649
Long COVID	415
Age (Years)	
16 - 17	14
18 - 24	128
25 - 34	184
35 - 44	223
45 - 54	262
55 - 64	188
65 - 74	49
75 - 84	15
85+	1
Gender	
Female	806
Male	244
Other	14
Countries	
UK	883
Non-UK	181

Categorical regression and objective memory scores

A categorical regression was used to assess the relative contributions of long COVID status (non-long COVID, diagnosed long COVID, or self-reported long COVID), age, number of COVID-19 infections and gender to memory scores in all participants. Table 2 shows that significant effects were found (all p-values < 0.001), in order of importance, for long COVID status (55.4%), followed by age (31.5%) and number of COVID-19 infections (13.3%). Gender did not have a significant effect on memory scores (p > 0.05).

Table 2. Categorical regression analysis and objective memory scores for non-long COVID, diagnosed long COVID and self-reported long COVID participants.

Adjusted regression coefficient R^2 = 0.114, Fisher's F-test = 13.479, p < 0.001				
Independent Variables	**Beta**	**F**	**p**	**Importance (Pratt)**
Long COVID status (non-long COVID, diagnosed long COVID, self-reported long COVID groups)	0.240	60.221	**< 0.001**	0.554
Age	-0.184	48.074	**< 0.001**	0.315
Number of COVID-19 infections	-0.110	7.686	**< 0.001**	0.133
Gender	0.011	0.402	0.669	-0.002

Significant p-values are given in bold.

A second categorical regression was performed for the two long COVID groups only (those who self-reported having long COVID or were diagnosed with long COVID) to assess the

additional contribution of long COVID duration on memory scores. Table 3 shows that the significant factors were, in order of importance: age (52.9%, $p < 0.001$), long COVID duration (26.0%, $p < 0.001$), and number of COVID-19 infections (15.5%, $p = 0.003$). Long COVID group (self-reported *versus* diagnosed long COVID) or gender did not have a significant effect on memory scores ($p > 0.05$).

Table 3. Categorical regression analysis and objective memory scores for diagnosed long COVID and self-reported long COVID participants only.

Adjusted regression coefficient R^2 = 0.079 Fisher's *F*-test= 3.207, **$p < 0.001$**				
Independent Variables	**Beta**	***F***	***p***	**Importance (Pratt)**
Age	-0.238	32.297	**<0.001**	0.529
Long COVID duration	-0.145	6.491	**<0.001**	0.260
Number of COVID infections	-0.139	4.678	**0.003**	0.155
Long COVID groups only (diagnosed long COVID, self-reported long COVID groups)	0.044	1.360	0.244	0.055
Gender	0.007	0.044	0.957	0.001

Significant *p*-values are given in bold.

Based on these categorical regressions (Tables 2 and 3), we selected the following main factors affecting objective memory scores and evaluated each of them independently: long COVID status, age, number of COVID-19 infections, and for the long COVID groups only long COVID duration.

Effect of long COVID status on working memory

Fig 2a shows that the objective memory scores are significantly reduced in the two long COVID groups combined (M = 14.15 ± 0.10) compared to the non-long COVID group (M = 15.02 ± 0.05, Mann-Whitney U = 167938.50, Z = -7.12, $p < 0.001$, r = 0.22). Next, we divided the long COVID group into sub-groups: those who suspected they may have long COVID (self-reported), and those who were diagnosed with long COVID by a medical professional. Fig 2b shows that objective memory scores were highest in the non-long COVID group but were significantly reduced in each of the long COVID sub-groups, with the lowest scores in the group diagnosed with long COVID. A Kruskal-Wallis test revealed a significant difference between the mean memory scores across groups [H(2) = 61.38, $p < 0.001$]. Dunn's pairwise comparison tests with Bonferroni correction revealed significant differences between the non-long COVID group (M = 15.02 ± 0.05) and the self-reported long COVID (M = 14.48 ± 0.14, $p = 0.0042$), and between the non-long COVID and diagnosed long COVID (M = 13.88 ± 0.15, $p < 0.001$) groups. In addition, there was a significant difference between the self-reported long COVID (M = 14.48 ± 0.14) and diagnosed (M = 13.88 ± 0.15, $p = 0.0031$) long COVID groups.

Fig 2. The effect of long COVID status on objective memory scores. a) Non-long COVID group compared to both long COVID groups combined (diagnosed plus self-reported long COVID), and **b)** Non-long COVID group compared to self-reported and diagnosed long COVID groups separately. Error bars represent standard error of the mean. *$p < 0.01$, **$p < 0.001$.

Effect of age on working memory

We found that objective memory scores decreased significantly with age for both the non-long COVID and the diagnosed long COVID groups but not for the self-reported long COVID group. A Kendall's tau-b correlation showed a negative relationship between mean memory score and age for both the non-long COVID (τb = –0.155, p < 0.001) and diagnosed long COVID (τb = –0.208, p < 0.001) groups across the nine age categories. The self-reported long COVID group did not have a significant relationship between age and mean memory scores (τb = –0.099, p = 0.098). Participants aged 16-17 years old were combined with those aged 18 to 24 years old, and all participants above 55 years old were combined into a single group to increase sample size due to the lower numbers of participants who completed the memory quiz in these age categories (see Table 1). Fig 3 shows the effect of age on memory scores for all three groups for the five age categories. A Kruskal-Wallis test was performed at each of the five age categories to determine differences between groups based on long COVID status. There was no significant effect of long COVID status on memory scores for the 16-24 year-old age category [H(2) = 4.201, p = 0.122], or the 25-34 year-old age category [H(2) = 0.004, p = 0.998]. However, there was a significant effect of long COVID status on memory scores for the 35-44 [H(2) = 18.539, p < 0.001], 45-54 [H(2) = 24.377, p < 0.001] and 55+ year-old categories [H(2) = 15.416, p < 0.001]. Dunn's pairwise comparisons (with Bonferroni correction for multiple tests) revealed that the memory scores in the diagnosed long COVID group were significantly reduced compared to the non-long COVID group in the 35-44, 45-54 and 55+ year-old categories (p < 0.001). The memory scores for the self-

reported long COVID group were significantly reduced compared to the non-long COVID group in the 35-44 age category ($p = 0.021$). In the 55+ age category, the memory scores for the diagnosed long COVID group were significantly smaller than for the self-reported long COVID group ($p = 0.034$).

Fig 3. The effect of age on objective memory scores in the non-long COVID, self-reported long COVID and diagnosed long COVID groups. The sample size is given in brackets for each data bar. Error bars represent the standard error of the mean. Asterisks indicate significant differences between groups for each age range category.   *$p < 0.05$, **$p < 0.001$.

Effect of the number of COVID-19 infections on working memory

Objective memory scores decreased significantly with increasing number of COVID-19 infections for the diagnosed long COVID group ($\tau b = -0.113$, $p = 0.043$). However, there was no significant correlation between the number of COVID-19 infections and memory scores for either the self-reported long COVID group ($\tau b = -0.040$, $p = 0.527$) or the non-long COVID group ($\tau b = 0.001$, $p = 0.970$). Next, we compared the impact of the number of COVID-19 infections on memory scores between the non-long COVID, self-reported long COVID and diagnosed long COVID groups. Since there were only 13 participants with four or more COVID-19 infections we combined these participants with those who had three COVID-19 infections (n=67) to give a total number of 80 participants in this category. Fig 4 shows the

effect of the number of COVID-19 infections on memory scores for all three long COVID

status groups for the four infection categories. A Kruskal-Wallis test was performed at each

of the four COVID-19 infection categories to determine differences between groups based

on long COVID status. When participants had one or two COVID-19 infections more than 12

weeks prior to completing the survey and memory quiz, there was a significant effect of

long COVID status on memory scores [one COVID-19 infection: H(2) = 23.046, p < 0.001; two

COVID-19 infections: H(2) = 25.352, p < 0.001]. There was no significant effect of long COVID

status on memory scores in participants with 3 or more infections [H(2) = 5.228, p = 0.073].

Dunn's pairwise comparisons (with Bonferroni correction for multiple tests) revealed that

the memory scores in the diagnosed long COVID group were significantly reduced compared

to the non-long COVID group in the one and two COVID-19 infection categories (p < 0.001).

The memory scores for the diagnosed long COVID group were significantly reduced

compared to the self-reported long COVID group in the two COVID-19 infection category (p

= 0.007). For the zero number of COVID-19 infection category, there was an insufficient

number of participants in the self-reported (n=5) and diagnosed long COVID (n=1) groups to

perform statistical analysis (these participants stated having long COVID despite an apparent

absence of COVID-19 infection).

Fig 4. The effect of COVID-19 infections on objective memory scores in the non-long

COVID, self-reported long COVID and diagnosed long COVID groups. The sample size is

given in brackets for each data bar. Error bars represent the standard error of the mean.

Asterisks indicate significant differences between groups for the number of COVID-19

infections. The dashed border on vertical bars (zero COVID-19 infections category) indicates

the small number of participants who stated having long COVID despite an apparent absence of COVID-19 infection. $*p < 0.01$, $**p < 0.001$

Effect of long COVID duration on working memory

As the number of months of having long COVID increased, the memory scores decreased significantly in the diagnosed long COVID group ($\tau b = -0.178$, $p < 0.001$) but not in the self-reported long COVID group ($\tau b = -0.021$, $p = 0.712$). Next, we compared the impact of long COVID-19 duration on memory scores between the self-reported long COVID and diagnosed long COVID groups. We combined participants into six-month long COVID-19 duration categories. Fig 5 shows the effect of long COVID duration on memory scores for the self-reported long COVID and diagnosed long COVID groups. A Mann-Whitney U test found no significant difference between the self-reported long COVID and diagnosed long COVID groups for any of the five long COVID duration categories (Bonferroni-corrected for five duration categories).

Fig 5. The effect of long COVID duration on objective memory scores in the self-reported and diagnosed long COVID groups. The sample size is given in brackets for each data bar. Note: Each bar represents a different sample of participants (between-subject design). Error bars represent the standard error of the mean.

COVID-19 variants and working memory

During the COVID-19 pandemic the SARS-CoV-2 virus evolved to produce different variants over time. Consequently, in the current study, those participants who developed long COVID early in the pandemic were likely to have been infected with a different variant than those who were infected later. It has been shown that participants infected when earlier variants were dominant have a higher probability of continuing long COVID symptoms, including cognitive impairment and poor memory [29]. Hence, our results indicating lower memory quiz scores with longer COVID duration (Figure 5) could have been influenced by the variant of SARS-CoV-2. When a participant's SARS-CoV-2 variant is unknown, previous studies have estimated the likely infection strain based on the dominant variant present at a given time period within a given country [29-31]. In a post-hoc analysis, we divided our long COVID participants from the United Kingdom only into groups according to the dominant variant at the time their long COVID symptoms began. This was achieved by subtracting the number of months since participants' long COVID symptoms started from the date they completed the online survey and memory quiz. Our variant groups were based on the time periods used previously for the United Kingdom [30, 31]: 1) wild-type, prior to December 2020; 2) alpha, from the beginning of December 2020 to the end of April 2021; 3) delta, from the beginning of May 2021 to the end of December 2021; 4) omicron, from the beginning of January 2022 onwards.

A Kendall's tau-b correlation showed a significant relationship between the SARS-CoV-2 variant that was dominant at the time at which long COVID symptoms started and the duration of long COVID symptoms for both the self-reported (τb = -0.740, $p < 0.001$) and

diagnosed (τb = -0.869, *p* < 0.001) long COVID groups. Fig 6 illustrates the objective memory

scores for each dominant SARS-CoV-2 variant in the self-reported and diagnosed long COVID

groups in United Kingdom participants only. For participants with diagnosed long COVID,

there was a significant difference in mean memory scores across the four SARS-CoV-2

variants (Kruskal-Wallis test, H(3) = 14.140, *p* = 0.003). For participants with diagnosed long

COVID, we found a significantly lower mean memory scores for the wild type (Dunn's

pairwise comparison corrected for multiple comparisons, *p* = 0.033) and alpha (*p* = 0.009)

variants compared to the omicron variant. There was no significant difference in mean

memory scores across the four SARS-CoV-2 variants for participants with self-reported long

COVID (Kruskal-Wallis test, H(3) = 1.236, *p* = 0.744).

Fig 6. The dominant SARS-CoV-2 variant and objective memory scores in the self-reported

and diagnosed long COVID groups in participants only from the United Kingdom. The long

COVID participants were divided into four groups (Wild type, Alpha, Delta and Omicron)

according to the dominant variant at the time when their long COVID symptoms began. The

sample size is given in brackets for each data bar. Error bars represent the standard error of

the mean. **p* < 0.05, ***p* < 0.01

Relationship between '*Brain fog*', '*Impact of long COVID symptoms on life*' and objective memory scores

There was a significant negative correlation between the subjective ratings for '*Brain fog*' and objective memory scores across all participants (τb = -0.185, *p* < 0.001). Fig 7a shows the mean subjective ratings for '*Brain fog*' versus objective memory scores for each of the non-long COVID, self-reported long COVID and diagnosed long COVID groups. A Kruskal-Wallis test revealed a significant relationship between long COVID status and '*Brain fog*' ratings [H(2) = 331.90, *p* < 0.001]. Dunn's pairwise comparisons (with Bonferroni correction for multiple tests) revealed that the subjective ratings of '*Brain fog*' in the self-reported and diagnosed long COVID groups were significantly more severe compared to the non-long COVID group (*p* < 0.001). Moreover, the subjective ratings of '*Brain fog*' in the diagnosed long COVID group were significantly more severe compared to the self-reported long COVID group (*p* = 0.011). There was a significant negative correlation between the subjective ratings for '*Impact of long COVID symptoms on life*' and objective memory scores across all participants with long COVID (τb = -0.169, *p* < 0.001). Fig 7b shows the mean subjective ratings of the '*Impact of long COVID symptoms on life*' versus objective memory scores for the self-reported and diagnosed long COVID groups. The subjective ratings of '*Impact of long COVID symptoms on life*' in the diagnosed long COVID group were significantly worse compared to the self-reported long COVID group (Mann-Whitney U test = 10372.00, Z = -9.61, *p* < 0.001, r=0.47). There was a significant correlation between the subjective ratings for '*Brain fog*' and '*Impact of long COVID symptoms on life*' in the long COVID (self-reported and diagnosed) group (τb = 0.405, *p* < 0.001).

Fig 7. Subjective ratings of a) 'Brain fog' and b) 'Impact of long COVID symptoms on life' versus memory scores. The vertical error bars represent the standard error of the mean for memory scores, and horizontal error bars indicate the standard error of the mean for 'Brain fog' or 'Impact of long COVID symptoms on life' ratings. Statistical significance values relate to ratings for 'Brain fog', and for 'Impact of long COVID symptoms on life' $*p < 0.01$, $**p < 0.001$.

Discussion

The main finding in our study was that long COVID negatively impacts upon working memory function. The most important factor affecting working memory scores was long COVID status (diagnosed long COVID, self-reported long COVID and non-long COVID) followed by age and number of COVID-19 infections. Using the same single rapid online working memory test as in our previous study [26], we found that the objective memory scores were significantly reduced in the long COVID groups (diagnosed with long COVID or self-reported having long COVID) compared to the non-long COVID group. These results are in line with findings from other studies showing memory deficits in individuals with continuing COVID-19 symptoms [9, 11, 19-22]. We found that the mean memory score for those diagnosed with long COVID (mean ± s.e.m. = 13.88 ± 0.15) was lower than for those in our previous study who had tested positive for COVID-19 but had not been hospitalized

(14.73 ± 0.06) and more comparable to people who had been hospitalized with COVID-19

(13.46 ± 0.41) [26]. Together, these findings indicate that working memory is particularly

impaired in those diagnosed with long COVID.

Our results provide evidence that working memory deficits may be linked to 'brain fog', one

of the most commonly reported symptoms in those with long COVID [6, 32-34]. We found

that for all participants, objective memory scores were negatively correlated with increasing

levels of brain fog, such that with those with lower memory scores reported the most

severe brain fog. The mean subjective ratings of brain fog were most severe for the

diagnosed long COVID group followed by the self-reported long COVID group and were

mildest in those who did not have long COVID. Our results align with those of a previous

study on long COVID, where memory impairments were self-reported in 66.2% of

participants with brain fog compared to 13.5% without brain fog [14]. The mean subjective

ratings of the '*Impact of long COVID symptoms on life*' were also most severe for the

diagnosed long COVID group with lower memory scores compared to the self-reported long

COVID group who had higher memory scores. Our results suggest that poorer working

memory may be one of the many symptoms that could contribute to the negative impact of

long COVID on daily life. This is further supported by our finding that subjective ratings for

'*Brain fog*' and '*Impact of long COVID symptoms on life*' were highly correlated. Previous

studies have also shown that aspects related to the lived experience of long COVID can

significantly reduce quality of life, including persistent physical symptoms, mental health

problems, ability to cope, reduced functional status and employment issues [7, 8, 19, 35-

37].

Previous studies on cognitive function in long COVID have typically classified participants into a long COVID group based on persistent symptomology rather than clinical diagnosis by a medical professional [21, 38-40]. One of the strengths of our study is that we included a group of long COVID participants who were clinically diagnosed. Medical professionals will diagnose patients with long COVID based on multiple sources of evidence, including case history, signs and symptoms, physical assessments/clinical tests, and may be supported by members of a multi-disciplinary team. In our study, we added a separate self-reported long COVID group to represent those who may have long COVID but were not clinically diagnosed. Furthermore, to guide participants who think they may have long COVID we provided in our survey an explicit definition from NHS England of long COVID namely, '*Signs and symptoms that develop during or after COVID-19 and continue for more than 12 weeks and are not explained by an alternative diagnosis.*'

We found that the objective memory scores decreased significantly with age in the non-long COVID control group. This finding aligns with the age-related effects observed in the control group of our previous study, which utilized the same working memory quiz [26]. It is also consistent with other research demonstrating that working memory declines as age increases [41, 42]. Importantly, the reduction of working memory scores with age was greater in the group diagnosed with long COVID compared to the non-long COVID group. Specifically, long COVID had a greater impact on memory scores for the older groups categories (35 years old and above) compared to the younger age categories (<35 years old) where there was no significant difference in memory score between the diagnosed and non-

long COVID groups. Our data indicate that working memory is impacted most by long COVID in adults aged 35 years and above. However, this interpretation must be contextualized by considering sample size, since the number of participants diagnosed with long COVID was lowest in the youngest age grouping (16-24 years) and higher in the older age groups. Nevertheless, our finding that memory problems in long COVID are more pronounced in older age groups aligns with the broader observation that older individuals experience greater memory impairments post-COVID-19 [15, 24-26, 43-45].

A study examining SARS-CoV-2 reinfection found that having two or more infections increased the risk of persistent symptoms in the post-acute phase (up to six months after infection) including neurological sequelae [46]. In our current investigation we found that objective memory scores decreased with increasing number of COVID-19 infections but only in the group diagnosed with long COVID. This finding supports the viewpoint that COVID-19 reinfections increase the severity of cognitive dysfunction in people with long COVID. There were a few participants who reported they had not had COVID-19 infection, but self-reported as having long COVID (n=5) or were diagnosed with long COVID (n=1). There are several possible reasons which could account for this, including the absence of COVID testing at the time of infection, being asymptomatic to the COVID-19 infection [47], a side effect of the COVID-19 vaccine [48], or simply an error when completing the questionnaire.

In our cross-sectional study, we found that memory scores decreased as the duration of long COVID symptoms persisted in those diagnosed with long COVID. This suggests that working memory function progressively deteriorates following COVID-19 infection in

individuals with long COVID. However, the effects of long COVID duration on working memory scores could be confounded with the effects of the different SARS-CoV-2 variants that infected participants. A previous investigation has indicated that variant type affects the severity of cognitive deficits, with earlier variants such as wild-type and alpha causing greater deficits than more recent delta and omicron variants [49]. In a post-hoc analysis, we divided participants from the United Kingdom with long COVID into four variant groups (wild-type, alpha, delta and omicron) based on dominant SARS-CoV-2 strains according to the time periods defined in previous studies [30, 31]. We found that the duration of long COVID symptoms was negatively correlated with the date order of dominant strains, 1) wild-type, 2) alpha, 3) delta and 4) omicron, in both self-reported and diagnosed long COVID groups. This suggests that individuals with longer duration symptoms were more likely to be infected with earlier variants of the virus. In the diagnosed long COVID group, those whose symptoms began when the wild-type and alpha variants were prevalent had lower memory scores compared to the later omicron variant. Although our results suggest that having long COVID for a protracted time is detrimental to working memory, this interpretation must be considered within the context of the prevalent SARS-CoV-2 strain. To further address the impact of long COVID duration on memory without the confound of SARS-CoV-2 variants, a future longitudinal approach is suggested where individuals' memory scores are assessed at multiple timepoints. In addition, individuals should be tested for variant type to extract the separate contribution of variant from duration.

We found that the group diagnosed with long COVID had lower scores compared to those who self-reported having long COVID indicating that working memory deficits were greater

in those diagnosed by a medical professional. Moreover, in the self-reported long COVID group, memory scores did not decrease significantly with age, number of COVID-19 infections, long COVID duration or probable SARS-CoV-2 variant. It is possible that participants were more likely to seek help and therefore be diagnosed with long COVID by medical professionals if they had more severe long COVID symptoms, including brain fog, memory problems and overall impact on their lives. It is possible that the self-reported group may have had higher memory scores than the diagnosed group because they had less severe symptoms and felt less of a need to consult a medical professional. Our study supports this hypothesis, as we found that the diagnosed long COVID group reported that they had more severe brain fog and that their symptoms had a greater impact on their life. A previous study has shown that more severe COVID-19 symptoms are linked to poorer cognitive function including memory deficits [21]. In addition, some participants in the self-reported long COVID group may have incorrectly diagnosed themselves based on their interpretation of the NHS single sentence definition of long COVID. Consequently, it is possible that these participants did not actually have long COVID and that their memory was less affected resulting in higher memory scores compared to the diagnosed group. The misdiagnosis of long COVID could affect the interpretation of results from prior studies that relied on self-reported data to classify the condition and may have led to an underestimation of the impact of long COVID on memory.

Several mechanisms have been proposed to explain the impact of long COVID on cognition including memory function. These include pathophysiological mechanisms that could impede brain function such as neuroinflammation, damage to neurons and astrocytes,

vascular alterations, neurotropic effects, dysautonomia, hypoxia, blood coagulation, and metabolic changes [15, 37, 50-54]. In addition, long COVID could affect psychological well-being, leading to conditions such as anxiety, depression, post-traumatic stress disorder, or sleep disturbances which in turn could contribute to memory deficits [35-37, 50, 55].

Since our main result demonstrates that long COVID can lead to problems with working memory, healthcare providers could incorporate a test which specifically assesses working memory when evaluating long COVID and to monitor changes over time following interventions. A number of rehabilitation approaches have been proposed to treat long COVID cognitive symptoms (with mixed effectiveness) including cognitive training, electromagnetic brain stimulation, psychosocial, pharmaceutical, natural supplements, psychoeducation/compensatory skills training, and physical rehabilitation such as hyperbaric oxygen therapy [53, 56, 57]. A personalized clinical treatment plan that considers working memory deficits may help address both the cognitive as well as physical symptoms of long COVID.

Conclusions

This study demonstrates that long COVID, particularly in clinically diagnosed cases, is associated with significant impairments in working memory. These memory impairments are exacerbated by factors such as age and the number of COVID-19 infections. Additionally, individuals reporting more severe brain fog and greater overall life impact from long COVID exhibited lower memory scores. These findings underscore the importance of implementing

targeted interventions for individuals with long COVID, with a focus on addressing cognitive deficits and improving quality of life.

Acknowledgments

We are grateful to Namitha Thottiyil for assistance in participant recruitment.

Author Contributions

Conceptualization: Aziz U. R. Asghar, Heidi A. Baseler, Abayomi Salawu, Murat Aksoy.

Data curation: Murat Aksoy, Aziz U. R. Asghar.

Formal analysis: Aziz U. R. Asghar, Hong Kiu Yuen, Heidi A. Baseler.

Investigation: Aziz U. R. Asghar, Hong Kiu Yuen, Murat Aksoy, Heidi A. Baseler

Methodology: Aziz U. R. Asghar, Hong Kiu Yuen, Murat Aksoy, Abayomi Salawu, Heidi A. Baseler.

Project administration, Aziz U. R. Asghar, Heidi A. Baseler.

Resources: Aziz U. R. Asghar, Murat Aksoy, Heidi A. Baseler.

Supervision: Aziz U. R. Asghar, Abayomi Salawu, Heidi A. Baseler.

Visualization: Aziz U. R. Asghar, Heidi A. Baseler.

Writing – original draft: Aziz U. R. Asghar, Heidi A. Baseler.

Writing – review & editing: Aziz U. R. Asghar, Hong Kiu Yuen, Murat Aksoy, Abayomi Salawu, Heidi A. Baseler.

References

1.   NICE. COVID-19 rapid guideline: managing the long-term effects of COVID-19: National Institute for Health and Care Excellence 2020 [updated 25 January 2024]. Available from: https://www.nice.org.uk/guidance/ng188/chapter/1-Identification.

2.   WHO. Post COVID-19 condition (Long COVID): World Health Organization; 2022 [updated 7 December 2022]. Available from: https://www.who.int/europe/news-room/fact-sheets/item/post-covid-19-condition.

3.   O'Mahoney LL, Routen A, Gillies C, Ekezie W, Welford A, Zhang A, et al. The prevalence and long-term health effects of Long Covid among hospitalised and non-hospitalised populations: A systematic review and meta-analysis. EClinicalMedicine. 2023;55:101762. Epub 2022/12/08. doi: 10.1016/j.eclinm.2022.101762. PubMed PMID: 36474804.

4.   ONS. Coronavirus (COVID-19) latest insights: Infections In: Statistics OfN, editor.: Government of the United Kingdom 2023.

5.   Cheetham NJ, Penfold R, Giunchiglia V, Bowyer V, Sudre CH, Canas LS, et al. The effects of COVID-19 on cognitive performance in a community-based cohort: a COVID symptom study biobank prospective cohort study. EClinicalMedicine. 2023;62:102086. Epub 2023/09/01. doi: 10.1016/j.eclinm.2023.102086. PubMed PMID: 37654669.

6.   Millet C, Narvaneni S, Roman S, Horani G, Chaudhry S, Michael P, et al. Symptoms persist in patients two years after COVID-19 infection: a prospective follow-up study. Clin Microbiol Infect. 2022;28(11):1505-7. Epub 2022/06/21. doi: 10.1016/j.cmi.2022.06.008. PubMed PMID: 35724867.

7.   Kim Y, Bae S, Chang HH, Kim SW. Long COVID prevalence and impact on quality of life 2 years after acute COVID-19. Sci Rep. 2023;13(1):11207. Epub 2023/07/12. doi: 10.1038/s41598-023-36995-4. PubMed PMID: 37433819.

8.   Ziauddeen N, Gurdasani D, O'Hara ME, Hastie C, Roderick P, Yao G, et al. Characteristics and impact of Long Covid: Findings from an online survey. PLoS One. 2022;17(3):e0264331. Epub 2022/03/09. doi: 10.1371/journal.pone.0264331. PubMed PMID: 35259179.

9.   Bertuccelli M, Ciringione L, Rubega M, Bisiacchi P, Masiero S, Del Felice A. Cognitive impairment in people with previous COVID-19 infection: A scoping review. Cortex. 2022;154:212-30. Epub 2022/07/06. doi: 10.1016/j.cortex.2022.06.002. PubMed PMID: 35780756.

10.   Zawilska JB, Kuczynska K. Psychiatric and neurological complications of long COVID. J Psychiatr Res. 2022;156:349-60. Epub 2022/11/04. doi: 10.1016/j.jpsychires.2022.10.045. PubMed PMID: 36326545.

11.   Zhao S, Toniolo S, Hampshire A, Husain M. Effects of COVID-19 on cognition and brain health. Trends Cogn Sci. 2023;27(11):1053-67. Epub 2023/09/02. doi: 10.1016/j.tics.2023.08.008. PubMed PMID: 37657964.

12.   Nordvig AS, Rajan M, Lau JD, Kingery JR, Mahmud M, Chiang GC, et al. Brain fog in long COVID limits function and health status, independently of hospital severity and preexisting conditions. Front Neurol. 2023;14:1150096. Epub 2023/05/30. doi: 10.3389/fneur.2023.1150096. PubMed PMID: 37251229.

13.   Lam GY, Damant RW, Ferrara G, Lim RK, Stickland MK, Ogando NS, et al. Characterizing long-COVID brain fog: a retrospective cohort study. J Neurol. 2023;270(10):4640-6. Epub 2023/08/09. doi: 10.1007/s00415-023-11913-w. PubMed PMID: 37555926.

14.   Jennings G, Monaghan A, Xue F, Duggan E, Romero-Ortuno R. Comprehensive Clinical Characterisation of Brain Fog in Adults Reporting Long COVID Symptoms. J Clin Med. 2022;11(12). Epub 2022/06/25. doi: 10.3390/jcm11123440. PubMed PMID: 35743516.

15.   Nalbandian A, Sehgal K, Gupta A, Madhavan MV, McGroder C, Stevens JS, et al. Post-acute COVID-19 syndrome. Nat Med. 2021;27(4):601-15. Epub 2021/03/24. doi: 10.1038/s41591-021-01283-z. PubMed PMID: 33753937.

16.   van der Feltz-Cornelis C, Turk F, Sweetman J, Khunti K, Gabbay M, Shepherd J, et al. Prevalence of mental health conditions and brain fog in people with long COVID: A systematic review and meta-analysis. Gen Hosp Psychiatry. 2024;88:10-22. Epub 2024/03/07. doi: 10.1016/j.genhosppsych.2024.02.009. PubMed PMID: 38447388.

17.   Baddeley AD, Hitch GJ, Allen RJ. From short-term store to multicomponent working memory: The role of the modal model. Mem Cognit. 2019;47(4):575-88. Epub 2018/11/28. doi: 10.3758/s13421-018-0878-5. PubMed PMID: 30478520.

18.   Repovs G, Baddeley A. The multi-component model of working memory: explorations in experimental cognitive psychology. Neuroscience. 2006;139(1):5-21. Epub 2006/03/07. doi: 10.1016/j.neuroscience.2005.12.061. PubMed PMID: 16517088.

19.   Graham EL, Clark JR, Orban ZS, Lim PH, Szymanski AL, Taylor C, et al. Persistent neurologic symptoms and cognitive dysfunction in non-hospitalized Covid-19 "long haulers". Ann Clin Transl Neurol. 2021;8(5):1073-85. Epub 2021/03/24. doi: 10.1002/acn3.51350. PubMed PMID: 33755344.

20.   Miskowiak KW, Johnsen S, Sattler SM, Nielsen S, Kunalan K, Rungby J, et al. Cognitive impairments four months after COVID-19 hospital discharge: Pattern, severity and association with illness variables. Eur Neuropsychopharmacol. 2021;46:39-48. Epub 2021/04/07. doi: 10.1016/j.euroneuro.2021.03.019. PubMed PMID: 33823427.

21.   Guo P, Benito Ballesteros A, Yeung SP, Liu R, Saha A, Curtis L, et al. COVCOG 2: Cognitive and Memory Deficits in Long COVID: A Second Publication From the COVID and Cognition Study. Front Aging Neurosci. 2022;14:804937. Epub 2022/04/05. doi: 10.3389/fnagi.2022.804937. PubMed PMID: 35370620.

22.   Miskowiak KW, Pedersen JK, Gunnarsson DV, Roikjer TK, Podlekareva D, Hansen H, et al. Cognitive impairments among patients in a long-COVID clinic: Prevalence, pattern and relation to illness severity, work function and quality of life. J Affect Disord. 2023;324:162-9. Epub 2023/01/01. doi: 10.1016/j.jad.2022.12.122. PubMed PMID: 36586593.

23.   Arbula S, Pisanu E, Bellavita G, Menichelli A, Lunardelli A, Furlanis G, et al. Insights into attention and memory difficulties in post-COVID syndrome using standardized neuropsychological tests and experimental cognitive tasks. Sci Rep. 2024;14(1):4405. Epub 2024/02/23. doi: 10.1038/s41598-024-54613-9. PubMed PMID: 38388708.

24.   Stavem K, Einvik G, Tholin B, Ghanima W, Hessen E, Lundqvist C. Cognitive function in non-hospitalized patients 8-13 months after acute COVID-19 infection: A cohort study in Norway. PLoS One. 2022;17(8):e0273352. Epub 2022/08/23. doi: 10.1371/journal.pone.0273352. PubMed PMID: 35994448.

25.   Cui R, Gao B, Ge R, Li M, Li M, Lu X, et al. The effects of COVID-19 infection on working memory: a systematic review. Curr Med Res Opin. 2024;40(2):217-27. Epub 2023/11/27. doi: 10.1080/03007995.2023.2286312. PubMed PMID: 38008952.

26.   Baseler HA, Aksoy M, Salawu A, Green A, Asghar AUR. The negative impact of COVID-19 on working memory revealed using a rapid online quiz. PLoS One. 2022;17(11):e0269353. Epub 2022/11/15. doi: 10.1371/journal.pone.0269353. PubMed PMID: 36374838.

27.   World Medical A. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. JAMA. 2013;310(20):2191-4. Epub 2013/10/22. doi: 10.1001/jama.2013.281053. PubMed PMID: 24141714.

28.   NHS. Post-COVID syndrome (long COVID): National Health Service England; 2021. Available from: https://www.england.nhs.uk/coronavirus/post-covid-syndrome-long-covid/.

29.   Diexer S, Klee B, Gottschick C, Xu C, Broda A, Purschke O, et al. Association between virus variants, vaccination, previous infections, and post-COVID-19 risk. Int J Infect Dis. 2023;136:14-21. Epub 2023/08/28. doi: 10.1016/j.ijid.2023.08.019. PubMed PMID: 37634619.

30.   Elliott P, Bodinier B, Eales O, Wang H, Haw D, Elliott J, et al. Rapid increase in Omicron infections in England during December 2021: REACT-1 study. Science. 2022;375(6587):1406-11. Epub 2022/02/09. doi: 10.1126/science.abn8347. PubMed PMID: 35133177.

31.   Atchison CJ, Davies B, Cooper E, Lound A, Whitaker M, Hampshire A, et al. Long-term health impacts of COVID-19 among 242,712 adults in England. Nat Commun. 2023;14(1):6588. Epub 2023/10/25. doi: 10.1038/s41467-023-41879-2. PubMed PMID: 37875536.

32.   Premraj L, Kannapadi NV, Briggs J, Seal SM, Battaglini D, Fanning J, et al. Mid and long-term neurological and neuropsychiatric manifestations of post-COVID-19 syndrome: A meta-analysis. J Neurol Sci. 2022;434:120162. Epub 2022/02/06. doi: 10.1016/j.jns.2022.120162. PubMed PMID: 35121209.

33.   Aghajani Mir M. Brain Fog: a Narrative Review of the Most Common Mysterious Cognitive Disorder in COVID-19. Mol Neurobiol. 2023. Epub 2023/10/24. doi: 10.1007/s12035-023-03715-y. PubMed PMID: 37874482.

34.   Kalak G, Jarjou'i A, Bohadana A, Wild P, Rokach A, Amiad N, et al. Prevalence and Persistence of Symptoms in Adult COVID-19 Survivors 3 and 18 Months after Discharge from Hospital or Corona Hotels. J Clin Med. 2022;11(24). Epub 2022/12/24. doi: 10.3390/jcm11247413. PubMed PMID: 36556030.

35.   Aiyegbusi OL, Hughes SE, Turner G, Rivera SC, McMullan C, Chandan JS, et al. Symptoms, complications and management of long COVID: a review. J R Soc Med. 2021;114(9):428-42. Epub 2021/07/16. doi: 10.1177/01410768211032850. PubMed PMID: 34265229.

36.   Kennelly CE, Nguyen ATP, Sheikhan NY, Strudwick G, Ski CF, Thompson DR, et al. The lived experience of long COVID: A qualitative study of mental health, quality of life, and coping. PLoS One. 2023;18(10):e0292630. Epub 2023/10/13. doi: 10.1371/journal.pone.0292630. PubMed PMID: 37831706.

37.   Natarajan A, Shetty A, Delanerolle G, Zeng Y, Zhang Y, Raymont V, et al. A systematic review and meta-analysis of long COVID symptoms. Syst Rev. 2023;12(1):88. Epub 2023/05/28. doi: 10.1186/s13643-023-02250-0. PubMed PMID: 37245047.

38.   Bungenberg J, Humkamp K, Hohenfeld C, Rust MI, Ermis U, Dreher M, et al. Long COVID-19: Objectifying most self-reported neurological symptoms. Ann Clin Transl Neurol. 2022;9(2):141-54. Epub 2022/01/22. doi: 10.1002/acn3.51496. PubMed PMID: 35060361.

39.   Lauria A, Carfi A, Benvenuto F, Bramato G, Ciciarello F, Rocchi S, et al. Neuropsychological measures of post-COVID-19 cognitive status. Front Psychol. 2023;14:1136667. Epub 2023/07/26. doi: 10.3389/fpsyg.2023.1136667. PubMed PMID: 37492442.

40.   Thronicke A, Hinse M, Weinert S, Jakubowski A, Grieb G, Matthes H. Factors Associated with Self-Reported Post/Long-COVID-A Real-World Data Study. Int J Environ Res Public Health. 2022;19(23). Epub 2022/12/12. doi: 10.3390/ijerph192316124. PubMed PMID: 36498197.

41.   Brockmole JR, Logie RH. Age-related change in visual working memory: a study of 55,753 participants aged 8-75. Front Psychol. 2013;4:12. Epub 2013/02/02. doi: 10.3389/fpsyg.2013.00012. PubMed PMID: 23372556.

42.   Park DC, Lautenschlager G, Hedden T, Davidson NS, Smith AD, Smith PK. Models of visuospatial and verbal memory across the adult life span. Psychol Aging. 2002;17(2):299-320. Epub 2002/06/14. PubMed PMID: 12061414.

43.   Blomberg B, Mohn KG, Brokstad KA, Zhou F, Linchausen DW, Hansen BA, et al. Long COVID in a prospective cohort of home-isolated patients. Nat Med. 2021;27(9):1607-13. Epub 2021/06/25. doi: 10.1038/s41591-021-01433-3. PubMed PMID: 34163090.

44.   Corbett A, Williams G, Creese B, Hampshire A, Hayman V, Palmer A, et al. Cognitive decline in older adults in the UK during and after the COVID-19 pandemic: a longitudinal analysis of PROTECT study data. Lancet Healthy Longev. 2023;4(11):e591-e9. Epub 2023/11/05. doi: 10.1016/S2666-7568(23)00187-3. PubMed PMID: 37924840.

45.   Finamore P, Arena E, Lupoi D, Savito L, Di Nunzio F, Furbatto M, et al. Long COVID Syndrome: A Narrative Review on Burden of Age and Vaccination. J Clin Med. 2024;13(16). Epub 2024/08/31. doi: 10.3390/jcm13164756. PubMed PMID: 39200898.

46.   Bowe B, Xie Y, Al-Aly Z. Acute and postacute sequelae associated with SARS-CoV-2 reinfection. Nat Med. 2022;28(11):2398-405. Epub 2022/11/11. doi: 10.1038/s41591-022-02051-3. PubMed PMID: 36357676.

47.   Gao Z, Xu Y, Sun C, Wang X, Guo Y, Qiu S, et al. A systematic review of asymptomatic infections with COVID-19. J Microbiol Immunol Infect. 2021;54(1):12-6. Epub 2020/05/20. doi: 10.1016/j.jmii.2020.05.001. PubMed PMID: 32425996.

48.   Finterer J, Scorza FA. A retrospective analysis of clinically confirmed long post-COVID vaccination syndrome. J Clin Transl Res. 2022;8(6):506-8. Epub 2022/12/02. PubMed PMID: 36452006.

49.   Hampshire A, Azor A, Atchison C, Trender W, Hellyer PJ, Giunchiglia V, et al. Cognition and Memory after Covid-19 in a Large Community Sample. N Engl J Med. 2024;390(9):806-18. Epub 2024/02/28. doi: 10.1056/NEJMoa2311330. PubMed PMID: 38416429.

50.   Davis HE, McCorkell L, Vogel JM, Topol EJ. Long COVID: major findings, mechanisms and recommendations. Nat Rev Microbiol. 2023;21(3):133-46. Epub 2023/01/14. doi: 10.1038/s41579-022-00846-2. PubMed PMID: 36639608.

51.   Bauer L, Laksono BM, de Vrij FMS, Kushner SA, Harschnitz O, van Riel D. The neuroinvasiveness, neurotropism, and neurovirulence of SARS-CoV-2. Trends Neurosci. 2022;45(5):358-68. Epub 2022/03/14. doi: 10.1016/j.tins.2022.02.006. PubMed PMID: 35279295.

52.   Turner S, Khan MA, Putrino D, Woodcock A, Kell DB, Pretorius E. Long COVID: pathophysiological factors and abnormalities of coagulation. Trends Endocrinol Metab. 2023;34(6):321-44. Epub 2023/04/21. doi: 10.1016/j.tem.2023.03.002. PubMed PMID: 37080828.

53.   Moller M, Borg K, Janson C, Lerm M, Normark J, Niward K. Cognitive dysfunction in post-COVID-19 condition: Mechanisms, management, and rehabilitation. J Intern Med. 2023;294(5):563-81. Epub 2023/09/28. doi: 10.1111/joim.13720. PubMed PMID: 37766515.

54.   Plantone D, Stufano A, Righi D, Locci S, Iavicoli I, Lovreglio P, et al. Neurofilament light chain and glial fibrillary acid protein levels are elevated in post-mild COVID-19 or asymptomatic SARS-CoV-2 cases. Sci Rep. 2024;14(1):6429. Epub 2024/03/19. doi: 10.1038/s41598-024-57093-z. PubMed PMID: 38499607.

55.   Tedjasukmana R, Budikayanti A, Islamiyah WR, Witjaksono A, Hakim M. Sleep disturbance in post COVID-19 conditions: Prevalence and quality of life. Front Neurol. 2022;13:1095606. Epub 2023/01/27. doi: 10.3389/fneur.2022.1095606. PubMed PMID: 36698905.

56.   Hawke LD, Nguyen ATP, Wang W, Brown EE, Xu D, Deuville S, et al. Systematic review of interventions for mental health, cognition and psychological well-being in long COVID. BMJ Ment Health. 2024;27(1). Epub 2024/10/10. doi: 10.1136/bmjment-2024-301133. PubMed PMID: 39384321.

57.   Gorenshtein A, Liba T, Leibovitch L, Stern S, Stern Y. Intervention modalities for brain fog caused by long-COVID: systematic review of the literature. Neurol Sci. 2024;45(7):2951-68. Epub 2024/05/02. doi: 10.1007/s10072-024-07566-w. PubMed PMID: 38695969.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7