1

Testosterone replacement therapy ameliorates spatial cognitive function in age-related hypogonadism (LOH) patients: analysis with a virtual three-dimensional maze

Suguru Kawato[1,2,3,4†], Hideo Mukai[5], Yusuke Kai[5], Masaki Kimura[6], Mari Ogiue-Ikeda[1,2,4], Mika Soma[1,2,4,7], Minoru Saito[4] and Shigeo Horie[1]

[1]Department of Urology, Graduate School of Medicine, Juntendo Univ., Tokyo, Japan,
[2]Department of Cognitive Neuroscience, Faculty of Pharma-Science, Teikyo Univ., Tokyo, Japan,
[3]Department of Biophysics and Life Sciences, Graduate School of Arts and Sciences, The University of Tokyo, Tokyo, Japan,
[4]Department of Biosciences, College of Humanities and Sciences, Nihon University, Tokyo, Japan,
[5]Department of Computer Science, Graduate School of Science and Technology, Meiji University, Kanagawa, Japan,
[6]Department of Urology, School of Medicine, Teikyo Univ., Tokyo, Japan,
[7] Department of Anatomy and Neurobiology, Nippon Medical School, Tokyo, Japan.

Correspondence: Suguru Kawato, e-mail : skkawato@gmail.com

Keywords: testosterone, spatial navigation, late onset hypogonadism, testosterone replacement therapy, hippocampus

Abstract

When comparing the behavioral levels of men and women using a maze navigation game created on a PC to investigate spatial cognitive function, many reports indicated that men are significantly faster than women at finding the exit of the maze.

Analysis with functional MRI neuroimaging indicates that neural activation at multiple sites were observed during a maze navigation task. Significantly different regions were identified in men and women, with men showing significantly increased activity in the hippocampus. In contrast, women showed increased activity in the Brodmann area of the prefrontal cortex. When differences in the activated regions were taken between men and women, the hippocampal activity showed the largest difference.

Since the hippocampal CA1 regions are known to be essentially responsible to spatial navigation ability, the reported difference in spatial ability between men and women could be due to higher testosterone (T) levels in mem than in women, because the application of T can increase hippocampal CA1 neural synapses as observed in rats, for example.

In the present study, we developed the original 3D maze navigation test on a PC to accurately analyze spatial cognitive function, and this maze was used to analyze differences in spatial cognitive function between young male and female volunteers. The results showed that males took a distinctly shorter time and path length to escape the maze than females. We applied this maze test to analyze the possible improvement effects of spatial cognitive function by T-replacement therapy (TRT) on male patients with late onset hypogonadism (LOH). LOH patients have significantly low levels of T as well as high Aging males' symptoms scale (AMS) scores when compared with healthy men. Upon TRT for 6 weeks, we revealed that the spatial cognitive ability of LOH patients was significantly improved along the increase of T levels and improvement of AMS scores.

The observed improvement of spatial cognitive ability of LOH patients could be interpreted as effects of elevation of T levels in the hippocampal CA1 region where androgen receptor (AR) is localized.

Introduction

Continuous visuospatial navigation in familiar and unfamiliar environments is a requirement of daily life.

Visuospatial navigation is one of the few cognitive functions for which a reliable gender-specific performance difference is well known (Astur et al 1998). For example, men were significantly faster than women at finding the way out of maze.

Navigation through mazes can be combined with functional neuroimaging for men and women. From the fMRI analysis, gender differences in brain neural activation was computed with maze navigation tasks for male and female subjects (Grön et al

2000). Although there was great overlap between men and women in brain regions that were significantly activated, there were brain regions that were activated exclusively by the male and female groups. From the comparison of the neural activity, men showed significantly increased activity in the hippocampus proper, however, women showed increased activity of the prefrontal cortex, in contrast with men. (Grön et al 2000)

In the hippocampus, the glutamatergic neurons in the CA1 region are responsible for spatial cognition (Tonegawa et al 1996). This is evidenced by the fact that spatial cognition is selectively inhibited in mice in which N-methyl-D-aspartate (NMDA) receptors of glutamatergic neurons in the CA1 region of the hippocampus are specifically knockout (Tsien et al 1996) (McHugh et al 1996).

Previous studies of sex hormones in rats have shown that the level of testosterone (T) in the hippocampus is much higher in males (approximately 17 nM) than in females (approxi. 0.05 nM) (Hojo et al 2009) (Ooishi et al 2012) (Kato et al 2013) . T activates neurons upon binding to androgen receptor (AR).   AR is significantly expressed in glutamatergic neurons of the hippocampal CA1 region (Tabori et al 2005) (Hatanaka et al 2015). In addition, T level is much higher in male hippocampus than female hippocampus (Kato et al 2013). Circulating T levels are qualitatively proportional to hippocampal T levels (Kato et al 2013) (Hojo et al 2014) (Hojo & Kawato 2018). In view of these situations, we hypothesize that "higher T levels in males and lower T levels in females may have a significant effect on the difference in spatial cognitive function between men and women".

In Section 1, based on this hypothesis, we created a virtual maze escape game on a PC in order to find the spatial cognitive ability that may depend on T levels.

In Section 2, we applied this maze whether this maze is able to clearly determine differences in spatial cognitive ability between young men and women. We obtained the results that young men were significantly faster than young women in escaping from the maze.

In Section 3, we applied this maze navigation test to male patients with late onset hypogonadism (LOH) in order to examine whether T-replacement therapy (TRT) is effective to improve spatial cognitive ability by increase of T levels. It should be noted that because LOH patients have low T levels, TRT can elevate T levels of LOH patients, and thereby can improve Aging males' symptoms scale (AMS) scores (Heinemann et al 2003) (Lunenfeld et al 2013) (Ide et al 2023).   However, there had not been any challenges to investigate how TRT could improve spatial cognitive ability of LOH patients.

Section 1: Creation of 3D maze for spatial maze navigation test

1.1   Technical specification of maze

A 3D virtual maze program was created using the 3D modeling software Unity 5.6.1
(Unity Technologies). C# was used as the programming language.
The PC environment for the maze program and operation is as follows
OS: Windows 10 or 11,   64bit
CPU: Intel Core i7-6700HQ 2.60GHz
RAM: 16.0GB
GPU: Intel HD Graphics 530
Monitor: 15.6 in wide
The size of the maze is 6 squares (vertical) x 6 squares (horizontal) for a total of 36
squares (Figure 1.1, 1.2).
Fifteen sets of maze (Maze Number 1 – 15) are prepared. One of the maze can be
selected by choosing the group number (1, 2 or 3) on the title screen (see below), and
then automatically one Maze will be randomly selected. Note that one group consists of
5 mazes depending on goal positions.
128

129
130 **Figure 1.1**   Aerial view of the maze. The maze is 6x6 squares in size and can be
131 presented in a choice of 15 different ways.
132
133 **Start and goal points**
134
The 15 mazes are divided into three groups: one with a goal in square (6, 6), one
with a goal in square (1, 6), and one with a goal in square (6, 1). Subject can choose
which maze with a goal position you want to generate by entering the maze number on
the title screen (see below).
The start point is randomly selected from the four corners of the maze except for the
goal. For example, in Figure 1.2, the goal is located at square (6, 6) (red square), so the
starting point is chosen from the squares (1, 1), (1, 6) and (6, 1) (blue square).

Figure 1.2 How the start (blue) and goal (red) points are selected. If the start point is
in a red square, the goal point is randomly selected from the blue squares. Note that
subjects cannot see the goal that is hidden from the start by a wall.

Operation of maze task

Start , top title screen

When the maze navigation program is started, the top title screen appears. On the
title screen, the subject enters his/her name and maze number in order to select the type
of maze. Two types of mazes are available: maze without landmarks or maze with three
landmarks.

Before starting the maze test at the first time, "Demo Video" should be selected on
the title screen in order to get an overview of the present navigation maze task.

When 'walking' through the maze, subjects have a first-person view. During the
maze test, the arrow keys on the keyboard can be used to move forward with the up
arrow key, backward with the down arrow key, and turn right and left with the right and
left arrow keys, respectively. Turning was possible within the full range of 360 degree.
[↑] forward , [↓] backward , [→] right turn, [←] left turn.

Figure 1.3 Top title screen. Press Return to next Selection screen.

181   *Selection of the time limit screen*
182
183

196
197   **Figure 1.4** Time limit selection screen. There are two time limits (300 and 600
198   seconds) that can be chosen. Normally 300 sec is chosen.
199
200   Once the time limit is selected, the maze test begins. When the time limit is
201   exceeded, the word "End" is displayed.

202
203   **Figure 1.5** Typical view inside the maze. During the control condition, the screen is
204   frozen at the last position of the active condition.
205
206   During the maze test, a subject proceeds in the first-person perspective and searches
207   for the correct path.
208
209
210   *Goal screen*
211   When a subject reaches the goal (the red wall) (Figure 1.6), a subject should presses
212   the return key to finish the task. Then the time taken to reach the goal and the distance
213   traveled during the maze task are recorded. At the same time, the screen will be back to
214   the title screen (see Fig 1.3).
215

216
217
218 **Figure 1.6** The "Goal" is displayed on the goal wall. Pressing the Enter key to finish.
219
220   Landmarks in each maze are placed at the simple turned in the load to give a hint of
221 location. Landmarks are not placed at intersection of the convey (that is, decision
222 points).
223

236
237 **Figure 1.7** Example of landmarks in the maze. Three types of landmarks(circle, star,
238 and triangle) are used. Only three walls have landmarks and each has different types of
239 landmarks.
240
241
242 **1.2   Assessment of Spatial Cognitive Ability**
243
Escape time
As a measure of spatial cognitive ability, the time taken to solve a maze test is
measured. This is called escape time.
The escape time is the number of seconds from the start of the maze test
(immediately after the time limit is selected) to the end of the maze test (when the
subject reaches the goal or the time limit is reached). The escape time is output to the
total score.csv file in the result folder directly under the C drive after the maze test is
completed.
252

Path Distance
As a measure of spatial cognitive ability other than escape time, the distance traveled
during the maze test is recorded. This is called the path distance.
The path distance, as well as the escape time, is output to the total score.csv file
located in the results folder directly under the C drive after the maze test is completed.
The Euclidean distance for each second during the maze test is taken, and the sum of
these Euclidean distances from start to goal is defined as the path distance (transit
distance).
261
262
263
264 **Section 2: Analysis of differences in spatial cognitive ability between young men**
265 **and women using a virtual maze navigation task**
266
267 **Materials and Methods**
268
Subjects
The subjects were 40 healthy students (20 men and 20 women) affiliated with Meiji
University. The ages of the men and women subjects were 21.6 ± 1.5 and 20.6 ± 1.2 years,
respectively. The experiment was notified to and approved by the Ethics Committee on
Safety and Human Subjects for Experimental Research on Genetic Recombination, School
of Science and Engineering, Meiji University (Approval No.: RIKEN 14-520). Subjects
were informed about the research in person in advance and their consent was obtained by
signing a research participation consent form.
277
278   **Experimental Procedures**
279   The experiment will be conducted according to the following procedures.
280
281 1. After explaining the outline of the experiment, show a demonstration video of the maze
282 test.
283 2. Ask the subjects to perform the maze escape task without landmarks.
284 3. Ask the subjects to perform the maze escape task with three landmarks.
285 4. Repeat steps 2 and 3 four times with an interval of one week.
286
287   When performing the maze test, maze number 1 should be entered in the first time,
288 maze number 2 in the second time, maze number 3 in the third time, and maze number 4 in
289 the fourth time, to avoid mazes with identical goal positions.
290
291 **Results**
292
293 Weekly comparison of maze navigation task, using maze without/with three landmarks.
294
In the maze navigation task, young men significantly outscored young women in the

maze without landmarks, however, this difference became smaller by the presence of
landmarks (Figure 2.1).
The escape time of young men are shorter than that of young women in both mazes
without and with landmarks. However, the presence of landmarks made the difference
smaller.
In terms of path distance, men showed significantly shorter path distance than
women in the maze without landmarks, however, by the presence of landmarks women
showed the same path distance as men (Figure 2.1). Differences between men and
women are large in the absence of landmarks. Significant decreasing trends in escape
time and path distance are observed only for women in the maze without landmarks.
Table 1.1 and Table 1.2 represent statistical analyses that showed a significance (p <
0.05) by comparison between gender (men or women), the presence and absence of
landmarks and weeks.
309

312 **Figure 2.1**   Weekly scores of escape time (Left) and path distance (Right) for young
313 men and women in maze tests. Vertical axis is escape time (second) and path distance
314 (arbitrary unit), respectively.
315 (●)Young men without landmarks, (○) Young Men with landmarks, (▲) Young
316 women without landmarks, (△) Young women with landmarks.
317

	Escape Time			Path Distance		
	F	p	η^2	F	p	η^2
Gender vs Presence/Absence of Landmarks	8.31	0.0064**	0.0176	6.59	0.0143*	0.0147
Gender vs Weeks	4.50	0.0051*	0.0344	1.41	0.244	0.0118
Gender vs Presence/Absence of Landmarks vs Week	0.789	0.503	0.00750	0.275	0.843	0.00270

* $p < 0.05$ ** $p < 0.01$

320 **Table 1.1**   A two-way analysis of variance test revealed an interaction between gender
321 (men or women) and presence/absence of landmarks for escape time and path distance,
322 and an interaction between gender and week for escape time.
323

	Escape Time			Path Distance		
	F	p	η^2	F	p	η^2
Effect of Gender in Maze without Landmarks	21.5	$< 1.00 \times 10^{-4}$ ***	0.160	3.87	0.0564	0.0331
Effect of Gender in Maze with Landmarks	7.18	0.0108 *	0.0598	0.252	0.618	0.00170
Effect of Presence/Absence of Landmarks in Male	0.0363	0.851	2.00×10^{-4}	0.0863	0.772	4.00×10^{-4}
Effect of Presence/Absence of Landmarks in Female	11.0	0.00370 **	0.0417	9.07	0.00720 *	0.0391
Effect of Gender in 0w	19.3	1.00×10^{-4} ***	0.224			
Effect of Gender in 1w	11.5	0.00160 **	0.131			
Effect of Gender in 2w	4.23	0.0467 *	0.0629			
Effect of Gender in 3w	1.85	0.182	0.0287			
Effect of Weeks in Male	1.99	0.126	0.0268			
Effect of Weeks in Female	4.86	0.00440 **	0.0741			

$* \ p < 0.05 \quad ** \ p < 0.01 \quad *** \ p < 0.001$

Table 1.2   Simple main effect tests are conducted on escape time and path distance. In the analysis of gender and presence/absence of landmarks, effects are shown of gender in maze without landmarks, gender in maze with landmarks, and presence/absence of landmarks in women in escape time.

Weekly comparison in order to check learning effects on the maze test

During repeating the maze test, possible learning effects (with familiarity factors) might improve the skill to escape the maze. If the subjects become accustomed to the maze, their scores could be improved. We observed the decrease in the escape time and the path distance by repeating women's maze tests without landmarks at different weeks (0th ,1st , 2nd, 3rd week trial) (see Figure 2.1). However, in women's maze tests with landmarks, these improvement effects by repetitive training were not observed.
On the other hand for men, the presence and absence of landmarks did not induce any difference in both escape time and path distance, suggesting that men use geometric cues (Grön et al 2000).

Figure 2.2 shows the examples of the recorded path taken in the maze test at first time (0th week) men follow the more efficient path than women toward the goal in the maze without landmarks, but in the presence of landmarks this gender difference became smaller.

Figure 2.2   From left to right, young men without landmarks , young men with landmarks, young women without landmarks, young women with landmarks. Start (Arrow) and Goal (Red) are indicated.

354
355
356 **Section 3   Testosterone-Replacement Therapy (TRT) improved spatial cognition**
357 **ability of LOH patients**
358

In Section 2, the visuospatial navigation test successfully demonstrated differences
in the spatial cognition ability between young men (with high T level) and young
women (with low T level). The hippocampus is probably responsible to this spatial
cognition ability from important previous reports. The spatial cognition ability of rats
mainly depends on CA1 hippocampal neurons in which androgen receptor AR is
localized (Tsien et al 1996) (McHugh et al 1996) (Tonegawa et al 1996) (Huerta et al
2000) (Tabori et al 2005) (Hatanaka et al 2015). Therefore, the observed difference in
spatial cognition ability may significantly depend on the much higher hippocampal T
levels in male than female as observed in rats (Hojo et al 2009) (Kato et al 2013). This
hypothesis is dependent on the observation that nearly 20-fold T levels (in the
hippocampus) and 100-fold T levels (in circulation) were observed in male rats than
female rats (Hojo et al 2009) (Ooishi et al 2012) (Kato et al 2013).
It is shown also that T application to hippocampal CA1 glutamatergic neurons
increased the number of synapses where memory is stored (Hatanaka et al 2015)
(Murakami et al 2018) (Hojo & Kawato 2018). It may be therefore possible to find
differences in maze navigation ability between individual men with high T and low T
levels.
We applied spatial maze navigation task (with three landmarks) to LOH patients
(with low T levels) in order to analyze a possible improvement in the spatial cognition
ability upon TRT. T supplementation was performed by intramuscular injection of
testosterone enanthate (TE, 250 mg) to LOH patients every 2 weeks, for 3 times.
380
381 **Materials and Methods**
382
383   **Subjects : LOH patients**
384

This study was applied to men with symptoms of LOH who visited Male Menopause
Outpatient Clinic of Teikyo University Hospital. They had at least one symptom of
LOH including lethargy, general fatigue, malaise, depression, insomnia, sweating, hot
flashes, dizziness, night sweat, sexual dysfunction, and decreased libido.
We assessed symptom scores with several specific questionnaires including the
Heineman's Aging males' symptoms scale (AMS) for LOH, the international index of
erectile function (IIEF-5) and self-rating depression scale (SDS) (Heinemann et al 2003)
(Lunenfeld et al 2013) (Ide et al 2023). Endocrinologic data, including total T and free
T were evaluated. All blood samples were collected between 09:00 and 11:00 to
monitor endocrinological variables.
LOH was categorized by the AMS score (above ~45), total T levels (< ~300 ng/dL) and
free T levels (< 8.5 ~ 11.8 pg/mL) (Lunenfeld et al 2013, Wu et al 2010) (Tsuru et al

2023) (Ide et al 2023) .
398
399   Thirteen males diagnosed with LOH participated in the study. The average age was
400 52.3 ± 5.8 years old). Therefore, they were not older men (60-80 years), but rather
401 middle- aged men.
402 The clinical study was approved by the Teikyo University Ethics Committee (Approval
403 No. Teirin 15-148).
404
Serum hormone assay
Serum total T and luteinizing hormone (LH) were determined in the SRL Inc. (Tokyo,
Japan) by Cobas 8000 electrochemiluminescence immunoassay (Hitachi High-Tech Co. Ltd., Tokyo,
Japan) , using ECLusys Testosterone Kit (Roche Diagnostics, Indianapolis, IN,USA). Serum free T
was determined in the SRL Inc. by RIA, using the DPC Free Testosterone Kit (Mitsubishi Kagaku
Iatron Inc., Tokyo, Japan) (Iwamoto et al 2009).
411
412
413   **Procedures of TRT and maze navigation test**
414
The task is conducted according to the following procedures (see Fig. 2.1).
1. After explaining the outline of the TRT for LOH, the subjects complete the AMS,
SDS and IIEF5.
2. Show a video demonstration of the maze test (first time only).
3. Conduct the maze navigation test with three landmarks.
The maze with three landmarks are used. This is because the maze with three
landmarks is easier to try for LOH patients whose spatial cognition ability is lower than
young men.
When performing the maze test, maze number 1 should be entered for the first time,
maze number 2 for the second time, and maze number 3 for the third time, to avoid
mazes with identical goal positions.
4. The subjects are given 250 mg of testosterone enanthate (ENARMON DEPOT, ASKA
Pharmaceutical Co. Ltd., Tokyo, Japan) by intramuscular injection for 3 times.
5. After a 2-week interval, repeat steps 1, 3 and 4.
429
430   The above procedure is repeated four times for each patient. The overall flow is
431 shown in Figure 3.1.
432
433

Figure 3.1 The overall flow of TRT and maze navigation test

T is supplemented by intramuscular injection three times every two weeks. Blood

samples are collected on the first and forth (final) maze test day in order to measure

total T, free T, luteinizing hormone (LH), Hb and hematocrit (Ht) levels.

Statistical Analysis
Data are expressed as mean ± SEM.
For analysis of navigation maze tests, we used a paired t-test. For analysis of AMS,
SDS and IIEF-5 we used a paired t-test. For analysis of serum hormone levels, we used
Wilcoxon signed rank sum test. A difference was considered significant at a value of $*p$
< 0.05, $**p < 0.001$.
448
449
450 **Results**
451
452 **Improvement of maze navigation task scores by TRT**
453
454 After 6 weeks TRT, with three times T injection every two weeks, both the maze
455 escape time and the path distance are significantly shortened (Figure 3.2 and Figure
456 3.3).
457 A paired t-test revealed significant differences in the maze test scores before (0
458 week) and after 6 week treatments for both escape time and path distance.
459

460
461
462 **Figure 3.2** Biweekly maze test scores of escape time along TRT for 6 weeks. LOH
463 patients (n = 13). Vertical axis is escape time (second).
464
465

466
467
468 **Figure 3.3**   Biweekly maze test scores of path distance along TRT for 6 weeks.
469 LOH patients (n = 13). Vertical axis is path distance (arbitrary unit).
470
471
472 Representative recorded paths of a typical LOH patient along TRT is shown in
473 Figure 3.4.

(A)	(B)	(C)	(D)

485
486
487 **Figure 3.4**   Recorded paths on the maze test obtained from one LOH patient.
488 (A) before T-replacement (0 week), (B) after T-replacement (2nd week), (C) after T-
489 replacement (4th week), (D) after T-replacement (6th week).
490 Start (Arrow) and Goal (Red) are indicated.
491
492
493
494   **Improvements of AMS, SDS and IIEF-5 scores after TRT**
495
496   AMS showed a significant improvement, indicating the restorative effects of Aging
497 males' symptoms after TRT (Figure 3.5).

498 Other important markers, SDS and IIEF5, also showed improvement after TRT (Figures
499 3.6 and 3.7), indicating the recovery of depression and erectile function.
500
501

502
503
504 **Figure 3.5**   Biweekly scores of AMS along TRT for 6 weeks.
505
506

507
508
509 **Figure 3.6**   Biweekly scores of SDS (self-rating depression scale) along TRT
510 for 6 weeks.
511

IIEF5

512
513
514 **Figure 3.7** Biweekly scores of IIEF-5 (index of erectile function) along TRT for 6
515 weeks.
516
517
518 **Changes in serum hormone levels**
519
520   The results of blood tests before and after TRT indicated the significant elevation of
521 total T and/or free T levels (Table 2).

	Before TRT	6 weeks after TRT	P value
LH (mIU/ml)	4.8±0.6	<0.1	
Total T (ng/dL)	329.0±48.2	616.0±46.3	P<0.01
Free T (pg/ml)	6.4±0.7	13.7±1.2	P<0.01
Hb (g/dL)	14.5±0.2	15.3±0.4	P<0.01
Ht (%)	43.1±0.7	46.3±1.0	P<0.01

522
523 **Table 2** List of serum hormone levels and some other factors.
524   Luteinizing hormone (LH), total T, free T, Hemoglobin (Hb) and hematocrit (Ht).
525
526
527
528

Discussion

In the present study, spatial navigation ability was shown to be improved by elevation of the T levels in LOH patients by TRT. This might be caused by increase in the T levels in the brain hippocampus including CA1 neurons in which AR is localized and which are responsible to spatial cognition (Tsien et al 1996) (Tonegawa et al 1996) (Tabori et al 2005) (Hatanaka et al 2015).

The present results might be consistent to the previous observations that (1) spatial navigation ability (that is hippocampus-dependent function) is one of the few cognitive functions for which male performance is superior to female performance (Grön et al 2000), and that (2) the much higher T levels in male hippocampus than that in female hippocampus (e.g., approximately 100-fold higher) could support the better performance of male than female subjects (Tsien et al 1996) (Tonegawa et al 1996) (Hojo et al 2009) (Kato et al 2013) (Hatanaka et al 2015) (Kuwahara et al 2021) (Spritzer et al 2021).

Since LOH patients have lower T levels than those in healthy men, the present TRT-induced T elevation in LOH patients probably improved the spatial navigation task scores (e.g., escape time and path length) (Figures 3.2 and 3.3).

So far, this kind of investigation, particularly focusing on spatial navigation task for LOH patients, has not been performed intensively. Besides spatial navigation, there are other several cognitive abilities, e.g. attention, memory, calculation, verbal ability, auditory and visual processing. However, many previous studies suggest that these general cognitive abilities did not show significant improvement in older men (60 – 80 years old) by long term TRT (for half a year or two years) (Borst et al 2014) (Lisco et al 2020) (Gregori et al 2021). Importantly, the LOH patients in the present study are not older men but rather middle-aged men (between 45 and 60 years old) who have more neural plasticity than older men, and this may be one reason why the present short term TRT (6 weeks) showed effective improvements.

LOH patients exhibit symptoms that include hot flashes, depression, fatigue, and decreased libido due to decreased total T and/or free T. The main symptoms of LOH are well evaluated by the AMS, SDS, and IIE5 scores. We observed significant improvement of AMS, SDS, and IIE5 scores by TRT (Figures 3.5, 3.6), which correlated well with the improvements of spatial navigation scores (escaped time and path length) (Figures 3.2, 3.3).

Importantly, in Section 3, subjects (LOH patients, middle-aged) did not show mild cognitive impairment (MCI) symptoms or dementia. MCI symptoms were examined using the Revised Hasegawa's Dementia Scale (HDS-R) that is a simplified Mini-Mental State Examination（MMSE) (Gong et al 2021) (Arevalo-Rodriguez et al 2015). LOH subjects in Section 3 showed no MCI with almost full score in HDS-R, and this is probably because subjects were younger than 61 years old.

Interestingly, many studies recently reported that the long term TRT (e.g., for 2 years) for older patients (aged 65 years or older) did not significantly improve their cognitive

572 impairment (as judged from MMSE or HDS-R test) , though TRT showed significant
573 improvement in muscle strength and physical performance (Lisco et al 2020).
574
In the first part of this study, it was important to select and create the suitable spatial
cognition test in order to examine the TRT effect on male LOH patients.
Many researches were conducted on the spatial cognitive functions of men and
women (Hyde 2016), and it is generally accepted that men outperform women. For
example, men show higher scores than women in spatial route-learning task, virtual
maze task, and visuo-spatial task (Galea & Kimura 1993) (Moffat et al 1998) (Bosco et
al 2004).
582
On the other hand, there are also other types of spatial cognition tests including the
Mental Rotation Test (MRT) (Feng et al 2007) (Neubauer et al 2010). In MRT, subjects
should determine which objects are identical by rotating several objects in their mind.
By using MRT, men showed superior results to women in spatial cognition. However,
little is known about specific brain regions and gender dependent regions which are
excited during MRT. Therefor we cannot further analyze deeply T-dependence of MRT.
Therefore in the present study, we chose spatial route-learning task which is suitable
to evaluate men's cognitive ability that depends on T levels. Fortunately, spatial
navigation solely depends on the hippocampal CA1 neural function and AR activity
which are closely correlated with T levels (Tsien et al 1996) (Tonegawa et al 1996)
(Hojo et al 2009) (Kato et al 2013) (Hatanaka et al 2015) (Spritzer et al 2021) .
594
595

Acknowledgement

597   Department of Cognitive Neuroscience, Teikyo Univ. is an endowment department,
598 supported with an unrestricted grant from Japan Tobacco Inc.

599
600

Author Contributions

602   SK, HM, MK and SH conceived and designed the study. HM, YK, MK, MO-I and
603 M Soma conducted the experiments and analysis of the data. SK, HM and YK wrote the
604 manuscript. SH, MO-I and M Saito contributed to discussions. All authors provided
605 feedback on the manuscript.

Conflict of Interest Statement

608   The authors declare no conflict of interest.
609
610
611

References

Arevalo-Rodriguez I, Smailagic N, Roque IFM, Ciapponi A, Sanchez-Perez E, et al. 2015.

Mini-Mental State Examination (MMSE) for the detection of Alzheimer's disease
and other dementias in people with mild cognitive impairment (MCI). *Cochrane*
Database Syst Rev 2015: CD010783

Astur RS, Ortiz ML, Sutherland RJ. 1998. A characterization of performance by men and
women in a virtual Morris water task: a large and reliable sex difference. *Behav*
Brain Res 93: 185-90

Borst SE, Yarrow JF, Fernandez C, Conover CF, Ye F, et al. 2014. Cognitive effects of
testosterone and finasteride administration in older hypogonadal men. *Clin Interv*
Aging 9: 1327-33

Bosco A, Longoni AM, Vecchi T. 2004. Gender effects in spatial orientation: cognitive
profiles and mental strategies. *Appl Cogn Psychol* 18: 519-32

Feng J, Spence I, Pratt J. 2007. Playing an action video game reduces gender differences
in spatial cognition. *Psychol Sci* 18: 850-5

Galea LAM, Kimura D. 1993. Sex differences in route-learning. *Personality and*
Individual Differences 14: 53-65

Gong Q, Ishii M, Numata O, Xie W, Hirata T. 2021. Utility of a shortened Hasegawa
Dementia Scale Revised questionnaire to rapidly screen and diagnose Alzheimer's
disease. *Aging Med (Milton)* 4: 109-14

Gregori G, Celli A, Barnouin Y, Paudyal A, Armamento-Villareal R, et al. 2021. Cognitive
response to testosterone replacement added to intensive lifestyle intervention in
older men with obesity and hypogonadism: prespecified secondary analyses of a
randomized clinical trial. *The American journal of clinical nutrition* 114: 1590-99

Grön G, Wunderlich AP, Spitzer M, Tomczak R, Riepe MW. 2000. Brain activation during
human navigation: gender-different neural networks as substrate of performance.
Nat Neurosci 3: 404-8

Hatanaka Y, Hojo Y, Mukai H, Murakami G, Komatsuzaki Y, et al. 2015. Rapid increase
of spines by dihydrotestosterone and testosterone in hippocampal neurons:
Dependence on synaptic androgen receptor and kinase networks. *Brain Res* 1621:
121-32

Heinemann LA, Saad F, Zimmermann T, Novak A, Myon E, et al. 2003. The Aging Males'
Symptoms (AMS) scale: update and compilation of international versions. *Health*
Qual Life Outcomes 1: 15

Hojo Y, Higo S, Ishii H, Ooishi Y, Mukai H, et al. 2009. Comparison between
hippocampus-synthesized and circulation-derived sex steroids in the
hippocampus. *Endocrinology* 150: 5106-12

Hojo Y, Kawato S. 2018. Neurosteroids in Adult Hippocampus of Male and Female

651   Rodents: Biosynthesis and Actions of Sex Steroids. *Front Endocrinol (Lausanne)*
652   9: 183

653   Hojo Y, Okamoto M, Kato A, Higo S, Sakai F, et al. 2014. Neurosteroid synthesis in adult
654   female rat hippocampus, including androgens and allopregnanolone. *J Steroids*
655   *and Hormonal Sci*: Article S4

656   Huerta PT, Sun LD, Wilson MA, Tonegawa S. 2000. Formation of temporal memory
657   requires NMDA receptors within CA1 pyramidal neurons. *Neuron* 25: 473-80

658   Hyde JS. 2016. Sex and cognition: gender and cognitive functions. *Curr Opin Neurobiol*
659   38: 53-6

660   Ide H, Akehi Y, Fukuhara S, Ohira S, Ogawa S, et al. 2023. Summary of the clinical
661   practice manual for late-onset hypogonadism. *Int J Urol* 30: 422-30

662   Iwamoto T, Yanase T, Horie H, Namiki M, Okuyama A. 2009. Late-onset hypogonadism
663   (LOH) and androgens: validity of the measurement of free testosterone levels in
664   the diagnostic criteria in Japan. *Int J Urol* 16: 168-74

665   Kato A, Hojo Y, Higo S, Komatsuzaki Y, Murakami G, et al. 2013. Female hippocampal
666   estrogens have a significant correlation with cyclic fluctuation of hippocampal
667   spines. *Front Neural Circuits* 7: 149

668   Kuwahara N, Nicholson K, Isaacs L, MacLusky NJ. 2021. Androgen Effects on Neural
669   Plasticity. *Androg Clin Res Ther* 2: 216-30

670   Lisco G, Giagulli VA, De Tullio A, De Pergola G, Guastamacchia E, Triggiani V. 2020.
671   Age-Related Male Hypogonadism and Cognitive Impairment in the Elderly:
672   Focus on the Effects of Testosterone Replacement Therapy on Cognition.
673   *Geriatrics (Basel)* 5

674   Lunenfeld B, Mskhalaya G, Kalinchenko S, Tishova Y. 2013. Recommendations on the
675   diagnosis, treatment and monitoring of late-onset hypogonadism in men - a
676   suggested update. *Aging Male* 16: 143-50

677   McHugh TJ, Blum KI, Tsien JZ, Tonegawa S, Wilson MA. 1996. Impaired hippocampal
678   representation of space in CA1-specific NMDAR1 knockout mice. *Cell* 87: 1339-
679   49

680   Moffat SD, Hampson E, Hatzipantelis M. 1998. Navigation in a "Virtual" Maze: Sex
681   Differences and Correlation With Psychometric Measures of Spatial Ability in
682   Humans. *Evolution and Human Behavior* 19: 73-87

683   Murakami G, Hojo Y, Kato A, Komatsuzaki Y, Horie S, et al. 2018. Rapid nongenomic
684   modulation by neurosteroids of dendritic spines in the hippocampus: Androgen,
685   oestrogen and corticosteroid. *Journal of neuroendocrinology* 30

Neubauer AC, Bergner S, Schatz M. 2010. Two- vs. three-dimensional presentation of

mental rotation tasks: Sex differences and effects of training on performance and
brain activation. *Intelligence* 38: 529-39

Ooishi Y, Kawato S, Hojo Y, Hatanaka Y, Higo S, et al. 2012. Modulation of synaptic
plasticity in the hippocampus by hippocampus-derived estrogen and androgen. *J*
Steroid Biochem Mol Biol 131: 37-51

Spritzer MD, Jaeger ECB, Guo JD. 2021. Testosterone and Spatial Memory: Rodent
Models and Clinical Applications. *Androgens: Clinical Research and*
Therapeutics 2: 275-93

Tabori NE, Stewart LS, Znamensky V, Romeo RD, Alves SE, et al. 2005. Ultrastructural
evidence that androgen receptors are located at extranuclear sites in the rat
hippocampal formation. *Neuroscience* 130: 151-63

Tonegawa S, Tsien JZ, McHugh TJ, Huerta P, Blum KI, Wilson MA. 1996. Hippocampal
CA1-region-restricted knockout of NMDAR1 gene disrupts synaptic plasticity,
place fields, and spatial learning. *Cold Spring Harb Symp Quant Biol* 61: 225-38

Tsien JZ, Huerta PT, Tonegawa S. 1996. The essential role of hippocampal CA1 NMDA
receptor-dependent synaptic plasticity in spatial memory. *Cell* 87: 1327-38

Tsuru T, Tsujimura A, Mizushima K, Kurosawa M, Kure A, et al. 2023. International
Prostate Symptom Score and Quality of Life Index for Lower Urinary Tract
Symptoms Are Associated with Aging Males Symptoms Rating Scale for Late-
Onset Hypogonadism Symptoms. *World J Mens Health* 41: 101-09

Wu FC, Tajar A, Beynon JM, Pye SR, Silman AJ, et al. 2010. Identification of late-onset
hypogonadism in middle-aged and elderly men. *N Engl J Med* 363: 123-35

709