1

Testosterone replacement therapy ameliorates spatial cognitive function in age-related hypogonadism (LOH) patients: analysis with a virtual three-dimensional maze

Suguru Kawato[1,2,3,4†], Hideo Mukai[5], Yusuke Kai[5], Masaki Kimura[6], Mari Ogiue-Ikeda[1,2,4], Mika Soma[1,2,4,7], Minoru Saito[4] and Shigeo Horie[1]

[1]Department of Urology, Graduate School of Medicine, Juntendo Univ., Tokyo, Japan,
[2]Department of Cognitive Neuroscience, Faculty of Pharma-Science, Teikyo Univ., Tokyo, Japan,
[3]Department of Biophysics and Life Sciences, Graduate School of Arts and Sciences, The University of Tokyo, Tokyo, Japan,
[4]Department of Biosciences, College of Humanities and Sciences, Nihon University, Tokyo, Japan,
[5]Department of Computer Science, Graduate School of Science and Technology, Meiji University, Kanagawa, Japan,
[6]Department of Urology, School of Medicine, Teikyo Univ., Tokyo, Japan,
[7] Department of Anatomy and Neurobiology, Nippon Medical School, Tokyo, Japan.

Correspondence: Suguru Kawato, e-mail : skkawato@gmail.com

Keywords: testosterone, spatial navigation, late onset hypogonadism, testosterone replacement therapy, hippocampus

28 **Abstract**
29
When comparing the behavioral levels of men and women using a maze navigation
game created on a PC to investigate spatial cognitive function, many reports indicated
that men are significantly faster than women at finding the exit of the maze.
Analysis with functional MRI neuroimaging indicates that neural activation at
multiple sites were observed during a maze navigation task. Significantly different
regions were identified in men and women, with men showing significantly increased
activity in the hippocampus. In contrast, women showed increased activity in the
Brodmann area of the prefrontal cortex. When differences in the activated regions were
taken between men and women, the hippocampal activity showed the largest difference.
Since the hippocampal CA1 regions are known to be essentially responsible to
spatial navigation ability, the reported difference in spatial ability between men and
women could be due to higher testosterone (T) levels in mem than in women, because
the application of T can increase hippocampal CA1 neural synapses as observed in rats,
for example.
In the present study, we developed the original 3D maze navigation test on a PC to
accurately analyze spatial cognitive function, and this maze was used to analyze
differences in spatial cognitive function between young male and female volunteers.
The results showed that males took a distinctly shorter time and path length to escape
the maze than females. We applied this maze test to analyze the possible improvement
effects of spatial cognitive function by T-replacement therapy (TRT) on male patients
with late onset hypogonadism (LOH). LOH patients have significantly low levels of T
as well as high Aging males' symptoms scale (AMS) scores when compared with
healthy men. Upon TRT for 6 weeks, we revealed that the spatial cognitive ability of
LOH patients was significantly improved along the increase of T levels and
improvement of AMS scores.
The observed improvement of spatial cognitive ability of LOH patients could be
interpreted as effects of elevation of T levels in the hippocampal CA1 region where
androgen receptor (AR) is localized.
58
59
60

61 **Introduction**
62
Continuous visuospatial navigation in familiar and unfamiliar environments is a
requirement of daily life.
Visuospatial navigation is one of the few cognitive functions for which a reliable
gender-specific performance difference is well known (Astur et al 1998). For example,
men were significantly faster than women at finding the way out of maze.
Navigation through mazes can be combined with functional neuroimaging for
men and women. From the fMRI analysis, gender differences in brain neural activation
was computed with maze navigation tasks for male and female subjects (Grön et al

2000). Although there was great overlap between men and women in brain regions that were significantly activated, there were brain regions that were activated exclusively by the male and female groups. From the comparison of the neural activity, men showed significantly increased activity in the hippocampus proper, however, women showed increased activity of the prefrontal cortex, in contrast with men. (Grön et al 2000)

In the hippocampus, the glutamatergic neurons in the CA1 region are responsible for spatial cognition (Tonegawa et al 1996). This is evidenced by the fact that spatial cognition is selectively inhibited in mice in which N-methyl-D-aspartate (NMDA) receptors of glutamatergic neurons in the CA1 region of the hippocampus are specifically knockout (Tsien et al 1996) (McHugh et al 1996).

Previous studies of sex hormones in rats have shown that the level of testosterone (T) in the hippocampus is much higher in males (approximately 17 nM) than in females (approxi. 0.05 nM) (Hojo et al 2009) (Ooishi et al 2012) (Kato et al 2013) . T activates neurons upon binding to androgen receptor (AR).   AR is significantly expressed in glutamatergic neurons of the hippocampal CA1 region (Tabori et al 2005) (Hatanaka et al 2015). In addition, T level is much higher in male hippocampus than female hippocampus (Kato et al 2013). Circulating T levels are qualitatively proportional to hippocampal T levels (Kato et al 2013) (Hojo et al 2014) (Hojo & Kawato 2018). In view of these situations, we hypothesize that "higher T levels in males and lower T levels in females may have a significant effect on the difference in spatial cognitive function between men and women".

In Section 1, based on this hypothesis, we created a virtual maze escape game on a PC in order to find the spatial cognitive ability that may depend on T levels.

In Section 2, we applied this maze whether this maze is able to clearly determine differences in spatial cognitive ability between young men and women. We obtained the results that young men were significantly faster than young women in escaping from the maze.

In Section 3, we applied this maze navigation test to male patients with late onset hypogonadism (LOH) in order to examine whether T-replacement therapy (TRT) is effective to improve spatial cognitive ability by increase of T levels. It should be noted that because LOH patients have low T levels, TRT can elevate T levels of LOH patients, and thereby can improve Aging males' symptoms scale (AMS) scores (Heinemann et al 2003) (Lunenfeld et al 2013) (Ide et al 2023).   However, there had not been any challenges to investigate how TRT could improve spatial cognitive ability of LOH patients.

Section 1: Creation of 3D maze for spatial maze navigation test

1.1   Technical specification of maze

A 3D virtual maze program was created using the 3D modeling software Unity 5.6.1
(Unity Technologies). C# was used as the programming language.
The PC environment for the maze program and operation is as follows
OS: Windows 10 or 11,   64bit
CPU: Intel Core i7-6700HQ 2.60GHz
RAM: 16.0GB
GPU: Intel HD Graphics 530
Monitor: 15.6 in wide
The size of the maze is 6 squares (vertical) x 6 squares (horizontal) for a total of 36
squares (Figure 1.1, 1.2).
Fifteen sets of maze (Maze Number 1 – 15) are prepared. One of the maze can be
selected by choosing the group number (1, 2 or 3) on the title screen (see below), and
then automatically one Maze will be randomly selected. Note that one group consists of
5 mazes depending on goal positions.
128

129
130   Figure 1.1   Aerial view of the maze. The maze is 6x6 squares in size and can be
131   presented in a choice of 15 different ways.
132

133   **Start and goal points**

134
The 15 mazes are divided into three groups: one with a goal in square (6, 6), one
with a goal in square (1, 6), and one with a goal in square (6, 1). Subject can choose
which maze with a goal position you want to generate by entering the maze number on
the title screen (see below).
The start point is randomly selected from the four corners of the maze except for the
goal. For example, in Figure 1.2, the goal is located at square (6, 6) (red square), so the
starting point is chosen from the squares (1, 1), (1, 6) and (6, 1) (blue square).

Figure 1.2   How the start (blue) and goal (red) points are selected. If the start point is
in a red square, the goal point is randomly selected from the blue squares. Note that
subjects cannot see the goal that is hidden from the start by a wall.
147
148
149 **Operation of maze task**
150
Start , top title screen
When the maze navigation program is started, the top title screen appears. On the
title screen, the subject enters his/her name and maze number in order to select the type
of maze. Two types of mazes are available: maze without landmarks or maze with three
landmarks.
Before starting the maze test at the first time, "Demo Video" should be selected on
the title screen in order to get an overview of the present navigation maze task.
When 'walking' through the maze, subjects have a first-person view. During the
maze test, the arrow keys on the keyboard can be used to move forward with the up
arrow key, backward with the down arrow key, and turn right and left with the right and
left arrow keys, respectively. Turning was possible within the full range of 360 degree.
[↑] forward ,   [↓] backward ,   [→] right turn,   [←] left turn.
163

178 Figure 1.3   Top title screen. Press Return to next Selection screen.
179
180

181 *Selection of the time limit screen*
182
183

196
197 Figure 1.4   Time limit selection screen. There are two time limits (300 and 600
198 seconds) that can be chosen. Normally 300 sec is chosen.
199
200   Once the time limit is selected, the maze test begins. When the time limit is
201 exceeded, the word "End" is displayed.

202
203 Figure 1.5: Typical view inside the maze. During the control condition, the screen is
204 frozen at the last position of the active condition.
205
206   During the maze test, a subject proceeds in the first-person perspective and searches
207 for the correct path.
208
209
210   *Goal screen*
211   When a subject reaches the goal (the red wall) (Figure 1.6), a subject should presses
212 the return key to finish the task. Then the time taken to reach the goal and the distance
213 traveled during the maze task are recorded. At the same time, the screen will be back to
214 the title screen (see Fig 1.3).
215

216
217
218 Figure 1.6   The "Goal" is displayed on the goal wall. Pressing the Enter key to finish.
219
220   Landmarks in each maze are placed at the simple turned in the load to give a hint of
221 location. Landmarks are not placed at intersection of the convey (that is, decision
222 points).
223

236
237 Figure 1.7   Example of landmarks in the maze. Three types of landmarks(circle, star,
238 and triangle) are used. Only three walls have landmarks and each has different types of
239 landmarks.
240
241
242 **1.2   Assessment of Spatial Cognitive Ability**
243
Escape time
As a measure of spatial cognitive ability, the time taken to solve a maze test is
measured. This is called escape time.
The escape time is the number of seconds from the start of the maze test
(immediately after the time limit is selected) to the end of the maze test (when the
subject reaches the goal or the time limit is reached). The escape time is output to the
total score.csv file in the result folder directly under the C drive after the maze test is
completed.
252

Path Distance

As a measure of spatial cognitive ability other than escape time, the distance traveled during the maze test is recorded. This is called the path distance.

The path distance, as well as the escape time, is output to the total score.csv file located in the results folder directly under the C drive after the maze test is completed. The Euclidean distance for each second during the maze test is taken, and the sum of these Euclidean distances from start to goal is defined as the path distance (transit distance).

Section 2: Analysis of differences in spatial cognitive ability between young men and women using a virtual maze navigation task

Materials and Methods

Subjects

The subjects were 40 healthy students (20 men and 20 women) affiliated with Meiji University. The ages of the men and women subjects were 21.6 ± 1.5 and 20.6 ± 1.2 years, respectively. The experiment was notified to and approved by the Ethics Committee on Safety and Human Subjects for Experimental Research on Genetic Recombination, School of Science and Engineering, Meiji University (Approval No.: RIKEN 14-520). Subjects were informed about the research in person in advance and their consent was obtained by signing a research participation consent form.

Experimental Procedures

The experiment will be conducted according to the following procedures.

1. After explaining the outline of the experiment, show a demonstration video of the maze test.
2. Ask the subjects to perform the maze escape task without landmarks.
3. Ask the subjects to perform the maze escape task with three landmarks.
4. Repeat steps 2 and 3 four times with an interval of one week.

When performing the maze test, maze number 1 should be entered in the first time, maze number 2 in the second time, maze number 3 in the third time, and maze number 4 in the fourth time, to avoid mazes with identical goal positions.

Results

Maze navigation task, maze without/with landmarks.

In the maze navigation task, young men significantly outscored young women in the

maze without landmarks, however, this difference became smaller by the presence of
landmarks (Figure 2.1).
The escape time of men are shorter than that of women in both mazes without and
with landmarks. However, the presence of landmarks made the difference smaller.
In terms of path distance, men showed significantly shorter path distance than young
women in the maze without landmarks, however, by the presence of landmarks women
showed the same path distance as men (Figure 2.1).
303
304   Figure 2.2 shows the examples of the recorded path taken in the maze test at first
305 time, men follow the more efficient path than women toward the goal in the maze
306 without landmarks, but in the presence of   landmarks this gender difference became
307 smaller.
308
309 *Weekly comparison in order to check learning effects on the maze test*
310
During repeating the maze test, possible learning effects (with familiarity factors)
might improve the skill to escape the maze. If the subjects become accustomed to the
maze, their scores could be improved. We observed the decrease in the escape time and
the path distance by repeating women's maze tests without landmarks at different weeks
(1st , 2nd, 3rd, 4th week trial) (see Figure 2.1). However, in women's maze tests with
landmarks, these improvement effects by repetitive training were not observed.
On the other hand for men, the presence and absence of landmarks did not induce any
difference in both escape time and path distance, suggesting that men use geometric
cues (Grön et al 2000).
320

321
322
323 Figure 2.1   Weekly scores of escape time (Left) and path distance (Right) for young
324 men and women in maze tests. Vertical axis is escape time (second) and path distance
325 (arbitrary unit), respectively.
326 (●)Young men without landmarks, (○) Young Men with landmarks, (▲) Young
327 women without landmarks, (△) Young women with landmarks.

Differences between men and women are large in the absence of landmarks. Significant decreasing trends in escape time and path are observed only for women in the maze without landmarks.

Examples of recorded path during the maze test for young men and women are shown in Figure 2.2

Figure 2.2   From left to right, young men without landmarks, young men with landmarks,
young women without landmarks, young men with landmarks.
Start (Arrow) and Goal (Red) are indicated.

Section 3   Testosterone-Replacement Therapy (TRT) improved spatial cognition ability of LOH patients

In Section 2, the visuospatial navigation test successfully demonstrated differences in the spatial cognition ability between young men (with high T level) and young women (with low T level). The hippocampus is probably responsible to this spatial cognition ability from important previous reports. The spatial cognition ability of rats mainly depends on CA1 hippocampal neurons in which androgen receptor AR is localized (Tsien et al 1996) (McHugh et al 1996) (Tonegawa et al 1996) (Huerta et al 2000) (Tabori et al 2005) (Hatanaka et al 2015). Therefore, the observed difference in spatial cognition ability may significantly depend on the much higher hippocampal T levels in male than female as observed in rats (Hojo et al 2009) (Kato et al 2013). This hypothesis is dependent on the observation that nearly 20-fold T levels (in the hippocampus) and 100-fold T levels (in circulation) were observed in male rats than female rats (Hojo et al 2009) (Ooishi et al 2012) (Kato et al 2013).

It is shown also that T application to hippocampal CA1 glutamatergic neurons increased the number of synapses where memory is stored (Hatanaka et al 2015) (Murakami et al 2018) (Hojo & Kawato 2018). It may be therefore possible to find differences in maze navigation ability between individual men with high T and low T levels.

We applied spatial maze navigation task (with three landmarks) to LOH patients (with low T levels) in order to analyze a possible improvement in the spatial cognition

367 ability upon TRT. T supplementation was performed by intramuscular injection of
368 testosterone enanthate (TE, 250 mg) to LOH patients every 2 weeks, for 3 times.
369
370 **Materials and Methods**
371
372   **Subjects : LOH patients**
373
This study comprised and applied men with symptoms of LOH who visited Male
Menopause Outpatient Clinic of Teikyo University Hospital. They had at least one
symptom of LOH including lethargy, general fatigue, malaise, depression, insomnia,
sweating, hot flashes, dizziness, night sweat, sexual dysfunction, and decreased libido.
We assessed symptom scores with several specific questionnaires including the
Heineman's Aging males' symptoms scale (AMS) for LOH, the international index of
erectile function (IIEF-5) and self-rating depression scale (SDS) (Heinemann et al 2003)
(Lunenfeld et al 2013) (Ide et al 2023). Endocrinologic data, including total T and free
T were evaluated. All blood samples were collected between 09:00 and 11:00 to
monitor endocrinological variables.
LOH was categorized by the AMS score (above ~45), total T levels (< ~300 ng/dL) and
free T levels (< 8.5 ~ 11.8 pg/mL) (Lunenfeld et al 2013, Wu et al 2010) (Tsuru et al
2023) (Ide et al 2023) .
387
388   Thirteen males diagnosed with LOH participated in the study. The average age.was
389 52.3 ± 5.8 years old). Therefore, they were not older men (60-80 years), but rather
390 middle- aged men.
391 The clinical study was approved by the Teikyo University Ethics Committee (Approval
392 No. Teirin 15-148).
393
394
395   **Procedures of TRT and maze navigation test**
396
The task will be conducted according to the following procedures (see Fig. 2.1).
1. After explaining the outline of the TRT for LOH, the subjects complete the AMS,
SDS and IIEF5.
2. Show a video demonstration of the maze test (first time only).
3. Conduct the maze navigation test with three landmarks.
The maze with three landmarks are used. This is because the maze with three
landmarks is easier to try for LOH patients whose spatial cognition ability is lower than
young men.
When performing the maze test, maze number 1 should be entered for the first time,
maze number 2 for the second time, and maze number 3 for the third time, to avoid
mazes with identical goal positions.
4. The subjects are given 250 mg of testosterone enanthate by intramuscular injection for
3 times.

5. After a 2-week interval, repeat steps 1, 3 and 4.
411
412   The above procedure is repeated four times for each patient. The overall flow is
413   shown in Figure 3.1.
414
415

416
417
418   Figure 3.1   The overall flow of TRT and maze navigation test
419
420   T is supplemented by intramuscular injection three times every two weeks. Blood
421   samples are collected on the first and forth (final) maze test day in order to measure
422   total T, free T, luteinizing hormone (LH), Hb and hematocrit (Ht) levels.
423
424   **Statistical Analysis**
425   Data are expressed as mean ± SEM. For analysis of navigation maze tests as well as ,
426   we used a paired t-test. A difference was considered significant at a value of $*p < 0.05$,
427    $**p < 0.001$.
428
429
430   **Results**
431
432   **Improvement of maze navigation task scores by TRT**
433
434   After 6 weeks TRT, with three times T injection every two weeks, both the maze
435   escape time and the path distance are significantly shortened (Figure 3.2 and Figure
436   3.3).
437   A paired t-test revealed significant differences in the maze test scores before (0
438   week) and after 6 week treatments for both escape time and path distance.
439

440
441
442   Figure 3.2 Biweekly maze test scores of escape time along TRT. LOH patients (n = 13)
443   Vertical axis is escape time (second).
444
445

446
447
448   Figure 3.3 Biweekly maze test scores of path distance along TRT.
449   LOH patients (n = 13).   Vertical axis is path distance (arbitrary unit).

457 Representative recorded paths of one LOH patient along TRT is shown in
458 Figure 3.4.

(A)   (B)   (C)   (D)

471
472 Figure 3.4. Recorded paths on the maze test obtained from one LOH patient.
473 (A) before T-replacement (0 week), (B) after T-replacement (2nd week), (C) after T-
474 replacement (4th week), (D) 6th week.
475 Start (Arrow) and Goal (Red) are indicated.

476
477

478 **Changes in blood hormone levels**

479
480   The results of blood tests before and after TRT indicated the significant elevation
481 of total T and/or free T levels (Table 1) .

482
483 Table 1 : List of blood hormone levels and some other factors.
484   Luteinizing hormone (LH), total T, free T, Hemoglobin (Hb) and hematocrit (Ht)

	Before TRT	6 weeks after TRT	P value
LH (mIU/ml)	4.8±0.6	<0.1	P<0.01
TST (ng/dL)	329.0±48.2	616.0±46.3	P<0.01
F-TST (pg/ml)	6.4±0.7	13.7±1.2	P<0.01
Hb (g/dL)	14.5±0.2	15.3±0.4	P<0.01
Ht (%)	43.1±0.7	46.3±1.0	P<0.01

488   **Improvements of AMS, SDS and IIEF-5 scores after TRT**
489
490   AMS showed a significant improvement, indicating the restorative effects of Aging
491   males' symptoms after TRT (Figure 3.5).
492   Other important markers, SDS and IIEF5, also showed improvement after TRT (Figures
493   3.6 and 3.7), indicating the recovery of depression and erectile function.
494
495

496
497
498   Figure 3.5   Biweekly scores of AMS along TRT for 6 weeks.
499
500

SDS

501
502
503   Figure 3.6   Biweekly scores of SDS (self-rating depression scale) along TRT
504   for 6 weeks.

506
507
508
509 Figure 3.7   Biweekly scores of IIEF-5 (index of erectile function) along TRT for 6
510 weeks.

511
512

Discussion

In the present study, spatial navigation ability was shown to be improved by elevation
of the T levels in LOH patients by TRT. This might be caused by increase in the T
levels in the brain hippocampus including CA1 neurons in which AR is localized and
which are responsible to spatial cognition (Tsien et al 1996) (Tonegawa et al 1996)
(Tabori et al 2005) (Hatanaka et al 2015).
The present results might be consistent to the previous observations that (1) spatial
navigation ability (that is hippocampus-dependent function) is one of the few cognitive
functions for which male performance is superior to female performance (Grön et al
2000), and that (2) the much higher T levels in male hippocampus than that in female
hippocampus (e.g., approximately 100-fold higher) could support the better
performance of male than female subjects (Tsien et al 1996) (Tonegawa et al 1996)
(Hojo et al 2009) (Kato et al 2013) (Hatanaka et al 2015) (Kuwahara et al 2021)
(Spritzer et al 2021).
Since LOH patients have lower T levels than those in healthy men, the present TRT-
induced T elevation in LOH patients probably improved the spatial navigation task
scores (e.g., escape time and path length) (Figures 3.2 and 3.3).
So far, this kind of investigation, particularly focusing on spatial navigation task for
LOH patients, has not been performed intensively. Besides spatial navigation, there are
other several cognitive abilities, e.g. attention, memory, calculation, verbal ability,
auditory and visual processing. However, many previous studies suggest that these

general cognitive abilities did not show significant improvement in older men (60 – 80
years old) by long term TRT (for half a year or two years) (Borst et al 2014) (Lisco et al
2020) (Gregori et al 2021). Importantly, the LOH patients in the present study are not
older men but rather middle-aged men (between 45 and 60 years old) who have more
neural plasticity than older men, and this may be one reason why the present short term
TRT (6 weeks) showed effective improvements.
LOH patients exhibit symptoms that include hot flashes, depression, fatigue, and
decreased libido due to decreased total T and/or free T. The main symptoms of LOH are
well evaluated by the AMS, SDS, and IIE5 scores. We observed significant
improvement of AMS, SDS, and IIE5 scores by TRT (Figures 3.5, 3.6), which
correlated well with the improvements of spatial navigation scores (escaped time and
path length) (Figures 3.2, 3.3).
547
Importantly, in Section 3, subjects (LOH patients, middle-aged) did not show mild
cognitive impairment (MCI) symptoms or dementia. MCI symptoms were examined
using the Revised Hasegawa's Dementia Scale (HDS-R) that is a simplified Mini-
Mental State Examination（MMSE) (Gong et al 2021) (Arevalo-Rodriguez et al 2015).
LOH subjects in Section 3 showed no MCI with almost full score in HDS-R, and this is
probably because subjects were younger than 61 years old.
Interestingly, many studies recently reported that the long term TRT (e.g., for 2 years)
for older patients (aged 65 years or older) did not significantly improve their cognitive
impairment (as judged from MMSE or HDS-R test) , though TRT showed significant
improvement in muscle strength and physical performance (Lisco et al 2020).
558
In the first part of this study, it was important to select and create the suitable spatial
cognition test in order to examine the TRT effect on male LOH patients.
Many researches were conducted on the spatial cognitive functions of men and
women (Hyde 2016), and it is generally accepted that men outperform women. For
example, men show higher scores than women in spatial route-learning task, virtual
maze task, and visuo-spatial task (Galea & Kimura 1993) (Moffat et al 1998) (Bosco et
al 2004).
566
On the other hand, there are also other types of spatial cognition tests including the
Mental Rotation Test (MRT) (Feng et al 2007) (Neubauer et al 2010). In MRT, subjects
should determine which objects are identical by rotating several objects in their mind.
By using MRT, men showed superior results to women in spatial cognition. However,
little is known about specific brain regions and gender dependent regions which are
excited during MRT. Therefor we cannot further consider T-dependence of MRT.
Therefore in the present study, we chose spatial route-learning task which is suitable
to evaluate men's cognitive ability that depends on T levels. Fortunately, spatial
navigation solely depends on the hippocampal CA1 neural function and AR activity
which are closely correlated with T levels (Tsien et al 1996) (Tonegawa et al 1996)
(Hojo et al 2009) (Kato et al 2013) (Hatanaka et al 2015) (Spritzer et al 2021) .

578
579
Acknowledgement
580
581   Department of Cognitive Neuroscience, Teikyo Univ. is an endowment department,
582 supported with an unrestricted grant from Japan Tobacco Inc.
583
584

585 **Author Contributions**
586   SK, HM, MK and SH conceived and designed the study. HM, YK, MK, MO-I and
587 M Soma conducted the experiments and analysis of the data. SK, HM and YK wrote the
588 manuscript. SH, MO-I and M Saito contributed to discussions. All authors provided
589 feedback on the manuscript.
590

591 **Conflict of Interest Statement**
592   The authors declare no conflict of interest.
593
594
595

References
596
597

Arevalo-Rodriguez I, Smailagic N, Roque IFM, Ciapponi A, Sanchez-Perez E, et al. 2015.
Mini-Mental State Examination (MMSE) for the detection of Alzheimer's disease
and other dementias in people with mild cognitive impairment (MCI). *Cochrane*
Database Syst Rev 2015: CD010783
Astur RS, Ortiz ML, Sutherland RJ. 1998. A characterization of performance by men and
women in a virtual Morris water task: a large and reliable sex difference. *Behav*
Brain Res 93: 185-90
Borst SE, Yarrow JF, Fernandez C, Conover CF, Ye F, et al. 2014. Cognitive effects of
testosterone and finasteride administration in older hypogonadal men. *Clin Interv*
Aging 9: 1327-33
Bosco A, Longoni AM, Vecchi T. 2004. Gender effects in spatial orientation: cognitive
profiles and mental strategies. *Appl Cogn Psychol* 18: 519-32
Feng J, Spence I, Pratt J. 2007. Playing an action video game reduces gender differences
in spatial cognition. *Psychol Sci* 18: 850-5
Galea LAM, Kimura D. 1993. Sex differences in route-learning. *Personality and*
Individual Differences 14: 53-65
Gong Q, Ishii M, Numata O, Xie W, Hirata T. 2021. Utility of a shortened Hasegawa
Dementia Scale Revised questionnaire to rapidly screen and diagnose Alzheimer's
disease. *Aging Med (Milton)* 4: 109-14
Gregori G, Celli A, Barnouin Y, Paudyal A, Armamento-Villareal R, et al. 2021. Cognitive
response to testosterone replacement added to intensive lifestyle intervention in
older men with obesity and hypogonadism: prespecified secondary analyses of a
randomized clinical trial. *The American journal of clinical nutrition* 114: 1590-99

Grön G, Wunderlich AP, Spitzer M, Tomczak R, Riepe MW. 2000. Brain activation during
human navigation: gender-different neural networks as substrate of performance.
Nat Neurosci 3: 404-8

Hatanaka Y, Hojo Y, Mukai H, Murakami G, Komatsuzaki Y, et al. 2015. Rapid increase
of spines by dihydrotestosterone and testosterone in hippocampal neurons:
Dependence on synaptic androgen receptor and kinase networks. *Brain Res* 1621:
121-32

Heinemann LA, Saad F, Zimmermann T, Novak A, Myon E, et al. 2003. The Aging Males'
Symptoms (AMS) scale: update and compilation of international versions. *Health*
Qual Life Outcomes 1: 15

Hojo Y, Higo S, Ishii H, Ooishi Y, Mukai H, et al. 2009. Comparison between
hippocampus-synthesized and circulation-derived sex steroids in the
hippocampus. *Endocrinology* 150: 5106-12

Hojo Y, Kawato S. 2018. Neurosteroids in Adult Hippocampus of Male and Female
Rodents: Biosynthesis and Actions of Sex Steroids. *Front Endocrinol (Lausanne)*
9: 183

Hojo Y, Okamoto M, Kato A, Higo S, Sakai F, et al. 2014. Neurosteroid synthesis in adult
female rat hippocampus, including androgens and allopregnanolone. *J Steroids*
and Hormonal Sci: Article S4

Huerta PT, Sun LD, Wilson MA, Tonegawa S. 2000. Formation of temporal memory
requires NMDA receptors within CA1 pyramidal neurons. *Neuron* 25: 473-80

Hyde JS. 2016. Sex and cognition: gender and cognitive functions. *Curr Opin Neurobiol*
38: 53-6

Ide H, Akehi Y, Fukuhara S, Ohira S, Ogawa S, et al. 2023. Summary of the clinical
practice manual for late-onset hypogonadism. *Int J Urol* 30: 422-30

Kato A, Hojo Y, Higo S, Komatsuzaki Y, Murakami G, et al. 2013. Female hippocampal
estrogens have a significant correlation with cyclic fluctuation of hippocampal
spines. *Front Neural Circuits* 7: 149

Kuwahara N, Nicholson K, Isaacs L, MacLusky NJ. 2021. Androgen Effects on Neural
Plasticity. *Androg Clin Res Ther* 2: 216-30

Lisco G, Giagulli VA, De Tullio A, De Pergola G, Guastamacchia E, Triggiani V. 2020.
Age-Related Male Hypogonadism and Cognitive Impairment in the Elderly:
Focus on the Effects of Testosterone Replacement Therapy on Cognition.
Geriatrics (Basel) 5

Lunenfeld B, Mskhalaya G, Kalinchenko S, Tishova Y. 2013. Recommendations on the
diagnosis, treatment and monitoring of late-onset hypogonadism in men - a
suggested update. *Aging Male* 16: 143-50

McHugh TJ, Blum KI, Tsien JZ, Tonegawa S, Wilson MA. 1996. Impaired hippocampal
representation of space in CA1-specific NMDAR1 knockout mice. *Cell* 87: 1339-
49

Moffat SD, Hampson E, Hatzipantelis M. 1998. Navigation in a "Virtual" Maze: Sex
Differences and Correlation With Psychometric Measures of Spatial Ability in
Humans. *Evolution and Human Behavior* 19: 73-87

Murakami G, Hojo Y, Kato A, Komatsuzaki Y, Horie S, et al. 2018. Rapid nongenomic
665   modulation by neurosteroids of dendritic spines in the hippocampus: Androgen,
666   oestrogen and corticosteroid. *Journal of neuroendocrinology* 30

667   Neubauer AC, Bergner S, Schatz M. 2010. Two- vs. three-dimensional presentation of
668   mental rotation tasks: Sex differences and effects of training on performance and
669   brain activation. *Intelligence* 38: 529-39

670   Ooishi Y, Kawato S, Hojo Y, Hatanaka Y, Higo S, et al. 2012. Modulation of synaptic
671   plasticity in the hippocampus by hippocampus-derived estrogen and androgen. *J*
672   *Steroid Biochem Mol Biol* 131: 37-51

673   Spritzer MD, Jaeger ECB, Guo JD. 2021. Testosterone and Spatial Memory: Rodent
674   Models and Clinical Applications. *Androgens: Clinical Research and*
675   *Therapeutics* 2: 275-93

676   Tabori NE, Stewart LS, Znamensky V, Romeo RD, Alves SE, et al. 2005. Ultrastructural
677   evidence that androgen receptors are located at extranuclear sites in the rat
678   hippocampal formation. *Neuroscience* 130: 151-63

679   Tonegawa S, Tsien JZ, McHugh TJ, Huerta P, Blum KI, Wilson MA. 1996. Hippocampal
680   CA1-region-restricted knockout of NMDAR1 gene disrupts synaptic plasticity,
681   place fields, and spatial learning. *Cold Spring Harb Symp Quant Biol* 61: 225-38

682   Tsien JZ, Huerta PT, Tonegawa S. 1996. The essential role of hippocampal CA1 NMDA
683   receptor-dependent synaptic plasticity in spatial memory. *Cell* 87: 1327-38

684   Tsuru T, Tsujimura A, Mizushima K, Kurosawa M, Kure A, et al. 2023. International
685   Prostate Symptom Score and Quality of Life Index for Lower Urinary Tract
686   Symptoms Are Associated with Aging Males Symptoms Rating Scale for Late-
687   Onset Hypogonadism Symptoms. *World J Mens Health* 41: 101-09

688   Wu FC, Tajar A, Beynon JM, Pye SR, Silman AJ, et al. 2010. Identification of late-onset
689   hypogonadism in middle-aged and elderly men. *N Engl J Med* 363: 123-35

690