

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27

Testosterone replacement therapy ameliorates spatial cognitive function in age-related hypogonadism (LOH) patients: analysis with a virtual three-dimensional maze

Suguru Kawato^{1,2,3,4†}, Hideo Mukai⁵, Yusuke Kai⁵, Masaki Kimura⁶, Mari Ogiue-Ikeda^{1,2,4}, Mika Soma^{1,2,4,7}, Minoru Saito⁴ and Shigeo Horie¹

¹Department of Urology, Graduate School of Medicine, Juntendo Univ., Tokyo, Japan,
²Department of Cognitive Neuroscience, Faculty of Pharma-Science, Teikyo Univ., Tokyo, Japan,
³Department of Biophysics and Life Sciences, Graduate School of Arts and Sciences, The University of Tokyo, Tokyo, Japan,
⁴Department of Biosciences, College of Humanities and Sciences, Nihon University, Tokyo, Japan,
⁵Department of Computer Science, Graduate School of Science and Technology, Meiji University, Kanagawa, Japan,
⁶Department of Urology, School of Medicine, Teikyo Univ., Tokyo, Japan,
⁷ Department of Anatomy and Neurobiology, Nippon Medical School, Tokyo, Japan.

Correspondence: Suguru Kawato, e-mail : skkawato@gmail.com

Keywords: testosterone, spatial navigation, late onset hypogonadism, testosterone replacement therapy, hippocampus

28 **Abstract**

29

30 When comparing the behavioral levels of men and women using a maze navigation
31 game created on a PC to investigate spatial cognitive function, many reports indicated
32 that men are significantly faster than women at finding the exit of the maze.

33 Analysis with functional MRI neuroimaging indicates that neural activation at
34 multiple sites were observed during a maze navigation task. Significantly different
35 regions were identified in men and women, with men showing significantly increased
36 activity in the hippocampus. In contrast, women showed increased activity in the
37 Brodmann area of the prefrontal cortex. When differences in the activated regions were
38 taken between men and women, the hippocampal activity showed the largest difference.

39 Since the hippocampal CA1 regions are known to be essentially responsible to
40 spatial navigation ability, the reported difference in spatial ability between men and
41 women could be due to higher testosterone (T) levels in men than in women, because
42 the application of T can increase hippocampal CA1 neural synapses as observed in rats,
43 for example.

44 In the present study, we developed the original 3D maze navigation test on a PC to
45 accurately analyze spatial cognitive function, and this maze was used to analyze
46 differences in spatial cognitive function between young male and female volunteers.
47 The results showed that males took a distinctly shorter time and path length to escape
48 the maze than females. We applied this maze test to analyze the possible improvement
49 effects of spatial cognitive function by T-replacement therapy (TRT) on male patients
50 with late onset hypogonadism (LOH). LOH patients have significantly low levels of T
51 as well as high Aging males' symptoms scale (AMS) scores when compared with
52 healthy men. Upon TRT for 6 weeks, we revealed that the spatial cognitive ability of
53 LOH patients was significantly improved along the increase of T levels and
54 improvement of AMS scores.

55 The observed improvement of spatial cognitive ability of LOH patients could be
56 interpreted as effects of elevation of T levels in the hippocampal CA1 region where
57 androgen receptor (AR) is localized.

58

59

60

61 **Introduction**

62

63 Continuous visuospatial navigation in familiar and unfamiliar environments is a
64 requirement of daily life.

65 Visuospatial navigation is one of the few cognitive functions for which a reliable
66 gender-specific performance difference is well known (Astur et al 1998). For example,
67 men were significantly faster than women at finding the way out of maze.

68 Navigation through mazes can be combined with functional neuroimaging for
69 men and women. From the fMRI analysis, gender differences in brain neural activation
70 was computed with maze navigation tasks for male and female subjects (Grön et al

71 2000). Although there was great overlap between men and women in brain regions that
72 were significantly activated, there were brain regions that were activated exclusively by
73 the male and female groups. From the comparison of the neural activity, men showed
74 significantly increased activity in the hippocampus proper, however, women showed
75 increased activity of the prefrontal cortex, in contrast with men. (Grön et al 2000)

76 In the hippocampus, the glutamatergic neurons in the CA1 region are responsible for
77 spatial cognition (Tonegawa et al 1996). This is evidenced by the fact that spatial
78 cognition is selectively inhibited in mice in which N-methyl-D-aspartate (NMDA)
79 receptors of glutamatergic neurons in the CA1 region of the hippocampus are
80 specifically knockout (Tsien et al 1996) (McHugh et al 1996).

81 Previous studies of sex hormones in rats have shown that the level of testosterone
82 (T) in the hippocampus is much higher in males (approximately 17 nM) than in females
83 (approx. 0.05 nM) (Hojo et al 2009) (Ooishi et al 2012) (Kato et al 2013) . T activates
84 neurons upon binding to androgen receptor (AR). AR is significantly expressed in
85 glutamatergic neurons of the hippocampal CA1 region (Tabori et al 2005) (Hatanaka et
86 al 2015). In addition, T level is much higher in male hippocampus than female
87 hippocampus (Kato et al 2013). Circulating T levels are qualitatively proportional to
88 hippocampal T levels (Kato et al 2013) (Hojo et al 2014) (Hojo & Kawato 2018). In
89 view of these situations, we hypothesize that “higher T levels in males and lower T
90 levels in females may have a significant effect on the difference in spatial cognitive
91 function between men and women”.

92
93 In Section 1, based on this hypothesis, we created a virtual maze escape game on a
94 PC in order to find the spatial cognitive ability that may depend on T levels.

95 In Section 2, we applied this maze whether this maze is able to clearly determine
96 differences in spatial cognitive ability between young men and women. We obtained the
97 results that young men were significantly faster than young women in escaping from the
98 maze.

99 In Section 3, we applied this maze navigation test to male patients with late onset
100 hypogonadism (LOH) in order to examine whether T-replacement therapy (TRT) is
101 effective to improve spatial cognitive ability by increase of T levels. It should be noted
102 that because LOH patients have low T levels, TRT can elevate T levels of LOH patients,
103 and thereby can improve Aging males’ symptoms scale (AMS) scores (Heinemann et al
104 2003) (Lunenfeld et al 2013) (Ide et al 2023). However, there had not been any
105 challenges to investigate how TRT could improve spatial cognitive ability of LOH
106 patients.

107
108
109

110 **Section 1: Creation of 3D maze for spatial maze navigation test**

111

112 **1.1 Technical specification of maze**

113

114 A 3D virtual maze program was created using the 3D modeling software Unity 5.6.1
115 (Unity Technologies). C# was used as the programming language.

116 The PC environment for the maze program and operation is as follows

117 OS: Windows 10 or 11, 64bit

118 CPU: Intel Core i7-6700HQ 2.60GHz

119 RAM: 16.0GB


120 GPU: Intel HD Graphics 530

121 Monitor: 15.6 in wide

122 The size of the maze is 6 squares (vertical) x 6 squares (horizontal) for a total of 36
123 squares (Figure 1.1, 1.2).

124 Fifteen sets of maze (Maze Number 1 – 15) are prepared. One of the maze can be
125 selected by choosing the group number (1, 2 or 3) on the title screen (see below), and
126 then automatically one Maze will be randomly selected. Note that one group consists of
127 5 mazes depending on goal positions.

128



129

130 **Figure 1.1** Aerial view of the maze. The maze is 6x6 squares in size and can be
131 presented in a choice of 15 different ways.

132


133 **Start and goal points**

134

135 The 15 mazes are divided into three groups: one with a goal in square (6, 6), one
136 with a goal in square (1, 6), and one with a goal in square (6, 1). Subject can choose
137 which maze with a goal position you want to generate by entering the maze number on
138 the title screen (see below).

139 The start point is randomly selected from the four corners of the maze except for the
140 goal. For example, in Figure 1.2, the goal is located at square (6, 6) (red square), so the
141 starting point is chosen from the squares (1, 1), (1, 6) and (6, 1) (blue square).

142



143

144 **Figure 1.2** How the start (blue) and goal (red) points are selected. If the start point is
145 in a red square, the goal point is randomly selected from the blue squares. Note that
146 subjects cannot see the goal that is hidden from the start by a wall.

147

148

149 **Operation of maze task**

150

151 *Start , top title screen*


152 When the maze navigation program is started, the top title screen appears. On the
153 title screen, the subject enters his/her name and maze number in order to select the type
154 of maze. Two types of mazes are available: maze without landmarks or maze with three
155 landmarks.

156 Before starting the maze test at the first time, “Demo Video” should be selected on
157 the title screen in order to get an overview of the present navigation maze task.

158 When ‘walking’ through the maze, subjects have a first-person view. During the
159 maze test, the arrow keys on the keyboard can be used to move forward with the up
160 arrow key, backward with the down arrow key, and turn right and left with the right and
161 left arrow keys, respectively. Turning was possible within the full range of 360 degree.
162 [↑] forward , [↓] backward , [→] right turn, [←] left turn.

163

164



177

178 **Figure 1.3** Top title screen. Press Return to next Selection screen.


179

180

181 *Selection of the time limit screen*

182

183




196

197 **Figure 1.4** Time limit selection screen. There are two time limits (300 and 600
198 seconds) that can be chosen. Normally 300 sec is chosen.

199

200 Once the time limit is selected, the maze test begins. When the time limit is
201 exceeded, the word "End" is displayed.



202

203 **Figure 1.5** Typical view inside the maze. During the control condition, the screen is
204 frozen at the last position of the active condition.

205

206 During the maze test, a subject proceeds in the first-person perspective and searches
207 for the correct path.

208

209

210 *Goal screen*

211 When a subject reaches the goal (the red wall) (Figure 1.6), a subject should presses
212 the return key to finish the task. Then the time taken to reach the goal and the distance
213 traveled during the maze task are recorded. At the same time, the screen will be back to
214 the title screen (see Fig 1.3).

215




216
217

Figure 1.6 The “Goal” is displayed on the goal wall. Pressing the Enter key to finish.

219
220
221
222
223

Landmarks in each maze are placed at the simple turned in the load to give a hint of location. Landmarks are not placed at intersection of the convey (that is, decision points).



236

Figure 1.7 Example of landmarks in the maze. Three types of landmarks(circle, star, and triangle) are used. Only three walls have landmarks and each has different types of landmarks.

240
241

1.2 Assessment of Spatial Cognitive Ability

243

Escape time

245
246

As a measure of spatial cognitive ability, the time taken to solve a maze test is measured. This is called escape time.

247
248

The escape time is the number of seconds from the start of the maze test (immediately after the time limit is selected) to the end of the maze test (when the subject reaches the goal or the time limit is reached). The escape time is output to the total score.csv file in the result folder directly under the C drive after the maze test is completed.

251
252

253 ***Path Distance***

254 As a measure of spatial cognitive ability other than escape time, the distance traveled
255 during the maze test is recorded. This is called the path distance.

256 The path distance, as well as the escape time, is output to the total score.csv file
257 located in the results folder directly under the C drive after the maze test is completed.
258 The Euclidean distance for each second during the maze test is taken, and the sum of
259 these Euclidean distances from start to goal is defined as the path distance (transit
260 distance).

261

262

263

264 **Section 2: Analysis of differences in spatial cognitive ability between young men
265 and women using a virtual maze navigation task**

266

267 **Materials and Methods**

268

269 **Subjects**

270 The subjects were 40 healthy students (20 men and 20 women) affiliated with Meiji
271 University. The ages of the men and women subjects were 21.6 ± 1.5 and 20.6 ± 1.2 years,
272 respectively. The experiment was notified to and approved by the Ethics Committee on
273 Safety and Human Subjects for Experimental Research on Genetic Recombination, School
274 of Science and Engineering, Meiji University (Approval No.: RIKEN 14-520). Subjects
275 were informed about the research in person in advance and their consent was obtained by
276 signing a research participation consent form.

277

278 **Experimental Procedures**

279 The experiment will be conducted according to the following procedures.

280

281 1. After explaining the outline of the experiment, show a demonstration video of the maze
282 test.

283 2. Ask the subjects to perform the maze escape task without landmarks.

284 3. Ask the subjects to perform the maze escape task with three landmarks.

285 4. Repeat steps 2 and 3 four times with an interval of one week.

286

287 When performing the maze test, maze number 1 should be entered in the first time,
288 maze number 2 in the second time, maze number 3 in the third time, and maze number 4 in
289 the fourth time, to avoid mazes with identical goal positions.

290

291 **Results**

292

293 Maze navigation task, maze without/with landmarks.

294

295 In the maze navigation task, young men significantly outscored young women in the

296 maze without landmarks, however, this difference became smaller by the presence of
297 landmarks (Figure 2.1).

298 The escape time of men are shorter than that of women in both mazes without and
299 with landmarks. However, the presence of landmarks made the difference smaller.

300 In terms of path distance, men showed significantly shorter path distance than young
301 women in the maze without landmarks, however, by the presence of landmarks women
302 showed the same path distance as men (Figure 2.1).

303

304 Figure 2.2 shows the examples of the recorded path taken in the maze test at first
305 time, men follow the more efficient path than women toward the goal in the maze
306 without landmarks, but in the presence of landmarks this gender difference became
307 smaller.

308


309 *Weekly comparison in order to check learning effects on the maze test*

310

311 During repeating the maze test, possible learning effects (with familiarity factors)
312 might improve the skill to escape the maze. If the subjects become accustomed to the
313 maze, their scores could be improved. We observed the decrease in the escape time and
314 the path distance by repeating women's maze tests without landmarks at different weeks
315 (1st, 2nd, 3rd, 4th week trial) (see Figure 2.1). However, in women's maze tests with
316 landmarks, these improvement effects by repetitive training were not observed.

317 On the other hand for men, the presence and absence of landmarks did not induce any
318 difference in both escape time and path distance, suggesting that men use geometric
319 cues (Grön et al 2000).

320



321

322

323 **Figure 2.1** Weekly scores of escape time (Left) and path distance (Right) for young
324 men and women in maze tests. Vertical axis is escape time (second) and path distance
325 (arbitrary unit), respectively.

326 (●) Young men without landmarks, (○) Young Men with landmarks, (▲) Young
327 women without landmarks, (△) Young women with landmarks.


328 Differences between men and women are large in the absence of landmarks.
329 Significant decreasing trends in escape time and path are observed only for women in
330 the maze without landmarks.

331

332 Examples of recorded path during the maze test for young men and women are
333 shown in Figure 2.2

334

335



336

337

338 **Figure 2.2** From left to right, young men without landmarks, young men with
339 landmarks,

340 young women without landmarks, young men with landmarks.

341 Start (Arrow) and Goal (Red) are indicated.

342

343

344

345 **Section 3 Testosterone-Replacement Therapy (TRT) improved spatial cognition** 346 **ability of LOH patients**

347

348 In Section 2, the visuospatial navigation test successfully demonstrated differences
349 in the spatial cognition ability between young men (with high T level) and young
350 women (with low T level). The hippocampus is probably responsible to this spatial
351 cognition ability from important previous reports. The spatial cognition ability of rats
352 mainly depends on CA1 hippocampal neurons in which androgen receptor AR is
353 localized (Tsien et al 1996) (McHugh et al 1996) (Tonegawa et al 1996) (Huerta et al
354 2000) (Tabori et al 2005) (Hatanaka et al 2015). Therefore, the observed difference in
355 spatial cognition ability may significantly depend on the much higher hippocampal T
356 levels in male than female as observed in rats (Hojo et al 2009) (Kato et al 2013). This
357 hypothesis is dependent on the observation that nearly 20-fold T levels (in the
358 hippocampus) and 100-fold T levels (in circulation) were observed in male rats than
359 female rats (Hojo et al 2009) (Ooishi et al 2012) (Kato et al 2013).

360

361 It is shown also that T application to hippocampal CA1 glutamatergic neurons
362 increased the number of synapses where memory is stored (Hatanaka et al 2015)
363 (Murakami et al 2018) (Hojo & Kawato 2018). It may be therefore possible to find
364 differences in maze navigation ability between individual men with high T and low T
365 levels.

365

366 We applied spatial maze navigation task (with three landmarks) to LOH patients
(with low T levels) in order to analyze a possible improvement in the spatial cognition

367 ability upon TRT. T supplementation was performed by intramuscular injection of
368 testosterone enanthate (TE, 250 mg) to LOH patients every 2 weeks, for 3 times.

369

370 **Materials and Methods**

371

372 **Subjects : LOH patients**

373

374 This study was applied to men with symptoms of LOH who visited Male Menopause
375 Outpatient Clinic of Teikyo University Hospital. They had at least one symptom of
376 LOH including lethargy, general fatigue, malaise, depression, insomnia, sweating, hot
377 flashes, dizziness, night sweat, sexual dysfunction, and decreased libido.

378 We assessed symptom scores with several specific questionnaires including the
379 Heineman's Aging males' symptoms scale (AMS) for LOH, the international index of
380 erectile function (IIEF-5) and self-rating depression scale (SDS) (Heinemann et al 2003)
381 (Lunenfeld et al 2013) (Ide et al 2023). Endocrinologic data, including total T and free
382 T were evaluated. All blood samples were collected between 09:00 and 11:00 to
383 monitor endocrinological variables.

384 LOH was categorized by the AMS score (above ~45), total T levels ($< \sim 300$ ng/dL) and
385 free T levels ($< 8.5 \sim 11.8$ pg/mL) (Lunenfeld et al 2013, Wu et al 2010) (Tsuru et al
386 2023) (Ide et al 2023) .

387

388 Thirteen males diagnosed with LOH participated in the study. The average age was
389 52.3 ± 5.8 years old). Therefore, they were not older men (60-80 years), but rather
390 middle- aged men.

391 The clinical study was approved by the Teikyo University Ethics Committee (Approval
392 No. Teirin 15-148).

393

394 **Serum hormone assay**

395 Serum total T and luteinizing hormone (LH) were determined in the SRL Inc. (Tokyo,
396 Japan) by Cobas 8000 electrochemiluminescence immunoassay (Hitachi High-Tech Co. Ltd., Tokyo,
397 Japan) , using ECLusys Testosterone Kit (Roche Diagnostics, Indianapolis, IN,USA). Serum free T
398 was determined in the SRL Inc. by RIA, using the DPC Free Testosterone Kit (Mitsubishi Kagaku
399 Iatron Inc., Tokyo, Japan) (Iwamoto et al 2009).

400

401

402 **Procedures of TRT and maze navigation test**

403

404 The task is conducted according to the following procedures (see Fig. 2.1).

405 1. After explaining the outline of the TRT for LOH, the subjects complete the AMS,
406 SDS and IIEF5.

407 2. Show a video demonstration of the maze test (first time only).

408 3. Conduct the maze navigation test with three landmarks.

409 The maze with three landmarks are used. This is because the maze with three

410 landmarks is easier to try for LOH patients whose spatial cognition ability is lower than
411 young men.

412 When performing the maze test, maze number 1 should be entered for the first time,
413 maze number 2 for the second time, and maze number 3 for the third time, to avoid
414 mazes with identical goal positions.

415 4. The subjects are given 250 mg of testosterone enanthate (ENARMON DEPOT, ASKA
416 Pharmaceutical Co. Ltd., Tokyo, Japan) by intramuscular injection for 3 times.


417 5. After a 2-week interval, repeat steps 1, 3 and 4.

418

419 The above procedure is repeated four times for each patient. The overall flow is
420 shown in Figure 3.1.

421

422



423

424

425 **Figure 3.1** The overall flow of TRT and maze navigation test.

426 T is supplemented by intramuscular injection three times every two weeks. Blood
427 samples are collected on the first and forth (final) maze test day in order to measure
428 total T, free T, luteinizing hormone (LH), Hb and hematocrit (Ht) levels.

429

430

431 **Statistical Analysis**

432 Data are expressed as mean \pm SEM. For analysis of navigation maze tests, we used a
433 paired t-test. For analysis of serum hormone levels, we used Wilcoxon test. A difference
434 was considered significant at a value of * $p < 0.05$, ** $p < 0.001$.

435

436

437 **Results**

438


439 **Improvement of maze navigation task scores by TRT**

440

441 After 6 weeks TRT, with three times T injection every two weeks, both the maze
442 escape time and the path distance are significantly shortened (Figure 3.2 and Figure
443 3.3).

444 A paired t-test revealed significant differences in the maze test scores before (0
445 week) and after 6 week treatments for both escape time and path distance.

446




447

448

449 **Figure 3.2** Biweekly maze test scores of escape time along TRT for 6 weeks. LOH
450 patients (n = 13). Vertical axis is escape time (second).

451

452



453
454

Figure 3.3 Biweekly maze test scores of path distance along TRT for 6 weeks. LOH patients (n = 13). Vertical axis is path distance (arbitrary unit).

456
457
458

Representative recorded paths of one LOH patient along TRT is shown in Figure 3.4.

459
460
461




Figure 3.4 Recorded paths on the maze test obtained from one LOH patient. (A) before T-replacement (0 week), (B) after T-replacement (2nd week), (C) after T-replacement (4th week), (D) after T-replacement (6th week). Start (Arrow) and Goal (Red) are indicated.

474
475
476
477
478
479

Changes in serum hormone levels

480
481
482
483
484

The results of blood tests before and after TRT indicated the significant elevation of total T and/or free T levels (Table 1).

485 **Table 1** List of serum hormone levels and some other factors.
486 Luteinizing hormone (LH), total T, free T, Hemoglobin (Hb) and hematocrit (Ht).

	Before TRT	6 weeks after TRT	P value
LH (mIU/ml)	4.8±0.6	<0.1	
Total T (ng/dL)	329.0±48.2	616.0±46.3	P<0.01
Free T (pg/ml)	6.4±0.7	13.7±1.2	P<0.01
Hb (g/dL)	14.5±0.2	15.3±0.4	P<0.01
Ht (%)	43.1±0.7	46.3±1.0	P<0.01

487

488

489

490 **Improvements of AMS, SDS and IIEF-5 scores after TRT**


491

492 AMS showed a significant improvement, indicating the restorative effects of Aging
493 males' symptoms after TRT (Figure 3.5).

494 Other important markers, SDS and IIEF5, also showed improvement after TRT (Figures
495 3.6 and 3.7), indicating the recovery of depression and erectile function.


496

497




498
499
500
501
502

Figure 3.5 Biweekly scores of AMS along TRT for 6 weeks.



503
504
505
506
507

Figure 3.6 Biweekly scores of SDS (self-rating depression scale) along TRT for 6 weeks.



508
509
510

511 **Figure 3.7** Biweekly scores of IIEF-5 (index of erectile function) along TRT for 6
512 weeks.

513
514

515 **Discussion**

516

517 In the present study, spatial navigation ability was shown to be improved by elevation
518 of the T levels in LOH patients by TRT. This might be caused by increase in the T
519 levels in the brain hippocampus including CA1 neurons in which AR is localized and
520 which are responsible to spatial cognition (Tsien et al 1996) (Tonegawa et al 1996)
521 (Tabori et al 2005) (Hatanaka et al 2015).

522 The present results might be consistent to the previous observations that (1) spatial
523 navigation ability (that is hippocampus-dependent function) is one of the few cognitive
524 functions for which male performance is superior to female performance (Grön et al
525 2000), and that (2) the much higher T levels in male hippocampus than that in female
526 hippocampus (e.g., approximately 100-fold higher) could support the better
527 performance of male than female subjects (Tsien et al 1996) (Tonegawa et al 1996)
528 (Hojo et al 2009) (Kato et al 2013) (Hatanaka et al 2015) (Kuwahara et al 2021)
529 (Spritzer et al 2021).

530 Since LOH patients have lower T levels than those in healthy men, the present TRT-
531 induced T elevation in LOH patients probably improved the spatial navigation task
532 scores (e.g., escape time and path length) (Figures 3.2 and 3.3).

533 So far, this kind of investigation, particularly focusing on spatial navigation task for
534 LOH patients, has not been performed intensively. Besides spatial navigation, there are
535 other several cognitive abilities, e.g. attention, memory, calculation, verbal ability,
536 auditory and visual processing. However, many previous studies suggest that these
537 general cognitive abilities did not show significant improvement in older men (60 – 80

538 years old) by long term TRT (for half a year or two years) (Borst et al 2014) (Lisco et al
539 2020) (Gregori et al 2021). Importantly, the LOH patients in the present study are not
540 older men but rather middle-aged men (between 45 and 60 years old) who have more
541 neural plasticity than older men, and this may be one reason why the present short term
542 TRT (6 weeks) showed effective improvements.

543 LOH patients exhibit symptoms that include hot flashes, depression, fatigue, and
544 decreased libido due to decreased total T and/or free T. The main symptoms of LOH are
545 well evaluated by the AMS, SDS, and IIE5 scores. We observed significant
546 improvement of AMS, SDS, and IIE5 scores by TRT (Figures 3.5, 3.6), which
547 correlated well with the improvements of spatial navigation scores (escaped time and
548 path length) (Figures 3.2, 3.3).

549
550 Importantly, in Section 3, subjects (LOH patients, middle-aged) did not show mild
551 cognitive impairment (MCI) symptoms or dementia. MCI symptoms were examined
552 using the Revised Hasegawa's Dementia Scale (HDS-R) that is a simplified Mini-
553 Mental State Examination (MMSE) (Gong et al 2021) (Arevalo-Rodriguez et al 2015).
554 LOH subjects in Section 3 showed no MCI with almost full score in HDS-R, and this is
555 probably because subjects were younger than 61 years old.
556 Interestingly, many studies recently reported that the long term TRT (e.g., for 2 years)
557 for older patients (aged 65 years or older) did not significantly improve their cognitive
558 impairment (as judged from MMSE or HDS-R test), though TRT showed significant
559 improvement in muscle strength and physical performance (Lisco et al 2020).

560
561 In the first part of this study, it was important to select and create the suitable spatial
562 cognition test in order to examine the TRT effect on male LOH patients.

563 Many researches were conducted on the spatial cognitive functions of men and
564 women (Hyde 2016), and it is generally accepted that men outperform women. For
565 example, men show higher scores than women in spatial route-learning task, virtual
566 maze task, and visuo-spatial task (Galea & Kimura 1993) (Moffat et al 1998) (Bosco et
567 al 2004).

568
569 On the other hand, there are also other types of spatial cognition tests including the
570 Mental Rotation Test (MRT) (Feng et al 2007) (Neubauer et al 2010). In MRT, subjects
571 should determine which objects are identical by rotating several objects in their mind.
572 By using MRT, men showed superior results to women in spatial cognition. However,
573 little is known about specific brain regions and gender dependent regions which are
574 excited during MRT. Therefore we cannot further analyze deeply T-dependence of MRT.

575 Therefore in the present study, we chose spatial route-learning task which is suitable
576 to evaluate men's cognitive ability that depends on T levels. Fortunately, spatial
577 navigation solely depends on the hippocampal CA1 neural function and AR activity
578 which are closely correlated with T levels (Tsien et al 1996) (Tonegawa et al 1996)
579 (Hojo et al 2009) (Kato et al 2013) (Hatanaka et al 2015) (Spritzer et al 2021).

580

581

582 **Acknowledgement**

583 Department of Cognitive Neuroscience, Teikyo Univ. is an endowment department,
584 supported with an unrestricted grant from Japan Tobacco Inc.

585

586

587 **Author Contributions**

588 SK, HM, MK and SH conceived and designed the study. HM, YK, MK, MO-I and
589 M Soma conducted the experiments and analysis of the data. SK, HM and YK wrote the
590 manuscript. SH, MO-I and M Saito contributed to discussions. All authors provided
591 feedback on the manuscript.

592

593 **Conflict of Interest Statement**

594 The authors declare no conflict of interest.

595

596

597

598

References

599

- 600 Arevalo-Rodriguez I, Smailagic N, Roque IFM, Ciapponi A, Sanchez-Perez E, et al. 2015.
601 Mini-Mental State Examination (MMSE) for the detection of Alzheimer's disease
602 and other dementias in people with mild cognitive impairment (MCI). *Cochrane*
603 *Database Syst Rev* 2015: CD010783
- 604 Astur RS, Ortiz ML, Sutherland RJ. 1998. A characterization of performance by men and
605 women in a virtual Morris water task: a large and reliable sex difference. *Behav*
606 *Brain Res* 93: 185-90
- 607 Borst SE, Yarrow JF, Fernandez C, Conover CF, Ye F, et al. 2014. Cognitive effects of
608 testosterone and finasteride administration in older hypogonadal men. *Clin Interv*
609 *Aging* 9: 1327-33
- 610 Bosco A, Longoni AM, Vecchi T. 2004. Gender effects in spatial orientation: cognitive
611 profiles and mental strategies. *Appl Cogn Psychol* 18: 519-32
- 612 Feng J, Spence I, Pratt J. 2007. Playing an action video game reduces gender differences
613 in spatial cognition. *Psychol Sci* 18: 850-5
- 614 Galea LAM, Kimura D. 1993. Sex differences in route-learning. *Personality and*
615 *Individual Differences* 14: 53-65
- 616 Gong Q, Ishii M, Numata O, Xie W, Hirata T. 2021. Utility of a shortened Hasegawa
617 Dementia Scale Revised questionnaire to rapidly screen and diagnose Alzheimer's
618 disease. *Aging Med (Milton)* 4: 109-14
- 619 Gregori G, Celli A, Barnouin Y, Paudyal A, Armamento-Villareal R, et al. 2021. Cognitive

- 620 response to testosterone replacement added to intensive lifestyle intervention in
621 older men with obesity and hypogonadism: prespecified secondary analyses of a
622 randomized clinical trial. *The American journal of clinical nutrition* 114: 1590-99
- 623 Grön G, Wunderlich AP, Spitzer M, Tomczak R, Riepe MW. 2000. Brain activation during
624 human navigation: gender-different neural networks as substrate of performance.
625 *Nat Neurosci* 3: 404-8
- 626 Hatanaka Y, Hojo Y, Mukai H, Murakami G, Komatsuzaki Y, et al. 2015. Rapid increase
627 of spines by dihydrotestosterone and testosterone in hippocampal neurons:
628 Dependence on synaptic androgen receptor and kinase networks. *Brain Res* 1621:
629 121-32
- 630 Heinemann LA, Saad F, Zimmermann T, Novak A, Myon E, et al. 2003. The Aging Males'
631 Symptoms (AMS) scale: update and compilation of international versions. *Health*
632 *Qual Life Outcomes* 1: 15
- 633 Hojo Y, Higo S, Ishii H, Ooishi Y, Mukai H, et al. 2009. Comparison between
634 hippocampus-synthesized and circulation-derived sex steroids in the
635 hippocampus. *Endocrinology* 150: 5106-12
- 636 Hojo Y, Kawato S. 2018. Neurosteroids in Adult Hippocampus of Male and Female
637 Rodents: Biosynthesis and Actions of Sex Steroids. *Front Endocrinol (Lausanne)*
638 9: 183
- 639 Hojo Y, Okamoto M, Kato A, Higo S, Sakai F, et al. 2014. Neurosteroid synthesis in adult
640 female rat hippocampus, including androgens and allopregnanolone. *J Steroids*
641 *and Hormonal Sci*: Article S4
- 642 Huerta PT, Sun LD, Wilson MA, Tonegawa S. 2000. Formation of temporal memory
643 requires NMDA receptors within CA1 pyramidal neurons. *Neuron* 25: 473-80
- 644 Hyde JS. 2016. Sex and cognition: gender and cognitive functions. *Curr Opin Neurobiol*
645 38: 53-6
- 646 Ide H, Akehi Y, Fukuhara S, Ohira S, Ogawa S, et al. 2023. Summary of the clinical
647 practice manual for late-onset hypogonadism. *Int J Urol* 30: 422-30
- 648 Iwamoto T, Yanase T, Horie H, Namiki M, Okuyama A. 2009. Late-onset hypogonadism
649 (LOH) and androgens: validity of the measurement of free testosterone levels in
650 the diagnostic criteria in Japan. *Int J Urol* 16: 168-74
- 651 Kato A, Hojo Y, Higo S, Komatsuzaki Y, Murakami G, et al. 2013. Female hippocampal
652 estrogens have a significant correlation with cyclic fluctuation of hippocampal
653 spines. *Front Neural Circuits* 7: 149
- 654 Kuwahara N, Nicholson K, Isaacs L, MacLusky NJ. 2021. Androgen Effects on Neural
655 Plasticity. *Androg Clin Res Ther* 2: 216-30

- 656 Lisco G, Giagulli VA, De Tullio A, De Pergola G, Guastamacchia E, Triggiani V. 2020.
657 Age-Related Male Hypogonadism and Cognitive Impairment in the Elderly:
658 Focus on the Effects of Testosterone Replacement Therapy on Cognition.
659 *Geriatrics (Basel)* 5
- 660 Lunenfeld B, Mskhalaya G, Kalinchenko S, Tishova Y. 2013. Recommendations on the
661 diagnosis, treatment and monitoring of late-onset hypogonadism in men - a
662 suggested update. *Aging Male* 16: 143-50
- 663 McHugh TJ, Blum KI, Tsien JZ, Tonegawa S, Wilson MA. 1996. Impaired hippocampal
664 representation of space in CA1-specific NMDAR1 knockout mice. *Cell* 87: 1339-
665 49
- 666 Moffat SD, Hampson E, Hatzipantelis M. 1998. Navigation in a “Virtual” Maze: Sex
667 Differences and Correlation With Psychometric Measures of Spatial Ability in
668 Humans. *Evolution and Human Behavior* 19: 73-87
- 669 Murakami G, Hojo Y, Kato A, Komatsuzaki Y, Horie S, et al. 2018. Rapid nongenomic
670 modulation by neurosteroids of dendritic spines in the hippocampus: Androgen,
671 oestrogen and corticosteroid. *Journal of neuroendocrinology* 30
- 672 Neubauer AC, Bergner S, Schatz M. 2010. Two- vs. three-dimensional presentation of
673 mental rotation tasks: Sex differences and effects of training on performance and
674 brain activation. *Intelligence* 38: 529-39
- 675 Ooishi Y, Kawato S, Hojo Y, Hatanaka Y, Higo S, et al. 2012. Modulation of synaptic
676 plasticity in the hippocampus by hippocampus-derived estrogen and androgen. *J*
677 *Steroid Biochem Mol Biol* 131: 37-51
- 678 Spritzer MD, Jaeger ECB, Guo JD. 2021. Testosterone and Spatial Memory: Rodent
679 Models and Clinical Applications. *Androgens: Clinical Research and*
680 *Therapeutics* 2: 275-93
- 681 Tabori NE, Stewart LS, Znamensky V, Romeo RD, Alves SE, et al. 2005. Ultrastructural
682 evidence that androgen receptors are located at extranuclear sites in the rat
683 hippocampal formation. *Neuroscience* 130: 151-63
- 684 Tonegawa S, Tsien JZ, McHugh TJ, Huerta P, Blum KI, Wilson MA. 1996. Hippocampal
685 CA1-region-restricted knockout of NMDAR1 gene disrupts synaptic plasticity,
686 place fields, and spatial learning. *Cold Spring Harb Symp Quant Biol* 61: 225-38
- 687 Tsien JZ, Huerta PT, Tonegawa S. 1996. The essential role of hippocampal CA1 NMDA
688 receptor-dependent synaptic plasticity in spatial memory. *Cell* 87: 1327-38
- 689 Tsuru T, Tsujimura A, Mizushima K, Kurosawa M, Kure A, et al. 2023. International
690 Prostate Symptom Score and Quality of Life Index for Lower Urinary Tract
691 Symptoms Are Associated with Aging Males Symptoms Rating Scale for Late-

692 Onset Hypogonadism Symptoms. *World J Mens Health* 41: 101-09
693 Wu FC, Tajar A, Beynon JM, Pye SR, Silman AJ, et al. 2010. Identification of late-onset
694 hypogonadism in middle-aged and elderly men. *N Engl J Med* 363: 123-35
695