Network Analysis of Rare Single-Nucleotide Polymorphisms Suggests a Central Role for Type 17 Immune Responses in Trauma-induced, Genotypically Associated Hypo-Inflammation and Critical Illness

Fayten El-Dehaibi[1,*], Ruben Zamora[1,*], Jinling Yin[1], Timothy R. Billiar[1], Yoram Vodovotz[1,2,†]

Running Title: Trauma-associated SNPs and Th17 Inflammation

[1]Department of Surgery, University of Pittsburgh, Pittsburgh, PA 15213

[2] Center for Inflammation and Regeneration Modeling, McGowan Institute for Regenerative Medicine, University of Pittsburgh, Pittsburgh, PA 15219

[*]Co-first authors

[†] **To whom correspondence should be addressed at**:
  Department of Surgery
  University of Pittsburgh
  W944 Starzl Biomedical Sciences Tower
  200 Lothrop St.
  Pittsburgh, PA 15213
  Tel.: 412-647-5609
  Fax: 412-383-5946
  E-mail: vodovotzy@upmc.edu

Keywords: trauma, SNP, Th17, hypo-inflammation, RORA

Word Count: 5009 words

Figure Count: 17 (11 figures, 6 tables)

Printed on July 8, 2025

ABSTRACT

35
Critical illness stemming from severe traumatic injury is a leading cause of morbidity and
mortality worldwide, involves the dysfunction of multiple organ systems, and is driven at least in
part by dysregulated inflammation that involves type 17 immunity. We and others have
demonstrated a genetic predisposition to adverse critical illness outcomes associated with single-
nucleotide polymorphisms (SNPs) in novel loci distinct from those which impact type 17
immune responses yet acting in concert with those responses. We have recently developed
SNPScanner, an algorithm that enables rapid scanning through a large SNP dataset and
associated inflammation biomarkers and clinical data, and in the present study leveraged this
algorithm in concert with existing bioinformatics tools to define networks of interaction among
rare SNPs (defined operationally as being present in 5-10% of patients due to the study cohort
size) assessed in critically ill trauma patients. RAR-related orphan receptor A (RORA), a
transcription factor central to the differentiation of Th17 cells, was inferred as a hub gene via its
SNP rs4774381; stratification of trauma patients based on homozygous genotype at this SNP
pointed to distinct organ dysfunction trajectories associated with network-defined hypo-
inflammation. Further, the SNP rs11919443 in the non-hub *TM4SF19* gene distinguished
genotype-associated systemic IL-17A trajectories, and associated hypo-inflammation with
adverse outcomes. In contrast, no Th17-related hub genes were identified in a control group of
SNPs associated algorithmically with adverse clinical outcomes but with genotypically indistinct
systemic inflammatory responses; the main inferred hub gene in this control group was solute
carrier family 2, facilitated glucose transporter member 1 (*SLC2A1*)/glucose transporter 1
(GLUT1). Secondary analysis showed several statistically significant differences in circulating
inflammatory mediators not including IL-17A, and also associated network-based hypo-
inflammation with adverse outcomes in this control group. This study thus extends our prior
work aimed at defining genetic predisposition to dysregulated inflammation and pathophysiology
in the context of critical illness, and points to a crucial role for type 17 immune responses.

1. INTRODUCTION

Trauma-induced critical illness, in which organ dysfunction is associated with dysregulated inflammation (i.e., an abnormal or imbalanced immune response with detrimental effects to the body), represents a major global health burden (1-3). Despite similar injury profiles and demographics, and the use of standardized protocols, critically ill trauma patients exhibit heterogeneous outcomes (4). We and others have proposed that genetic variation among patients may account for some of this heterogeneity (5). Previous studies have identified distinct dynamic inflammatory networks after severe trauma as a function of demographics (e.g., age) (6), injury timing (7) and characteristics (8-11), or clinical outcomes (12, 13). We have also characterized endotypes of blunt trauma patients based on early inflammatory patterns (14, 15), multiple organ dysfunction severity (16), or multi-'omics patterns (17, 18). A key immune pathway that emerged from these studies involves interleukin-17A (IL-17A), suggesting a novel, early role for type 17 immune responses in trauma-associated critical illness (13, 15, 19-21).

Our studies have also highlighted a role for genetic polymorphisms in the disordered systemic inflammation that is characteristic of critically ill trauma patients (19, 22-25). These studies, particularly those showing that specific single-nucleotide polymorphisms (SNPs) are linked to sub-acute mortality in trauma patients with early divergent inflammatory trajectories associated with the Multiple Organ Dysfunction Syndrome (MODS) (13, 24), imply that genetic factors may influence the variability in patient outcomes and acute inflammation in critically ill trauma patients.

We have previously developed novel computational approaches to aid in the search for novel, yet common (defined as present in 20% or more of the patients in our study cohort) SNPs associated with altered trajectories of systemic inflammation and subsequent clinical outcomes (24), and have combined this discovery program with existing computational tools that facilitate the discovery of SNPs in networks of genes acting in concert (25). In the present study, we leveraged these two approaches to scan a dataset of >500,000 SNPs for networks of both rare and common SNPs associated with adverse outcomes in critically ill trauma patients. In these analyses, we focused on defining hub genes and assessing clinical outcomes and inflammatory profiles in patient subgroups stratified based on genotypes at those hub genes. Our results suggest a key role for Th17 differentiation and associate this with systemic hypo-inflammation.

2.   MATERIALS AND METHODS

2.1 *Trauma patient cohort*

Studies were carried out using a cohort of 380 blunt trauma survivors with roughly equivalent distributions of homozygous and alternative homozygous genotypes at any of >500,000 SNPs described previously (24). In brief, patients were enrolled by screening after presentation to the emergency department of the UPMC Presbyterian hospital (a Level 1 trauma center) between May 5, 2004, to May 2, 2012. Patients eligible for enrollment in the study were at least 18 years of age, admitted to the intensive care unit (ICU) after being resuscitated, and, per treating physician, were expected to live more than 24h. Exclusion criteria were isolated head injury, pregnancy, and penetrating trauma. Informed consent was obtained from each patient or next of kin as per the University of Pittsburgh Institutional Review Board (IRB; Protocol No. MOD08010232-19 / PRO08010232) and in accordance with the Declaration of Helsinki. Blood samples for DNA and inflammatory biomarker analysis were obtained upon admission to the trauma bay, at first 24 h of hospitalization, and daily thereafter for 7 days.

Clinical data, including injury severity score (ISS), abbreviated injury scale (AIS) score, Marshall Multiple Organ Dysfunction (MOD) score, admission Shock Index, ICU LOS, hospital LOS, and days on mechanical ventilation were collected from the hospital inpatient electronic and trauma registry database. ISS (26) and AIS (27) were calculated for each patient by a single trauma surgeon after attending radiology evaluations were finalized. The ISS is based on an anatomical scoring system that provides an overall score for patients with multiple injuries. Each injury is assigned an AIS score, allocated to one of six body regions: head, face, chest, abdomen, extremities (including pelvis), and external. As an index of organ dysfunction, the MOD score (28) ranging from 0 to 24 was calculated. In brief, six variables were obtained from the electronic trauma data registry including (a) the respiratory system (PO_2/FIO_2 ratio); (b) the renal system (serum creatinine concentration); (c) the hepatic system (serum bilirubin concentration); (d) the hematologic system (platelet count); (e) the central nervous system (Glasgow Coma Scale); and (f) the cardiovascular system- the pressure-adjusted heart rate (PAR).

The overall characteristics of the patients in the trauma cohort were as follows (mean ± SD): (ISS) = 19.0 ± 10.1 [min: 1; max: 54]; mean age: 49.3 ± 19.2 years (min: 18y; max: 90y); 119 women, 261 men; 96.84% whites, 2.63% blacks and 0.53% others.

2.2 *Genomic analyses*

For the blunt trauma patient cohorts, DNA was prepared from whole blood samples and analyzed using Illumina® arrays as described previously (19, 22, 23). Whole blood samples were collected into EDTA-treated tubes and DNA was extracted using the QIAamp® DNA Blood Midi Kit (QIAGEN, Valencia, CA) as per manufacturer's specifications. Single nucleotide polymorphism genotyping was performed with 200 ng of genomic DNA input using the Human Core Exome-24 v1.1 BeadChip (Illumina, San Diego, CA) following the manufacturer's Infinium® HTS Assay protocol. This array interrogated 551,839 SNPs, including both SNPs previously implicated in genome-wide association studies as well as additional coding and non-coding SNPs spaced along the human genome. Briefly, DNA was denatured in 0.1N NaOH and neutralized prior to isothermal amplification. Amplified DNA was fragmented and then hybridized to locus-specific 50mers that make up the array for 16-24 h with rocking at 48°C. After removal of unbound or non-specifically annealed DNA, single base extension of the 50mer

135   oligonucleotides was performed with labeled nucleotides, which were scanned using an Illumina
136   iScan with autoloader 2.x. Data analysis was performed using Illumina Genome Studio 2.0.
137
2.3 *Assays of inflammation biomarkers*
Whole-blood samples were drawn in EDTA-treated tubes, which were kept on ice and
centrifuged to obtain plasma and then stored at -80°C until assays were performed. The
Luminex™ 100 IS analyzer (Luminex, Austin, TX) and Human Cytokine/Chemokine
MILLIPLEX™ Panel kit (Millipore Corporation, Billerica, MA) were used to measure plasma
levels of Eotaxin (CCL11), interleukin (IL)-1β, IL-1 receptor antagonist (IL-1RA), IL-2, soluble
IL-2 receptor-α (sIL-2Rα), IL-4, IL-5, IL-6, IL-7, IL-8 (CCL8), IL-10, IL-13, IL-15, IL-17A,
interferon (IFN)-α, IFN-γ, IFN-γ inducible protein (IP)-10 (CXCL10), monokine induced by
gamma interferon (MIG; CXCL9), macrophage inflammatory protein (MIP)-1α (CCL3), MIP-1β
(CCL4), monocyte chemotactic protein (MCP)-1 (CCL2), granulocyte-macrophage colony
stimulating factor (GM-CSF) and tumor necrosis factor alpha (TNF-α). Human Th17
MILLIPLEX™ Panel kit (Millipore Corporation, Billerica, MA) was used to measure IL-9, IL-
21, IL-22, IL-23, IL-17E/25, and IL-33. NO2-/NO3 levels were measured by a Griess Reagent
colorimetric assay (Cayman Chemical, Ann Arbor, MI). IL-1 receptor-like 1/ suppression of
tumorigenicity-2 (ST2) was measured by a sandwich ELISA assay (R&D Systems, Minneapolis,
MN). All cytokine/chemokine concentrations are given in pg/ml; NO2-/NO3- concentrations are
in μM. Experimental data are shown as mean ± SEM.
155
2.4 *FLAME and associated bioinformatics analyses*
Functional Enrichment Analysis was performed using the web application FLAME (v2.0)
(29). Two lists of SNPs— the first associated with trajectories of systemic inflammation typical
that did not differ based on genotype (which served as controls) vs. SNPs associated with
genotype-specific differential systemic inflammatory responses as determined in our prior study
(24) —were separately entered into the SNP tab of the Input page (Table 1). To reduce potential
artifacts in this analysis, 15 SNPs associated with 12 or more genotype-specific differential
systemic inflammatory responses as determined in our prior study (22) were selected out of 7482
SNPs which met the initial criterion of having at least one statistically different biomarker
difference as a function of SNP genotype in our original study (22). In the control group for
evenly distributed genotypes, SNPs were extracted from the group of 567 SNPs with zero
significantly different mediators as a function of genotype at each given SNP (described initially
in (22). To decrease the likelihood of artifactual associations among SNPs and to obtain
meaningful information about associations of SNPs with inflammation, we imposed an
additional condition of having at least 3-4 significant differences in clinical outcomes as a
function of SNP genotype which is to find SNPs while having no significant differences in
inflammation biomarkers. This led to down-selection from 567 to 16 SNPs.
To examine less-common SNPs (referred to as 'rare' for our purposes) we performed the
process described to extract differential inflammatory SNPs from the human trauma group but
used a 5–10% to 90–95% ratio of homozygous prevalence (minimum 20 patients) to select 7212
SNPs from the original >500,000. The previous method was applied to run Mann-Whitney U
analysis on the same 30 inflammatory biomarkers on patients from each group. Given the
population size disparity between the homozygotes of the rare SNP set, we sought to reduce the

likelihood of artifactual results by considering p-values less than 10-5 as significant, as opposed
to 0.05 used earlier. This stringency provided 131 SNPs associated with at least one statistically
significant biomarker, and 15 SNPs with up to two. The omission of X-linked SNPs yielded 12
rare somatic SNPs with at least two significantly different inflammatory biomarkers. The rare
SNP subset was also checked for control SNPs in the same manner described for the evenly
distributed SNPs. Screening clinical outcomes with Mann-Whitney U reduced 433 rare SNPs
with zero significantly different mediators ($p < 0.05$) down to 10 with at least 3-4 statistically
different outcomes.
Each list of corresponding genes found was separately inputted into FLAME's Functional
Enrichment page for whole genome search, with the enrichment tools set to gProfiler (30) and
enrichR (31) and the significance threshold set to 0.05. The following data sources were selected:
Molecular Function, Cellular Component and Biological Function from the gene ontology
knowledgebase (GO)(32), and the Kyoto Encyclopedia of Genes and Genomics (KEGG) (33).
192
2.5 *Statistical analyses*
Kruskal-Wallis Analysis of Variance (ANOVA) on Ranks, One-Way and Two-Way
ANOVAs were conducted to analyze the differences in ISS, AIS, Marshall MODScore, and
time-dependent changes in plasma biomarkers in AA and BB patients using SigmaPlot (Systat
Software, San Jose, CA) as appropriate. Two-Way ANOVA was followed by the Holm-Sidak
post hoc test to isolate which group(s) differ from the others. Chi-square analysis of the
proportions of observations for patients with rs10404939 or rs11117564 genotype was performed
using SigmaPlot (Systat Software, San Jose, CA).
201
2.6 *Network analysis*
Dynamic Network Analysis (DyNA) was carried out to define, in a granular fashion, the
central inflammatory network nodes as a function of both time and patient subgroup. DyNA
networks were created over six consecutive time periods (D1-D2, D2-D3, D3-D4, D4-D5, D5-
D6, D6-D7) using MATLAB® software as described previously(34-36). Connections, defined as
the number of inflammatory biomarkers that were positively or negatively correlated across time
intervals, were created if the Pearson correlation coefficient between any two nodes (biomarkers)
at the same time-interval was greater or equal to a threshold of an absolute value of 0.95, as
appropriate. The network complexity for each time-interval was calculated using the following
formula: Sum $(N_1 + N_2 + \ldots + N_n)/(n - 1)$, where N represents the number of connections for each
biomarker and n is the total number of biomarkers analyzed.
213
2.7 *Th17 Cell Subset Inference/Correlation Analysis*
Spearman's correlation carried out to measure the strength of the association between the
Luminex™ data for two different mediators (IL-17A and each of the following: GM-CSF, IL-10,
TNF-α, and IL-22) was generated using MATLAB® software (Natick, MA).

3. RESULTS

3.1 Bioinformatic analysis defines networks of interaction among rare SNPs in blunt
trauma patients and implicates Th17 differentiation and systemic hypo-inflammation. We
have previously searched for common SNPs (defined operationally as being present in 20% or
more of our study cohort) that might be both unlikely and associated with dysregulated
inflammation and adverse clinical outcomes in the context of critical illness, and found one such
SNP in the *LYPD4* gene (out of >500,000 assessed SNPs) using a Python-based algorithm we
term *SNPScanner* (24).
In the present study, we sought to study less common SNPs and focused on SNPs present
in 5-10% of the population to allow for a sufficient number of patients in the various comparison
groups. Herein, we refer to these SNPs as "rare" to distinguish them from the "common" *LYPD4*
SNP identified previously (24), acknowledging that most SNPs are found in between 0.5 and 5%
of the global population, and SNPs found in a greater proportion of individuals are unlikely (37).
Our search yielded 12 somatic SNPs associated with two highly statistically significantly
different inflammation biomarkers (based on the initial Mann-Whitney U-test performed in the
SNPScanner algorithm), with a p-value < 0.00001 between subgroups of patients homozygous
for the main allele at a given SNP (referred to as AA) vs. the alternative homozygous allele at
that SNP (referred to as BB). We then carried out Functional Enrichment Analysis using the web
application FLAME (29). When the *SNPScanner* results were entered into FLAME, one of these
SNPs, three major and one minor hub gene were identified.
One major hub gene, with three connections, was identified via the SNP rs4774381 in the
gene encoding RAR-related orphan receptor A (RORA), which is involved in Th17
differentiation (38) (Figure 1). When segregating trauma patients based on AA vs. BB genotype
at this SNP, statistically significant differences were noted in circulating IL-17A along with IFN-
γ, IL-1RA, IL-5, IL-7, IL-10, MIG/CXCL9, sIL-2rα, and sST2 (Figure 2A). This was associated
with statistically significant differences in MODS trajectories (Figure 2B) but not in admission
Shock Index between AA and BB (p=0.799). Paradoxically, rs4774381^{AA} exhibited significantly
(p < 0.001) lower Marshall MODScores as compared to rs4774381^{BB} patients despite the
elevated systemic inflammation in rs4774381^{AA} vs. rs4774381^{BB} patients (Figure 2B). This
finding raised the possibility that pathological hypo-inflammation might be associated with
adverse outcomes, as suggested in our prior work (15). In support of this hypothesis, we note the
early elevation of systemic IL-10, a key anti-inflammatory cytokine, in rs4774381^{BB} patients as
compared to rs4774381^{AA} patients.
We next utilized Dynamic Network Analysis (DyNA) (39) to define networks of
interaction among all systemic inflammatory mediators in rs4774381^{AA} vs. rs4774381^{BB} patients
(Figure 3). This analysis demonstrated striking differences in network patterns between these
two patient subgroups. Specifically, the dynamic networks of systemic inflammation inferred in
rs4774381^{AA} (Figure 3A, 3C) were much sparser at early time points post-hospitalization than
those inferred in s4774381^{BB} patients (Figure 3B, 3C). In addition, both rs4774381^{AA} and
rs4774381^{BB} patients exhibited oscillating systemic inflammation networks over the first seven
days following hospitalization, with the amplitude of oscillation appearing to manifest a "phase
shift" when comparing the two patient subgroups (Figure 3C).
The overall network connectivity difference between homozygous subgroups was also
reflected in the network connectivity of individual inflammatory mediators (Figure 3D). Multiple

inflammatory mediators that had zero connections in networks inferred in rs4774381[AA] patients
exhibited high connectivity in rs4774381[BB] patients; this included IL-17A as well as IL-5, GM-
CSF, sIL-2rα, and IL-6 (Figure 3D). In contrast, Eotaxin and IFN-α exhibited the opposite
pattern, with zero network connections inferred in rs4774381[BB] patients and high connectivity in
rs4774381[AA] patients (Figure 3D). Overall, the most connected mediators were MIP-1β and IL-2,
both inferred in rs4774381[BB] patients (Figure 3D).
Since RORA is a key transcription factor associated with Th17 differentiation (38), we
next carried out a correlation analysis aimed at inferring Th17 cell subsets as in prior studies (13,
15, 19), noting that identification of Th17 cells in trauma patients using this correlation-based
analysis was supported by more direct methods in studies carried out by others (40). This
analysis involves correlation of circulating IL-17A with circulating GM-CSF, IL-10, TNF-α, and
IL-22 to infer the presence of pathogenic Th17 cells (IL-17A vs. GM-CSF), non-pathogenic
Th17 cells (IL-17A vs. IL-10), memory/effector T cells (IL-17A vs. TNF-α), and δγ T cells (IL-
17A vs. IL-22), respectively (Figure 4), and suggested distinct patterns of Th17 subpopulations
based on statistically significant inter-mediator correlations. Specifically, we found a significant
correlation between IL-17A and IL-10 only in rs4774381[AA] patients (compare Figure 4B and
4F), suggesting the predominance of non-pathogenic (non-inflammatory (38)) Th17 cells in these
patients. Both rs4774381[AA] and rs4774381[BB] patients exhibited statistically significant
correlations between IL-17A and TNF-α (compare Figures 4C and 4G) as well as IL-22
(compare Figures 4D and 4H), suggesting similar levels of memory/effector and δγ T cell
responses, respectively. Finally, neither patient subgroup exhibited an upregulation of pathogenic
Th17 cells as inferred from non-significant correlations between IL-17A and GM-CSF (compare
Figures 4A and 4E). This analysis suggests that upregulation of non-pathogenic Th17 cells is
associated with hypo-inflammation following severe injury.
The SNP rs7918717 in the *RPP38* gene, which codes for a subunit of RNAse P (41) was
the second major hub gene identified by FLAME, with four connections. Interestingly, the SNP
rs7918717 in the *NMT2* gene was identified as a minor hub gene with one connection also to
rs7918717. Patients stratified based on genotype (AA vs. BB) at this SNP exhibited significantly
different circulating GM-CSF (p=0.014) as well as MODS trajectories (p=0.024), but not in ICU
LOS, total LOS, days on mechanical ventilation, ISS, and AIS (p>0.05, not shown). There was
no statistically significant difference in Shock Index between AA and BB (p=0.866).
The SNP rs261537 in the *UBE2D2* gene was the third major hub gene identified by
FLAME, with 2 connections. The AA and BB subgroups at this SNP exhibited statistically
significant differences in both circulating IFN-γ (p=0.026) and GM-CSF (p=0.042). However,
we found no significant differences in any of the clinical characteristics analyzed (not shown).
297
3.2 rs11919443 in the *TM4SF19* gene is associated with distinct trajectories of multiple
systemic inflammatory mediators including IL-17A and hypo-inflammation, but not with
distinct Th17 cell subsets. We next hypothesized the trauma patient subgroups segregated based
on non-hub, rare SNPs associated with genotypically distinct systemic inflammation profiles
would also exhibit differences in circulating IL-17A. One such SNP was rs11919443 in the gene
coding for transmembrane 4 L six family member 19 (*TM4SF19*). In support of this hypothesis,
rs11919443[AA] patients exhibited significantly elevated levels of circulating IL-17A as compared
to rs11919443[BB] patients (Figure 5A). In addition, the following circulating inflammatory

mediators were also significantly different between the two groups by Two-Way ANOVA: IL-6
(p=0.005), IL-7 (p=0.007), IL-8 (p=0.001), IL-10 (p=0.017), MCP-1 (p<0.001) and sST2
(p<0.001). These differences were associated with significantly different clinical outcomes,
including longer ICU stay (Figure 5C) and longer time on mechanical ventilation (Figure 5D) in
rs11919443^{AA} patients as compared to rs11919443^{BB} patients. In contrast, there was no
statistically significant difference in Shock Index between AA and BB (p=0.207),
Similarly to the analysis of patients stratified based on genotype at rs4774381, inference
of dynamic networks in rs11919443^{AA} patients as compared to rs11919443^{BB} patients suggested
a striking hypo-inflammatory phenotype, with rs11919443^{BB} patients exhibiting zero
inflammatory network connectivity while rs11919443^{AA} patients exhibited fairly robustly
connected networks (Figure 6A) (Supplementary Figure 2). The most connected mediators (with
10 connections or more) in rs11919443^{AA} patients were, in descending order, IL-5, IL-7, IL-2,
IL-15, IL-1β, GM-CSF, IL-1RA, IL-17A, IFN-γ, and IFN-α (Figure 6B). In agreement with the
less important role of IL-17A in these dynamic networks, and in contrast with the analysis
carried out on patients stratified based on homozygous genotype rs4774381 at the hub gene
RORA, rs11919443 patient subgroups exhibited no major inferred differences in their Th17
subsets (Figure 7).
323
3.3 Analysis of SNP networks in a control group defined algorithmically as having
genotypically associated differences in clinical outcomes but not circulating inflammatory
mediators implicates *SLC2A1*. The 443 rare SNPs with zero statistically significant mediators
were entered into FLAME, which returned 261 SNP-gene permutations. When we added the
requirement that the control SNPs must have 3-4 statistically different post-traumatic outcomes,
the number of rare control SNPs fell to 10 SNPs. Running these SNPs through FLAME resulted
in five genes associated with a given SNP within the set, of which four gave results for
functional enrichment (Supplementary Table 2). The four genes are *SLC2A1* (associated with
rs841852), *IQSEC2* (rs1409117), *ABI3BP* (rs7609779), and *RGS21* (rs7514892). Of these, only
SLC2A1 (solute carrier family 2, facilitated glucose transporter member 1/glucose transporter 1
[GLUT1]), indicated by its SNP rs841852, was a hub gene (Figure 8A-B). Patients segregated
based on their genotype at rs841852 (AA vs. BB) did not differ with regard to injury severity
(ISS as well as AIS) and Shock Index (p=0.066) but did exhibit statistically significant
differences in (Figure 8C-G): ICU length of stay (p< 0.002), total hospital length of stay (p=
0.046), and Marshall MODScore (p< 0.001). Circulating inflammatory mediators in the two
genotypes differed significantly (Two-Way ANOVA) in IL-1RA (p= 0.019), IL-7 (p = 0.032),
IL-13 (p< 0.001), IL-15 (p= 0.006), MIG (p= 0.024), and sIL-2Ra (p= 0.003) (Figure 9).
Similarly to the hypo-inflammation inferred in dynamic networks observed in the other
SNP comparison groups, trauma patients segregated based on alternative homozygous genotype
at rs841852 exhibited overall low inflammatory network connectivity except in rs841852^{AA}
patients at time points from day 4-7 post-hospitalization; rs841852^{BB} patients exhibited zero
network connections at all time points (Figure 10A). Mediators with 10 or more connections in
rs841852^{AA} patients were, in descending order, MIP-1α, IL-4, IL-5, IFN-γ, IL-7, IL-13, IL-17A,
IL-1RA, and IL-33 (Figure 10B; Supplementary Figure 3). Consistent with the hypothesis of
hypo-inflammation in rs841852^{BB} patients was a mild inferred upregulation of non-pathogenic

Th17 cells while other inferred Th17 subsets were generally similar between patient subgroups
(Figure 11).

4. DISCUSSION

The present study extends multiple prior studies linking specific genetic polymorphisms
to outcomes of trauma-associated critical illness (19, 22, 23, 25, 42, 43). Specifically, we
leveraged novel bioinformatics tools and approaches to implicate the Th17-related RAR-related
orphan receptor A (RORA) transcription factor (54) as a major hub gene in networks of SNPs
associated with dysregulated systemic inflammation in blunt trauma patients. Segregating blunt
trauma survivors based on alternative homozygous genotypes at rs11919443, a non-hub, intronic
SNP in the *TM4SF19* gene also identified in this search, yielded subpopulations of patients with
distinct trajectories of systemic inflammatory mediators including IL-17A. Furthermore, these
subpopulations also differed in key critical illness outcomes but not in core demographics or
injury/hemorrhage characteristics, supporting the notion that the differences we identified are
due to genetic causes. Our study also suggested *SLC2A1/GLUT1* as a hub gene in the responses
of trauma patients whose systemic inflammation was not characterized by extensive,
genotypically associated differences while still exhibiting genotypically related differences in
clinical outcomes. Finally, the present study also highlighted the presence and possible
pathophysiologic role of systemic hypo-inflammation, and implicated differential IL-17A
production in this process.
Interleukin-17A is a pro-inflammatory cytokine that is produced by Th17 cells, innate
lymphoid cells, $\gamma\delta$ T cells, and natural killer cells among others (44-50). Interleukin-17A and
Th17 cells were studied initially in the context of chronic inflammatory/autoimmune diseases
such as rheumatoid arthritis, Crohn's disease, psoriasis, and systemic lupus erythematosus (SLE)
(49, 51). More recently, this pathway has been implicated in acute inflammatory states such as
sepsis (52, 53), trauma/hemorrhagic shock (13, 15, 20), COVID-19 (54), wound healing (55),
and acute allograft rejection (56).
Interleukin-17A and Th17 cells have also been implicated in the dysregulated
inflammation that accompanies adverse outcomes in the context of critical illness, based on
initial studies in animal models of sepsis (52, 53) and subsequent studies in human trauma
patients (13, 40). Specifically, mortality in trauma-associated critical illness (13) was associated
with a predominance of so-called pathogenic Th17 cells (57, 58). In these initial studies
documenting an early role for IL-17A in dynamic networks of self-sustaining inflammation, we
inferred an upregulation of pathogenic Th17 cells and a concomitant downregulation of
nonpathogenic Th17 cells (13). Our initial study associating seven novel SNPs with mortality
extended these initial findings and suggested that this genetic predisposition to mortality in the
context of trauma-associated critical illness was not determined solely by these SNPs but also
required a predominant pathogenic Th17 phenotype that was in turn associated with an overall
elevated systemic inflammatory response (19). These results pointed to related, but distinct, roles
for genetic predisposition as well as systemic inflammation characterized by distinct Th17
subsets, and this was confirmed in subsequent studies (22). In the present study, we extend these
results further with the observation that the flip side of pathogenic Th17-driven, self-sustaining
hypo-inflammation is a genetically controlled predisposition to hypo-inflammation characterized
by an upregulation of non-pathogenic Th17 cells. We note the early upregulation of systemic IL-
10 and inferred upregulation of non-pathogenic Th17 cells in rs4774381[BB] patients as compared
to rs4774381[AA] patients. We have suggested a similar phenotype in patients stratified based on a

SNP in the *CD55* gene (25). This hypothesis needs to be tested in larger patient cohorts as well
as via mechanistic perturbations in appropriate experimental models.
A related hypothesis concerns the IL-10 family cytokine IL-22 (59) and the correlation
between IL-17A and IL-22. In this and related studies, we have inferred the presence of γδ T
cells based on a positive correlation between IL-17A and IL-22. However, this supposition
would need to be tested directly by examining cellular profiles and other markers specific for any
of the inferred cell subpopulations, as noted above. It is also possible that the positive correlation
between IL-17A and IL-22 reflects an IL-17A-induced upregulation of IL-22 by other cell types.
In this regard, recent studies have implicated epithelial-derived IL-22 and related tissue-
protective cytokines such as IL-21 and IL-17E/IL-25 in the context of the response to traumatic
injury (60). It is intriguing to speculate that IL-17A, derived from any of the cell populations that
can produce this cytokine, induces IL-22 from epithelial cells in various tissues and that this
process, alone or in concert with upregulation of γδ T cells as well as increases in circulating IL-
10, may contribute to systemic hypo-inflammation in some patients. Given the difficulty in
teasing apart these processes (especially given the need to assess IL-22 in tissues of human
trauma patients), this hypothesis may require studies in animal models or advances in the ability
to detect fluxes in cell populations in humans *in situ* (61).
Another intriguing hypothesis that will need to be tested further is that of the apparent
"phase shift" in an oscillatory systemic inflammatory response inferred via DyNA when
stratifying trauma patients based on homozygous genotype at rs4774381. In unpublished studies,
we have inferred similar dynamic network oscillations when examining the systemic
inflammatory responses of spinal cord injury patients segregated based on the location of their
spinal cord injury (T6 and above vs. below T6; R. Zamora, E. Robbins, W. Huang, Y. Vodovotz,
and G. Sowa, unpublished), which raises the possibility of neural involvement in this
phenomenon. In support of this hypothesis, we have previously documented a role for the
chemokine IP-10/CXCL10 as a driver of IL-10 and systemic hypo-inflammation when
comparing trauma patients with vs. without spinal cord injury (62).
rs11919443 in the *TM4SF19* gene was another key SNP inferred to be related to
dysregulated systemic inflammation (including distinct IL-17A time courses as a function of
genotype at this SNP). Specifically, patients with the homozygous AA genotype of rs11919443
exhibited significantly longer ICU length of stay and requirement for mechanical ventilation
(along with a trend towards longer total hospital length of stay) as well as higher circulating IL-
17A levels as compared to patients with the alternative homozygous (BB) genotype.
Transmembrane 4 L six family member 19 (TM4SF19) is a lysosomal protein that suppresses
acidification; its inactivation was found to block the increase in accumulation of macrophages in
a murine adipose tissue and thus was associated with reduced inflammation (63). *In vitro* studies
suggested that TM4SF19 exacerbates the effects of bacterial lipopolysaccharide on human
venous endothelial cells by impacting the expression of cell junction proteins (64). Thus,
TM4SF19 appears to exert pro-inflammatory effects, albeit indirectly.
One such indirect mechanism appears to involve the chemokine CXCL2, based on a
recent bioinformatics study linking TM4SF19 and CXCL2 in the context of childhood obesity
(65). Interleukin-17A induced CXCL2 among its many actions (66), and bioinformatics analyses
have therefore placed CXCL2 in the IL-17A signaling pathway (67, 68). Furthermore, IL-17A

synergizes with TNF-α to drive endothelial expression of CXCL2 and subsequent neutrophil
recruitment (69).
439   A key aspect of the present study is the focus on a control group of SNPs selected based
on the following criteria: 1) absence of statistically significant differences in circulating
inflammatory mediators between subgroups of trauma patients stratified on the basis of their
homozygous genotype at a given SNP, and 2) presence of statistically significant differences in
clinical outcomes in these same patients. Once SNPs that met these criteria were selected,
FLAME analysis was used to define hub genes. This analysis identified *SLC2A1* as the main hub
gene. SLC2A1 (solute carrier family 2, facilitated glucose transporter member 1/glucose
transporter 1 [GLUT1]) is the main glucose transporter (70), in which there has been extensive
interest due to the emerging role of glucose metabolism in autoimmunity (71, 72), cancer (73,
74), as well as sepsis-induced critical illness (75). Our findings support a role for differential
glucose transport in organ dysfunction in the context of trauma-induced critical illness as well.
450   Our study has several limitations. These include the need to assess more SNPs and other
genetic variants in larger, more diverse patient cohorts; to measure more mediators to better
define the impact of the specific SNPs assessed herein on the immuno-inflammatory response;
and to assess other 'omics layers to better delineate the host's pathophysiological response in the
context of genetic predisposition. Computational modeling – adding mechanistic modeling to
the present study's data-driven modeling approaches (61) – would also serve an important role in
any attempt to connect our findings to the host's pathophysiology following severe traumatic
injury and would also help address the network of interaction among SNPs (76, 77). Finally, our
conclusions are based on associations rather than direct mechanistic work, though our initial
study documenting the role of IL-17A in the injury response included supportive mechanistic
studies involving neutralization of IL-17A in a mouse model of trauma/hemorrhage (13).
461   Despite these limitations, the present study suggests a clear role for nuanced Th17
response in the context of trauma-associated systemic inflammation and critical illness. Multiple
cell types can express IL-17A, including Th17 cells, δγ T cells, and innate lymphoid cells (ILCs)
(44-50). In the context of trauma-associated critical illness or experimental models of
trauma/hemorrhage, IL-17A production is associated with morbidity and mortality (13, 15, 19-
21). While studies have implicated both Th17 cells (13, 40) and ILCs (78) in this
pathophysiology, the present study points to RORA and Th17 differentiation as a key genetically
regulated aspect of this response; future studies will hopefully address the relative contributions
of these distinct cell populations.

REFERENCES
1.   Horiguchi H, Loftus TJ, Hawkins RB, Raymond SL, Stortz JA, Hollen MK, et al. Innate immunity in the Persistent nflammation, mmunosuppression, and Catabolism Syndrome and its implications for therapy. Frontiers in immunology. 2018;9:595.
2.   Lord JM, Midwinter MJ, Chen YF, Belli A, Brohi K, Kovacs EJ, et al. The systemic immune response to trauma: an overview of pathophysiology and treatment. Lancet. 2014;384(9952):1455-65.
3.   Huber-Lang M, Lambris JD, Ward PA. Innate immune responses to trauma. Nature immunology. 2018;19(4):327-41.
4.   Lamparello AJ, Namas RA, Constantine G, McKinley TO, Elster E, Vodovotz Y, et al. A conceptual time window-based model for the early stratification of trauma patients. Journal of internal medicine. 2019.
5.   Hildebrand F, Mommsen P, Frink M, van Griensven M, Krettek C. Genetic predisposition for development of complications in multiple trauma patients. Shock (Augusta, Ga). 2011;35(5):440-8.
6.   Lamparello AJ, Namas RA, Abdul-Malak O, Vodovotz Y, Billiar TR. Young and aged blunt trauma patients display major differences in circulating inflammatory mediator profiles after severe injury. Journal of the American College of Surgeons. 2019;228:148-60.e7.
7.   Zaaqoq AM, Namas RA, Abdul-Malak O, Almahmoud K, Barclay D, Yin J, et al. Diurnal variation in systemic acute inflammation and clinical outcomes following severe blunt trauma. Frontiers in immunology. 2019;10(2699).
8.   Almahmoud K, Namas RA, Zaaqoq AM, Abdul-Malak O, Namas R, Zamora R, et al. Prehospital hypotension is associated with altered inflammation dynamics and worse outcomes following blunt trauma in humans. Critical care medicine. 2015;43:1395-404.
9.   Almahmoud K, Namas RA, Abdul-Malak O, Zaaqoq AM, Zamora R, Zuckerbraun BS, et al. Impact of injury severity on dynamic inflammation networks following blunt trauma. Shock (Augusta, Ga). 2015;44:105-9.
10.   Abdul-Malak O, Vodovotz Y, Zaaqoq A, Guardado J, Almahmoud K, Yin J, et al. Elevated admission base deficit is associated with a complex dynamic network of systemic inflammation which drives clinical trajectories in blunt trauma patients. Mediators of inflammation. 2016;2016:7950374. http://dx.doi.org/10.1155/2016/.
11.   Almahmoud K, Abboud A, Namas RA, Zamora R, Sperry J, Peitzman AB, et al. Computational evidence for an early, amplified systemic inflammation program in polytrauma patients with severe extremity injuries. PloS one. 2019;14(6):e0217577.
12.   Namas RA, Vodovotz Y, Almahmoud K, Abdul-Malak O, Zaaqoq A, Namas R, et al. Temporal patterns of circulating inflammation biomarker networks differentiate susceptibility to nosocomial infection following blunt trauma in humans. Ann Surg. 2016;263:191-8.
13.   Abboud AN, Namas RA, Ramadan M, Mi Q, Almahmoud K, Abdul-Malak O, et al. Computational analysis supports an early, type 17 cell-associated divergence of blunt trauma survival and mortality. Crit Care Med. 2016;44:e1074-e81.
14.   Namas R, Almahmoud K, Mi Q, Ghuma A, Namas R, Zaaqoq A, et al. Individual-specific principal component analysis of circulating inflammatory mediators predicts early organ dysfunction in trauma patients. Journal of critical care. 2016;36:146-53.

15.   Schimunek L, Lindberg H, Cohen M, Namas RA, Mi Q, Yin J, et al. Computational derivation of core, dynamic human blunt trauma inflammatory endotypes. Frontiers in immunology. 2021;11(3481).

16.   Liu D, Namas RA, Vodovotz Y, Peitzman AB, Simmons RL, Yuan H, et al. Unsupervised clustering analysis based on MODS severity identifies four distinct organ dysfunction patterns in severely injured blunt trauma patients. Frontiers in medicine. 2020;7(46).

17.   Wu J, Vodovotz Y, Abdelhamid S, Guyette FX, Yaffe MB, Gruen DS, et al. Multi-omic analysis in injured humans: Patterns align with outcomes and treatment responses. Cell Reports Medicine. 2021;2(12).

18.   Wu J, Cyr A, Gruen DS, Lovelace TC, Benos PV, Das J, et al. Lipidomic signatures align with inflammatory patterns and outcomes in critical illness. Nature communications. 2022;13(1).

19.   Schimunek L, Namas RA, Yin J, Liu D, Barclay D, El-Dehaibi F, et al. An enrichment strategy yields seven novel single nucleotide polymorphisms associated with mortality and altered TH17 responses following blunt trauma. Shock (Augusta, Ga). 2018;49:259-68.

20.   Zamora R, Forsberg JA, Shah AM, Unselt D, Grey S, Lisboa FA, et al. Central role for neurally dysregulated IL-17A in dynamic networks of systemic and local inflammation in combat casualties. Scientific reports. 2023;13(1).

21.   Shah AM, Zamora R, Vodovotz Y. Interleukin-17 as a spatiotemporal bridge from acute to chronic inflammation: Novel insights from computational modeling. WIREs Mechanisms of Disease. 2023;e1599.

22.   Schimunek L, Namas RA, Yin J, Barclay D, Liu D, el-Dehaibi F, et al. MPPED2 polymorphism is associated with altered systemic inflammation and adverse trauma outcomes. Frontiers in genetics. 2019;10(1115).

23.   Lamparello AJ, Namas RA, Schimunek L, Cohen M, El-Dehaibi F, Yin J, et al. An aging-related single-nucleotide polymorphism is associated with altered clinical outcomes and distinct inflammatory profiles in aged blunt trauma patients. Shock (Augusta, Ga). 2020;53:146-55.

24.   El-Dehaibi F, Zamora R, Radder J, Yin J, Shah AM, Namas RA, et al. A common single nucleotide polymorphism is associated with inflammation and critical illness outcomes. iScience. 2023;26(12):108333.

25.   El-Dehaibi F, Zamora R, Yin J, Namas RA, Billiar TR, Vodovotz Y. Network analysis of single-nucleotide polymorphisms associated with aberrant inflammation in trauma patients suggests a role for vesicle-associated inflammatory programs involving CD55. Shock (Augusta, Ga). 2024:10.1097/SHK.0000000000002448.

26.   BAKER SP, O'NEILL B, HADDON WJ, LONG WB. The injury severity score: A method for describing patients with multiple injuries and evaluating emergency care. Journal of Trauma and Acute Care Surgery. 1974;14(3):187-96.

27.   Gennarelli TA, Wodzin E. AIS 2005: a contemporary injury scale. Injury. 2006;37(12):1083-91.

28.   Marshall JC, Cook DJ, Christou NV, Bernard GR, Sprung CL, Sibbald WJ. Multiple organ dysfunction score: a reliable descriptor of a complex clinical outcome. Critical care medicine. 1995;23(10):1638-52.

29.   Karatzas E, Baltoumas FA, Aplakidou E, Kontou PI, Stathopoulos P, Stefanis L, et al.
Flame (v2.0): advanced integration and interpretation of functional enrichment results from
multiple sources. Bioinformatics. 2023;39(8).
30.   Kolberg L, Raudvere U, Kuzmin I, Adler P, Vilo J, Peterson H. g:Profiler-interoperable
web service for functional enrichment analysis and gene identifier mapping (2023 update).
Nucleic Acids Res. 2023;51(W1):W207-W12.
31.   Xie Z, Bailey A, Kuleshov MV, Clarke DJB, Evangelista JE, Jenkins SL, et al. Gene Set
Knowledge Discovery with Enrichr. Curr Protoc. 2021;1(3):e90.
32.   Aleksander S, Balhoff J, Carbon S, Cherry J, Drabkin H, Ebert D, et al. The Gene
Ontology knowledgebase in 2023., 2023, 224, p. iyad031. DOI: https://doi
org/101093/genetics/iyad031 PMID: https://www ncbi nlm nih gov/pubmed/36866529.
33.   Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids
Res. 2000;28(1):27-30.
34.   Ziraldo C, Vodovotz Y, Namas RA, Almahmoud K, Tapias V, Mi Q, et al. Central role
for MCP-1/CCL2 in injury-induced inflammation revealed by in vitro, in silico, and clinical
studies. PLoS One. 2013;8(12):e79804.
35.   Abboud A, Namas RA, Ramadan M, Mi Q, Almahmoud K, Abdul-Malak O, et al.
Computational Analysis Supports an Early, Type 17 Cell-Associated Divergence of Blunt
Trauma Survival and Mortality. Crit Care Med. 2016;44(11):e1074-e81.
36.   Zamora R, Korff S, Mi Q, Barclay D, Schimunek L, Zucca R, et al. A computational
analysis of dynamic, multi-organ inflammatory crosstalk induced by endotoxin in mice. PLoS
Comput Biol. 2018;14(11):e1006582.
37.   Auton A, Abecasis GR, Altshuler DM, Durbin RM, Abecasis GR, Bentley DR, et al. A
global reference for human genetic variation. Nature. 2015;526(7571):68-74.
38.   Korn T, Bettelli E, Oukka M, Kuchroo VK. IL-17 and Th17 Cells. Annu Rev Immunol.
2009;27:485-517.
39.   Mi Q, Constantine G, Ziraldo C, Solovyev A, Torres A, Namas R, et al. A dynamic view
of trauma/hemorrhage-induced inflammation in mice: Principal drivers and networks. PloS one.
2011;6:e19424.
40.   Seshadri A, Brat GA, Yorkgitis BK, Keegan J, Dolan J, Salim A, et al. Phenotyping the
immune response to trauma: A multiparametric systems immunology approach. Critical care
medicine. 2017;45:1523-30.
41.   Kovrigina E, Wesolowski D, Altman S. Coordinate inhibition of expression of several
genes for protein subunits of human nuclear RNase P. Proceedings of the National Academy of
Sciences of the United States of America. 2003;100(4):1598-602.
42.   Brown D, Namas RA, Almahmoud K, Zaaqoq A, Sarkar J, Barclay DA, et al. Trauma *in*
silico: individual-specific mathematical models and virtual clinical populations. Science
translational medicine. 2015;7:285ra61.
43.   Sperry JLZ, S.; Zuckerbraun, B.S.; Vodovotz, Y.; Namas, R.; Ferrell, R.E.; Rosengart,
M.R.; Peitzman, A.B.; Billiar, T.R. . X-chromosome linked IRAK1 polymorphism is a strong
predictor of multiple organ failure and mortality post-injury. Ann Surg. 2014;260:698-703.
44.   Lillehoj HS, Lillehoj EP, Weinstock D, Schat KA. Functional and biochemical
characterizations of avian T lymphocyte antigens identified by monoclonal antibodies. European
Journal of Immunology. 1988;18(12):2059-65.

45.   Park H, Li Z, Yang XO, Chang SH, Nurieva R, Wang Y-H, et al. A distinct lineage of
CD4 T cells regulates tissue inflammation by producing interleukin 17. Nature Immunology.
2005;6(11):1133-41.
46.   Chan MM, Chen CL, Ager LL, Cooper MD. Identification of the avian homologues of
mammalian CD4 and CD8 antigens. J Immunol. 1988;140(7):2133-8.
47.   Harrington LE, Hatton RD, Mangan PR, Turner H, Murphy TL, Murphy KM, et al.
Interleukin 17–producing CD4+ effector T cells develop via a lineage distinct from the T helper
type 1 and 2 lineages. Nature Immunology. 2005;6(11):1123-32.
48.   McGeachy MJ, Cua DJ, Gaffen SL. The IL-17 Family of Cytokines in Health and
Disease. Immunity. 2019;50(4):892-906.
49.   Skopelja-Gardner S, Tai J, Sun X, Tanaka L, Kuchenbecker JA, Snyder JM, et al. Acute
skin exposure to ultraviolet light triggers neutrophil-mediated kidney inflammation. Proc Natl
Acad Sci U S A. 2021;118(3).
50.   Cua DJ, Tato CM. Innate IL-17-producing cells: the sentinels of the immune system. Nat
Rev Immunol. 2010;10(7):479-89.
51.   Beringer A, Noack M, Miossec P. IL-17 in Chronic Inflammation: From Discovery to
Targeting. Trends in Molecular Medicine. 2016;22(3):230-41.
52.   Flierl MA, Rittirsch D, Gao H, Hoesel LM, Nadeau BA, Day DE, et al. Adverse
functions of IL-17A in experimental sepsis. Faseb j. 2008;22(7):2198-205.
53.   Bosmann M, Ward PA. Therapeutic potential of targeting IL-17 and IL-23 in sepsis. Clin
Transl Med. 2012;1(1):4.
54.   Xu Z, Shi L, Wang Y, Zhang J, Huang L, Zhang C, et al. Pathological findings of
COVID-19 associated with acute respiratory distress syndrome. Lancet Respir Med.
2020;8(4):420-2.
55.   Konieczny P, Xing Y, Sidhu I, Subudhi I, Mansfield KP, Hsieh B, et al. Interleukin-17
governs hypoxic adaptation of injured epithelium. Science. 2022;377(6602):eabg9302.
56.   Shah AM, Aral AM, Zamora R, Gharpure N, El-Dehaibi F, Zor F, et al. Peripheral nerve
repair is associated with augmented cross-tissue inflammation following vascularized composite
allotransplantation. Frontiers in immunology. 2023;14.
57.   Bettelli E, Carrier Y, Gao W, Korn T, Strom TB, Oukka M, et al. Reciprocal
developmental pathways for the generation of pathogenic effector TH17 and regulatory T cells.
Nature. 2006;441(7090):235-8.
58.   Lee Y, Awasthi A, Yosef N, Quintana FJ, Xiao S, Peters A, et al. Induction and
molecular signature of pathogenic TH17 cells. Nature immunology. 2012;13(10):991-9.
59.   Ouyang W, O'Garra A. IL-10 family cytokines IL-10 and IL-22: from basic science to
clinical translation. Immunity. 2019;50(4):871-91.
60.   Cai J, McKinley T, Billiar I, Zenati MS, Gaski G, Vodovotz Y, et al.
Protective/reparative cytokines are suppressed at high injury severity in human trauma. Trauma
surgery & acute care open. 2021;6(1):e000619.
61.   Vodovotz Y. Towards systems immunology of critical illness at scale: from single cell
'omics to digital twins. Trends Immunol. 2023;44:345-55.
62.   Zaaqoq AM, Namas R, Almahmoud K, Azhar N, Mi Q, Zamora R, et al. Inducible
protein-10, a potential driver of neurally-controlled IL-10 and morbidity in human blunt trauma.
CritCare Med. 2014;42:1487-97.

63.   Choi C, Jeong YL, Park KM, Kim M, Kim S, Jo H, et al. TM4SF19-mediated control of
lysosomal activity in macrophages contributes to obesity-induced inflammation and metabolic
dysfunction. Nature communications. 2024;15(1):2779.
64.   Ding L, Li LM, Hu B, Wang JL, Lu YB, Zhang RY, et al. TM4SF19 aggravates LPS-
induced attenuation of vascular endothelial cell adherens junctions by suppressing VE-cadherin
expression. Biochemical and biophysical research communications. 2020;533(4):1204-11.
65.   He E, Chang K, Dong L, Jia M, Sun W, Cui H. Identification and validation of CXCL2
as a key gene for childhood obesity. Biochem Genet. 2024;62(4):2743-65.
66.   Onishi RM, Gaffen SL. Interleukin-17 and its target genes: mechanisms of interleukin-17
function in disease. Immunology. 2010;129(3):311-21.
67.   Cao M, Wang Y. Bioinformatic analyses determine the importance of CXCL2 and
CXCL8 in atypical meningioma development and reoccurrence. Turk Neurosurg.
2022;32(3):430-41.
68.   Sun Y, Cai D, Hu W, Fang T. Identifying hub genes and miRNAs in Crohn's disease by
bioinformatics analysis. Frontiers in genetics. 2022;13:950136.
69.   Griffin GK, Newton G, Tarrio ML, Bu DX, Maganto-Garcia E, Azcutia V, et al. IL-17
and TNF-α sustain neutrophil recruitment during inflammation through synergistic effects on
endothelial activation. Journal of immunology (Baltimore, Md : 1950). 2012;188(12):6287-99.
70.   Brown GK. Glucose transporters: structure, function and consequences of deficiency. J
Inherit Metab Dis. 2000;23(3):237-46.
71.   Palmer CS, Hussain T, Duette G, Weller TJ, Ostrowski M, Sada-Ovalle I, et al.
Regulators of glucose metabolism in CD4(+) and CD8(+) T cells. Int Rev Immunol.
2016;35(6):477-88.
72.   Di Dedda C, Vignali D, Piemonti L, Monti P. Pharmacological targeting of GLUT1 to
control autoreactive T cell responses. International journal of molecular sciences. 2019;20(19).
73.   Gonzalez-Menendez P, Hevia D, Mayo JC, Sainz RM. The dark side of glucose
transporters in prostate cancer: Are they a new feature to characterize carcinomas? Int J Cancer.
2018;142(12):2414-24.
74.   Alves AP, Mamede AC, Alves MG, Oliveira PF, Rocha SM, Botelho MF, et al.
Glycolysis inhibition as a strategy for hepatocellular carcinoma treatment? Curr Cancer Drug
Targets. 2019;19(1):26-40.
75.   Deane AM, Rayner CK, Keeshan A, Cvijanovic N, Marino Z, Nguyen NQ, et al. The
effects of critical illness on intestinal glucose sensing, transporters, and absorption. Critical care
medicine. 2014;42(1):57-65.
76.   Pairo-Castineira E, Rawlik K, Bretherick AD, Qi T, Wu Y, Nassiri I, et al. GWAS and
meta-analysis identifies 49 genetic variants underlying critical COVID-19. Nature.
2023;617(7962):764-8.
77.   Yang D, Jin Y, He X, Dong A, Wang J, Wu R. Inferring multilayer interactome networks
shaping phenotypic plasticity and evolution. Nature communications. 2021;12(1):5304.
78.   Manson J, Hoffman R, Chen S, Ramadan MH, Billiar TR. Innate-like lymphocytes are
immediate participants in the hyper-acute immune response to trauma and hemorrhagic shock.
Frontiers in immunology. 2019;10:1501.

79.   El-Dehaibi F, Zamora R, Yin J, Billiar TR, Vodovotz Y. medRxiv. 2025.01.28.25321184
[Preprint]. Available at https://www.medrxiv.org/content/10.1101/2025.01.28.25321184v2
(Accessed June 13, 2025).

ABBREVIATIONS

ABI3BP: ABI family member 3 binding protein
AIS: Abbreviated Injury Scale
ANOVA: Analysis of Variance
DyNA: Dynamic Network Analysis
GLUT1: glucose transporter 1
GM-CSF: granulocyte-macrophage colony stimulating factor
GO: Gene Ontology Knowledgebase
ICU: intensive care unit
IFN: interferon
IL: Interleukin
ILC: innate lymphoid cell
IP-10: IFN-γ inducible protein (CXCL10)
IL: interleukin
IQSEC2: IQ motif and Sec7 domain ArfGEF 2
ISS: Injury Severity Score
KEGG: Kyoto Encyclopedia of Genes and Genomics
LOS: length of stay
LYPD4: LY6/PLAUR domain containing 4
MCP: monocyte chemotactic protein
MIG: monokine induced by gamma interferon (CXCL9)
MIP: macrophage inflammatory protein
MODS: Multiple Organ Dysfunction Syndrome
NMT2: N-myristoyltransferase 2
RGS21: regulator of G protein signaling 21
RPP38: ribonuclease P/MRP subunit p38
RORA: RAR-related orphan receptor A
SLC2A1: solute carrier family 2, facilitated glucose transporter member 1
SNP: single nucleotide polymorphism
sST2: soluble suppression of tumorigenicity-2
TM4SF19: transmembrane 4 L six family member 19
TNF-α: tumor necrosis factor α
UBE2D2: ubiquitin conjugating enzyme E2 D2

FUNDING
This work was supported by Department of Defense grant W81XWH-18-2-0051 and Defense Advanced Research Projects Agency grant D20AC00002.

ACKNOWLEDGMENTS
This work was made possible by the Department of Defense and the Defense Advanced Research Projects Agency. The manuscript was first published as a preprint to medRxiv.org.

AUTHOR CONTRIBUTIONS
FE: Writing, software development, data analysis, investigation, visualization. RZ: Writing, data analysis, visualization. JY: Inflammatory mediator arrays, genomic analysis. TB: Curation of patient data, review and editing. YZ: Conceptualization, writing, review and editing, project administration, funding acquisition. This manuscript has been read and approved by all authors.

CONFLICT OF INTEREST
The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

FIGURE CAPTIONS
Figure 1. FLAME tree plot describing molecular function connectivity of 'rare' SNPs
implicated in inflammatory biomolecules disparity by *SNPScanner*. These results were found
in the Gene Ontology knowledgebase with the web tool *enrichR*. SNP rs4774381 corresponds to
the gene for RORA, which is a component of the mTORc1 complex in a Th17 differentiation
pathway.
745
746 **Figure 2. Distinct trajectories of circulating inflammatory mediator mediators as a function**
747 **of homozygous genotype at rs4774381. (A)** Statistically distinct (*p < 0.05 by Two-Way
748 ANOVA) circulating inflammatory biomarker concentrations from 4 hours to 7 days past
749 admission between patients with rs4774381^{AA} and rs4774381^{BB}. **(B)** Marshall MODS seven-day
750 time course between rs4774381^{AA} and rs4774381^{BB} patients (*p < 0.001 by Two-Way ANOVA).
751
Figure 3. Dynamic network analysis (DyNA) suggests distinct inflammatory programs as a
function of homozygous genotype at rs4774381. DyNA results for inflammatory biomolecules
from 4h-D7 of rs4774381^{AA} patients **(A)** and rs4774381^{BB} patients **(B)**. A representative DyNA
output showing all inflammatory mediators in the networks is also shown to easily identify the
mediators. **(C)** Graph of dynamic network complexity over the 4h-D7 time course between
rs4774381^{AA} and rs4774381^{BB}. **(D)** Table of the total number of connections for each
biomolecule for both rs4774381 genotypes.
759
760 **Figure 4.** Spearman's correlations of IL-17A and Th17 differentiation-associated biomolecules:
761 GM-CSF **(A and E)**, IL-10 **(B and F)**, TNF-α **(C and G)** and IL-22 **(D and H)**. Panels **(A-D)** are
762 correlations performed for rs4774381^{AA} and panels **(E-H)** are those for rs4774381^{BB}.
763
Figure 5. Distinct IL-17A trajectories and clinical outcomes as a function of homozygous
genotype at rs11919443. (A-C) Post-admission comparison of systemic inflammation and
clinical outcomes between rs11919443^{AA} and rs11919443^{BB}. **(A)** rs11919443^{AA} patients
exhibited a significant spike in IL-17A concentration from 20 hours to 2 days post-injury
compared to rs11919443^{BB} patients. **(B-C)** Patients with rs11919443^{AA} had significantly longer
ICU stays and required more days on mechanical ventilation than rs11919443^{BB}.
770
771 **Figure 6. Dynamic network analysis suggests distinct inflammatory programs and the**
772 **presence of hypo-inflammation as a function of homozygous genotype at rs11919443. (A)**
773 Graph of dynamic network complexity over the 4h-D7 time course between rs11919443 ^{AA} and
774 rs11919443 ^{BB}. **(B)** Table of the total number of connections for each biomolecule for both
775 rs11919443 genotypes.
776
777 **Figure 7. Correlation-based inference of Th17 cell subsets as a function of homozygous**
778 **genotype at rs11919443.** Spearman's correlations for rs11919443^{AA} **(A-D)** and rs11919443^{BB}
779 **(E-H)** of IL-17A and Th17 differentiation-associated biomolecules: GM-CSF **(A and E)**, IL-10
780 **(B and F)**, TNF-α **(C and G)**, and IL-22 **(D and H)**.
781

Figure 8. (A-B) FLAME tree plot describing molecular function connectivity of 'rare' SNPs
with no significant inflammatory biomarker disparity, according to *SNPScanner*. These
results were found in the Gene Ontology knowledgebase with the web tools *gProfiler* **(A)** and
enrichR **(B)**. Clinical parameter analysis by two-way ANOVA failed to reveal differences in ISS
(C) and AIS **(D)** between rs841852^{AA} and rs841852^{BB}, but implicated significant differences ($p <$
0.05) for days in the ICU **(E)**, total hospital days **(F)**, and Marshall MODS **(G)**.

Figure 9. Distinct trajectories of circulating inflammatory mediator mediators as a function
of homozygous genotype at rs841852. Statistically significantly different ($p < 0.05$ by two-way
ANOVA) biomarkers: IL-1RA **(A)**, IL-7 **(B)**, IL-13 **(C)**, IL-15 **(D)**, MIG **(E)**, and sIL-2Ra **(F)**.

Figure 10. Dynamic network analysis suggests distinct inflammatory programs and the
presence of hypo-inflammation as a function of homozygous genotype at rs841852. (A)
Network complexity of inflammatory biomolecules from 4h-D7 for rs841852 homozygous
genotypes. **(B)** Table of total network connections for all inflammatory biomolecules for both
genotypes.

Figure 11. Spearman's correlations for rs841852 ^{AA} **(A-D)** and rs841852 ^{BB} **(E-H)** of IL-17A and
Th17 differentiation-associated biomolecules: GM-CSF **(A and E)**, IL-10 **(B and F)**, TNF-α **(C**
and G), and IL-22 **(D and H)**.

Figure 1

Genotype-specific dysregulated inflammation ('rare SNPs')

802
803

Figure 2

804
805

Figure 2

B

807
808

Figure 3

A

AA

809
810

Figure 3

B

BB

811
812

Figure 3

C

D

Total Connectivity	AA 0.95	BB 0.95
MIP-1b	7	47
IL-2	16	45
IL-1b	15	34
IL-15	17	33
IL-5	0	31
GM-CSF	0	29
IL-17A	0	25
TNFa	6	24
IL-4	19	20
IL-7	0	16
IL-13	20	15
IL-1RA	0	14
IFNg	4	8
sIL-2ra	0	7
IL-6	0	6
MIP-1a	0	3
IL-33	0	2
Eotaxin	14	0
IFNa	24	0
IL-8	0	1
IL-10	0	1
IL-17E/IL-25	0	1
IL-22	0	1
MCP-1	0	1
IL-9	0	0
IL-21	0	0
IL-23	0	0
IP-10	0	0
MIG	0	0
NO2-/NO3-	0	0
ST2	0	0

813
814

Figure 5

817
818

Figure 6

819
820

Figure 7

821
822
823

Figure 8

A gProfiler

SLC2A1
(rs841852)

GO:1990350
glucose transporter complex

B enrichR

GO:0015149
hexose transmembrane transporter activity

SLC2A1
(rs841852)

GO:0005355
glucose transmembrane transporter activity

GO:0005085
guanyl-nucleotide exchange factor activity

IQSEC2
(rs1409117)

824
825

Figure 8

826
827

Figure 9

828
829

Figure 10

A

B

Total Connectivity	AA 0.95	BB 0.95
MIP-1a	20	0
IL-4	15	0
IL-5	15	0
IFNg	12	0
IL-7	11	0
IL-13	11	0
IL-17A	11	0
IL-1RA	10	0
IL-33	10	0
IL-1b	9	0
IL-17E/IL-25	9	0
IL-21	9	0
sIL-2ra	9	0
GM-CSF	8	0
IL-22	7	0
IL-23	7	0
IL-2	6	0
IL-10	2	0
MIP-1b	1	0
Eotaxin	0	0
IFNa	0	0
IL-6	0	0
IL-8	0	0
IL-9	0	0
IL-15	0	0
IP-10	0	0
MCP-1	0	0
MIG	0	0
NO2-/NO3-	0	0
ST2	0	0
TNFa	0	0

Figure 11

831
832

833   **TABLES**
834
835   Table 1. 'Rare' SNPs entered into FLAME, their number of statistically significant inflammatory
836   biomolecules found by *SNPScanner*, and their corresponding gene/entry found by FLAME (if
837   applicable).

'Rare' Inflammatory Disparity SNPs		
SNP	Number of Distinct Biomolecules ($p < 0.00001$)	Gene/Entry
rs10489239	2	ENSG00000231424
rs6603352	2	KLHL13
rs4774381	2	RORA
rs11782445	2	---
rs7782688	2	---
rs1010779	2	---
rs6603357	2	KLHL13
rs12009735	2	KLHL13
rs6726331	2	LINC01807
rs7258387	2	ENSG00000269110
rs261537	2	UBE2D2
rs13007274	2	FSHR, ENSG00000282890
rs13253492	2	CSMD1
rs7918717	2	RPP38, NMT2
rs11919443	2	TM4SF19, TM4SF19-AS1, TM4SF19-DYNLT2B
'Rare' Control SNPs		
SNP	Number of Distinct Biomolecules ($p < 0.05$)	Gene/Entry
rs1409117	0	IQSEC2
rs2388252	0	ENSG00000235281
rs12450000	0	---
rs4144013	0	PRRG1, ENSG00000250349
rs6609469	0	---
rs9558839	0	---
rs7609779	0	ABI3BP
rs767478	0	---
rs841852	0	SLC2A1
rs7514892	0	RGS21, ENSG00000285280

838
839
840   Table 2. Clinical summary of rs4774381 patients by genotype.

	rs4774381 Genotypes		
	AA	AB	BB
Patient Count	223	136	21
Age	50.55 ± 19.08 (min: 18; max: 90)	46.26 ± 18.70 (min: 18; max: 87)	54.90 ± 20.39 (min: 21; max: 85)
Sex			
Male (%)	154 (69.06%)	91 (66.91%)	16 (76.19%)
Female (%)	69 (30.94%)	45 (33.09%)	5 (23.81%)
Ethnicity			
White (%)	220 (98.65%)	130 (95.59%)	18 (85.71%)
African American (%)	1 (0.45%)	6 (4.41%)	3 (14.29%)
Asian (%)	2 (0.90%)	0	0
ISS	19.65 ± 10.32 (min: 1; max: 54)	18.26 ± 10.15 (min: 1; max: 50)	17.38 ± 7.38 (min: 5; max: 30)
ICU Length of Stay (days)	6.51 ± 7.03 (min: 1; max: 36)	5.64 ± 7.10 (min: 1; max: 36)	7.62 ± 8.81 (min: 1; max: 35)
Total Hospital Length of Stay (days)	11.86 ± 9.37 (min: 1; max: 48)	11.44 ± 9.73 (min: 2; max: 50)	13.71 ± 8.11 (min: 2; max: 35)
Days on Mechanical Ventilation	3.26 ± 6.17 (min: 0; max: 34)	2.50 ± 5.74 (min: 0; max: 37)	3.90 ± 6.39 (min: 0; max: 26)
Shock Index*	0.85 ± 0.70 (min: 0.07; max: 8.67)	0.84 ± 0.85 (min: 0.29; max: 9.00)	0.75 ± 0.24 (min: 0.26; max: 1.42)

842   Table 3. Clinical summary of rs11919443 patients by genotype.

	rs11919443 Genotypes		
	AA	AB	BB
Patient Count	20	111	249
Age	57.95 ± 21.10 (min: 23; max: 90)	46.04 ± 17.81 (min: 18; max: 87)	49.99 ± 19.31 (min: 18; max: 89)
Sex			
Male (%)	14 (70%)	88 (79.28%)	159 (63.86%)
Female (%)	6 (30%)	23 (20.72%)	90 (36.14%)
Ethnicity			
White (%)	18 (90%)	108 (97.30%)	242 (97.19%)
African American (%)	1 (5%)	2 (1.80%)	7 (2.81%)
Asian (%)	1 (5%)	1 (0.90%)	0

ISS	18.35 ± 10.23 (min: 5; max: 48)	20.06 ± 10.91 (min: 1; max: 54)	18.62 ± 9.75 (min: 1; max: 50)
ICU Length of Stay (days)	9.85 ± 9.02 (min: 1; max: 33)	6.57 ± 7.91 (min: 1; max: 36)	5.83 ± 6.56 (min: 1; max: 36)
Total Hospital Length of Stay (days)	16.40 ± 13.04 (min: 2; max: 48)	12.63 ± 9.54 (min: 1; max: 45)	11.08 ± 8.92 (min: 2; max: 50)
Days on Mechanical Ventilation	4.40 ± 4.47 (min: 0; max: 14)	3.50 ± 7.54 (min: 0; max: 37)	2.70 ± 5.31 (min: 0; max: 28)
Shock Index*	0.68 ± 0.15 (min: 0.49; max: 1.04)	0.87 ± 0.81 (min: 0.07; max: 8.67)	0.84 ± 0.74 (min: 0.30; max: 9.00)

844   Table 4. Clinical summary of rs7918717 patients by genotype.

	rs7918717 Genotypes		
	AA	AB	BB
Patient Count	242	117	21
Age	49.78 ± 19.15 (min: 18; max: 90)	46.76 ± 18.79 (min: 18; max: 86)	57.10 ± 19.06 (min: 20; max: 83)
Sex			
Male (%)	168 (69.42%)	78 (66.67%)	15 (71.43%)
Female (%)	74 (30.58%)	29 (24.79%)	6 (28.57%)
Ethnicity			
White (%)	239 (98.76%)	114 (97.44%)	15 (71.43%)
African American (%)	1 (0.41%)	3 (2.56%)	6 (28.57%)
Asian (%)	2 (0.82%)	0	0
ISS	18.70 ± 10.00 (min: 1; max: 50)	19.40 ± 10.35 (min: 1; max: 54)	20.71 ± 10.48 (min: 4; max: 50)
ICU Length of Stay (days)	5.73 ± 6.69 (min: 1; max: 33)	7.00 ± 7.72 (min: 1; max: 36)	8.19 ± 8.70 (min: 1; max: 35)
Total Hospital Length of Stay (days)	11.15 ± 8.84 (min: 1; max: 50)	13.30 ± 10.67 (min: 2; max: 46)	11.19 ± 7.95 (min: 2; max: 35)
Days on Mechanical Ventilation	2.59 ± 5.28 (min: 0; max: 27)	3.74 ± 7.41 (min: 0; max: 37)	3.95 ± 5.40 (min: 0; max: 17)
Shock Index*	0.82 ± 0.75 (min: 0.07; max: 9.00)	0.89 ± 0.79 (min: 0.38; max: 8.67)	0.73 ± 0.22 (min: 0.41; max: 1.22)

846   Table 5. Clinical summary of rs261537 patients by genotype.

	rs261537 Genotypes		
	AA	AB	BB

Patient Count	223	136	21
Age	49.47 ± 19.19 (min: 18; max: 86)	48.20 ± 19.11 (min: 18; max: 90)	53.86 ± 18.72 (min: 23; max: 85)
Sex			
Male (%)	154 (69.06%)	91 (66.91%)	16 (76.19%)
Female (%)	69 (30.94%)	45 (33.09%)	5 (23.81%)
Ethnicity			
White (%)	219 (98.2%)	130 (95.59%)	19 (90.48%)
African American (%)	2 (0.90%)	6 (4.41%)	2 (9.52%)
Asian (%)	2 (0.90%)	0	0
ISS	18.53 ± 10.23 (min: 1; max: 54)	20.15 ± 10.20 (min: 4; max: 50)	17.86 ± 8.28 (min: 5; max: 42)
ICU Length of Stay (days)	6.24 ± 7.18 (min: 1; max: 36)	6.62 ± 7.41 (min: 1; max: 35)	4.14 ± 5.05 (min: 1; max: 24)
Total Hospital Length of Stay (days)	11.97 ± 10.00 (min: 2; max: 50)	11.76 ± 8.79 (min: 1; max: 45)	10.48 ± 7.15 (min: 2; max: 33)
Days on Mechanical Ventilation	2.91 ± 5.80 (min: 0; max: 34)	3.43 ± 6.66 (min: 0; max: 37)	1.67 ± 3.68 (min: 0; max: 17)
Shock Index*	0.86 ± 0.70 (min: 0.30; max: 9.00)	0.83 ± 0.85 (min: 0.07; max: 8.67)	0.68 ± 0.19 (min: 0.30; max: 0.99)

848   Table 6. Clinical summary of rs841852 patients by genotype. * Five patients had no recorded
849   Shock Index value, and four outliers were observed in the available patients' data.

	rs841852 Genotypes		
	AA	AB	BB
Patient Count	21	109	250
Age	57.90 ± 15.83 (min: 23; max: 85)	49.59 ± 19.25 (min: 18; max: 87)	48.38 ± 19.22 (min: 18; max: 90)
Sex			
Male (%)	12 (57.14%)	80 (73.39%)	169 (67.60%)
Female (%)	9 (42.86%)	29 (26.61%)	81 (32.40%)
Ethnicity			
White (%)	19 (90.48%)	104 (95.41%)	245 (98.00%)
African American (%)	2 (9.52%)	5 (4.59%)	3 (1.20%)
Asian (%)	0	0	2 (0.80%)
ISS	15.86 ± 7.83 (min: 1; max: 30)	17.93 ± 10.23 (min: 1; max: 50)	19.77 ± 10.19 (min: 1; max: 54)

ICU Length of Stay (days)	3.86 ± 7.19 (min: 1; max: 35)	7.12 ± 7.86 (min: 1; max: 33)	6.08 ± 6.81 (min: 1; max: 36)
Total Hospital Length of Stay (days)	8.43 ± 7.20 (min: 2; max: 35)	12.96 ± 10.24 (min: 2; max: 48)	11.60 ± 9.17 (min: 1; max: 50)
Days on Mechanical Ventilation	1.00 ± 2.58 (min: 0; max: 12)	3.58 ± 6.31 (min: 0; max: 31)	2.95 ± 6.10 (min: 0; max: 37)
Shock Index*	1.02 ± 1.80 (min: 0.07; max: 9.00)	0.88 ± 0.94 (min: 0.42; max: 8.67)	0.81 ± 0.42 (min: 0.26; max: 5.4)

850