Asymmetric limits on timely interventions from noisy epidemic data

Kris V. Parag[1,2,*], Ben Lambert[3,4], Christl A. Donnelly[3,4] and Sandor Beregi[1]

[1]MRC Centre for Global Infectious Disease Analysis, Imperial College London, London, UK.

[2]NIHR HPRU in Behavioural Science and Evaluation, University of Bristol, Bristol, UK.

[3]Department of Statistics, University of Oxford, Oxford, UK.

[4]Pandemic Sciences Institute, University of Oxford, Oxford, UK.

*For correspondence: k.parag@imperial.ac.uk.

Abstract

Deciding on when to initiate or relax an intervention in response to an emerging infectious disease is both difficult and important. Uncertainties from noise in epidemiological surveillance data must be hedged against the potentially unknown and variable costs of false alarms and delayed actions. Here we clarify and quantify how case under-reporting and latencies in case ascertainment, which are predominant surveillance noise sources, can restrict the timeliness of decision-making. Decisions are modelled as binary choices between responding or not that are informed by reported case curves or transmissibility estimates from those curves. Optimal responses are triggered by thresholds on case numbers or estimate confidence levels, with thresholds set by the costs of the various choices. We show that, for growing epidemics, both noise sources induce additive delays on hitting any case-based thresholds and multiplicative reductions in our confidence in estimated reproduction numbers or growth rates. However, for declining epidemics, these noise sources have counteracting effects on case data and limited cumulative impact on transmissibility estimates. We find this asymmetry persists even if more sophisticated feedback control algorithms that consider the longer-term effects of interventions are employed. Standard surveillance data therefore provide substantially weaker support for deciding when to initiate a control action or intervention than for determining when to relax it. This information bottleneck during epidemic growth may justify proactive intervention choices.

Keywords: infectious diseases; decision theory; epidemic growth; surveillance noise; non-pharmaceutical interventions; reproduction numbers.

Introduction

Rapidly and reliably detecting the key dynamics of infectious diseases is a recurrent challenge in epidemiology with critical consequences and trade-offs [1–3]. Early warnings of upcoming epidemic growth or decline can provide valuable signals for mobilising or relaxing interventions [4–6] and even enhance the effectiveness of decisions to act [7]. However, epidemic data are often scarce or uncertain in early periods [8,9] and the economic and other costs of premature action (i.e., false alarms) may be detrimental. Choosing to wait for data to accumulate and

improve the evidence base for decision-making can also be costly as resulting delays to intervene (i.e., missed actions) may mean lost opportunities to minimise epidemic burden or relax interventions when infections are, respectively, rising or waning [10–12].

Although the decision to act or not at any time commonly involves many external factors (e.g., political and sociocultural precedents) [13,14], at the core of these choices is the complex but fundamental trade-off between uncertainty in epidemic dynamics and uncertainty in the costs of deciding to intervene (and hence modify those dynamics). While integrated epidemiological and economic modelling frameworks [15–18] have been proposed to balance uncertainties and optimise decision-making, these approaches require knowledge or at least assumptions about these uncertainties, which can be difficult to quantify accurately. Even if this uncertainty is well-described, recommended actions may be sensitive to parameter, model structure and implementation choices [7,19]. The complexity of integrated models may further constrain the ability to validate the robustness of outputs, precluding generalised insights [9,20,21].

Given these challenges, a complementary set of studies have instead aimed to use simplified models, which are easier to parametrise and validate, to uncover the fundamental limits that uncertainties impose on public health decision-making [22]. While these studies have yielded valuable insights into the robustness and timeliness of intervention choices and into how we can optimally detect shifts in epidemic dynamics [6,7,23–25], few works have directly coupled these to costs of missed actions and false alarms (though they do examine proxies e.g., the times over which interventions are sustained) or characterised how noise in surveillance data impacts detection and hence decision timepoints, which then feed back onto those costs. Here we aim to resolve these gaps by adapting optimal Bayesian detection theory [26,27], with the goal of extracting general guidelines about how noise constrains cost-optimal action points.

We show that the costs of missed actions and false alarms, even when not known accurately, imply a threshold criterion that we must compare with our evidence to act. We then describe how the information supporting this decision to act accumulates above this threshold with time. The costs of erroneous actions are therefore directly balanced by the aggregate of information that supports action i.e., larger costs require more evidence and longer wait-times to act are justifiable [26]. For growing and waning epidemics, this evidence, respectively, informs the initiating and relaxing of interventions. Our central contribution is to reformulate this decision-cost framework to understand how under-reporting of infections and delays in reporting modify the optimal thresholds for supporting action. Interestingly, we find these ubiquitous sources of noise intrinsically but asymmetrically limit our capacity to respond to epidemics.

When an epidemic is growing, delays and under-reporting introduce additive lags on achieving any threshold based on the incidence of infections or related curves such as those of cases and deaths, and multiplicative reductions in information available for estimating transmissibility i.e., uncertainties around estimates of reproduction numbers and growth rates are amplified by both noise sources. However, when an epidemic is waning, the noise sources counteract, with reporting delays offsetting the impact of under-reporting when hitting thresholds and both having minimal effect on transmissibility estimate uncertainties. This asymmetry results from the directionality of delays and under-reporting, is practically magnified because data are often scarcer during early epidemic growth stages [23] and persists even if sophisticated decision-algorithms [28] that optimise intervention timings and their longer-term impact are applied. We argue that markedly more restrictive bottlenecks to optimal decision-making during emergent epidemics fundamentally support proactive outbreak intervention choices [9].

Results

Detection thresholds and the costs of intervening

We start by reworking classical Bayesian decision theory to better understand what a decision may involve and how the impact of costs (even if not accurately known) can be included [29]. We examine binary decision or hypothesis testing problems where, given information X_t, on the dynamics of an infectious disease, we must decide if it is optimal to act now (at time t) or wait for future information. Let H_0 denote our null hypothesis, which defines the situation that we should do nothing. Alternatively, H_1 represents the case that we should act. For a growing epidemic this may be to initiate non-pharmaceutical interventions (NPIs) and for a declining one this may involve relaxing existing NPIs. These hypotheses are reassessed sequentially with time and delineate optimal decision points for action. While we consider binary decision-making only, the expressions below are known to generalise to complex decision problems involving multiple hypotheses. Solutions to those problems are qualitatively similar [30].

To determine what optimal means, we must attach costs to actions. We define c_{ij} as the cost of acting according to H_i when H_j is true. A false alarm (false positive, FP), which is interpreted as acting prematurely to initiate or relax an intervention, is then $c_{FP} = c_{10}$. A missed action i.e., acting too slowly (false negative, FN) is $c_{FN} = c_{01}$. A true positive (TP) or true negative (TN), which indicates when an action (or inaction) is appropriately taken, has cost of $c_{TP} = c_{11}$ or $c_{TN} = c_{00}$. The expected cost of a decision i, $\mathbf{E}[c_i]$, and the associated optimal action $i_t^* =$

$\min_{i\in\{0,1\}} \mathbf{E}[c_i]$, follow as in **Eq. (1)**. There $\mathbf{P}(H_j|X_t)$ is the posterior distribution of the evidence for hypothesis $j \in \{0,1\}$ given available information X_t, with $\mathbf{1}(.)$ as an indicator function [30].

$$\mathbf{E}[c_i] = \sum_{j\in\{0,1\}} c_{ij}\mathbf{P}(H_j|X_t), \qquad i_t^* = \mathbf{1}\left(\frac{\mathbf{P}(H_1|X_t)}{\mathbf{P}(H_0|X_t)} \geq \epsilon = \frac{c_{FP} - c_{TN}}{c_{FN} - c_{TP}}\right). \quad (1)$$

Eq. (1) states that the optimal action depends on a threshold or decision boundary ϵ that solely depends on the costs. This standard result from decision theory is already interesting as even when we do not know these costs accurately, we can still assess their likely effects via different choices of ϵ. Moreover, if we include prior distributions on each hypothesis, **Eq. (1)** defines a likelihood ratio test. **Eq. (1)** also generalises to multiple hypotheses [29], where we would find different thresholds for every hypothesis. We can rearrange our decision rule from **Eq. (1)** by recognising that $\mathbf{P}(H_0|X_t) = 1 - \mathbf{P}(H_1|X_t)$ to obtain **Eq. (2)**, with $\eta = \frac{\epsilon}{1+\epsilon}$.

$$i_t^* = \mathbf{1}\left(\mathbf{P}(H_1|X_t) \geq \eta = \frac{c_{FP} - c_{TN}}{(c_{FP} + c_{FN}) - (c_{TP} + c_{TN})}\right), \qquad t^* = \underset{t\geq 0}{\operatorname{argmin}}(i_t^* = 1). \quad (2)$$

This expression reveals that our decision to initiate (or relax) an intervention in the face of a potentially growing (or waning) epidemic rests on comparing the posterior evidence for the need to act against the proportion of the total excess cost of making incorrect choices that results from false alarms. The time at which we are statistically justified in acting is t^* under this model i.e., the first time when the evidence for action, $\mathbf{P}(H_1|X_t)$, exceeds η. Larger $c_{FP} - c_{TN}$ can only be counterbalanced by stronger evidence for the need to act (i.e., to reject H_0) and increases t^*. **Eq. (2)** is noteworthy as whatever the costs of action or inaction, ultimately optimal decision-making involves testing a threshold η against our evidence, which (assuming no change in the epidemic state) will accumulate with time. If false alarms (or missed actions) are prohibitively expensive then η rises to 1 (or falls to 0) and t^* becomes infinite (or 0).

Since we cannot control (or sometimes even know) the value of η, we focus on understanding the factors that regulate the posterior evidence. If H_1 is true, then we expect that any threshold will eventually be crossed. However, for fixed costs, this lag to crossing is regulated by at least two key factors. The first is the choice of information source X_t. In the next section we consider two reasonable and elementary metrics – the reported count of symptomatic cases over time and the inferred transmissibility parameters underpinning those cases. Second, the evidence for initiating (or relaxing) an intervention will deteriorate with the level of noise in our epidemic

surveillance data. We examine and quantify, for the first time to our knowledge, how key noise sources reduce the ability of the above metrics to inform decision thresholds.

We also reinterpret this Bayesian detection problem from an information theoretic perspective. Effectively, we want to communicate one bit of information to a decision maker i.e., a 1 or 0 to indicate the need for action or inaction respectively. When H_1 is true, then $\mathbf{P}(H_1|X_t) = p_t$ will accrue with time and is the probability that we should act given the epidemic state (which we estimate from X_t). The Shannon entropy $\mathcal{H}_{p_t} = p_t \log_2 \frac{1}{p_t} + (1 - p_t) \log_2 \frac{1}{1-p_t}$ describes the uncertainty for a distribution over two choices with probabilities p_t and $1 - p_t$ [31]. Logarithms are to the base 2 so $0 \leq \mathcal{H}_{p_t} \leq 1$ is in bits. In **Eq. (3)** we compute the negative derivative of this function with p_t, showing how this uncertainty reduces as we accrue evidence to act.

$$\frac{d\mathcal{H}_{p_t}}{dp_t} = \log_2 \left(\frac{1 - \mathbf{P}(H_1|X_t)}{\mathbf{P}(H_1|X_t)} \right), \qquad i_t^* = \mathbf{1}\left(-\frac{d\mathcal{H}_{p_t}}{dp_t} \geq \log_2 \epsilon \right). \quad (3)$$

Consequently, our decision-making process compares the loss of randomness or increase in certainty that H_1 is true against a cost-based threshold. This is characterised by $-\frac{d\mathcal{H}_{p_t}}{dp_t}$, which is defined by a logit function. We can apply an analogous procedure to verify that symmetrical reasoning holds for selecting H_0. The speed at which we cross our decision threshold, i.e., the cost-adjusted amount of time t^* that we should wait before acting, is controlled by how quickly p_t rises with time [32]. For any given cost structure, this time depends on the noise in epidemic surveillance, the effects of which we explore next.

Responding to epidemic growth and decline using reported incidence

The most timely and visible indicator of the likely state of an epidemic is the reported incidence of symptomatic cases, C_t at time t. This serves as a proxy for the count of new infections, I_t, which is rarely observable. Case incidence also measures potential epidemic burden because the incidence of hospitalisations or deaths may be described as delayed and resized versions of C_t [33,34]. A common approach to epidemic detection involves sequentially comparing the reported incidence against some baseline threshold [6,25,35,36]. We denote such a baseline by a and define our decision problem with the posterior evidence of $\mathbf{P}(H_1|X_t) \equiv \mathbf{P}(I_t \geq a|C_1^t)$ for a growing epidemic and $\mathbf{P}(H_1|X_t) \equiv \mathbf{P}(I_t \leq a|C_1^t)$ for a waning or controlled epidemic. We use Y_i^j for some variable Y to represent the time series or vector $\{Y_i, Y_{i+1}, \ldots, Y_{j-1}, Y_j\}$. Our evidence is based on the likely number of infections given observed cases.

We start by deriving analytic insight from a related deterministic model of epidemic spread. As epidemics often grow exponentially, we may describe the true infection incidence I_t and the reported case incidence C_t as in **Eq. (4)**, with r_0 as the growth rate, ρ the (fixed) proportion of infections reported, I_0 the initial infection number and τ as the (constant) reporting delay. The quantity ρ includes effects such as under-reporting and asymptomatic spread while τ encodes latencies due to surveillance limitations and lags in diagnosing symptomatic patients.

$$I_t = I_0 \exp(r_0 t), \qquad C_t = \rho I_0 \exp(r_0(t - \tau)), \qquad r_0 = \gamma(R_0 - 1). \quad (4)$$

The last expression in **Eq. (4)** provides the version of this growth process that corresponds to the dynamics of an SIR or SEIR model when the fraction of susceptible individuals is constant, with γ as the duration of infectiousness and R_0 as the basic reproduction number. This relation can be used to approximate epidemic dynamics more generally as well [37].

We assume that neither the reporting fraction nor the delay are known. This is common during the early stages of an emerging outbreak but may remain problematic into the later epidemic stages due to interventions or new pathogen variants changing the value of these and related epidemiological variables [38]. Consequently, we may often have to make decisions by simply choosing some alert threshold a and finding the first time that this value is exceeded by the epidemic curve. Such exceedance approaches have been practically used [6] but, as far as we can tell, the heterogeneous impacts of major surveillance noise sources on these decision-making problems has not been quantified analytically. We consider both growing and waning epidemics with $\Delta_{\mathrm{grow}} = t_{C \geq a} - t_{I \geq a}$ as the lag in exceeding our alert threshold for a growing epidemic and $\Delta_{\mathrm{wane}} = t_{I \leq a} - t_{C \leq a}$ as the related time lag for a waning epidemic.

We calculate these crossing times by equating expressions from **Eq. (4)** with the alert value i.e., we solve for $a = I_0 \exp(r_0 t_{I \geq a}) = \rho I_0 \exp(r_0(t_{C \geq a} - \tau))$. This yields **Eq. (5)** with $r_0 > 0$ ($r_0 < 0$) for growing (waning) epidemics and τ_ρ as an effective lag caused by ρ.

$$\Delta_{\mathrm{grow}} = \tau + \frac{1}{|r_0|}\log\frac{1}{\rho}, \qquad \Delta_{\mathrm{wane}} = \tau - \frac{1}{|r_0|}\log\frac{1}{\rho}, \qquad \rho \equiv \exp(-|r_0|\tau_\rho). \quad (5)$$

While **Eq. (5)** is trivial to derive it embodies some key insights. First, we can treat the influence of incomplete reporting as an effective shift of τ_ρ time units. Second, τ_ρ adds to the reporting delay τ for growing epidemics but subtracts from it when aiming to detect waning epidemics. Accordingly, early detection of emerging epidemics is substantially more difficult than the early detection of waning epidemics. For example, if $\tau_\rho \approx \tau$, then $\Delta_{\mathrm{grow}} \approx 2\tau$ but $\Delta_{\mathrm{wane}} \approx 0$.

This asymmetry is striking and, importantly, we find this insight holds for more realistic models. Specifically, we simulate epidemic case incidence according to the noisy renewal model from **Eq. (10)** of the Methods, which includes stochasticity both in the incidence generation and the reporting and delay distributions. Using parameters for COVID-19 we showcase the expected asymmetry for hitting alert thresholds in **Fig 1**. Our results do not depend on a being constant so we can also allow it to vary with time. In this setting, it can theoretically encode stochastic, time-varying and seasonal baselines for exceedance alerts.

Fig 1: Extrinsic detection case threshold performance under noisy surveillance. We plot the lag in crossing alert threshold a for 1000 epidemics simulated with COVID-19 parameters under a renewal process (see Methods). Left panels are upward crossing lag times $\Delta_{grow} = t_{C \geq a} - t_{I \geq a}$ during the growth stage of the simulated epidemics. Right panels show downward crossing lags $\Delta_{wane} = t_{I \leq a} - t_{C \leq a}$ for waning epidemics. We plot lags for cases corrupted by only under-reporting, delays and then for both noise sources combined.

The histograms of **Fig 1** summarise the lags in times at which cases C_t exceed a relative to infections I_t i.e., Δ_{grow} and Δ_{wane}. These inform on our evidence criteria $\mathbf{P}(I_t \geq a | C_1^t) \geq \eta$ for growing epidemics and $\mathbf{P}(I_t \leq a | C_1^t) \geq \eta$ for declining ones. We directly use case counts instead of hypothesis probabilities as cases provide the clearest extrinsic signals that may be used to decide if to act or not [3]. Note that we can infer infections from cases and other proxies by simply upsizing the case counts by the noise probabilities (under a popular binomial model) [34,39]. Our main insight is that irrespective of the values of η and a, practical noise sources induce a substantial asymmetry that makes timely responses to resurgences markedly more difficult e.g., in **Fig 1** at every a we find that $\mathbf{E}[\Delta_{\text{grow}}] > 6\mathbf{E}[\Delta_{\text{wane}}]$. Next, we test if these asymmetries hold when intrinsic transmissibility signals are used instead.

Responding to epidemic growth and decline using transmissibility estimates

The reproduction number, R, is a popular measure of infectious disease transmissibility that is frequently among the metrics used to inform public health policymaking [40]. The value of R is compared to a threshold of 1 to indicate whether the epidemic will grow or wane. Because R is a latent parameter underlying the epidemic, it needs to be estimated from observed data. The incidence of new infections, I_t, contains the most information for inferring R, but practical surveillance biases mean that we frequently can only observe new cases, C_t. This necessarily results in a loss in information about R, which leads to an increase in our estimate uncertainty. We can quantify these losses by computing the Fisher information of R from the incidence of infections $\mathbf{I}(R | I_1^t)$ and cases $\mathbf{I}(R | C_1^t)$ [31,34]. The Fisher information is an important measure because it defines the smallest asymptotic uncertainty achievable by any unbiased estimator.

We may use estimates of R to inform actions by setting our decision problem according to the posterior evidence $\mathbf{P}(H_1 | X_t) \equiv \mathbf{P}(R \geq b | C_1^t)$ and $\mathbf{P}(H_1 | X_t) \equiv \mathbf{P}(R \leq b | C_1^t)$ for growing and waning epidemics respectively, with some threshold b. As we aim to derive analytic insights about our ability to solve these decision problems generally, we consider asymptotic limits at which the Fisher information measures the uncertainty around our R estimates. At these limits the posterior distribution of R is Gaussian with mean at its maximum likelihood estimate (which converges to the true R) and variance σ^2 inversely proportional to the Fisher information. The b that we should act on, given η, relates to the quantile function $\sqrt{2}\,\sigma \text{erf}^{-1}(2\eta - 1)$ (directly for $\mathbf{P}(R \leq b | C_1^t)$ or its complement for $\mathbf{P}(R \geq b | C_1^t)$), with erf as the error function.

Consequently, for any decision cost-threshold η the Fisher information plays a central role. We model epidemics using renewal processes as in the Methods and **Fig 1** and assume that

cases derive from infections with a reporting proportion of ρ_s at time s and cumulative delay probability F_{t-s} for reports delayed by at most $t-s$ time units. We then compute the Fisher information about R from case data $\mathbf{I}(R|C_1^t)$ as on the left side of **Eq. (6)**. This follows from the Methods and the framework developed in [34,41] and involves summing along the period over which R is constant. If we set $\rho_s = 1$ for all s (perfect reporting) and $F_0 = 1$ (no delays) then we obtain the Fisher information from the unobservable infection counts $\mathbf{I}(R|I_1^t)$.

$$\mathbf{I}(R|C_1^t) = \frac{1}{R}\sum_{s=1}^{t} \rho_s F_{t-s}\Lambda_s, \qquad \frac{\mathbf{I}(R|C_1^t)}{\mathbf{I}(R|I_1^t)} = \frac{\sum_{s=1}^{t} \rho_s F_{t-s}\Lambda_s}{\sum_{s=1}^{t}\Lambda_s} \leq 1. \quad (6)$$

We cannot compute these Fisher information values without knowledge of the noise sources corrupting our surveillance data, but we can extract key insights by taking the ratio of the case to infection Fisher information, $\frac{\mathbf{I}(R|C_1^t)}{\mathbf{I}(R|I_1^t)}$, as on the right of **Eq. (6)**. This reveals that the losses in information about R due to noise in the cases depend multiplicatively on noise terms $\rho_s F_{t-s}$ weighted by $\Lambda_s (\sum_{s=1}^{t}\Lambda_s)^{-1}$ across time, which scales with the number of active infections.

As delays are always forward in time and induce more information loss towards the present t (i.e., $F_0 = \min F_{t-s}$ as it is cumulative) we find an inherent asymmetry in our ability to make decisions for growing versus waning epidemics. Growing epidemics have an increasing Λ_s so $\Lambda_s (\sum_{s=1}^{t}\Lambda_s)^{-1}$ rises as $s \to t$, contributing more to the sum in **Eq. (6)**. Consequently, the noise from the delay has a magnified effect on the overall estimate uncertainty, limiting our evidence $\mathbf{P}(H_1|X_t)$. The converse occurs for declining epidemics. In **Fig 2** we plot Fisher information ratios from the simulations underlying **Fig 1** (which use realistic noise parameters) and expose this performance gap, with growing epidemics having ratios that are approximately 3 times smaller. Note that as we shrink the noise towards perfect reporting (not shown), both our ratios tend to 1 (as expected) but the convergence during the growth phase is slower.

Fig 2: Intrinsic reproduction number or growth rate detection performance under noisy surveillance. We simulate epidemics according to COVID-19 parameters via renewal models (see Methods) subject to under-reporting and delays. This results in 1000 case curves, which we plot in various colours in the bottom panel against the true infections (black). The first half of the curve is generated under a supercritical R (red, dashed and overlaid) and the latter half under a subcritical R (blue, dashed and overlaid). Top panels (vertical dashed matches the supercritical to subcritical transition in bottom panels) compute ratios of the Fisher information in noisy cases $I(R|C_1^t)$ from those simulations to that from the true infections $I(R|I_1^t)$ for the growing (red) and waning (blue) phases of the epidemic. Solid lines show mean values while shaded ribbons give the full range of the information ratios.

The main alternative to R is the epidemic growth rate r, which has similar properties [40,42] and indicates resurgence or control based on whether it is positive or negative. Assuming a gamma generation time distribution with shape α and scale β^{-1} parameters (see Methods) we can compute Fisher information values as on the left side of **Eq. (7)**.

$$\mathbf{I}(r|C_1^t) = \frac{\alpha}{\beta}\left(1 + \frac{r}{\beta}\right)^{\frac{\alpha-2}{2}}\sum_{s=1}^{t}\rho_s F_{t-s}\Lambda_s\,, \qquad \frac{\mathbf{I}(r|C_1^t)}{\mathbf{I}(r|I_1^t)} = \frac{\mathbf{I}(R|C_1^t)}{\mathbf{I}(R|I_1^t)}. \quad (7)$$

Interestingly, we observe that the noise terms appear in the same way as **Eq. (6)** and that the remaining terms are independent of the noise. As a result, when we take the ratio of case to infection data Fisher information values, as in the right of **Eq. (7)**, we obtain precisely the same result as in **Eq. (6)**. The noise induced asymmetry observed for R therefore also affects r and likely will not be overcome by changing our measure of epidemic transmissibility. We confirm this further by noting that recent metrics from [43] that generalise Λ_s to remove its dependence on the generation time distribution have similar Fisher information formulae and will also show this skew. Consequently, optimal decision thresholds for growing epidemics are substantially more difficult to resolve than equivalent thresholds for waning epidemics.

Asymmetry persists under complex (feedback) decision frameworks

In previous sections we demonstrated that surveillance noise induces asymmetric limits to optimal decision-making when comparing growing and waning epidemics. Since we aimed to extract generalisable insights, the models we investigated, while commonly used to study real epidemics [40], were somewhat simplified. Here we confirm that the asymmetry we uncovered remains an intrinsic barrier to performance even when cost-optimal decision frameworks that leverage feedback control theory [44,45] and Bayesian optimisation are employed. Using the same parameters from **Fig 2** (see Methods), we simulated COVID-19 epidemics and applied the model predictive control algorithm from [28] to optimise the timing of decisions.

This algorithm aligns with the framework of **Eq. (2)** but explicitly evaluates and propagates the costs of both action and inaction at every decision time (weekly). We detail these costs in **Eq. (13)** of the Methods, including terms that account for the disruption (both economic and other types) of initiating an NPI and penalties for the trajectory of the epidemic (e.g., peak and endemic infection loads induce costs linked to healthcare burdens) expected to occur due to our NPI decisions. We model a lockdown as a multiplicative reduction in R, using parameters consistent with [46]. We find the optimal time for initiating (relaxing) a lockdown given a growing (waning) epidemic, by minimising **Eq. (13)** across a projected horizon that considers longer-term feedback or rebound effects resulting from action or inaction.

We present results of this algorithm in **Fig 3** for simulated COVID-19 epidemics. We compute optimal decision times for the perfect (noiseless) case and for scenarios featuring realistic surveillance noise (under-reporting and delays in line with previous sections). Analysing the start, end and length of lockdowns (histograms in **Fig 3**), we observe that noisy surveillance

causes lags of 2-4 weeks when initiating an NPI. However, this same noise induces delays of only under 1 week when relaxing that NPI. The growth-waning asymmetry is well-defined and persistent even under sophisticated decision algorithms. In **Fig 4** we show equivalent analyses for epidemics simulated under Ebola virus parameters (see Methods for details), verifying the consistency of our claims. Noise causes delays of 4-5 weeks when initiating lockdowns but only 1-2 weeks when relaxing them. Absolute times are longer for Ebola virus due to its slower timescale of spread (its mean generation time [47] is more than double that of COVID-19).

Fig 3: Explicitly optimised NPI initiation and relaxation under noisy surveillance. We simulate epidemics under COVID-19 parameters (see Methods) and renewal models and then apply the model predictive control (MPC) algorithm from [28]. This approach projects the costs of action or inaction forward in time together with the outcomes of those choices (e.g., larger epidemic peaks or endemic infection loads) and optimises timing based on minimising those costs over the projections (see Methods). Left panels show that the optimal lockdown (LD) or NPI decision time and properties for a growing epidemic are substantially affected by imperfect surveillance. In some subfigures multiple LD actions are visible because the MPC algorithm continually optimises costs but we focus only on the statistics of the first decision (in line with **Figs 1-2**). This is characterised by accompanying histograms plotting the distributions of LD start, end and duration for the perfect (blue) and noisy (green) scenarios. The difference in the

corresponding histograms shows the impact of the surveillance noise. In contrast, right panels demonstrate that the optimal relaxing of lockdown or another related NPI is minimally affected by imperfect surveillance with the histograms also showing smaller discordance.

Fig 4: Persistent noise-induced asymmetry under optimal NPI initiation and relaxation. We repeat the simulation of **Fig 3** but under Ebola virus transmission and surveillance noise parameters (see Methods for details). We apply the MPC algorithm from [28] to optimise the timing of initiating and relaxing a lockdown (LD) by minimising joint projected costs from the NPI and its associated epidemiological outcomes. The LD start, end and duration (see the histograms displaying these statistics) become much worse due to realistic surveillance noise when initiating (left panels, compare blue and green histograms) versus relaxing the LD (right panels, compare blue and green histograms). The asymmetry in **Fig 3** therefore persists.

Discussion

Policymakers often have to make critical public health decisions from uncertain and unreliable data. While research has shown how this uncertainty can markedly impact (often in unintuitive ways) the timing, selection and success of interventions [7,12,16,18,28], much is still unknown about how real-time decision-making can best incorporate or mitigate this uncertainty. Recent studies have proposed integrating uncertainty within rigorous decision theory frameworks [48–

50]. Here we support this proposal but instead of applying complex modelling to guide specific action or inaction, we concentrated on extracting generalisable insights about how uncertainty fundamentally limits decision-making. We tackled this problem by exploring how predominant forms of surveillance uncertainty or noise, relating to under-reporting and delays in reporting cases, present bottlenecks for real-time decisions based on two of the most common outbreak indicators – the incidence of cases and the reproduction number (or related growth rate).

We discovered that there is a surprising, intrinsic and important asymmetry induced by these sources of noise that results in substantially reduced performance of the solutions to optimal decision problems for growing epidemics, relative to the performance on equivalent problems for waning epidemics. This asymmetry remains irrespective of whether we base decisions on extrinsic proxies of infection incidence (**Fig 1**) or intrinsic estimates of transmissibility (**Fig 2**). We found theoretical justification for this asymmetry using information and Bayesian detection theory and then confirmed this asymmetry remained when complex algorithms were used to explicitly optimise decisions according to the costs of action as well as the opportunity costs of inaction (**Fig 3**-**4**). While surveillance noise is expected to restrict actionable information, it is not obvious that the bottleneck it imposes should be notably more restrictive during growth.

There are several important ramifications of this asymmetry. First, as this innate performance restriction on responding to epidemic growth is likely further compounded by scarcer data and additional unknowns (e.g., transmissibility can be poorly specified during epidemic emergence [12]), proactive interventions may be necessary to achieve timely control [7,23]. In contrast, a more reactive approach to relaxing interventions may be sufficient. Second, if the surveillance noise can be reduced by allocating resources (e.g., rapid diagnostic tests), then deployment should prioritise earlier stages of outbreaks. Third, when NPI effectiveness is retrospectively assessed it may be valuable to contextualise performance against the limits imposed by the bottlenecks we have explored to gain a more objective evaluation. This provides a clearer reference for effectiveness, ensuring we are not comparing against unrealistic ideals.

Although the asymmetry we uncovered is fundamental to noisy epidemic surveillance, there are limitations to our analyses. We only considered homogeneous (well-mixed) transmission models. The impact of realistic heterogeneities in spread due to geography, demography and other characteristics may influence what an optimal decision should be and hence the level of asymmetry. This is an open question that we aim to explore in future. Additionally, we have not examined how auxiliary epidemic data could help us overcome these bottlenecks. Recent initiatives [51] have proposed integrating wastewater surveys and genomic sequencing within standard surveillance schemes. While promising, the potential of these data for enhancing

early-warnings of outbreaks is still being assessed. Should these data feature some latency and under-ascertainment [34], we expect decision-making asymmetries to re-emerge.

Lastly, we constrained our analyses to binary decisions between action and inaction based on data available in real time. This was both to embody the common situation where policymakers must make choices from the latest (updating) data and to allow for analytic and generalisable insights to be extracted. However, this framework neglects that optimal decisions could involve more complex trade-offs among multiple, simultaneous NPI options with differing benefits and costs informed by both real-time and historical data from other outbreaks or past experience with those NPIs. Although some complex decision problems can reduce to generalisations of the binary case we study, yielding multiple decision thresholds [26,30], it remains unknown if and when these complexities could modify the performance asymmetry that we uncovered, particularly when combined with the heterogeneities and auxiliary data above.

These limitations notwithstanding, the consistent asymmetry we found, using both elementary mathematical arguments and sophisticated (predictive) algorithms, underscores a meaningful directionality in how surveillance imperfections shape and bound decision-making in the face of uncertainty. Knowing these asymmetric performance bounds is not only helpful for deciding when interventions need to be proactive or reactive, but also for diagnosing how our responses to outbreaks may fail and for justifying why faster and more stringent measures are necessary. With mounting calls for better integration of formal decision frameworks [16,18,45,48] together with enhanced and multimodal surveillance data [38,51], epidemic models will only become more complex and difficult to interrogate. Having generalisable insights into performance limits can help anticipate failures in surveillance-driven decisions and improve their robustness.

Methods

Renewal models with practical surveillance

The renewal branching process [52] is widely used to model infectious disease epidemics of COVID-19, Ebola virus, pandemic influenza and many others [53]. It simulates the incidence of new infections I_t at some time t in terms of the disease reproduction number R and past incidence $I_1^{t-1} \equiv \{I_1, I_2, \dots, I_{t-1}\}$ as in the left of **Eq. (10)** with **Pois** as Poisson noise. Here Λ_t is the total infectiousness and encodes the impact of past incidence i.e., $\Lambda_t \stackrel{\text{def}}{=} \sum_{x=1}^{t-1} w_{t-x} I_x$. The w_{t-x} terms are the probability of an infection being transmitted in $t - x$ time units and define the generation time distribution of the disease with $\sum_{u=1}^{\infty} w_u = 1$ [54].

$$I_t \sim \mathbf{Pois}(\Lambda_t R), \qquad C_t \sim \mathbf{Pois}\left(R \sum_{s=1}^{t} \rho_s \, \delta_{t-s} \, \Lambda_s \right). \quad (10)$$

Normally infections are not observable, so the standard renewal model is modified to describe the incidence of symptomatic cases, C_t at time t, instead. This requires including noise terms such as reporting fractions, ρ_s at time s, and the probability, δ_{t-s} of a reporting delay of $t - s$ time units. This leads to the right side of **Eq. (10)** [34,39]. In all of these equations we assume a constant R until time t but note that they remain valid for time-varying reproduction numbers. If we have no delay and perfect reporting then $\rho_s = 1$, $\delta_0 = 1$, $\delta_{x>0} = 0$ and $C_t = I_t$.

In **Fig 1-3** we simulate under established COVID-19 transmission and surveillance noise parameters from the literature. We use the generation time distribution from [10] (a gamma distribution with mean of 6.5 days) and model delays in reporting as done in [55] (a negative binomial distribution with mean of 10.8 days) with the fraction of infections reported as cases based on [56] (a beta distribution with mean of 0.38, implying under-reporting of mean 0.72). In **Fig 4** we perform equivalent simulations to **Fig 3** but now under Ebola virus parameters. We apply the generation time distribution from [47] (a gamma distribution with a mean of 15.3 days) and reporting noise from [57,58] (delays are follow a negative binomial distribution with mean of 11.8 days and under-reporting conforms to a beta distribution with mean of 0.4). For the precise distributions underlying both our COVID-19 and Ebola virus analyses (as well as code to reproduce these figures) see the Data and code availability section.

Fisher information given noisy observations

The minimum (asymptotic) uncertainty around estimates of R derived from the renewal model of **Eq. (10)** can be quantified using the Fisher information $\mathbf{I}(.)$. This can be computed from the expected curvature of the log-likelihood of the statistical models from **Eq. (10)**. This leads to the relations in **Eq. (11)**, which provide the information from infections and cases respectively, and are adapted from the frameworks introduced in [34,41]. Here $F_{t-s} = \sum_{x=0}^{t-s} \delta_x$ describes the cumulative reporting delay probability and $\mathbf{I}(R|I_1^t) \geq \mathbf{I}(R|C_1^t)$ (noise reduces information).

$$\mathbf{I}(R|I_1^t) = \frac{1}{R}\sum_{s=1}^{t} \Lambda_s\,, \qquad \mathbf{I}(R|C_1^t) = \frac{1}{R}\sum_{s=1}^{t} \rho_s F_{t-s} \Lambda_s. \quad (11)$$

In the main text we examine ratios of these information terms and also derive related ratios for epidemic growth rates r. These growth rates have a mapping to R described by the Euler-Lotka equation [54]. This equation depends on the generation time distribution of the disease. Commonly, a gamma-distributed generation time distribution is assumed [42,53] i.e., the w_u probabilities describe a shape-scale $\mathbf{Gam}(\alpha, \beta^{-1})$ distribution. Under this setting the Euler-Lotka relationship $f(.)$ takes the form of the left side of **Eq. (12)**.

$$R = f(r) = \left(1 + \frac{r}{\beta}\right)^{\alpha}, \qquad \mathbf{I}(r|X_1^t) = \mathbf{I}(f(r)|X_1^t)\left(\frac{df}{dr}\right)^2. \quad (12)$$

For a given generation time distribution, R is a smooth function of r, $f(r)$, and we may apply the Fisher information change of variables formula on the right side of **Eq. (12)** to convert the expressions for R in **Eq. (11)** to ones for the growth rate. Here X_1^t may be the time series of cases C_1^t or infections I_1^t as required. In the main text we use the above equations to derive Fisher information ratios for both reproduction numbers and growth rates.

Cost-optimal feedback control of epidemics

We have focussed on extracting general insights about the asymmetries in response times to growing or waning epidemics under arbitrary cost thresholds. This required tractable modelling approaches that facilitate analytic results. However, it is possible to instead construct complex algorithms [15] that directly integrate competing costs from interventions and disease burden (e.g., epidemic peaks and infection endemic levels), as well as consider longer-term, feedback effects from chosen interventions [45]. We adopt such an algorithm from [28], which uses model predictive control (MPC) [44] to balance the component costs of $\psi(t)$ in **Eq. (13)**.

$$\psi(t) = \alpha|I_t - I_{\text{end}}| + \beta\mathbf{1}\big(I_t > I_{\text{pk}}\big) + \phi(\text{NPI}_t). \quad (13)$$

This MPC algorithm considers costs due to (i) the difference between infections and a practical endemic goal I_{end} (weighted by α), (ii) exceeding a peak value I_{pk}, which for example models the level of infections beyond which healthcare resources are overrun (weighted by β) and (iii) the actual economic or other penalties from enforcing an NPI (wrapped in a flexible function $\phi(.)$). Heuristically, applying a more stringent NPI increases (iii) but can decrease (i) and (ii). Hence, we penalise both action and inaction. This aligns and expands on the framework from **Eq. (2)** and implies a threshold for action based on how the components of (i)-(iii) balance.

The MPC approach of [28] finds the optimal time to initiate or relax an NPI by minimising the long-term costs of those actions over a horizon h i.e., $\sum_{s=0}^{h} \gamma^s \psi(t+s)$, with γ as a discount factor. This is done by projecting the consequences of those choices using a generalised form of **Eq. (10)** and then applying Bayesian optimisation to select the cost-optimal choice. See [28] for complete details of this procedure. In our analyses NPI_t is 0 when inactive at time t. When active it is 1 and models a lockdown or stay-at-home order as a multiplicative reduction in R (though it can be easily modified to model other intervention types). The cost $\phi(1)$ that is associated with this action is set based on analyses in [46]. Late NPI relaxation increases

(iii), while late NPI initiation increases (i)-(ii). When surveillance is noisy, we replace I_t with cases C_t and apply appropriate delay and under-reporting parameters from the literature.

Bibliography

1. Wagner MM, Tsui FC, Espino JU, Dato VM, Sittig DF, Caruana RA, et al. The emerging science of very early detection of disease outbreaks. J Public Health Manag Pract. 2001;7: 51–59. doi:10.1097/00124784-200107060-00006

2. Buckee C. Improving epidemic surveillance and response: big data is dead, long live big data. Lancet Digit Health. 2020;2: e218–e220. doi:10.1016/S2589-7500(20)30059-5

3. Jackson ML, Baer A, Painter I, Duchin J. A simulation study comparing aberration detection algorithms for syndromic surveillance. BMC Med Inform Decis Mak. 2007;7: 6. doi:10.1186/1472-6947-7-6

4. Brett TS, Drake JM, Rohani P. Anticipating the emergence of infectious diseases. J R Soc Interface. 2017;14. doi:10.1098/rsif.2017.0115

5. Yuan M, Boston-Fisher N, Luo Y, Verma A, Buckeridge DL. A systematic review of aberration detection algorithms used in public health surveillance. J Biomed Inform. 2019;94: 103181. doi:10.1016/j.jbi.2019.103181

6. Sonesson C, Bock D. A review and discussion of prospective statistical surveillance in public health. J Royal Statistical Soc A. 2003;166: 5–21. doi:10.1111/1467-985X.00256

7. Morris DH, Rossine FW, Plotkin JB, Levin SA. Optimal, near-optimal, and robust epidemic control. Commun Phys. 2021;4: 78. doi:10.1038/s42005-021-00570-y

8. Metcalf CJE, Andriamandimby SF, Baker RE, Glennon EE, Hampson K, Hollingsworth TD, et al. Challenges in evaluating risks and policy options around endemic establishment or elimination of novel pathogens. Epidemics. 2021;37: 100507. doi:10.1016/j.epidem.2021.100507

9. Metcalf CJE, Lessler J. Opportunities and challenges in modeling emerging infectious diseases. Science. 2017;357: 149–152. doi:10.1126/science.aam8335

10. Ferguson N, Laydon D, Nedjati Gilani G, Imai N, Ainslie K, Baguelin M, et al. Report 9: Impact of non-pharmaceutical interventions (NPIs) to reduce COVID19 mortality and healthcare demand. Imperial College London. 2020; doi:10.25561/77482

11. Karatayev VA, Anand M, Bauch CT. Local lockdowns outperform global lockdown on the far side of the COVID-19 epidemic curve. Proc Natl Acad Sci USA. 2020;117: 24575–24580. doi:10.1073/pnas.2014385117

12. Probert WJM, Jewell CP, Werkman M, Fonnesbeck CJ, Goto Y, Runge MC, et al. Real-time decision-making during emergency disease outbreaks. PLoS Comput Biol. 2018;14: e1006202. doi:10.1371/journal.pcbi.1006202

13. Greer SL, King EJ, da Fonseca EM, Peralta-Santos A. The comparative politics of COVID-19: The need to understand government responses. Glob Public Health. 2020;15: 1413–1416. doi:10.1080/17441692.2020.1783340

14. Sebhatu A, Wennberg K, Arora-Jonsson S, Lindberg SI. Explaining the homogeneous diffusion of COVID-19 nonpharmaceutical interventions across heterogeneous countries. Proc Natl Acad Sci USA. 2020;117: 21201–21208. doi:10.1073/pnas.2010625117

15. Haw DJ, Morgenstern C, Forchini G, Johnson R, Doohan P, Smith PC, et al. Data needs for integrated economic-epidemiological models of pandemic mitigation policies. Epidemics. 2022; 100644. doi:10.1016/j.epidem.2022.100644

16. Barnett M, Buchak G, Yannelis C. Epidemic responses under uncertainty. Proc Natl Acad Sci USA. 2023;120: e2208111120. doi:10.1073/pnas.2208111120

17. Keogh-Brown MR, Jensen HT, Edmunds WJ, Smith RD. The impact of Covid-19, associated behaviours and policies on the UK economy: A computable general equilibrium model. SSM Popul Health. 2020;12: 100651. doi:10.1016/j.ssmph.2020.100651

18. Shea K, Borchering RK, Probert WJM, Howerton E, Bogich TL, Li S-L, et al. Multiple models for

outbreak decision support in the face of uncertainty. Proc Natl Acad Sci USA. 2023;120: e2207537120. doi:10.1073/pnas.2207537120

19. Li S-L, Bjørnstad ON, Ferrari MJ, Mummah R, Runge MC, Fonnesbeck CJ, et al. Essential information: Uncertainty and optimal control of Ebola outbreaks. Proc Natl Acad Sci USA. 2017;114: 5659–5664. doi:10.1073/pnas.1617482114

20. Ferguson NM, Keeling MJ, Edmunds WJ, Gani R, Grenfell BT, Anderson RM, et al. Planning for smallpox outbreaks. Nature. 2003;425: 681–685. doi:10.1038/nature02007

21. Carrasco LR, Jit M, Chen MI, Lee VJ, Milne GJ, Cook AR. Trends in parameterization, economics and host behaviour in influenza pandemic modelling: a review and reporting protocol. Emerg Themes Epidemiol. 2013;10: 3. doi:10.1186/1742-7622-10-3

22. Fraser C, Riley S, Anderson RM, Ferguson NM. Factors that make an infectious disease outbreak controllable. Proc Natl Acad Sci USA. 2004;101: 6146–6151. doi:10.1073/pnas.0307506101

23. Parag KV, Donnelly CA. Fundamental limits on inferring epidemic resurgence in real time using effective reproduction numbers. PLoS Comput Biol. 2022;18: e1010004. doi:10.1371/journal.pcbi.1010004

24. Dehning J, Zierenberg J, Spitzner FP, Wibral M, Neto JP, Wilczek M, et al. Inferring change points in the spread of COVID-19 reveals the effectiveness of interventions. Science. 2020;369. doi:10.1126/science.abb9789

25. Unkel S, Farrington CP, Garthwaite PH, Robertson C, Andrews N. Statistical methods for the prospective detection of infectious disease outbreaks: a review. J Royal Statistical Soc A. 2012;175: 49–82. doi:10.1111/j.1467-985X.2011.00714.x

26. Johnson P, Moriarty J, Peskir G. Detecting changes in real-time data: a user's guide to optimal detection. Philos Trans A, Math Phys Eng Sci. 2017;375. doi:10.1098/rsta.2016.0298

27. Kay SM. Fundamentals of Statistical Signal Processing: Detection theory. reprint. Prentice-Hall PTR; 1998.

28. Beregi S, Parag KV. Optimal algorithms for controlling infectious diseases in real time using noisy infection data. medRxiv. 2024; doi:10.1101/2024.05.24.24307878

29. Berger JO. Statistical decision theory and bayesian analysis. New York, NY: Springer New York; 1985. doi:10.1007/978-1-4757-4286-2

30. Hardt M, Recht B. [2102.05242] Patterns, predictions, and actions: A story about machine learning. arXiv. 2021;

31. Cover T, Thomas J. Elements of Information Theory Second Edition. John Wiley and Sons; 2006.

32. Gabriele T. Information criteria for threshold determination (Corresp.). IEEE Trans Inform Theory. 1966;12: 484–486. doi:10.1109/TIT.1966.1053918

33. Goldstein E, Dushoff J, Ma J, Plotkin JB, Earn DJD, Lipsitch M. Reconstructing influenza incidence by deconvolution of daily mortality time series. Proc Natl Acad Sci USA. 2009;106: 21825–21829. doi:10.1073/pnas.0902958106

34. Parag KV, Donnelly CA, Zarebski AE. Quantifying the information in noisy epidemic curves. Nat Comput Sci. 2022;2: 584–594. doi:10.1038/s43588-022-00313-1

35. Salmon M, Schumacher D, Stark K, Others. Bayesian outbreak detection in the presence of reporting delays. Biometr J. 2015;57: 1051–67.

36. Southall E, Brett TS, Tildesley MJ, Dyson L. Early warning signals of infectious disease transitions: a review. J R Soc Interface. 2021;18: 20210555. doi:10.1098/rsif.2021.0555

37. Bettencourt LMA, Ribeiro RM. Real time bayesian estimation of the epidemic potential of emerging infectious diseases. PLoS One. 2008;3: e2185. doi:10.1371/journal.pone.0002185

38. Kraemer MUG, Pybus OG, Fraser C, Cauchemez S, Rambaut A, Cowling BJ. Monitoring key epidemiological parameters of SARS-CoV-2 transmission. Nat Med. 2021;27: 1854–1855. doi:10.1038/s41591-021-01545-w

39. Azmon A, Faes C, Hens N. On the estimation of the reproduction number based on misreported epidemic data. Stat Med. 2014;33: 1176–1192. doi:10.1002/sim.6015

40. Anderson R, Donnelly C, Hollingsworth D, Keeling M, Vegvari C, Baggaley R. Reproduction number (R) and growth rate (r) of the COVID-19 epidemic in the UK: methods of. The Royal Society. 2020;

41. Parag KV, Donnelly CA. Adaptive estimation for epidemic renewal and phylogenetic skyline models. Syst Biol. 2020;69: 1163–1179. doi:10.1093/sysbio/syaa035

42. Parag KV, Thompson RN, Donnelly CA. Are epidemic growth rates more informative than reproduction numbers? J Royal Statistical Soc A. 2022; doi:10.1111/rssa.12867

43. Parag KV, Cowling BJ, Lambert BC. Angular reproduction numbers improve estimates of transmissibility when disease generation times are misspecified or time-varying. Proceedings of the Royal Society B: Biological Sciences. 2023;290. doi:10.1098/rspb.2023.1664

44. Schwenzer M, Ay M, Bergs T, Abel D. Review on model predictive control: an engineering perspective. Int J Adv Manuf Technol. 2021;117: 1327–1349. doi:10.1007/s00170-021-07682-3

45. Parag KV. How to Measure the Controllability of an Infectious Disease? Physical Review X. 2024;

46. Haw DJ, Forchini G, Doohan P, Christen P, Pianella M, Johnson R, et al. Optimizing social and economic activity while containing SARS-CoV-2 transmission using DAEDALUS. Nat Comput Sci. 2022;2: 223–233. doi:10.1038/s43588-022-00233-0

47. Van Kerkhove MD, Bento AI, Mills HL, Ferguson NM, Donnelly CA. A review of epidemiological parameters from Ebola outbreaks to inform early public health decision-making. Sci Data. 2015;2: 150019. doi:10.1038/sdata.2015.19

48. Runge MC, Shea K, Howerton E, Yan K, Hochheiser H, Rosenstrom E, et al. Scenario design for infectious disease projections: Integrating concepts from decision analysis and experimental design. Epidemics. 2024;47: 100775. doi:10.1016/j.epidem.2024.100775

49. Shearer FM, Moss R, McVernon J, Ross JV, McCaw JM. Infectious disease pandemic planning and response: Incorporating decision analysis. PLoS Med. 2020;17: e1003018. doi:10.1371/journal.pmed.1003018

50. Swallow B, Birrell P, Blake J, Burgman M, Challenor P, Coffeng LE, et al. Challenges in estimation, uncertainty quantification and elicitation for pandemic modelling. Epidemics. 2022;38: 100547. doi:10.1016/j.epidem.2022.100547

51. Cori A, Lassmann B, Nouvellet P. Data needs for better surveillance and response to infectious disease threats. Epidemics. 2023;43: 100685. doi:10.1016/j.epidem.2023.100685

52. Fraser C. Estimating individual and household reproduction numbers in an emerging epidemic. PLoS One. 2007;2: e758. doi:10.1371/journal.pone.0000758

53. Cori A, Ferguson NM, Fraser C, Cauchemez S. A new framework and software to estimate time-varying reproduction numbers during epidemics. Am J Epidemiol. 2013;178: 1505–1512. doi:10.1093/aje/kwt133

54. Wallinga J, Lipsitch M. How generation intervals shape the relationship between growth rates and reproductive numbers. Proc R Soc B. 2007;274: 599–604.

55. Irons NJ, Raftery AE. Estimating SARS-CoV-2 infections from deaths, confirmed cases, tests, and random surveys. Proc Natl Acad Sci USA. 2021;118. doi:10.1073/pnas.2103272118

56. Pullano G, Di Domenico L, Sabbatini CE, Valdano E, Turbelin C, Debin M, et al. Underdetection of cases of COVID-19 in France threatens epidemic control. Nature. 2021;590: 134–139. doi:10.1038/s41586-020-03095-6

57. Dalziel BD, Lau MSY, Tiffany A, McClelland A, Zelner J, Bliss JR, et al. Unreported cases in the 2014-2016 Ebola epidemic: Spatiotemporal variation, and implications for estimating transmission. PLoS Negl Trop Dis. 2018;12: e0006161. doi:10.1371/journal.pntd.0006161

58. Team WER. Ebola Virus Disease in West Africa – The First 9 Months of the Epidemic and Forward Projections. N Engl J Med. 2014;371: 1481–95.

Funding

KVP and SB acknowledge support (Reference No. MR/X020258/1) from the MRC Centre for Global Infectious Disease Analysis funded by the UK Medical Research Council. This UK-funded grant is carried out in the frame of the Global Health EDCTP3 Joint Undertaking. The funders played no role in study design, data collection and analysis, decision to publish, or manuscript preparation.

Data and code availability

All data and code to reproduce the analyses and figures of this work are freely available (in MATLAB) at: https://github.com/kpzoo/asymmetricDetection.