1 **Multi-ancestral genome-wide association study of clinically defined nicotine dependence**
2 **reveals strong genetic correlations with other substance use disorders and health-**
3 **related traits.**
4
5 Running head: GWAS of nicotine dependence
6

Emma C Johnson[1]*, Dongbing Lai[2]*, Alex P Miller[3], Alexander S Hatoum[1], Joseph D Deak[4,7],
Jared V Balbona[1], David AA Baranger[5], Marco Galimberti[6,7], Kittipong Sanichwankul[8], Thorgeir
Thorgeirsson[9], Sarah MC Colbert[10], Sandra Sanchez-Roige[11,12,13], Keyrun Adhikari[6,7], Anna
Docherty[14], Louisa Degenhardt[15], Tobias Edwards[16], Louis Fox[1], Alexandros Giannelis[16], Paul
Jeffries[1], Tellervo Korhonen[17], Claire Morrison[18,19], Yaira Z Nunez[6,7], Teemu Palviainen[17], Mei-
Hsin Su[20], Pamela N Romero Villela[18,19], Leah Wetherill[2], Emily A Willoughby[16], Stephanie
Zellers[17], Laura Bierut[1], Jadwiga Buchwald[17], William Copeland[21], Robin Corley[19], Naomi P.
Friedman[18,19], Tatiana M Foroud[2], Nathan A Gillespie[20,22], Ian R Gizer[25], Andrew C Heath[1], Ian B
Hickie[26], Jaakko A Kaprio[17], Matthew C Keller[18,19], James L Lee[16], Penelope A Lind[24,27,28],
Pamela A Madden[1], Hermine HM Maes[20,23,29], Nicholas G Martin[22], Matt McGue[16], Sarah E
Medland[24,30,31], Elliot C Nelson[1], John V Pearson[24,32], Bernice Porjesz[33], Michael Stallings[18,19],
Scott Vrieze[16], Kirk C Wilhelmsen[34,35,36], Raymond K Walters[37,38,39], Renato Polimanti[4,40,41],
Robert T Malison[4,42], Hang Zhou[4,7], Kari Stefansson[9,43], Marc N Potenza[4,41,44,45,46], Apiwat
Mutirangura[47], Vorasuk Shotelersuk[48,49], Rasmon Kalayasiri[50,51], Howard J Edenberg[2,52], Joel
Gelernter[4,6,7,53], Arpana Agrawal[1]
22
23
24 *These authors made equal contributions.
25
[1] Department of Psychiatry, Washington University School of Medicine in St Louis, Saint Louis,
MO, USA
[2] Department of Medical and Molecular Genetics, Indiana University School of Medicine,
Indianapolis, IN, USA
[3] Department of Psychiatry, Indiana University School of Medicine, Indianapolis, IN, USA
[4] Department of Psychiatry, Yale University School of Medicine, New Haven, CT, USA
[5] Department of Psychological and Brain Sciences, Washington University in St Louis, Saint
Louis, MO, USA
[6] Department of Human Genetics, Yale University School of Medicine, New Haven, CT, USA
[7] Department of Psychiatry, Veterans Affairs Connecticut Healthcare Center, West Haven, CT,
USA
[8] Suanprung Psychiatric Hospital, Chiang Mai, Thailand
[9] deCODE Genetics/Amgen, Sturlugata 8, IS-101, Reykjavik, Iceland
[10] Department of Genetics and Genomic Sciences, Icahn School of Medicine at Mount Sinai,
New York, NY, USA
[11] Department of Psychiatry, University of California San Diego, La Jolla, CA, USA
[12] Institute for Genomic Medicine, University of California San Diego, La Jolla, CA, USA
[13] Division of Genetic Medicine, Vanderbilt University Medical Center, Nashville, TN, USA
[14] Department of Psychiatry, University of Utah School of Medicine, Salt Lake City, UT, USA
[15] National Drug and Alcohol Research Centre, University of New South Wales, Sydney,
Australia
[16] Department of Psychology, University of Minnesota Twin Cities, Minneapolis, MN, USA
[17] Institute for Molecular Medicine Finland, University of Helsinki, Helsinki, Finland
[18] Department of Psychology and Neuroscience, University of Colorado Boulder, Boulder, CO,
USA

[19] Institute for Behavioral Genetics, University of Colorado Boulder, Boulder, CO, USA
[20] Virginia Institute for Psychiatric and Behavioral Genetics, Virginia Commonwealth University,
Richmond, VA, USA
[21] Department of Psychiatry, College of Medicine, University of Vermont, Burlington, VT, USA
[22] QIMR Berghofer Medical Research Institute, Brisbane, Queensland, Australia
[23] Department of Human and Molecular Genetics, Virginia Commonwealth University,
Richmond, VA, USA
[24] Psychiatric Genetics, QIMR Berghofer Medical Research Institute, Brisbane, Queensland,
Australia
[25] Department of Psychological Sciences, University of Missouri, Columbia, MO, USA
[26] Brain and Mind Institute, University of Sydney, New South Wales, Sydney, Australia
[27] School of Biomedical Sciences, Faculty of Medicine, University of Queensland, Brisbane,
QLD, Australia
[28] School of Biomedical Sciences, Queensland University of Technology, Brisbane, QLD,
Australia
[29] Department of Psychiatry, Virginia Commonwealth University, Richmond, VA, USA
[30] School of Psychology and Counselling, Queensland University of Technology, Brisbane,
Australia
[31] School of Psychology, University of Queensland, Brisbane, Australia
[32] Faculty of Medicine, The University of Queensland, Brisbane, Australia
[33] Henri Begleiter Neurodynamics Laboratory, Department of Psychiatry, SUNY Downstate
Health Science University, Brooklyn, NY, USA
[34] Department of Neurology and Genetics and the Bowles Center of Alcohol Studies, University
of North Carolina, Chapel Hill, NC, USA
[35] Rockefeller Neuroscience Institute, West Virginia University, Morgantown, WV, USA
[36] Renaissance Computing Institute, Chapel Hill, NC, USA
[37] Analytic and Translational Genetics Unit, Massachusetts General Hospital, Boston, MA, USA
[38] Harvard Medical School, Boston, MA, USA
[39] Stanley Center for Psychiatric Research, Broad Institute of Harvard and MIT, Cambridge, MA,
USA
[40] Cooperative Studies Program Clinical Epidemiology Research Center (CSP-CERC), VA
Connecticut Healthcare System, West Haven, CT, USA
[41] Wu Tsai Institute, Yale University, New Haven, CT, USA
[42] The Clinical Neuroscience Research Unit, Connecticut Mental Health Center, New Haven,
CT, USA
[43] Faculty of Medicine, University of Iceland, IS-101, Reykjavik, Iceland
[44] Child Study Center, Yale School of Medicine, New Haven, CT, USA
[45] Connecticut Council on Problem Gambling, Wethersfield, CT, USA
[46] Connecticut Mental Health Center, New Haven, CT, USA
[47] Department of Anatomy, Faculty of Medicine, Chulalongkorn University, King Chulalongkorn
Memorial Hospital, Bangkok, Thailand
[48] Center of Excellence for Medical Genomics, Department of Pediatrics, Faculty of Medicine,
Chulalongkorn University, Bangkok, Thailand
[49] Excellence Center for Genomics and Precision Medicine, King Chulalongkorn Memorial
Hospital, the Thai Red Cross Society, Bangkok, Thailand
[50] Department of Psychiatry, Epidemiology of Psychiatric Disorders and Mental Health Research
Unit, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand
[51] Department of Psychiatry, King Chulalongkorn Memorial Hospital, Bangkok, Thailand
[52] Department of Biochemistry and Molecular Biology, Indiana University School of Medicine,
Indianapolis, IN, USA

[53] Department of Neuroscience, Yale University School of Medicine, New Haven, CT, USA
102
103

104 **Conflicts of Interest**

105 J.G. is named as an inventor on PCT patent application #15/878,640 entitled: "Genotype-guided
106 dosing of opioid agonists," filed January 24, 2018. There are no other conflicts of interest to
107 disclose.
108
109

ABSTRACT

Genetic research on nicotine dependence has utilized multiple assessments that are in weak
agreement. We conducted a genome-wide association study of nicotine dependence defined
using the Diagnostic and Statistical Manual of Mental Disorders (DSM-NicDep) in 61,861
individuals (47,884 of European ancestry, 10,231 of African ancestry, 3,746 of East Asian
ancestry) and compared the results to other nicotine-related phenotypes. We replicated the
well-known association at the *CHRNA5* locus (lead SNP: rs147144681, p =1.27E-11 in
European ancestry; lead SNP = rs2036527, p = 6.49e-13 in cross-ancestry analysis). DSM-
NicDep showed strong positive genetic correlations with cannabis use disorder, opioid use
disorder, problematic alcohol use, lung cancer, material deprivation, and several psychiatric
disorders, and negative correlations with respiratory function and educational attainment. A
polygenic score of DSM-NicDep predicted DSM-5 tobacco use disorder and 6 of 11 individual
diagnostic criteria, but none of the Fagerström Test for Nicotine Dependence (FTND) items, in
the independent NESARC-III sample. In genomic structural equation models, DSM-NicDep
loaded more strongly on a previously identified factor of general addiction liability than did a
"problematic tobacco use" factor (a combination of cigarettes per day and nicotine dependence
defined by the FTND). Finally, DSM-NicDep was strongly genetically correlated with a GWAS of
tobacco use disorder as defined in electronic health records, suggesting that combining the
wide availability of diagnostic EHR data with nuanced criterion-level analyses of DSM tobacco
use disorder may produce new insights into the genetics of this disorder.
130

INTRODUCTION
Nicotine use and nicotine dependence (or tobacco use disorder) are influenced by both
genetic liability and environmental factors[1,2]. Several different smoking-related phenotypes have
been studied, including smoking initiation[3], cigarettes per day[3], the Fagerström Test for Nicotine
Dependence (FTND)[4], problematic tobacco use[5], tobacco use disorder defined by codes from
the International Classification of Diseases (ICD-TUD)[6], and nicotine dependence[7] defined by
the Diagnostic and Statistical Manual of Mental Disorders (DSM-NicDep) (**Table 1**).
Genome-wide association studies (GWAS) of nicotine use behaviors have largely
focused on phenotypes that are easily ascertained through self-report questionnaires (e.g.,
when did you start smoking?, *are you a current smoker?*, and *how many cigarettes do you*
smoke in a day?). The largest genome-wide association studies (GWAS) of tobacco smoking
behaviors to date identified 140 loci associated with cigarettes per day (CPD; N = 784,353)[3].
GWAS of nicotine dependence phenotypes have generally been smaller. Some GWAS
of nicotine dependence-related phenotypes have focused on data collected from short

questionnaires such as the Fagerström Test for Nicotine Dependence (FTND[4]). The FTND is a
6-item questionnaire which includes cigarettes smoked per day as an ordinal indicator. The
largest GWAS of FTND (defined as 0–3 for mild, 4–6 for moderate, and 7–10 for severe
dependence) identified 5 loci[4]. Given the strong genome-wide genetic correlation between CPD
and FTND (r_g=0.95)[4], these GWAS were combined into a single phenotype, problematic
tobacco use (PTU), in another study[5] using multi-trait analysis of GWAS (MTAG). A recent
GWAS meta-analysis of ICD-TUD from electronic health records (EHR; see **Table 1**) reached
nearly 900,000 samples and identified 88 loci[6], all of which had been implicated in prior
smoking-related GWAS. ICD-TUD showed modest genetic correlations with cigarettes per day
($r_g = 0.44$)[6] and FTND ($r_g = 0.63$).

Table 1. Definitions of different smoking-related phenotypes discussed in this
manuscript, the acronyms used to represent them, and the relevant GWASs used in
analyses.

Smoking phenotype	Shorthand name used in current manuscript	Phenotype definition	Relevant GWAS PubMed ID
Smoking initiation	SmkInit	Whether someone has ever been a regular smoker.	36477530
Cigarettes per day	CPD	Number of cigarettes smoked per day.	36477530
Fagerström Test for Nicotine Dependence	FTND	A six-item questionnaire assessing the quantity of cigarettes smoked and compulsion to use, resulting in an ordinal measure of nicotine dependence.	33144568
Tobacco use disorder	ICD-TUD	Binary diagnosis defined using International Classification of Disease (ICD) codes (ICD 10 code for TUD: F17.200).	38632388
Nicotine dependence	DSM-NicDep	Seven criteria designed to assess clinical features of dependence, including tolerance and problems caused by using cigarettes; at least 3 of the 7 criteria must be endorsed to reach a diagnosis of nicotine dependence.	NA
Problematic tobacco use	PTU	A combination of FTND and CPD GWAS, produced using MTAG.	34750568

Our study was motivated by epidemiological and clinical data supporting nosological
distinctions between FTND and ICD or DSM-based diagnoses, including some studies that
suggest qualitative and quantitative differences in associations between DSM- and FTND-
defined nicotine dependence and some psychopathology[8,9]. The FTND is brief, and thus easily
and frequently collected. It has been especially prioritized in clinical trials of tobacco
cessation[10,11], likely because FTND scores correlate well with relapse and treatment response
and the scale places a great deal of emphasis on physiological aspects of dependence (e.g.,
items related to tolerance and withdrawal[12]). On the other hand, both ICD- and DSM-based
nicotine dependence include criteria related to physical and psychological (and social, in DSM-
5) impairment due to nicotine/tobacco use, as well as behaviors directed at seeking and using
nicotine to the exclusion of other activities. Neither FTND nor the ICD-TUD diagnostic
classification maps perfectly to DSM-NicDep[13,14].

While many large GWAS meta-analyses of substance use disorders have relied on
cases defined using clinical criteria recommended by DSM or ICD classification, the few GWAS
of DSM-NicDep to date have been relatively small[7]. We conducted a large meta-analysis of
DSM-NicDep, combining data across 16 cohorts and multiple genetic ancestries. The largest
analyses of psychiatric traits have focused on individuals whose genetics most closely resemble
the European ancestry subset of the 1000 genomes project, so we assessed the genetic
correlations between DSM-NicDep and other substance use phenotypes, psychiatric disorders,
and related phenotypes in that subset. We also compared the correlations of the various
nicotine dependence related measures (DSM-NicDep, ICD-TUD, and PTU) and other substance
use disorders (cannabis use disorder, problematic alcohol use, and opioid use disorder) by
fitting a previously published common factor model (the Addiction-risk-factor[15]) to the data using
genomic structural equation modeling. Finally, we tested whether a polygenic score for DSM-
NicDep was associated with DSM-5 tobacco use disorder and its 11 diagnostic criteria, and 4 of
the 6 FTND criteria, in a large, independent sample, the National Epidemiologic Survey on
Alcohol and Related Conditions-III (NESARC-III) cohort.
188
METHODS
Genome-wide association study of DSM-defined nicotine dependence (DSM-NicDep)
We performed meta-analyses of GWASs of DSM-defined nicotine dependence
(hereafter referred to as "DSM-NicDep") across a total of 18 cohorts, three of which included
samples of multiple ancestries, using a sample size-weighted meta-analysis implemented in
METAL[16]. We used nicotine-exposed controls where possible (see **Supplemental Materials** for
more details on each cohort). Genetic ancestry similarity was inferred by comparing an
individual's genome to the genomes from global reference populations using statistical methods
such as principal components analysis, although exact methods varied somewhat across
cohorts. There were 16 cohorts with samples that were most genetically similar to European
ancestry global reference populations, 4 with samples that were most genetically similar to
African ancestry global reference populations, and one cohort whose participants were most
genetically similar to East Asian ancestry global reference populations (hereafter referred to as
European, African, or East Asian ancestries). All GWAS controlled for age, sex, and 10 principal
components; more details, including cohort-specific covariates, are provided in **Supplemental
Table 1**. Twelve cohorts provided summary statistics, while five cohorts provided individual
genotype and phenotype data for analysis (**Supplemental Table 1**).
We used FUMA v1.5.2[17] to identify independent, genome-wide significant risk loci,
annotate variants, and perform gene-wise analyses via MAGMA[18]. We used the default FUMA
parameters to define "independent significant SNPs" as those which reached genome-wide
significance ($p < 5e-8$) and were independent of each other at r2 < 0.6, and "lead SNPs" as
those SNPs that were independent of each other at r2 < 0.1. "Genomic risk loci" were defined
by merging linkage disequilibrium (LD) blocks of independent significant SNPs within a 250 kb
distance. We performed gene mapping in FUMA using positional mapping (based on
ANNOVAR annotations), expression quantitative trait locus (eQTL) mapping (using GTEx V8[19],
CommonMind[20], and Braineac[21] data), and chromatin interaction mapping. We also performed
the MAGMA gene, gene-set, and gene expression analyses (using GTEx V8 data).

We performed follow-up analyses described below (genetic correlations, genomic
structural equation modeling, and polygenic scores) only in the European ancestry data,
because the African and East Asian ancestry subsets were under-powered.
219
Genetic correlations with other substance use disorders, psychiatric disorders, and other relevant phenotypes

We used linkage disequilibrium score regression (LDSC[22,23]) to estimate the SNP-
heritability of DSM-NicDep and the genetic correlations between DSM-NicDep and other
substance use phenotypes, using published GWASs of problematic alcohol use (PAU[24]),
Fagerström Test for Nicotine Dependence (FTND[4]), ICD-based tobacco use disorder (ICD-
TUD[6]), problematic tobacco use (PTU, a combination of FTND and cigarettes per day[15]),
cannabis use disorder (CanUD[25]), opioid use disorder (OUD[26]), smoking initiation [3], smoking
cessation [3], and cigarettes per day[3]. We also estimated genetic correlations between DSM-
NicDep and other phenotypes including psychiatric disorders, behavioral traits, respiratory
health, and socioeconomic status-related phenotypes. Details on the individual GWAS used in
genetic correlation analyses are provided in the **Supplemental Methods**. We further tested
whether genetic correlations for DSM-NicDep and FTND were different from each other using a
block-jackknife method[23,27].
234
Genomic structural equation modeling

We applied confirmatory factor analysis to the covariance matrix generated by LDSC
using genomic structural equation modeling (genomic SEM[28]) with weighted least squares
estimation. As in Hatoum et al.[5], the indicators were allowed to load freely on a single latent
factor (*Addiction-Risk-Factor*), but updated the OUD[26], PAU[24], and CanUD[25] GWAS. We
compared *Addiction-Risk-Factor* models with DSM-NicDep, PTU, or ICD-TUD as the tobacco-
related indicator. The variance of the common latent factor was scaled to 1.0.
242
Polygenic score analyses

We created polygenic scores for DSM-NicDep and FTND in the European ancestry
subset of the NESARC-III sample. NESARC-III was genotyped using the Affymetrix Axiom®
Exome Array[29], which limited our ability to impute SNPs due to lack of appreciable non-exonic
coverage and resulted in some regions with low SNP densities. Details of QC and imputation
are available in the **Supplemental Materials.** We used PRS-CS[30], a Bayesian method that
uses continuous shrinkage priors to weight SNP effect sizes, and used the 'auto' function, which
allows the global scaling parameter to be automatically learned from the data. We then used the
'score' function in Plink 1.9[31] to create polygenic scores for DSM-NicDep and FTND in
NESARC-III. The associations between DSM-5 tobacco use disorder, endorsement of the 11
individual diagnostic criteria, and 4 of the 6 FTND criteria (excluding cigarettes per day and
smoking when ill) in NESARC-III (N = 12,482; N_{cases} = 4,205) and the polygenic scores for DSM-
NicDep and FTND were estimated using logistic regression models in R, covarying for age, sex,
and 10 within-ancestry principal components. Both polygenic scores were included jointly as
predictors in the same model.
258

RESULTS

Genome-wide association study of DSM-defined nicotine dependence

There were a total of 15 cohorts included in the European ancestry (EUR) meta-analysis (N_{cases} = 20,923, $N_{controls}$ = 26,961), four cohorts included in the African ancestry (AFR) meta-analysis (N_{cases} = 5,293, $N_{controls}$ = 4,938), and one cohort included in the East Asian ancestry (EAS) GWAS (N_{cases} = 2,007, $N_{controls}$ = 1,739), for a total cross-ancestry sample size of N = 61,861 (28,223 cases; **Supplemental Table 1**). We identified one genome-wide significant locus, near the cholinergic receptor nicotinic alpha 5 subunit (*CHRNA5*) gene on chromosome 15 (lead SNP: rs147144681, p-value =1.27E-11 in European ancestry; lead SNP = rs2036527, p-value = 6.49e-13 in cross-ancestry analysis; **Supplemental Tables 2 & 3**). Using a lifetime prevalence of 24%[32], we estimated the liability scale SNP-heritability of DSM-defined nicotine dependence (DSM-NicDep) to be 0.07 (SE = 0.01) in the European ancestry samples.

In the MAGMA gene-based analysis of the European ancestry data, three genes were significant after multiple testing corrections: *CHRNA5, CHRNA3,* and *IREB2* (**Supplemental Table 4**). Three gene-sets were significantly associated: "GOBP_TRANSCRIPTION_BY_RNA_POLYMERASE_III", "GOMF_PRE_MRNA_5_SPLICE_SITE_BINDING", and "REACTOME_HIGHLY_CALCIUM_PERMEABLE_NICOTINIC_ACETYLCHOLINE_RECEPTORS". None of the 30 tissue types from GTEx v8 were significant.

Genetic correlations with substance use and other phenotypes

In the EUR data, DSM-NicDep was strongly correlated with the other measures of nicotine dependence (r_gs with FTND, PTU, and ICD-TUD ranged from 0.77 to 1.01). The genetic correlations between DSM-NicDep and published GWASs of substance use disorders were of moderate-to-high magnitude: cannabis use disorder (CanUD), problematic alcohol use (PAU), and opioid use disorder (OUD; r_g = 0.64 – 0.84). Overall, DSM-NicDep was significantly correlated with 23 of the 26 phenotypes tested (**Figure 1; Supplemental Table 5**). Compared to the other tobacco-related phenotypes, DSM-NicDep showed the strongest correlations with many traits, albeit with wider confidence intervals due to smaller sample size. When correcting for 24 comparisons, the genetic correlations between DSM-NicDep and smoking initiation, cannabis ever-use, CanUD, and the Townsend deprivation index were significantly larger (p < 0.002) than the corresponding genetic correlations between FTND and these phenotypes.

Figure 1. Comparing genetic correlations (r_g) for DSM-NicDep, FTND, ICD-TUD, and PTU
with other traits in European ancestry data. Traits include other substance use disorders
(CanUD = cannabis use disorder[25]; OUD = opioid use disorder[26]; PAU = problematic alcohol
use[24], ICD-TUD = ICD-based tobacco use disorder[6]), substance use behaviors (CanUse =
cannabis ever-use[33]; DPW = drinks per week[3]; SmkInit = smoking initiation[3], SmkCessation =
smoking cessation[3], CPD = cigarettes per day[3]), psychiatric disorders and other mental health

phenotypes (ADHD = attention deficit hyperactivity disorder[34]; PTSD = post-traumatic stress
disorder[35]), biomarkers (Cot+HC = cotinine + 3-hydroxycotinine[36]), lung health-related traits
(FEV1 = forced expiratory volume in 1 second), risk tolerance[37], socioeconomic status-related
traits (Edu attainment = educational attainment[38]; TDI = Townsend deprivation index), executive
function (EF[39]), and anthropometric measures (BMI = body mass index[40]; height[41]). * indicates
r_gs that significantly differ between DSM-NicDep and FTND at   = 0.002 (Bonferroni correction
for 24 comparisons).
305
Genomic structural equation models of broad addiction liability
In the EUR data, we found that a common genetic factor model with DSM-NicDep, PAU,
OUD, and CanUD as indicators, similar to the *Addiction-Risk-Factor* presented in Hatoum et
al.[4], fit the data well (**Figure 2A; Supplemental Table 6**). There were two differences between
the Hatoum et al. *Addiction-Risk-Factor* and our modified model: (1) we used larger, more
recent versions of the OUD[26], PAU[24], and CanUD[25] GWASs; and (2) we removed the residual
correlation between PAU and OUD, as this path was no longer significant. We compared the
model with DSM-NicDep as the tobacco-related indicator (**Figure 2A**) to one where DSM-
NicDep was substituted by the original problematic tobacco use (PTU) GWAS meta-analysis
(FTND + CPD) from Hatoum et al.[4] (**Figure 2B; Supplemental Table 7**), and one where
NicDep was substituted by ICD-TUD[6] (**Figure 2C; Supplemental Table 8**). Each of the
modified models fit the data well, but the loading for DSM-NicDep was the largest of the three
tobacco-related indicators, almost 3-fold larger than the loading for PTU in the equivalent model
(0.86 vs. 0.30).
320
321 **A.**

322
323 **B.**

324
325
326 **C.**

327
328
Figure 2. A modified Addiction-Risk-Factor model. This model is patterned upon the
common factor model in Figure 1A of Hatoum *et al.*, 2022, but updated with new, larger versions
of the OUD[26], PAU[24,25], and CanUD GWAS and using 3 different phenotypes for tobacco
GWAS. **Panel A:** DSM-NicDep. **Panel B:** PTU[5] GWAS. **Panel C:** ICD-TUD[6]. Significant
loadings at $p < 0.05$ are represented by *. Addiction-rf = The Addiction-Risk-Factor; OUD =
opioid use disorder; PAU = problematic alcohol use; CanUD = cannabis use disorder; DSM-
NicDep = nicotine dependence; ICD-TUD = ICD-based tobacco use disorder.
336

Polygenic scores for DSM-NicDep and FTND

A polygenic score (PGS) for DSM-NicDep was significantly associated with DSM-5 tobacco use disorder in the European ancestry subset of the NESARC-III sample (beta = 0.07, SE = 0.02, p-value = 5.11e-3; N_{cases} = 4,205), as was a polygenic score for FTND (beta = 0.07, SE = 0.02, p-value = 3.11e-3). Results from item-level logistic regression models are shown in **Figure 3.** The DSM-NicDep PGS was associated with 6 of 11 criteria, as was the FTND PGS, although only 3 of these criteria were associated with both (TimeSpent, Problems, and Withdrawal). The FTND PGS was associated with 3 of the 4 FTND items, as expected, while the DSM-NicDep PGS was not associated with any FTND items. Failure to fulfill major role obligations (Fail) and Use of larger amounts or for longer than intended (Larger) were unrelated to both PGS after multiple testing correction. Despite these apparent disparities, estimates were not significantly different for the DSM-NicDep and FTND PGS.

Figure 3. Polygenic scores (PGS) for FTND and DSM-NicDep predict individual DSM-5 nicotine use disorder and FTND criteria in the European ancestry subset of the NESARC-III sample. Filled circles represent estimates that were significant after FDR correction, while open circles represent estimates that were not significant after FDR correction. Hazardous = Recurrent use in physically hazardous situations; Fail = Recurrent use resulting in failure to fulfill major role obligations at work, school or home; Tolerance = Marked need for increased amount to get same effect, or diminished effect of same amount; TimeSpent = Great deal of time spent in activities necessary to obtain or use; GiveUp = Important recreational, social or occupational activities given up or reduced; Problems = Use despite knowledge of persistent/recurrent physical/psychological problems; Larger = Taken over larger amounts/longer periods than intended; Withdrawal = Withdrawal syndrome or use to relieve/avoid syndrome; Cutdown = Persistent desire or unsuccessful attempts to cutdown or control use; Crave = Craving, or

strong urge or desire to use; Social = Persistent use despite recurring social/interpersonal
problems caused or exacerbated by use; FTND1_within30min = How soon after you wake up
do you smoke your first cigarette?; FTND2_prohibited = Do you find it difficult to refrain from
smoking in places where it is forbidden?; FTND3_morning = Which cigarette would you hate
most to give up?; FTND5_waking = Do you smoke more frequently during the first hours after
waking than during the rest of the day?
369
370
DISCUSSION
In this first cross-ancestry GWAS of DSM-NicDep (N = 61,861), we replicated the well-
known *CHRNA5* risk locus. DSM-NicDep was genetically correlated with other substance use
disorders, smoking-related phenotypes, and psychiatric disorders. DSM-NicDep was more
closely related to the genetic underpinnings of other DSM- or ICD-defined substance use
disorders than was PTU (problematic tobacco use, a phenotype that combined FTND and
cigarettes per day).
ICD and DSM criteria focus on physical and psychological (and social, in DSM-5)
impairment due to substance use, and are, for the most part, identical across substances.
FTND, on the other hand, places greater emphasis on physiological aspects of dependence
(e.g., items related to tolerance and withdrawal[12]). Because most GWAS of substance use
disorders utilize ICD and DSM criteria, we hypothesized that this discrepancy might underlie the
lower genetic correlation observed between nicotine dependence and other substance use
disorders in prior twin and genome-wide correlation studies that used FTND[6]. Indeed, the
loading for DSM-NicDep in the Addiction-Risk-Factor genomic SEM was nearly 3 times that of
the loading for PTU (0.83 vs. 0.30) (**Figure 1**). Notably, the loadings for OUD and PAU also
increased in our model (0.83 to 1.0, and 0.58 to 0.81, respectively), likely due to the larger
sample sizes of these more recent GWAS. We also noted no significant residual correlation
between OUD and PAU in our analysis.
Polygenic score (PGS) analyses of individual criteria in NESARC-III revealed an
interesting pattern of associations. First, none of the FTND-specific items were correlated with
the DSM-NicDep PGS, suggesting that the DSM-NicDep PGS captured genetic variance that
was distinct from the FTND items. Second, even though DSM-IV did not include a diagnosis of
nicotine abuse, DSM-5 includes 3 abuse criteria in its diagnosis of Tobacco Use Disorder – of
these, two were unrelated to the DSM-NicDep PGS (Social, Fail) while the third (Hazard) was
significantly associated. Notably, Fail was unrelated to both PGS and was amongst the least
commonly endorsed criteria. Third, while the FTND is presumed to evaluate aspects of both
tolerance and withdrawal, the DSM Tolerance criterion was unrelated to the FTND PGS. In
contrast, Withdrawal as assessed by the DSM was associated with both PGS, while Time to first
cigarette was solely associated with FTND. Taken together, these item-level analyses suggest
that the DSM-NicDep GWAS may have indexed genetic liability to a distinct set of tobacco-
related problems than the FTND GWAS. However, many confidence intervals on these
estimates were wide. Future item-level GWAS using novel structural equation modeling
methods that bring together DSM and FTND items might be of high value at parsing whether
genetic and clinical heterogeneity align (**Figure 3**).

We expected that DSM-NicDep would be more genetically correlated with
psychopathology than FTND, as has been found in clinical and epidemiological studies[8]. The
point estimates of our genetic correlation analysis suggested that PTU (CPD + FTND) was the
least associated with most indices of psychopathology, while both DSM-NicDep and ICD-TUD
were more strongly associated with psychosocial indices. DSM-NicDep appeared to additionally
index material deprivation more strongly than the other traits, while all four tobacco GWAS
(DSM-NicDep, ICD-TUD, FTND and PTU) were equivalently related to respiratory (e.g., FEV1,
Lung cancer) and metabolic markers of tobacco exposure.
The largest GWAS of CPD[3] identified 140 genomic risk loci, compared with 5 in the
largest GWAS of FTND[4] and 1 in the current study of DSM-NicDep. This likely reflects the much
larger sample size in the CPD study. Despite a similar sample size, the FTND GWAS identified
more loci than the current DSM-NicDep GWAS, possibly because that study used a linear
model to analyze a categorical 3-level (mild-moderate-severe) measure while our analyses
relied on binary diagnoses. A recent GWAS of ICD-TUD that had a much larger sample size
than ours (N = 898,680) identified 88 loci[6]. In the current study, DSM-NicDep was highly
correlated with ICD-TUD. Given their high genetic correlation, future efforts may consider
combining these diagnostic modalities as sources of information on tobacco use disorder.
We did not find any significant loci in the much smaller African or East Asian ancestry
analyses, due to the lack of power. The lack of diversity in genetic data is unfortunate as
nicotine dependence and tobacco use disorder are leading contributors to mortality in worldwide
populations[2,42]. A related limitation was our inability to conduct sex- and birth cohort-stratified
analyses; rates of tobacco use vary markedly according to both sex and birth cohort, which may
in turn modify the extent to which genetic liability influences risk for DSM-NicDep. Amassing
more genetic datasets with DSM-derived nicotine dependence would enable such valuable
analyses.
In summary, our analyses highlight the importance of considering diagnostic assessment
in genetic studies of substance use disorders. We found that DSM-NicDep was more closely
related to a general addiction liability factor compared to a "problematic tobacco use" phenotype
that combined FTND and CPD. The DSM-NicDep PGS was not significantly associated with
FTND criteria in an independent sample. Compared to FTND, DSM-NicDep was more strongly
genetically correlated with ICD-TUD, smoking initiation, cannabis ever-use, and material
deprivation. Given the strong genetic correlation between DSM-NicDep and ICD-TUD, future
analyses may consider combining data from DSM-NicDep and ICD-based studies of TUD to
maximize sample size for gene discovery.
440

441   **Author Contributions**
442   ECJ, DL, HJE, JG, and AA conceived the study idea. ECJ, DL, APM, ASH, JDD, JVB, MG, KS,
443   TT, TE, PJ, TK, CM, YN, TP, MS, PNRV, LW, and IRG contributed to data analysis. ECJ, DL,
444   and AA drafted the manuscript, and all authors edited, reviewed, and approved the final
445   manuscript.
446

Funding
We acknowledge the following sources of support:
K01DA051759 (ECJ); T32DA007261 (JVB); K01DA058807 (JDD); K01AA031724 (APM);
R01DA054869 (AA, HJE, JG); DP1DA054394, T32IR5226 (SSR); R01DA042755,

U01DA042217, R01AG077742, R01DA054087, R01DA044283, DA05147, DA13240, DA02441,
AA09367, AA11886, MH066140 (TE, AG, EAW, SV, MM, JJL); R01DA030976 (KCW, IRG);
R01MH100141 (PNRV); R01MH123489, R01MH123619, R01MH134284 (ARD); AUS NHMRC
464914 (IBH, NGM); R00DA023549 (NAG); HSRI 67-095 (VS). LD acknowledges support from
NHMRC Investigator Grant L3 (2016825), NHMRC Senior Principal Research Fellowship
(1135991), the National Drug and Alcohol Research Centre (NDARC), UNSW funded by the
Australian Government Department of Health. The views expressed in this publication do not
necessarily represent the position of the Australian Government.
The AGDS was primarily funded by National Health and Medical Research Council (NHMRC) of
Australia (Grant No. APP1086683) to N.G.M. This work was further supported by NHMRC
grants (No. 1145645, 1078901 and 1087889). N.G.M. is supported by a NHMRC Investigator
Grant (No. APP1172990).
GBP: Data collection was funded and data analysis was supported by the Australian National
Health and Medical Research Council (No. APP1138514) to S.E.M. S.E.M. is supported by a
National Health and Medical Research Council Investigator Grant (No. APP1172917).
The Psychiatric Genomics Consortium's Substance Use Disorders (PGC-SUD) Working Group
receives support from the National Institute on Drug Abuse via R01DA054869. We gratefully
acknowledge prior support from the National Institute on Alcohol Abuse and Alcoholism and
support from National Institutes of Mental Health to the overall PGC. Statistical analyses were
carried out on the NL Genetic Cluster Computer (http://www.geneticcluster.org) hosted by
SURFsara.
472

Acknowledgements

We gratefully acknowledge our contributing studies and the participants in those studies without
whom this effort would not be possible.
AGDS: We thank all the participants for giving their time to contribute to this study. We wish to
thank all the people who helped in the conception, implementation, beta testing, media
campaign and data cleaning. We would specifically like to acknowledge Ken Kendler, Patrick
Sullivan, Andrew McIntosh and Cathryn Lewis for input on the questionnaire; Lorelle Nunn,
Mary Ferguson, Lucy Winkler and Natalie Garden for data and sample collection; Natalia
Zmicerevska, Alissa Nichles and Candace Brennan for participant recruitment support.
Jonathan Davies, Luke Lowrey and Valeriano Antonini for support with IT aspects; Vera Morgan
and Ken Kirkby for help with the media campaign.
GBP: We thank the participants for giving their time and support for this project. We
acknowledge and thank M. Steffens for her generous donations in loving memory of J. Banks.
Funding support for the **Comorbidity and Trauma Study (CATS)** (dbGAP accession number:
phs000277.v1.p1) was provided by the National Institute on Drug Abuse (R01 DA17305);
GWAS genotyping services at the CIDR at The Johns Hopkins University were supported by the
National Institutes of Health (contract N01-HG-65403). The **Christchurch Health and
Development Study (CHDS**: dbGAP in progress) has been supported by funding from the
Health Research Council of New Zealand, the National Child Health Research Foundation
(Cure Kids), the Canterbury Medical Research Foundation, the New Zealand Lottery Grants
Board, the University of Otago, the Carney Centre for Pharmacogenomics, the James Hume
Bequest Fund, US NIH grant MH077874, and NIDA grant ''A developmental model of gene-

environment interplay in SUDs'' (R01DA024413) 2007–2012. The **Collaborative Study on the**
Genetics of Alcoholism (COGA), Principal Investigators B. Porjesz, V. Hesselbrock, A.
Agrawal; Scientific Director, A. Agrawal; Translational Director, D. Dick, includes ten different
centers: University of Connecticut (V. Hesselbrock); Indiana University (H.J. Edenberg, T.
Foroud, Y. Liu, M.H. Plawecki); University of Iowa Carver College of Medicine (S. Kuperman, J.
Kramer); SUNY Downstate Health Sciences University (B. Porjesz, J. Meyers, C. Kamarajan, A.
Pandey); Washington University in St. Louis (L. Bierut, J. Rice, K. Bucholz, A. Agrawal, S.
Hartz); University of California at San Diego (M. Schuckit); Rutgers University (J. Tischfield, D.
Dick, R. Hart, J. Salvatore); The Children's Hospital of Philadelphia, University of Pennsylvania
(L. Almasy); Icahn School of Medicine at Mount Sinai (A. Goate, P. Slesinger); and Howard
University (D. Scott). Other COGA collaborators include: L. Bauer (University of Connecticut); J.
Nurnberger Jr., L. Wetherill, X., Xuei, D. Lai, S. O'Connor, (Indiana University); G. Chan
(University of Iowa; University of Connecticut); D.B. Chorlian, J. Zhang, P. Barr, S. Kinreich, G.
Pandey, Z. Neale (SUNY Downstate); N. Mullins (Icahn School of Medicine at Mount Sinai); A.
Anokhin, S. Hartz, E. Johnson, V. McCutcheon, S. Saccone (Washington University); J. Moore,
F. Aliev, Z. Pang, S. Kuo (Rutgers University); A. Merikangas (The Children's Hospital of
Philadelphia and University of Pennsylvania); H. Chin and A. Parsian are the NIAAA Staff
Collaborators. We continue to be inspired by our memories of Henri Begleiter and Theodore
Reich, founding PI and Co-PI of COGA, and also owe a debt of gratitude to other past
organizers of COGA, including Ting- Kai Li, P. Michael Conneally, Raymond Crowe, and Wendy
Reich, for their critical contributions. This national collaborative study is supported by NIH Grant
U10AA008401 from the National Institute on Alcohol Abuse and Alcoholism (NIAAA) and the
National Institute on Drug Abuse (NIDA). Support for the **Study of Addiction: Genetics and**
Environment (SAGE) was provided through the NIH Genes, Environment and Health Initiative
[GEI; U01 HG004422; dbGaP study accession phs000092.v1.p1]. SAGE is one of the genome-
wide association studies funded as part of the Gene Environment Association Studies
(GENEVA) under GEI. Assistance with phenotype harmonization and genotype cleaning, as
well as with general study coordination, was provided by the GENEVA Coordinating Center
[U01 HG004446]. Assistance with data cleaning was provided by the National Center for
Biotechnology Information. Support for collection of datasets and samples was provided by the
Collaborative Study on the Genetics of Alcoholism [COGA; U10 AA008401], the Collaborative
Genetic Study of Nicotine Dependence [COGEND; P01 CA089392; see also phs000404.v1.p1],
and the Family Study of Cocaine Dependence [FSCD; R01 DA013423, R01 DA019963].
Funding support for genotyping, which was performed at the Johns Hopkins University Center
for Inherited Disease Research (CIDR), was provided by the NIH GEI [U01HG004438], the
National Institute on Alcohol Abuse and Alcoholism, the National Institute on Drug Abuse, and
the NIH contract "High throughput genotyping for studying the genetic contributions to human
disease" [HHSN268200782096C]. The **GSMS** project (phs000852.v1.p1) was supported by the
National Institute on Drug Abuse (U01DA024413, R01DA11301), the National Institute of Mental
Health (R01MH063970, R01MH063671, R01MH048085, K01MH093731 and K23MH080230),
NARSAD, and the William T. Grant Foundation. We are grateful to all the GSMS and CCC study
participants who contributed to this work. The following grants supported data collection and
analysis of **CADD** (dbGAP in progress): DA011015, DA012845, DA021913, DA021905,
DA032555, and DA035804. **ADAA** was funded by NIH grant R01 AA017444. **Brisbane**

Longitudinal Twin Study (BLTS) was supported by the United States National Institute on
Drug Abuse (R00DA023549), and by the Australian Research Council (DP0343921,
DP0664638, 464914, 619667, FT110100548). **Yale-Penn** (phs000425.v1.p1; phs000952.v1.p1)
was supported by National Institutes of Health Grants RC2 DA028909, R01 DA12690, R01
DA12849, R01 DA18432, R01 AA11330, and R01 AA017535 and the Veterans Affairs
Connecticut and Philadelphia Veterans Affairs Mental Illness Research, Educational, and
Clinical Centers. **Australian Alcohol and Nicotine studies (OZ-ALC-NAG**; phs000181.v1.p1)
were supported by National Institutes of Health Grants AA07535,AA07728, AA13320, AA13321,
AA14041, AA11998, AA17688, DA12854, and DA019951; by Grants from the Australian
National Health and Medical Research Council (241944, 339462, 389927,389875, 389891,
389892, 389938, 442915, 442981, 496739, 552485,and 552498); by Grants from the Australian
Research Council(A7960034, A79906588, A79801419, DP0770096, DP0212016, and
DP0343921); and by the 5th Framework Programme (FP-5) GenomEUtwin Project (QLG2-CT-
2002-01254). Genome-wide association study genotyping at Center for Inherited Disease
Research was supported by a Grant to the late Richard Todd, M.D., Ph.D., former Principal
Investigator of Grant AA13320. The **Finnish Twin Cohort/Nicotine Addiction Genetics-
Finland** study was supported by Academy of Finland (grants # 213506, 129680 to JK), NIH
DA12854 (PAFM), Global Research Award for Nicotine Dependence / Pfizer Inc. (JK),
Wellcome Trust Sanger Institute, UK and the European Community's Seventh Framework
Programme ENGAGE Consortium (HEALTH-F4-2007- 201413). In **Finntwin12**, support for data
collection and genotyping has come from National Institute of Alcohol Abuse and Alcoholism
(grants AA-12502, AA-00145, and AA-09203 to RJR and AA15416 and K02AA018755 to DMD),
the Academy of Finland (grants 100499, 205585, 118555, 141054 and 264146 to JK) &
Wellcome Trust Sanger Institute, UK.

563
564
References

1.   MAES, H. H. *et al.* A twin study of genetic and environmental influences on tobacco
initiation, regular tobacco use and nicotine dependence. *Psychol Med* **34**, 1251–1261
(2004).
2.   Le Foll, B. *et al.* Tobacco and nicotine use. *Nat Rev Dis Primers* **8**, 19 (2022).
3.   Saunders, G. R. B. *et al.* Genetic diversity fuels gene discovery for tobacco and alcohol
use. *Nature* **612**, 720–724 (2022).
4.   Quach, B. C. *et al.* Expanding the genetic architecture of nicotine dependence and its
shared genetics with multiple traits. *Nat Commun* **11**, 5562 (2020).
5.   Hatoum, A. S. *et al.* The addiction risk factor: a unitary genetic vulnerability characterizes
substance use disorders and their associations with common correlates.
Neuropsychopharmacology **47**, 1739–1745 (2022).
6.   Toikumo, S. *et al.* Multi-ancestry meta-analysis of tobacco use disorder identifies 461
potential risk genes and reveals associations with multiple health outcomes. *Nat Hum
Behav* **8**, 1177–1193 (2024).
7.   Loukola, A. *et al.* Genome-wide association study on detailed profiles of smoking
behavior and nicotine dependence in a twin sample. *Mol Psychiatry* **19**, 615–624 (2014).

8.   Breslau, N. & Johnson, E. O. Predicting smoking cessation and major depression in
nicotine-dependent smokers. *Am J Public Health* **90**, 1122–1127 (2000).

9.   Brook, J. S., Koppel, J. & Pahl, K. Predictors of DSM and Fagerström-Defined Nicotine
Dependence in African American and Puerto Rican Young Adults. *Subst Use Misuse* **44**,
809–822 (2009).

10. Tashkin, D. P. *et al.* Effects of Varenicline on Smoking Cessation in Patients With Mild to
Moderate COPD: A Randomized Controlled Trial. *Chest* **139**, 591–599 (2011).

11. Ramon, J. M., Morchon, S., Baena, A. & Masuet-Aumatell, C. Combining varenicline and
nicotine patches: a randomized controlled trial study in smoking cessation. *BMC Med* **12**,
1–9 (2014).

12. Baker, T. B., Breslau, N., Covey, L. & Shiffman, S. DSM criteria for tobacco use disorder
and tobacco withdrawal: a critique and proposed revisions for DSM-5. *Addiction* **107**,
263–275 (2012).

13. Moolchan, E. T. *et al.* The Fagerstrom Test for Nicotine Dependence and the Diagnostic
Interview Schedule: Do they diagnose the same smokers? *Addictive Behaviors* **27**, 101–
113 (2002).

14. Mwenifumbo, J. C. & Tyndale, R. F. DSM-IV, ICD-10 and FTND: discordant tobacco
dependence diagnoses in adult smokers. *J Addict Res Ther* **2**, 105 (2011).

15. Hatoum, A. S. *et al.* Multivariate genome-wide association meta-analysis of over 1 million
subjects identifies loci underlying multiple substance use disorders. *Nature Mental Health*
1, 210–223 (2023).

16. Willer, C. J., Li, Y. & Abecasis, G. R. METAL: fast and efficient meta-analysis of
genomewide association scans. *Bioinformatics* **26**, 2190–2191 (2010).

17. Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and
annotation of genetic associations with FUMA. *Nat Commun* **8**, 1826 (2017).

18. de Leeuw, C. A., Mooij, J. M., Heskes, T. & Posthuma, D. MAGMA: Generalized Gene-
Set Analysis of GWAS Data. *PLoS Comput Biol* **11**, (2015).

19. Aguet, F. *et al.* Genetic effects on gene expression across human tissues. *Nature* **550**,
204–213 (2017).

20. Huckins, L. M. *et al.* Gene expression imputation across multiple brain regions provides
insights into schizophrenia risk. *Nat Genet* **51**, 659–674 (2019).

21. Trabzuni, D. *et al.* Quality control parameters on a large dataset of regionally dissected
human control brains for whole genome expression studies. *J Neurochem* **119**, 275–282
(2011).

22. Bulik-Sullivan, B. K. *et al.* LD Score regression distinguishes confounding from
polygenicity in genome-wide association studies. *Nat Genet* **47**, 291–295 (2015).

23. Bulik-Sullivan, B. K. *et al.* An atlas of genetic correlations across human diseases and
traits. *Nat Genet* **47**, 1236 (2015).

24. Zhou, H. *et al.* Multi-ancestry study of the genetics of problematic alcohol use in over 1
million individuals. *Nat Med* **29**, 3184–3192 (2023).

25. Levey, D. F. *et al.* Multi-ancestry genome-wide association study of cannabis use
disorder yields insight into disease biology and public health implications. *Nat Genet* **55**,
2094–2103 (2023).

26.   Deak, J. D. *et al.* Genome-wide association study in individuals of European and African
ancestry and multi-trait analysis of opioid use disorder identifies 19 independent genome-
wide significant risk loci. *Mol Psychiatry* **27**, 3970–3979 (2022).
27.   Coleman, J. R. I. *et al.* Genome-wide gene-environment analyses of major depressive
disorder and reported lifetime traumatic experiences in UK Biobank. *Mol Psychiatry* **25**,
1430–1446 (2020).
28.   Grotzinger, A. D. *et al.* Genomic structural equation modelling provides insights into the
multivariate genetic architecture of complex traits. *Nat Hum Behav* **3**, 513–525 (2019).
29.   Zhang, H. *et al.* Strong and weak cross-inheritance of substance use disorders in a
nationally representative sample. *Mol Psychiatry* **27**, 1742–1753 (2022).
30.   Ge, T., Chen, C.-Y., Ni, Y., Feng, Y.-C. A. & Smoller, J. W. Polygenic prediction via
Bayesian regression and continuous shrinkage priors. *Nat Commun* **10**, 1–10 (2019).
31.   Chang, C. C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer
datasets. *Gigascience* **4**, 7 (2015).
32.   Breslau, N., Johnson, E. O., Hiripi, E. & Kessler, R. Nicotine Dependence in the United
States: Prevalence, Trends, and Smoking Persistence. *Arch Gen Psychiatry* **58**, 810–816
(2001).
33.   Pasman, J. A. *et al.* GWAS of lifetime cannabis use reveals new risk loci, genetic overlap
with psychiatric traits, and a causal influence of schizophrenia. *Nat Neurosci* **21**, 1161–
1170 (2018).
34.   Demontis, D. *et al.* Genome-wide analyses of ADHD identify 27 risk loci, refine the
genetic architecture and implicate several cognitive domains. *Nat Genet* **55**, 198–208
(2023).
35.   Nievergelt, C. M. *et al.* International meta-analysis of PTSD genome-wide association
studies identifies sex- and ancestry-specific genetic risk loci. *Nat Commun* **10**, 4558
(2019).
36.   Buchwald, J. *et al.* Genome-wide association meta-analysis of nicotine metabolism and
cigarette consumption measures in smokers of European descent. *Mol Psychiatry* **26**,
2212–2223 (2021).
37.   Linnér, R. K. *et al.* Genome-wide association analyses of risk tolerance and risky
behaviors in over 1 million individuals identify hundreds of loci and shared genetic
influences. *Nat Genet* **51**, 245 (2019).
38.   Lee, J. J. *et al.* Gene discovery and polygenic prediction from a genome-wide association
study of educational attainment in 1.1 million individuals. *Nat Genet* **50**, 1112–1121
(2018).
39.   Hatoum, A. S. *et al.* Genome-wide association study shows that executive functioning is
influenced by GABAergic processes and is a neurocognitive genetic correlate of
psychiatric disorders. *Biol Psychiatry* **93**, 59–70 (2023).
40.   Yengo, L. *et al.* Meta-analysis of genome-wide association studies for height and body
mass index in ~700000 individuals of European ancestry. *Hum Mol Genet* **27**, 3641–
3649 (2018).
41.   Yengo, L. *et al.* A saturated map of common genetic variants associated with human
height. *Nature* **610**, 704–712 (2022).

42.   Reitsma, M. B. *et al.* Spatial, temporal, and demographic patterns in prevalence of
smoking tobacco use and attributable disease burden in 204 countries and territories,
1990–2019: a systematic analysis from the Global Burden of Disease Study
2019. *The Lancet* **397**, 2337–2360 (2021).