Diarrheal disease attributed to *Shigella* spp. and Enteroinvasive *Escherichia coli* among children at households in rural Haïti: A case-control study

Short title: Diarrheal disease attributed to *Shigella* spp./Enteroinvasive *Escherichia coli*

Authors
Ishae Sriguha[1,2*†], Saradiya R Kuyt[2†], Youseline Cajusma[2], Emilee Cato[2], Lindsey Brinkley[2], Md Abu Sayeed[1,2], Stace Maples[3], Valery Madsen Beau De Rochars[4], Daniel T Leung[5], Anthony T Maurelli[1], Chantale Baril[6], Molly B Klarman[1,2], Eric J Nelson[1,2*]

Affiliations
[1] Department of Environmental and Global Health, University of Florida, Gainesville, FL,
[2] Department of Pediatrics, University of Florida, Gainesville, FL
[3] Stanford Geospatial Center, Stanford University, Stanford, CA
[4] Department of Health Services Research, Management and Policy, University of Florida, Gainesville, FL
[5] Department of Medicine, University of Utah, Salt Lake City, UT
[6] Université d'État d'Haïti- Faculté de Médecine et de Pharmacie, Port-au-Prince, Haïti

† **Co-first authors**

* **Corresponding authors**
Ishae Sriguha (isriguha@ufl.edu), Eric Nelson (eric.nelson@ufl.edu)

Keywords: shigella, shigellosis, EIEC, diarrhea, diarrhoea, GEMS, MAL-ED, Haïti

ABSTRACT

Infections from *Shigella* spp. and Enteroinvasive *Escherichia coli* (EIEC) are considered leading causes of symptomatic diarrheal disease, globally. However, there is a paucity of case-control studies from Caribbean nations to guide regional public health priorities and interventions. A case-control study was conducted within a larger cross-sectional healthcare study in Haïti. Participant households were identified using a geospatially randomized method; families with children under 5 years were consented and enrolled. Rectal swabs from child participants were tested for *Shigella* spp./EIEC by qPCR using the *ipaH* target. Two case-definitions were used: 'diarrheal symptom' (DS) cases were defined as those reporting diarrheal symptoms ≤7 days ago; 'acute diarrhea' (AD) cases were defined as those reporting diarrheal symptoms ≤7 days ago with ≥3 loose stools in the past 24 hours and onset <7 days ago. Of 868 households screened, 568 were enrolled with 794 participating children; samples from 732 children were analyzed. Rates of *Shigella* spp./EIEC. detection among DS cases and controls were 11.4% (22/193) and 6.1% (33/539), respectively. Rates of detection among AD cases and controls were 18.6% (8/43) and 6.8% (47/689), respectively. The adjusted odds of having DS increased by 84% (aOR=1.84; 95%CI 1.02 to 3.27) and having AD increased by 183% (aOR=2.83; 95%CI 1.14 to 6.36) when *Shigella* spp./EIEC was detected. The attributable fractions for DS and AD were 5.62% (95%CI 0.44% to 10.9%) and 12.6% (95%CI 0% to 25.3%), respectively. Rates of bloody diarrhea (dysentery) were minimal (<1%, 6/732). Within this case-control study in Haïti, *Shigella* spp./EIEC detection was common and attributed to symptomatic disease. These results align with prior global health studies. *Shigella* spp./EIEC represent an important public health target for intervention once the security situation in Haïti stabilizes.

AUTHOR SUMMARY

Global studies outside the Caribbean have identified a group of closely related enteric

pathogens (*Shigella* spp., Enteroinvasive *Escherichia coli* (EIEC)) as a leading cause of

diarrheal disease. This group can be associated with short-term (e.g., bloody diarrhea)

or long-term (e.g., malnutrition, stunting) health impacts. Within the Caribbean, there is

a lack of information to guide public health policy on enteric pathogens, especially in

Haïti. Therefore, we conducted a case-control study where we identified households

with children under 5 years, gathered data on diarrhea symptoms, and collected stool

swabs. We tested the stools using a sensitive molecular assay (*ipaH* qPCR). *Shigella*

spp./EIEC detection was common among both children with and without diarrhea, yet

more common among, and attributed to, those with symptomatic diarrheal disease.

These results align with prior global health studies and represent an opportunity for

high-impact public health action when the security situation in Haïti stabilizes.

INTRODUCTION

Diarrheal diseases are the second leading cause of death among children between one month and 5 years of age worldwide with at least 1.1 billion cases and 446,000 deaths annually (1). Morbidity has both acute components and more complex bi-directional correlates that include malnutrition, stunting, and neuro-cognitive delay (2-5). *Shigella* spp./EIEC are a group of closely related bacterial pathogens estimated to cause 111 million cases and 63,100 deaths annually among children under 5 years (5-8).

The pathophysiologic consequences of infection vary. Upon fecal-oral transmission, *Shigella* spp. infection may manifest in symptomatic diarrheal disease with or without blood in the stool (9). Fluid loss leading to dehydration occurs at the level of the small intestine (9). Invasive disease leading to enterocolitis generally occurs at the level of the large intestine in a subset of patients (10). While shigellosis can be self-limited, antibiotics are indicated for severe disease and vulnerable sub-populations (11). The infectious dose can be as low as 10-100 bacilli which likely contributes to large outbreaks when hygiene and sanitation infrastructure are compromised (7, 12). The Pan American Health Organization (PAHO) considers *Shigella* spp. to be one of the most dangerous antibiotic resistant pathogens in Latin America and the Caribbean, exhibiting increased rates of resistance to antibiotics such as ciprofloxacin and azithromycin (4, 13, 14).

A landmark multisite case-control study ('GEMS') found that *Shigella* spp./EIEC were the leading enteric pathogens detected. The GEMS study found the attributable fraction for *Shigella* spp./EIEC to range from 3.9% to 83.2% depending on age stratum and study site (15). *Shigella* spp./EIEC was the most attributable or one of the most

attributable pathogens in infants, toddlers, and older children (15). A second landmark

multisite birth cohort study ('MAL-ED'), among children ≤ 2 years with and without

diarrheal disease who were not seeking care, found an attributable fraction of 10.2% for

Shigella spp./EIEC (16). While both were global in reach, there was only a single site in

Latin America (Peru), and there remains a paucity of insight from case-control studies

from the Caribbean, including Haïti, to set regional public health priorities.

The taxonomy of this group of pathogens is complicated because approaches were

established in the pre-genomics era. Phylogenetics revealed *Shigella* spp. and EIEC

are closely related (17). *Shigella* spp. are Gram-negative, nonmotile, facultatively

anaerobic, non-spore-forming, rod-shaped bacteria (18). They are differentiated from

Escherichia coli by pathophysiology and metabolism (e.g., inability to ferment lactose)

(18). The Shigella genus is divided into four serogroups with multiple serotypes: A (*S.*

dysenteriae, 12 serotypes); B (*S. flexneri*, 6 serotypes); C (*S. boydii*, 18 serotypes); and

D (*S. sonnei*, 1 serotype); these pathogens can readily be detected by PCR/qPCR using

primers for the invasion plasmid antigen H gene (*ipaH*) (9, 19). Given that *ipaH* is also

found among Enteroinvasive *Escherichia coli* (EIEC) it is difficult to disaggregate these

pathogens without cost-prohibitive molecular and culture analyses (19). Both the GEMS

and MAL-ED studies chose to treat *ipaH* positive samples as a singleton entity when

assessing attributions to symptomatic infection.

We hypothesized that among children in Haïti, *Shigella* spp./EIEC would be detected at

rates similar to the GEMS and MAL-ED studies and the odds of *Shigella* spp./EIEC

detection would be higher among, and attributed more often to, those with diarrheal

disease. To test these hypotheses, we conducted a case-control study among children

at the household level who were enrolled within a larger cross-sectional study in Haïti, a

setting with progressively deteriorating public health infrastructure, security and

governance (20). Despite limited data at the household level, diarrheal diseases have

been studied among care seeking children in Haïti (21-23). In 2012, *Shigella* spp. were

detected in 21.5% and 18.2% of children under 5 years presenting at a cholera

treatment center and an oral rehydration point in Port au Prince, respectively (23). Our

findings were consistent with our guiding hypothesis and aligned well with both the

GEMS and MAL-ED studies (15, 16).

METHODS

Ethics statement. Ethical approvals were obtained from the Institutional Review Board

at the University of Florida (IRB201703246) and the Comité National de Bioéthique

(National Bioethics Committee of Haïti, Ref:1718-35).

Study population and setting. A case-control study was conducted within a larger

cross-sectional study investigating healthcare-seeking intention and behavior in Haïti

(20). The study population resides within a deteriorating healthcare infrastructure due to

socio-political unrest (24). The under-five mortality rate is approximately 56 deaths per

1000 live births and the stunting rate is 20% (25). In 2023, rates of vaccination of 12

vaccines in the WHO/UNICEF schedule ranged from only 41-75%, with some vaccine

coverage lower than in 2013, possibly due to the state of unrest (26). Participant

households were located in the communes of Gressier and Leogane Haïti (located 12

and 27 miles respectively southwest of the capital Port-au-Prince). The enrollment

strategy aimed to equitably enroll families across rural, peri-urban and urban areas. The

study area (477 km^2) was divided into grid cells two square kilometers in size and each

cell was classified by density; "low" (0-112 structures), "medium" (113-397 structures),

or "high" (398-4664 structures). Six grid cells of each density were randomly selected

for sampling. Each of the selected grid cells was divided into 18, 24, or 53 Thiessen

polygons based on their designation as low, medium, or high density, respectively. The

study team navigated to a targeted polygon and screened the first house they

encountered. This process was repeated until at least one house in each polygon was

enrolled or all households were screened. Enrollment criteria included a head of

household who consented to complete a household survey and had at least one child

under five years old for whom consent was obtained to collect a stool swab. The

surveys were conducted to ascertain the socioeconomic status of families and the

recent diarrhea history of child participants. Although not part of the criteria, participants

were not actively seeking care at the time of enrollment.

Clinical procedures. Anthropomorphic measurements were obtained including height,

weight, and mean upper arm circumference (MUAC). Temperature was measured and

dehydration status was assessed using WHO IMCI methods for children reporting

current fever or diarrhea symptoms, respectively (27).

Laboratory procedures

Sample collection. Stool samples were obtained via rectal swabs, immediately placed

on ice in transport tubes with 3 mL of normal saline and brought to a laboratory in

Gressier. The contents of the transport tubes were divided into two 2 mL tubes and

stored at -80°C. Samples were transported on dry ice to the Emerging Pathogens

Institute at University of Florida for analysis.

Molecular assays. Standard methods for nucleic acid extraction were performed (28,

29). NEB Luna Universal qPCR Master Mix, nuclease free water, and primers (*ipaH* or

16S) were combined as a master mix. The *ipaH* primer was chosen for the detection of

Shigella spp./EIEC and the *16S* primer was used as a *eubacteria* control (30, 31). Each

well of a Hard-Shell® 96-Well PCR plate was loaded with 20 µL of master mix along

with either 1 µL of sample or 1 µL of control. The plates were set onto a thermocycler

(Bio Rad CFX96 Touch Real-Time PCR Detection System) to denature at 95°C for one

minute, then 35 cycles of denaturing at 95°C for 15 seconds, and annealing and

extension at 60°C for 30 seconds. Two operators conducted two biological replicates for

each of nine 96-well plates and switched between two thermocycler machines (except

for plate 2) to control for machine or operator bias. One positive control of *Shigella* spp.

genomic DNA and two negative controls of nuclease-free water were included on each

plate.

Analytic strategy

Definitions. Two definitions for cases and controls were established based on survey

responses to questions about history of diarrheal symptoms (Fig S1). Cases for

'diarrheal symptoms' (DS) were defined as answering 'YES' to the question "Diarrheal

symptoms within the last 7 days?". Controls for 'diarrheal symptoms' were defined as

answering 'NO' to the same question. Cases for 'acute diarrhea' (AD) were defined as

answering 'YES' to the previous question as well as the questions "Three or more loose

stools in the past 24 hours?" and "Onset less than 7 days ago?". Controls for 'acute

diarrhea' were defined as answering 'NO' to any or all of the three previous questions.

Analysis of molecular data. Results from the qPCR reactions were analyzed

quantitatively and qualitatively. Blank readings indicative of no detection of *ipaH* were

set to the maximum possible Ct value of 35. Qualitatively, samples were ruled positive if

the *ipaH* Ct was less than 28 and the *16S* Ct value was less than 26. The *ipaH*

threshold was based on standard curves and positive/negative controls; the *16S*

threshold was based on distributions among experimental samples and

positive/negative controls. Samples negative for *16S* were excluded from further

analysis.

Statistical analysis. A socio-economic status (SES) index was previously generated

using components from the household survey (20). Data were described by proportions

for categorical variables and medians with interquartile ranges for continuous variables.

Characteristics of cases were compared to controls within each case definition using the

two-sided non-paired Mann-Whitney U test, Chi-square test for independence, and

Fisher's exact test. For the primary analysis, an odds ratio was used to compare odds of

Shigella spp./EIEC detection among cases and controls for both DS and AD case

definitions. Additionally, adjusted odds ratios were calculated to account for age, sex,

and socioeconomic status. Attributable fractions were calculated with the graphPAF

package in R using the Bruzzi formula with 1,000 bootstrap replications (32, 33). A two-

sided non-paired Mann-Whitney U test was used to compare differences in Ct values

between cases and controls for each definition for all samples and only *ipaH* positive

samples. All analyses were performed using R Statistical Software (v4.4.1) (34, 35).

Figures were created using R Statistical Software (v4.4.1) and GraphPad Prism

(v10.4.1) (34, 36).

RESULTS

Participant characteristics. Of the 868 households screened, 568 were enrolled with

794 children. Among the enrolled children, 732 met inclusion criteria for the case-control

analysis (Fig 1). Over half of the participants were between 24-59 months (57%, n=416)

and the median age was 26 months (IQR 12-42 months) (Tables 1 and 2). Participation

by sex was approximately equal (51%, n=372 female). The median MUAC was 146 mm

(IQR 139-155 mm) and based on these measurements 1% (n=5) were severely

malnourished (Fig 2) (37). The majority of both male and female participants were within

3 z-scores of their age-specific median weight (98%, 708; Fig S2) and median height

(93%, 681; Fig S3) (38). Approximately 10% (n=71) of participants reported a current

fever, 1% (n=6) reported blood in stool and none of the participants with acute diarrhea

(0/43) were dehydrated (Table 3).

Characteristics of cases and controls. One quarter of participants (n=193) fit the

case definition for 'diarrheal symptoms' (Table 1) and 6% (n=43) fit the more restrictive

case definition for 'acute diarrhea' (Table 2). The DS and AD cases were slightly

younger than the respective DS (P<.001) and AD (P=.022) controls and had slightly

lower MUAC measurements than the respective DS (P=<.001) and AD (P=.010)

controls. The proportion of cases who were female was similar to that of controls for

both case definitions. There was no significant difference between cases and controls in

the proportions of underweight or severely malnourished participants for either DS or

AD case definitions. The SES distribution of DS and AD cases was not significantly

different than their corresponding DS and AD controls.

Pathogen detection. The overall rate of *ipaH* detection was 7.5% (n=55). Among the

DS cases, 22 (11.4%) were *ipaH* positive, representing an increased odds of 97%

(OR=1.97; 95%CI 1.11 to 3.46) for detection among participants with diarrheal disease

(Table 1). Among the AD cases, 8 (5.9%) were *ipaH* positive, representing an increased

odds of 212% (OR=3.12; 95% CI 1.29 to 6.81) for detection among participants with

diarrheal disease (Table 2). When controlling for age, sex, and SES, detection of *ipaH*

was associated with an increased odds of diarrheal disease for both case definitions;

DS (aOR=1.84; 95% CI 1.02 to 3.27) and AD (aOR=2.83; 95% CI 1.14 to 6.36). Among

participants with positive *ipaH* Ct values, the median value for DS cases (25.9; IQR

22.4-26.8) compared to DS controls (24.6; IQR 22.3-26.5) was not statistically different

(p=0.23). The median *ipaH* Ct value among AD cases (26.4; IQR 24.3-26.9; p=.03)

compared to AD controls (24.6; IQR 21.3-26.6) was slightly higher (p=0.03; Fig S4). The

unadjusted attributable fractions of *Shigella* spp./EIEC for diarrheal disease were 5.62%

(95% CI 0.44% to 10.9%) for DS and 12.6% (95% CI 0% to 25.3%) for AD. After

adjusting for confounders, the attributable fractions of *Shigella* spp./EIEC to diarrheal

disease were 5.2% (95% CI 0% to 10.5%) for DS and 12.0% (95% CI 0% to 24.8%) for

AD.

DISCUSSION

In this case-control study, detection of *Shigella* spp./EIEC was common and attributed

to diarrhea symptoms among children under five years of age in the Caribbean nation of

Haïti. Participants with *Shigella* spp./EIEC were twice as likely to report diarrheal

symptoms (DS) within the seven-day period prior to rectal swab collection. Among

participants meeting the more stringent acute diarrheal (AD) disease definition, the odds

of having AD increased to three times as likely. When adjusting for the potential

confounding variables of sex, age category, and SES, similar findings were made.

Attributable fractions of *Shigella* spp./EIEC detection and symptomatic disease were

more similar to the MAL-ED study than GEMS (15, 16). The GEMS study, stratified by

age, found that *Shigella* spp./EIEC were the attributable pathogen in 3.9% to 15.8% of

cases in infants, 15.4% to 64.9% of cases in toddlers, and 17.8% to 83.2% in older

children across all sites (15). The difference in proportions compared to our findings

may be due to differences in participant inclusion criteria, as those included in the

GEMS study were actively seeking care (15). The MAL-ED study, while focusing on

children under 2 years of age, included participants at the household level not actively

seeking care. Our attributable fraction of 12.9% is close to their findings of 10.2% of

cases that were attributed to *Shigella* spp./EIEC.

This study found negligible qPCR differences (<1 Ct) between cases and controls

for both DS and AD definitions. This contrasts with the MAL-ED study that found

participants with diarrhea attributed to *Shigella* spp./EIEC had lower Ct values (higher

abundance) compared to controls as would be expected (15, 16, 39, 40). There are

multiple explanations for our findings including study design (e.g., lower sample size),

procedure (e.g. possible differences in extraction efficiencies) and biological variation

(possible differences in target copy number between cases and controls). Importantly,

our attributable fractions for *Shigella* spp./EIEC align well with the attributable fraction

found in the MAL-ED study for diarrheal disease. Lastly, our study had low rates of

reported blood in stool overall (< 1%). Whereas, in MAL-ED, 4.7% of participants with

Shigella-specific diarrhea reported blood in stool (16). An underlying pathophysiologic

explanation for this difference is unknown and would require additional laboratory

studies to assess differences in underlying gut health between the two populations.

Our findings should be considered within the context of the study limitations.

First, enrollment bias may have been present in the larger public health study (20)

because recruitment occurred Monday to Friday during daytime hours when potentially

eligible head of households and school aged children were likely outside of the home.

Second, the case definitions for DS and AD relied on survey responses, which are not

as accurate as data ascertained clinically. Third, there was a procedural delay of

approximately five years between sample collection and analysis which poses a risk of

sample degradation. To minimize this, *16S* primers were used as a control to confirm

that expected rates of total bacteria were present in the stool sample and all *16S*

negative samples were excluded. Further, the *ipaH* gene is present in both *Shigella* spp.

and EIEC.(41) While EIEC infection is uncommon (42, 43), especially compared to

other diarrheagenic pathogens including other *E. coli* pathogens and *Shigella* spp., they

cannot be differentiated from *Shigella* spp. without further analysis (44). Serotyping can

differentiate *Shigella* spp. from EIEC, however, antibodies to O antigens of *Shigella* spp.

have cross-reactivity with EIEC O antigens (45, 46).

CONCLUSION

Despite study limitations, *Shigella* spp./EIEC detection was common and associated

with symptomatic disease among participants within the community. The results indicate

the need for monitoring and control of *Shigella* spp./EIEC infections in Haïti as an

etiological agent for diarrheal disease among children under five years. Intervention

would reduce rates of acute disease as well as reduce nutritional and development

sequalae associated with *Shigella* spp./EIEC.

Acknowledgements: We thank the participants for joining in this study as well as the clinical and laboratory teams who collected and analyzed the samples. We are grateful to those who collaborated on the original clinical study in which the samples analyzed herein were collected. We are also grateful to R. Autrey, B. Johnson and K. Berquist for their administrative expertise at the University of Florida. We also thank Dr. James Platts-Mills for his guidance on computing attributable fractions. EJN was the PI and obtained IRB approval at the University of Florida. Associated protocol numbers/registrations are: University of Florida IRB 201703246 and the Comité National de Bioéthique (National Bioethics Committee of Haïti) Ref:1718-35. This collective research infrastructure and support was invaluable to the success of this study.

Funding: This work was supported by the National Institutes of Health [DP5OD019893] to EJN and [R01 AI135115] to DTL as well as the Children's Miracle Network. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Conceptualization: MAS, DTL, ATM, CB, MBK, EJN

Methodology: IS, SRK, EC, YC, LB, MAS, DTL, SM, ATM, MBK, EJN

Investigation: IS, SRK, YC, EC, LB, CB, MBK, EJN

Visualization: IS, SRK, MK, EJN

Funding acquisition: DTL, EJN

Project administration: YC, MBK, EJN

Supervision: YC, MAS, VBR, CB, MBK, EJN

Formal analysis: IS, MBK, EJN

Resources: MAS, VBR, DTL, ATM, EJN

Data Curation: IS, SRK, YC, EC, LB, MBK

Writing – original draft: IS, SRK, MBK, EJN

Writing – review & editing: YC, EC, LB, MAS, SM, VBR, DTL, ATM, CB, EJN

Competing interests: Authors declare that they have no competing interests.

Data and materials availability: All data produced in the present study are available

upon reasonable request to the authors.

Code availability: The computer code to reproduce data analyses is available upon

request.

FIGURE 1

Fig 1. Participant enrollment and exclusions.

FIGURE 2

Fig 2. Sex- specific MUAC measurements and Z-scores for all participants as well as stratifications for DS and AD case definitions. A MUAC <115 mm is considered severely malnourished. DS=Diarrheal symptoms; answered 'YES' to "diarrheal symptoms within the last 7 days". AD=Acute diarrhea; answered 'YES' "diarrheal symptoms within the last 7 days", and "three or more loose stools in the past 24 hours", and "onset less than 7 days ago". MUAC=mid-upper arm circumference.

9 **TABLE 1**

10 **Table 1. Identification of factors associated with diarrhea symptoms (DS)**

	Cases	Controls	OR[a] (95% CI)	P	aOR[b] (95% CI)	P
Diarrheal symptoms (DS)[c]	193 (26%)	539 (74%)				
ipaH **positive**	22 (11%)	33 (6%)	**1.973 (1.106-3.457)**	**0.019**	**1.838 (1.017-3.265)**	**0.040**
Age; median months (IQR)[d]	21 (10–36)	30 (14.5-46)	**0.976 (0.966-0.986)**	**<0.001**	-	-
Age						
0-11 months	59 (31%)	103 (19%)	REF	REF	REF	REF
12-23 months	49 (25%)	105 (20%)	0.815 (0.510-1.300)	0.390	0.844 (0.523-1.358)	0.485
24-59 months	85 (44%)	331 (61%)	**0.448 (0.301-0.669)**	**<0.001**	**0.456 (0.305-0.683)**	**<0.001**
Sex; Female	90 (47%)	282 (52%)	0.796 (0.572-1.107)	0.175	0.830 (0.591-1.163)	0.278
SES[e] Index						
1st quintile	41 (21%)	99 (18%)	REF	REF	REF	REF
2nd quintile	36 (19%)	117 (22%)	0.743 (0.440-1.251)	0.264	0.793 (0.464-1.35)	0.395
3rd quintile	43 (22%)	101 (19%)	1.028 (0.617-1.714)	0.915	1.120 (0.663-1.896)	0.672
4th quintile	40 (21%)	108 (20%)	0.894 (0.534-1.496)	0.670	0.907 (0.535-1.535)	0.715
5th quintile	33 (17%)	114 (21%)	0.699 (0.409-1.187)	0.187	0.680 (0.392-1.170)	0.164
MUAC (IQR)	143 (135-150)	146 (141-155)	**0.967 (0.952-0.981)**	**<0.001**	-	-

[a] OR=odds ratio.
[b] aOR= adjusted odds ratio. Age (Continuous) and MUAC were not accounted for when computing the adjusted odds ratios.
[c] DS=Diarrheal symptoms; answered 'YES' to "diarrheal symptoms within the last 7 days".
[d] IQR=Inter quartile range.
[e] SES=Socioeconomic status. See methods for components of the index.

TABLE 2

Table 2. Identification of factors associated with acute diarrhea (AD)

	Cases	Controls	OR[a] (95% CI)	P	aOR[b] (95% CI)	P
Acute diarrhea (AD)[c]	43 (6%)	689 (94%)				
ipaH positive	8 (19%)	47 (7%)	**3.122 (1.286-6.813)**	**0.007**	**2.832 (1.140-6.400)**	**0.016**
Age; median months (IQR)[d]	20 (9.5-29)	27 (13-43)	**0.977 (0.958-0.996)**	**0.019**	-	-
Age						
0-11 months	14 (33%)	148 (22%)	REF	REF	REF	REF
12-23 months	14 (33%)	140 (20%)	1.057 (0.483-2.314)	0.888	1.050 (0.467-2.360)	0.905
24-59 months	15 (35%)	401 (58%)	**0.395 (0.185-0.848)**	**0.016**	**0.412 (0.192-0.892)**	**0.023**
Sex; Female	19 (44%)	353 (51%)	0.753 (0.401-1.397)	0.371	0.826 (0.433-1.558)	0.555
SES[e] index						
1st quintile	11 (26%)	129 (19%)	REF	REF	REF	REF
2nd quintile	8 (19%)	145 (21%)	0.647 (0.244-1.648)	0.365	0.744 (0.275-1.940)	0.547
3rd quintile	10 (23%)	134 (19%)	0.875 (0.353-2.144)	0.769	1.040 (0.409-2.629)	0.934
4th quintile	4 (9%)	144 (21%)	0.326 (0.089-0.979)	0.060	0.335 (0.090-1.025)	0.070
5th quintile	10 (23%)	137 (20%)	0.856 (0.346-2.097)	0.732	0.841 (0.331-2.115)	0.711
MUAC	142 (135-150)	146 (140-155)	**0.967 (0.941-0.993)**	**0.015**	-	-

[a] OR=odds ratio.
[b] aOR= adjusted odds ratio.
[c] AD=Acute diarrhea; answered 'YES' "Diarrheal symptoms within the last 7 days", and "Three or more loose stools in the past 24 hours", and "Onset less than 7
days ago".
[d] IQR=Inter quartile range.
[e] SES=Socioeconomic status. See methods for components of the index.
Age (Continuous) and MUAC were not accounted for when computing the adjusted odds ratios.

TABLE 3

6 **Table 3. Report of fever or blood in stool stratified by diarrheal symptom (DS) and acute diarrhea (AD) case**
7 **definitions.**

	All n=732	DS[a] Cases n=193	DS[a] Controls n=539	AD[bc] Cases n=43	AD[bc] Controls n=689
Fever; n (%)	71 (10%)	12 (6%)	59 (11%)	1 (2%)	70 (10%)
Blood in stool; n (%)	6 (1%)	6 (3%)	0	0	6 (1%)

8 [a] DS=Diarrheal Symptoms; answered 'YES' to "diarrheal symptoms within the last 7 days".
9 [b] AD=Acute Diarrhea; answered 'YES' "diarrheal symptoms within the last 7 days", and "three or more loose stools in the past 24
10 hours", and "onset less than 7 days ago".
11 [c] Children with acute diarrhea were assessed for dehydration and all had 'no' dehydration.

REFERENCES

1. Liu L, Oza S, Hogan D, Chu Y, Perin J, Zhu J, et al. Global, regional, and national causes of under-5 mortality in 2000–15: an updated systematic analysis with implications for the Sustainable Development Goals. The Lancet. 2016;388(10063):3027-35.

2. Pinkerton R, Oriá RB, Lima AA, Rogawski ET, Oriá MO, Patrick PD, et al. Early childhood diarrhea predicts cognitive delays in later childhood independently of malnutrition. Am J Trop Med Hyg. 2016;95(5):1004-10.

3. Troeger C, Colombara DV, Rao PC, Khalil IA, Brown A, Brewer TG, et al. Global disability-adjusted life-year estimates of long-term health burden and undernutrition attributable to diarrhoeal diseases in children younger than 5 years. The Lancet Global Health. 2018;6(3):e255-e69.

4. Hussen S, Mulatu G, Yohannes Kassa Z. Prevalence of *Shigella* species and its drug resistance pattern in Ethiopia: a systematic review and meta-analysis. Annals of Clinical Microbiology and Antimicrobials. 2019;18(1):22.

5. Libby TE, Delawalla MLM, Al-Shimari F, MacLennan CA, Vannice KS, Pavlinac PB. Consequences of *Shigella* infection in young children: a systematic review. Int J Infect Dis. 2023;129:78-95.

6. Anderson JDIV, Bagamian KH, Muhib F, Amaya MP, Laytner LA, Wierzba T, et al. Burden of enterotoxigenic *Escherichia coli* and *Shigella* non-fatal diarrheal infections in 79 low-income and lower middle-income countries: a modelling analysis. The Lancet Global Health. 2019;7(3):e321-e30.

7. Njuguna HN, Cosmas L, Williamson J, Nyachieo D, Olack B, Ochieng JB, et al. Use of Population-based surveillance to define the high incidence of shigellosis in an urban slum in Nairobi, Kenya. PLOS ONE. 2013;8(3):e58437.

8. Kotloff KL, Winickoff JP, Ivanoff B, Clemens JD, Swerdlow DL, Sansonetti PJ, et al. Global burden of *Shigella* infections: implications for vaccine development and implementation of control strategies. Bull World Health Organ. 1999;77(8):651-66.

9. Deer DM, Lampel KA. Development of a multiplex real-time PCR assay with internal amplification control for the detection of *Shigella* species and enteroinvasive *Escherichia coli*. J Food Prot. 2010;73(9):1618-25.

10. Kotloff KL, Riddle MS, Platts-Mills JA, Pavlinac P, Zaidi AKM. Shigellosis. The Lancet. 2018;391(10122):801-12.

11. Baker S, Scott TA. Antimicrobial-resistant *Shigella*: where do we go next? Nature Reviews Microbiology. 2023;21(7):409-10.

12. Lampel KA, Formal SB, Maurelli AT. A brief history of *Shigella*. EcoSal Plus. 2018;8(1):10.1128/ecosalplus.ESP-0006-2017.

13. Ranjbar R, Farahani A. *Shigella*: antibiotic-resistance mechanisms and new horizons gor treatment. Infect Drug Resist. 2019;12:3137-67.

14. Pan American Health Organization, World Health Organization. Epidemiological alert: Emergence and spread of *Shigella sonnei* with extreme resistance to antibiotics. Potential risk for Latin America and the Caribbean. Washington DC. 2022. [Available from: https://www.paho.org/en/documents/epidemiological-alert-emergence-and-spread-shigella-sonnei-extreme-resistance-antibiotics].

15. Liu J, Platts-Mills JA, Juma J, Kabir F, Nkeze J, Okoi C, et al. Use of quantitative molecular diagnostic methods to identify causes of diarrhoea in children: a reanalysis of the GEMS case-control study. The Lancet. 2016;388(10051):1291-301.

16. Platts-Mills JA, Liu J, Rogawski ET, Kabir F, Lertsethtakarn P, Siguas M, et al. Use of quantitative molecular diagnostic methods to assess the aetiology, burden, and clinical characteristics of diarrhoea in

children in low-resource settings: a reanalysis of the MAL-ED cohort study. The Lancet Global Health.
2018;6(12):e1309-e18.
17. Hazen TH, Leonard SR, Lampel KA, Lacher DW, Maurelli AT, Rasko DA. Investigating the
relatedness of enteroinvasive *Escherichia coli* to other *E. coli* and *Shigella* isolates by using comparative
genomics. Infect Immun. 2016;84(8):2362-71.
18. Schroeder GN, Hilbi H. Molecular pathogenesis of *Shigella* spp.: controlling host cell signaling,
invasion, and death by type III secretion. Clin Microbiol Rev. 2008;21(1):134-56.
19. Venkatesan MM, Buysse JM, Kopecko DJ. Use of *Shigella flexneri ipaC* and *ipaH* gene sequences
for the general identification of *Shigella* spp. and enteroinvasive *Escherichia coli*. J Clin Microbiol.
1989;27(12):2687-91.
20. Klarman M, Schon J, Cajusma Y, Maples S, Beau de Rochars VEM, Baril C, et al. Opportunities to
catalyse improved healthcare access in pluralistic systems: a cross-sectional study in Haïti. BMJ Open.
2021;11(11):e047367.
21. Diaz JN, Iannotti LL, Louis Dulience SJ, Vie S, Jiang X, Grigura V, et al. Prevalence of diarrheagenic
Escherichia coli and impact on child health in Cap-Haïtien, Haïti. PLOS Global Public Health.
2023;3(5):e0001863.
22. Beau De Rochars VEM, Alam MT, Telisma T, Masse R, Chavannes S, Anilis MG, et al. Spectrum of
outpatient illness in a school-based cohort in Haïti, with a focus on diarrheal pathogens. Am J Trop Med
Hyg. 2015;92(4):752-7.
23. Charles M, Delva GG, Boutin J, Severe K, Peck M, Mabou MM, et al. Importance of Cholera and
other etiologies of acute diarrhea in post-earthquake Port-au-Prince, Haïti. Am J Trop Med Hyg.
2014;90(3):511-7.
24. Dyer O. Haiti's health system collapsing under weight of doctors' strike. BMJ. 2016;354:i3939.
25. Estimation UNI-AGfCM. Levels & Trends in Child Mortality [Internet]. 2023 [Available from:
https://childmortality.org/wp-content/uploads/2024/03/UNIGME-2023-Child-Mortality-Report.pdf].
26. United Nations Children's Fund, World Health Organization. Immunization country profiles.
World Health Organization. 2024. [Available from: https://data.unicef.org/resources/immunization-
country-profiles/].
27. World Health Organization, United Nations Children's Fund. Model IMCI handbook: integrated
management of childhood illness. Geneva: World Health Organization. 2005. [Availabel from:
http://apps.who.int/iris/bitstream/10665/42939/1/9241546441.pdf].
28. Lappan R, Henry R, Chown SL, Luby SP, Higginson EE, Bata L, et al. Monitoring of diverse enteric
pathogens across environmental and host reservoirs with TaqMan array cards and standard qPCR: a
methodological comparison study. The Lancet Planetary Health. 2021;5(5):e297-e308.
29. Madi N, Cato ET, Abu Sayeed M, Creasy-Marrazzo A, Cuénod A, Islam K, et al. Phage predation,
disease severity, and pathogen genetic diversity in cholera patients. Science. 2024;384(6693):eadj3166.
30. Grembi JA, Mayer-Blackwell K, Luby SP, Spormann AM. High-Throughput Multiparallel
Enteropathogen Detection via Nano-Liter qPCR. Frontiers in Cellular and Infection Microbiology.
2020;10.
31. Maeda H, Fujimoto C, Haruki Y, Maeda T, Kokeguchi S, Petelin M, et al. Quantitative real-time
PCR using TaqMan and SYBR Green for *Actinobacillus actinomycetemcomitans*, *Porphyromonas
gingivalis*, *Prevotella intermedia*, *tetQ gene* and total bacteria. FEMS Immunol Med Microbiol.
2003;39(1):81-6.
32. Ferguson J, O'Connell M. Estimating and displaying population attributable fractions using the R
package: graphPAF. Eur J Epidemiol. 2024;39(7):715-42.
33. Bruzzi P, Green SB, Byar DP, Brinton LA, Schairer C. Estimating the population attributable risk
for multiple risk factors using case-control data. Am J Epidemiol. 1985;122(5):904-14.

34.   R Core Team. R: A Language and Environment for Statistical Computing. 2024. [Available from:
https://www.R-project.org/].
35.   Posit Team. RStudio: Integrated Development Environment for R. 2024. [Available from:
http://www.posit.co/].
36.   GraphPad Software. GraphPad Prism Version 10.4.1. (n.d.). [Available from:
www.graphpad.com].
37.   de Onis M, Yip R, Mei Z. The development of MUAC-for-age reference data recommended by a
WHO Expert Committee. Bull World Health Organ. 1997;75(1):11-8.
38.   WHO Multicentre Growth Reference Study Group. WHO Child Growth Standards based on
length/height, weight and age. Acta Paediatr Suppl. 2006;450:76-85.
39.   Vu DT, Sethabutr O, Von Seidlein L, Tran VT, Do GC, Bui TC, et al. Detection of *Shigella* by a PCR
assay targeting the *ipaH* gene suggests increased prevalence of shigellosis in Nha Trang, Vietnam. J Clin
Microbiol. 2004;42(5):2031-5.
40.   Huang Z, He Z, Wei Z, Wang W, Li Z, Xia X, et al. Correlation between prevalence of selected
enteropathogens and diarrhea in children: a case–control study in China. Open Forum Infectious
Diseases. 2021;8(10).
41.   Løbersli I, Wester AL, Kristiansen Å, Brandal LT. Molecular differentiation of *Shigella* Spp. from
enteroinvasive E. Coli. Eur J Microbiol Immunol (Bp). 2016;6(3):197-205.
42.   Giannella RA. Chapter 107 - infectious enteritis and proctocolitis and bacterial food poisoning.
In: Feldman M, Friedman LS, Brandt LJ, editors. Sleisenger and Fordtran's Gastrointestinal and Liver
Disease (Ninth Edition). Philadelphia: W.B. Saunders; 2010. p. 1843-87.e7.
43.   American Society of Microbiology. What's the significance of enteroinvasive *E. coli* (EIEC),
enteropathogenic *E. coli* (EPEC), and enteroaggregative *E. coli* (EAEC) in Stool? American Society of
Microbiology: American Society of Microbiology; 2019 [Available from:
https://asm.org/articles/2019/march/whats-the-significance-of-enteroinvasive-e-coli.
44.   Vieira N, Bates SJ, Solberg OD, Ponce K, Howsmon R, Cevallos W, et al. High prevalence of
enteroinvasive *Escherichia coli* isolated in a remote region of northern coastal Ecuador. Am J Trop Med
Hyg. 2007;76(3):528-33.
45.   Sahl JW, Morris CR, Emberger J, Fraser CM, Ochieng JB, Juma J, et al. Defining the
phylogenomics of *Shigella* species: a pathway to diagnostics. J Clin Microbiol. 2015;53(3):951-60.
46.   Kim HJ, Ryu JO, Song JY, Kim HY. Multiplex polymerase chain reaction for identification of
Shigellae and four *Shigella* species using novel genetic markers screened by comparative genomics.
Foodborne Pathog Dis. 2017;14(7):400-6.