Failing maternal-fetal tolerance in Systemic Lupus Erythematosus (FaMaLE): a prospective

cohort study for finding the molecular mechanisms behind pregnancy complications

W. Dankers[1,2,3,4,*], J.F. van Ruitenbeek[1,2], S.A. Germe[1,2,3], A.R. Parra Sanchez[1,2,3], M.F.H.M. van Gaal[1,2,5],

K. Cramer[6], D.C. Rohrich-Heldens[6], M.A. de Boer[4,5], L.G.M. van Baarsen[1,2,3], I.E.M. Bultink[1,3,4,*]

[1]Amsterdam UMC, Department of Rheumatology and Clinical Immunology, Amsterdam, the

Netherlands

[2]Amsterdam UMC, Department of Experimental Immunology, Amsterdam, the Netherlands,

[3]Amsterdam UMC, Amsterdam Institute for Infection and Immunity, Amsterdam, the Netherlands,

[4]Amsterdam UMC, Amsterdam Reproduction and Development, Amsterdam, the Netherlands,

[5]Amsterdam UMC, Department of Obstetrics and Gynaecology, Amsterdam, the Netherlands

[6]Patient research partner, Amsterdam, the Netherlands

* To whom correspondence should be addressed: Wendy Dankers w.dankers@amsterdamumc.nl,

Irene E.M. Bultink iem.bultink@amsterdamumc.nl.

Keywords: pregnancy, immunology, immunophenotyping, research design, rheumatology

Word count: 2377

Subject areas: rheumatology (primary), obstetrics & gynaecology, immunology

ABSTRACT

Introduction

Pregnant women with systemic lupus erythematosus (SLE) have an increased risk of maternal

complications and adverse fetal outcomes. These include preeclampsia, preterm birth and fetal

growth restriction. Interestingly, this increased risk persists in subsequent pregnancies, whereas it

decreases in healthy women due to the development of maternal-fetal tolerance. As maternal-fetal

tolerance is crucial for a healthy pregnancy, we hypothesize that its failure contributes to the

increased risk of pregnancy complications in women with SLE. Therefore, we initiated the FaMaLE

study to investigate the failure of maternal-fetal tolerance in pregnant women with SLE.

Methods and analysis

In the FaMaLE study, women with SLE and healthy women are included in their first trimester of

pregnancy (< 14 weeks gestational age (GA)) at Amsterdam UMC. Throughout the pregnancy, data on

SLE disease activity, pregnancy course, and medication use are collected. Peripheral blood is collected

once per trimester, within 48 hours before delivery and 5-12 weeks post-partum. In addition, the

placenta is collected after delivery. Whole blood, peripheral blood mononuclear cells (PBMC) and

placenta samples are freshly analyzed by flow cytometry to assess immune cell composition. The

resulting data are analyzed in relation to SLE disease course, pregnancy course and pregnancy

outcomes.

Ethics and dissemination

The study has been approved by the Amsterdam UMC Medical Ethics Committee and all participating
women will be asked to provide informed consent. The findings will be disseminated through peer-
reviewed publications, presentations at scientific meetings and via patient organizations.
46
STRENGTHS AND WEAKNESSES
- A unique prospective longitudinal study design, featuring the collection of serum, plasma and
PBMC throughout and after pregnancy, alongside placental cells and biopsies from the same
participants. This is complemented by detailed clinical data on SLE disease course and
pregnancy course, and outcomes.
- Fresh flow cytometry analyses allow immediate assessment of cell composition in blood and
placenta, without freeze/thawing effects
- The study does not include pre-pregnancy collection of serum, plasma and PBMC; however
detailed clinical data are collected during this period.
56
57
INTRODUCTION
Systemic lupus erythematosus (SLE) is a chronic autoimmune disease with the potential to affect all
organ systems in the body. It is characterized by a disruption in immunological tolerance, leading to
the production of autoantibodies targeting nuclear self-antigens[1,2]. SLE affects 1 in 1000 people
worldwide, with a higher prevalence among individuals of non-Caucasian descent, and predominantly
affects women during their childbearing years[1]. Unfortunately, up to 50% of pregnant women with
SLE experience complications, such as pre-eclampsia, fetal growth restriction and preterm birth[3,4].
Importantly, almost 50% of preterm births have a spontaneous onset, the other preterm births are
induced mostly because of pre-eclampsia or fetal growth restriction[5]. Partially due to (the increased
risk of) pregnancy complications, 60% of SLE patients have fewer children than desired[6,7].
Despite the significant impact on the health and wellbeing of pregnant women with SLE and their
children, the exact underlying cellular and molecular mechanisms driving these complications are still
poorly understood. Understanding of these biological processes is crucial to enable prediction, early
detection and novel treatments for placenta-related pregnancy complications in SLE.
One of the requirements for a healthy pregnancy is a well-functioning maternal immune system,
which facilitates proper placentation, triggers the labor process at the end of pregnancy, and ensures
that the semi-allogeneic fetus is not immunologically rejected during pregnancy[8]. This latter
phenomenon is called maternal-fetal tolerance, and is thought to be the reason why generally the
risk of pregnancy complications decreases in subsequent pregnancies with the same father.
Interestingly, this protective effect is absent in women with SLE[3], suggesting that impaired maternal-
fetal tolerance may play a central role in the increased risk for pregnancy complications in women
with SLE.
Previous data suggest an altered immune response in pregnant women with SLE who experience
pregnancy complications[9]. Furthermore, there are some studies reporting an increased deposition of
neutrophils extracellular traps (NETs) and higher numbers of natural killer (NK) cells and extravillous
trophoblasts in the placentas of women with SLE. However, if and how this dysregulation contributes
to pregnancy complications in women with SLE is currently unknown. Therefore, to address these
knowledge gaps, we have initiated the FaMaLE study in which we aim to delineate the differences in
cellular composition and function at the maternal-fetal interface in women with SLE, both with and
without pregnancy complications, compared to healthy pregnant women.

METHODS AND ANALYSIS

Study overview and design

This single-center prospective cohort study was initiated in May 2022 and will include 60 pregnant SLE patients and 18 healthy pregnant controls, all aged 18 years or older, who receive their antenatal care from the first trimester onwards at Amsterdam UMC. Women with SLE must be diagnosed according to the ACR 1997, SLICC and/or ACR/EULAR 2019 criteria and must be under treatment for their disease at Amsterdam UMC. The treating gynaecologist and rheumatologist will screen potentially suitable participants during their first trimester of pregnancy at the Department of Obstetrics and Gynaecology and the Department of Rheumatology and Clinical Immunology of Amsterdam UMC. Eligible subjects will be informed about the aim, procedures and potential risk of the study by members of the research team. Participants can be enrolled before 14 weeks gestational age after providing informed consent. An overview of the study protocol is shown in *figure 1*.

Figure 1. FaMaLE study flowchart. *Abbreviations: SLE-CP, SLE complicated pregnancy; SLE-NCP, SLE non-complicated pregnancy; HC, healthy control. Created in BioRender. Dankers, W. (2025) .*
https://BioRender.com/t19u876

Data collection includes demographic information (age, ethnicity, highest level of education), medical history (smoking status, alcohol/drug use, general and SLE-specific medical history, current and prior medication use) and obstetric history (number of previous pregnancies, any previous miscarriages, course and outcome of previous pregnancies). SLE disease activity will be assessed using the SLEPDAI during pregnancy[10], and SLEDAI-2K[11] before pregnancy and after delivery. Preconceptional disease-related damage is assessed by the Systemic Lupus International Collaborating Clinics/American College of Rheumatology (SLICC/ACR) Damage Index[12]. Additionally, gestational age at delivery and gender, weight and delivery outcomes for the newborn will be recorded. Apart from routine

laboratory investigations for regular clinical care in pregnant women with SLE, at each visit an additional 45 ml of blood is collected in 4 heparin tubes and 1 coagulation tube. From these serum, plasma and PBMC are isolated. The blood sample from time point 4 (T4) is collected within 48 hours before delivery, when the woman is admitted at the delivery room. After delivery, the placenta will be collected for further analysis.

Medical data will be recorded by the treating gynaecologist and rheumatologist. Histopathological examination of all placentas will be performed by one assigned pathologist according to the Amsterdam Placental Workshop Group Consensus Statement[13]. Pregnancies are classified as complicated if they meet any of the following criteria: preterm delivery (<37 weeks gestational age), pregnancy induced hypertension (*de novo* or superimposed), pre-eclampsia (ISSHP criteria 2018[14]), small for gestational age child (<p10), placental abruption and/or intrauterine fetal death (>16 weeks gestational age).

Inclusion and exclusion criteria

Inclusion criteria for SLE patients
- Age 18 years or older
- Gestational age at inclusion <14 weeks
- Antenatal care from first trimester onwards in the Department of Obstetrics and Gynaecology of Amsterdam UMC
- Diagnosed with SLE, fulfilling ACR 1997, SLICC and/or ACR/EULAR 2019 criteria
- Signed informed consent

Inclusion criteria for healthy pregnant women
- Age 18 years or older
- Gestational age at inclusion <14 weeks
- Antenatal care and delivery in the Department of Obstetrics and Gynaecology of Amsterdam UMC
- Signed informed consent

Exclusion criteria for SLE patients
- Diagnosed with a systemic inflammatory disease other than SLE
- Diagnosed with a malignant disease in the previous 5 years, except for basal cell or squamous epithelial cell skin cancers
- Multifetal pregnancy

Exclusion criteria for healthy pregnant women
- Diagnosed with a systemic inflammatory disease
- Diagnosed with a malignant disease in the previous 5 years, except for basal cell or squamous epithelial cell skin cancers
- Multifetal pregnancy
- Treatment with immunosuppressive agents
- Pre-existing hypertensive disorder
- Obstetric history of hypertensive disorder of pregnancy, fetal growth restriction (birth weight < p10), preterm birth, placental abruption or intra-uterine fetal death

160 - Obesity (BMI > 30)
161 - Any disease for which the woman receives specialist treatment other than obstetric care
162

Sample size calculation

163
The rate of pregnant SLE patients to be included in this study is expected to be 15-20 per year based
on our experience over the last >15 years in the multidisciplinary SLE pregnancy care team in
Amsterdam UMC. Pregnancy complications due to placental insufficiency as defined above are
expected in 30% of SLE patients as previously published by our research group[3]. In the present study
women are categorized into 3 groups; healthy control (HC), SLE complicated pregnancy (SLE-CP), or
SLE not complicated pregnancy (SLE-NCP).
In the analysis, differences in cell composition in whole blood, PBMC and placental tissue between
these groups will be compared. We aim to detect differences greater than 10%, so at least 16 patients
per group are needed for 80% power to detect differences (with an α of 0.05). Accounting for a 10%
dropout rate throughout this study, 18 subjects per group need to be included. Considering an
expected 30% incidence of pregnancy complications in SLE patients, 60 pregnant women with SLE are
needed to include at least 18 patients with complications. Additionally 18 healthy pregnant women
will be included.
177

Study aims

178
The primary aims of this study are (1) to compare the cellular profile of placental cells from women in
the HC, SLE-CP, SLE-NCP groups; (e.g. surface markers, transcription factors, gene expression), (2) to
analyze the cellular function, cell-cell interactions and the development of tolerance at the *in vitro*
maternal-fetal interface, and (3) to identify potential peripheral blood biomarkers associated with
pregnancy complications.
As secondary objectives, the study aims to establish an *in vitro* organoid-based model for the
maternal-fetal interface in SLE and to compare the cellular profile and different proteins in the
peripheral blood of women in the three groups.
187

Study procedures

188
Blood sampling
From the blood drawn at the five timepoints throughout pregnancy, serum and plasma will be
isolated within four hours after blood draw from anti-coagulation and heparin tubes, respectively.
They are stored at -80°C for later use for detection of biologic markers (antibodies, cytokines,
(disease-related) proteins, cell-free RNA). 100 μl of whole blood is saved for analysis using flow
cytometry, and then peripheral blood mononuclear cells (PBMC) are isolated within 24 hours from
the heparin tubes, using Lymphoprep™ (Serumwerk, Bernburg, Germany) density gradient
centrifugation. 2-4 million PBMC are analyzed using flow cytometry, and remaining PBMC are frozen
per 5-10 million cells in freezing medium (10% DMSO (VWR, Amsterdam, The Netherlands) and 90%
Iscove's Modified Dulbecco's Medium (IMDM; Gibco) supplemented with 30% FCS, 0.05mg/ml
gentamicin (Gibco) and 0.00036 (v/v)% β-mercaptoethanol.
200

Placenta sampling
After delivery the placenta will be collected by the research team and stored in 0,9% NaCl at 4°C until
processing, which starts within 2 hours after delivery. Since most current protocols for processing
placenta focus on obtaining either immune cells or trophoblasts, a customized protocol was used to

capture both cell types based on previously published protocols[15,16]. The umbilical cord is removed
from the placenta, leaving the insertion intact. 4 cm of umbilical cord is stored in formalin for
histopathological examination and classification by the assigned pathologist, as well as 1/3 of the
chorioamniotic membranes. The remaining decidua parietalis and chorion (PC) are separated from
the amnion, cut off and placed in 1x PBS. Once the membranes and umbilical cord are removed, the
placenta weight is determined. 3 biopsies from random locations at the placenta are taken using an
8-mm biopsy punch, covered in O.C.T., snap-frozen in liquid nitrogen and stored at -80°C. The
placenta is then cut into slices of ~1-1.5 cm thick, to assess the percentage of calcifications
throughout the tissue. At least two blocks of 2x1 cm, of which one includes the umbilical cord
insertion site, and both blocks covering the entire depth of the placenta, are harvested and stored in
formalin for histopathological examination and classification by the assigned pathologist.
Subsequently, the basal plate (BP; decidua basalis and ~5mm of villi) is dissected and put in PBS. Now,
both PC and BP are cut into small fragments (~2-3 mm diameter) and washed with 1x PBS until the
PBS is clear. After washing with Roswell Park Memorial Institute (RPMI) medium, tissues are
transferred to C-tubes (Miltenyi Biotec, Bergisch Gladbach, Germany) and resuspended in digestion
mix (200mg/ml collagenase and 10mg/ml DNase (Roche) in RPMI). Tissue is then dissociated on
GentleMACS (BP: 1x m-heart-02.01, PC: 2x h-tumor-02.01) and left to digest for 1 hour at 37°C in a
shaking water bath. After 30 minutes there is another round of dissociation (all: 1x m-heart-02.01).
Digested tissues are filtered through 100μm cell strainers (Miltenyi Biotec), after which red blood cells
(RBC) are lysed using 1x RBC lysis buffer (BioLegend). Finally, cells are filtered through a 70μm cell
strainer, after which ~5 million cells are analyzed using flow cytometry, 1 million cells are plated for
subsequent culture and remaining cells are frozen as described for PBMC.
227
228   *Flow cytometry*
229   2-4 million PBMC, 5 million cells from BP and PC, and 100 μl whole blood are freshly analyzed for
230   cellular composition using five flow cytometry panels: myeloid cells, B cells, T cells, ILC and non-
231   immune cells (***Table 1-5***). With each sample, a reference control sample is taken along to account for
232   batch effects during analysis.
233
234   **Table 1. Flow cytometry panel for PBMC and placenta: Myeloid cells.** CD = cluster of differentiation,
235   DC = dendritic cell, NK cell = natural killer cell, BDCA2 = blood dendritic cell antigen 2. Viability is
236   measured using fixable viability dye.

Target	Fluorochrome	Clone	Company	Purpose
BDCA2	APC	201A	BioLegend	DC differentiation
CD3	PE-Dazzle594	UCHT1	BioLegend	T cells
CD11b	BUV395	ICRF44	BD Biosciences	DC differentiation, macrophages, neutrophils
CD11c	AlexaFluor700	Bu15	BioLegend	DC differentiation, macrophages
CD14	FITC	61D3	eBioscience	DC- and monocyte differentiation, macrophages
CD15	BV605	W6D3	BioLegend	Neutrophils
CD16	APC-eFluor780	CB16	eBioscience	Monocyte differentiation, Neutrophils, NK cells
CD19	BV650	HIB19	BioLegend	B cells
CD45	PerCP-Cy5.5	HI30	BioLegend	Immune cells

Target	Fluorochrome	Clone	Company	Purpose
CD56	PE	NCAM 16	BD Biosciences	NK(T) cells
CD103	BV421	Ber-ACT8	BioLegend	CD8[+] T cell-, DC- and NK cell differentiation
CD123	PE-Cy7	6H6	eBioscience	DC differentiation
Viability	eF506	-	Invitrogen	Cell viability

Table 2. Flow cytometry panel for PBMC and placenta: B cells. CD = cluster of differentiation, IgD = immunoglobulin D, IgM = immunoglobulin M. Viability is measured using fixable viability dye.

Target	Fluorochrome	Clone	Company	Purpose
CD19	BV650	HIB19	BioLegend	B cells
CD21	PE-Cy7	B-ly4	BD Biosciences	B cell differentiation
CD24	BV421	ML5	BioLegend	B cell differentiation
CD27	APC-Fire750	O323	BioLegend	B- and CD8[+] T cell differentiation
CD38	PE	HIT2	BioLegend	B cell differentiation
CD45	PerCP-Cy5.5	HI30	BioLegend	Immune cells
CD138	APC	MI15	BioLegend	B cell differentiation
IgD	BV785	IA6-2	BioLegend	B cell differentiation
IgM	FITC	G20-127	BD Biosciences	B cell differentiation
Viability	eF506	-	Invitrogen	Cell viability

241 **Table 3. Flow cytometry panel for PBMC and placenta: T cells.** CD = cluster of differentiation, TCRγδ
242 = T cell receptor γδ. Viability is measured using fixable viability dye.

Target	Fluorochrome	Clone	Company	Purpose
CCR7	PE-Cy7	3D12	BD Biosciences	CD4$^+$-, and CD8$^+$ T cell differentiation
CD3	APC-R700	UCHT1	BD Biosciences	T cells
CD4	BUV563	SK3	BD Biosciences	CD4$^+$ T cells
CD8	BV785	RPA-T8	BioLegend	CD8$^+$ T cells
CD25	APC	M-A251	BioLegend	CD4$^+$ T cell differentiation
CD45	PerCP-Cy5.5	HI30	BioLegend	Immune cells
CD45RA	eFluor450	HI100	eBioscience	γδ- CD4$^+$-, CD8$^+$ T cell differentiation
CD45RO	PE	UCHL1	BioLegend	γδ- CD4$^+$-, CD8$^+$ T cell differentiation
CD56	BV605	HCD56	BioLegend	γδ differentiation, NK(T) cells
CD127	APC-eFluor780	eBioRDR5	eBioscience	Non-NK ILC
TCRγδ	FITC	B1	BD Biosciences	γδ T cells
Viability	eF506	-	Invitrogen	Cell viability

244 **Table 4. Flow cytometry panel for PBMC and placenta: innate lymphoid cells (ILC).** CD = cluster of
245 differentiation, *RPA-2.10, OKT3, 61D3, CB16, HIB19, TULY56, HIR2. Viability is measured using
246 fixable viability dye.

Target	Fluorochrome	Clone	Company	Purpose
CD45	PerCP-Cy5.5	HI30	BioLegend	Immune cells
CD56	BV605	HCD56	BioLegend	NK(T) cells
CD103	BV421	Ber-ACT8	BioLegend	CD8$^+$ T cell-, DC- and NK cell differentiation
CD117	PE	104D2	eBioscience	Non-NK ILC differentiation
CD127	PE-Cy7	A019D5	BioLegend	Non-NK ILC
Lineage	FITC	Multiple*	eBioscience	Identify non-ILC
NKp44	AlexaFluor647	P44-8	BioLegend	Non-NK ILC differentiation
Viability	eF506	-	Invitrogen	Cell viability

248 **Table 5. Flow cytometry panel for placenta: Non-immune cells.** CD = cluster of differentiation, CK =
249 Cytokeratin, HLA = human leukocyte antigen, * = intracellular staining. Viability is measured using
250 fixable viability dye.

Target	Fluorochrome	Clone	Company	Purpose
CD31	APC-eF780	WM-59	eBioscience	Endothelial cells
CD45	PerCP-Cy5.5	HI30	BioLegend	Immune cells
CD235a	PE-Cy5	HIR2	BioLegend	Red blood cells
CK7*	AF647	W16155A	BioLegend	Trophoblasts
HLA-DR	BV711	L243	BioLegend	Non-syncytiotrophoblasts
HLA-G	PE	MEM-G/9	Exbio	Cytotrophoblasts, extravillous trophoblast, syncytiotrophoblasts

Viability	eF506	-	Invitrogen	Cell viability
Vimentin*	AF488	O91D3	BioLegend	Decidual stromal cells, trophoblasts

Whole blood is directly stained with the myeloid panel after which red blood cells are lysed using 1x RBC lysis buffer (BioLegend). PBMC and placenta cells are stained with the respective panels in PBA (PBS supplemented with 0.5% BSA, 0.01% NaN_3), and then with fixable viability dye eF506 (Invitrogen, Carlsbad, CA, USA) in PBS. After staining, cells are fixed using Fix/Perm buffer (eBioscience, ThermoFisher Scientific). For the immune panels the cells are then measured at the LSR Fortessa (BD biosciences). For the non-immune panel, cells are stained with the intracellular antibodies in Perm buffer (eBioscience, ThermoFisher) and measured at the Sony SP6800. Data is analyzed using FlowJo v10.9.0.

Statistical analysis

All statistical tests will be performed with a two-sided approach, using a 5% type I error threshold. Confidence intervals (CIs) will be set at 95%, unless otherwise indicated. Missing values will not be imputed. Collected data from patients lost to follow-up will be included in analysis where possible. Variables will be reported as either means with standard error of the mean (SEM; normal distribution), medians with 95% CI (skewed distribution) or frequencies depending on the type and distribution of the data.

Ethics and dissemination

This study is conducted according to the principles of the Declaration of Helsinki (version October 2013), and in accordance with the Medical Research Involving Human Subjects Act (WMO). This study was approved by the Medical Ethics Committee of Amsterdam UMC under registration number 2021.0774. Eligible women with SLE are given a verbal and written explanation of the study and the procedures involved, by their treating physician. Eligible healthy controls are asked by their treating physician whether they are interested in participating in the study, and subsequently are given verbal and written explanation of the study procedures by a member of the research team. Subjects who show interest in participation are contacted by a certified member of the research team. An independent physician is available to answer any question the subject may have. If the subject decides to participate written informed consent will be obtained before performance of any study-related activity.

Through identifying mechanisms underlying failing maternal-fetal tolerance, this study will identify potential therapeutic targets and biomarkers for development of pregnancy complications in women with SLE. In the long term, this may improve the development of preventive and therapeutic measures to reduce the incidence of pregnancy complications in SLE. Additionally, these findings may also be relevant to understand the pathogenesis of pregnancy complications in the general population.

The burden and risks of participation are negligible, considering the placenta is a waste product after child-birth and the blood collection will be performed simultaneously with routine clinical care as much as possible. There is no individual benefit from participation in this study.

294   Data will be disseminated through peer-reviewed publications in scientific journals, scientific
295   presentations and through articles published in patient magazines and presentations for patient
296   organizations.
297   Results from additional analyses performed on the stored biomaterials in a later stage, such as single-
298   cell RNA sequencing, will be deposited in the appropriate data repositories (e.g. GEO Datasets).

REFERENCES

1.   Arnaud L, Van Vollenhoven R. Advanced Handbook of Systemic Lupus Erythematosus [Internet]. Cham: Springer International Publishing; 2018 [cited 2024 Oct 29]. Available from: http://link.springer.com/10.1007/978-3-319-43035-5

2.   Tsokos GC. The immunology of systemic lupus erythematosus. Nat Immunol. 2024 Aug;25(8):1332–43.

3.   Kroese SJ, Abheiden CNH, Blomjous BS, van Laar JM, Derksen R, Bultink IEM, et al. Maternal and Perinatal Outcome in Women with Systemic Lupus Erythematosus: A Retrospective Bicenter Cohort Study. J Immunol Res. 2017;2017:8245879.

4.   He WR, Wei H. Maternal and fetal complications associated with systemic lupus erythematosus: An updated meta-analysis of the most recent studies (2017-2019). Medicine (Baltimore). 2020 Apr;99(16):e19797.

5.   Abheiden CNH, Blomjous BS, Slaager C, Landman AJEMC, Ket JCF, Salmon JE, et al. Systemic lupus erythematosus is associated with an increased frequency of spontaneous preterm births: systematic review and meta-analysis. American Journal of Obstetrics and Gynecology. 2024 Oct;231(4):408-416.e21.

6.   Blomjous BS, de Vries JIP, Zijlstra E, Cramer K, Voskuyl AE, Bultink IEM. Desire to have children and preferences regarding to pre-pregnancy counselling in women with SLE. Rheumatology (Oxford). 2021 Jun 18;60(6):2706–13.

7.   Clowse ME, Chakravarty E, Costenbader KH, Chambers C, Michaud K. Effects of infertility, pregnancy loss, and patient concerns on family size of women with rheumatoid arthritis and systemic lupus erythematosus. Arthritis Care Res (Hoboken). 2012 May;64(5):668–74.

8.   Orefice R. Immunology and the immunological response in pregnancy. Best Pract Res Clin Obstet Gynaecol. 2021 Oct;76:3–12.

9.   Hong S, Banchereau R, Maslow BL, Guerra MM, Cardenas J, Baisch J, et al. Longitudinal profiling of human blood transcriptome in healthy and lupus pregnancy. J Exp Med. 2019 May 6;216(5):1154–69.

10.   Buyon JP, Kalunian KC, Ramsey-Goldman R, Petri MA, Lockshin MD, Ruiz-Irastorza G, et al. Assessing disease activity in SLE patients during pregnancy. Lupus. 1999;8(8):677–84.

11.   Buyon JP, Kim MY, Guerra MM, Laskin CA, Petri M, Lockshin MD, et al. Predictors of Pregnancy Outcomes in Patients With Lupus: A Cohort Study. Ann Intern Med. 2015 Aug 4;163(3):153–63.

12.   Gladman D, Ginzler E, Goldsmith C, Fortin P, Liang M, Sanchez-Guerrero J, et al. The development and initial validation of the systemic lupus international collaborating clinics/American college of rheumatology damage index for systemic lupus erythematosus. Arthritis & Rheumatism. 1996 Mar;39(3):363–9.

13.   Khong TY, Mooney EE, Ariel I, Balmus NCM, Boyd TK, Brundler MA, et al. Sampling and Definitions of Placental Lesions: Amsterdam Placental Workshop Group Consensus Statement. Arch Pathol Lab Med. 2016 Jul;140(7):698–713.

14.   Brown MA, Magee LA, Kenny LC, Karumanchi SA, McCarthy FP, Saito S, et al. The hypertensive disorders of pregnancy: ISSHP classification, diagnosis & management recommendations for international practice. Pregnancy Hypertens. 2018 Jul;13:291–310.

15.   Garcia-Flores V, Xu Y, Pusod E, Romero R, Pique-Regi R, Gomez-Lopez N. Preparation of single-cell suspensions from the human placenta. Nat Protoc. 2023 Mar;18(3):732–54.

16.   Krop J, van der Zwan A, Ijsselsteijn ME, Kapsenberg H, Luk SJ, Hendriks SH, et al. Imaging mass
cytometry reveals the prominent role of myeloid cells at the maternal-fetal interface. iScience. 2022
Jul 15;25(7):104648.

AUTHOR CONTRIBUTIONS
WD, MB, LB and IB designed the study, with input on patient information and study setup from KC
and DCRH. WD and JR wrote the manuscript. WD and APS designed the study protocols. SG and MG
collect clinical data. All authors provided input on the manuscript and approved the final version.
355

356 **FUNDING STATEMENT**
357 This work was supported by research grants from FOREUM Foundation for Research in
358 Rheumatology, Amsterdam UMC and the NVLE Foundation.

360 **COMPETING INTERESTS STATEMENT**
361 The authors declare no competing interests.
362

363 **PATIENT AND PUBLIC INVOLVEMENT**
364 Two patient research partners were involved in the design of the study, critically reviewed the patient
365 information folders and contributed to this manuscript.
366

367 **ACKNOWLEDGEMENTS**
368 We would like to thank the NVLE and Lupus Europe for their input and support for the FaMaLE study.
369 In addition, we acknowledge the European Reference Network on Rare and Complex Connective
370 Tissue Diseases (ERN ReCONNET) for declaring the Department of Rheumatology and Clinical
371 Immunology of our centre as a member.