Single-cell RNA sequencing of peripheral blood mononuclear cells in patients with acutely decompensated cirrhosis reveals a specific monocyte subset associated with an increased risk of progression to ACLF

Authors:

Theresa H. Wirtz[1]†, Sara Palomino-Echeverria[2,3,4]†*, Maike R. Pollmanns[1], Estefanía Huergo[2], Felix Schreibing[5], Johanna Reißing[1], Cristina Sánchez-Garrido[6], Cristina López-Vicario[6,7,8], Ana M. Aransay[8,9], Maurizio Baldassare[10], Giacomo Zaccherini[10,11], Enrico Pompili[10,11], Martin Schulz[12], Frank E. Uschner[12], Sabine Klein[12], Wenyi Gu[12], Robert Schierwagen[12], Shantha Valainathan[13], Annabelle Verbeeck[13], Daniela Campion[14], Ilaria Giovo[14], Alexander Koch[1], Rafael Kramann[5,15], Tony Bruns[1], Narsis Kiani[16], Paolo Caraceni[10,17], Carlo Alessandria[14], Richard Moreau[6,13,18], Jonel Trebicka[6,12], Joan Clària[6,7,8,19], Núria Planell[20]*, Pierre-Emmanuel Rautou[6,13,18#]*, Christian Trautwein[1,21,22#]*, David Gomez-Cabrero[2,23#]*

Affiliations:

[1]Medical Department III, University Hospital RWTH Aachen, Aachen, Germany

[2]Translational Bioinformatics Unit (TransBio), Navarrabiomed, Navarra Health Department (CHN), Public University of Navarra (UPNA), Navarra Institute for Health Research (IdiSNA), Pamplona, Spain

[3]Research Program in Systems Oncology, University of Helsinki, Helsinki, Finland

[4]Institute for Molecular Medicine Finland, Helsinki Institute for Life Sciences, University of Helsinki, Helsinki, Finland

[5]Department of Medicine 2 (Nephrology, Rheumatology, Clinical Immunology and Hypertension), RWTH Aachen University, Medical Faculty, Aachen, Germany

[6]European Foundation for the Study of Chronic Liver Failure (EF CLIF) and Grifols Chair, Barcelona, Spain

[7]Biochemistry and Molecular Genetics Service, Hospital Clinic-IDIBAPS, Barcelona, Spain

[8]CIBERehd, Madrid, Spain

[9]Center for Cooperative Research in Biosciences (CIC bioGUNE), Derio, Bizkaia, Spain

[10]Unit of Semeiotics, Liver and Alcohol-related Diseases, IRCCS Azienda Ospedaliero-Universitaria di Bologna, Bologna, Italy

[11]Department of Medical and Surgical Sciences (DIMEC), Alma Mater Studiorum, University of Bologna, Bologna, Italy

[12]Department of Internal Medicine B, University Hospital Muenster, Muenster, Germany

[13]AP-HP, Hôpital Beaujon, Service d´Hepatologie, DMU DIGEST, Centre de Référence des Maladies Vasculaires du Foie, FILFOIE, ERN RARE-LIVER, Clichy, France

[14]Division of Gastroenterology and Hepatology, A.O.U. Città della Salute e della Scienza di Torino, Torino, Italy

[15]Department of Internal Medicine, Nephrology and Transplantation, Erasmus Medical Center, Rotterdam, The Netherlands

[16]Algorithmic Dynamics lab, Department of Oncology-Pathology, Karolinska Institutet, Solna, Sweden

[17]IRCCS Azienda Ospedaliera-Universitaria di Bologna, Bologna, Italy

[18]Université Paris-Cité, Inserm, Centre de recherche sur l´inflammation, UMR 1149, Paris, France

[19]Department of Biomedical Sciences, University of Barcelona, Spain

[20]Universidad de Navarra, Centro de Investigación Médica Aplicada (CIMA), Computational Biology Program, Instituto de Investigación Sanitaria de Navarra (IdiSNA), Pamplona, Spain

[21]Leibniz Research Centre for Working Environment and Human Factors at the TU Dortmund (IfADo), Dortmund, Germany

[22]Liver Center, Department of Gastroenterology, Hepatology and GI-Oncology, Klinikum Stuttgart, Stuttgart, Germany

[23]Bioscience Program, Biological and Environmental Sciences and Engineering Division (BESE), King Abdullah University of Science and Technology KAUST, Thuwal, Saudi Arabia

†: These authors contributed equally to this work and share first authorship. Either name may be listed first when citing.

#: These authors contributed equally to this work and share last authorship.

*: **Corresponding authors:** sara.palominoecheverria@helsinki.fi, nplanellpic@unav.es, pierre-emmanuel.rautou@inserm.fr, trautwein.christian@gmx.net, david.gomez.cabrero@navarra.es

Abstract

Background: Patients with acute decompensation (AD) of cirrhosis are at high risk of developing acute-
on-chronic liver failure (ACLF), a syndrome characterized by multiple organ failure and high short-term
mortality. This study aimed to analyse alterations in peripheral blood mononuclear cells (PBMCs) in
patients with AD using single-cell technologies and to explore their implications for risk stratification
and prognosis.

Methods: A total of 63 AD patients and 15 healthy donors were enrolled at five European centers, with
blood samples collected upon hospital admission. Patients were monitored for 90 days after study
inclusion, focusing on hospital readmission related to AD and the development of ACLF. Cellular
indexing of transcriptomes and epitopes by sequencing (CITE-seq) was performed on PBMCs from 16
patients (n=6 without readmission, n=6 with readmission, and n=4 with ACLF development) along with
4 healthy controls. The whole blood of the same 16 patients was assessed by bulk RNA-seq. The whole
blood transcriptional profiles from two independent prospective multicentre cohorts of patients with
AD (PREDICT (n=689) and ACLARA (n=521)) were included as validation cohorts.

Results: Cell proportion analysis revealed a significant increase of classical monocytes in patients with
subsequent ACLF development. Notably, classical monocytes represented the cell type with the
highest transcriptional alterations. Within the classical monocyte population, a specific subcluster,
termed "C2", was identified and found to be enriched during progression towards ACLF. This
monocytic subpopulation was primarily characterized by decreased expression of genes encoding
enzymes relevant to cellular energy supply, such as oxidative phosphorylation and ATP production.
Pathway analysis confirmed a marked impairment of energy metabolism pathways within the C2
subcluster. A gene signature derived from that monocytic subpopulation was significantly increased in
patients with bacterial infection, ACLF development and non-survivors of two independent
multicenter cohorts of patients with AD.

Conclusion: Patients with acute decompensation of cirrhosis who subsequently develop ACLF are
characterized by a subpopulation of classical monocytes exhibiting features of impaired energy
metabolism pathways.

Introduction

Acute decompensation (AD) is the most frequent cause of hospitalization among individuals diagnosed with cirrhosis (1). It involves the emergence of one or more clinically relevant complications, such as ascites, hepatic encephalopathy, or gastrointestinal bleeding (2–4). Patients with AD are at high risk of developing acute-on-chronic liver failure (ACLF) – a clinical syndrome marked by the acute deterioration of liver function in individuals with AD accompanied by the failure of one or more major organs or systems (5,6). Depending on the severity and number of organ failures, ACLF is associated with poor short-term mortality ranging from 15% to 80% (7).

The transition from decompensated cirrhosis to ACLF is a dynamic and multifactorial process driven by a complex interplay of systemic inflammation (SI), immune dysfunction, and organ crosstalk (8,9). Significant progress has been made in understanding the underlying pathophysiological mechanisms as well as identifying the pivotal mediators that initiate and regulate these processes. Particularly, SI was identified to gradually increase with the progression of portal hypertension and to induce organ dysfunction as part of ACLF transition (10,11). Respective studies have identified increased circulating levels of pro-inflammatory cytokines and the activation of innate immune pathways, including Toll-like receptor (TLR) signaling, as key contributors (12,13). Furthermore, the translocation of microbial products from the gut into systemic circulation, a consequence of intestinal barrier dysfunction, has been implicated in exacerbating SI and immune activation (14). In parallel, cellular studies have revealed immune cells to exhibit phenotypic and functional alterations that promote SI and impair immune response. Dysfunctional neutrophils and monocytes have been identified to impair innate immunity and increase susceptibility to bacterial infection – a common trigger of ACLF (15–17).

Despite these advances, the relative contributions of different immune cell subsets during disease progression are incompletely defined, and the identification of functional cellular states associated with ACLF development remains an area of investigation. Single-cell technologies, including single-cell RNA sequencing (scRNA-seq), have emerged as transformative tools for dissecting cellular heterogeneity and capturing the dynamic immune landscape in complex diseases. Hence, recent studies have used scRNA-seq to elucidate pathophysiological characteristics of peripheral immune cells in patients with ACLF compared to healthy controls (18–21). They identified distinct immunometabolic cellular states contributing to ACLF outcome, with the most significant changes in pro-inflammatory monocytes and T-cells (18–21). However, those publications focused on patients who already transitioned to ACLF. Yet, no prospective, high-resolution baseline analysis of functional alterations of peripheral immune cells in patients with AD who subsequently develop ACLF is available.

We, therefore, conducted a prospective multicenter study involving patients admitted due to AD at five different European centers. The aim of the study was to characterize the PBMCs at single-cell resolution. Our data provide insights into changes within peripheral immune cell subpopulations associated with AD progression, as well as molecular mechanisms driving ACLF development.

Material and Methods

Study design

A prospective multicenter study was conducted in 5 European centers (Hôpital Beaujon, Université Paris-Cité, France; Goethe University Frankfurt, Germany; RWTH Aachen University, Germany; University of Bologna, Italy; University of Turin, Italy) between October 2020 and July 2021. Patients who were admitted to the hospital due to AD (ascites, overt hepatic encephalopathy, new onset of non-obstructive jaundice, gastrointestinal hemorrhage, and/or bacterial infections) met the inclusion criterion and were screened for exclusion criteria (**Supplementary Table 1**). A total of 63 patients with AD and 15 healthy controls were recruited. Written informed consent was obtained from the patient, her/his spouse, or the appointed legal guardian upon hospital admission. Clinical data were collected, and sample acquisition was performed within 72 hours after hospital admission in all included patients. Patients were followed up for 90 days, and readmission due to AD, development of ACLF, and outcome were documented (for detailed study design and visit schedule, **Supplementary Figure 1** and **Supplementary Table 2**). The study protocol was approved by the local ethics committee of each participating center (Paris, Frankfurt, Aachen, Bologna, Turin) and conducted in accordance with the Declaration of Helsinki.

Sample collection and PBMC isolation

After study inclusion, peripheral blood was drawn from all participants from the *discovery* cohort, using BD vacutainer CPT tubes (BD Biosciences; (22)) for PBMC isolation and TempusTM tubes (Applied Biosystems) for whole blood assessment. After density gradient centrifugation at 1500 RCF for 20 minutes at 20°C PBMCs were aspirated and then cryopreserved using fetal calf serum with 10% DMSO. The cells were gradually frozen and stored at -80°C for several weeks until library preparation and sequencing or flow cytometry analysis, respectively.

Group allocation and patient selection strategy for single-cell analysis

90 days after study inclusion, patients were classified according to their clinical course as no readmission due to AD ("stable decompensated cirrhosis", SDC, n=46), readmission due to AD ("unstable decompensated cirrhosis", UDC, n=8) or ACLF development within 28 days ("pre-ACLF", n=10). The diagnostic criteria of ACLF were according to the EF-CLIF definition (5). Of the 63 recruited patients, samples from 4 pre-ACLF, 6 UDC, and 6 SDC patients, along with four healthy controls, were selected to represent the entire study cohort and subjected to CITE-seq profiling (detailed in **Supplementary Material** and **Supplementary Tables 3, 4, and 5**). For baseline characteristics analysis, categorical variables were summarized as counts (percentages), and continuous variables as median (interquartile range, IQR). The Wilcoxon–Mann–Whitney test was used for continuous variables, and the chi-square test for categorical variables.

CITE-seq

Library preparation, sequencing, and demultiplexing

The isolated PBMCs were recovered by gently thawing and immediately diluting in pure FCS. Next, cells were centrifuged at 500 RCF for 7 minutes at 20°C, and the supernatant was removed. PBMCs were then resuspended in 4 ml 5% FCS in PBS and filtered through a 20 µm pluriStrainer® provided by pluriSelect (Leipzig, Germany) to obtain a single-cell solution. After the PBMCs were passed through a strainer, a second centrifugation was performed. The supernatant was removed, and cells were

diluted in 100 µl 5% FCS in PBS. 5 µl Human TruStain FcX (BioLegend) was added and incubated for 10
minutes at 4°C.

A panel of 23 TotalSeq-C antibodies (BioLegend) was pooled, using 0.5 µg of each antibody
(**Supplementary Table 6**). Samples were incubated with the TotalSeq-C antibody pool for 30 min at
4°C. Cells were washed two times using 3 ml 5% FCS in PBS and rediluted in 1.5 ml of sorting buffer
(5% FCS in PBS).

For the detection of dead cells, DAPI was added at a final concentration of 0.5 µg/ml. Up to 300,000
cells from each sample were sorted on a Sony Cell Sorter by life/dead sorting. Sorted viable cells were
resuspended in 10 µl 1% BSA in PBS, counted, and ~20,000 cells in 38.7 µl were applied to single cell
partitioning and barcoding *via* Chromium Controller (10x Genomics).

Each participant's sample was prepared into two distinct libraries: (I) 5' Gene Expression Library (GEX),
designed for scRNA-seq to capture transcriptome data; and (II) Surface Protein Library, which utilized
TotalSeq-C barcodes to profile surface proteins, aligning with the CITE-seq methodology.

Cell Ranger (v6.01) was used for read alignment, filtering, barcode processing, and UMI counting with
default settings (23). GRCh38-2020 was used as the transcriptome reference. Specifically, STAR was
used in FASTQs alignment (24). Cell barcodes were then automatically determined based on the
distribution of the UMI count.

Pre-processing
We used the R package *Seurat* v4.3.0 (25) to preprocess the gene-barcode matrix and the surface
protein matrix of both patients and healthy controls and to perform further analysis. For each sample,
genes not observed in at least three cells were dropped. Low-quality or damaged cells were excluded
using a combination of a number of expressed genes (cells with <300 and >3500 genes) and
mitochondrial transcript percentages (cells with >10% mitochondrial genes). Each sample dataset was
normalized using the *'SCTransform'* function (26) from the *Seurat* package and regression of
mitochondrial RNA and cell cycle genes.

Integration, clustering, and cell type identification
The filtered and cleaned single-cell dataset underwent a process to identify highly variable genes and
remove unwanted sources of variation. This was done by calculating and extracting a subset of
features that show high cell-to-cell variation in the dataset using the '*FindVariableFeatures'* Seurat's
function. Additionally, the *'FeatureScatter'* plot from *Seurat* was applied, and we removed the
detected outliers when plotting feature-feature relationships (27). Harmony was used to correct for
batch effects (28). For each sample where the protein surface markers data was available, we selected
the most variable protein sub-space and applied feature-wise centered log-ratio transformation (CLR)
(29). Data were visualized using UMAP. Data were scaled, and the dimensionality was reduced by PCA.
To identify cell clusters, a shared nearest neighbor (SNN) graph was built using distances in PCA space
(top 1 to 20 components) through Seurat's *'FindNeighbors'* function. The Louvain algorithm was then
employed to iteratively cluster neighboring cells using Seurat's *'FindClusters'* function, using a
resolution parameter of 0.5. Cell type annotation was done by combining three parallel evaluations:
gene expression of canonical marker genes (**Supplementary Figure 2**), surface protein markers

(**Supplementary Figure 3 A, B**), and Azimuth results (a reference-based approach using an external
well-annotated dataset of PBMCs (25)).

Cluster Composition Analysis: changes in cell proportions between groups
Differences in cell type proportion between conditions were assessed by comparing the proportional
differences in cell populations between AD patients and healthy controls for both gene expression and
protein surface markers data, utilizing the *'sc_utils'* function from *scProportionTest* package with
default parameters (30). The estimations of the p-value and the confidence interval of the magnitude
of the difference are computed respectively using a permutation test and bootstrapping.

Differential expression analysis, pathway analysis, and trajectory analysis
Seurat *'FindAllMarkers'* function with the default Wilcoxon rank sum test based on the normalized
data was used to identify the cluster markers used for cluster annotation. The '*FindMarkers'* function
was employed to identify markers for each cell cluster to measure differential gene expression
between conditions, using the default Wilcoxon rank sum test method. Genes detected in less than
50% of both evaluated groups were discarded. We considered differentially expressed genes (DEG) to
be statistically significant if the adjusted p-value < 0.05 (Benjamini-Hochberg (BH)) and absolute Fold
Change (FC) > 1.5. '*VlnPlot'*, '*FeaturePlot*,' and '*Dotplot'* functions were applied to visualize gene
expression in each cell type or group.

Next, Gene Set Enrichment Analysis (GSEA) was performed using the R package *ClusterProfiler* (31),
considering the comprehensive functions included in Gene Ontology (GO). Biological pathways in
which genes appeared at a different frequency than expected were considered enriched. The results
were visualized using default enrichment plots. Enrichment scores with p-value adjusted < 0.05 were
considered significant. The differences in the pathways among patient groups were computed using
the non-parametric Kruskal Wallis test.

Single-cell trajectory analyses were computed using the *'learn_graph'* function from the *Monocle3*
package with default parameters (32). We then plot the cells based on their *pseudotime*.

Module scores for specific gene sets
We utilized cell module scores to quantify the extent of individual cell expression of specific
predefined gene sets. For this purpose, we used the '*AddModuleScore'* function from *Seurat* with
default settings. The lists of genes defining this module were prepared based on the intersection of
upregulated DEGs resulting from various differential expression analyses (DEA). Differences in module
scores were assessed using the non-parametric Kruskal Wallis test.

Bulk RNA-seq

Library preparation, sequencing and demultiplexing
Bulk RNA-seq was performed in collaboration with the Genomics platform of the Center for
Cooperative Research in Biosciences (CIC bioGUNE), Derio, Bizkaia, Spain. RNA was extracted from
whole blood collected in Tempus™ tubes from the *discovery* cohort using the Extraction Tempus™
Spin RNA Isolation Kit (Applied Biosystems). The extracted RNA was treated with DNase. RNA quality
was assessed using Agilent RNA 6000 Nano and Pico Chips (Agilent Technologies). RNA concentration
was assessed using RNA HS Assay Kit in a Qubit 2.0 fluorometer (Thermo Fisher Scientific).

Sequencing libraries were prepared using the NuGEN UniversalPlus mRNA-Seq with NuQuant kit
(Tecan Life Sciences, Cat.#0520-24) and following the user guide (M01485v6.1andv7). The kit is
intended to prepare sequencing libraries depleted of globin. Briefly, starting from 500 ng of total RNA,
rRNA, and globin mRNA were depleted, and the remaining RNA was purified, fragmented, and primed
for cDNA synthesis. cDNA first strand was synthesized with SuperScript-II Reverse Transcriptase
(Thermo Fisher Scientific, Cat.#18064-014) for 10 min at 25°C, 15 min at 42°C, 15 min at 70°C and
pause at 4°C. The second strand of cDNA was synthesized with Illumina reagents at 16°C for 1 hour.
Then, A-tailing and adaptor ligation were performed. Finally, enrichment of libraries was achieved by
PCR (30 s at 98°C; 15 cycles of 10 s at 98°C, 30 s at 60°C, 30 s at 72°C; 5min at 72°C and pause at 4°C).

Afterward, libraries were visualized on an Agilent 2100 Bioanalyzer using Agilent High Sensitivity DNA
kit (Agilent Technologies, Cat. # 5067-4626), quantified using Qubit dsDNA HS DNA Kit (Thermo Fisher
Scientific, Cat. # Q32854) and sequenced in a NovaSeq-6000 (Illumina Inc.) by at least 40 million
paired-end 100nt reads.

Raw reads were demultiplexed via bclfastq2 Conversion Software v2.20 (Illumina) to convert base call
(BCL) files into FastQ files. Quality control of FastQ files was performed using FastQC (Bioinformatics
Babraham Institute) (33), and reads were trimmed for contaminating sequence adapters and poor-
quality bases using Trimmomatic (34). Sequencing reads were aligned to the Human (GRCh38) genome
with STAR (2.6.1) setting the parameters to default values (24). Uniquely aligned reads were assigned
to GENCODE18 gene annotations using HTSeq v0.11 and quantified with htseq-counts run in stranded
mode with default parameters (35).

Bulk RNA-seq downstream analysis
R v4.3.0 was employed in the downstream analysis of RNA-seq data. Genes exhibiting more than 1
count per million (CPM) in at least three samples were selected for the analysis. Utilizing the *limma-*
voom workflow (36), RNA-seq count data was normalized, and variance estimation was performed.
PCA was applied to the normalized data to assess sample similarity.

Module scores
A module score was calculated for each evaluated sample group using the Gene Set Variation Analysis
(*GSVA*) R package v1.42.0 (37) in the single-cell paired bulk RNA-seq, as well as in the PREDICT and the
ACLARA cohorts. This gene signature was calculated using previously selected genes to compute the
module score at the single-cell level.

Flow Cytometry Analysis
Flow cytometry of peripheral CD14[+] monocytes was conducted. For this purpose, additional PBMC
aliquots from the discovery cohort were used. First, PBMCs were thawed gently, as described above.
Subsequently, CD14[+] cells were enriched by magnetic cell separation using positive selection for
human CD14 (Miltenyi Biotec, Bergisch Gladbach, Germany) for in vitro experiments.

For surface marker expression analysis, CD14[+] peripheral monocytes were incubated with
fluorochrome-conjugated primary antibodies at optimal dilutions at 4°C for 30 min in PBS containing
0.5% fetal calf serum and 2 mmol/L EDTA. Single-color staining, isotype-matched controls (IMCs), and
fluorescence minus one (FMO) controls were applied. The antibodies used were CD14 (clone HCD14),
CD16 (3G8), and human leukocyte antigen-DR (HLA-DR) (G46-6).

Cell viability was determined with the LIVE/DEAD™ Fixable Aqua Dead Cell Stain Kit from Thermo
Fisher and IMC mouse IgG1 κ and IgG2a κ isotype controls from BD Pharmingen (BD Biosciences,
Franklin Lakes, NJ, USA). Flow cytometry analysis was performed using a FACS Canto II flow cytometer
and FlowJo (v10) software (BD Biosciences, Franklin Lakes, NJ, USA). Delta mean fluorescence intensity
(ΔMFI) was calculated by subtracting the mean MFI count in the SDC group from single MFI counts in
the four study groups. The Wilcoxon–Mann–Whitney test was used to test statistical differences
within the four study groups.

Validation cohorts
Validation studies were conducted using whole blood bulk RNA-seq data from two prospective,
multicenter European and Latin America cohorts, PREDICT and ACLARA (7,38). The inclusion and
exclusion criteria of both cohorts are shown in **Supplementary Tables 7 and 8**. Both cohorts collected
various measures, including clinical, pharmacological, and outcome data from AD patients upon
hospital admission and during follow-up visits (**Supplementary Tables 9 and 10**). The follow-up period
was 90 days for PREDICT and 28 days for ACLARA (**Supplementary Figures 5 and 6**). 90 days after study
inclusion, AD patients from *PREDICT* cohort were categorized into SDC/UDC and pre-ACLF (7).
ACLARA patients were classified either into AD or pre-ACLF. The diagnostic criteria of ACLF in both
cohorts were according to the EF-CLIF definition (5).

The criteria for inclusion in our investigation were AD at hospital admission, availability of data on
short-term outcomes (liver transplant, ACLF development, or death), clinical data, treatment, and
availability of biological samples (buffy coat, tempus, serum, and plasma). Based on these criteria, 768
AD patients from PREDICT were chosen for analysis, with 738 samples passing quality control after
sequencing. According to the definition of ACLF within the discovery cohort, only patients who
developed ACLF within 28 days after hospital admission were included (n=61, **Supplementary Figure
7**). For the ACLARA cohort, the same criteria applied, with the added requirement that samples be
available at the Barcelona EFCLIF biobank by the end of March 2021 due to COVID-19-related delays
in shipping from South America. From ACLARA, 549 AD cirrhosis patients were selected, with 32
lacking buffy coat samples and 27 failing quality control after sequencing (**Supplementary Figure 8)**.
In total, 689 patients from PREDICT and 521 patients from ACLARA were included to validate the
single-cell results.

PREDICT & ACLARA bulk RNA-seq
Whole blood RNA-seq from 1,210 patients with AD included in the PREDICT (n=689) (7) and the
ACLARA (n=521)(38) studies were conducted according to the above-described bulk RNA-seq protocol.

A module score was calculated for each evaluated sample group, obtaining a gene signature score, as
described in Material and Methods: Module scores. Pearson (parametric) and Spearman (non-
parametric) correlation coefficients were computed to assess the relationships between the module
score and clinical variables. The Area Under the Receiver Operating Characteristic (AUC-ROC) curve
using the "*pROC*" R package (39) was utilized to evaluate the relationship between the module score
and outcomes (40). The comparison of ROC curves was performed using DeLong's Test. The analysis
of mean differences in the computed module scores among patient groups was assessed using the
ANOVA test when more than two groups were compared and the Wilcoxon Signed-Rank test
otherwise.
314

315 *Statistical Methods*

316 All analyses were performed in the R Computing Environment version 4.0.3 (41). Statistical methods

317 are described in detail above in each subsection. In all statistical analyses, we considered p-value <

318 0.05 as statistically significant. In case an adjusted p-value for multiple testing is considered, it is

319 described in the text.

Results

Single-cell study cohort and sample description

In total, 63 patients with AD and 15 healthy controls were prospectively recruited in 5 European centers. After 90 days, patients were classified as having no readmission due to AD (SDC), readmission due to AD (UDC), or ACLF development within the first 28 days (pre-ACLF). Of those, 16 patients (6 SDC, 6 UDC, 4 pre-ACLF) and four healthy controls were selected for single-cell analysis (see Material and Methods; **Figure 1 A**). Respective PBMC samples subsequently underwent CITE-seq; bulk RNA-seq on whole blood from the same patients was also performed (**Figure 1 B**). The median age at the time of hospital admission was 60 years in the SDC, 57 years in the UDC, and 74 years in the pre-ACLF group (**Table 1**). Of all patients, 75% were males, and 50% had been diagnosed with alcohol-related liver cirrhosis. Previous episodes of acute decompensation were common in the UDC (4/6 patients) and pre-ACLF (3/4 patients) groups. The primary reason for hospital admission in all groups was ascites. Bacterial infection was present in 2 out of 6 SDC patients, and spontaneous bacterial peritonitis (SBP) was evident in 2 out of 4 pre-ACLF patients. MELD and CLIF-C AD scores were significantly higher in the pre-ACLF group compared to the SDC and the UDC groups (20 and 59 points, respectively). Treatment at blood draw was not significantly different between the groups. All pre-ACLF patients developed ACLF during their first hospital admission. Within the pre-ACLF group, one patient died, and one patient underwent a liver transplant during follow-up. Additional baseline characteristics are presented in **Supplementary Tables 3, 4, and 5**.

An increase in circulating classical monocytes is associated with ACLF development

For the single-cell transcriptional analysis, 130,846 cells, with a median of 1,632 genes per cell, were recovered from the 20 individuals included in this study (**Figure 1 A**). We identified 11 cell clusters corresponding to classical monocytes, non-classical monocytes, CD4$^+$ T cells, CD8$^+$ T cells, B cells, NK cells, MAIT cells, dendritic cells, regulatory T cells, dnT cells, and gdT cells (**Figure 2 A, Supplementary Figures 2**). Cell annotation was confirmed by cell surface marker profiling (**Supplementary Figure 3 A, B**).

Aiming to identify changes in cellular proportions associated with disease progression, we compared the cell composition among healthy controls, SDC, UDC, and pre-ACLF groups. We observed marked proportion variations when investigating the most prevalent cell subtypes (those representing at least 1% of the total cells) (**Figure 2 B, C**). As expected, we found a significant increase of classic monocytes in AD patients compared to healthy controls, with pre-ACLF exhibiting the highest fold change, followed by UDC and SDC. Non-classical monocytes also increased significantly in the pre-ACLF patients but to a lesser extent. Regarding T cell populations, CD8$^+$ T cells in AD patients decreased compared to healthy controls. Similar trends were observed in cell proportions derived from surface protein markers´ data (**Supplementary Figure 3 C, D**).

To examine the transcriptional changes within each cell population, we conducted a differential expression analysis (DEA) across all groups of AD patients and healthy controls. We identified varying numbers of differentially expressed genes (DEGs) in each cell type and contrast (SDC vs. healthy controls, UDC vs. healthy controls, pre-ACLF vs. healthy controls) (**Figure 2 D**). Classical monocytes showed the highest number of DEGs in both the total of all comparisons and each comparison

individually. In detail, pre-ACLF comparisons with SDC and UDC, respectively, showed that they exhibited the highest number of DEGs, especially in classical monocytes (**Figure 2 E and Supplementary Figure 9**).

Altogether, we identified perturbations in the peripheral immune cell landscape in AD patients compared to healthy controls, with a notable accentuation of the monocytic population, especially in the pre-ACLF patients.

Identification of classical monocyte subpopulations and their association with AD progression

Given the observed alterations in both the cell proportion and gene expression of classical monocytes in AD patients compared to healthy controls, we aimed to further explore the role of this cell population in disease progression by identifying distinct classical monocytic subpopulations. To this end, we applied unsupervised clustering to the classical monocytes, identifying three major cell subclusters labelled as C0, C1, and C2 (**Figure 3 A**). C0 constituted the highest proportion of classical monocytes, followed by C1 and C2.

Next, we aimed to investigate differences in the identified monocytic subclusters within the four clinical groups. Similar to the overall monocyte population, an increase in the three monocyte subpopulations was observed in AD patients compared to healthy controls (**Figure 3 B, Supplementary Figure 10**). Specifically, C0 increased significantly in all AD groups, C1 was significantly enriched in UDC and pre-ACLF patients, and C2 only in the pre-ACLF patients (**Figure 3 C**). Interestingly, within the pre-ACLF group, the proportions of C1 and C2 monocytes increased according to the latency of ACLF onset (**Figure 3 B)**. To further investigate the dynamic processes underlying the monocyte compartment, trajectory analysis was complemented, which identified a continuous process from C0 to C2, with C1 representing a transitional state between C0 and C2 monocytes (**Figure 3 D**).

In the next step, we aimed to elucidate the functional and pathophysiological characteristics of the C2 subset. We first explored the transcriptional profile of monocytic markers with known biological function in these cells (**Figure 3 E**). In contrast to the C0 subset, we observed lower expression of *CD14*, as well as of genes encoding HLA-DR subunits (*HLA-DRA*) or isoforms (*HLA-DRB1*) in the C2 cluster. Moreover, the expression of genes encoding pro-inflammatory mediators from the S100 protein family, as well as antibacterial agents such as lysozyme (*LYZ*), were decreased within the C2 subset. On the other hand, genes encoding proteins with regulatory functions in inflammatory and infection-driven processes, such as *MERTK*, *VCAN*, *IRAK3*, *HIF1α*, and *TLR2*, showed increased expression compared to C0. The expression of *CD15* was also higher in C2. C1 monocytes showed an intermediate state between C0 and C2.

We then characterized transcriptional alterations by DEA of C2 subcluster vs. C0 and C1 subclusters. The exploration of the top 100 up- and downregulated genes confirmed the intermediate state of C1 (**Figure 3 F, Supplementary Figure 10**). Interestingly, the upregulated genes illustrated a rather heterogeneous biological pattern. In contrast, among the most downregulated genes, we observed several genes that encode different proteins/enzymes involved in processes relevant to energy production and metabolism. These genes included ATP-production genes (i.e. gene: *ATP5F1E*/protein: ATP synthase F1 subunit epsilon**;** *ATP5MG*/ATP synthase membrane subunit g) and oxidative phosphorylation genes (*UQCR11*/ubiquinol-cytochrome c reductase, complex III sub-unit XI;

COX7C/cytochrome c oxidase subunit 7C; *COX8A*/cytochrome c oxidase subunit 8A). In total, three genes that encode ATP synthase membrane subunits and six that encode cytochrome c oxidase subunits were among the 50 most downregulated genes in the C2 subset compared to C0/C1. Moreover, among the downregulated genes, we found numerous S100 proteins described as important regulators of stress and inflammatory responses (**Supplementary Figure 10**).

Given the transcriptional profile of C2 classical monocytes, showing a decreased expression of the surface receptor *HLA-DR*, we wondered whether this translates at the protein level. Hence, we assessed the HLA-DR profile by flow cytometry on classical monocytes enriched from PBMCs from the same AD and healthy control individuals**.** To analyze differences of the HLA-DR expression between the groups, the delta of the mean fluorescence intensity (ΔMFI) of HLA-DR was calculated by subtracting the mean MFI count in the SDC group from single MFI counts in the four study groups. Even if the entire population of classical monocytes was considered in this approach, we still observed a statistical tendency towards a decreased relative HLA-DR expression in patients with pre-ACLF (**Figure 3 G**).

Thus, we identified a distinct monocytic whose abundance is associated with future ACLF development. This subcluster is predominantly characterized by decreased expression of genes involved in cellular energy supply.

Functional characterization of the classical monocyte subsets

To further elucidate the biological functions of the monocyte subsets identified, we performed gene set enrichment analysis (GSEA) comparing C2 and C0, with C1 considered an intermediate state. Our analysis revealed a significant global downregulation of pathways related to energy metabolism, including "*oxidative phosphorylation (OXPHOS)*" and "*aerobic respiration*" – both representing the most overall regulated pathways in this data set–, as well as "*ATP biosynthetic process*" in the C2 subset (**Figure 4 A** and **Supplementary Table 11**). Additionally, we investigated how these biological processes were regulated in the different subclusters (C0, C1, and C2) (**Figure 4 B**) and in the different patient subgroups (healthy controls, SDC, UDC, and pre-ACLF) (**Figure 4 C**). Importantly, our findings indicated that C2 had significantly lower signature scores than C0 and C1 subclusters independently of the clinical course.

Since bacterial infection represents a complication of cirrhosis-associated immune dysfunction and a critical driver of ACLF progression, we also examined pathways related to the immune response in the C2 monocytes (**Figure 4 D**). Although no significantly regulated pathways were identified based on adjusted p-value, we observed a tendency towards a reduced activity of pathways related to the immune response in C2 monocytes. Among them, we identified "*positive regulation of inflammatory response*", "*defense response to bacterium*", and "*cytokine production*" (**Figure 4 D** and **Supplementary Table 11**).

Hence, C2 monocytes exhibit an extensive impairment of pathways related to cellular energy supply and a tendency towards a dysfunctional immune response.

Identification and translation of a C2 monocytic gene signature to whole blood profiles

Given the identification of a specific population of C2 monocytes expanded in pre-ACLF patients by CITE-seq of the PBMCs and the availability of transcriptional data from the whole blood of the same patients (**Figure 1 B**), we aimed to investigate whether a C2-associated gene signature could still be detected in whole blood in the included patients. First, based on scRNA-seq data, we selected genes to define such a signature by combining three criteria: (i) they were upregulated in classical monocytes when compared with the rest of the cell types; (ii) they were upregulated when comparing C2 with the other subtypes of classical monocytes; and (iii) they were upregulated within C2 when comparing pre-ACLF and healthy controls/SDC/UDC. As a result, we identified 96 upregulated genes that characterize the C2 monocytes subset in pre-ACLF patients (**Figure 5 A**).

Next, we analyzed the expression profiles of these 96 genes at the whole blood level. 94 genes were expressed at the bulk level. Then, we selected those genes with a fold change higher than 25% in bulk RNA-seq, equivalent to log2 FC> 0.32, for the three comparisons: healthy controls/SDC/UDC vs. pre-ACLF. As a result, we identified 18 genes defining the C2-signature at whole blood level: *SLC11A1*, *NAMPT*, *SMAP2*, *CD55*, *ACSL1*, *NEAT1*, *DLEU2*, *IRAK3*, *KYNU*, *TBXAS1*, *FKBP5*, *NUMB*, *PHF21A*, *AOAH*, *ZFAND3*, *LYN*, *SLCO3A1*, and *PTEN* (**Figure 5 B**).

Subsequently, we computed an activity score associated with the defined C2 signature to assess its profile in both single-cell PBMC and whole-blood RNA-seq data (**Figure 5 C, D**). We computed such activity value using a signature score (see Material and Methods: Module scores), in which higher values indicate high activity of the gene sets in a given sample or cell. As expected, we observed a significant progressive increase in the C2 signature activity score from healthy controls to SDC and UDC, culminating in the pre-ACLF group, which showed the highest levels at both the single-cell (PBMCs) (**Figure 5 C**) and bulk (whole blood) levels (**Figure 5 D**).

These findings highlight the efficient translation of the C2 monocytic signature from the cellular to the systemic level and its enrichment in pre-ACLF patients.

Validation of C2 gene signature in two prospective multicenter cohorts of patients with AD

To validate the clinical significance of the defined C2 gene signature, we evaluated its relevance in two large and independent prospective multicenter cohorts of patients with AD from Europe (PREDICT; n=689) and from Latin America (ACLARA; n=521) (**Figure 6 A**). In both cohorts, whole blood samples were collected at hospital admission, and bulk RNA-seq was performed. In PREDICT, patients were followed up for 90 days and classified as SDC, UDC or pre-ACLF based on their clinical trajectory (**Supplementary Figure 4, Supplementary Table 9**). In the ACLARA cohort, patients were followed up for 28 days and classified as either AD or pre-ACLF (**Supplementary Figure 5, Supplementary Table 10**). Detailed description of patient selection from both cohorts are provided in the Material and Methods section and **Supplementary Figures 6 and 7**.

After computing the C2 signature activity score in both the PREDICT and ACLARA cohorts, we observed a significant positive correlation with indicators of liver disease severity, including total bilirubin, MELD, and CLIF-C AD score in patients from both cohorts at hospital admission (**Figure 6 B and Supplementary Tables 12, 13**). In line, it was significantly enriched in the pre-ACLF group as well as in

non-survivors in both cohorts (**Figure 6 C, D**), emphasizing the identified classical monocyte subtype
(C2) to be associated with disease severity, AD progression, and fatal outcome in patients with AD.

In the next step, we asked which pathophysiological processes might primarily be associated with an
increased expression of the C2 gene signature in patients with AD. We performed additional
correlation analysis between the C2 gene signature score and clinical and laboratory parameters of
AD patients from both cohorts at hospital admission. We found a significant positive correlation of the
C2 score with white blood cell count and C-reactive protein in both cohorts (**Supplementary Figure 11**
A and Supplementary Tables 12, 13). Moreover, upon dividing the cohorts into patients with and
without bacterial infection, we observed additionally higher C2 signature activity scores in patients
with bacterial infection compared to those without infection in patients from the SDC and UDC groups
of the PREDICT cohort (**Supplementary Figure 11 B**).

The C2 signature activity score had an AUROC for ACLF or death in both PREDICT and ACLARA cohorts
around 0.65 (**Figure 6 E**). However, combining the newly identified C2 signature score with the CLIF-C
AD score did not significantly improve the latter´s predictive capability regarding outcomes of ACLF
and death (**Supplementary Figure 12**).

In summary, we demonstrated that the C2 signature can be detected in whole blood RNA-seq data
from patients with AD and, most importantly, is associated with the progression to ACLF and mortality
in two independent cohorts. Additionally, we showed that the C2 signature provides predictive value
beyond the presence of bacterial infection.

Discussion

In this European multicenter study, we prospectively investigated the transcriptomic alterations of peripheral immune cells at single-cell resolution in patients with AD. We found that classical monocytes showed the most pronounced transcriptional regulation among the investigated immune cells. A monocytic subpopulation, termed C2, was identified that correlated with the onset of ACLF. This subpopulation was characterized by impaired cellular energy supply. Notably, the transcriptomic signature of these cells remained significantly associated with bacterial infection, ACLF development, and mortality in whole-blood bulk RNA-seq data of two large prospective multicenter cohorts of patients with AD.

While previous studies represent case-control studies focusing on ACLF (18–20), our study is the first to prospectively analyze changes in peripheral immune cells in patients with AD *before* ACLF development. Classical monocytes were identified to show the highest transcriptional regulation of all investigated immune cells during AD. Various studies have consistently reported an overrepresentation of monocytic gene signatures among regulated genes in the blood of patients with manifest ACLF (17,18). Our analysis revealed that those significant transcriptional alterations identified in classical monocytes, in fact, can be observed much earlier in the course of the disease, even in the early stages of AD. These findings indicate monocytes to play a pivotal role in AD and support their involvement in disease progression.

In-depth exploration of classical monocytes identified three major subclusters: C0, C1, and C2. The C2 subcluster showed an expanded presence in pre-ACLF patients (up to 10%) and correlated with latency towards ACLF. C1 appeared as an intermediate state, illustrating the progressive alterations in monocyte populations as cirrhosis advances. Consequently, we focused our analysis on the differences between the C0 and C2 subclusters. Interestingly, the C2 subset was not only detectable in patients with pre-ACLF but – to a lesser extent – already in healthy controls and patients from the other AD groups. This suggests that the C2 monocytic subpopulation already exists in physiological conditions and expands as the disease progresses.

In 2018, Bernsmeier et al. described an expansion of mononuclear myeloid-derived suppressor cells (M-MDSCs, CD14$^+$HLADR$^{lo/-}$CD15$^-$) in ACLF patients associated with impaired antimicrobial response (42). We questioned whether our newly identified C2 subcluster shares features with M-MDSCs. While the expression of *CD14, HLA-DR, CD163, CX3CR1,* and *CCR5* was decreased and pointed in the same direction, the expression of *CD15, MERTK*, and *CD86* was increased, in contrast to the M-MDSC phenotype. Furthermore, the expression of *S100A8* and *S100A9*, which encode for two pro-inflammatory mediators preferentially secreted by M-MDSCs (42), were downregulated in the C2 subset. Thus, we propose that the C2 monocyte subpopulation represents a previously undescribed monocyte subset associated with AD progression.

Additional transcriptomic profiling was conducted to further characterize the C2 subpopulation using marker genes of monocyte differentiation and expansion during cirrhosis progression. Various genes encoding proteins related to inflammatory and immune responses, including interleukin-1 receptor-associated kinase 3 (*IRAK3*), hypoxia-inducible factor 1α (*HIF-1α*), and toll-like receptor 2 (*TLR2*), were upregulated in the C2 subset. In contrast, we observed several genes encoding proteins associated with *suppressed* immune responses to be increasingly expressed in C2 monocytes. For example,

MERTK, a tyrosine kinase receptor observed to be increasingly expressed on immunoregulatory monocytes in patients with ACLF, was upregulated in the C2 subset (43). Further upregulated genes included *VCAN*, which encodes versican and was found to be associated with an immunosuppressive phenotype of monocytes in sepsis (44). On the other hand, distinct genes encoding pro-inflammatory S100 proteins, including *S100A8, S100A9,* and *S100A12*, were downregulated in the C2 subset. Finally, genes relevant to bacterial immune response, such as *CD68*, which is relevant for monocyte adhesion, as well as *LYZ*, encoding for anti-bacterial lysozymes, were decreasingly expressed. In summary, the C2 monocyte subcluster identified in this study is characterized by a distinct transcriptional profile that partially reflects enhanced expression of genes related to immune stimulation. However, genes associated with immunosuppression and, especially, impaired bacterial response were likewise regulated. We hypothesized that the C2 subset reflects evidence of early immunological dysfunction that might favor severe bacterial infections as pivotal precipitators of ACLF. Our hypothesis was validated in whole blood bulk RNA-seq data from two large and independent cohorts, including more than 1200 AD patients (PREDICT and ACLARA). Here, the C2 signature not only positively correlated with liver disease severity but was significantly enriched in the pre-ACLF groups of both cohorts. But, in accordance with our hypothesis that metabolic disturbances within the C2 monocytes favour bacterial infection, the C2 signature was even more evident in AD patients with simultaneous bacterial infections. These findings highlight the importance of the C2 signature not only in cirrhosis progression but also in precipitating events that lead to ACLF.

To explore our findings in more detail, we conducted a pathway analysis uncovering the biological functions of the newly identified C2 monocyte subset. While our analysis showed a clear trend of reduced immune function within the C2 subset, immune-related pathways were not the most significantly regulated. Instead, our analysis revealed a dominant downregulation of genes that encode key enzymes in cellular energy supply. This aligns with reports from several studies, which describe a progressive worsening of metabolic disturbances in patients with compensated cirrhosis, AD, and ACLF, likely driven by systemic inflammation (19,45–48). We here identified the specific monocytic subset that predominantly is characterized by bioenergetic failure and early signs of immune exhaustion, reflecting the inflammation-associated immunometabolic changes associated with cirrhosis progression (49,50).

Our findings are of translational relevance as they underline the hypothesis of systemic inflammation-associated energetic exhaustion in patients before ACLF development. Systemic inflammation in patients with cirrhosis is, in part, assumed to be driven by bacterial translocation. Hence, current approaches targeting gut barrier function and intestinal microbiota diversity, such as rifaximin or fecal microbiota transfer, are investigated to prevent AD or ACLF development in patients with chronic liver disease (51,52). The fact that we identified a monocytic subset that is characterized by bioenergetic failure and expands before ACLF development further supports therapeutic strategies that aim to prevent inflammation-induced bioenergetic failure in patients with AD. Moreover, the identified monocytic subset could serve as a cellular model to develop and test such therapeutic strategies that aim to reduce systemic inflammation-induced immune cell exhaustion.

Despite the valuable insights gained from this study, complementary omics or functional assays are needed to mechanistically validate our findings and pinpoint potential intervention targets. Moreover,

longitudinal single-cell analyses would be beneficial to help understand the dynamic changes in
monocyte subsets over the disease course.

In summary, our analysis presents strong evidence supporting the involvement of monocytes,
particularly a novel subset referred to as C2, in cirrhosis progression and ACLF development. We offer
valuable insights into transcriptional alterations in these cells, highlighting a state of immuno-
metabolic exhaustion that may increase susceptibility to bacterial infections. Our findings support the
idea that preventing the bioenergetic failure of immune cells during cirrhosis progression represents
a promising therapeutic target.
587

Acknowledgements: We thank all participating centres for their efforts in patient recruitment and fruitful collaboration. We thank the Genomics Facility of the Interdisciplinary Center for Clinical Research (IZKF) Aachen for sequencing experiments. Special thanks go to Jasmin Hübner and Anna Rudzinski for their extraordinary help in sequencing the samples.

Conflicts of interest: Jonel Trebicka has received speaking and/or consulting fees from Astra-Zeneca, Gore, Boehringer-Ingelheim, Falk, Grifols, Genfit, and CSL Behring. T Bruns received consulting fees from Intercept/Advanz Pharma, Grifols, Sobi, and SmartDyeLivery. Paolo Caraceni has received speaking and/or consulting fees from Grifols SA , Octapharma SA, CSL Behring, and Gilead.

Data availability statement: Researchers who provide a methodology sound proposal can apply for the data, as far as the proposal is in line with the research consented by the patients. These proposals should be requested through https://www.clifresearch.com/decision/Home.aspx. Data requestors will need to sign a data transfer agreement. Single-cell data is currently being uploaded to the European Genome-phenome Archive (EGA).

Ethics approval statement: The studies involving human participants were reviewed and approved by the Ethical Board of all five participating centres in Europe. The patients/participants provided written informed consent to participate in this study.

Funding:
This project has received funding from the European Union's Horizon 2020 research and innovation program under grant agreement No 847949. This research was made possible through access to data generated by the PREDICT and ACLARA studies promoted and funded by the European Foundation for the Study of Chronic Liver Failure (EF-CLIF), a private, non-profit organization supported by unrestricted grants from Grifols and Genfit.

T.H.W. received financing from the START program of the RWTH Aachen University.

N.P.P. was funded by a Ramón y Cajal fellow (RYC2021-032197-I) from the MCIN/AEI/10.13039/501100011033 and European Union "NextGenerationEU"/PRTR and by a Juan de la Cierva-formación fellow (FJC2019-042304-I) from the Spanish Ministry of Science and Innovation (MCIN).

T.B. was supported by the German Research Foundation (SFB1383 Project ID 403224013/B07).

The Genome Analysis Platform in CIC bioGUNE is supported by the Basque Department of Industry, Tourism and Trade (Etortek, Elkartek and Emaitek Programs), the Innovation Technology Department of Bizkaia County, CIBERehd Network, and Spanish MINECO the Severo Ochoa Excellence Accreditation (CEX2021-001136-S).

JC laboratory is supported by Ministerio de Ciencia e Innovación/Agencia Estatal Investigación MCIN/AEI (PID2022-138970OB-I00 10.13039/501100011033/FEDER, UE). C.T. was supported by the DFG grants TR285/10-2 and SFB CRC1382, A08 (ID 403224013).

PER research laboratory is supported by the Fondation pour la Recherche Médicale (FRM EQU202303016287), "Institut National de la Santé et de la Recherche Médicale" (ATIP AVENIR), the "Agence Nationale pour la Recherche" (ANR-18-CE14-0006-01, RHU QUID-NASH, ANR-18-IDEX-0001,

ANR-22-CE14-0002) by « Émergence, Ville de Paris », by Fondation ARC and by France 2030 RHU LIVER-
TRACK (ANR-23-RHUS-0014).

D.G.-C. research laboratory is supported by King Abdullah University of Science and Technology.

Authors´ contributions: THW, RK, JC, PER, CT, and DGC designed the study. THW, MRP, MB, GZ, DB,
MS, WG, RS, SV, AV, DC, and IG supervised patient recruitment and sample collection and assessed
clinical data. THW, MRP, FS, AMA and JR performed experiments. SPE led and performed data
assessment, analysis, and visualization. THW, DGC, NP, CLV and EH supported data assessment,
analysis, and visualization. THW, SPE, MRP, DGC, NP, and EH performed data interpretation. THW,
SPE, NP, and DGC wrote the manuscript. EH, MRP, FS, JR, AK, RK, TB, PC, CA, NK, JT, JC, PER, and CT
revised the manuscript. All authors provided intellectual input and approved the final version of the
manuscript.

References

1. Ginès P, Krag A, Abraldes JG, Solà E, Fabrellas N, Kamath PS. Liver cirrhosis. The Lancet. 2021 Oct;398(10308):1359–76.

2. Mansour D, McPherson S. Management of decompensated cirrhosis. Clin Med (Lond). 2018 Apr 1;18(Suppl 2):s60–5.

3. Engelmann C, Clària J, Szabo G, Bosch J, Bernardi M. Pathophysiology of decompensated cirrhosis: Portal hypertension, circulatory dysfunction, inflammation, metabolism and mitochondrial dysfunction. Vol. 75, Journal of Hepatology. Elsevier B.V.; 2021. p. S49–66.

4. D'Amico G, Morabito A, D'Amico M, Pasta L, Malizia G, Rebora P, et al. Clinical states of cirrhosis and competing risks. J Hepatol. 2018 Mar;68(3):563–76.

5. Moreau R, Jalan R, Gines P, Pavesi M, Angeli P, Cordoba J, et al. Acute-on-chronic liver failure is a distinct syndrome that develops in patients with acute decompensation of cirrhosis. Gastroenterology. 2013 Jun;144(7):1426–37, 1437.e1-9.

6. Gustot T, Stadlbauer V, Laleman W, Alessandria C, Thursz M. Transition to decompensation and acute-on-chronic liver failure: Role of predisposing factors and precipitating events. J Hepatol. 2021 Jul;75 Suppl 1:S36–48.

7. Trebicka J, Fernandez J, Papp M, Caraceni P, Laleman W, Gambino C, et al. The PREDICT study uncovers three clinical courses of acutely decompensated cirrhosis that have distinct pathophysiology. J Hepatol. 2020 Oct 1;73(4):842–54.

8. Albillos A, Martin-Mateos R, Van der Merwe S, Wiest R, Jalan R, Álvarez-Mon M. Cirrhosis-associated immune dysfunction. Nat Rev Gastroenterol Hepatol. 2022 Feb;19(2):112–34.

9. Arroyo V, Angeli P, Moreau R, Jalan R, Clària J, Trebicka J, et al. The systemic inflammation hypothesis: Towards a new paradigm of acute decompensation and multiorgan failure in cirrhosis. J Hepatol. 2021 Mar;74(3):670–85.

10. Clària J, Arroyo V, Moreau R. Roles of systemic inflammatory and metabolic responses in the pathophysiology of acute-on-chronic liver failure. JHEP Reports. 2023 Sep;5(9):100807.

11. Clària J, Stauber RE, Coenraad MJ, Moreau R, Jalan R, Pavesi M, et al. Systemic inflammation in decompensated cirrhosis: Characterization and role in acute-on-chronic liver failure. Hepatology. 2016 Oct;64(4):1249–64.

12. Engelmann C, Sheikh M, Sharma S, Kondo T, Loeffler-Wirth H, Zheng YB, et al. Toll-like receptor 4 is a therapeutic target for prevention and treatment of liver failure. J Hepatol. 2020 Jul;73(1):102–12.

13. Clària J, Arroyo V, Moreau R. The Acute-on-Chronic Liver Failure Syndrome, or When the Innate Immune System Goes Astray. J Immunol. 2016 Nov 15;197(10):3755–61.

14. Trebicka J, Bork P, Krag A, Arumugam M. Utilizing the gut microbiome in decompensated cirrhosis and acute-on-chronic liver failure. Nat Rev Gastroenterol Hepatol. 2021 Mar;18(3):167–80.

15. Wasmuth HE, Kunz D, Yagmur E, Timmer-Stranghöner A, Vidacek D, Siewert E, et al. Patients with acute on chronic liver failure display "sepsis-like" immune paralysis. J Hepatol. 2005 Feb;42(2):195–201.

16. Martin-Mateos R, Alvarez-Mon M, Albillos A. Dysfunctional Immune Response in Acute-on-Chronic Liver Failure: It Takes Two to Tango. Front Immunol. 2019;10:973.

17. Weiss E, de la Grange P, Defaye M, Lozano JJ, Aguilar F, Hegde P, et al. Characterization of Blood Immune Cells in Patients With Decompensated Cirrhosis Including ACLF. Front Immunol. 2021 Feb 5;11.

18. Yao J, Liu T, Zhao Q, Ji Y, Bai J, Wang H, et al. Genetic landscape and immune mechanism of monocytes associated with the progression of acute-on-chronic liver failure. Hepatol Int. 2023 Jun 10;17(3):676–88.

688   19.   Feio-Azevedo R, Boesch M, Radenkovic S, van Melkebeke L, Smets L, Wallays M, et al. Distinct
689   immunometabolic signatures in circulating immune cells define disease outcome in acute-on-
690   chronic liver failure. Hepatology. 2024 May 16;

691   20.   Yao J, Ji Y, Liu T, Bai J, Wang H, Yao R, et al. Single-Cell RNA Sequencing Shows T-Cell Exhaustion
692   Landscape in the Peripheral Blood of Patients with Hepatitis B Virus-Associated Acute-on-
693   Chronic Liver Failure. Gut Liver. 2024 May 15;18(3):520–30.

694   21.   Zhang P, Li H, Zhou C, Liu K, Peng B, She X, et al. Single-Cell RNA Transcriptomics Reveals the
695   State of Hepatic Lymphatic Endothelial Cells in Hepatitis B Virus-Related Acute-on-Chronic Liver
696   Failure. J Clin Med. 2022 May 20;11(10):2910.

697   22.   Ruitenberg JJ, Mulder CB, Maino VC, Landay AL, Ghanekar SA. VACUTAINER CPT and Ficoll
698   density gradient separation perform equivalently in maintaining the quality and function of
699   PBMC from HIV seropositive blood samples. BMC Immunol. 2006 May 25;7:11.

700   23.   Zheng GXY, Terry JM, Belgrader P, Ryvkin P, Bent ZW, Wilson R, et al. Massively parallel digital
701   transcriptional profiling of single cells. Nat Commun. 2017 Jan 16;8(1):14049.

702   24.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal
703   RNA-seq aligner. Bioinformatics. 2013 Jan 1;29(1):15–21.

704   25.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, et al. Integrated analysis of
705   multimodal single-cell data. Cell. 2021 Jun;184(13):3573-3587.e29.

706   26.   Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq data
707   using regularized negative binomial regression. Genome Biol. 2019 Dec 23;20(1):296.

708   27.   Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell transcriptomic data
709   across different conditions, technologies, and species. Nat Biotechnol. 2018 May 2;36(5):411–
710   20.

711   28.   Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K, et al. Fast, sensitive and accurate
712   integration of single-cell data with Harmony. Nat Methods. 2019 Dec 18;16(12):1289–96.

713   29.   Mulè MP, Martins AJ, Tsang JS. Normalizing and denoising protein expression data from
714   droplet-based single cell profiling. Nat Commun. 2022 Apr 19;13(1):2099.

715   30.   Miller SA, Policastro RA, Sriramkumar S, Lai T, Huntington TD, Ladaika CA, et al. LSD1 and
716   Aberrant DNA Methylation Mediate Persistence of Enteroendocrine Progenitors That Support
717   *BRAF* -Mutant Colorectal Cancer. Cancer Res. 2021 Jul 15;81(14):3791–805.

718   31.   Wu T, Hu E, Xu S, Chen M, Guo P, Dai Z, et al. clusterProfiler 4.0: A universal enrichment tool
719   for interpreting omics data. The Innovation. 2021 Aug;2(3):100141.

720   32.   Qiu X, Mao Q, Tang Y, Wang L, Chawla R, Pliner HA, et al. Reversed graph embedding resolves
721   complex single-cell trajectories. Nat Methods. 2017 Oct 21;14(10):979–82.

722   33.   Andrews S. http://www.bioinformatics.babraham.ac.uk/projects/fastqc. 2010. FastQC: a
723   quality control tool for high throughput sequence data.

724   34.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data.
725   Bioinformatics. 2014 Aug 1;30(15):2114–20.

726   35.   Anders S, Pyl PT, Huber W. HTSeq—a Python framework to work with high-throughput
727   sequencing data. Bioinformatics. 2015 Jan 15;31(2):166–9.

728   36.   Law CW, Chen Y, Shi W, Smyth GK. voom: precision weights unlock linear model analysis tools
729   for RNA-seq read counts. Genome Biol. 2014;15(2):R29.

730   37.   Hänzelmann S, Castelo R, Guinney J. GSVA: gene set variation analysis for microarray and RNA-
731   Seq data. BMC Bioinformatics. 2013 Dec 16;14(1):7.

732   38.   Farias AQ, Curto Vilalta A, Momoyo Zitelli P, Pereira G, Goncalves LL, Torre A, et al. Genetic
733   Ancestry, Race, and Severity of Acutely Decompensated Cirrhosis in Latin America.
734   Gastroenterology. 2023 Sep;165(3):696–716.

735   39.   Robin X, Turck N, Hainard A, Tiberti N, Lisacek F, Sanchez JC, et al. pROC: an open-source
736   package for R and S+ to analyze and compare ROC curves. BMC Bioinformatics. 2011 Dec
737   17;12(1):77.

40. Hajian-Tilaki K. Receiver Operating Characteristic (ROC) Curve Analysis for Medical Diagnostic
Test Evaluation. Caspian J Intern Med. 2013;4(2):627–35.

41. R Core Team. R: A language and environment for statistical computing. Vienna, Austria: R
Foundation for Statistical Computing; 2021.

42. Bernsmeier C, Triantafyllou E, Brenig R, Lebosse FJ, Singanayagam A, Patel VC, et al. CD14+
CD15- HLA-DR- myeloid-derived suppressor cells impair antimicrobial responses in patients
with acute-on-chronic liver failure. Gut. 2018 Jun;67(6):1155–67.

43. Bernsmeier C, Pop OT, Singanayagam A, Triantafyllou E, Patel VC, Weston CJ, et al. Patients
with acute-on-chronic liver failure have increased numbers of regulatory immune cells
expressing the receptor tyrosine kinase MERTK. Gastroenterology. 2015 Mar;148(3):603-
615.e14.

44. Yao RQ, Zhao PY, Li ZX, Liu YY, Zheng LY, Duan Y, et al. Single-cell transcriptome profiling of
sepsis identifies HLA-DRlowS100Ahigh monocytes with immunosuppressive function. Mil Med
Res. 2023 Jun 19;10(1):27.

45. Li J, Liang X, Jiang J, Yang L, Xin J, Shi D, et al. PBMC transcriptomics identifies immune-
metabolism disorder during the development of HBV-ACLF. Gut. 2022 Jan;71(1):163–75.

46. Maheshwari D, Kumar D, Jagdish RK, Nautiyal N, Hidam A, Kumari R, et al. Bioenergetic Failure
Drives Functional Exhaustion of Monocytes in Acute-on-Chronic Liver Failure. Front Immunol.
2022 Jun 3;13.

47. Langer MM, Eickelmann C, Kleinbongard P, Lange CM. Impaired mitochondrial complex IV
respiration in peripheral blood mononuclear cells discriminates acute-on-chronic liver failure
from acute decompensation. J Hepatol. 2022 Sep;77(3):874–6.

48. Zhang IW, Curto A, López-Vicario C, Casulleras M, Duran-Güell M, Flores-Costa R, et al.
Mitochondrial dysfunction governs immunometabolism in leukocytes of patients with acute-
on-chronic liver failure. J Hepatol. 2022 Jan;76(1):93–106.

49. Wasmuth HE, Kunz D, Yagmur E, Timmer-Stranghöner A, Vidacek D, Siewert E, et al. Patients
with acute on chronic liver failure display "sepsis-like" immune paralysis. J Hepatol. 2005
Feb;42(2):195–201.

50. Bernsmeier C, van der Merwe S, Périanin A. Innate immune cells in cirrhosis. J Hepatol. 2020
Jul;73(1):186–201.

51. Bajaj JS, Salzman NH, Acharya C, Sterling RK, White MB, Gavis EA, et al. Fecal Microbial
Transplant Capsules Are Safe in Hepatic Encephalopathy: A Phase 1, Randomized, Placebo-
Controlled Trial. Hepatology. 2019 Nov;70(5):1690–703.

52. Trebicka J, Macnaughtan J, Schnabl B, Shawcross DL, Bajaj JS. The microbiota in cirrhosis and
its role in hepatic decompensation. J Hepatol. 2021 Jul;75 Suppl 1(Suppl 1):S67–81.

Figures

777
778
779
Figure 1. Study design.
(A) Schematic diagram of the study design. PBMC samples from 63 patients admitted with AD and 15 HC were
collected. After 90 days of follow-up (d90), AD patients were classified as SDC, UDC, or pre-ACLF due to their
clinical course. Of those, 16 patients and 4 HC were selected for scRNA-seq analysis. **(B)** Representation of the
multi-omics approach along with the associated clinical metadata, including study group, CLIF-C AD score,
gender, and age. *Abbreviations:* AD, acute decompensation; HC, healthy controls; PBMCs, peripheral blood
mononuclear cells; pre-ACLF, pre-acute-on-chronic liver failure; SDC, stable decompensated cirrhosis; UDC,
unstable decompensated cirrhosis.

Figure 2. Single-cell transcriptional profiling of PBMCs from healthy controls and patients with SDC, UDC, and pre-ACLF.

(A) UMAP projection of the 143,952 PBMCs analyzed, colored by the 11 detected cell populations. **(B)** Barplot showing the variability of immune cell subsets between the different groups. **(C)** Cleveland plot showing the log2 fold-change for the eight main cellular subpopulations. For each cell type, pairwise comparisons were made: SDC *vs.* HC (yellow), UDC *vs.* HC (orange), and pre-ACLF *vs.* HC (dark red). Cell types were considered differentially expressed when the fold change (log2 FC; estimated using a bootstrapping confidence interval analysis) was > 0.58 (dashed vertical lines) and the adjusted p-value was < 0.05 (represented by the dot size; estimated using a permutation analysis). **(D)** Sum of the number of DEGs across cell types; a gene is considered significant if adjusted p-value < 0.05 (Benjamini-Hochberg) and absolute Fold Change > 1.5. **(E, F)** Volcano plots representing the fold change (log2 FC) and statistical significance (adjusted p-value) for different immune cell populations: B cells, classical monocytes, non-classical monocytes, CD4+ T cells, CD8+ T cells, NK cells, MAIT cells, and Tregs. The x-axis shows the log2 FC, while the y-axis represents the -log10 of the adjusted p-values for each gene. Points in red and blue indicate statistically significant changes (based on adjusted p-values) with positive or negative log2FC, respectively, while gray points indicate non-significant changes. **E,** illustrates the comparison of pre-ACLF against SDC, and **F,** illustrates the comparison of pre-ACLF against UDC. *Abbreviations:* AD, acute decompensation; HC, healthy controls; PBMC, peripheral blood mononuclear cells; pre-ACLF, pre-acute-on-chronic liver failure; NK cells, natural killer cells; MAIT cells, mucosal-associated invariant T cells; SDC, stable decompensated cirrhosis; Tregs, regulatory T cells; UDC, unstable decompensated cirrhosis.

809
810

811
Figure 3. Identification of a classical monocyte subpopulation associated with ACLF development.
(A) UMAP of all cell types with a zoom on classical monocytes with cells colored based on the subclusters C0,
C1, and C2. **(B)** Relative percentages of each monocyte subcluster in each individual. **(C)** Cleveland plot showing
the log2 fold-change in cell subtype prevalence for the 3 monocyte subclusters. For each subcluster, pairwise
comparisons were made: SDC vs. HC, UDC vs. HC, and pre-ACLF vs. HC. Log2FC and FDR were estimated,
respectively, using bootstrapping and permutation analysis. **(D)** UMAP with the pseudo-time trajectory on
classical monocytes subclusters using Monocle3. **(E)** Dot plot indicating the expression of selected genes in each
monocytic subcluster. **(F)** Heatmap displaying the scaled expression values of the top 50 up and downregulated
genes in each subcluster. Please find extensive gene annotation in Supplementary Figure 10. **(G)** Violin plots
illustrating the distribution, median, and IQR of the ΔMIF (subtraction of the mean MFI count in the SDC group
from single MFI counts in the four study groups) of surface HLA-DR expression on classical monocytes. Samples
were drawn from the single-cell study cohort, including healthy controls (n=4) and patients with SDC (n=6), UDC
(n=6), and pre-ACLF (n=4). The Wilcoxon-Mann-Whitney test was used to test statistically significant differences
in relative HLA-DR expression within the four study groups. *Abbreviations:* Avg, average; HC, healthy controls;
Log2FC, logarithmic 2 Fold Change; pre-ACLF, pre-acute-on-chronic liver failure; SDC, stable decompensated
cirrhosis; UDC, unstable decompensated cirrhosis; ΔMIF, delta of mean fluorescence intensity.

829
Figure 4. GSEA of key biological processes defining monocyte subpopulations.
(A) GSEA results for the three biological processes: oxidative phosphorylation, aerobic respiration, and ATP
biosynthetic process. Running Enrichment Score (green line) depicts the cumulative enrichment score as genes
are ranked by expression level. Ranked List Metric (bottom grey plot) shows the correlation of genes with the
phenotype of interest across the ranked dataset. The heatmap (middle colored bar) represents the distribution
of gene ranks—with red for positive correlation, blue for negative correlation, and white for a midpoint. **(B)**
Boxplots showing the mean pathway signature score for the three processes for each monocyte subset and **(C)**
for each group. Differences in module scores were assessed using Kruskal Wallis Test.***adjusted *p*-value
<0.001; n.s. not significant**. (D)** Average gene expression trends across monocytic subsets related to Positive
regulation of inflammatory response, Defense response to bacterium, and Cytokine production involved in
immune response. Asterisks indicate reaching a significant nominal *p*-value, whereas the adjusted *p*-value was
not significant. Differences in module scores were assessed using Kruskal Wallis Test. *nominal *p*-value <0.05.
Abbreviations: n.s. not significant.
843
844

Figure 5. Identification of the C2 signature in whole blood RNA-seq data.
(A) Venn diagram illustrates the overlap of DEGs at single-cell resolution among three comparisons: classical monocytes versus all other cell types (blue), C2 versus Clusters 0 and 1 (yellow), and C2 pre-ACLF versus C2 in all groups (HC, SDC, and UDC (red). **(B)** Heatmap showing the scaled log2 FC levels of the selected DEGs from different comparatives in bulk RNA-seq: HC/SDC/ UDC vs. pre-ACLF. The color scale represents the log2 fold-change, with blue indicating downregulation and red indicating upregulation. Significant DEGs with log2 FC > 0.32 and adjusted p-value < 0.05 are marked with an asterisk; genes with a tendency (log2 FC > 0.32) are marked with a cross. **(C)** UMAP shows the distribution of the expression of the C2 signature. The inset panels show the distribution of C2 signature across different groups. **(D)** Boxplot displaying the module score for the C2 signature across the paired bulk RNA-seq data, categorized by group.

Figure 6. C2 monocytic subcluster validation in two independent AD cohorts.

(A) C2 signature was assessed in whole blood RNA-seq data collected at day 0 (d0) from two independent cohorts
of AD patients. PREDICT cohort (top) accounted for data from 689 patients who were labeled 90 days after HA
as SDC (n=449), UDC (n=179), or pre-ACLF (n=61). ACLARA (bottom) accounted for data from 521 patients who
were labeled 28 days after HA as AD (n=456) or as pre-ACLF (n=65). **(B)** Correlation analysis of C2 signature values
with CLIF-C AD and MELD across different patient groups in PREDICT (left) and ACLARA (right) cohorts. Top row:
C2 signature vs. CLIF-C AD count. Bottom row: C2 signature vs. MELD levels. Each point represents an individual,
colored by patient group: SDC, yellow; UDC, orange; pre-ACLF, dark red for the PREDICT cohort, and AD, yellow
and pre-ACLF, dark red for the ACLARA cohort. **(C)** Boxplots display the average expression of the C2 signature
among the different groups in PREDICT (left) and ACLARA (right) cohorts. Wilcoxon Signed-Rank Test was used
to assess the differences between groups. **(D)** Boxplots display the average expression of the C2 signature
between patients that died (pink) vs. patients that did not die (turquoise) in PREDICT (left) and ACLARA (right)
cohorts Wilcoxon Signed-Rank Test was used to assess the differences between groups**. (E)** ROC curves showing
the predictive capacity of the C2 signature regarding outcomes ACLF (top) and mortality (bottom) for the
PREDICT (red) and the ACLARA (blue) cohorts. *** Adjusted p-value < 0.001; ** Adjusted p-value < 0.01; *
Adjusted p-value < 0.05. *Abbreviations:* AD, acute decompensation; AUC, area under the curve; d0, day 0; HC,
healthy controls; PBMC, peripheral blood mononuclear cells; pre-ACLF, pre-acute-on-chronic liver failure; ROC,
Receiver-operating characteristic curve; SDC, stable decompensated cirrhosis; UDC, unstable decompensated
cirrhosis.

Table 1. Baseline characteristics of single-cell study cohort.

	HC (n=4)	SDC (n=6)	UDC (n=6)	pre-ACLF (n=4)	*p*-value
Characteristics					
Age, median (range)	56 (52 - 64)	60 (54 - 81)	57 (51- 60)	74 (60 - 79)	0.063
Female Sex, yes, n (%)	2 (50)	2 (33)	0	2 (50)	0.248
Previous decompensations in the last 2 years, yes, n (%)	-	0	4 (67)	3 (75)	**0.023**
Cause of cirrhosis, n (%)					
Alcohol-related	-	1 (17)	3 (50)	2 (50)	0.606
Viral (HBV/HCV)	-	1 (17)	1 (17)	0	0.259
Alcohol-related + MASLD	-	1 (17)	1 (17)	0	0.259
MASLD	-	2 (50)	1 (17)	1 (25)	0.801
cryptogenic	-	0	0	1 (25)	0.202
other	-	1 (17)	0	0	0.411
Data at admission, n (%)					
Ascites	-	5 (83)	5 (83)	4 (100)	0.683
SBP	-	0	0	2 (50)	**0.032**
Other bacterial infection	-	2 (33)	0	0	0.149
Variceal bleeding	-	0	1 (17)	0	0.411
HE	-	1 (17)	2 (33)	1 (25)	0.801
Alcoholic hepatitis		1 (17)	0	1 (25)	0.467
Acute kidney injury	-	0	1 (17)	0	0.411
Laboratory parameters					
Bilirubin, total [mg/dl], median, (range)	-	2.31 (1.3-3.4)	2.14 (1.37-4.60)	3.91 (2.61-5.85)	0.217
AST [U/l], median, (range)		44 (28-64)	61 (35-361)	54 (34-125)	0.294
ALT [U/l], median, (range)		33 (11-38)	32 (17-76)	36 (18-57)	0.709
INR, median, (range)		1.38 (1.20-1.75)	1.46 (1.31-1.77)	1.69 (1.26-2.95)	0.516
Creatinine [mg/dl], median, (range)		0.72 (0.54-0.92)	0.80 (0.60-1.97)	1.36 (1.17-1.62)	**0.012**
Platelets [x10^3/L], median (range)		79 (39-176)	75 (41-130)	81 (37-106)	0.972
WBC [x10^9/L], median (range)	4.8 (4.0-6.3)	4.1 (2.1-5.7)	5.4 (2.9-7.5)	5.5 (3.4-9.4)	0.476
CRP [mg/L], median (range)	-	16.25 (2.22-76)	13.7 (1.78-45.2)	8.55 (1.23-37.10)	0.567
Severity scores					
Child-Pugh, median, range	-	7.5 (5 - 12)	10 (7 - 11)	11 (8 - 13)	0.126
MELD, median, range	-	12 (8 - 18)	13 (12 - 23)	20 (17 - 27)	**0.033**
CLIF-C AD, median, range	-	47 (41 - 49)	47.5 (41 - 59)	61 (55 - 74)	**0.025**
Treatment at blood draw					
Diuretics, yes, n (%)	-	3 (50)	4 (67)	3 (75)	0.701
Antibiotics, yes, n (%)	-	1 (17)	1 (17)	1 (25)	0.934
Non-selective ß-blockers, yes, n (%)	-	1 (17)	2 (33)	3 (75)	0.169
Albumin, yes, n (%)	-	1 (17)	0	1 (25)	0.467
HE treatment, yes, n (%)	-	1 (17)	3 (50)	3 (75)	0.176
Follow-up and Outcome					

90-day mortality, yes, n (%)	-	0	0	1 (25)	0.229
90-day OLT, yes, n (%)	-	0	0	1 (25)	0.229

HE treatment includes lactulose and rifaximin. Baseline characteristics are depicted as frequencies or median (IQR). Values of *p*-value are based on Wilcoxon–Mann–Whitney test for continuous and chi-square test for categorical variables. *Abbreviations:* SDC, stable decompensated cirrhosis; UDC, unstable decompensated cirrhosis; pre-ACLF, pre-acute-on-chronic liver failure; HBV: Hepatitis B virus; HCV: Hepatitis C virus; MASLD: metabolic-dysfunction associated steatotic liver disease; SBP: spontaneous bacterial peritonitis; HE: hepatic encephalopathy; AST, aspartate aminotransferase; ALT, alanine aminotransferase; WBC: white blood cell count; CRP: C-reactive protein; OLT: orthotopic liver transplantation.