Exposure to trauma in pregnant women and its association with previous perinatal complications,

IPV and antenatal service satisfaction in rural Ethiopia: a cross-sectional facility-based study

Raquel Catalao[1]

Lelina Kebede[2]

Adiyam Mulushoa[3]

Tigist Eshetu[3]

Girmay Medhin[4]

Ahmed Abdella[5]

Atalay Alem[6]

Roxanne C. Keynejad[7, 8]

Jane Sandall[9]

Louise M Howard[7]

Martin Prince[10]

Charlotte Hanlon[11, 3, 6*]

*Corresponding author: Prof Charlotte Hanlon (chanlon@ed.ac.uk)

Affiliations

[1] Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King´s

College London, United Kingdom

[2] St Barnabas Hospital, Bronx, New York, USA

[3] Centre for Innovative Drug Development and Therapeutic Trials for Africa (CDT Africa), College of

Health Sciences, Addis Ababa University, Addis Ababa, Ethiopia

[4] Aklilu-Lema Institute of Pathobiology, Addis Ababa University, Addis Ababa, Ethiopia

[5] Department of Obstetrics and Gynaecology, School of Medicine, College of Health Sciences, Addis

Ababa University, Addis Ababa, Ethiopia

[6] Department of Psychiatry, WHO Collaborating Centre for Mental Health Research and Capacity

Building, School of Medicine, College of Health Sciences, Addis Ababa University, Addis Ababa,

Ethiopia

[7] Section of Women's Mental Health, Department of Health Service and Population Research,

Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United

Kingdom

[8] Department of Women and Children's Health, School of Life and Population Sciences, King's

College, London, United Kingdom

[9] Centre for Global Mental Health, Health Service and Population Research Department, Institute of

Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom

[10] King's Global Health Institute, King's College London, London, United Kingdom

[11] Division of Psychiatry, Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh,

Scotland, United Kingdom

*Corresponding author: Prof Charlotte Hanlon (chanlon@ed.ac.uk)

Affiliations

[1] Department of Psychological Medicine, Institute of Psychiatry, Psychology and Neuroscience, King's

College London, United Kingdom

[2] St Barnabas Hospital, Bronx, New York, USA

[3] Centre for Innovative Drug Development and Therapeutic Trials for Africa (CDT Africa), College of

Health Sciences, Addis Ababa University, Addis Ababa, Ethiopia

[4] Department of Obstetrics and Gynaecology, School of Medicine, College of Health Sciences, Addis

Ababa University, Addis Ababa, Ethiopia

[5] Department of Psychiatry, WHO Collaborating Centre for Mental Health Research and Capacity

Building, School of Medicine, College of Health Sciences, Addis Ababa University, Addis Ababa,

Ethiopia

[6] Section of Women's Mental Health, Department of Health Service and Population Research,

Institute of Psychiatry, Psychology and Neuroscience, King's College London, London, United

Kingdom

[7] Department of Women and Children's Health, School of Life and Population Sciences, King's

College, London, United Kingdom

[8] Centre for Global Mental Health, Health Service and Population Research Department, Institute of

Psychiatry, Psychology and Neuroscience, King's College London, London, United Kingdom

[9] King's Global Health Institute, King's College London, London, United Kingdom

[10] Division of Psychiatry, Centre for Clinical Brain Sciences, University of Edinburgh, Edinburgh,

Scotland, United Kingdom

Abstract

Background

We aimed to describe the prevalence of exposure to traumatic events and post-traumatic stress

disorder (PTSD) in pregnant women attending antenatal care (ANC) in rural Ethiopia. We

hypothesised that antenatal PTSD symptoms would be associated with previous obstetric

complications and intimate partner violence (IPV) and impact negatively on women's satisfaction

with ANC.

Methods

The design was a facility-based cross-sectional study in primary health centres providing ANC in

southern Ethiopia. Trauma events were assessed using the Life Events Checklist (LEC) and PTSD

checklist for DSM-5 (PCL-5). Previous obstetric complications were extracted from clinical records.

IPV was measured using the 'Non-Graphic Language' screening test and ANC satisfaction was

measured using a locally validated adapted version of the Mental Health Service Satisfaction Scale.

Generalized linear mixed-effects regression models were used to calculate prevalence ratios

between PTSD, IPV and ANC satisfaction.

Results

Out of 2079 interviewed women, 52.3% (n=1,087) reported one or more traumatic life events on the

LEC. Physical assault was the most common traumatic event experienced (n=485; 23.3%) and

witnessed (n=1,176; 56.6%) but only 289 (13.9%) screened positive for IPV. One hundred and six

women (5.1%) met DSM-5 criteria for PTSD. Women meeting diagnostic criteria for PTSD had five

times increased prevalence of IPV in their current pregnancy [prevalence ratio (PR) 4.34 95%CI 3.01-

6.30; p<0.001]. Only twenty-six women had a record of previous obstetric complications (0.01%).

Overall, women with PTSD reported less satisfaction with antenatal care.

Conclusions

Despite high exposure to traumatic life events, particularly physical violence, among pregnant

women attending ANC in Southern Ethiopia, the prevalence of PTSD is relatively low. Previous

obstetric complications and IPV were under-reported, relative to known prevalence estimates. Our

study highlights the challenges of detection of psychosocial needs in the ANC setting and the need

for targeted interventions to support women's disclosure of difficulties in maternity care settings.

Introduction

Despite the high prevalence and adverse impacts of mental health conditions (MHCs) during the

perinatal period (from conception up to two years post-partum), maternal health programmes

continue to neglect women's mental health needs (1). In low- and middle-income countries (LMICs),

it is estimated 1 in 4 women experience depression in the perinatal period (2). The prevalence of

perinatal depressive and anxiety symptoms is higher among the most marginalized women, with

least access to health and social care (3). Beyond adverse birth outcomes, perinatal MHCs have

transgenerational effects, negatively impacting children's health and development (4). Inadequate

recognition and low availability of treatment of perinatal MHCs in LMICs results in women remaining

undiagnosed and untreated, impacting their quality of life and the physical and emotional health of

their children and families (5).

Most research on perinatal MHCs in Africa has focused on common mental disorders (CMDs),

namely depressive and anxiety disorders (6). Little is known about the prevalence of perinatal post-

traumatic stress disorder (PTSD) in LMICs and its consequences for women's health and future

pregnancies. A systematic review of studies mostly conducted in high income countries (HICs),

reported PTSD prevalence of 3.3% antenatally and 4.0% postpartum (7). However, the prevalence of

PTSD was higher in women with a history of childhood maltreatment: 18.9% antenatally and 18.5%

postpartum (7). Antenatal PTSD has been reported following traumatic life events such as accidents,

interpersonal violence and natural disasters, while birth trauma is a common precipitant for

postpartum PTSD (8). Disrespectful and abusive care during childbirth is widespread in Ethiopia (9),

as in other countries. In HICs, there is growing awareness of the needs of women exposed to trauma,

the potential for re-traumatisation in maternity care and the requirement for therapeutic

relationships and trauma-informed care (10). Obstetric complications and IPV are commoner among

women in LMICs (11) but few studies have investigated the associations between trauma exposure,

PTSD, and ANC satisfaction among pregnant women in low resource settings. There is growing

evidence from both HICs and LMICs that person-centered maternity care, which is respectful and

responsive, supports women´s autonomy and engages the women in their care leads to improved

outcomes (12).

In this study we aimed to describe the prevalence of exposure to traumatic events and PTSD

symptoms in pregnant women attending ANC in rural Ethiopia and their associations with previous

obstetric complications, IPV and ANC satisfaction. We hypothesised that antenatal PTSD symptoms

would be associated with previous obstetric complications and current IPV and would impact

negatively on women´s satisfaction with antenatal care received.

Materials and Methods

Study Design

We conducted secondary analysis of data collected in a facility-based cross-sectional study that

formed part of a larger study (ASSET) of health system strengthening in sub-Saharan Africa (13).

ASSET was a multi-country consortium in Ethiopia, Sierra Leone, South Africa, and Zimbabwe which

aimed to improve access to high quality care across three health care platforms: integrated

continuing care for non-communicable diseases including mental health conditions (NCD/MH),

maternal and newborn care, and surgical care. The data for this analysis was part of the maternal

and newborn care platform in Ethiopia (14)

Study Setting

The study was conducted in eight purposively selected health centres in Meskan and Sodo districts,

in the Gurage Zone of the Southern Nations, Nationalities and People's Region (SNNPR) of Ethiopia,

recently renamed Central Ethiopia Regional State. According to census projections, in 2019 SNNPR

had a total population of 20.1 million people, of whom 52.7% were women aged 15-49 years. The

median age in the region is 20 years, 84.3% of the population live in rural areas and the total fertility

rate is 3.8 births per woman (15).

Study Population

A total of 2079 pregnant women attending health centers consecutively for ANC, who were able to

converse in Amharic and provide written informed consent, were included in this study. Acutely ill

women who required emergency medical attention were excluded. Data collection took place from

18th July 2019 to 9th January 2020. Women were reimbursed the equivalent of $2 for their time: a

locally appropriate fee.

Data Collection

Data collection was carried out in a private room within the health facility. All data collectors were

female and comprised both lay data collectors (with a minimum of high school education) and

clinical data collectors (with a minimum qualification of a diploma in nursing). Data were collected

162 with electronic tablets using Open Data Kit (ODK) software (16). Trained lay data collectors

163 administered fully structured questionnaires. Research nurses extracted information from medical

164 records for each woman using a bespoke template developed for the study(14).

166 Measures

168 Clinical Characteristics of pregnancy

169 Data on clinical characteristics of the current and any past pregnancies were collected from clinical

170 records using structured forms. Our exposure of interest was previous obstetric complications,

171 defined as one or more of the following during a previous pregnancy or delivery:

172 spontaneous/induced abortion, ectopic pregnancy/ ruptured ectopic pregnancy, hospital admission

173 for pre-eclampsia/ eclampsia, pre-term labour, intra-uterine foetal death, obstructed labour; fistula,

174 Instrumental delivery, delivery by Caesarean section, postpartum haemorrhage, previous baby with

175 a congenital anomaly, and previous baby born weighing less than 2500g or more than 4500g

177 Trauma exposure and symptoms

178 Exposure to traumatic events and trauma symptoms were assessed using the Life Events Checklist

179 (LEC) (17) and PTSD checklist for DSM-5 (PCL-5)(18), translated into Amharic and adapted for the

180 rural Ethiopia context(19). In the LEC, women were asked if they had experienced traumatic life

181 events, witnessed them or learned about them through social interactions or their occupation. The

182 PCL-5 checklist consists of 20 statements rated from "not at all" (score 0) to "extremely" (score 4).

183 Each item rated 2 ("moderately") or higher was considered endorsement of a symptom. PCL-5

184 divides PTSD symptoms, with questions 1-5 reflecting criterion B (intrusion) symptoms, questions 6-

7 reflecting criterion C (avoidance) symptoms, questions 8-14 reflecting criterion D (negative

alterations in cognitions and mood) symptoms, and questions 15-20 reflecting criterion E (alterations

in arousal and reactivity) symptoms. Women met DSM-5 criteria for PTSD if they scored 31 or above

on PCL-5 and endorsed at least one criterion B, C, D or E symptom.

Depressive and Anxiety Symptoms

Depressive symptoms were measured using the Patient Health Questionnaire (PHQ-9). In a previous

validation study of PHQ-9 (20) in health centres in the study area, a cut-off of 5 or more provided the

best discrimination between people with and without major depressive disorder but had low

positive predictive value (<25%). We categorised cases as 'probable depression' if the individual

scored 5 or more on the PHQ-9 and reported that it was very difficult or extremely difficult for them

to function in daily life. This accorded with international diagnostic criteria for depressive disorder

which require impaired functioning alongside symptoms. Anxiety symptoms were measured using

the generalised anxiety disorder scale (GAD-7)(21). Women were categorised as having low anxiety

symptoms if they scored 5-9, moderate anxiety symptoms if they scored 10-14 and severe anxiety

symptoms if they scored 15 or above.

Intimate Partner Violence (IPV)

IPV was measured using the 'non-graphic language' (NGL) IPV screening test (22) based on the

Conflict Tactics Scale-Revised (23). The NGL scale comprises 5 questions on experience of physical or

psychological violence with an intimate partner. It was previously found to be acceptable in the study

population, where it had convergent validity with the more extensive World Health Organization

(WHO) IPV questionnaire (24). We categorised participants scoring 2 or more on questions 1

(working out arguments), 3 (partner treatment) or 4 (feeling safe) as experiencing marital discord

and possible IPV.

Service satisfaction

Satisfaction with ANC was measured using the adapted version of the Mental Health Service

Satisfaction scale, validated for Ethiopian health centres and used in the ASSET diagnostic phase

(25). The modified MHSS scale had 21 items, each with four Likert response categories (1: strongly

disagree, 2: disagree, 3: agree and 4: strongly agree). Most items assessed generic aspects of

outpatient care relevant to the context, such as health worker communication, waiting times,

privacy, usefulness of care, cleanliness of the facility. For this study we analysed total satisfaction

scores and responses to all specific questions. We dichotomised results for individual items into

satisfied (answering 'agree' or 'strongly agree') and non-satisfied (answering 'disagree' or 'strongly

disagree').

Covariates

Sociodemographic characteristics including age, marital status, place of residence, educational level

and religion was gathered using structured measures in an interview format by the same lay data

collectors. Social support was assessed using the Oslo Social Support scale used previously in the

study setting and found to have convergent validity (26).

Ethical Considerations

We obtained ethical approval from the Addis Ababa University College of Health Sciences

Institutional Review Board (Reference number: 028/18/Psy) and King's College London Research

Ethics Committee (Reference number: HR-17/18-6570). The study was conducted according to the

Declaration of Helsinki. All participants provided written informed consent.

Analysis Plan

We used descriptive statistics to calculate socio-demographic, clinical characteristics and satisfaction

scores for the total sample and those participants meeting criteria for PTSD. We used Chi-squared

and Fisher exact tests to compare socio-demographic characteristics and previous obstetric

complications in those with and without PTSD. To account for the effect of the number of data

collection days in each health centre and average ANC attendance per health centre per day

(calculated over a 6-month period), we included these variables and used health centre as the

clustering variable in all the following regression analyses. We used Poisson mixed-effect regression

to calculate prevalence ratios of co-morbid depression and anxiety in participants meeting criteria

for PTSD. We used a univariate and multivariate Poisson mixed-effects regression model, adjusted

for *a priori* identified confounders: parity, education, place of residence and age, to calculate

prevalence ratios between PTSD and IPV. We performed a univariate and multivariate linear mixed-

effects regression test adjusted for the same confounders, to test the association between PTSD

detection and ANC satisfaction. We conducted a univariate and multivariate logistic mixed-effects

regression model for the association between PTSD and dichotomised satisfaction on each MHSS

item. We tested differences in responses to each MHSS item between those with and without PTSD

using Fisher exact tests.

Results

Socio-demographic characteristics

256 See Table 1. Out of 2079 participants, 1,024 (49.3%) were attending their first ANC appointment.

257 Participants' mean age was 26.0 years (standard deviation (SD) 4.9; range: 18 to 50 years). The

258 majority of women (n=2013; 96.8%) were married and had some formal education (n=1,468; 70.6%,

259 see Table 1).

262   **Table 1. Participant socio-demographic characteristics.**

Socio-demographic characteristics	Total sample n (%)	Probable PTSD[1] n=106 n (%)	Without probable PTSD n=1,973 n (%)	Statistical test for significance
Age (years)				
18-24	751 (36.1%)	26 (24.5%)	725 (36.7%)	Chi-squared test for trend X^2 (1, 2,079) = 3.75 p=0.0527
25-34	1144 (55.0%)	71 (67.0%)	1,073 (54.4%)	
35-50	184 (8.9%)	9 (8.5%)	175 (8.9%)	
Education				
No formal education	611 (29.4%)	25 (23.6%)	586 (29.7%)	Chi-squared test for trend X^2 (1, 2,079) = 0.05 p=0.8225
Primary education	1119 (53.8%)	69 (65.1%)	1,050 (53.2%)	
Secondary education	290 (14.0%)	11 (10.4%)	279 (14.1%)	
Post-secondary	59 (2.8 %)	1 (0.9%)	58 (2.9%)	
Residence				

Rural	1115 (53.6%)	52 (49.1%)	1,063 (53.9%)	Chi-squared test X² (1, 2,079) = 0.94, p=0.332
Marital Status				
Married	2013 (96.8%)	1,912 (96.9%)	101 (95.3%)	Fisher's exact test p=0.271
Single	37 (1.8%)	37 (1.8%)	2 (1.9%)	
Separated, divorced or widowed	29 (1.4%)	26 (1.3%)	3 (2.8%)	
Parity (n=2,076)				
0	595 (28.7%)	29 (27.4%)	566 (28.7%)	Chi-squared test for trend X² (1, 2,076) = 0.003 p=0.8684
1	485 (23.4%)	27 (25.5%)	458 (23.2%)	
2-4	777 (37.4%)	41 (38.7%)	736 (37.4%)	
5 or more	219 (10.6%)	9 (8.5%)	210 /10.7%)	
Pregnancy intention				
Wanted	1632 (78.5%)	76 (71.7%)	1,556 (78.9%)	Fisher's exact test p=0.094
Initially unwanted, now wanted	409 (19.7%)	4 (3.8%)	34 (1.7%)	

Unwanted	38 (1.8%)	26 (24.5%)	383 (19.4%)	
Social support				
Poor	533 (25.6%)	510 (25.9%)	23 (21.7%)	Chi-squared test for trend X^2 (1, 2,079) = 0.39 p=0.5300
Intermediate	1023 (49.2%)	958 (48.5%)	65 (61.3%)	
Strong	523 (25.2%)	505 (25.6%)	18 (17.0%)	

[1] Using DSM-5 criteria

Current and Previous Pregnancy Characteristics

Most participants were in their second trimester (54.3%, n=1,120) of a wanted pregnancy (78.5%,

n=1,632). Median parity was 1 (interquartile range: 0-3); for 28.7% of women (n=595) this was their

first pregnancy. A history of co-morbid physical health problems was documented in the clinical

records of four (n=4) participants.

A total of 26 (0.01%) women had previous obstetric complications documented in their clinical

records: recurrent miscarriage (n=15), stillbirth (n=12), previous Caesarean section (n=4),

hospitalisation for pre-eclampsia/eclampsia (n=2), fistula (n=1), and previous post-partum

haemorrhage (n=1). Ectopic pregnancy, assisted delivery, previous baby with a congenital anomaly,

and previous baby born weighing less than 2500g or more than 4500g were not documented for any

participants.

Trauma exposure and PTSD symptoms

A total of 1,087 (52.3%) women reported one or more traumatic life events on the LEC (range: 0 to

10; see Figure1). Twenty-seven percent of women (n=551) reported one traumatic event and 13.2%

(n=275) reported three or more traumatic life events. Physical assault was the most common

traumatic event experienced (n=485; 23.3%) and witnessed (n=1,176; 56.6%), whereas sexual

violence was infrequently reported (2.7% (n=57) reported sexual assault and 1.9%(n=40) reported

other unwanted sexual experiences; see Table 2). Direct exposure to armed combat was reported by

five (0.2%) women and witnessed by 90 (4.3%). Experience of a life-threatening injury or illness or

severe suffering were reported by 567 women (27.3%). One hundred and six women (5.1%) met

DSM-5 criteria for PTSD ('probable PTSD').

293 **Table 2. Types of Traumatic Life Events directly experienced and witnessed by women.**

Life event	Directly experienced (n, %)	Witnessed (n, %)
Physical assault	485 (23.3%)	1,176 (56.6%)
Life threatening injury/illness	396 (19.1%)	846 (40.7%)
Natural Disaster	221 (10.6%)	556 (26.7%)
Fire	182 (8.8%)	809 (38.9%)
Other very stressful event	175 (8.4%)	288 (13.8%)
Severe human suffering	171 (8.2%)	646 (31.1%)
Motor vehicle accident	99 (4.8%)	520 (25.0%)
Captivity	85 (4.1%)	391 (18.8%)
Sudden unexpected death of loved one	80 (3.8%)	661 (31.8%)
Other serious accident	64 (3.1%)	231 (11.1%)
Sexual assault	57 (2.7%)	110 (5.3%)
Assault with a weapon	51 (2.5%)	312 (15.0%)
Sudden violent death	50 (2.4%)	495 (23.8%)
Other unwanted sexual experience	40 (1.9%)	58 (2.8%)
Caused serious injury/ death of another	18 (0.9%)	155 (7.5%)

Exposure to toxic substance	11 (0.5%)	41 (2.0%)
Combat	5 (0.2%)	90 (4.3%)

There were no significant differences in education level, residence, marital status and parity

between women with and without probable PTSD (see Table 1). Women meeting diagnostic criteria

for PTSD were more likely to be aged 25 to 34 years old rather than younger and to report lower

levels of social support than women not meeting criteria for PTSD.

Forty-four percent (n=47) of those with probable PTSD also experienced moderate to severe anxiety

symptoms compared to 1.5% (n=29) of women without PTSD. Nearly a quarter of participants with

probable PTSD experienced co-morbid probable depression (n=25; 23.6%), compared to 4.3% (n=85)

of those without PTSD. Women with probable PTSD were significantly more likely to experience co-

morbid probable depression (prevalence ratio (PR): 5.50, 95% CI: 3.99-7.59, p<0.001) and severe

anxiety symptoms (PR: 20.80, 95% CI: 13.87-31.20, p<0.001) in unadjusted analyses.

Associations between previous obstetric complications and trauma symptoms

Very few participants' clinical records documented obstetric complications in a previous pregnancy

(n=24; 0.01 % in women not meeting criteria for PTSD and n=2 in women meeting criteria for PTSD);

the association was not statistically significant (p=0.502).

Association between trauma symptoms and IPV

A larger proportion of women with probable PTSD screened positive for probable IPV (41.5%; n=44)

compared to women without PTSD (n=245; 12.4%). Women meeting diagnostic criteria for PTSD had

four times greater risk of reporting IPV in their current pregnancy in both the unadjusted model (PR:

4.30, 95% CI: 2.25-8.21, p<0.001) and after adjustment for parity, age, education and area of

residence (PR 4.20 95%CI 2.05- 8.61; p<0.001). Of those screening positive for IPV in the total

sample (n=289, 13.9%), 133 (46.0%) reported directly experiencing physical assault and 19 (6.6%)

sexual assault.

Association between trauma symptom and service satisfaction

The mean total satisfaction with ANC score was 57.59 (SD=8.42) for women without probable PTSD

and 56.40 (SD=6.27) in women with probable PTSD. The mean difference was 1.19 (95% CI: -0.44 to -

2.82) which when accounting for clustering by health center was found to be significant in both

unadjusted analyses (regression coefficient: -0.73 (95% CI: -1.35 - -0.11, p=0.022) and after

adjustment for parity, age, education and area of residence [adjusted regression coefficient (adj RC):

-0.73 (95% CI: -1.36 - -0.10, p=0.022] .

Women meeting criteria for PTSD were more likely to report that ANC waiting times were

acceptable [adjusted Odds Ratio (adj OR) =2.35, 95% CI: 1.64-3.38, p<0.001)], that their privacy was

respected (adj OR =2.94, 95% CI: 1.91-4.52, p<0.001), that their information was kept confidential

(adj OR =2.14, 95% CI: 1.61-2.85, p <0.001) and that referral to other services was possible (adj OR

=3.64, 95% CI: 1.82-13.00, p=0.047) . Participants with probable PTSD were significantly less likely to

report that the health worker involved their family helpfully (adj OR =0.29, 95% CI: 0.12-0.67,

p<0.001) and that they can afford to attend the health facility for treatment (adj OR 0.60, 95% CI

0.37-0.99, o=0.047); see Table 3). There were significant differences in satisfaction responses

between women with and without probable PTSD for all ANC satisfaction items. Women fulfilling

criteria for PTSD were more likely to answer "agree" rather than "strongly agree" to most items,

despite similar overall satisfaction scores (table 4).

Table 3. Logistic mixed-effects regression models of associations between PTSD and ANC satisfaction.

		Probable PTSD (n=106)		Not probable PTSD (n=1,973)		Univariate logistic mixed-effects regression model (Odds ratio (OR) 95% confidence interval (CI)	Multivariate logistic mixed-effects regression model adjusted for parity, age, education and residence Adjusted OR (95%CI)
		Satisfied	Not Satisfied	Satisfied	Not satisfied		
1	The health worker treated me with courtesy	102 (96.2%)	4 (3.8%)	1,891 (95.8%)	82 (4.2%)	1.01 (0.26-4.81)	1.09 (0.24-4.98)
2	The health worker listened to me carefully	104 (98.1%)	2 (1.9%)	1,892 (95.9%)	81 (4.1%)	2.01 (0.62- 6.50)	2.10 (0.58 – 7.53)

3	The health worker explained things to me in a way I understood	93 (87.7%)	13 (12.3%)	1,693 (85.8%)	280 (14.2%)	1.10 (0.54- 2.28)	1.17 (0.57- 2.40)
4	The health facility was clean	98 (92.4%)	8 (7.6%)	1,882 (95.4%)	91 (4.6%)	0.63 (0.29-1.39)	0.57 (0.27-1.21)
5	The waiting room was clean	98 (92.4%)	8 (7.6%)	1,799 (91.2%)	174 (8.8%)	1.02 (0.76-1.37)	1.04 (0.78-1.39)
6	The latrine was clean	75 (70.8%)	31 (29.3%)	1,298 (65.8%)	675 (34.2%)	1.18 (0.83- 1.67)	1.21 (0.88-1.71)
7	The waiting time was acceptable	88 (83.0%)	18 17.0%)	1,312 (66.5%)	661 (33.5%)	2.35 (1.64-3.37)	2.35 (1.64-3.38)
8	I had enough time to discuss with health worker	98 (92.4%)	8 (7.6%)	1,794 (90.9%)	179 (9.1%)	1.12 (0.53- 2.39)	1.06 (0.46- 2.45)

9	I was given information in a way I understood	94 (88.7%)	12 (11.3%)	1,651 (83.7%)	322 (16.3%)	1.46 (0.68-3.25)	1.38 (0.60- 3.17)
10	I received helpful advice	85 (80.2%)	21 (19.8%)	1,410 (71.5%)	563 (28.5%)	1.59 (0.92-2.77)	1.58 (0.93-.2.68)
11	The administrative staff treated me with courtesy and respect	102 (96.2%)	4 (3.8%)	1,839 (93.2%)	134 (6.8%)	1.65 (0.61- 4.49)	1.76 (0.61-5.14)
12	The health worker involved my family helpfully	20 (18.9%)	86 (81.1%)	1,103 (55.9%)	870 (44.1%)	0.29 (0.12- 0.67)	0.29 (0.12- 0.67)
13	My privacy was respected	98 (92.4%)	8 (7.6%)	1,582 (80.2%)	391 (19.8%)	2.91 (1.88 -4.52)	2.94 (1.91 – 4.52)
14	I have the opportunity for follow up with the same	21 (19.8%)	85 (80.2%)	481 (24.4%)	1,492 (75.6%)	0.82 (0.55- 1.24)	0.83 (0.54- 1.23)

	health worker					
15	My personal information is kept confidential	89 (84.9%)	17 (16.0%)	1,428 (72.4%)	545 (27.6%)	2.07 (1.53- 2.79)	2.14 (1.61-2.85)
16	Referral to specialist is possible	104 (98.1%)	2 (1.9%)	1,847 (93.6%)	126 (6.4%)	3.50 (1.00-12.05)	3.64 (1.82- 13.00)
17	It is possible to see the health worker when needed	99 (93.4%)	7 (6.6%)	1,766 (89.5%)	207 (10.5%)	1.65 (0.54- 5.07)	1.72 (0.53 – 5.52)
18	It was easy to attend the health facility	91 (85.8%)	15 (14.2%)	1,705 (86.4%)	268 (13.6%)	0.98 (0.70- 1.39)	0.93 (0.63- 1.36)
19	I had enough time to attend the health facility	98 (92.4%)	8 (7.6%)	1,794 (90.9%)	179 (9.1%)	0.69 (0.42- 1.13)	0.67 (0.40 –1.13)
20	I could afford to attend the	61	45	1,384	589	0.61	0.60

	health facility for treatment	(57.5%)	(42.5%)	(70.2%)	(29.8%)	(0.36- 1.01)	(0.37 – 0.99)
21	I would advise my family members to come to this facility for treatment if they were pregnant	104 (98.1%)	2 (1.9%)	1,889 (95.7%)	85 (4.3%)	2.15 (0.57- 8.08)	2.03 (0.46- 8.92)

Table 4. ANC satisfaction in women with and without probable PTSD

		Women with probable PTSD (n=106)				Women without probable PTSD (n=1,973)				Fisher exact test of differences in category of responses
		Strongly agree	Agree	Disagree	Strongly Disagree	Strongly agree	Agree	Disagree	Strongly Disagree	
1	The health worker treated me with courtesy	13 (12.3%)	89 (84.0%)	3 (2.8%)	1 (0.9%)	472 (23.9%)	1,419 (71.9%)	64 (3.2%)	18 (0.9%)	P=0.026
2	The health worker listened to me carefully	9 (8.5%)	95 (89.6%)	1 (0.9%)	1 (0.9%)	403 (20.4%)	1,489 (75.5%)	66 (3.3%)	15 (0.8%)	P=0.004

3	The health worker explained things to me in a way I understood	8 (7.6%)	85 (80.2%)	7 (6.6%)	6 (5.6%)	330 (16.7%)	1,363 (69.1%)	218 (11.0%)	62 (3.1%)	P=0.008
4	The health facility was clean	7 (6.6%)	91 (85.9%)	8 (7.5%)	0 (0%)	299 (15.4%)	1,583 (80.2%)	74 (3.5%)	17 (0.9%)	P=0.017
5	The waiting room was clean	5 (4.7%)	93 (87.7%)	7 (6.6%)	1 (0.9%)	259 (13.1%)	1,540 (78.1%)	10.8 (5.5%)	66 (3.3%)	P=0.020
6	The latrine was clean	4 (3.8%)	71 (67.0%)	25 (23.6%)	6 (5.6%)	216 (10.9%)	459 (23.3%)	1,208 (61.2%)	90 (4.6%)	P=0.355
7	The waiting time was acceptable	5 (4.8%)	83 (78.3%)	12 (11.3%)	6 (5.6%)	224 (11.4%)	1,088 (55.1%)	509 (25.8%)	172 (7.7%)	P<0.001
8	I had enough time to	8	90	5	3	347	1,447	147	32	P=0.012

#	Statement									P-value
	discuss with health worker	(7.6%)	(84.9%)	(4.7%)	(2.8%)	(17.6%)	(73.3%)	(7.5%)	(1.6%)	
9	I was given information in a way I understood	8 (7.6%)	86 (81.1%)	9 (8.5%)	3 (2.8%)	298 (15.1%)	1,353 (68.6%)	250 (12.7%)	72 (3.6%)	P=0.049
10	I received helpful advice	7 (6.6%)	78 73.6%	17 (16.0%)	4 (3.8%)	277 (14.0%)	1,133 (57.4%)	422 (21.4%)	141 (7.2%)	P=0.009
11	The administrative staff treated me with courtesy and respect	6 (5.6%)	96 (90.6%)	2 (1.9%)	2 (1.9%)	342 (1.9%)	1,497 (75.9%)	96 (4.9%)	38 (1.9%)	P=0.002
12	The health worker involved my family helpfully	4 (3.8%)	16 (15.1%)	75 (70.8%)	11 (10.4%)	244 (12.4%)	626 (31.7%)	770 (39.0%)	333 (16.9%)	P<0.001

13	My privacy was respected	7 (6.6%)	91 (85.9%)	2 (1.9%)	6 (5.7%)	341 (17.3%)	1,241 (62.9%)	197 (10.0%)	194 (9.8%)	P<0.001
14	I have the opportunity for follow up with the same health worker	1 (0.9%)	20 (18.9%)	70 (66.0%)	1 (0.9%)	148 (7.5%)	333 (16.9%)	945 (47.9%)	547 (27.7%)	P<0.001
15	My personal information is kept confidential	6 (5,7%)	83 (78.3%)	10 (9.4%)	7 (6.6%)	326 (16.5%)	1,102 (55.9%)	315 (16.0%)	230 (11.7%)	P<0.001
16	Referral to specialist is possible	9 (8.5%)	95 (89.6%)	0 (0%)	2 (1.9%)	405 (20.5%)	1,442 (73.1%)	60 (3.0%)	66 (3.4%)	P=0.001
17	It is possible to see the health worker	8	91	5	2	358	1, 408	133	74	P=0.009

#	Statement									P-value
	when needed	(7.6%)	(85.8%)	(4.7%)	(1.9%)	(18.1%)	(71.4%)	(6.7%)	(3.8%)	
18	It was easy to attend the health facility	6 (5.7%)	85 (80.2%)	11 (10.4%)	4 (3.8%)	405 (20.5%)	1,300 (65.9%)	169 (8.6%)	99 (5.0%)	P<0.001
19	I had enough time to attend the health facility	6 (5.7%)	83 (78.3%)	10 (9..4%)	7 (6.6%)	383 (19.4%)	1,358 (68.8%)	97 (4.9%)	135 (6.8%)	P<0.001
20	I could afford to attend the health facility for treatment	2 (1.9%)	59 (55.7%)	32 (30.2%)	13 (12.3%)	203 (10.3%)	1,181 (59.9%)	379 (19.2%)	210 (10.6%)	P<0.001
21	I would advise my family members to come to this facility for treatment if they were pregnant	10 (9.4%)	94 (88.7%)	0 (0%)	2 (1.9%)	516 (26.2%)	1,373 (69.6%)	44 (2.2%)	40 (2.0%)	P<0.001

Discussion

Our facility-based cross-sectional study identified high levels of exposure to life-threatening injury, illness and suffering but paradoxically a low prevalence of antenatal PTSD (5.1%) (6), similar to previous survey of PTSD symptoms conducted in Nigeria (27). Over half the participants had directly experienced trauma, with physical violence victimisation especially common. Nearly half of those reporting physical violence screened positive for IPV. Very few women disclosed sexual trauma. There may be several reasons for nondisclosure. Stigma associated with sexual assault deters many victim-survivors from speaking about their experiences (28). Sexual discrimination against girls is common in Ethiopia (29) and worldwide. A multiregional study conducted by World Health Organization estimated the prevalence of lifetime sexual violence among women to be 30% (30). In the Ethiopian Demographic and Health Survey (EDHS) 2016, 10% of women aged 15–49 years were reported to have experienced sexual violence (31) Furthermore, previous studies have found that many women and girls in Ethiopia do not disclose their experience of domestic violence (32), despite known high prevalence (33). Reasons include shame, embarrassment and fear of disclosure related consequences. Less than 15% of our sample disclosed IPV, despite the estimated prevalence of antenatal IPV in Ethiopia ranging between 21.0% (34) and 26.1% (35). IPV has been associated with post-partum depressive symptoms in the Ethiopian context (34). Our study found that IPV was significantly associated with antenatal PTSD symptoms.

Among participants who had experienced a previous pregnancy, less than 1% of clinical records documented obstetric complications. This is not consistent with national estimates of around 20% (36). Prolonged labour is the leading cause of major direct obstetric morbidity in Ethiopia, accounting for 23.4%. Hypertensive disorders are the second most frequent cause of major direct maternal morbidity with a prevalence of around 11% (36). Reasons behind low recording of obstetric complications need to be further explored. Women with previous obstetric complications may be

less prepared or less likely to attend ANC in subsequent pregnancies, or feel unable to disclose previous complications to health workers, for example in case it negatively impacts their care. It is also likely health workers may not actively ask about previous complications or do so in a manner that does not promote disclosure. Unfortunately, disrespectful and abusive care during childbirth is widespread in Ethiopia (9), as in other countries. Reported prevalence of disrespect and abuse during childbirth in Ethiopia ranges from 21.1% to 98.9% (9). Qualitative research has described how fear of using health facilities is a common consequence of experiencing disrespectful care (37).

Women with antenatal PTSD showed a lower overall satisfaction with ANC compared to women without antenatal PTSD but paradoxically scored certain aspects of care higher than women without PTSD. Indeed, they endorsed more items regarding respect for their privacy and confidentiality. This finding should be interpreted with caution due to potential for response bias if women believe their answers may influence further care. Women with probable PTSD were less likely than women without probable PTSD to report that the health worker engaged their family helpfully and that they could afford to attend antenatal care. As IPV was more prevalent among participants with probable PTSD, training on sensitive ways to involve family members and ensure women are supported to attend ANC is a particular priority. Women with probable PTSD were less likely to endorse "strongly agree" responses for most aspects of care provided, indicating a systematic difference in reporting and highlighting the need to develop sensitive and validated scales to adequately capture the actual experiences of women accessing care.

Our results highlight the importance of person-centred maternity care to improve outcomes and experience of care for women. Previous studies from LMICs indicate that higher ratings of ANC dignity, respect, and supportiveness are associated with fewer newborn complications and increased willingness to access ANC for future pregnancies (38). The absence of person-centred maternity care has been associated with birth and post-partum complications (39). In 2019, the Federal Ministry of

Health of Ethiopia introduced the Ethiopian Primary Health care Clinical Guidelines (EPHCG) promoting integrated, person-centred and evidence-based care in health centres (40). In order to promote person-centred maternity care, systemic changes are required in tandem with health worker training and supervision. The ASSET team collaborated with the Ministry of Health to adapt training resources in person-centred care for maternal care in Ethiopia(41) Beyond training, services need to empower women with greater understanding of their rights and instil trust to facilitate disclosure of trauma; however, this requires cultural changes at the organisational level, including support and supervision for health workers, manageable caseloads and conductive clinical environments (42). Previous research in South Africa highlighted that efforts to improve detection of mental health conditions and IPV need to occur concurrently with development of pathways for referral and treatment/support for women to be confident their needs will be listened to and acted upon (42).

Despite high levels of high trauma exposure, only a small proportion of women in our sample met criteria for PTSD. Qualitative research is required to explore how women respond to and cope with traumatic life events in this context. While perinatal depression is widely researched, less is known about the impacts of PTSD and trauma exposure on woman and child outcomes (43). Our finding of high comorbidity between perinatal PTSD, depressive symptoms, and IPV highlights the need for studies to measure a range of CMDs and aetiologically relevant social determinants. Further research into perinatal PTSD and trauma symptoms, especially their association with birth and other obstetric trauma. Such studies in representative community samples will aid the development of trauma-informed, context-appropriate interventions for women across the perinatal period.

Strengths and Limitations

This is the first study in Ethiopia and one of few from LMICs, to investigate the relationships between trauma symptoms, IPV, obstetric complications and ANC satisfaction using validated measures in a large sample. Limitations include potential under detection of traumatic events, PTSD and IPV despite use of validated measures and highlights need for further studies how trauma symptoms may manifest differently in this setting and development of targeted interventions to elicit sensitive disclosure. ANC satisfaction may have been impacted by response bias; attention is needed to confidential ways of providing women with safe spaces to candidly disclose their experiences. The low recording of obstetric complications, inconsistent with national prevalence estimates, highlights how health system strengthening initiatives need to address the quality of data recording by clinicians. Future research should explore maternal health care workers' and other relevant stakeholders', including intimate partners' perspectives on high quality ANC and person-centred care.

Conclusions

Pregnant women attending ANC in Ethiopia report frequent exposure to traumatic life events, particularly physical violence. However, the prevalence of detected probable PTSD is low. IPV is associated with perinatal PTSD symptoms. Obstetric complications and IPV were under-reported, relative to previous studies in Ethiopia. ANC satisfaction was lower in women with probable PTSD. Our study highlights the importance of woman-centred approaches for improving recognition of trauma exposure, antenatal PTSD and the quality of ANC.

Funding acknowledgments

The research underpinning the findings presented in this paper was funded by the National Institute of Health and Care Research (NIHR) Global Health Research Unit on Health System Strengthening in

Sub-Saharan Africa (ASSET), King's College London (GHRU 16/136/54) using UK aid from the UK

Government. CH and MP are funded by an NIHR global health research group on homelessness and

mental health in Africa (HOPE; NIHR134325). The views expressed in this publication are those of the

authors and not necessarily those of the NHS, the National Institute for Health and Care Research or

the Department of Health and Social Care, England. CH is also funded by the Wellcome Trust through

grants 222154/Z20/Z (SCOPE) and 223615/Z/21/Z (PROMISE).

Acknowledgements

For the purpose of open access, the authors have applied a Creative Commons Attribution (CC BY)

licence to any Author Accepted Author Manuscript version arising from this submission.

We thank all the women who kindly agreed to participate in this study.

Author Contributions

RC- Conceptualisation; Data Curation; Methodology Formal analysis; Writing-original draft

preparation; LK- Conceptualisation; Formal analysis; Writing- Reviewing & Editing; AM-

investigation; Writing- Review & Editing; TE- Investigation; Project Administration; Resources;

Writing- Reviewing & Editing; AA- Project Administration; Investigation; AA- Writing- Reviewing &

Editing; RCK- Writing- Reviewing &Editing; JS- Writing – Reviewing & Editing; LMH- Writing-

Reviewing & Editing; MP- Writing- Reviewing & Editing; CH- Conceptualisation; Supervision;

Methodology; Funding acquisition; Witing- Reviewing & Editing. -

References:

1.   McNab SE, Dryer SL, Fitzgerald L, Gomez P, Bhatti AM, Kenyi E, et al. The silent burden: a landscape analysis of common perinatal mental disorders in low- and middle-income countries. BMC Pregnancy Childbirth. 2022; 22(1):1–14. doi:10.1186/s12884-022-04589-z

2.   Roddy Mitchell A, Gordon H, Lindquist A, Walker SP, Homer CSE, Middleton A, et al. Prevalence of Perinatal Depression in Low- and Middle-Income Countries: A Systematic Review and Meta-analysis. JAMA Psychiatry. 2023;80(5):425–31. doi:10.1001/jamapsychiatry.2023.0069

3.   Fisher J, Cabral de Mello M, Patel V, Rahman A, Tran T, Holton S, et al. Prevalence and determinants of common perinatal mental disorders in women in low- and lower-middle-income countries: a systematic review. Bull World Health Organ. 2012;90(2):139G-149G. doi: 10.2471/BLT.11.091850

4.   Howard LM, Khalifeh H. Perinatal mental health: a review of progress and challenges. World Psychiatry. 2020;19(3):313. doi: 10.1002/wps.20769

5.   McNab S, Fisher J, Honikman S, Muvhu L, Levine R, Chorwe-Sungani G, et al. Comment: silent burden no more: a global call to action to prioritize perinatal mental health. BMC Pregnancy Childbirth. 2022;22(1). doi: 10.1186/s12884-022-04645-8

6.   Sawyer A, Ayers S, Smith H. Pre- and postnatal psychological wellbeing in Africa: A systematic review. Journal of Affective Disorders. 2010; 123(1-3):17-29. doi: 10.1016/j.jad.2009.06.027.

7.   Yildiz PD, Ayers S, Phillips L. The prevalence of posttraumatic stress disorder in pregnancy and after birth: A systematic review and meta-analysis. J Affect Disord. 2017 ;208:634–45. doi: 10.1016/j.jad.2016.10.009.

8.   Anniverno R, Bramante A, Mencacci C, Durbano F. Anxiety Disorders in Pregnancy and the Postpartum Period. In: New Insights into Anxiety Disorders. 2013. doi: 10.5772/52786

9.   Mengesha MB, Desta AG, Maeruf H, Hidru HD. Disrespect and Abuse during Childbirth in Ethiopia: A Systematic Review. Biomed Res Int .2020; :8186070. doi: 10.1155/2020/8186070

10.   Sperlich M, Seng JS, Li Y, Taylor J, Bradbury-Jones C. Integrating Trauma-Informed Care Into Maternity Care Practice: Conceptual and Practical Issues. Journal of Midwifery and Women's Health. 2017; 62(6):661-672. doi: 10.1111/jmwh.12674.

11.   Goldenberg RL, McClure EM, Saleem S. Improving pregnancy outcomes in low- and middle-income countries. Reproductive Health. 2018; 15. doi: 10.1186/s12978-018-0524-5.

12.   Sudhinaraset M, Afulani P, Diamond-Smith N, Bhattacharyya S, Donnay F, Montagu D. Advancing a conceptual model to improve maternal health quality: The Person-Centered Care Framework for Reproductive Health Equity. Gates Open Research 2017 1:1 doi: 10.12688/gatesopenres.12756.1

13.   Seward N, Hanlon C, Abdella A, Abrahams Z, Alem A, Araya R, et al. HeAlth System StrEngThening in four sub-Saharan African countries (ASSET) to achieve high-quality, evidence-informed surgical, maternal and newborn, and primary care: protocol for pre-implementation phase studies. Glob Health Action. 2022;15(1): 1987044. doi: 10.1080/16549716.2021.1987044.

14.   Eshetu T, Fekadu E, Abdella A, Mulushoa A, Medhin G, Belina M, et al. Towards person-centred maternal and newborn care in Ethiopia: a mixed method study of satisfaction and experiences of care. ResearchSquare [Preprint] 2024 [cited 22 Oct 2024]. Available from: https://assets-eu.researchsquare.com/files/rs-3972651/v1/0bb39d54-185f-42b5-87ea-e74ad6d9d446.pdf?c=1709224168 . doi: 10.21203/rs.3.rs-3972651/v1

15.   CSA. Central Statistical Agency Population; Population Projections for Ethiopia 2007-2037. Central Statistical Agency Population, Ethiopia. 2013. Available from: https://www.statsethiopia.gov.et/wp-content/uploads/2019/05/ICPS-Population-Projection-2007-2037-produced-in-2012.pdf

16.   Hartung C, Anokwa Y, Brunette W, Lerer A, Tseng C, Borriello G. Open data kit: Tools to build information services for developing regions. In: ACM International Conference Proceeding Series. 2010. doi:10.1145/2369220.2369236.

17.   Girma E, Ametaj A, Alemayehu M, Milkias B, Yared M, Misra S, et al. Measuring Traumatic Experiences in a Sample of Ethiopian Adults: Psychometric Properties of the Life Events Checklist-5. European journal of trauma & dissociation, ;6(4):100298. doi: 10.1016/j.ejtd.2022.100298.

18.   Blevins CA, Weathers FW, Davis MT, Witte TK, Domino JL. The Posttraumatic Stress Disorder Checklist for DSM-5 (PCL-5): Development and Initial Psychometric Evaluation. J Trauma Stress. 2015 Dec 1;28(6):489–98. doi: 10.1002/jts.22059.

19.   Ng LC, Serba EG, Dubale BW, Fekadu A, Hanlon C. Posttraumatic stress disorder intervention for people with severe mental illness in a low-income country primary care setting: a randomized feasibility trial protocol. Pilot and Feasibility Studies. 2021;7(149):1–15. doi:10.1186/s40814-021-00883-3

20.   Hanlon C, Medhin G, Selamu M, Breuer E, Worku B, Hailemariam M, et al. Validity of brief screening questionnaires to detect depression in primary care in Ethiopia. J Affect Disord. 2015;186:32–9. doi: 10.1016/j.jad.2015.07.015.

21.   Habtamu K, Birhane R, Medhin G, Hanlon C, Fekadu A. Psychometric properties of screening questionnaires to detect depression in primary healthcare setting in rural Ethiopia. BMC Primary Care. 2022;23(138):1–14. doi:10.1186/s12875-022-01755-2.

22.   Zink T, Levin L, Putnam F, Beckstrom A. Accuracy of five domestic violence screening questions with nongraphic language. Clin Pediatr (Phila). 2007;46(2):127–34, doi: 10.1177/0009922806290029.

23.   Straus MA, Hamby SL, Boney-McCoy S, Sugarman DB. The revised conflict tactics scales (CTS2) development and preliminary psychometric data. J Fam Issues. 1996;17(3):283–316. doi: 10.1177/019251396017003001.

24.   Bitew T, Hanlon C, Kebede E, Medhin G, Fekadu A. Antenatal depressive symptoms and maternal health care utilisation: A population-based study of pregnant women in Ethiopia. BMC Pregnancy Childbirth. 2016;16(301). doi: 10.1186/s12884-016-1099-1.

25.   Mayston R, Habtamu K, Medhin G, Alem A, Fekadu A, Habtamu A, et al. Developing a measure of mental health service satisfaction for use in low income countries: a mixed methods study. BMC Health Serv Res. 2017;17(183). doi: 10.1186/s12913-017-2126-2 .

26.   Fekadu A, Medhin G, Selamu M, Hailemariam M, Alem A, Giorgis TW, et al. Population level mental distress in rural Ethiopia. BMC Psychiatry. 2014;14(194):1–13. doi: 10.1186/1471-244X-14-194.

27.   Adewuya AO, Ologun YA, Ibigbami OS. Post-traumatic stress disorder after childbirth in Nigerian women: Prevalence and risk factors. Vol. 113, BJOG: An International Journal of Obstetrics and Gynaecology. 2006; 113(3):284-8. doi:10.1111/j.1471-0528.2006.00861.x.

28.   Adinew YM, Hagos MA. Sexual violence against female university students in Ethiopia. BMC Int Health Hum Rights. 2017;17(19). doi: 10.1186/s12914-017-0127-1.

29.   Koohi-kamali F. Intrahousehold Inequality and Child Gender Bias in Ethiopia. World Bank Policy Research Working Paper. 2008. Available from: http://documents.worldbank.org/curated/en/996101468251996959/Intrahousehold-inequality-and-child-gender-bias-in-Ethiopia

30.   World Health Organization. Global and regional estimates of violence against women. Geneva; 2018. Available from: https://www.who.int/publications/i/item/9789241564625

31.   Central Statistical Agency Addis Ababa E. Ethiopian Demographic and Health Survey 2016: Key Indicators Report. Ethiopians Water Sector Development Program. Maryland, USA; 2016. Available from: https://dhsprogram.com/pubs/pdf/fr328/fr328.pdf.

32.   Bifftu BB, Dachew BA, Tiruneh BT, Gezie LD, Guracho YD. Domestic violence related disclosure among women and girls in Ethiopia: A systematic review and meta-Analysis. Reprod Health. 2019;16(1):1–10. doi: 10.1186/s12978-019-0845-z .

33.   Chernet AG, Cherie KT. Prevalence of intimate partner violence against women and associated factors in Ethiopia. BMC Womens Health. 2020;20(1). doi: 10.1186/s12905-020-0892-1.

34.   Belay S, Astatkie A, Emmelin M, Hinderaker SG. Intimate partner violence and maternal depression during pregnancy: A community-based cross-sectional study in Ethiopia. PLoS One. 2019;14(7):e0220003. doi: 10.1371/journal.pone.0220003.

35.   Alebel A, Kibret GD, Wagnew F, Tesema C, Ferede A, Petrucka P, et al. Intimate partner violence and associated factors among pregnant women in Ethiopia: A systematic review and meta-analysis. Reproductive Health. 2018; 15. doi: 10.1186/s12978-018-0637-x.

36.   Geleto A, Chojenta C, Taddele T, Loxton D. Magnitude and determinants of obstetric case fatality rate among women with the direct causes of maternal deaths in Ethiopia: A national cross sectional study. BMC Pregnancy Childbirth. 2020;20(1). doi: 10.1186/s12884-020-2830-5.

37.   Werdofa HM, Lindahl AK, Lulseged B, Thoresen L. Women's perspectives on disrespect and abuse experiences during childbirth in a teaching hospital in Southwest Ethiopia: a qualitative study. Sex Reprod Health Matters. 2022;30(1): 2088058. doi: 10.1080/26410397.2022.2088058.

38.   Sudhinaraset M, Landrian A, Afulani PA, Diamond-Smith N, Golub G. Association between person-centered maternity care and newborn complications in Kenya. International Journal of Gynecology and Obstetrics. 2020; 148(1):27-34. doi: 10.1002/ijgo.12978.

39.   Raj A, Dey A, Boyce S, Seth A, Bora S, Chandurkar D, et al. Associations Between Mistreatment by a Provider during Childbirth and Maternal Health Complications in Uttar Pradesh, India. Matern Child Health J. 2017;21(9):1821-1833. doi: 10.1007/s10995-017-2298-8.

40.   Feyissa YM, Hanlon C, Emyu S, Cornick RV, Fairall L, Gebremichael D, et al. Using a mentorship model to localise the Practical Approach to Care Kit (PACK): From South Africa to Ethiopia. BMJ Glob Health. 2018; 3:e001108.

41. CDT-Africa. ASSET. CSS Training Manuals. 2024 [cited 2024 Oct 22]. Available from: https://www.cdt-africa.org/index.php/projets/completed/asset

42. Abrahams Z, Boisits S, Schneider M, Honikman S, Lund C. Facilitators and barriers to detection and treatment of depression, anxiety and experiences of domestic violence in pregnant women. Sci Rep. 2023;13(1). doi: 10.1038/s41598-023-36150-z.

43. Giannandrea SAM, Cerulli C, Anson E, Chaudron LH. Increased risk for postpartum psychiatric disorders among women with past pregnancy loss. J Womens Health. 2013;22(9) :760-8. doi: 10.1089/jwh.2012.4011.