Cervicovaginal inflammation and HIV target cell activation in adolescent girls and young women with *Chlamydia trachomatis* infections

Smritee Dabee,[1,2]* Shaun Barnabas,[1,2]* Brian Kullin,[1,3] Bart Versteeg,[4] Nonhlanhla Mkhize,[5] Etienne Muller,[5] Venessa Maseko,[5] David A. Lewis,[6,7] Janan Dietrich,[8] Glenda Gray,[8,9] Katherine Gill,[1,3] Linda-Gail Bekker[1,3], Musalula Sinkala,[1] Darren P. Martin,[1] Sylvia M. Bruisten,[10,11] Heather B. Jaspan,[1,12,13] Jo-Ann S. Passmore[1,3,14#]

[1]Institute of Infectious Disease and Molecular Medicine (IDM), Division of Medical Virology, Dept. Pathology, University of Cape Town, Cape Town, South Africa; [2]Desmond Tutu HIV Centre, University of Cape Town, South Africa; [3]DST-NRF CAPRISA Centre of Excellence in HIV Prevention, University of Cape Town, Cape Town, South Africa; [4]Knowledge institute of the Dutch association of Medical Specialists, Utrecht, Netherlands; [5]Centre for HIV and STIs, National Institute for Communicable Disease, National Health Laboratory Service, South Africa; [6]Western Sydney Sexual Health Centre, Western Sydney Local Health District, Parramatta, Australia; [7]Sydney Infectious Diseases Institute & Sydney Medical School-Westmead, University of Sydney, Sydney, Australia; [8]Perinatal HIV Research Unit, Faculty of Health Sciences, University of the Witwatersrand, Diepkloof, Johannesburg, South Africa; [9]South African Medical Research Council, Cape Town, South Africa; [10]Department of Infectious Diseases, Public Health Service Amsterdam, Amsterdam, the Netherlands; [11]Amsterdam Infection and Immunity Institute, Academic Medical Center, University of Amsterdam, Amsterdam, the Netherlands; [12]Seattle Children's Research Institute, Seattle, Washington, USA; [13]University of Washington, Seattle, Washington, USA; [14]National Health Laboratory Service, Groote Schuur Hospital, Cape Town, South Africa

#Address correspondence to Professor Jo-Ann S. Passmore, Institute of Infectious Disease and Molecular Medicine, University of Cape Town, Medical School, Anzio Road, Observatory, 7925 Cape Town, South Africa. E-mail: Jo-ann.Passmore@uct.ac.za, Tel: +27-21-650 7963

Word count: 2933; Abstract:293

Figures: 4; Tables: 1

Running title:
Chlamydia-associated inflammation in adolescents [49 characters, with spaces]

Abstract

Background: Adolescent girls and young women (AGYW) in sub-Saharan Africa are disproportionately affected by *Chlamydia trachomatis* (CT) infections, a leading cause of pelvic inflammatory disease (PID), infertility, and increased HIV susceptibility. CT-induced genital inflammation disrupts mucosal barriers and activates HIV target cells, compounding reproductive and immunological risks. We investigated the impact of CT infection in AGYW from two South African regions, focusing on genital inflammation and immune activation.

Methods: An observational cohort of 298 sexually active AGYW (16–22 years) from Cape Town and Johannesburg was enrolled. Participants in Johannesburg attended a single study visit, while those in Cape Town were followed longitudinally for up to three visits over 6–8 months. Cytokine profiling of cervicovaginal samples assessed inflammatory responses, while cervical cytobrush-derived CD4+ T cells were analyzed for CD38, Ki67 and CCR5 expression. CT *ompA* genotyping examined strain diversity.

Results: CT prevalence was 2.4 times higher in Cape Town (41.6%) than in Johannesburg (17.4%). Infection with CT led to an upregulation of inflammatory cytokines, including IL-1β, IL-6, and TNF-α, with a stronger inflammatory response observed in Johannesburg. Genotypic analysis revealed regional variation: genovar D, predominant in Cape Town, was associated with lower IFN-γ concentrations, whereas genovar E, more common in Johannesburg, was associated with higher IFN-γ levels. AGYW who experienced CT infection across multiple study visits did not exhibit elevated genital tract cytokine levels compared to those with a single infection. However, they did show an increase in activated and proliferating cervical CD4+ T cells, which may contribute to heightened HIV susceptibility.

Conclusions: CT genotypic diversity and regional variation are associated with differences in immune responses in AGYW, alongside important contributions from host immune history.

These findings support the need for integrated sexual health services and inform future regionally relevant vaccine strategies.

Keywords: Adolescent girls and young women, AGYW, Chlamydia trachomatis, ompA, genital inflammation, Sexual and reproductive health

1   Introduction

Young women in sub-Saharan Africa face disproportionately high risks of HIV infection,

driven by both biological and socio-behavioural vulnerabilities (1). Genital inflammation is a

critical factor that increases the risk of HIV acquisition (2). Sexually transmitted infections

(STIs), including *Chlamydia trachomatis* (CT) are major drivers of genital inflammation. CT

infections are frequently asymptomatic in women, leading to under-treatment despite

associations with pelvic inflammatory disease (PID), infertility, ectopic pregnancy, and

increased HIV risk due to subclinical inflammation (3).

CT infection induces genital cytokine responses that play a key role in pathogenesis

(4). Pro-inflammatory mediators such as IL-1β, IL-6, and IP-10 can also exacerbate tissue

damage and increase HIV susceptibility through mucosal barrier disruption and CD4+ T cell

activation (2,5,6). Persistent CT infection can further amplify these inflammatory responses,

contributing to immune dysregulation and impaired clearance (6).

CT exhibits substantial genetic diversity that influences transmission dynamics, tissue

tropism, and host immune responses. Urogenital CT infections in women, including AGYW

in sub-Saharan Africa, are dominated by a limited number of *ompA*-defined genotypes,

particularly E, D, and F, with genotypes J, G, and K contributing smaller but consistent

proportions across regions (7–8). Variation within and between *ompA* genotypes has been

associated with differences in immune activation, persistence, and clinical outcomes,

underscoring the relevance of characterizing circulating CT genotypes in high-burden

African AGYW populations (9).

AGYW are particularly vulnerable to CT infection due to socio-behavioral,

anatomical, and hormonal factors. The adolescent cervix has a larger area of ectropion (10),

exposing columnar epithelium that is more permissive to bacterial adhesion and invasion than

the stratified squamous epithelium in mature women. Together with delayed diagnosis and

reinfection, these factors increase the risk of prolonged inflammation and reproductive

complications in AGYW (11).

Given these challenges, it is critical to investigate the complex interplay between CT

genotypes, host inflammatory responses, and mucosal immune activation in AGYW. We

examined how CT genotypes influence genital inflammation and immune activation in

AGYW from two South African regions.

Materials and Methods

Cohort description

Between November 2013 and February 2015, AGYW (16–22 years) were recruited from

Cape Town (n=149) and Johannesburg (n=149) through the Desmond Tutu HIV Centre

(DTHC) and the Perinatal HIV Research Unit (PHRU), respectively (12). The study was

cross-sectional in Johannesburg while the cohort from Cape Town arm were followed

longitudinally for up to three visits, using a targeted enrolment strategy with predefined

sample sizes (Barnabas et al., 2018). For the longitudinal cohort, study visits were 3 monthly

unless participants were using Net-EN, in which case it was 2 monthly. AGYW using the

injectable contraceptives (DMPA or Net-EN) had visits scheduled two weeks post-injection,

while non-users were scheduled during the luteal phase (days 14–28) of their menstrual

cycle. Participants were excluded if they were HIV-positive, pregnant, sexually inactive,

douched or used spermicides in the past two days, or took antibiotics in the past two weeks.

Research Ethics Committees of the Universities of Cape Town and Witwatersrand approved

the study (#267/2013 and #M130745, respectively). Written informed consent was obtained

from participants 18+ years and assent was provided by those <18 years, with written consent

from a parent/legal guardian.

Sample collection

At enrolment, participants completed a socio-demographic questionnaire (12). At each visit,

cervicovaginal fluid was collected using a Softcup® menstrual cup, inserted for one hour.

During the clinical speculum exam, a Dacron vulvovaginal swab was taken for STI testing, a

FLOQSwab™ collected lateral vaginal wall samples for BV screening (Nugent scoring; 13),

and cervical mononuclear cells were obtained using a Digene® endocervical cytobrush.

Testing for STIs and BV

Vulvovaginal samples were tested for discharge-causing STIs (CT, *Neisseria gonorrhoeae*

[NG], *Trichomonas vaginalis* [TV], *Mycoplasma genitalium* [MG]), as well as ulcer-causing

STIs (HSV-1, HSV-2, *Haemophilus ducreyi* [HD], *Treponema pallidum*), using real-time

multiplex PCRs (14). BV was diagnosed by Nugent score (13). AGYW positive for CT were

treated with Azithromycin (1g orally; single dose), those with NG given Ceftriaxone (250mg

intramuscular injection; single dose), and those positive for BV (Nugent 7-10) or TV were

given Metronidazole (400mg orally, twice daily; 7 days), according to South African national

guidelines at the time. AGYW with CT or NG were given partner notification letters and

referred to the local Department of Health clinic.

Chlamydia *ompA* genotyping

OmpA genovar diversity of CT was analyzed by MLST genotyping (8) (15). Sequence data

were analysed using the CT database hosted on PubMLST

(https://pubmlst.org/bigsdb?db=pubmlst_chlamydiales_seqdef). The *ompA* genovars were

assigned by comparison to an offline reference database described previously (8) (15).

Measurement of cytokine concentrations

Softcups were collected (in 50ml Falcon tubes), maintained at 4^{o}C after collection (as

previously described; 16, 17), and processed at a site-adjacent laboratory within 4 hours for

storage at -80^{o}C. Processing and elution of cervicovaginal fluid included inverting the softcup

membrane using a sterile wooden disposable spatula in a BSL2 hood and centrifuging (1500

rpm, 10 minutes). The volume of secreted fluid was calculated, as previously described (17).

The softcup was discarded. Assuming 1g=1mL, secretions were diluted 10 times with PBS

and stored immediately at -80oC in aliquots. All softcup samples were transported to the

UCT laboratory on dry ice, aliquots thawed in batches, and cytokines measured by Luminex

using the same lot number of kits (16). Prior to Luminex, all samples were filtered by

centrifugation (1950g, 10 min, 4°C) using SPIN-X® 0.2µm filters. Two different kits were

used: the Human Cytokine Group I 27-plex (FGF-basic, Eotaxin, G-CSF, GM-CSF, IFN-γ,

IL-1β, IL-1RA, IL-2, IL-4, IL-5, IL-6, IL-7, IL-8, IL-9, IL-10, IL-12(p70), IL-13, IL-15, IL-

17, IP-10, MCP-1, MIP-1α, MIP-1β, PDGF-BB, RANTES, TNF-α, and VEGF) and the

Human Cytokine 21-plex kit (IL-1α, IL-2RA, IL-3, IL-12(p40), IL-16, IL-18, CTACK,

GRO-α, HGF, IFN-α2, LIF, MCP-3, M-CSF, MIF, MIG, NGF, SCF, SCGF-β, SDF-1α,

TNF-β, and TRAIL) (Bio-Rad). Experiments followed manufacturer instructions. Samples

from both sites were randomly allocated on the different assay plates, with the same ten

samples included in all plates: Inter- and intra-plate correlations (Spearman correlations)

between these ten samples were carried out for each cytokine amongst all the plates. IL-15

was excluded as the values did not pass the QC. The cytokines IL-2, IL-5 and RANTES were

not included in the analysis as 76.0%, 64.2% and 52.6% of the values respectively were

below detectable range. For the rest, detection limits ranged from 1-307 pg/ml, and

concentrations were determined using 5PL regression. Values below detection limits were

recorded as half the lowest measured concentration.

Flow cytometry

CMC samples from Cape Town participants were collected and analyzed via flow cytometry

at each visit. Cervical cytobrush samples were processed as previously described (16), with

the cell pellet resuspended in PBS, transferred to a 96-well plate, centrifuged at 300g for 3

minutes at 4°C, and washed before staining. An eight-color flow cytometry panel was used to

assess CCR5 (APC; BD), activation markers CD38 (PE-Cy7; eBioscience) and HLA-DR

(PE; BD), and proliferation marker Ki67 (FITC; BD) in CD4+CD3+ (APC-H7; BD) and

CD8+CD3+ (QDot605, Invitrogen) T cells. CD8+ T cells is not shown. A dump channel

(LIVE/DEAD® Fixable Violet Dead Cell Stain [Invitrogen], CD14 [PacBlue, BD], CD19

[Pac Blue, Invitrogen]) excluded dead cells, B-cells, and monocytes, respectively. Cells were

first incubated with the live/dead marker for 20 minutes at room temperature, washed twice,

and stained with extracellular markers. After 30-minute ice incubation, cells were washed,

fixed and permeabilized using Cytofix/Cytoperm (BD). Intracellular markers CD3 and Ki67

were added, and cells incubated for 30 minutes on ice. The pellet was resuspended in BD

CellFIX^{TM}, transferred to a FACS tube, wrapped in foil, and stored at 4°C until acquisition on

the BD LSRFortessa^{TM}. Data were acquired within 48 hours, with compensation tubes

prepared for each run. FlowJo (v10.0.8; Treestar) was used for analysis, with gating based on

FMO controls.

Statistical analysis

Statistical analyses and linear regressions were carried out using STATA 12.0 and R v3.6.0.

Graphs and figures were generated using Prism 6 and R. Unpaired and paired continuous

variables were compared using the Mann–Whitney U and Wilcoxon signed-rank tests,

respectively, and categorical variables using two-sided chi-squared tests. Analyses were

adjusted for multiple comparisons using false discovery rate step down procedures.

Results

A total of 298 sexually active AGYW (median age: 18 years; IQR: 17–20) from Cape Town and Johannesburg were included (12) (16). Johannesburg participants had a single-visit study, while Cape Town participants were followed over three visits spanning 6–8 months. Contraceptive use differed by site: in Johannesburg, 63.8% (77/121) of AGYW reported condom use alone, whereas in Cape Town 70.5% (105/149) used injectable hormonal contraception (DMPA or Net-EN). CT was the most prevalent STI (29.5%, 88/298), with CT prevalence being significantly higher in Cape Town (41.6%, 62/149; 95% CI: 33.6–49.9%) than in Johannesburg (17.4%, 26/149; 95% CI: 11.8–24.3%; $p<0.0001$). Among those with CT, 71.6% (63/88) had another STI or BV.

Impact of CT infection on genital inflammation in AGYW

The genital cytokine responses to CT infection in AGYW at baseline, were assessed by multivariate linear regressions, adjusted for contraceptive use, co-infections with STIs and BV (Figure 1A). AGYW infected with CT exhibited significant upregulation of cytokines across all functional classes compared to uninfected individuals. These included IL-1β (β-coefficient=0.36; $p=0.022$), IL-6 (β=0.21; $p=0.038$), TNF-α (β=0.23; $p=0.015$), IP-10 (β=0.29; $p=0.035$), MIG (β=0.27; $p<0.0001$), G-CSF (β=0.28; $p=0.018$), HGF (β=0.27; $p=0.011$), SCF (β=0.13; $p=0.034$), IFN-γ (β=0.17; $p=0.025$), and IL-4 (β=0.17; $p=0.015$), after adjusting for multiple comparisons. A focused analysis on cytokines implicated in CT pathogenesis (IL-1α, IL-1β, IL-6, IL-8, TNF-α, TNF-β) and protection (IFN-γ), including AGYW with CT infection alone (n=26; no co-infections or BV) compared to uninfected controls (n=62; no STIs or BV) (Figure 1B), further confirmed that CT infection significantly elevated concentrations of inflammatory cytokines, including IL-1α ($p=0.03$), IL-6 ($p=0.01$),

IL-8 (p=0.05), IP-10 (p=0.002), and TNF-α (p=0.01). Although protective IFN-γ levels were

higher in CT cases compared to controls, this was not significant.

Chlamydia diversity and associated inflammation by study site

CT infection was more inflammatory in Johannesburg than in Cape Town at baseline (Figure

2A). In Cape Town, CT was linked to MIG upregulation (not significant after multiple

comparisons), while in Johannesburg, it was associated with increased concentrations of

34/44 cytokines. Among AGYW with CT alone (no other STIs or BV), inflammatory

cytokines tended to be higher in Johannesburg (n=4 CT+; no other STIs/BV) than Cape

Town (n=22 CT+; no other STIs/BV), although numbers were small (Figure 2B).

To assess whether CT strain diversity contributed to differences in inflammatory

responses, *ompA* genotyping was performed on 128 samples with sufficient DNA (Figure

2C), including 20/26 cases from Johannesburg and 108/121 cases from Cape Town (spanning

all three study visits). Eighteen samples were excluded due to low DNA quantity. The

distribution of *ompA* genovars did not differ significantly between sites (p = 0.1; χ^2 test), and

globally predominant urogenital genotypes were represented at both sites. In Cape Town,

genovar D was the most prevalent (21%, 13/62), followed by genovars E (18%, 11/62) and F

(18%, 11/62), while genovar G was detected in 5% of cases (3/62). In contrast, in

Johannesburg, genovars E and G were most common (23% each, 6/26), while genovar D was

detected in 19% (5/26) of cases and genovar F was not detected. Less common genovars (I, J,

K) were observed at low frequencies at both sites. No significant associations were observed

between individual CT genovars and inflammatory cytokines, after adjustment for multiple

comparisons (Figure 2D). However, the limited number of CT-positive samples, particularly

within specific genovars and at the Johannesburg site, constrained statistical power and

precluded site-specific comparisons of genovar-associated inflammatory profiles.

Inflammation associated with persistent CT infections

In Cape Town, where AGYW were followed longitudinally, 33.9% (43/127) were infected

with CT at visit 2, and 18.2% (16/88) were infected at visit 3 (Table 1). Among participants

with repeat CT detection, genotyping indicated that the infecting genovar was most often the

same across visits: 10 women were CT-positive at all three study visits, of whom 7/10

harbored the same genovar at each visit; 10 were CT-positive at visits 1 and 2 only, all of

whom (10/10) had the same genovar at both visits; and 5 were CT-positive at visits 1 and 3

but not visit 2, of whom 4/5 had the same genovar at both positive visits. These findings

suggest that repeated detection of CT in this cohort was more frequently associated with

persistence or re-exposure to the same strain, which may reflect ongoing exposure from the

same sexual partner or incomplete clearance, rather than acquisition of a new genotype. To

investigate whether persistent or repeat CT infections were associated with increased

inflammation, cervicovaginal cytokine concentrations were compared among women infected

at one, two, or three visits (Figure 3). The analysis revealed no significant differences in the

concentrations of any measured cytokines across these groups, regardless of the duration of

infection.

Genital tract HIV target-cell activation associated with CT infections

Cervical cytobrush-derived CD3+CD4+ T cells were used to assess the expression of CCR5

(HIV entry receptor on CD4+ cells), CD38 (a marker of cellular activation), and Ki67 (a

marker of T-cell proliferation) (Cape Town only). Figure 4A shows a representative figure of

the gating strategy. The frequency of activated cervical CD38$^+$ cells among CD3+CD4$^+$ T

cells was significantly higher in CT cases compared to controls (p=0.05; Figure 4B, top

panel). However, there were no significant differences in the frequencies of CD3+CD4$^+$ T

cells or those expressing CCR5 or Ki67. As a technical observation, CT+ cytobrush

specimens yielded higher total acquired live events under standardized acquisition settings

(Figure 4B, bottom panel).

We further compared CD3+CD4$^+$ T cell events in women infected with CT at one,

two, or three study visits (Figure 4C). The frequency of activated or proliferating cervical

CD3+CD4$^+$ T cells did not differ significantly between women infected at multiple time

points and those infected at a single visit. In a paired analysis of women infected at two study

visits (n=32), there was no significant difference in the frequency of T cells measured at the

second positive visit compared to the first positive visit, and frequencies did not correlate

significantly (CD3+CD4+ R^2=0.002 p=0.8; CD3+CD4+CD38+ R^2=0.02 p=0.5;

CD3+CD4+Ki67+ R^2=0.02 p=0.5). This suggests that inflammation did not resolve or

intensify with persistent or repeated infections (Figure 4C, lower panel).

Discussion

Understanding how asymptomatic CT infection contributes to HIV susceptibility in AGYW remains an important challenge for prevention. Here, we examined the intersection between CT infection, genital inflammation, and HIV target cell availability across two South African settings, highlighting a complex interplay between pathogen diversity, host immune history, and local mucosal context. Rather than implicating a single explanatory factor, our findings indicate that inflammatory and cellular immune responses to CT reflect multiple determinants, including strain-level variation, prior exposure, and site-specific host or vaginal milieu factors (microbiome), helping to explain why inflammatory outcomes may differ across populations despite exposure to common CT lineages.

Genotyping revealed few differences in ompA genovar distribution between sites, with genovar D predominating in Cape Town and genovars E and G more frequent in Johannesburg. Despite this, AGYW with CT infection in Johannesburg exhibited a stronger overall inflammatory signature, even though genovar G—enriched at this site—was associated with lower cytokine concentrations. As this conclusion is based on the markers measured, site-level inflammatory differences are unlikely to be explained by genotype alone and may instead reflect additional unmeasured inflammatory or regulatory pathways and broader geographic, host, or vaginal milieu factors.

While variation in ompA genovars was explored, emerging evidence indicates that host immune history is likely a more dominant determinant of inflammation. In the same cohort, prior CT exposure and the quality of CT-specific CD4$\square$ T cell responses strongly influenced genital immune profiles, with robust systemic Th1 responses inversely associated with mucosal inflammatory mediators (18). Women with untreated or recurrent CT infection showed increased genital T cell activation alongside impaired CT-specific T cell function, consistent with immune dysregulation following repeated exposure. Together, these findings

suggest that adaptive immune history may outweigh strain-level variation, and that genovar

associations should be regarded as hypothesis-generating rather than causal.

Strain-specific associations nonetheless highlight heterogeneity in CT

immunogenicity. The association of genovar D with lower IFN-γ and genovar E with higher

IFN-γ supports a role for pathogen genetic variation in shaping host responses. IFN-γ can

drive CT into a persistent state via induction of indoleamine-2,3-dioxygenase (IDO),

promoting immune evasion and chronic infection (19). Supporting this, Van Duynhoven et al.

(20) reported greater lower abdominal pain among women infected with F or G group

genotypes. Collectively, these findings may reflect variation in immunodominant T-cell

epitopes within the major outer membrane protein (MOMP) (21), reinforcing the value of

strain-level characterization for understanding CT pathogenesis and informing vaccine

design.

This study has several limitations. Longitudinal data were only available from Cape

Town, limiting analyses of repeat CT infections, and cellular activation measurements were

similarly restricted. Partner treatment data were not systematically collected, and post-

treatment cure testing was not performed. Longitudinal ompA genotyping in Cape Town

showed that repeat CT detection usually involved the same genovar, suggesting persistence

or re-exposure to the same strain, although reinfection from the same partner or incomplete

clearance cannot be excluded. Genotyping was based on ompA sequencing rather than

whole-genome sequencing, which may not capture within-genotype diversity (21) or

recombination events (22). The small number of CT-positive participants in Johannesburg

may also limit generalizability of site-specific genotype associations.

The high prevalence of CT-associated inflammation in AGYW highlights the need for

improved STI screening and integrated sexual and reproductive health services in high HIV-

burden settings (23). Regional differences in CT immunogenicity support tailored prevention

approaches, including regionally informed vaccine strategies. In conclusion, our findings

show that CT-associated genital immune responses in AGYW reflect the interplay of strain

diversity, regional context, and host immune history.

Acknowledgements

We would like to thank the WISH study team, especially Ms Penelope Ngcobo (participant

recruitment at DTHF), Sr Janine Nixon (sample collection), and all the young women

enrolled in the study. We also thank Ms Michelle Himschoot from GGD Amsterdam, who

assisted with the MLST lab work. This research was funded by the EDCTP Strategic Primer

grant (SP.2011.41304.038; PI J. Passmore, UCT) and a self-initiated grant from the South

African Medical Research Council (2017-2020; PI J. Passmore, UCT).

Conflict of Interest Statement:

The authors declare no potential conflicts of interest with respect to this research, authorship,

and/or publication of this article. The study was funded by the EDCTP Strategic Primer grant

and a self-initiated grant from the South African Medical Research Council. The funders had

no role in study design, data collection and analysis, decision to publish, or preparation of the

manuscript.

References

1. de Oliveira T, Kharsany AB, Graf T, Cawood C, Khanyile D, Grobler A, *et al.* Transmission networks and risk of HIV infection in KwaZulu-Natal, South Africa: a community-wide phylogenetic study. **Lancet HIV** 2017;4:e41–50. doi:10.1016/S2352-3018(16)30186-2

2. Masson L, Passmore JA, Liebenberg LJ, Werner L, Baxter C, Arnold KB, *et al.* Genital inflammation and the risk of HIV acquisition in women. **Clin Infect Dis** 2015;61:260–9. doi:10.1093/cid/civ298

3. Pillay J, Wingert A, MacGregor T, Gates M, Vandermeer B, Hartling L. Screening for chlamydia and/or gonorrhea in primary health care: systematic reviews on effectiveness and patient preferences. **Syst Rev** 2021;10:118. doi:10.1186/s13643-021-01658-w

4. Redgrove KA, McLaughlin EA. The role of the immune response in Chlamydia trachomatis infection of the male genital tract: a double-edged sword. **Front Immunol** 2014;5:534. doi:10.3389/fimmu.2014.00534

5. Arnold KB, Burgener A, Birse K, Romas L, Dunphy LJ, Shahabi K, *et al.* Increased inflammatory cytokines in the female reproductive tract are associated with altered mucosal barrier proteins and an influx of HIV-susceptible target cells. **Mucosal Immunol** 2016;9:194–205. doi:10.1038/mi.2015.51

6. Lijek RS, Helble JD, Olive AJ, Seiger KW, Starnbach MN. Pathology after Chlamydia trachomatis infection is driven by nonprotective immune cells distinct from protective populations. **Proc Natl Acad Sci U S A** 2018;115:2216–21. doi:10.1073/pnas.1711356115

7. Smelov V, Vrbanac A, van Ess EF, Noz MP, Wan R, Eklund C, *et al.* Chlamydia trachomatis strain types have diversified regionally and globally with evidence for recombination across geographic divides. **Front Microbiol** 2017;8:2195. doi:10.3389/fmicb.2017.02195

8. Versteeg B, Dubbink JH, Bruisten SM, *et al*. High-resolution multilocus sequence typing reveals novel urogenital Chlamydia trachomatis strains in women in Mopani district, South Africa. **Sex Transm Infect** 2015;91:510–12. doi:10.1136/sextrans-2014-051891

9. Lowry K, Apadinuwe S, Starr M, Starr S, Lynch KD, Solomon AW, *et al.* Population-based concordance of Chlamydia trachomatis genotypes between ocular and urogenital samples in Nauru. **Microbiol Spectr** 2025;13:e03205-24. doi:10.1128/spectrum.03205-24

10. Cagle AJ, Hu SY, Sellors JW, Bao YP, Lim JM, Li SM, *et al*. Use of an expanded gold standard to estimate the accuracy of colposcopy and visual inspection with acetic acid. **Int J Cancer** 2010;126:156–61. doi:10.1002/ijc.24719

11. Hoenderboom BM, van Benthem BHB, van Bergen J, Dukers-Muijrers N, Götz HM, Hoebe C, et al. Relation between Chlamydia trachomatis infection and pelvic inflammatory disease, ectopic pregnancy and tubal factor infertility. **Sex Transm Infect** 2019;95:300–6. doi:10.1136/sextrans-2018-053778

12. Barnabas SL, Dabee S, Passmore JS, Jaspan HB, Lewis DA, Jaumdally SZ, *et al*. Converging epidemics of sexually transmitted infections and bacterial vaginosis in southern African female adolescents. **Int J STD AIDS** 2018;29:531–9. doi:10.1177/0956462417740487

13. Nugent RP, Krohn MA, Hillier SL. Reliability of diagnosing bacterial vaginosis is improved by a standardized method of gram stain interpretation. **J Clin Microbiol** 1991;29:297–301. doi:10.1128/jcm.29.2.297-301.1991

14. Lewis DA, Muller E, Steele L, Sternberg M, Radebe F, Lyall M, *et al.* Prevalence and associations of genital ulcer pathogens in men presenting with genital ulcer syndrome in South Africa. **Sex Transm Dis** 2012;39:880–5. doi:10.1097/OLQ.0b013e318269cf90

15. Bom RJ, Christerson L, Schim van der Loeff MF, Coutinho RA, Herrmann B, Bruisten SM. Evaluation of high-resolution typing methods for Chlamydia trachomatis. **J Clin Microbiol** 2011;49:2844–53. doi:10.1128/JCM.00128-11

16. Dabee S, Barnabas SL, Lennard KS, Jaumdally SZ, Gamieldien H, Balle C, *et al.* Defining genital health in South African adolescent girls at high risk for HIV. **PLoS One** 2019;14:e0213975. doi:10.1371/journal.pone.0213975

17. Jaumdally SZ, Masson L, Jones HE, Dabee S, Hoover DR, Gamieldien H, *et al.* Lower genital tract cytokine profiles in South African women living with HIV. **Sci Rep** 2018;8:12203. doi:10.1038/s41598-018-30587-1

18. Bunjun R, Lurie M, Dabee S, Barnabas S, Maseko V, Jaumdally SZ, *et al.* Chlamydia trachomatis-specific T cell immunity reflects widespread exposure in South African adolescents and young women. **J Infect Dis** 2025;:jiaf595. doi:10.1093/infdis/jiaf595

19. Wong WF, Chambers JP, Gupta R, Arulanandam BP. Chlamydia and its immune evasion strategies. J Pathol 2019;2019:8604958. doi:10.1155/2019/8604958

20. van Duynhoven YT, Ossewaarde JM, Derksen-Nawrocki RP, van der Meijden WI, van de Laar MJ.
Chlamydia trachomatis genotypes and clinical manifestations. **Clin Infect Dis** 1998;26:314–22.
doi:10.1086/516291
21. Su H, Morrison RP, Watkins NG, Caldwell HD. T helper cell epitopes of the major outer membrane
protein of Chlamydia trachomatis. **J Exp Med** 1990;172:203–12. doi:10.1084/jem.172.1.203
22. Harris SR, Clarke IN, Seth-Smith HM, Solomon AW, Cutcliffe LT, Marsh P, *et al*. Whole-genome
analysis of Chlamydia trachomatis. **Nat Genet** 2012;44:413–9. doi:10.1038/ng.2214
23. Kachale F, Mahaka I, Mhuriro F, Mugambi M, Murungu J, Ncube B, *et al*. Integration of HIV
prevention and sexual and reproductive health. **Gates Open Res** 2021;5:145.
doi:10.12688/gatesopenres.13330.2

Table 1. Characteristics for study participants

Characteristics	All women (n=298)	Johannesburg (n=149)	Cape Town Visit 1 (n=149)	Cape Town Visit 2 (n=128*)	Cape Town Visit 3 (n=93*)	p-value#
Age (years, range)	19 (18-20)	19 (17-20)	19 (18-19)	-	-	0.4
Condoms (always)	69/267 (25.8%)	49/138 (36%)	20/129 (16%)	-	-	0.0002
Condoms (last sex)	190/267 (71.2%)	94/138 (68%)	96/129 (74%)	-	-	0.3
Sex w HIV+ partner	20/267 (7.5%)	7/138 (5%)	13/129 (10%)	-	-	0.2
Transactional sex	1/267 (0.4%)	0/138 (0%)	1/129 (1%)	-	-	0.5
Contraceptive:						
None	14/298 (4.7%)	14/149 (9.4%)	0/149 (0%)	-	-	<0.0001
Condoms only	77/298 (31.9%)	77/149 (63.8%)	0/149 (0%)	-	-	<0.0001
COCP	16/298 (5.4%)	9/149 (6.0%)	7/149 (4.7%)	-	-	0.8
Implant	10/298 (3.4%)	1/149 (0.7%)	9/149 (6.0%)	-	-	0.02
Net-EN	129/298 (43.3%)	24/149 (16.1%)	105/149 (70.5%)	-	-	<0.0001
DMPA	38/298 (12.8%)	11/149 (7.4%)	27/149 (18.1%)	-	-	0.009
CCVR	1/298 (0.3%)	0/149 (0%)	1/149 (0.7%)	-	-	1.0
Ever pregnant	64/267 (24%)	30/138 (22%)	34/129 (26%)	-	-	0.4
Anal sex (ever)	11/267 (4.1%)	3/139 (2%)	8/129 (6%)	-	-	0.1
STI frequency:						
Any STI	221/298 (74.2%)	101/149 (67.8%)	120/149 (80.5%)	97/128 (75.8%)	70/93 (75.3%)	0.02
C. trachomatis	88/298 (29.5%)	26/149 (17.4%)	62/149 (41.6%)	43/127 (33.9%)	16/88 (18.2%)	<0.0001
N. gonorrhoea	24/298 (8.1%)	7/149 (4.7%)	17/149 (11.4%)	8/127 (6.3%)	7/88 (7.9%)	0.05
T. vaginalis	16/298 (5.4%)	5/149 (3.4%)	11/149 (7.4%)	10/127 (7.9%)	6/88 (6.8%)	0.2
M. genitalium	11/298 (3.7%)	5/149 (3.4%)	6/149 (4.0%)	7/127 (5.5%)	6/88 (6.8%)	1.000
Multiple STIs	30/298 (10.1%)	6/143 (4.0%)	24/149 (16.1%)	44/128 (34.4%)	23/88 (26.1%)	0.001
BV (Nugent 7-10)	135/292 (46.2%)	64/144 (44.4%)	71/148 (48.0%)	62/126 (49.2%)	39/84 (46.4%)	0.6
Any STIs+BV	111/298 (37.2%)	51/149 (34.2%)	60/149 (40.3%)	46/128 (35.9%)	31/93 (33.3%)	0.4

344   *Of the 149 AGYW enrolled in Cape Town, 128/149 returned for visit 2 and 93/149 returned for visit 3. Denominators and
345   percentages for these visits have been adjusted to reflect those lost to follow up. #P-values reflect comparison of Cape Town
346   (visit 1) to Johannesburg (single visit) prevalence; Chi-squared test. COCP: Combined oral contraceptive pill; DPMA: depot
347   medroxyprogesterone acetate; Net-EN: norethisterone enanthate; STI: sexually transmitted infection; BV: bacterial vaginosis
348

Figure legends:

Figure 1. Impact of CT infection on cervicovaginal cytokine profiles in South African AGYW at baseline (only visit 1 for Cape Town AGYW). (A) Multivariate linear regression analysis illustrating associations between cervicovaginal cytokine levels and CT infection status, adjusted for participant age, hormonal contraceptive use, other co-infections, BV, and semen exposure. Each association is represented by a β-coefficient, with error bars indicating the 95% confidence intervals. Statistically significant associations ($p \leq 0.05$ after correction for multiple comparisons) are denoted with filled symbols. (B) Concentrations of cervicovaginal cytokines associated with inflammation (IL-1α, IL-1β, IL-6, IL-8, IP-10, TNF-α, TNF-β) and protection (IFN-γ) are compared between AGYW with CT infections (n = 26; with no other STIs or BV) and those without CT (n = 62; no STIs or BV). Group comparisons were performed using Mann-Whitney U-tests with statistical significance set at $p \leq 0.05$ after adjustment for multiple comparisons. For women in Cape Town for which multiple visits were available, only the baseline visit was included (no repeated measures).

Fig 2. Investigating the impact of CT genovar on inflammatory responses in AGYW from Cape Town and Johannesburg at baseline. (A) Multivariate linear regression analysis comparing cervicovaginal cytokine levels associated with CT infection in AGYW from Cape Town (left panel) and Johannesburg (right panel), adjusted for participant age, hormonal contraceptive use, other co-infections, BV, and semen exposure. Associations are represented by β-coefficients, with error bars indicating 95% confidence intervals. Statistically significant associations ($p \leq 0.05$ after correction for multiple comparisons) are marked with filled symbols. (B) Cervicovaginal cytokine concentrations associated with inflammation (IL-1α, IL-1β, IL-6, IL-8, IP-10, TNF-α, TNF-β) and protection (IFN-γ) are compared between AGYW with CT infections in Cape Town (CPT) and Johannesburg (JHB) (n = 26; with no other STIs or BV). Group comparisons were conducted using Mann-Whitney U-tests, with significance defined as $p \leq 0.05$ after adjustment for multiple comparisons. For women in Cape Town for which multiple visits were available, only the baseline visit was included (no repeated measures). (C) Distribution (n, percentage) of CT ompA genotypes in AGYW from Johannesburg (left pie chart) and Cape Town (right pie chart). Genovar D is represented in blue, E in green, F in yellow, G in orange, I in bright red, J in dark red, K in purple, and unknown genotypes in white. (D) Multivariate linear regression analysis of associations between cervicovaginal cytokine levels and CT ompA genovars (D to K), adjusted for participant age, hormonal contraceptive use, other co-infections, BV, and semen exposure. Associations are

represented by β-coefficients with 95% confidence intervals. None of the associations were Statistically

significant (p ≤ 0.05) after correction for multiple comparisons.

Figure 3. Cervicovaginal cytokine concentrations in AGYW infected with CT over one, two, or three study

visits. Each dot represents the cytokine response of an individual participant, with black lines indicating the

median values for each group. Group comparisons were conducted using Mann-Whitney U-tests, with statistical

significance defined as p ≤ 0.05.

Figure 4. Impact of prevalent or persistent CT Infections on cervical cytobrush-derived CD4+ T cell activation.

(A) Representative gating strategy for flow cytometry analysis of cervical cytobrush-derived T cells, focusing

on live CD3+CD4+ cells expressing activation marker CD38+, proliferation marker Ki67+, or HIV co-receptor

CCR5. (B) The frequency (top panel) and absolute number of events (bottom panel) are shown for total

CD3+CD4+ T cells (far left), CD3+CD4+ T cells expressing CD38 (middle left), CCR5 (middle right), and

Ki67 (far right). Data were collected at visit 1 (study entry) from N = 36 controls and N = 18 CT cases. (C)

Longitudinal analysis of cervicovaginal CD3+CD4+ T cell activation marker expression is presented for

AGYW with CT infections spanning one, two, or three study visits (top panel). Matched analyses for

participants with repeat CT infections across multiple visits are shown in the lower panel. Statistical

comparisons were performed using Mann-Whitney U-tests for unpaired analyses (B) and Wilcoxon Ranks test

for paired data (C), with significance defined as p ≤ 0.05.

Figure 1A and 1B

A

B

C