Conservative surgery vs hysterectomy for placenta accreta spectrum disorders: a

systematic review and meta-analysis

Running title: Conservative surgery for placenta accreta spectrum disorders: a

systematic review

Gabriel Moreira Lino[1], Pauliana Valéria Machado Galvão[2], Valda Lúcia Moreira

Luna[1], George Alessandro Maranhão Conrado[1]

[1]Universidade de Pernambuco, Serra Talhada, Brazil

[2]Faculdade de Ciências Médicas, Universidade de Pernambuco, Recife, Brazil

Corresponding author: **Pauliana Valéria Machado Galvão**

Address: R. Arnóbio Marquês, 310 - Santo Amaro, Recife - PE, 50100-130

Telephone number: +55 81 996604662

Email address: pauliana.galvao@upe.br

DECLARATIONS

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and materials

The data can be shared upon request

Competing interests

The authors declare that they have no competing interests

Funding

No specific funding was available for the project

Acknowledgments

Not applicable.

ABSTRACT

Objective

Resection and reconstruction of the uterus may be an alternative to hysterectomy

in placenta accreta spectrum disorders. We aimed to review whether surgical and maternal

outcomes were comparable between these techniques.

Data sources

Medline, Embase, and the Cochrane Library databases were searched until 1

January 2025.

Study selection

We included randomized controlled trials (RCT) and nonrandomized studies

(NRSI) comparing conservative surgery, involving resection and reconstruction of the

myometrium without placental removal, to hysterectomy in patients with placenta

accreta, increta, or percreta.

Data collection

Two authors were responsible for independently screening the title and abstract,

and the full texts of all identified records. We assessed the risk of bias with ROB-2 or

ROBINS-I, and the quality of evidence with GRADE. The main analysis only included

studies without a high risk of bias. A secondary non-inferiority Bayesian analysis was

reported alongside the results.

Data synthesis

We included one RCT and eleven nonrandomized studies (1586 women).

Compared to hysterectomy, conservative surgery may result in a similar number of

transfused packs of red blood cells (MD -0.70 packs, 95% CI -1.28 to -0.12; 1 RCT and

4 NRSI, 649 women; $I^2 = 0\%$; low-certainty evidence), and a similar risk of bladder injury

(RR 0.36, 95% CI 0.12 to 1.07; 1 RCT and 4 NRSI, 649 women; $I^2 = 0\%$; low-certainty

evidence). Excluding rare outcomes, the probability that conservative surgery is non-

inferior to hysterectomy was greater than 90.0%.

Conclusions

On the basis of low evidence, conservative surgery did not increase the risk of

clinically important harm compared to hysterectomy, with the benefit of uterus

preservation. Further RCTs are needed to reach definitive conclusions.

Keywords: Placenta accreta. Surgery. Fertility. One-step surgery. Triple-P procedure.

Conservative management.

Registration: PROSPERO CRD42024601117.

INTRODUCTION

Until recently, hysterectomy was virtually the only management option for

placenta accreta spectrum disorder (PAS). However, as the prevalence of PAS has

increased to affect approximately 1 in every 313 women undergoing cesarean section, the

growing number of women wishing to preserve fertility has prompted investigation into

alternative strategies for uterine preservation[1]. Currently, leaving the placenta in situ is

one of the most utilized alternatives to hysterectomy, although it requires high-resource

settings and continuous long-term monitoring[2,3]. In this context, uterus-preserving

surgery (also referred to as one-step surgery and triple P procedure, according to

variations in the technique proposed by individual authors) has been developed to provide

immediate surgical resolution of PAS during cesarean delivery[4].

This technique involves removing the invaded myometrium and reconstructing

the uterine wall, with or without preventive surgical devascularization[2,4]. However, it

remains to be proven whether surgical outcomes are superior or similar to hysterectomy,

specifically because of the challenging surgical anatomy and the inherent risk of

intrapartum hemorrhage and visceral organ injury. Therefore, this review aims to

summarize the available evidence on this conservative surgical technique, comparing it

to hysterectomy in patients diagnosed with PAS who wish to preserve fertility.

METHODS

Protocol and registration

This systematic review was registered prospectively in the PROSPERO database,

identifier CRD42024601117. We followed the PRISMA 2020 guidelines for the reporting

of systematic reviews and meta-analyses[5].

Criteria for considering studies for this review

We considered randomized controlled trials (RCT) and nonrandomized studies of

interventions comparing conservative surgery to hysterectomy in patients with placenta

accreta, increta, or percreta. To be included, the surgical technique must have involved

resection of abnormally invasive placenta and myometrium and reconstruction of the

uterine wall. Therefore, we excluded studies that provided fertility preservation by

manual removal of the placenta followed by vascular control procedures, such as uterine

compressive sutures, tourniquets, and uterine devascularization.

The following critical outcomes were considered: intraoperative blood loss,

number of transfused packs of red blood cells or women needing transfusion,

intraoperative visceral organ injury (bladder injury, bowel injury, ureter injury, and

vascular injury), reoperation, long-term complications of surgery (fistula, premature

ovarian insufficiency, urinary dysfunction, dysmenorrhea). The secondary outcomes of

the review were: duration of surgery, duration of hospital stay, admission to the intensive

care unit, and postoperative infection. Studies had to report at least one of the primary

outcomes to be eligible for the review.

Search methods for identification of studies

We searched for published studies up to January 2025, on Medline, Embase, and

the Cochrane Library using the terms: (one step conservative surgery OR uterine

conservative-resective surgery OR uterus resection OR uterine resective surgery OR

uterine conservative surgery OR uterus conservation surgery OR uterine preserving

surgery OR fertility sparing surgery OR fertility preserving surgery OR triple p procedure

OR triple p) and (placenta accreta spectrum OR placenta increta OR placenta percreta OR

abnormally invasive placenta OR morbidly adherent placenta). There were no language

restrictions.

Data collection and analysis

Two authors were responsible for independently screening the title and abstract

of all identified records. When one of the authors considered the record eligible, both

assessed the full text. Thereafter, they proceeded with selecting and extracting

information about study characteristics, which were summarized on a cross-validated file.

Risk of bias assessment in included studies

For the risk of bias assessment, two authors independently used the Cochrane risk-

of-bias tool for randomized trials (RoB 2), version of August 2019[6]. The ROBINS-I V2

tool, 2024, was used instead for nonrandomized studies of interventions[7]. We evaluated

the overall risk of bias in each study based on three primary outcomes (blood loss,

transfusion, and intraoperative visceral organ injury). Dropouts, group conversions, or

loss to follow-up below 5% were judged as low risk of bias.

Synthesis methods

We used an inverse-variance random effects model, reporting dichotomous data

as Relative Risk (RR) with 95% Confidence intervals (CI) and continuous outcomes as

mean differences (MD) and 95% CI. Statistical heterogeneity was assessed by calculating

tau (DerSimonian-Laird estimator and Jackson method for confidence interval of tau^2),

I^2, and Cochran's Q; visual I^2 variation that included clinically important differences

indicated serious heterogeneity[8]. We investigated publication bias with funnel plots and

Egger's test.

Studies that reported medians and interquartile ranges were transformed using an

internal function of the meta package, following what was described by Luo and

collaborators[9]. We stratified our analyses based on the risk of bias. Primary results in

every section only included studies with randomized design or that were not at high risk

of bias. Subsequent analyses reported the estimated effect including the remaining

studies, except when the results of both analyses were uncertain, making it unnecessary

to report them separately.

Analyses were conducted in R, with the packages meta and metafor[10,11].

Sensitivity analyses

To explore the certainty of the main results, we conducted a planned secondary

analysis using a Bayesian framework. Specifically, we assessed how integrating prior

knowledge from observational studies not at serious or critical risk of bias would

influence the interpretation of the randomized studies, which had limited statistical power

due to modest sample sizes.

We first synthesized the selected nonrandomized studies using a Bayesian meta-

analysis with a non-informative prior (mean = 0, SD = 10,000), generating a posterior

distribution that served as an informative prior for the subsequent analysis of the trial

data. We then combined this prior with the likelihood derived from the randomized

controlled trial to obtain a posterior estimate reflecting both sources of evidence.

This reanalysis focused on critical clinical outcomes: number of packs of red

blood cells transfused, intraoperative visceral injuries (including bladder and ureter), need

for reoperation, and long-term surgical complications.

Certainty of the evidence assessment

The overall certainty of the evidence for each outcome was assessed by two

authors using the Grading of Recommendations, Assessment, Development and

Evaluation (GRADE) handbook[12]. The quality of the evidence was summarized in a

Summary of Findings table generated in GRADEpro GDT. For the evaluation of

imprecision, the update provided in 2022 was followed[13]. The Review Information Size

was determined using conventional power analysis ($\alpha = 0.05$; $\beta = 0.2$), taking into account

the range of standard deviations from randomized controlled trials and medium to low

risk of bias studies.

RESULTS

Results of the search

We initially identified 640 records in our search. Of these, 106 were deemed

potentially eligible, and 12 (only one RCT) were ultimately included in the review. The

primary reasons for exclusion were inadequate surgical techniques (manual placental

removal or failure to report surgical methods), study design (such as case series, reports,

reviews, editorials, and letters), absence of relevant outcome data specified in the review

protocol, and not comparing conservative surgery to hysterectomy (such as comparing to

manual removal of the placenta).

Included studies

Table 1 summarizes the characteristics of the included studies. One RCT and

eleven non-randomized studies were included, four of which had prospective designs.

Moreover, some studies reported the PAS FIGO clinical grading[14], with patients of every

grade in both arms. The average maternal age ranged from 30 to 37 years, with most

women having had one or two previous cesarean sections.

Preventive vascular interventions before myometrial resection were utilized in

three of the included studies. Further descriptions of methods, eligibility criteria, and

participants are available in Supplementary Table S1. In general, the most common

eligibility criteria for uterine conservative surgery were: (1) at least 2 cm of healthy

myometrium in the anterior uterine wall, (2) more than 50% of the uterine circumference

without PAS involvement, and (3) the ability to safely dissect the bladder from the uterus.

Most studies followed the surgical technique described as one-step surgery[4].

Table 1 – Summary of characteristics of included studies

Author (Year)	Country	Study design	Sample size (CS;HT)	PAS FIGO grading (n)		Maternal age (y)		Previous cesarean deliveries (n)		Preventive vascular intervention	Risk of bias
				CS	HT	CS	HT	CS	HT		
Nieto-Calvache 2024[15]	Multicenter	RCT, not blinded	60 (29;31)	Grade 1 2 Grade 2 18 Grade 3A 3 Grade 3B-C 4	1 20 0 3	33.45 ±4.56	33.13 ±4.26	-	-	No	Moderate risk
Nieto-Calvache 2023[16]	Multicenter	NRSI, prospective design	75 (64;11)	-	-	33 (26–38)	31 (29–37)	-	-	No	Moderate risk
Paping 2024[17]	Multicenter	NRSI, retrospective design, prospectively-collected data	224 (38;186)	Grade 1 10 Grade 2 13 Grade 3A 14 Grade 3B-C 1	21 46 60 59	34 (30–37)	35 (31–39)	2 (1–3)	2 (1–2)	No	Moderate risk

Study	Center	Design	N (subgroups)	Grade		Age 1	Age 2	Col1	Col2		Risk
Abouda 2024[18]	Single center	NRSI, prospective design	12 (6;6)	-	-	37.17±2.93	33.67±3.07	4±0.89	2.83±0.98	No	Moderate risk
van Beekhuizen 2021[19]	Multicenter	NRSI, retrospective design, prospectively collected data	278 (26;252)	Grade 1 3 Grade 2 2 Grade 3A 18 Grade 3B-C 3	31 52 102 66	-	-	-	-	No	Moderate risk
Vuong 2024[20]	Single center	NRSI, retrospective design	296 (217;79)			33.17 ± 4.77	33.68 ± 4.94	-	-	No	High risk
Aryananda 2023[21]	Single center	NRSI, retrospective design	126 (91;35)	Grade 1 16 Grade 2 54 Grade 3A 15 Grade 3B-C 6	0 7 10 18	33 (22–46)	33 (22–42)	2 (1–3)	2 (1–3)	No	High risk

Aryananda 2022[22]	Single center	NRSI, retrospective design	234 (200;34)	-	-	33 (19–44)	35 (26–41)			Yes: internal iliac artery ligation	High risk
Jafari 2022[23]	Single center	NRSI, retrospective design	68 (44;24)	-	-	33.23 ± 4.72	32.54 ± 5.69	-	-	No	High risk
Durukan 2021[24]	Single center	NRSI, retrospective design	65 (23;42)	-	-	33.1 ± 5.4	34.1 ± 5.4	2.0 ± 1.1	2.1 ± 0.9	No	High risk
Üstünyurt 2020[25]	Single center	NRSI, retrospective design	97 (30;67)	-	-	31.1± 5.4	33.2±4.7	-	-	Yes: Intrauterine balloon tamponade	High risk
Kilicci 2018[26]	Single center	NRSI, retrospective design	51 (22;29)	-	-	32.3 ±5.3	32.45± 4.7	1.6 ±0.9	1.8 ±0.6	No	High risk

227   RCT: randomized controlled trial; NRSI: nonrandomized study of intervention.

Risk of bias of included studies

An overview of the risk of bias is provided in Table 1, and the detailed description

is presented in Table S2. The included RCT was classified as having a moderate risk of

bias due to the lack of blinding and important patient crossover/drop-out. Nearly all non-

randomized studies were considered at high risk of bias, as patient selection was often at

serious risk of bias. Additionally, few studies reported potential confounding variables or

adjusted their analyses as necessary.

An investigation into publication bias revealed that retrospective studies, which

were at high risk of bias, generally reported higher mean differences with larger standard

errors compared to studies with moderate to low risk of bias (Figure S1). However,

Egger's Test did not detect significant publication bias affecting the analysis (p-value =

0.20).

Synthesis of results

1.1 Intraoperative blood loss

Conservative surgery may result in similar intraoperative blood loss compared to

hysterectomy (MD -320.89 mL, 95% CI -574.95 to -66.82; 1 RCT and 3 NRSI, 637

women; I^2= 14%; low-certainty evidence). By including studies at high risk of bias, the

difference in blood loss is larger, but of lower certainty (MD -497.33 mL, 95% CI -755.37

to -226.17; 1 RCT and 7 NRSI; I^2= 73%; very low-certainty evidence)(Figure 2).

1.2 Number of transfused packs of red blood cells (pRBCs) and number of women needing

transfusion

The number of transfused pRBCs per patient may be similar in conservative

surgery compared to hysterectomy (MD -0.70 packs, 95% CI -1.28 to -0.12; 1 RCT and

4 NRSI, 649 women; I^2= 0%; low-certainty evidence). This benefit could be slightly

larger (MD -1.15 packs, 95% CI -1.64 to -0.65; 1 RCT and 9 NRSI; I^2= 20%; very low-

certainty evidence)(Figure 3). Further, assuming a non-inferiority margin of 0.5 pRBCs,

the probability that conservative surgery is non-inferior to hysterectomy is 95.4%

(Figures S2 and S3).

Moreover, there may be a similar number of patients needing blood transfusion

(RR 0.87, 95% CI 0.67 to 1.13; 1 RCT and 1 NRSI, 135 women; I^2= 0%; low-certainty

evidence) or fewer women will require transfusion (RR 0.82, 95% CI 0.74 to 0.91; 1 RCT

and 4 NRSI; I^2= 0%; very low-certainty evidence).

1.3 Intraoperative visceral organ injury

Few cases of bladder injury were reported, therefore, we found similar event rates

for both groups (RR 0.36, 95% CI 0.12 to 1.07; 1 RCT and 4 NRSI, 649 women; I^2= 0%;

low-certainty evidence). Potentially, the risk of bladder injury could be lower with

conservative surgery (RR 0.32, 95% CI 0.19 to 0.54; 1 RCT and 6 NRSI; I^2= 0%; very

low-certainty evidence)(Figure 4). Due to the low event rate, the non-inferiority analysis

was uninformative (Figures S4 and S5).

In the two studies that investigated ureter injuries, no cases were observed in either

arm (2 NRSI, 87 women; very low-certainty evidence). Likewise, no cases of bowel

injury were reported with conservative surgery, and seven were reported with

hysterectomy (RR 0.17, 95% CI 0.02 to 1.65; 3 NRSI; I^2= 17%; very low-certainty

evidence).

1.4 Reoperation

Very few cases of reoperation were reported in both groups, hence, the pooled

estimate was very wide (RR 0.50, 95% CI 0.07 to 3.51; 1 RCT and 3 NRSI; I^2= 0%; very

low-certainty evidence)(Figure S6). The non-inferiority analysis was uninformative due

to low event rates (Figures S7 and S8).

1.5 Long-term complications of surgery

Vesicovaginal fistula formation was the only long-term complication reported

across studies. As previously observed, this outcome was rare in both groups, leading to

a wide effect estimate (RR 1.40, 95% CI 0.16 to 12.08; 1 RCT and 1 NRSI, 294 women;

I^2 = 0%; very low-certainty evidence)(Figure S9). The non-inferiority analysis was

uninformative due to low event rates (Figures S10 and S11).

1.6 Duration of surgery

There was little to no difference in the duration of surgery (MD -36.31 minutes,

95% CI -73.06 to 0.43; 1 RCT and 2 NRSI, 147 women; I^2= 54%; low-certainty

evidence). Very uncertainly, conservative surgery could be shorter, however, high

heterogeneity limits the confidence in this possibility (MD -26.45 minutes, 95% CI -45.35

to -7.56; 1 RCT and 6 NRSI; I^2= 89%; very low-certainty evidence)(Figure 5).

1.7 Duration of hospital stay

There was little to no difference in the duration of hospital stay (MD -0.39 days,

95% CI -0.87 to 0.08; 1 RCT and 4 NRSI; I^2=33%; very low-certainty evidence). This

could depend on institutional protocols and may not relate to the surgeries investigated.

1.7 Admission to the intensive care unit (ICU)

There was little to no difference in the risk of ICU admission (RR 1.11, 95% CI

0.83 to 1.47; 1 RCT and 4 NRSI; I^2= 0%; very low-certainty evidence).

Certainty of evidence

The certainty of the evidence ranged from low to very low, according to the

GRADE assessment (Table 2). Whenever the analysis included nonrandomized studies,

the evidence was limited to low. To overcome these limitations on evidence, the Review

Information Size (RIS) necessary to not downgrade for imprecision, for equally divided

randomized controlled trials, would be: 166 to 594 participants for pRBC, and 210 to

4908 participants for blood loss, assuming moderate to small effects for the outcomes.

Conversely, more rare outcomes (visceral organ injury and ICU admission) would

need far greater sample sizes, which might not be feasible.

DISCUSSION

Summary of main results

Conservative surgery has the advantage of preserving the uterus with surgical

outcomes that were similar to hysterectomy in our analyses, in light of low-certainty

evidence. Visceral organ injury, reoperation, and long-term complications were fairly

rare, while blood loss was similar. In this regard, there is a high probability that

conservative surgery is non-inferior to hysterectomy. However, patients had to have a

specific PAS topography to be eligible for conservative surgery, with generally at least 2

cm of healthy myometrium on the anterior uterine wall, >50% of the uterine

circumference free of PAS, without vesicouterine fibrosis, or extensive invasion of the

lower anterior uterine wall.

Limitations of the evidence included in the review

The most important limitation of our study is the inclusion of nonrandomized

studies of interventions. Major limitations that future trials on PAS would face are the

low incidence of PAS, which requires longer recruitment and results in smaller samples,

the long-term follow-up needed to determine patient-centered outcomes, and the specific

placenta topography that limits the number of eligible women to be recruited.

Fundamentally, nonrandomized studies are better at evaluating potential harm than

potential benefit[27].

Inherent consequences of results drawn from these studies are uncontrolled

confounding and suboptimal outcome reporting. Even after excluding high-risk studies

from the main analysis, the confidence in the estimates provided was low. If only the data

of the randomized trial were analyzed, the small sample size and wide confidence

intervals would make it unreasonable to assume any given effect for conservative surgery.

Additionally, not all patients can be considered for conservative surgery, as there

are topographical constraints to this technique. Another limitation was the lack of patient-

driven outcomes in the included studies. However, it could be argued that uterus

preservation (with reasonable safety) would be the most prominent outcome for patients.

Agreements and disagreements with other studies or reviews

The success rate of conservative surgery in preserving fertility was reviewed

elsewhere, and hence, was not one of our outcomes of interest[28]. Notably, subsequent

pregnancy is possible, however, adverse maternal outcomes or recurrence of PAS was

reported in nearly 20% of patients, and hysterectomy was necessary for around 2% of all

women[28,29]. Such findings should be interpreted in the context of very low-certainty

evidence due to the limitations of these studies.

Our results add substantially to the initial assessment from FIGO in 2018, in which

it was suggested that leaving the placenta in situ was a valid management option for

women desiring to preserve fertility while conservative surgery remained unclear[2]. Here

we present both randomized and prospective nonrandomized data that suggest the safety

of conservative surgery in some outcomes. In theory, considering both nonsurgical and

surgical conservative management for PAS would be ideal for women desiring to

preserve fertility, as these approaches have different shortcomings and eligibility criteria.

Moreover, there are many variations of conservative surgery, which include

prophylactic balloon occlusion or major vascular intervention (such as the triple P

procedure)[30]. However, many of these studies did not compare their interventions to

hysterectomy and, therefore, were excluded from our review. The studies that were

eventually included were not randomized, and almost all had a high risk of bias, making

it impossible to derive any conclusions about prophylactic vascular interventions. In

previous investigations, meta-analyses and randomized controlled trials reported

conflicting results for prophylactic balloon occlusion in women undergoing hysterectomy

for PAS[31,32].

In terms of intraoperative safety, the management of PAS requires careful

planning due to the complexity of the procedures and the high risk of visceral organ

injuries. Urological complications occur in approximately 20% of hysterectomies

performed for PAS[33], and massive blood loss remains both common and difficult to

predict[34]. In our analysis, injury and transfusion rates appeared non-inferior between

groups; however, strategies to effectively mitigate these intraoperative challenges remain

elusive.

Moreover, in the non-inferiority analysis, we found a high probability that

conservative surgery is non-inferior to hysterectomy in terms of blood loss. However, we

could not make the same claim for other critical outcomes, such as intraoperative visceral

organ injury and long-term complications. This limitation is largely attributable to the

low prevalence of these events, which resulted in underpowered analyses. Nonetheless, it

is unlikely that a potential increase in harm associated with conservative surgery would

be detected.

Implications for practice

Conservative surgery in women diagnosed with placenta accreta spectrum

disorders, and with sufficiently healthy lower anterior uterine wall and uterine

circumference, may not lead to higher blood loss, number of packs of red blood cells

transfused, intraoperative visceral organ injury, and duration of surgery compared to

hysterectomy. Therefore, uterus preservation is possible without the risk of substantial

harm in critical outcomes likely increasing. This surgical procedure is not possible in all

women diagnosed with PAS, especially in the presence of vesicouterine fibrosis.

Implications for research

Further research should seek to increase the total sample size of randomized trials

comparing conservative surgery to hysterectomy in order to minimize imprecision on

confidence interval boundaries and review information size. Additionally, it would be

ideal to determine whether there are specific variations of conservative surgery, especially

involving prophylactic balloon occlusion, that result in better surgical outcomes. At

present, our results suggest that conservative management does not result in substantially

increased harm to women compared to hysterectomy, but more precise estimates would

allow the determination of possible benefits.

REFERENCES

1.   Matsuzaki S, Mandelbaum RS, Sangara RN, et al. Trends, characteristics, and outcomes of

placenta accreta spectrum: a national study in the United States. *Am J Obstet Gynecol*.

2021;225(5):534.e1-534.e38. doi:10.1016/j.ajog.2021.04.233

2.   Sentilhes L, Kayem G, Chandraharan E, Palacios-Jaraquemada J, Jauniaux E, FIGO

Placenta Accreta Diagnosis and Management Expert Consensus Panel. FIGO consensus

guidelines on placenta accreta spectrum disorders: Conservative management. *Int J*

Gynaecol Obstet Off Organ Int Fed Gynaecol Obstet. 2018;140(3):291-298.

doi:10.1002/ijgo.12410

3.   Sentilhes L, Seco A, Azria E, et al. Conservative management or cesarean hysterectomy for

placenta accreta spectrum: the PACCRETA prospective study. *Am J Obstet Gynecol*.

2022;226(6):839.e1-839.e24. doi:10.1016/j.ajog.2021.12.013

4.   Palacios Jaraquemada JM, Pesaresi M, Nassif JC, Hermosid S. Anterior placenta percreta:

surgical approach, hemostasis and uterine repair. *Acta Obstet Gynecol Scand*.

2004;83(8):738-744. doi:10.1111/j.0001-6349.2004.00517.x

5.   Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: An updated

guideline for reporting systematic reviews. *PLOS Med*. 2021;18(3):e1003583.

doi:10.1371/journal.pmed.1003583

6.   Sterne JAC, Savović J, Page MJ, et al. RoB 2: a revised tool for assessing risk of bias in

randomised trials. *BMJ*. 2019;366:l4898. doi:10.1136/bmj.l4898

7.   Sterne JA, Hernán MA, Reeves BC, et al. ROBINS-I: a tool for assessing risk of bias in

non-randomised studies of interventions. *BMJ*. 2016;355:i4919. doi:10.1136/bmj.i4919

8.   Guyatt G, Zhao Y, Mayer M, et al. GRADE guidance 36: updates to GRADE's approach to

addressing inconsistency. *J Clin Epidemiol*. 2023;158:70-83.

doi:10.1016/j.jclinepi.2023.03.003

9. Luo D, Wan X, Liu J, Tong T. Optimally estimating the sample mean from the sample size,

median, mid-range, and/or mid-quartile range. *Stat Methods Med Res*. 2018;27(6):1785-

1805. doi:10.1177/0962280216669183

10. Viechtbauer W. Conducting meta-analyses in {R} with the {metafor} package. *J Stat Softw*.

2010;36(3).

11. Balduzzi S, Rücker G, Schwarzer G. How to perform a meta-analysis with R: a practical

tutorial. *Evid Based Ment Health*. 2019;22(4):153-160. doi:10.1136/ebmental-2019-300117

12. Schünemann H, Brożek J, Guyatt G, Oxman A, editor(s). Handbook for grading the quality

of evidence and the strength of recommendations using the GRADE approach (updated

October 2013). Published online 2013.

https://gdt.gradepro.org/app/handbook/handbook.html

13. Zeng L, Brignardello-Petersen R, Hultcrantz M, et al. GRADE Guidance 34: update on

rating imprecision using a minimally contextualized approach. *J Clin Epidemiol*.

2022;150:216-224. doi:10.1016/j.jclinepi.2022.07.014

14. Jauniaux E, Ayres-de-Campos D, Langhoff-Roos J, Fox KA, Collins S, FIGO Placenta

Accreta Diagnosis and Management Expert Consensus Panel. FIGO classification for the

clinical diagnosis of placenta accreta spectrum disorders. *Int J Gynaecol Obstet Off Organ*

Int Fed Gynaecol Obstet. 2019;146(1):20-24. doi:10.1002/ijgo.12761

15. Nieto-Calvache ÁJ, Aryananda RA, Palacios-Jaraquemada JM, et al. One-step conservative

surgery vs hysterectomy for placenta accreta spectrum: a feasibility randomized controlled

trial. *Am J Obstet Gynecol MFM*. 2024;6(6):101333. doi:10.1016/j.ajogmf.2024.101333

16. Nieto-Calvache AJ, Palacios-Jaraquemada JM, Aryananda R, et al. How to perform the

one-step conservative surgery for placenta accreta spectrum move by move. *Am J Obstet*

Gynecol MFM. 2023;5(2):100802. doi:10.1016/j.ajogmf.2022.100802

17. Paping A, Bluth A, Al Naimi A, et al. Opportunities for, and barriers to, uterus-preserving

surgical techniques for placenta accreta spectrum. *Acta Obstet Gynecol Scand*. Published

online May 2, 2024. doi:10.1111/aogs.14855

18. Abouda HS, Aloui H, Azouz E, et al. New surgical technique for managing placenta accreta

spectrum and pilot study of the "CMNT PAS" study. *AJOG Glob Rep*. 2025;5(1):100430.

doi:10.1016/j.xagr.2024.100430

19. van Beekhuizen HJ, Stefanovic V, Schwickert A, et al. A multicenter observational survey

of management strategies in 442 pregnancies with suspected placenta accreta spectrum.

Acta Obstet Gynecol Scand. 2021;100 Suppl 1(Suppl 1):12-20. doi:10.1111/aogs.14096

20. Vuong ADB, Pham TH, Pham XTT, et al. Modified one-step conservative uterine surgery

(MOSCUS) versus cesarean hysterectomy in the management of placenta accreta spectrum:

A single-center retrospective analysis based on 619 Vietnamese pregnant women. *Int J*

Gynaecol Obstet Off Organ Int Fed Gynaecol Obstet. 2024;165(2):723-736.

doi:10.1002/ijgo.15220

21. Aryananda RA, Duvekot JJ, Van Beekhuizen HJ, Cininta NI, Ariani G, Dachlan EG.

Transabdominal and transvaginal ultrasound findings help to guide the clinical management

of placenta accreta spectrum cases. *Acta Obstet Gynecol Scand*. 2024;103(1):93-102.

doi:10.1111/aogs.14715

22. Aryananda RA, Aditiawarman A, Gumilar KE, et al. Uterine conservative-resective surgery

for selected placenta accreta spectrum cases: Surgical-vascular control methods. *Acta*

Obstet Gynecol Scand. 2022;101(6):639-648. doi:10.1111/aogs.14348

23. Mohammad Jafari R, Najafian M, Barati M, Saadati N, Jalili Z, Poolad A. Comparison of

uterine preservation versus hysterectomy in women with placenta accreta: A cross-sectional

study. *Int J Reprod Biomed*. 2022;20(9):739-744. doi:10.18502/ijrm.v20i9.12063

24. Durukan H, Durukan ÖB, Yazıcı FG. Placenta accreta spectrum disorder: a comparison

between fertility-sparing techniques and hysterectomy. *J Obstet Gynaecol J Inst Obstet*

Gynaecol. 2021;41(3):353-359. doi:10.1080/01443615.2020.1755629

25. Üstünyurt E. Local uterine resection with Bakri balloon placement in placenta accreta

spectrum disorders. *Turk J Obstet Gynecol*. 2020;17(2):108-114.

doi:10.4274/tjod.galenos.2020.82652

26. Kilicci C, Ozkaya E, Eser A, et al. Planned cesarean hysterectomy versus modified form of

segmental resection in patients with placenta percreta. *J Matern-Fetal Neonatal Med Off J*

Eur Assoc Perinat Med Fed Asia Ocean Perinat Soc Int Soc Perinat Obstet.

2018;31(22):2935-2940. doi:10.1080/14767058.2017.1359535

27. Reeves B, Deeks J, Higgins J, Shea B, Tugwell P, Wells G. Chapter 24: Including non-

randomized studies on intervention effects. Cochrane Handbook for Systematic Reviews of

Interventions version 6.4 (updated August 2023). Published online 2023. Accessed March

2, 2025. https://training.cochrane.org/handbook/current/chapter-24

28. Javinani A, Qaderi S, Hessami K, et al. Delivery outcomes in the subsequent pregnancy

following the conservative management of placenta accreta spectrum disorder: a systematic

review and meta-analysis. *Am J Obstet Gynecol*. 2024;230(5):485-492.e7.

doi:10.1016/j.ajog.2023.10.047

29. Zhao H, Liu C, Fu H, Abeykoon SDI, Zhao X. Subsequent pregnancy outcomes and risk

factors following conservative treatment for placenta accreta spectrum: a retrospective

cohort study. *Am J Obstet Gynecol MFM*. 2023;5(12):101189.

doi:10.1016/j.ajogmf.2023.101189

30. Piñas-Carrillo A, Chandraharan E. Conservative surgical approach: The Triple P procedure.

Best Pract Res Clin Obstet Gynaecol. 2021;72:67-74. doi:10.1016/j.bpobgyn.2020.07.009

31. Bonsen LR, Sleijpen K, Hendriks J, et al. Prophylactic Radiologic Interventions for

Postpartum Hemorrhage Control in Women With Placenta Accreta Spectrum Disorder.

Obstet Gynecol. 2024;144(3):315-327. doi:10.1097/AOG.0000000000005662

32. Yu SCH, Cheng YKY, Tse WT, et al. Perioperative prophylactic internal iliac artery

balloon occlusion in the prevention of postpartum hemorrhage in placenta previa: a

randomized controlled trial. *Am J Obstet Gynecol*. 2020;223(1):117.e1-117.e13.

doi:10.1016/j.ajog.2020.01.024

33. Lucidi A, Jauniaux E, Hussein AM, et al. Urological complications in women undergoing

Cesarean section for placenta accreta spectrum disorders: systematic review and meta-

analysis. *Ultrasound Obstet Gynecol*. 2023;62(5):633-643. doi:10.1002/uog.26299

34. Wright JD, Pri-Paz S, Herzog TJ, et al. Predictors of massive blood loss in women with

placenta accreta. *Am J Obstet Gynecol*. 2011;205(1):38.e1-6.

doi:10.1016/j.ajog.2011.01.040

Figure 1 PRISMSA 2020 flow diagram of study selection

Figure 2 Comparison of uterine conservative surgery vs hysterectomy on intraoperative

blood loss in women with placenta accreta spectrum disorders

Figure 3 Comparison of uterine conservative surgery to hysterectomy on the number of

transfused packs of red blood cells per patient in women with placenta accreta spectrum

disorders. RCT, randomized controlled trial; IV, inverse variance; NRSI,

nonrandomized study of intervention.

Figure 4 Comparison of uterine conservative surgery to hysterectomy on bladder injury

in women with placenta accreta spectrum disorders. RCT, randomized controlled trial;

IV, inverse variance; NRSI, nonrandomized study of intervention.

Figure 5 Comparison of uterine conservative surgery to hysterectomy on the duration of

surgery in women with placenta accreta spectrum disorders. RCT, randomized

controlled trial; IV, inverse variance; NRSI, nonrandomized study of intervention.