Estimating the impact of decreasing vaccination response times for outbreaks of vaccine-preventable diseases in low and middle-income countries

Dominic Delport, MSc[1,2] Alina M. Muellenmeister, MSc[1]; Jane Greig, PhD[1]; Romesh G. Abeysuriya, PhD[1,2,*]; Nick Scott, PhD[1,2,*]

[1] Burnet Institute, Melbourne, VIC 3004, Australia

[2] Department of Epidemiology and Preventive Medicine, Monash University, Melbourne, VIC 3004, Australia

[*] Equal senior author

Corresponding author: Dominic Delport, dominic.delport@burnet.edu.au, 85 Commercial Rd, Melbourne, VIC 3004, Australia

Abstract

Background

Globally, infectious disease outbreaks cause a major health and economic burden. For infectious diseases which have existing vaccines, reactive vaccination programs during an outbreak are a powerful tool for reducing disease transmission — preventing infections and deaths. However, despite awareness, monitoring, and accessible stockpiles of vaccines, it can take months for vaccines to be delivered once outbreaks are detected in low-resource settings.

The 7-1-7 targets are gaining traction as measurable targets for assessing a country's outbreak readiness. The targets are outbreak detection within seven days of emergence; notification to health authorities within one day; and key early response actions commenced within another seven days. For outbreaks of measles, cholera, yellow fever, and meningococcal meningitis, we aim to estimate the impact of initiating outbreak response immunisation (ORI) within 15 days of outbreak emergence, relative to the mean ORI response time for each disease in low and middle-income countries (LMICs) since 2000. While only one component of outbreak response, initiating ORI within 15-days of outbreak emergence aligns with 7-1-7 targets and supports outbreak containment.

Methods

Using calibrated agent-based models for four diseases, a status-quo and series of 'Faster response' scenarios were compared for simulated outbreaks of each disease, with a 15-day ORI response time as the minimum.

Results

In a synthetic model population, a 15-day ORI response could avert: 80% of cases from

cholera outbreaks relative to a historical response time of 105 days; 35% of cases from

meningococcal meningitis outbreaks relative to a historical response time of 75 days; 0 –

35% of cases from yellow fever outbreaks relative to a historical response time of 105 days

(depending on routine vaccine coverage and environmental suitability); and 0 – 55% of

cases from measles outbreaks relative to a historical response time of 120 days (depending

on routine vaccine coverage).

Conclusions

Improvements made to ORI response time could reduce disease burden and decrease the

risk of large outbreaks of vaccine-preventable diseases in LMICs. Efforts to improve ORI

timeliness should be prioritised to higher risk settings, and it was clear that even a slow

vaccination response was beneficial relative to no response at all.

Keywords: Public Health; Infectious Diseases; Outbreaks; Vaccination; Measles; Cholera;

Yellow Fever; Meningococcal Meningitis

List of abbreviations

AIDS – acquired immunodeficiency syndrome

COVID-19 – coronavirus disease 2019

IMD – invasive meningococcal disease

LMIC – low and middle-income country

ORI – outbreak response immunisation

SDG – Sustainable Development Goals

VPD – vaccine-preventable disease

Background

The COVID-19 pandemic and 2013 – 2016 West African Ebola epidemic are two stark, recent

examples of how vital it is to have effective systems of detection, notification, and response

to combat disease outbreaks, and how disruptive and devastating they can become if the

systems are unprepared. The 7-1-7 targets developed by Resolve to Save Lives are gaining

traction as metrics for assessing a country's outbreak readiness; they propose that

outbreaks be detected within seven days; notification occur within one day; and key early

response actions be started within another seven days.[1] One of the seven early response

actions is to initiate appropriate public health countermeasures in affected communities,

which can include starting immunisation. Relatively few countries have adopted these

metrics yet, and of 41 public health events assessed across five low and middle-income

countries (LMICs) between 2018 – 2022, only 27% fully satisfied the 7-1-7 targets.[2]

Logistical challenges (such as cold-chain constraints for vaccines), weak healthcare

infrastructure, and understaffed workforces are some of the primary hurdles to overcome

when improving outbreak response timeliness.[3] Ideally, use of the 7-1-7 targets with

bottleneck and enabler identification can make every outbreak an opportunity for public

health system performance improvement, and are a tool for clear communication of

improvement to stakeholders.[2]

Outbreaks of vaccine-preventable diseases (VPDs) frequently occur in LMICs, often requiring

outbreak response immunisation (ORI) programs and other containment interventions.[4, 5]

Immunisation is a highly effective tool available for prevention of and response to VPD

outbreaks.[6-8] However, despite awareness, monitoring, and accessible stockpiles of

vaccines,[9] it can still take weeks or months for vaccines to be delivered once disease

outbreaks are detected.[4, 8, 10, 11] Governments and organisations which fund outbreak

response programs seek metrics by which to assess the timeliness of these responses, as a

measure of success and to identify areas for improvement, but in LMICs the lack of

consistent reporting makes benchmarking and target setting challenging.[2, 10, 12-14] The

Sustainable Development Goals (SDGs) Target 3.3 calls for the end of epidemics of very

high-burden infectious diseases such as HIV, tuberculosis, and malaria; and the combatting

of communicable diseases more broadly.[15] Unfortunately, there is a great deal of work

left to do to reduce the burden of infectious diseases to the levels indicated by SDG 3.3, and

the current approach does not appear to be working fast enough.[16]

It is expected that non-pharmaceutical interventions (e.g., contact tracing or public health

messaging) can typically be initiated quickly in outbreak contexts by utilising existing

workforces and systems, but accessing vaccine stockpiles can be slowed by logistical

challenges (e.g., physical distance or cold-chain constraints). Therefore, there is clear scope

for improvement in a critical element of outbreak response, and mathematical models can

be used to quantify the impact of improving ORI timeliness for policy makers.[17]

Previous modelling work found that for outbreaks of four diseases — measles,

meningococcal meningitis, yellow fever, and cholera — the time between outbreak

detection and immunisation response had considerable impact on the proportion of disease

burden averted by ORI.[8] Faster immunisation responses (typically around one month)

consistently prevented a higher proportion of disease burden than outbreaks with slower

responses (typically three to six months).[8] Despite this, the typical immunisation response

time for outbreaks of these diseases in LMICs is two to four months, with the Immunization

Agenda 2030 scorecard indicating that less than 30% of outbreaks had a timely detection

and response between 2021 – 2023.[18] To our knowledge, there are a limited number of

publications investigating the relationship between ORI timeliness and disease burden for

outbreaks of VPDs using mathematical modelling techniques,[19-29] and none which

estimate the impacts of achieving it within a period that supports 7-1-7 targets.

In this study, we estimate the impact of improving ORI timeliness for outbreaks of measles,

cholera, yellow fever, and meningococcal meningitis in LMICs, down to a minimum ORI

response time of 15 days to assess the incremental benefits of ORI timeliness gains in the

context of the 7-1-7 targets. For the purposes of this analysis, we considered 'ORI response

time' to be the time between an outbreak beginning and the initiation of an immunisation

response delivered to the population at-risk, as this captures the 15-day period covered by

the 7-1-7 targets.[1]

Methods

Data

We used a previously formulated dataset, which contains relevant epidemiological and

programmatic response data for 51 measles, 40 cholera, 24 meningococcal meningitis, and

88 yellow fever outbreaks in LMICs between 2000 – 2023 to calibrate our disease models.[8,

30] The mean ORI response time and daily vaccination rate from the set of outbreaks from

each disease were used to inform *Baseline* scenario parameters. The dataset is available in a

Zenodo repository.[30]

Models

We used previously developed agent-based models for measles, cholera, meningococcal

meningitis (serogroups C, W, X, Y), and yellow fever,[8] which were built using the *Starsim*

toolbox.[31] *Starsim* is an open-source agent-based framework developed by the Institute

for Disease Modeling at the Bill and Melinda Gates Foundation in collaboration with the

Burnet Institute and other partners.[31, 32] The framework is written in Python and allows

for rapid development of agent-based disease models that can be customised to arbitrary

levels of detail. The *Starsim* framework provides several key components that can be used

to build models:

• *Disease 'modules'*: capturing different health states and disease progression within

individuals.

• *Contact networks*: to simulate interactions between agents that can give rise to

transmission. Multiple networks can be used to allow for different mechanisms of

transmission and different settings (e.g., household networks, school networks,

random community contacts).

• *Interventions*: allow changes to be made to the model during simulations, such as

delivering vaccines, performing contact tracing, or changing transmission levels due

to non-pharmaceutical interventions brought in once an outbreak is detected.

We used the *Starsim* framework to build the models in this study to provide a single

common software and analysis architecture for all models, while simultaneously having the

flexibility to account for very different modes of transmission across the diseases being

modelled.

<u>Disease models:</u>

Each of the four disease models (i.e., *Starsim* disease modules) handles transmission events

differently:

- 154   The measles model simulates transmission through random human-human contacts
- 155   via networks generated randomly at each time step;

- 156   The meningococcal meningitis model simulates transmission through random
- 157   human-human contacts (regenerated each time step) and static, fully connected
- 158   household networks;

- 159   The cholera model simulates transmission through random contacts with tainted
- 160   water sources (calculated each time step based on cholera prevalence in the water)
- 161   and human-human contact in static, fully connected household networks;

- 162   The yellow fever model simulates transmission through random mosquito-human
- 163   and human-mosquito contacts (calculated each time step based on prevalence in
- 164   both populations).

165   Table 1 contains a summary of key model characteristics for each disease considered, and

166   Table 2 describes the key parameters and assumptions for each disease model and vaccine

167   used in the ORI programs. Figures S1, S7, S11, and S16 in 'Additional file 1' show the

168   schematic representation of each disease model. Additional details of each model can be

169   found in the 'Model overview' sections of Sections 2 – 5 in 'Additional file 1'.

	Modelled population	Transmission	Routine/ preventative vaccination	Asymptomatic cases	Non-pharmaceutical interventions	Mediating parameters for simulation
Measles	0 − 5	Direct	Y	N	N	Routine vaccine coverage
Cholera	All ages	Direct + Environmental (water)	N	Y	Y	
Yellow fever	All ages	Vector-borne	Y	Y	N	Routine vaccine coverage, environmental transmission level
Meningitis	All ages	Direct (inc. asymptomatic)	N (MenA outbreaks not included)*	Y	N	Asymptomatic carrier prevalence

Table 1: Summary of key model characteristics for each disease considered. *Pentavalent conjugate vaccines for serogroups ACWYX were prequalified and recommended for inclusion in routine vaccination programs by the WHO for countries in the African meningitis belt,[33, 34] however uptake is not yet widespread and has not been captured in this analysis.[35, 36]

Parameter	Value	Source
Meningococcal meningitis		
Average duration of exposed period	2 days	Model uses a lognormal distribution fit to results of analysis by Bosis et al. (2015).[37]
Average duration of asymptomatic infection	90 days	Mean is an estimate from Blakebrough et al. (1982),[38] and aligns with other recent estimates.[39, 40] Model uses a negative binomial distribution of to achieve a long tail as carriage durations vary significantly in the literature.
Average duration of symptomatic infection (to recovery)	7 days	Aligns with Karachaliou et al. (2015) meningitis A compartmental model.[41] Model assumes a normal distribution with a standard deviation of 1 day.
Average duration of symptomatic infection (to death)	8 days	Model uses a normal distribution fit to range of durations discussed by Sharew et al. (2020).[42]
Probability of invasive meningococcal disease (IMD)	~6%	The model uses age specific probabilities for IMD based on age-based risk estimates from Rivero-Calle et al. (2022),[43] and estimates of invasive disease incidence by age from the US Centers for Disease Control and Prevention.[44] However, the average probability of IMD was calibrated to match outbreak data.
Probability of death given symptomatic disease	7%	Calibrated value.[8] The estimated case fatality rate is 5 − 15%.[45-47]
Conjugate vaccine protection against IMD	90%	Effectiveness across A, C, Y, W-135 estimated to be 85 − 100% according to Martinez et al. (2013),[48] and 90% according to Daugla et al. (2014).[49]
Conjugate vaccine protection against asymptomatic carriage	41%	Aligned with value from Hadley et al. (2023).[50]
Cholera		
Average duration of exposed period	2·8 days	Model uses a lognormal distribution fit to results of systematic review by Azman et al. (2013).[51]
Average duration of asymptomatic infection	5·5 days	Model uses a uniform distribution of 1 − 10 days, based on World Health Organisation databook on cholera.[52]

Average duration of symptomatic infection (to recovery)	5 days	Commonly used estimate in modelling literature, according to review by Fung (2014).[53] Model uses a lognormal distribution fit to 5-day mean and range of $2-14$ days, based on values in Table 1 of the paper[53].
Average duration of symptomatic infection (to death)	1 day	According to Somboonwit et al. (2017),[54] death can occur within $6-12$ hours of symptoms for severe disease. Minimum timestep in the model is one day, and it assumes a lognormal distribution with mean of 1 day and range of $1-3$ days.
Duration of hyperinfectious bacteria	1 day	New infected agents shed hyperinfectious bacteria for first day, with 50x lower threshold for infection. Based on approach by Hartley et al. (2006) and other modelling studies.[55-57]
Probability of death given symptomatic infection	0·5%	Calibrated value,[8] typical case fatality rate is less than 1% for treated cholera.[58]
Probability of symptomatic cholera	50%	Calibrated assumption from upper range of ~25% and 57% estimates from Jackson et al. (2013).[8, 59] Literature reports wide range of estimates for asymptomatic disease.[60]
Relative transmissibility from asymptomatic infection	10%	Assumption, based on reduced bacteria shedding from asymptomatic cases reported by Nelson et al. (2009).[60]
Vaccine protection against infection	One dose: 52·7%	Based on single dose effectiveness study by Malembaka et al. (2024).[61] Maximum protection is reached 10 days after vaccination, based on $7-10$ day estimate after two doses from Song et al. (2021).[62]
Vaccine protection against death	50%	Used the estimate for protection against severe disease as a proxy, from single dose oral cholera vaccine efficacy estimates by Song et al. (2021).[62]
Measles		
Probability of death, given severe disease	0·5%	Calibrated value.[8]
Average duration of exposed period	$9-12$ days	Measles is infectious 4 days before the rash onset, which follows $2-4$ days after prodromal symptoms that appear $11-12$ days from exposure.[63]
Average duration of infection	8 days	Measles is infectious 4 days before and 4 days after the rash onset.[63]
Measles Testing Probability	10%	Assumption.
Isolation Factor	0·7	Assumption based on noted difficulties for measles isolation, particularly in low and middle-income countries.[64]
Vaccine protection against infection	83%; one dose	Based on vaccine efficacy for children over nine months old. Vaccine efficacy increases linearly over time and reaches peak of 83% after 10 days.[65]
Maternal Immunity	Age calculated in the model	Age calculated as weighted average of 3·78 and 0·97 months based on proportion of naturally infected and vaccinated mothers.[66]
Yellow fever		
Probability of death, given severe disease	20%	Calibrated value.[8]
Probability of mild or asymptomatic infection	33% + 55% (mild + asymptomatic)	Probability of yellow fever virus infections being asymptomatic, mild or severe.[67]

Probability of developing severe disease	12%	
Average duration of exposed period	3 − 6 days	Model uses a uniform distribution of 3 − 6 days, based on WHO data book on yellow fever.[68]
Average duration of mild infection (to recover)	3 − 4 days	Initial symptoms usually disappear after 3 − 4 days. Model uses a uniform distribution of 3 − 4 days, based on WHO data book on yellow fever.[68]
Average time to develop severe disease	4 − 5 days	Within 24 hours of recovering from initial symptoms, a small percentage of patients enter a second, more toxic phase. Therefore, the model uses a uniform distribution of 4 − 5 days (duration of mild disease 3 − 4 days + 24hrs), based on WHO data book on yellow fever.[68]
Average duration of severe infection (to recovery/death)	7 − 10 days	Proportion of patients who enter the toxic phase die within 7 − 10 days. Model uses a uniform distribution of 7 − 10 days, based on WHO data book on yellow fever.[68]
Vaccine protection against infection	97·5%	Based on random-effects meta-analysis and pooled estimate of serological response. Vaccine protection increases linearly over time; it reaches 80% after 10 days and 97·5% after 30 days. Time courses from WHO.[68] Peak vaccine protection from Jean et al. (2016).[69]

Table 2: Disease model and vaccine parameters for meningococcal meningitis, cholera, measles, and yellow
fever.

<u>Contact networks</u>:

Each network in the models can have a different structure, with agents being connected to

their contacts randomly or agents being grouped into disconnected household clusters.

Each agent in the models (except the yellow fever model, which has no human-human

transmission) has a specified number of contacts in at least one network. For those who

have a non-zero number of contacts in a particular network, if the contact network is

random, then their number of contacts per day is drawn from a Poisson distribution with a

mean of 11, 9, and 3·5 for meningococcal meningitis, cholera, and measles, respectively. For

each disease, these means were estimated from Prem et al. (2017),[70] with values

averaged over all non-household contact rates and all ages, for all countries with outbreaks

in the outbreak dataset.[30]

If the contact network is a household, then the size of each cluster is drawn from a synthetic

household size distribution generated by summing over household size data (from the

United Nations, Department of Economic and Social Affairs, Population Division)[71] for

each country with an outbreak included in the outbreak dataset.[30]

<u>Vaccine characteristics</u>:

For each modelled disease there was a vaccine intervention included in the simulations, for

use during ORI. The vaccine effects varied by disease (with relevant characteristics and

efficacies detailed in Table 2), but all acted to at least moderately protect vaccinated agents

in the outbreak simulations against infection. For all vaccines, the implementation within

the models assumed a linear increase in protection against infection over time, achieving

52.7%, 83.0%, and 97.5% for cholera, measles, and yellow fever, respectively. Both the

polysaccharide and conjugate meningitis vaccines provide 90% protection against invasive

disease, but the conjugate vaccines also provide 41% protection against asymptomatic

infection. The times to maximal protection were 10 days for the cholera and measles

vaccines[62, 65]; eight days for the polysaccharide and conjugate meningitis vaccines[48];

and 30 days for the yellow fever vaccine[68].

Model populations

The disease models were used to run simulations of future outbreaks in a synthetic model

population intended to be representative of LMIC contexts where outbreaks have typically

occurred, with ORI response parameters based on the mean values from the outbreak

dataset (unless varied as a part of a scenario analysis).[30] Each disease model simulated a

population of 50,000 agents, intended to be large enough to capture widespread

transmission in an outbreak area but small enough to limit computational intensity. The age

distributions of the agents in each model were generated by summing over data from

countries where outbreaks occurred in our dataset, informed by single-year age distribution

data from United Nations, Department of Economic and Social Affairs, Population

Division.[71]

Calibration

The model calibration was unchanged from previous work, which has been published in

detail elsewhere.[8] In brief, each model was calibrated to a set of historical outbreaks in

LMICs which received ORI as a part of the outbreak response. The calibration ensured that

model simulations could reproduce the range of outbreak sizes and durations observed in

the dataset.[30]

The calibration method controlled for epidemiological and ORI programmatic parameters

such as response time, daily vaccination rate, routine vaccine coverage, the mosquito-to-

human transmission modifier, or the asymptomatic carriage prevalence.[8] The transmission

modifier parameter was used to capture the impact of temperature and rainfall on

mosquito population size and biting behaviours on yellow fever transmission in outbreak

settings. Asymptomatic carriage prevalence was a parameter used to modify the initial

growth rate of an outbreak as an epidemic season begins in meningococcal meningitis

outbreak settings. Further details about the calibration process have been published

previously,[8] and a summary can be found in Section 1 in 'Additional file 1'. Further,

additional details about the specific calibrations for each model are available in Sections 2 –

5 in 'Additional file 1'.

Unless varied during a scenario (i.e., response time, achieved coverage, or vaccination rate)

almost all disease and intervention parameters in the four models are informed by

estimates from the literature or were determined during calibration (detailed in Table 2).

The only three exceptions to this were the routine vaccine coverage, asymptomatic carriage

prevalence, or mosquito-to-human transmission modifier parameters, which were used to

investigate outbreaks in settings with different levels of risk. The primary parameters for

each disease which were determined during calibration were the overall transmissibility; the

probability of death, given severe infection; and the number of seed infections (except for

meningitis, which was triggered by asymptomatic carriage in the population). For the

cholera model, the probability of symptomatic disease was also estimated during

calibration.

Outbreak detection and response

For cholera, measles, and yellow fever, the threshold for outbreak declaration in the models

was defined as one case. According to the WHO, measles outbreaks are typically declared if

there are five or more epidemiologically linked cases.[72] Cholera outbreaks are considered

'suspected', 'probable', or 'confirmed' for a series of thresholds, beginning with two

suspected cases or one suspected case with a positive rapid diagnostic test.[73] The WHO

also states that outbreaks are declared in the 'presence of at least one confirmed case of

yellow fever'.[74] As the models do not differentiate between suspected and confirmed

cases, the assumption for the threshold is optimistic for measles; however for cholera and

yellow fever, the thresholds used in the model are identical or similar to declaration

thresholds stated by the WHO. The case is assumed to be detected on the first day of the

simulation, a one-day delay is assumed for notification, and the ORI begins after a further

(N-2)-days, where N is the response time used for each scenario.

For meningococcal meningitis, the threshold for outbreak declaration in the model is five

cases of invasive disease within one week. This threshold aligns with the typical threshold of

10 cases per 100,000 population used for districts with a population greater than

30,000.[75] Once this threshold is crossed, it is detected within one simulation day and the

timeline for response is the same as for the other disease models.

These detection and notification timelines are optimistic, but the purpose of this analysis is

to consider the impact of achieving a set of very ambitious timeliness targets.

Scenarios

To estimate the impact of ORI response time on outbreak outcomes for each disease we ran

two main scenarios and two sensitivity analyses:

• *Baseline*; response time and daily vaccination rate were based on the mean value

from the outbreak dataset for each disease, and ORI was assumed to achieve 75%

coverage in the target population (except for measles, which we assumed achieved

100% coverage);

• *Faster response time*; the ORI response time was varied between 15 days and the

Baseline response time in steps of 15 days;

• *Alternate coverage* (sensitivity); the ORI coverage achieved was increased to 100%

(except for measles, which was varied between 60% and 90% coverage in steps of

10%) in the target population. For meningococcal meningitis we also ran an

additional sub-scenario with broadened age-eligibility and 75% coverage;

• *Alternate vaccination rate* (sensitivity); the daily vaccination rate was varied to

produce either a one-day campaign or a one-month campaign for each disease.

Each scenario analysed comprised 1000 stochastic simulations. Table 3 describes the

parameter values varied in the scenarios for each disease, and Figure 1 represents the

scenario impacts on the outbreak simulations. Specific justifications (and sources) for the

parameter values chosen for each disease under each modelled scenario are available in

Sections 2 − 5 in 'Additional file 1'.

Disease	Response time (days)		Achieved coverage (%)		Vaccination rate (vaccines/day)	
	Baseline	Faster response times	Baseline	Alternate coverage	Baseline	Alternate vaccination rate
Measles	120	15 − 105	100%	60 − 90%	4360	47,812**; or 1300^^
Cholera	105	15 − 90	75%	100%	5100	48,997**; or 1300^^
Meningococcal meningitis	75	15 − 60	75%*	100%*; or 75%^	4300	33,153**; or 800^^
Yellow fever	105	15 − 90	75%	100%	6560	48,544**; or 1300^^

Table 3: Parameter values used for each scenario analysed for measles, cholera, meningococcal meningitis,
and yellow fever. * targeting ages 1 − 29 years; ^ targeting ages 1+ years; ** vaccinating the eligible population
in one day; ^^ vaccinating the eligible population in one month.

Figure 1: Representation of how the modelled scenarios impact the outbreak simulations. Outbreak response
immunisation (ORI) implemented earlier in an outbreak is expected to reduce the total size and duration of the
outbreak.

Setting archetypes

As an additional dimension of this analysis, we aimed to explore the impact of ORI response time for different 'setting archetypes' in which transmission was expected to be relatively higher or lower for each disease. In previous work, key parameters were identified for the measles, meningococcal meningitis, and yellow fever models, which varied by outbreak setting and influenced the expected size of outbreaks in our models.[8] For each scenario considered in this analysis, we simulated outbreaks over a range of values for these parameters (routine vaccine coverage, asymptomatic carriage prevalence, or mosquito-to-human transmission modifier), which represent outbreaks occurring in settings with different levels of risk that the outbreak may grow particularly large.

Impact estimation and outbreak size thresholds

For each disease, setting archetype, and scenario, we recorded the distribution of cumulative cases and deaths which occurred across the outbreak simulations. The mean difference in cumulative cases and deaths between scenarios and associated uncertainty in the mean were estimated from these collections of simulations using bootstrap resampling (i.e., for each outbreak to produce estimates of cases averted with a faster or further reaching response).

We produced a distribution of outbreak sizes for each disease, setting archetype, and scenario, by defining a series of outbreak size thresholds for each disease and assigned each simulation to its appropriate threshold range. This allows us to estimate the frequency with which outbreaks exceeding certain total case thresholds occur, given a response time and achieved coverage by the ORI, based on methods adapted from previous analyses of COVID-19[76] and Ebola.[77] The impact of ORI on reducing the risk of large outbreaks was

315   estimated by comparing the distribution of cumulative cases across simulations in the

316   *Baseline* scenario against the distributions in each of the *Faster response time* scenarios. All

317   impacts were disaggregated by setting archetypes, where relevant.

Results

Figure 2: Area plots representing the proportion of outbreak simulations which exceed certain case thresholds.
These proportions are plotted as a function of outbreak response immunisation (ORI) response time;
disaggregated for different diseases (rows) and for worst case (left column) and best case (right column)
setting archetypes considered. For each panel, the righthand stacked bar represents the distribution of
outbreak sizes if no outbreak response immunisation is delivered, and the dashed line represents a 10%
threshold for visualization. Panels a) and b) are meningococcal meningitis for settings with 10% and 0·25%

initial meningococcal meningitis carriage prevalence, respectively. Panel c) is for cholera (only one setting
archetype considered). Panels d) and e) are measles for settings with 50% and 100% routine measles vaccine
coverage. Panels f) and g) are yellow fever for settings with 40% and 80% routine yellow fever vaccine
coverage (BC) and with a mosquito-human transmission modifier of 1·5 and 0·5, respectively.

Across the four diseases, improvements to ORI response times reduced the proportion of

simulations exceeding different outbreak size thresholds in higher risk setting archetypes

(i.e., those with lower routine vaccine coverage, higher initial prevalence, or higher

transmission modifiers) (Figure 2, left column). This behaviour was not observed (or the

relationship was weaker) in the lower risk settings (Figure 2, right column). Additional

examples of these figures for different setting archetypes are in Sections 2, 4, and 5 in

'Additional file 1', however those in Figure 2 were chosen as the 'worst' and 'best'

archetypes for each disease (except cholera) which show the highest and lowest overall risks

for large outbreaks.

Each subplot in Figure 2 also has a secondary stacked bar graph next to the area graphs

showing the distribution of model simulations in which outbreaks exceed the defined case

thresholds. The stacked bar plot shows the same distribution for a set of model simulations

in which ORI was not implemented during the outbreaks. In the high-risk setting archetypes

(and the single cholera subplot), the proportion of simulations which exceeded the highest

case thresholds when no ORI was used was always higher than for the slowest response

time simulations. This was also observable in the figures for different setting archetypes in

Sections 2, 4, and 5 in 'Additional file 1'. In these modelled settings, even a slow ORI

response was beneficial relative to no response at all.

Meningococcal meningitis

The expected size of the meningococcal meningitis outbreaks was strongly dependent on

the initial carriage prevalence (Figure S4, Figure S5 in Section 2 in 'Additional file 1'). When

the initial prevalence was low outbreaks tended to be quite small and ORI response times

had minimal impact on expected outbreak size. However, when the initial prevalence was

high the expected outbreak size was larger, and response time had a greater impact on the

proportion of simulations which exceeded different case thresholds (Figure 2 and Figure S6

in Section 2 in 'Additional file 1').

While the ORI response time was more critical for absolute numbers of cases in settings

with higher initial prevalence, the relative impact of a faster ORI response time was

consistent across the choice of initial prevalence (Figure 3). For example, a 15-day response

time was estimated to avert ~35% of the case burden compared to the mean historical

response time (75 days) between 2011 – 2022 regardless of initial prevalence. Similarly, a

30-day response time averted ~25% of the cases, a 45-day response averted ~15%, and a

60-day response averted ~7% (Table S2 in Section 2 in 'Additional file 1').

Cholera

The expected size of the cholera outbreaks was strongly impacted by the ORI response time (Figure S9 in Section 3 in 'Additional file 1'), and the proportion of simulations exceeding certain case thresholds also increased with ORI response time (Figure 2 and Figure S10 in Section 3 in 'Additional file 1'). For example, slower response times noticeably increased the proportion of simulations which exceeded 400 or 500 cumulative cases. When comparing outcomes with faster ORI response times to outcomes with the mean historical response time (105 days) between 2015 – 2023, a 15-day response time was estimated to avert an additional ~80% of the cases. Similarly, a 30, 45, 60, 75, or 90-day response averted an additional ~70%, ~55%, ~40%, ~30%, or ~25% of the cases, respectively (Figure 4 and Table S3 in Section 3 in 'Additional file 1').

Figure 4: Proportional reduction in the mean cumulative cases and deaths for the simulated cholera outbreaks. Results include outbreaks with response times of 15, 30, 45, 60, 75 and 90 days, relative to the set of outbreaks with a response time of 105 days, estimated using bootstrap resampling. The values stated at the

385 bottom of the column are the mean cumulative cases or deaths produced by cholera outbreaks which received
386 ORI after 105 days, i.e., the value serving as the denominator for the proportional reduction estimate.

Measles

The expected size of the measles outbreaks was strongly dependent on the routine vaccine

coverage (Figure S13, Figure S14 in Section 4 in 'Additional file 1'), and the proportion of

simulations exceeding certain high case thresholds decreased with routine coverage and

increased with ORI response time (Figure 2 and Figure S15 in Section 4 in 'Additional file 1').

When the routine vaccine coverage was high (i.e., in a low-risk setting archetype), outbreaks

tended to be quite small, even with longer ORI response times. However, in high-risk

settings, the expected outbreak size increased and ORI response time had a greater impact

on outcomes, with slower ORI response time scenarios having an increased proportion of

simulations which exceeded 5000 cumulative cases.

ORI response time was more critical for absolute numbers of cases in higher risk setting

archetypes, and the proportional impact of a faster ORI response time was also greater for

outbreaks in these settings. Interestingly, the estimated impacts of faster response times

decreased to 0% as the routine vaccine coverage approached 100% (Figure 5), which

occurred because high routine vaccine coverage settings had smaller outbreaks and saw less

benefit from fast ORI response times due to slower transmission. For example, in a setting

with 50% routine vaccination coverage, a 15-day response time was estimated to avert

~55% of the cases compared to the mean historical response time (120 days) between 2000

– 2022, while a 105-day response time averted ~13%. However, in a setting with 90%

routine vaccine coverage, a 15-day response time was estimated to avert ~8% of the cases,

while a 105-day response time averted no additional cases compared to a 120-day response

time (Table S4 in Section 4 in 'Additional file 1').

Figure 5: Proportional reduction in the mean cumulative cases and deaths for the simulated measles outbreaks. Results include outbreaks with response times of 15, 30, 45, 60, 75, 90 and 105 days relative to the set of outbreaks with a response time of 120 days, estimated using bootstrap resampling. Proportional impacts are grouped vertically by routine measles vaccine coverage, and the values stated at the bottom of each column are the mean cumulative cases or deaths produced by measles outbreaks which received ORI after 120 days, i.e., the value serving as the denominator for the proportional reduction estimate.

Yellow fever

The expected size of the yellow fever outbreaks was strongly dependent on both the routine vaccine coverage (Figure S17, Figure S19 in Section 5 in 'Additional file 1') and the mosquito-human transmission modifier (Figure S18, Figure S20 in Section 5 in 'Additional file 1'). The proportion of simulations exceeding certain high case thresholds decreased with routine coverage, increased with ORI response time (Figure 2 and Figure S21 in Section 5 in 'Additional file 1'), and increased with the transmission modifier (Figure S22 in Section 5 in 'Additional file 1'). When the routine vaccine coverage was high or the transmission modifier was low (i.e., in a low-risk setting archetype), outbreaks tended to be quite small even with longer response times. However, in a high-risk setting archetype, the expected

outbreak size increased and slower response time scenarios increased the proportion of

simulations which exceeded 500 or 1000 cumulative cases.

ORI response time was more critical for absolute numbers of cases in high-risk setting

archetypes, and the proportional impact of a faster response time was also greater for

outbreaks in these settings (Figure 6 and Figure S23 in Section 5 in 'Additional file 1'). For

example, when the transmission modifier was fixed at 1·0, in a 50% routine vaccination

setting, a 15-day response time was estimated to avert ~25% of the cases compared to the

mean historical response time (105 days) between 2000 – 2022, while a 90-day response

time averted no additional cases. While in an 80% coverage setting, a 15-day response time

was estimated to avert ~11% of the cases, and a 90-day response time averted no additional

cases compared to a 105-day response time (Tables S5, Table S6 in Section 5 in 'Additional

file 1').

Figure 6: Proportional reduction in the mean cumulative cases and deaths for the simulated yellow fever
outbreaks. Results include outbreaks with response times of 15, 30, 45, 60, 75, and 90 days relative to the set
of outbreaks with a response time of 105 days, estimated using bootstrap resampling and assuming a
transmission modifier of 1·0 for the mosquito vectors. Proportional impacts are grouped vertically by routine
yellow fever vaccine coverage, and the values stated at the bottom of each column are the mean cumulative
cases or deaths produced by outbreaks which received ORI after 105 days, i.e., the value serving as the
denominator for the proportional reduction estimate.

Sensitivity analysis

The results of the alternate ORI coverage sensitivity analysis can be seen in Figures S24 – 40

in Section 6 in 'Additional file 1'. For meningococcal meningitis and cholera outbreaks,

increasing achieved ORI coverage decreased the cumulative cases by up to 20 – 25%. For

measles outbreaks, we found that when the achieved ORI coverage was low, outbreaks

grew quite large even with rapid ORI response times, because large proportions of the

population were missed by the response. The effect varied by setting archetype, and we

observed less impact from gains in ORI coverage achieved if the routine coverage level was

higher.

The results of the daily vaccination rate sensitivity analysis can be seen in Figures S41 – 48 in

Section 6 in 'Additional file 1'. Meningococcal meningitis, cholera, and measles outbreaks

showed little sensitivity to increasing the daily vaccination rate to deliver all doses in a single

day, as the rates used in the primary scenarios were already high (varying by disease).

However, yellow fever outbreaks were more sensitive to this change, with total cases

decreasing by up to 40% across varying levels of response time and routine vaccine

coverage, for a transmission modifier value of 1·5. For lower transmission modifier values,

the decrease was smaller but still noticeable. This is likely due to the relatively longer time

for the immunity to fully develop within the vaccinated agents (30 days for yellow fever[68],

relative to ten days or fewer for the other diseases[48, 62, 65]), making earlier vaccine

delivery relatively more impactful than for the other diseases. The impact of decreasing the

vaccination rate such that the doses are delivered over the course of a month varied by

disease, and within each disease varied by response time or by setting archetype. The

cholera, yellow fever, and measles ORI were sensitive to the decrease, particularly for faster

ORI response times and settings with lower routine vaccine coverage (for measles). Yellow

fever outbreaks in the highest risk setting archetypes were up to three times larger if the

ORI took one month to vaccinate the population instead of one week – which also likely

results from the slower immunity development – however such a slow rollout would be

atypical.

Discussion

We used agent-based models to estimate the impact of faster ORI response times for

outbreaks of measles, cholera, yellow fever, and meningococcal meningitis in synthetic LMIC

populations, stratified into setting archetypes. We found that ORI responses initiated within

the timeframe of the 7-1-7 framework (15 days of the outbreak beginning) would avert a

larger proportion of cases and deaths than the average response times for recent historical

outbreaks in LMICs.

In the models, a 15-day ORI response time reduced the expected burden of cases and

deaths from meningococcal meningitis outbreaks by approximately 35%, cholera outbreaks

by 80%, and from measles and yellow fever outbreaks in settings with low routine vaccine

coverage by up to 55% and 35%, respectively. This is consistent with previous modelling

analyses, which estimated that implementing vaccination responses faster for outbreaks of

measles, Ebola, cholera, yellow fever, poliovirus, and meningococcal meningitis reduced

disease burden.[8, 19-26, 28, 29] This also supports the push for frameworks like 7-1-7,

because the diseases considered here are responsible for a large health burden in

LMICs,[78] and the time taken to initiate ORI in LMICs is typically on the order of months.[8,

11] If ORI timeliness can be improved then delivering the same number of vaccines will have

a much greater protective impact for the population at-risk, which could help accelerate

progress towards targets like SDG 3.3. While these results are not granular enough and the

models used not detailed enough to directly inform how ORI timeliness could be improved,

ideally these impact estimates could motivate bottleneck analyses and other studies which

could directly inform investment priorities.

Caution is needed when focusing on ambitious targets like 7-1-7, particularly when they are

optimistic compared to current timeliness metrics for ORI in LMICs,[4, 10] because it may be

viewed as unrealistic to achieve in practice. In the case of the diseases modelled here,

achieving the 7-1-7 targets for the ORI component of an outbreak response reflects a more

than 80% reduction in response times relative to the historical averages used in the *Baseline*

simulations, which may be challenging to achieve in many low-resource settings.

In high-income settings, robust reporting principles, established public health surveillance

systems, and strong infrastructure with a well-trained workforce facilitate rapid detection of

outbreaks and rapid initiation of appropriate response measures. In contrast, in LMICs,

delays to each stage of detection, notification, and response are more common because of

the relatively weaker systems, infrastructure, and workforce training, among other causes.

Even when these delays are known, the lack of consistent reporting during outbreaks in

LMICs can make benchmarking and identifying areas for improvement a challenge.[2, 10,

12-14] It should be noted, however, that these results still indicate that even a slow ORI

response is noticeably better than no response at all (Figure 2). Reactive vaccination is an

effective intervention, even if delayed, but our results indicate that improving timeliness

could be a clear method for maximising the protective impact achievable during an

outbreak.

In low-resource settings, a more nuanced approach, with incremental targets or a graded

assessment of progress towards specified targets is preferable to an 'all-or-nothing' goal,

and is more in-line with the intent of the 7-1-7 Alliance and Resolve to Save Lives to identify

and address bottlenecks rapidly in outbreak response.[1, 2] For example, consider a given

year in which a country had no outbreak responses achieve 7-1-7, but 90% of its responses

achieved 7-1-37 (for a total time of 45 days). If those responses were estimated to have

averted a large proportion of disease burden relative to the country's average response

time from previous years, then the level of progress should be noted and a subsequent goal

perhaps set towards 30 days, rather than being assessed as making no progress toward 7-1-

7. Our results indicate that achieving a 15-day ORI response time could avert an additional

35 – 80% of cases and deaths, but we also examined the benefits of incrementally faster ORI

response times, finding that any improvements relative to the average historical response

times reduces both the expected outbreak size and negative health outcomes.

The benefits of incrementally improving ORI response times varied widely across diseases

and setting archetypes, suggesting that there may be utility in having context-specific

timeliness targets, and that efforts to improve ORI response times could be focused for

increased impact. When outbreaks were simulated in higher risk settings (i.e., those with

lower routine vaccine coverage, higher initial prevalence, or higher transmission modifiers)

we consistently found a stronger relationship between ORI response time and outbreak size

than in lower risk settings. By computing risk profiles for setting archetypes (as in Figure 2),

context-dependent timeliness targets could be designed which minimise the risk of large

outbreaks but are evidence-based and support prioritisation of limited resources for

greatest benefit. For example, in a setting where the meningococcal meningitis carriage

prevalence in the lead-up to an epidemic season was suspected to be high (e.g., 10% in a

district), a 30-day ORI response time may be adequate to keep the risk of an outbreak larger

than 500 cases below 10%. For yellow fever, in low-risk outbreak settings, ORI response

time may make minimal difference to outcomes and so efforts for improvement may be

better directed elsewhere. This information allows decision makers to prioritise efforts

where incremental improvements have the greatest impact.

Another factor to consider when designing ambitious targets for response time is the

potential for diminishing returns. While it seems clear that a shorter response time would

not lead to worse outcomes during an outbreak, our results indicate that it may not always

produce better outcomes. In low-risk setting archetypes the impact of improving ORI

response time on both outbreak size and proportional reductions was typically quite small,

but the reasons for this are clear. High routine vaccine coverage reduces both the pool of

susceptible people and the rate of transmission during an outbreak of measles or yellow

fever, for example. What is more interesting is that there may be diminishing returns in

high-risk settings as well, depending on the disease. For measles outbreaks in settings with

low routine vaccine coverage there were benefits from each incremental improvement to

ORI response time from the historical average of 120 days down to 60 days, with each 15-

day improvement to this time reducing total cases by $10 - 15\%$, however for even faster

responses there was almost no additional benefit. In contrast, for meningococcal meningitis,

the expected size of an outbreak was modified by the initial carriage prevalence in the

population before an epidemic season, but the proportional impact of each improvement in

ORI response time was largely independent of this. We found that each 15-day

improvement reduced the total cases by 7 – 10% for each level of initial carriage.

One of our sensitivity analyses considered the impact of achieving varying levels of ORI

coverage during an outbreak. Achieving higher coverage with ORI averted a higher

proportion of cases, as would be expected, with a relatively smaller effect size observed

when response times were longer. Interesting implications can be drawn from the results of

the measles ORI coverage sensitivity analysis specifically, as it varied both ORI coverage and

response time. Figure S27 in Section 6 in 'Additional file 1' shows that the expected size of

outbreaks in the measles model was more strongly impacted by the level of ORI coverage

achieved than by ORI response time (over the parameter ranges tested in our simulations).

This indicates that for measles, it may be preferable to prioritise a comprehensive ORI

campaign that reaches as many people as possible, even if it takes slightly longer to

implement, as this would help close immunity gaps that allow continued transmission.

Future modelling could explore this potential trade-off between response speed and

coverage in more depth and across other diseases. This is an important consideration in the

context of reaching unvaccinated children, particularly those who live in remote

communities. The logistical and sociological difficulties of reaching these subpopulations to

try and achieve high levels of vaccine coverage in a geographic region are well known,[79]

and while the assumption of achieving 100% coverage in the measles analysis may be overly

optimistic, hopefully the results presented here can help to motivate additional efforts to

reach these vulnerable populations.

This analysis focused on outbreaks of four vaccine preventable diseases — meningococcal

meningitis, cholera, measles, and yellow fever — and found that the relationship between

ORI response time and impact differs for all of them. However, there are some lessons that

can be generalised across outbreaks of VPDs in LMICs. Where there are known factors which

affect outbreak risk or the rate of transmission, ORI response time is typically more critical

in higher risk settings. In the face of slowing and declining global vaccination rates

(particularly since the COVID-19 pandemic) and the ongoing impacts of climate change,

many countries may begin to face increasing outbreak risks[80, 81]. As such, developing and

maintaining systems of disease surveillance to assess these risks could help to inform the

importance of rapid outbreak response. Strong surveillance systems would facilitate faster

and more sensitive outbreak detection, a core tenet of 7-1-7, making rapid responses more

achievable. Through development of technology and investments in health systems, the

timeliness for both detection and response has improved over recent decades, but there is

still room for growth.[82, 83] We believe there is potential for future mathematical

modelling work to meaningfully inform the benefits of enhancements to surveillance

systems on the impact of outbreak responses using models such as ours.

Limitations

This was a broad modelling analysis, using four different agent-based models in synthetic

populations, and both the models and methodology are subject to limitations. Specific

limitations for each disease model are described in Sections 2 – 5 in 'Additional file 1'.

Broadly, the study methodology was limited by the nature of our calibration process, and by

the assumptions which informed our scenarios. The calibration of our models was

performed collectively on outbreak datasets of different sizes,[8] and while each model

reproduces the range of outbreak sizes observed in the data, they are biased towards the

most frequently observed outbreak sizes in LMICs. For the purposes of this prospective

analysis, it means that the model simulations represented outbreaks similar to the average

outbreak observed in LMICs between 2000 – 2023 for each disease, but future outbreaks

may behave differently. One additional factor which may influence this is that the period

between 2000 – 2023 covers the COVID-19 pandemic, which has had measurable impacts

on routine vaccine coverage for measles and yellow fever,[84] and may have affected

changes in human behaviour and contact networks. Our exploration of different vaccine

coverage scenarios provides insight into the first impact, but it is unclear to what extent

contact patterns may have changed and our models therefore cannot account for these

changes in potential future outbreaks. As our models and synthetic populations were

designed to focus on outbreaks in LMICs, these results are not generalisable outside of an

LMIC context.

As our analysis only considered the impact of ORI programs on cases and deaths accrued

over the course of a single outbreak for each set of simulations, we do not account for the

mid to long-term protective impacts which may be imparted by the vaccines. It is unclear

how this would change the estimated relative impacts of improving ORI response time

which we presented, but it means that we are likely underestimating the overall protective

impact imparted by the ORI campaigns. Particularly in the high-risk setting archetypes

where the risk was driven by low routine vaccine coverage.

We aimed to understand the effects of the key parameter assumptions in our scenarios by

showing simulations over a range of setting archetypes, and by performing a sensitivity

analysis for the daily vaccination rate and achieved ORI coverage, but outbreaks and

responses which exist outside of the considered range of the parameters may not be well

represented by our results. While the models capture a lot of stochastic uncertainty

associated with outbreaks, we did not capture uncertainty in some of our model

parameters, such as vaccine efficacy or the probability of symptomatic disease for cholera

and meningococcal meningitis, using point estimates from the literature.

Finally, our analysis only considered the impact of improvements to ORI timeliness, and no

other components of outbreak response, noting ORI is only one aspect amongst the seven

key early response actions within the 7-day response component of the 7-1-7 targets. We

also did not disaggregate the ORI response times by detection-notification-response as

defined by 7-1-7. As such we do not capture impacts from faster implementation of

programs such as contact tracing, or distinguish whether timeliness improvements were

made for detection, notification, or response.

Conclusion

The results of this analysis indicate that achieving a 15-day vaccine response to outbreaks of

measles, cholera, yellow fever, or meningococcal meningitis in LMICs, as per the 7-1-7

targets, could prevent a large disease burden. Furthermore, we found that there are

benefits to be gained from incremental improvements to response time even when 7-1-7 is

not achieved, and we support focusing on incremental improvements in timeliness

irrespective of whether the 7-1-7 target is routinely achievable. We found a risk-associated

variability in the impact of achieving faster responses, where response time gains were

more impactful in higher risk settings than lower risk settings. Therefore, we expect that a

graded approach paired with context-specific risk assessments could also facilitate resource

prioritisation decisions if multiple outbreaks occur in a short timeframe.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

Model code and outputs are published on Zenodo at
<https://doi.org/10.5281/zenodo.15825915> and will be publicly available indefinitely. The
outbreak data previously used for model calibration are available on Zenodo at
<https://doi.org/10.5281/zenodo.15751150>.

Competing interests

The authors declare that they have no competing interests. No author has been paid to
write this article by a pharmaceutical company or other agency.

Funding

D.D. has received funding for a PhD scholarship from the National Health and Medical
Research Council in Australia (Award Number: 2021/GNT2014056), which partially funded
this work. The funding source was not involved in the study or decision to submit the
manuscript for publication.

Authors' contributions

Conceptualisation, D.D., R.G.A. and N.S.; formal analysis, D.D.; data curation, D.D. and
A.M.M.; investigation (modelling), D.D.; methodology, D.D., R.G.A. and N.S.; project
administration, N.S.; software, D.D., A.M.M. and R.G.A.; supervision, R.G.A. and N.S.;

validation, D.D., A.M.M., J.G., R.G.A. and N.S.; visualisation, D.D.; writing—original draft,

D.D.; writing—review and editing, D.D., A.M.M., J.G., R.G.A. and N.S. Authors were not

precluded from accessing data in the study, and they accept responsibility to submit for

publication.

Acknowledgements

The authors thank Gabrielle MacKechnie, Dr Stefanie Vaccher, Dr Todi Mengistu, and Dr Dan

Hogan for their previous contributions to the outbreak dataset and models used in this

analysis. The authors also thank Fenella McAndrew for her advice when creating some of

the graphics.

References

1.   Frieden TR, Lee CT, Bochner AF, Buissonniere M, McClelland A. 7-1-7: an organising
principle, target, and accountability metric to make the world safer from pandemics. Lancet.
2021;398(10300):638-40.
2.   Bochner AF, Makumbi I, Aderinola O, Abayneh A, Jetoh R, Yemanaberhan RL, et al.
Implementation of the 7-1-7 target for detection, notification, and response to public health
threats in five countries: a retrospective, observational study. Lancet Glob Health.
2023;11(6):e871-e9.
3.   Crealock-Ashurst B, Levanita S, Frischer SR, Adhikari S, Mwebia W, Kadri-Alabi Z, et
al. PEARLES challenges and solutions to the implementation of clinical research responses to
epidemics and pandemics: a scoping review. (2589-5370 (Electronic)).
4.   Impouma B, Roelens M, Williams GS, Flahault A, Codeco CT, Moussana F, et al.
Measuring Timeliness of Outbreak Response in the World Health Organization African
Region, 2017-2019. Emerg Infect Dis. 2020;26(11):2555-64.
5.   Realizing Africa CDC's Autonomy: Making Measurable Impact in Africa's Health
Security. Addis Ababa, Ethiopia: Africa Centres for Disease Control and Prevention; 2024.
6.   Carter A, Msemburi W, Sim SY, Gaythorpe KAM, Lambach P, Lindstrand A, et al.
Modeling the impact of vaccination for the immunization Agenda 2030: Deaths averted due
to vaccination against 14 pathogens in 194 countries from 2021 to 2030. Vaccine. 2024;42
Suppl 1:S28-S37.
7.   Shattock AJ, Johnson HC, Sim SY, Carter A, Lambach P, Hutubessy RCW, et al.
Contribution of vaccination to improved survival and health: modelling 50 years of the
Expanded Programme on Immunization. Lancet. 2024;403(10441):2307-16.
8.   Delport D, Muellenmeister AM, MacKechnie G, Vaccher S, Mengistu T, Hogan D, et
al. Estimating the historical impact of outbreak response immunisation programmes across
210 outbreaks in low and middle-income countries. BMJ Global Health. 2025;10(7):e016887.

9.   International Coordinating Group (ICG) on Vaccine Provision - About us: World
Health Organization;[accessed 2025-01-06]. Available from:
https://www.who.int/groups/icg.
10.   Fieldhouse JK, Randhawa N, Fair E, Bird B, Smith W, Mazet JAK. One Health
timeliness metrics to track and evaluate outbreak response reporting: A scoping review.
EClinicalMedicine. 2022;53:101620.
11.   Pezzoli L. Global oral cholera vaccine use, 2013–2018. Vaccine. 2020;38:A132-A40.
12.   Crawley AW, Divi N, Smolinski MS. Using Timeliness Metrics to Track Progress and
Identify Gaps in Disease Surveillance. Health Secur. 2021;19(3):309-17.
13.   Smolinski MS, Crawley AW, Olsen JM. Finding Outbreaks Faster. Health Secur.
2017;15(2):215-20.
14.   Fieldhouse J, Nakiire L, Kayiwa J, Brindis CD, Mitchell A, Makumbi I, et al. How
feasible or useful are timeliness metrics as a tool to optimise One Health outbreak
responses? BMJ Glob Health. 2024;9(7).
15.   Transforming Our World: The 2030 Agenda for Sustainable Development: United
Nations;[accessed 25/03/2024]. Available from: https://sdgs.un.org/2030agenda
16.   The Sustainable Development Goals Report 2024. New York: United Nations; 2024.
17.   Shankar M, Hartner A-M, Arnold CR, Gayawan E, Kang H, Kim J-H, et al.
Mathematical modelling to inform outbreak response vaccination. 2024.
18.   Immunization Agenda 2030 Scorecard SP5.1: Outbreak Detection and
Response;[accessed 2024-11-25]. [cited 2024-11-25]. Available from:
https://scorecard.immunizationagenda2030.org/sp5.1.
19.   Grais RF, Conlan AJ, Ferrari MJ, Djibo A, Le Menach A, Bjornstad ON, et al. Time is of
the essence: exploring a measles outbreak response vaccination in Niamey, Niger. J R Soc
Interface. 2008;5(18):67-74.
20.   Yagci Sokat K, Armbruster B. Modeling and Controlling Epidemic Outbreaks: The Role
of Population Size, Model Heterogeneity and Fast Response in the Case of Measles.
2020;8(11):1892.
21.   Kellerborg K, Brouwer W, van Baal P. Costs and benefits of early response in the
Ebola virus disease outbreak in Sierra Leone. Cost Eff Resour Alloc. 2020;18:13.
22.   Trotter CL, Cibrelus L, Fernandez K, Lingani C, Ronveaux O, Stuart JM. Response
thresholds for epidemic meningitis in sub-Saharan Africa following the introduction of
MenAfriVac(R). Vaccine. 2015;33(46):6212-7.
23.   Cooper LV, Ronveaux O, Fernandez K, Lingani C, Goumbi K, Ihekweazu C, et al.
Spatiotemporal Analysis of Serogroup C Meningococcal Meningitis Spread in Niger and
Nigeria and Implications for Epidemic Response. J Infect Dis. 2019;220(220 Suppl 4):S244-
S52.
24.   Ferrari MJ, Fermon F, Nackers F, Llosa A, Magone C, Grais RF. Time is (still) of the
essence: quantifying the impact of emergency meningitis vaccination response in Katsina
State, Nigeria. Int Health. 2014;6(4):282-90.
25.   Zhao S, Stone L, Gao D, He D. Modelling the large-scale yellow fever outbreak in
Luanda, Angola, and the impact of vaccination. PLoS Negl Trop Dis. 2018;12(1):e0006158.
26.   Kalkowska DA, Pallansch MA, Wassilak SGF, Cochi SL, Thompson KM. Serotype 2 oral
poliovirus vaccine (OPV2) choices and the consequences of delaying outbreak response.
Vaccine. 2023;41 Suppl 1(Suppl 1):A136-A41.

27. Laydon DJ, Cauchemez S, Hinsley WR, Bhatt S, Ferguson NM. Impact of proactive and
reactive vaccination strategies for health-care workers against MERS-CoV: a mathematical
modelling study. Lancet Glob Health. 2023;11(5):e759-e69.
28. Reyburn R, Deen JL, Grais RF, Bhattacharya SK, Sur D, Lopez AL, et al. The case for
reactive mass oral cholera vaccinations. PLoS Negl Trop Dis. 2011;5(1):e952.
29. Azam JM, Saitta B, Bonner K, Ferrari MJ, Pulliam JRC. Modelling the relative benefits
of using the measles vaccine outside cold chain for outbreak response. Vaccine.
2021;39(40):5845-53.
30. Delport D, Muellenmeister A, MacKechnie G, Vaccher S, Mengistu T, Hogan D, et al.
Estimating the historical impact of outbreak response immunization programmes across 210
outbreaks in low and middle-income countries. Zenodo; 2024.
31. Starsim github repository.
32. Starsim: Bill and Melinda Gates Foundation;[accessed 13/05/2025]. Available from:
https://docs.idmod.org/projects/starsim/en/latest/.
33. Meningococcal vaccines: WHO position paper on the use of multivalent
meningococcal conjugate vaccines in countries of the African meningitis belt, January 2024:
World Health Organization; 2024;[accessed 15/10/2025]. Available from:
https://www.who.int/publications/i/item/who-wer9901-02-1-10.
34. Geddes L. Five things you need to know about the world's latest weapon against
bacterial meningitis: Gavi the Vaccine Alliance; 2024;[accessed 29/10/2025]. Available from:
https://www.gavi.org/vaccineswork/five-things-you-need-know-about-worlds-latest-
weapon-against-bacterial-meningitis.
35. In world first, Nigeria introduces new 5-in-1 vaccine against meningitis: World Health
Organization; 2024;[accessed 15/10/2025]. Available from:
https://www.who.int/news/item/12-04-2024-in-world-first--nigeria-introduces-new-5-in-1-
vaccine-against-meningitis.
36. Immunization coverage: World Health Organization; 2025;[accessed 29/10/2025].
Available from: https://www.who.int/news-room/fact-sheets/detail/immunization-
coverage.
37. Bosis S, Mayer A, Esposito S. Meningococcal disease in childhood: epidemiology,
clinical features and prevention. J Prev Med Hyg. 2015;56(3):E121-4.
38. Blakebrough IS, Greenwood BM, Whittle HC, Bradley AK, Gilles HM. The
epidemiology of infections due to Neisseria meningitidis and Neisseria lactamica in a
northern Nigerian community. J Infect Dis. 1982;146(5):626-37.
39. Basta NE, Berthe A, Keita M, Onwuchekwa U, Tamboura B, Traore A, et al.
Meningococcal carriage within households in the African meningitis belt: A longitudinal pilot
study. J Infect. 2018;76(2):140-8.
40. MenAfriCar C. Household transmission of Neisseria meningitidis in the African
meningitis belt: a longitudinal cohort study. Lancet Glob Health. 2016;4(12):e989-e95.
41. Karachaliou A, Conlan AJ, Preziosi MP, Trotter CL. Modeling Long-term Vaccination
Strategies With MenAfriVac in the African Meningitis Belt. Clin Infect Dis. 2015;61 Suppl
5(Suppl 5):S594-600.
42. Sharew A, Bodilsen J, Hansen BR, Nielsen H, Brandt CT. The cause of death in
bacterial meningitis. BMC Infect Dis. 2020;20(1):182.
43. Rivero-Calle I, Raguindin PF, Pardo-Seco J, Martinon-Torres F. Risk Analysis by Age on
the Burden of Meningococcal Disease in Spain. Vaccines (Basel). 2022;10(4).

44.   Yaesoubi R, Trotter C, Colijn C, Yaesoubi M, Colombini A, Resch S, et al. The cost-
effectiveness of alternative vaccination strategies for polyvalent meningococcal vaccines in
Burkina Faso: A transmission dynamic modeling study. PLoS Med. 2018;15(1):e1002495.
45.   Pace D, Pollard AJ. Meningococcal disease: clinical presentation and sequelae.
Vaccine. 2012;30 Suppl 2:B3-9.
46.   Rouphael NG, Stephens DS. Neisseria meningitidis: biology, microbiology, and
epidemiology. Methods Mol Biol. 2012;799:1-20.
47.   Agier L, Deroubaix A, Martiny N, Yaka P, Djibo A, Broutin H. Seasonality of meningitis
in Africa and climate forcing: aerosols stand out. J R Soc Interface. 2013;10(79):20120814.
48.   M.J. Fresnadillo Martínez EGM, E. García Sánchez, J.E. García Sánchez, A. Martín del
Rey, G. Rodríguez Sánchez. A Mathematical Model to Study the Meningococcal Meningitis.
Procedia Computer Science. 2013;18:2492-5.
49.   Daugla DM, Gami JP, Gamougam K, Naibei N, Mbainadji L, Narbe M, et al. Effect of a
serogroup A meningococcal conjugate vaccine (PsA-TT) on serogroup A meningococcal
meningitis and carriage in Chad: a community study [corrected]. Lancet. 2014;383(9911):40-
7.
50.   Hadley L, Karachaliou Prasinou A, Christensen H, Ramsay M, Trotter C. Modelling the
impact of COVID-19 and routine MenACWY vaccination on meningococcal carriage and
disease in the UK. Epidemiol Infect. 2023;151:e98.
51.   Azman AS, Rudolph KE, Cummings DA, Lessler J. The incubation period of cholera: a
systematic review. J Infect. 2013;66(5):432-8.
52.   Cholera: World Health Organization; 2023;[accessed 18/01/2024]. Available from:
https://www.who.int/news-room/fact-sheets/detail/cholera.
53.   Fung IC. Cholera transmission dynamic models for public health practitioners. Emerg
Themes Epidemiol. 2014;11(1):1.
54.   Somboonwit C, Menezes LJ, Holt DA, Sinnott JT, Shapshak P. Current views and
challenges on clinical cholera. Bioinformation. 2017;13(12):405-9.
55.   Hartley DM, Morris JG, Jr., Smith DL. Hyperinfectivity: a critical element in the ability
of V. cholerae to cause epidemics? PLoS Med. 2006;3(1):e7.
56.   Wang J. Mathematical Models for Cholera Dynamics-A Review. Microorganisms.
2022;10(12).
57.   Mukandavire Z, Liao S, Wang J, Gaff H, Smith DL, Morris JG, Jr. Estimating the
reproductive numbers for the 2008-2009 cholera outbreaks in Zimbabwe. Proc Natl Acad Sci
U S A. 2011;108(21):8767-72.
58.   Fournier JM, Quilici ML. [Cholera]. Presse Med. 2007;36(4 Pt 2):727-39.
59.   Jackson BR, Talkington DF, Pruckler JM, Fouche MDB, Lafosse E, Nygren B, et al.
Seroepidemiologic survey of epidemic cholera in Haiti to assess spectrum of illness and risk
factors for severe disease. Am J Trop Med Hyg. 2013;89(4):654-64.
60.   Nelson EJ, Harris JB, Morris JG, Jr., Calderwood SB, Camilli A. Cholera transmission:
the host, pathogen and bacteriophage dynamic. Nat Rev Microbiol. 2009;7(10):693-702.
61.   Malembaka EB, Bugeme PM, Hutchins C, Xu H, Hulse JD, Demby MN, et al.
Effectiveness of one dose of killed oral cholera vaccine in an endemic community in the
Democratic Republic of the Congo: a matched case-control study. Lancet Infect Dis. 2024.
62.   Song KR, Lim JK, Park SE, Saluja T, Cho SI, Wartel TA, et al. Oral Cholera Vaccine
Efficacy and Effectiveness. Vaccines (Basel). 2021;9(12).

63.   Measles Rubeola: Center for Disease Control and Prevention (CDC); 2021;[accessed
08/01/2024]. Available from: https://www.cdc.gov/measles/downloads/Measles-fact-sheet-
508.pdf.
64.   Gastañaduy PA, Banerjee E, DeBolt C, Bravo-Alcántara P, Samad SA, Pastor D, et al.
Public health responses during measles outbreaks in elimination settings: Strategies and
challenges. (2164-554X (Electronic)).
65.   Nic Lochlainn LM, de Gier B, van der Maas N, Strebel PM, Goodman T, van Binnendijk
RS, et al. Immunogenicity, effectiveness, and safety of measles vaccination in infants
younger than 9 months: a systematic review and meta-analysis. Lancet. 2019;19(11):1235-
45.
66.   Leuridan E, Hens N, Hutse V, Ieven M, Aerts M, Van Damme P. Early waning of
maternal measles antibodies in era of measles elimination: longitudinal study. BMJ.
2010;340(7756).
67.   Johansson MA, Vasconcelos PFC, Staples JE. The whole iceberg: estimating the
incidence of yellow fever virus infection from the number of severe cases. Transactions of
The Royal Society of Tropical Medicine and Hygiene. 2014;108(8):482-7.
68.   Yellow fever: World Health Organization; 2023;[accessed 02/02/2024]. Available
from: https://www.who.int/news-room/fact-sheets/detail/yellow-fever.
69.   Jean K, Donnelly CA, Ferguson NM, Garske T. A Meta-Analysis of Serological
Response Associated with Yellow Fever Vaccination. The American Journal of Tropical
Medicine and Hygiene. 2016;95(6):1435-9.
70.   Prem K, Cook AR, Jit M. Projecting social contact matrices in 152 countries using
contact surveys and demographic data. PLoS Comput Biol. 2017;13(9):e1005697.
71.   World Population Prospects 2022, Online Edition: United Nations, Department of
Economic and Social Affairs, Population Division (2022); 2022;[accessed 03/12/2023].
Available from: https://population.un.org/wpp/Download/Standard/Population/.
72.   Measles Outbreak Toolkit: World Health Organization; 2022;[accessed 08/01/2024].
Available from: https://www.who.int/emergencies/outbreak-toolkit/disease-outbreak-
toolboxes/measles-outbreak-toolbox.
73.   Cholera Outbreak Toolbox: World Health Organization; 2024. [cited 24/10/2025].
Available from: https://www.who.int/emergencies/outbreak-toolkit/disease-outbreak-
toolboxes/cholera-outbreak-toolbox.
74.   Yellow Fever Outbreak Toolbox: World Health Organization; 2024;[accessed
28/03/2024]. Available from: https://www.who.int/emergencies/outbreak-toolkit/disease-
outbreak-toolboxes/yellow-fever-outbreak-toolbox.
75.   Revised guidance on meningitis outbreak response in sub-Saharan Africa. Wkly
Epidemiol Rec. 2014;89(51-52):580-6.
76.   Abeysuriya RG, Delport D, Stuart RM, Sacks-Davis R, Kerr CC, Mistry D, et al.
Preventing a cluster from becoming a new wave in settings with zero community COVID-19
cases. BMC Infect Dis. 2022;22(1):232.
77.   Delport D, Sanderson B, Sacks-Davis R, Vaccher S, Dalton M, Martin-Hughes R, et al.
A Framework for Assessing the Impact of Outbreak Response Immunization Programs.
Diseases. 2024;12(4).
78.   Ferrari AJ, Santomauro DF, Aali A, Abate YH, Abbafati C, Abbastabar H, et al. Global
incidence, prevalence, years lived with disability (YLDs), disability-adjusted life-years
(DALYs), and healthy life expectancy (HALE) for 371 diseases and injuries in 204 countries

and territories and 811 subnational locations, 1990-2021: a systematic analysis for the
Global Burden of Disease Study 2021. The Lancet. 2024;403(10440):2133-61.
79.   Hogan D, Gupta A. Why Reaching Zero-Dose Children Holds the Key to Achieving the
Sustainable Development Goals. LID - 10.3390/vaccines11040781 [doi] LID - 781. (2076-
393X (Print)).
80.   Collaborators GBDVC. Global, regional, and national trends in routine childhood
vaccination coverage from 1980 to 2023 with forecasts to 2030: a systematic analysis for the
Global Burden of Disease Study 2023. Lancet. 2025.
81.   Mora C, McKenzie T, Gaw IM, Dean JM, von Hammerstein H, Knudson TA, et al. Over
half of known human pathogenic diseases can be aggravated by climate change. Nat Clim
Chang. 2022;12(9):869-75.
82.   Houlihan CF, Whitworth JA. Outbreak science: recent progress in the detection and
response to outbreaks of infectious diseases. Clin Med (Lond). 2019;19(2):140-4.
83.   Kluberg SA, Mekaru SR, McIver DJ, Madoff LC, Crawley AW, Smolinski MS, et al.
Global Capacity for Emerging Infectious Disease Detection, 1996-2014. Emerg Infect Dis.
2016;22(10):E1-6.
84.   Gaythorpe KA-O, Abbas K, Huber JA-O, Karachaliou A, Thakkar N, Woodruff KA-O, et
al. Impact of COVID-19-related disruptions to measles, meningococcal A, and yellow fever
vaccination in 10 countries. LID - 10.7554/eLife.67023 [doi] LID - e67023. (2050-084X
(Electronic)).