# Genetic-epigenetic interactions (meQTLs) in orofacial clefts etiology

Machado-Paula, LA[1], Romanowska, J[2]; Lie, RT[2]; Hovey, L[1], Doolittle, B[1]; Awotoye, W[1], Dunlay, L[1], Xie, XJ[1]; Zeng, E[1]; Butali A[1,3]; Marazita, ML[4], Murray, JC[3]; Moreno-Uribe, LM[1], *Petrin, AL[1]

1. University of Iowa College of Dentistry and Dental Clinics, Iowa City, IA, USA
2. University of Bergen, Bergen, Norway
3. University of Iowa Carver College of Medicine, Iowa City, IA, USA
4. University of Pittsburgh, Pittsburgh, PA, USA

## ABSTRACT

**Objectives**: Nonsyndromic orofacial clefts (OFCs) etiology involves multiple genetic and environmental factors with over 60 identified risk loci; however, they account for only a minority of the estimated risk. Epigenetic factors such as differential DNA methylation (DNAm) are also associated with OFCs risk and can alter risk for different cleft types and modify OFCs penetrance. DNAm is a covalent addition of a methyl (CH3) group to the nucleotide cytosine that can lead to changes in expression of the targeted gene. DNAm can be affected by environmental influences and genetic variation via methylation quantitative loci (meQTLs). We hypothesize that aberrant DNAm and the resulting alterations in gene expression play a key role in the etiology of OFCs, and that certain common genetic variants that affect OFCs risk do so by influencing DNAm. **Methods:** We used genotype from 10 cleft-associated SNPs and genome-wide DNA methylation data (Illumina 450K array) for 409 cases with OFCs and 456 controls and identified 23 cleft-associated meQTLs. We then used an independent cohort of 362 cleft-discordant sib pairs for replication. We used methylation-specific qPCR to measure methylation levels of each CpG site and combined genotypic and methylation data for an interaction analysis of each SNP-CpG pair using the R package MatrixeQTL in a linear model. We also performed a Paired T-test to analyze differences in DNA methylation between each member of the sibling pairs.
**Results:** We replicated 9 meQTLs, showing interactions between rs13041247 (*MAFB*) - cg18347630 (*PLCG1*) (P=0.04); rs227731 (*NOG*) - cg08592707 *(PPM1E)* (P=0.01); rs227731 (*NOG*) - cg10303698 (*CUEDC1*) (P=0.001); rs3758249 (*FOXE1*) - cg20308679 (*FRZB*) (P=0.04); rs8001641 (*SPRY2*) - cg19191560 (*LGR4*) (P=0.04); rs987525(8q24) - cg16561172(*MYC*) (P=0.00000963); rs7590268(*THADA*) - cg06873343 (*TTYH3*) (P=0.04); rs7078160 (*VAX1*) - cg09487139 (P=0.05); rs560426 (*ABCA4/ARHGAP29*) - cg25196715 (*ABCA4/ARHGAP29*) (P=0,03). Paired T-test showed significant differences for cg06873343 (*TTYH3*) (P=0.04); cg17103269 (*LPIN3*) (P=0.002), and cg19191560 (*LGR4*) (P=0.05). **Conclusions:** Our results confirm previous evidence that some of the common non-coding variants detected through GWAS studies can influence the risk of OFCs via epigenetic mechanisms, such as DNAm, which can ultimately affect and regulate gene expression. Given the large prevalence of non-coding SNPs in most OFCs genome wide association studies, our findings can potentially address major knowledge gaps, like missing heritability, reduced penetrance, and variable expressivity associated with OFCs phenotypes.

**\*Corresponding author:**
Aline L. Petrin, College of Dentistry and Dental Clinics, University of Iowa, Iowa City, IA, 52242, USA, aline-petrin@uiowa.edu

## Introduction

Nonsyndromic orofacial clefts (OFCs) are the most common craniofacial birth defects in humans, affecting approximately 1 in 700 individuals worldwide [1]. OFCs are sub-divided into cleft lip occurring with or without cleft palate (CL/P) and isolated cleft palate (CPO) [2, 3]. The identification of causal factors for OFCs has been a challenge due to its complex etiology arising from multiple genetic and environmental contributions. Family history is observed in about 23% of cases showing a strong genetic component; however, monozygotic twins have a concordance rate of about 50% [4] and amongst those that are concordant there are often discernable differences in OFCs phenotypic expression indicating that other factors are at play beyond genetic risk.

Together, GWAS [5-12], GWAS meta-analyses [13, 14], linkage [15] and replications studies have identified over 60 risk loci for OFCs; however, they account for a minority of the estimated risk, and many of them reside in non-coding regions with unclear functional relevance so far. Thus, the challenge of translating statistical associations into biological mechanisms still limits our knowledge on OFCs etiology, and thus to gain further insight it is necessary to expand beyond the single-omics analyses.

Epigenetic modification, such as DNA methylation (DNAm) is a likely mechanism through which environmental factors or genetic variation can modify gene expression. DNAm consists of the covalent addition of a methyl (CH3) group to a nucleotide cytosine followed by a guanine, a CpG site, by a family of enzymes called methyltransferases (DNMTs) [16-18]. Changes in DNAm can lead to differential transcriptional activity of a target gene and have important roles in multiple processes during human development and throughout life such as genomic imprinting, maintenance of X-inactivation and genomic stability [18-22].

Multiple studies have shown that genetic variants can impact a significant proportion of the human DNA methylome [23-33]. Some studies indicate that epigenetic factors such as differential DNA methylation (DNAm) are associated with OFCs risk [34-43], also that differentially methylated regions tend to cluster around gene pathways previously linked to palatogenesis, and that some of these differential DNAm regions seem to be influenced by genetic variation [34, 37, 44, 45]. However, the overall role of epigenetics in OFCs etiology remains underexplored [46].

Individual genotypes at methylation quantitative trait loci (meQTLs) can modify DNA methylation patterns. meQTLs can be divided into *cis* or *trans* based on the proximity of the genetic variant to the CpG site. Cis-meQTLs are genetic variants near to or proximal to the target CpG site, and trans-meQTLs are separated by one or more Mbp from the target CpG or located on different chromosomes [47-49]. Studies on blood DNA and commercial methylation arrays (Illumina Infinium MethylationEPIC array) show that 34% to 45% of the CpGs [24, 50-53] are influenced by SNPs, with 98% of the effects cis-acting or within 1Mb of the tested CpG sites [33]. Additionally, both meQTLS SNPs and target CpGs are frequently found in enhancers [33, 54-56]. Literature has shown meQTL SNPs in various tissues like brain [57, 58], adipose [59], and buccal tissue samples [60]. Further, studies have shown that there is a significant overlap of meQTLs between ancestral groups, developmental stages and tissue types [61].

Differential DNAm may alter risk for different cleft types and modify OFCs penetrance [34]. Our group recently showed that differential methylation can contribute to phenotypic variability observed in patients with Van der Woude syndrome, including monozygotic twins with discordant phenotypes despite carrying the same causal mutation in the *IRF6* gene [27, 29]. These findings support the role of DNA methylation as a modifier influencing phenotypic severity.

Therefore, analyses that combine known genetic risk factors with epigenetic data to explore methylation quantitative loci (meQTLs) function significantly contribute to understanding such loci by addressing the causal alleles, their pathogenic consequences, and the biological mechanisms through which they influence OFCs risk.

Discordant sib pairs are valuable cohorts for mapping complex human traits. The use of discordant sibling pairs in methylation studies offers several advantages: siblings share a significant portion of their genetic makeup, which helps control for genetic variation and isolates environmental influences on methylation. Additionally, siblings often experience similar early-life environments, aiding in the study of

environmental impacts on DNA methylation[62]. Comparing affected and unaffected siblings (discordant pairs) can reveal disease-related methylation changes, and the reduced variability from genetic and environmental similarities enhances the statistical power and reliability of the study.

Based on an analysis of subset of cases and controls from previous studies [63, 64], we found that genotypes of 10 cleft-associated SNPs were highly correlated with the methylation levels of certain CpG sites, indicating that these SNPs function as methylation quantitative trait loci (meQTLs). This genetic-epigenetic association was replicated using a unique sample of same-sex siblings discordant for OFCs.

**Methods**

Samples

The first cohort was composed of a subset of subset of unrelated cases with cleft lip with or without cleft palate and unaffected controls from a previous published studies [63, 64]. We used genotype from 10 cleft-associated SNPs and genome-wide DNA methylation data (obtained with Illumina 450K array) for 409 cases with OFCs and 456 controls, all with DNA from blood. The second independent cohort was composed of 362 pairs of same-sex siblings, discordant for cleft lip with or without cleft palate (one sibling affected-case, one unaffected-control); of these, 166 pairs had DNA extracted from blood and 196 pairs had DNA extracted from saliva from saliva. All samples were obtained in accordance with prior study protocols, following their respective approval by the University of Iowa and locals Institutional Review Boards (IRBs). Informed consent was provided by patients, parents or guardians prior obtaining samples and clinical information.

Genotyping

As mentioned above, for cohort 1, we extracted the genetic data for the selected 10 SNPs (top loci for OFCs from previous GWAS and candidate gene studies) (Table 1) from a previous published studies [63] [64]. All the details of the data collection, quality control, and preprocessing are available in the Appendix of the cited work. For cohort 2, we genotyped the 10 SNPs in the discordant sib-pairs using Taqman assays on a Fluidigm (Fluidigm Corp., South San Francisco, CA, USA) nanofluidic platform. Genotype calling was done using the Fluidigm SNP genotyping software, version 4.1.2, with the default settings. The quality check included setting the confidence threshold to 65% for the genotype calling algorithm and then visually checking all the genotyping plots.

Bisulfite conversion and DNA methylation profiling

All samples were bisulfite converted prior measuring DNA methylation levels. Briefly, DNA quality was assessed QubitTM dsDNA High Sensitivity Range Assay Kit (Thermo Fisher Scientific) and 1.5% agarose gel. After quantification of each sample, 500 ng of each genomic DNA sample was submitted to bisulfite conversion using the EZ DNA Methylation Kit (Zymo Research) according to manufacturer's protocol.

For cohort 1, DNA methylation levels were obtained using the Illumina Human Methylation 450K BeadChip. DNA methylation level was then estimated at 485,577 CpG sites and the R package illuminaio (ver. 0.18)3 was used to extract the raw probe intensity values. These raw values were preprocessed using ENmix R package4. The following criteria were used to identify low-quality samples: (1) average intensity value across internal control probes less than 5500, (2) more than 5% of CpG probes having low-quality data (Illumina detection P value > $10^{-6}$, read from less than 3 beads, or outlier value for the probe in the dataset), and (3) clear outliers based on visual inspection of a density plot of total intensity. Next, the low quality CpG probes were defined as follows: (1) more than 5% low-quality data; (2) common SNP within the probe's sequence (minor allele frequency $\geq$ 0:05 in Europeans based on 1000 Genomes Project data), or probes mapping to multiple genomic locations, or CpGs on X or Y chromosomes; (3) CpGs with multiple mode distributions identified with ENmix. More details about the filtering steps can be found in the cited work[63].

For cohort 2, after bisulfite conversion, genomic DNA samples were amplified by fluorescence-based, real time quantitative PCR (Methylight) [65]. Quantitative MethyLight assays (e.g., EpiTect

MethyLight Assays), consists of two probes, one methylation specific, and the other nonmethylation specific, which be used in a single real-time PCR reaction. This enables highly accurate quantitative methylation analysis, due to the simultaneous detection of methylated and unmethylated DNA. We used locus-specific PCR primers flanking an oligonucleotide probe with a 5' fluorescent reporter dye to detect methylated (SUN) and unmethylated (FAM) alleles and a 3' quencher dye (3IABkFQ). The primers and fluorescent probes were designed against bisulfite-converted DNA sequence and quantitative information was obtained in real time. Serial dilutions of the EpiTect Control DNAs (Qiagen) were included on each plate to generate a standard curve and to verify plate to plate consistency. The PCR amplification was performed in a 384-well plate format and each sample was ran in duplicate. EpiTect MethyLight assays enable the direct quantification of the methylation degree in a sample by taking the threshold cycles ($C_T$) determined in the SUN channel with the probe detecting methylated DNA or in the FAM channel with the probe detecting unmethylated DNA. Ten nanograms of bisulfite converted, methylated and unmethylated human control DNA, or defined mixtures of both DNAs were used for methylation quantification. The methylation degree of each sample was calculated from the average of $C_T$ values in SUN ($C_{T(CG)}$) and FAM ($C_{T(TG)}$) channel, obtained in quantitative real-time PCR using the formula described in [66]: $C_{meth} = 100/[1+2^{(C_{T(CG)} - C_{T(TG)})}]\%$; where $C_{T(CG)}$ is the methylated signal obtained by the threshold cycle of the CG reporter (SUN channel), and $C_{T(TG)}$ is the unmethylated signal obtained by the threshold cycle of the CG reporter (FAM channel).

## Statistical analysis

The meQTL analyses were performed using the R package MatrixEQTL version 2.1.1 [67], which was designed to perform ultra-fast calculations of associations between SNPs and gene expression data (eQTLs). Due to similarity between the gene expression and methylation data (both are numeric big matrices), the method has been widely applied to methylation quantitative loci as well [48, 68-74]. In addition to replicating the meQTLs, we used data obtained from GeneHancer [75]— a regulatory element database containing genomic coordinates of known enhancer and promoter elements, including those active during craniofacial development [76, 77]—to (1) investigate whether the validated CpG sites overlapped with any regulatory elements and (2) prioritize the most likely CpG-target genes in epithelial and mesenchymal tissues.

For cohort 1, all 10 SNPs were analyzed against the 407,513 CpGs that passed QC. Linear regression was used to assess the relationship between each CpG methylation and each SNP genotype (0, 1, or 2 copies of the minor allele) using cleft status (affected or unaffected) as covariate. The linear model analyzes the effect that the genotype (additive effect) has on the methylation based on the cleft status (case or control). It tests for the significance of the genotype and the cleft status over the methylation level. The linear model was tested separately for each of the 3 groups, blood, saliva and combined.

Matrix eQTL performs a separate test for each SNP-CpG pair and corrects for multiple comparisons by

**Table 1: Most significant meQTLs resulting from case-control study**

| SNP | SNP gene | CpG | CpG target gene | p-value |
|---|---|---|---|---|
| rs227731 | NOG | cg08592707 | SKA2 | 1.59E-05 |
| rs4752028 | VAX1 | cg08319991 | UCHL1 | 2.83E-08 |
| rs7590268 | THADA | cg06873343 | N/A | 7.72E-08 |
| rs7078160 | VAX1 | cg08319991 | UCHL1 | 1.11E-07 |
| rs3758249 | FOXE1 | cg20308679 | FRZB | 1.29E-07 |
| rs4752028 | VAX1 | cg11876012 | AFAP1L2 | 1.41E-07 |
| rs8001641 | SPRY2 | cg19191560 | LGR4 | 2.88E-08 |
| rs1873147 | TPM1 | cg11936410 | TPM1 | 5.41E-04 |
| rs560426 | ABCA4/ARHGAP29 | cg25196715 | BCAR3/MIR760 | 6.72E-04 |
| rs987525 | 8q24 | cg16561172 | MYC | 7.04E-04 |
| rs560426 | ABCA4/ARHGAP29 | cg00405232 | ABCA4 | 7.93E-04 |
| rs227731 | NOG | cg01964121 | PPM1E | 8.11E-04 |
| rs227731 | NOG | cg10303698 | CUEDC1 | 8.74E-04 |
| rs1873147 | TPM1 | cg24483493 | HERC1 | 9.14E-04 |
| rs4752028 | VAX1 | cg08329473 | POLR3K | 9.29E-04 |
| rs13041247 | MAFB | cg18347630 | CHD6 | 1.02E-03 |
| rs7078160 | VAX1 | cg08329473 | AFAP1L2 | 1.06E-03 |
| rs1873147 | TPM1 | cg16659880 | TPM1 | 1.11E-03 |
| rs7078160 | VAX1 | cg09487139 | N/A | 1.38E-03 |
| rs13041247 | MAFB | cg00514723 | LPIN3 | 1.39E-03 |
| rs742071 | PAX7 | cg20940024 | N/A | 1.56E-03 |
| rs1873147 | TPM1 | cg19631779 | TPM1 | 1.67E-03 |
| rs13041247 | MAFB | cg17103269 | PLCG1 | 1.78E-03 |

calculating Benjamini-Hochberg FDR. From this first analysis, we selected 23 SNP-CPG associations with p≤ 0.002 and FDR<0.4 for replication with an independent cohort (cohort 2).

For cohort 2, we used MatrixEQTL to run each SNP-CpG pair in the 362 pairs of siblings discordant for OFCs, following the same parameters above. Further, to test for associations between DNA methylation levels independent of SNP effect within sibling pairs, we used paired Student's T-test analyses for each tissue subgroups (blood, saliva and combined).

<u>Bioinformatic analysis</u>

We used bioinformatic methods to prior regions and select the most likely target gene for each CpG site. We obtained data from GeneHancer—a regulatory element database containing genomic coordinates of known enhancer and promoter elements [75], including those active during craniofacial development [76, 77]— and compared the coordinates to our CpG sites to (1) investigate whether the validated CpG sites overlapped with any regulatory elements and (2) prioritize the most likely CpG-target genes in relevant tissues (e.g., epithelial and mesenchymal tissues) based on GeneHancer *in silico* prediction tools.

**Results**

The analysis with the first cohort (409 cases, 456 controls) correlated the genotype of 10 cleft-associated SNPs [5, 8, 63, 64, 78-82] with genome-wide DNAm patterns, identifying 23 significant SNP-CpG interactions (meQTLs). All 10 SNPs are non-coding (and may be located near/in regulatory elements)— supporting the need for novel approaches to elucidate their pathological consequence—and several have been identified as meQTLs in other tissues and phenotypes (Table 1, SNPs in *Italics*) [83]. We then followed-up with an independent cohort of 362 discordant sib pairs and replicated 9 out of the 23 cleft-associated SNP-CpG associations. In Table 1, we show the result of the first analysis for the case-control cohort and the best replication result obtained with the sib pairs cohort for each pair of SNP-CpG associations.

Table 2 shows results for the SNP-CpG linear model for each tissue group. For blood analysis we detected interaction between

**Table 2: Replicated SNP-CpG associations (meQTLs) from linear analysis in all groups**

| Sample | Affected /Unaffected | SNPs | SNP gene | CpG | CpG gene | pvalue |
|---|---|---|---|---|---|---|
| All | 353/353 | rs13041247 | | cg18347630 | | 2.13E-01 |
| Blood | 160/160 | rs13041247 | MAFB | cg18347630 | PLCG1 | **3.93E-02** |
| Saliva | 193/193 | rs13041247 | | cg18347630 | | 3.38E-01 |
| All | 354/354 | rs227731 | | cg08592707 | | **1.54E-02** |
| Blood | 163/163 | rs227731 | NOG | cg08592707 | PPM1E | 3.66E-01 |
| Saliva | 191/191 | rs227731 | | cg08592707 | | 5.34E-01 |
| All | 300/300 | rs227731 | | cg10303698 | | **1.50E-03** |
| Blood | 110/110 | rs227731 | NOG | cg10303698 | CUEDC1 | 5.08E-01 |
| Saliva | 190/190 | rs227731 | | cg10303698 | | **5.95E-02** |
| All | 318/318 | rs3758249 | | cg20308679 | | 1.10E-01 |
| Blood | 131/131 | rs3758249 | FOXE1 | cg20308679 | FRZB | 2.92E-01 |
| Saliva | 187/187 | rs3758249 | | cg20308679 | | **3.80E-02** |
| All | 340/340 | rs8001641 | | cg19191560 | | **4.70E-02** |
| Blood | 158/158 | rs8001641 | SRPY2 | cg19191560 | LGR4 | **9.33E-02** |
| Saliva | 182/182 | rs8001641 | | cg19191560 | | **3.72E-02** |
| All | 305/305 | rs987525 | | cg16561172 | | **2.78E-05** |
| Blood | 138/138 | rs987525 | 8q24 | cg16561172 | MYC | 6.12E-01 |
| Saliva | 167/167 | rs987525 | | cg16561172 | | **9.63E-06** |
| All | 356/356 | rs7590268 | | cg06873343 | | 3.08E-01 |
| Blood | 165/165 | rs7590268 | THADA | cg06873343 | TTYH3 | **3.91E-02** |
| Saliva | 191/191 | rs7590268 | | cg06873343 | | 9.14E-01 |
| All | 333/333 | rs7078160 | | cg09487139 | | 7.30E-01 |
| Blood | 157/157 | rs7078160 | VAX1 | cg09487139 | N/A | **4.60E-02** |
| Saliva | 176/176 | rs7078160 | | cg09487139 | | 2.75E-01 |
| All | 297/297 | rs560426 | | cg25196715 | | **3.16E-02** |
| Blood | 126/126 | rs560426 | ABCA4/ ARHGAP29 | cg25196715 | ABCA4/ ARHGAP29 | 3.88E-01 |
| Saliva | 171/171 | rs560426 | | cg25196715 | | **3.81E-02** |

p-values in bold are boderline significant (p<0.09); FDR=False Discovery Rate

Beta value ranges: 0-0.2 = fully unmethylated; 0.21-0.60 = partially methylated; 0.61-1.00 = fully methylated

rs13041247/*MAFB* - cg18347630/*PLCG1* (P=0.04); rs7590268/*THADA* - cg06873343/*TTYH3* (P=0.04); rs7078160/*VAX1* - cg09487139 (P=0.05). Analysis for saliva revealed significant associations between rs3758249/*FOXE1* - cg20308679/*FRZB/* (P=0.04). The combined analysis (blood and saliva together) showed associations between rs227731/*NOG* - cg08592707/*PPM1E* (P=0.01) and rs227731/*NOG* - cg10303698/*CUEDC1* (P=0.001). Three meQTLs had significant associations in both saliva and combined analysis (blood and saliva), being rs8001641/*SPRY2* - cg19191560/*LGR4* (P=0.04-saliva, P=0.05-all); rs987525/8q24 - cg16561172/*MYC* (P= P=9.6E-06-saliva, P= 2.8E-05-all); rs560426*ABCA4/ARHGAP29* - cg25196715/*ABCA4/ARHGAP29* (P=0.04-saliva, P=0.03-all).

For the 10 CpG sites analyzed, the differential methylation analysis (independent of SNP effect) between sibling pairs and subdivided by groups through Paired T-test (Table 3) showed significant results for cg06873343 (*TTYH3*) for blood (P=0.04), saliva (P=0.05) and all (P= 0.004); cg17103269 (*LPIN3*) (P=0.002 – blood) and cg19191560 (*LGR4*) (P=0.05 – all). The most significant result was found for the cg17103269 (*LPIN3*), followed by cg06873343 (*TTYH3*).We had borderline results for cg19191560 (*LGR4*) (P=0.06-blood) and cg25196715 (*ABCA4/ARHGAP29*) (P=0.07-blood).

**Table 3: Differential methylation levels between sibling pairs and subdivided by groups**

| CpG site | Chr position | Closest Gene | # of sib pairs | Group | p-value |
|---|---|---|---|---|---|
| cg06873343 | chr7:2,667,608 | *TTYH3* | 358 | All | **0.004** |
| | | | 165 | Blood | **0.04** |
| | | | 193 | Saliva | **0.05** |
| cg17103269 | chr20:39,972,305 | *LPIN3* | 360 | All | 0.25 |
| | | | 166 | Blood | **0.0017** |
| | | | 194 | Saliva | 0.87 |
| cg19191560 | chr11:27,492,758 | *LGR4* | 340 | All | **0.05** |
| | | | 158 | Blood | **0.06** |
| | | | 182 | Saliva | 0.26 |
| cg25196715 | chr1:94,489,792 | *ABCA4/ARHGAP29* | 297 | All | 0.13 |
| | | | 126 | Blood | **0.07** |
| | | | 171 | Saliva | 0.56 |

p-values in bold are borderline significant (p<0.09)

## Discussion

The integration of emerging epigenetic data for orofacial clefts with existing GWAS hits may elucidate functional mechanisms that underlie statistical significance. Moreover, enrichment analysis of epigenetic data also has the potential to reveal pathways not implicated by genetic findings. Our hypothesis was that certain common genetic variants that affect OFCs risk do so by influencing DNAm which in turn can modulate gene expression of target genes that are key to palatogenesis. Using discordant sibling pairs, we were able to refine our results and control for heterogeneity, population stratification and environmental exposures while still being able to detect differences in genetic and epigenetic influences

Our top SNP-CpG association after replication included rs987525 (p=9.63E[-06]), known to be the lead SNP in the OFCs-associated 8q24 region [5, 10, 13, 14, 78, 84-87]. While OFCs-associated variants continue to be identified at the 8q24 locus (one of them, rs72728755, is in high linkage disequilibrium [$r^2$>0.81 in Caucasians] with our meQTL SNP rs987525) the mechanism behind the significant association remains unclear [88-90]. Our analysis shows that rs987525 interacts with a CpG site, cg16561172, located upstream of the *MYC* gene. Thus, our data suggests genetically driven abnormal DNAm in 8q24 as a potential pathogenic mechanism contributing to increased OFCs risk. A second replicated association was rs227731 and cg08592707 (p=2.64E[-02]). While rs227731 was one of the 9 SNPs in our meQTL analysis—owing to its identification as a lead SNP in a previous GWAS study— there is prior evidence that rs227731 likely serves as a sentinel for a cleft-associated allele in complete LD with rs227731, rs227727. Importantly, the cleft-associated allele in rs227727 disrupts an epithelial enhancer located near the *NOG* transcription start site in GMSM-K cells [82]. Intriguingly, *PPM1E,* the gene nearest the associated CpG (cg08592707) has been shown to be epigenetically regulated and controls osteoblast proliferation [91-95], providing rationale for its influence on OFCs. While biologically compelling, functional studies are a prerequisite and a necessary next step for mechanistically link these associations.

We also found significative association between rs7078160 (*VAX1*) and cg09487139 (N/A gene) (p=0.05-blood). A study carried out by Zhang et al. (2018) [96], which included 302 nonsyndromic cleft lip with or without palate (OFCs) in Western Han Chinese population, confirmed that allele A at rs7078160 at *VAX1* gene was a risk factor for OFCS and was over-transmitted among OFCs group.

Research using an animal model with *Vax1* knockout mice demonstrated that this gene is related to palate development, showing its role in the etiology of OFCs[97]. We know that this gene contains a homeobox and has significant expression in brain development.[98] Some independent GWAS studies conducted in different populations have shown that SNPs in VAX1 are associated with OFCs [97 8 99 100 101]. The CpG site cg09487139 maps to a long intergenic non-protein coding RNA 2626 (LINC002626) which has been associated to deficit hyperactivity disorder (ADHD) & bipolar disorder[102]. However, neither a craniofacial candidate gene nor enhancer region proximal to cg09487139 has been characterized, thus precluding any assertions regarding its association with target genes or enhancers implicated in craniofacial development.

Another significant association was found between rs13041247/*MAFB* and cg18347630 (*PLCG1*) (p=0,04-blood). *MAFB* is a basic leucine zipper transcription factor that acts as a transcriptional activator or repressor [103]. It plays a crucial role in regulating lineage-specific hematopoiesis [104] and is important for the differentiation of monocytes, macrophages, osteoclasts, podocytes, and islet beta cells [105-107].

SNPs near *MAFB* were first associated with OFCs in a cleft GWAS using case-parent trios. The authors also demonstrated through an expression study conducted in mouse that the *Mafb* gene plays a role in palatal development. Asian families showed stronger evidence for association with *MAFB* and *ABCA4* [5]. These results were confirmed in different populations, with Asian families showing a strong association of these genes with CL/P, although some results were conflicting [5, 108-111].

Although the relationship between *MAFB* and *PLCG1* is unclear, it is known that PLCG1 is a gene related to cell growth and migration, as well as cancer[112]. In addition to being expressed in immune cells, it is also expressed in various organs and tissues, such as the skin, brain, kidneys, gastrointestinal tract, and muscles [113]. *PLCG1* also plays critical roles in maintaining immune homeostasis[114].

Additionally, our replication study has identified other associations such as rs7590268/*THADA* and cg06873343/*TTYH3*, and rs560426/*ABCA4/ARHGAP29* with cg25196715/*ABCA4/ARHGAP29*. *THADA* and *ABCA4-ARHGAP29*, show a strong link to CLP but not CLO [115]and *ARHGAP29* is recognized as causative for nonsyndromic OFCs[116, 117].

Regarding the *TTYH3* gene found near cg06873342, it is known that this gene encodes a large-conductance chloride channel [118]. A study conducted with *ttyh3 gene in* Xenopus laevis, showed expression during embryonic development in the nervous system and in somites, suggesting important role both in early proliferation and differentiation[119]. However, it is still uncertain whether this gene is similarly expressed in humans [120]. Research related to cancer shows its high expressivity in cervical cancer [121]. At present, its relationship with orofacial clefts is unknown. As for *ARHGAP29,* Liu et al. (2017) [122] applied an experimental pipeline to test the hypothesis that functional SNPs in the 1p22 regions lie within enhancers that drive the expression of *ARHGAP29* in one or more oral tissues, and that the risk alleles alter *ARHGAP29* expression. The study included the application of *in vitro* and *in vivo* enhancer assays, chromatin conformation capture, and genome-editing to provide evidence of functional SNPs. In addition, chromatin immunoprecipitation analysis indicated that the risk associated SNPs affect the activity of *ARHGAP29* enhancers. A recent study found four new pathogenic variants in *ARHGAP29*, which lead to loss of function, such as the loss of the important RhoGap functional domain, showing strong evidence of its relationship with oral clefts [123].

Finally, we also present association between rs3758249 located in *FOXE1* and cg20308679 close to *FRZB* (p=0.04) in the saliva group. Both genes are known to be associated with orofacial clefts. According to a study conducted by Conte et al. (2016), patients with OFCs presented deletions in *FRZB* and *SPRY1* genes[124]. Animal models showed that *FRZB* is expressed in oral tissues, has an important role in the ossification of craniofacial regions by interacting with the WNT pathway. Activation of the

Wnt signaling pathway is necessary for development of many organs of the body [125-129] and also for palatogenesis[130].

We also conducted a differential DNA methylation analysis via paired T tests (Table 3), which shows significant results for three CpG sites, cg006873343 near *TTYH3* gene, cg17103269 near *LPIN3* and cg19191560 *LGR4*. These three genes have unknow function on craniofacial development as of yet. Although the cg25196715 has shown borderline result, it is of particular interest, as it is located close to genes highly associated with clefting in Asian population (*ABCA4/ARHGAP29)* [10]. Mice animal models showed *Arghgap29* gene expression in craniofacial development[117].

Methylome analysis is a valuable post-GWAS resource, giving insights into regulatory genomic potential of GWAS signals, especially the non-coding variants, and helping to prioritize loci to further follow-up. Analyses that combine known genetic risk factors with epigenetic data to explore methylation quantitative loci (meQTLs) function, significantly contribute to understanding such loci by addressing the causal alleles, their pathogenic consequences, and the biological mechanisms through which they influence disease.

Our study indicates the potential for the identification of new cleft related gene networks that can be disrupted via aberrant DNA methylation. A better understanding of genetic-epigenetic interactions underlying OFCs etiology has significant implications in risk prediction and can uncover biological mechanisms that could potentially be targeted for prevention strategies for OFCs and other craniofacial conditions. Given the large prevalence of non-coding SNPs in most OFCs genome wide association studies, our findings can potentially address major knowledge gaps, like missing heritability, reduced penetrance, and variable expressivity associated with OFCs phenotypes.

**Funding and Acknowledgments**

This work was supported by the National Institute of Dental and Craniofacial Research, Eunice Kennedy Shriver National Institute of Child Health and Human Development grants K01DE027995, R37DE08559, U01DE020057, R01DE012472, R21DE016930, R01DE014667, R01DE028300).

**References**

1.      Rahimov F, Jugessur A, Murray JC. Genetics of nonsyndromic orofacial clefts. Cleft Palate Craniofac J. 2012;49(1):73-91. Epub 20110505. doi: 10.1597/10-178. PubMed PMID: 21545302; PMCID: PMC3437188.

2.      Fraser FC. Thoughts on the etiology of clefts of the palate and lip. Acta Genet Stat Med. 1955;5(4):358-69. PubMed PMID: 13339079.

3.      Marazita ML. The evolution of human genetic studies of cleft lip and cleft palate. Annual review of genomics and human genetics. 2012;13:263-83. Epub 2012/06/19. doi: 10.1146/annurev-genom-090711-163729. PubMed PMID: 22703175; PMCID: PMC3760163.

4.      Grosen D, Bille C, Pedersen JK, Skytthe A, Murray JC, Christensen K. Recurrence risk for offspring of twins discordant for oral cleft: a population-based cohort study of the Danish 1936-2004 cleft twin cohort. American journal of medical genetics Part A. 2010;152a(10):2468-74. Epub 2010/08/28. doi: 10.1002/ajmg.a.33608. PubMed PMID: 20799319; PMCID: PMC2946514.

5.      Beaty TH, Murray JC, Marazita ML, Munger RG, Ruczinski I, Hetmanski JB, Liang KY, Wu T, Murray T, Fallin MD, Redett RA, Raymond G, Schwender H, Jin SC, Cooper ME, Dunnwald M, Mansilla MA, Leslie E, Bullard S, Lidral AC, Moreno LM, Menezes R, Vieira AR, Petrin A, Wilcox AJ, Lie RT, Jabs EW, Wu-Chou YH, Chen PK, Wang H, Ye X, Huang S, Yeow V, Chong SS, Jee SH, Shi B, Christensen K, Melbye M, Doheny KF, Pugh EW, Ling H, Castilla EE, Czeizel AE, Ma L, Field LL, Brody L, Pangilinan F, Mills JL, Molloy AM, Kirke PN, Scott JM, Arcos-Burgos M, Scott AF. A genome-wide association study of cleft lip with and without cleft palate identifies risk variants near MAFB and ABCA4. Nat Genet. 2010;42(6):525-9. Epub 2010/05/04. doi: 10.1038/ng.580. PubMed PMID: 20436469; PMCID: PMC2941216.

6.      Birnbaum S, Ludwig KU, Reutter H, Herms S, Steffens M, Rubini M, Baluardo C, Ferrian M, Almeida de Assis N, Alblas MA, Barth S, Freudenberg J, Lauster C, Schmidt G, Scheer M, Braumann B, Berge SJ, Reich RH, Schiefke F, Hemprich A, Potzsch S, Steegers-Theunissen RP, Potzsch B, Moebus S, Horsthemke B, Kramer FJ, Wienker TF, Mossey PA, Propping P, Cichon S, Hoffmann P, Knapp M, Nothen MM, Mangold E. Key susceptibility locus for nonsyndromic cleft lip with or without cleft palate on chromosome 8q24. Nat Genet. 2009;41(4):473-7. Epub 2009/03/10. doi: ng.333 [pii] 10.1038/ng.333. PubMed PMID: 19270707.

7.      Leslie EJ, Liu H, Carlson JC, Shaffer JR, Feingold E, Wehby G, Laurie CA, Jain D, Laurie CC, Doheny KF, McHenry T, Resick J, Sanchez C, Jacobs J, Emanuele B, Vieira AR, Neiswanger K, Standley J, Czeizel AE, Deleyiannis F, Christensen K, Munger RG, Lie RT, Wilcox A, Romitti PA, Field LL, Padilla CD, Cutiongco-de la Paz EM, Lidral AC, Valencia-Ramirez LC, Lopez-Palacio AM, Valencia DR, Arcos-Burgos M, Castilla EE, Mereb JC, Poletta FA, Orioli IM, Carvalho FM, Hecht JT, Blanton SH, Buxo CJ, Butali A, Mossey PA, Adeyemo WL, James O, Braimah RO, Aregbesola BS, Eshete MA, Deribew M, Koruyucu M, Seymen F, Ma L, de Salamanca JE, Weinberg SM, Moreno L, Cornell RA, Murray JC, Marazita ML. A Genome-wide Association Study of Nonsyndromic Cleft Palate Identifies an Etiologic Missense Variant in GRHL3. Am J Hum Genet. 2016;98(4):744-54. Epub 2016/03/29. doi: 10.1016/j.ajhg.2016.02.014. PubMed PMID: 27018472; PMCID: PMC4833215.

8.      Mangold E, Ludwig KU, Birnbaum S, Baluardo C, Ferrian M, Herms S, Reutter H, de Assis NA, Chawa TA, Mattheisen M, Steffens M, Barth S, Kluck N, Paul A, Becker J, Lauster C, Schmidt G, Braumann B, Scheer M, Reich RH, Hemprich A, Potzsch S, Blaumeiser B, Moebus S, Krawczak M, Schreiber S, Meitinger T, Wichmann HE, Steegers-Theunissen RP, Kramer FJ, Cichon S, Propping P, Wienker TF, Knapp M, Rubini M, Mossey PA, Hoffmann P, Nothen MM. Genome-wide association study identifies two susceptibility loci for nonsyndromic cleft lip with or without cleft palate. Nat Genet. 2010;42(1):24-6. Epub 2009/12/22. doi: ng.506 [pii] 10.1038/ng.506. PubMed PMID: 20023658.

9.    Camargo M, Rivera D, Moreno L, Lidral AC, Harper U, Jones M, Solomon BD, Roessler E, Velez JI, Martinez AF, Chandrasekharappa SC, Arcos-Burgos M. GWAS reveals new recessive loci associated with non-syndromic facial clefting. European journal of medical genetics. 2012;55(10):510-4. Epub 2012/07/04. doi: 10.1016/j.ejmg.2012.06.005. PubMed PMID: 22750566; PMCID: PMC3521569.

10.    Grant SF, Wang K, Zhang H, Glaberson W, Annaiah K, Kim CE, Bradfield JP, Glessner JT, Thomas KA, Garris M, Frackelton EC, Otieno FG, Chiavacci RM, Nah HD, Kirschner RE, Hakonarson H. A genome-wide association study identifies a locus for nonsyndromic cleft lip with or without cleft palate on 8q24. J Pediatr. 2009;155(6):909-13. Epub 2009/08/07. doi: 10.1016/j.jpeds.2009.06.020. PubMed PMID: 19656524.

11.    Sun Y, Huang Y, Yin A, Pan Y, Wang Y, Wang C, Du Y, Wang M, Lan F, Hu Z, Wang G, Jiang M, Ma J, Zhang X, Ma H, Ma J, Zhang W, Huang Q, Zhou Z, Ma L, Li Y, Jiang H, Xie L, Jiang Y, Shi B, Cheng J, Shen H, Wang L, Yang Y. Genome-wide association study identifies a new susceptibility locus for cleft lip with or without a cleft palate. Nat Commun. 2015;6:6414. doi: 10.1038/ncomms7414. PubMed PMID: 25775280.

12.    Wolf ZT, Brand HA, Shaffer JR, Leslie EJ, Arzi B, Willet CE, Cox TC, McHenry T, Narayan N, Feingold E, Wang X, Sliskovic S, Karmi N, Safra N, Sanchez C, Deleyiannis FW, Murray JC, Wade CM, Marazita ML, Bannasch DL. Genome-wide association studies in dogs and humans identify ADAMTS20 as a risk variant for cleft lip and palate. PLoS genetics. 2015;11(3):e1005059. Epub 2015/03/24. doi: 10.1371/journal.pgen.1005059. PubMed PMID: 25798845; PMCID: PMC4370697.

13.    Beaty TH, Taub MA, Scott AF, Murray JC, Marazita ML, Schwender H, Parker MM, Hetmanski JB, Balakrishnan P, Mansilla MA, Mangold E, Ludwig KU, Noethen MM, Rubini M, Elcioglu N, Ruczinski I. Confirming genes influencing risk to cleft lip with/without cleft palate in a case-parent trio study. Hum Genet. 2013;132(7):771-81. Epub 20130320. doi: 10.1007/s00439-013-1283-6. PubMed PMID: 23512105; PMCID: PMC3707506.

14.    Ludwig KU, Mangold E, Herms S, Nowak S, Reutter H, Paul A, Becker J, Herberz R, AlChawa T, Nasser E, Bohmer AC, Mattheisen M, Alblas MA, Barth S, Kluck N, Lauster C, Braumann B, Reich RH, Hemprich A, Potzsch S, Blaumeiser B, Daratsianos N, Kreusch T, Murray JC, Marazita ML, Ruczinski I, Scott AF, Beaty TH, Kramer FJ, Wienker TF, Steegers-Theunissen RP, Rubini M, Mossey PA, Hoffmann P, Lange C, Cichon S, Propping P, Knapp M, Nothen MM. Genome-wide meta-analyses of nonsyndromic cleft lip with or without cleft palate identify six new risk loci. Nat Genet. 2012;44(9):968-71. Epub 2012/08/07. doi: 10.1038/ng.2360. PubMed PMID: 22863734; PMCID: PMC3598617.

15.    Moreno LM, Mansilla MA, Bullard SA, Cooper ME, Busch TD, Machida J, Johnson MK, Brauer D, Krahn K, Daack-Hirsch S, L'Heureux J, Valencia-Ramirez C, Rivera D, Lopez AM, Moreno MA, Hing A, Lammer EJ, Jones M, Christensen K, Lie RT, Jugessur A, Wilcox AJ, Chines P, Pugh E, Doheny K, Arcos-Burgos M, Marazita ML, Murray JC, Lidral AC. FOXE1 association with both isolated cleft lip with or without cleft palate, and isolated cleft palate. Hum Mol Genet. 2009;18(24):4879-96. Epub 2009/09/26. doi: 10.1093/hmg/ddp444. PubMed PMID: 19779022; PMCID: PMC2778374.

16.    Song J, Rechkoblit O, Bestor TH, Patel DJ. Structure of DNMT1-DNA complex reveals a role for autoinhibition in maintenance DNA methylation. Science. 2011;331(6020):1036-40. Epub 20101216. doi: 10.1126/science.1195380. PubMed PMID: 21163962; PMCID: PMC4689315.

17.    Bourc'his D, Xu GL, Lin CS, Bollman B, Bestor TH. Dnmt3L and the establishment of maternal genomic imprints. Science. 2001;294(5551):2536-9. Epub 20011122. doi: 10.1126/science.1065848. PubMed PMID: 11719692.

18.    van Eijk KR, de Jong S, Boks MPM, Langeveld T, Colas F, Veldink JH, de Kovel CGF, Janson E, Strengman E, Langfelder P, Kahn RS, van den Berg LH, Horvath S, Ophoff RA. Genetic analysis of

DNA methylation and gene expression levels in whole blood of healthy human subjects. BMC Genomics. 2012;13(1):636. doi: 10.1186/1471-2164-13-636.

19.     Vilain A, Bernardino J, Gerbault-Seureau M, Vogt N, Niveleau A, Lefrançois D, Malfoy B, Dutrillaux B. DNA methylation and chromosome instability in lymphoblastoid cell lines. Cytogenet Cell Genet. 2000;90(1-2):93-101. doi: 10.1159/000015641. PubMed PMID: 11060456.

20.     Maunakea AK, Nagarajan RP, Bilenky M, Ballinger TJ, D'Souza C, Fouse SD, Johnson BE, Hong C, Nielsen C, Zhao Y, Turecki G, Delaney A, Varhol R, Thiessen N, Shchors K, Heine VM, Rowitch DH, Xing X, Fiore C, Schillebeeckx M, Jones SJM, Haussler D, Marra MA, Hirst M, Wang T, Costello JF. Conserved role of intragenic DNA methylation in regulating alternative promoters. Nature. 2010;466(7303):253-7. doi: 10.1038/nature09165.

21.     Sharp AJ, Stathaki E, Migliavacca E, Brahmachary M, Montgomery SB, Dupre Y, Antonarakis SE. DNA methylation profiles of human active and inactive X chromosomes. Genome Res. 2011;21(10):1592-600. Epub 20110823. doi: 10.1101/gr.112680.110. PubMed PMID: 21862626; PMCID: PMC3202277.

22.     Pervjakova N, Kasela S, Morris AP, Kals M, Metspalu A, Lindgren CM, Salumets A, Mägi R. Imprinted genes and imprinting control regions show predominant intermediate methylation in adult somatic tissues. Epigenomics. 2016;8(6):789-99. Epub 20160323. doi: 10.2217/epi.16.8. PubMed PMID: 27004446; PMCID: PMC5066126.

23.     Alvizi L, Ke X, Brito LA, Seselgyte R, Moore GE, Stanier P, Passos-Bueno MR. Differential methylation is associated with non-syndromic cleft lip and palate and contributes to penetrance effects. Sci Rep. 2017;7(1):2441. Epub 20170526. doi: 10.1038/s41598-017-02721-0. PubMed PMID: 28550290; PMCID: PMC5446392.

24.     Banovich NE, Lan X, McVicker G, van de Geijn B, Degner JF, Blischak JD, Roux J, Pritchard JK, Gilad Y. Methylation QTLs are associated with coordinated changes in transcription factor binding, histone modifications, and gene expression levels. PLoS Genet. 2014;10(9):e1004663. Epub 20140918. doi: 10.1371/journal.pgen.1004663. PubMed PMID: 25233095; PMCID: PMC4169251.

25.     Chundru VK, Marioni RE, Prendergast JGD, Lin T, Beveridge AJ, Martin NG, Montgomery GW, Hume DA, Deary IJ, Visscher PM, Wray NR, McRae AF. Rare genetic variants underlie outlying levels of DNA methylation and gene-expression. Hum Mol Genet. 2023;32(11):1912-21. doi: 10.1093/hmg/ddad028. PubMed PMID: 36790133; PMCID: PMC10196672.

26.     Juriloff DM, Harris MJ, Mager DL, Gagnier L. Epigenetic mechanism causes Wnt9b deficiency and nonsyndromic cleft lip and palate in the A/WySn mouse strain. Birth Defects Res A Clin Mol Teratol. 2014;100(10):772-88. Epub 2014/09/27. doi: 10.1002/bdra.23320. PubMed PMID: 25257647.

27.     Petrin AL, Zeng E, Thomas MA, Moretti-Ferreira D, Marazita ML, Xie XJ, Murray JC, Moreno-Uribe LM. DNA methylation differences in monozygotic twins with Van der Woude syndrome. Front Dent Med. 2023;4. Epub 20230217. doi: 10.3389/fdmed.2023.1120948. PubMed PMID: 36936396; PMCID: PMC10019782.

28.     Plamondon JA, Harris MJ, Mager DL, Gagnier L, Juriloff DM. The clf2 gene has an epigenetic role in the multifactorial etiology of cleft lip and palate in the A/WySn mouse strain. Birth Defects Res A Clin Mol Teratol. 2011;91(8):716-27. Epub 2011/03/09. doi: 10.1002/bdra.20788. PubMed PMID: 21384535.

29.     Seaberg A, Awotoye W, Qian F, Machado-Paula LA, Dunlay L, Butali A, Murray J, Moreno-Uribe L, Petrin AL. DNA Methylation Effects on Van der Woude Syndrome Phenotypic Variability. Cleft Palate Craniofac J. 2024:10556656241269495. Epub 20240807. doi: 10.1177/10556656241269495. PubMed PMID: 39109995.

30.     Sharp GC, Stergiakouli E, Sandy J, Relton C. Epigenetics and Orofacial Clefts: A Brief Introduction. Cleft Palate Craniofac J. 2018;55(6):795-7. Epub 2017/01/14. doi: 10.1597/16-124. PubMed PMID: 28085511.

31.     Stefansson OA, Sigurpalsdottir BD, Rognvaldsson S, Halldorsson GH, Juliusson K, Sveinbjornsson G, Gunnarsson B, Beyter D, Jonsson H, Gudjonsson SA, Olafsdottir TA, Saevarsdottir S, Magnusson MK, Lund SH, Tragante V, Oddsson A, Hardarson MT, Eggertsson HP, Gudmundsson RL, Sverrisson S, Frigge ML, Zink F, Holm H, Stefansson H, Rafnar T, Jonsdottir I, Sulem P, Helgason A, Gudbjartsson DF, Halldorsson BV, Thorsteinsdottir U, Stefansson K. The correlation between CpG methylation and gene expression is driven by sequence variants. Nat Genet. 2024. Epub 20240724. doi: 10.1038/s41588-024-01851-2. PubMed PMID: 39048797.

32.     Villicaña S, Bell JT. Genetic impacts on DNA methylation: research findings and future perspectives. Genome biology. 2021;22(1):127. Epub 2021/05/02. doi: 10.1186/s13059-021-02347-6. PubMed PMID: 33931130; PMCID: PMC8086086.

33.     Villicaña S, Castillo-Fernandez J, Hannon E, Christiansen C, Tsai PC, Maddock J, Kuh D, Suderman M, Power C, Relton C, Ploubidis G, Wong A, Hardy R, Goodman A, Ong KK, Bell JT. Genetic impacts on DNA methylation help elucidate regulatory genomic processes. Genome biology. 2023;24(1):176. Epub 20230731. doi: 10.1186/s13059-023-03011-x. PubMed PMID: 37525248; PMCID: PMC10391992.

34.     Alvizi L, Ke X, Brito LA, Seselgyte R, Moore GE, Stanier P, Passos-Bueno MR. Differential methylation is associated with non-syndromic cleft lip and palate and contributes to penetrance effects. Scientific reports. 2017;7(1):2441. doi: 10.1038/s41598-017-02721-0.

35.     Gonseth S, Shaw GM, Roy R, Segal MR, Asrani K, Rine J, Wiemels J, Marini NJ. Epigenomic profiling of newborns with isolated orofacial clefts reveals widespread DNA methylation changes and implicates metastable epiallele regions in disease risk. Epigenetics. 2019;14(2):198-213. Epub 2019/03/15. doi: 10.1080/15592294.2019.1581591. PubMed PMID: 30870065.

36.     Joubert BR, Felix JF, Yousefi P, Bakulski KM, Just AC, Breton C, Reese SE, Markunas CA, Richmond RC, Xu CJ, Kupers LK, Oh SS, Hoyo C, Gruzieva O, Soderhall C, Salas LA, Baiz N, Zhang H, Lepeule J, Ruiz C, Ligthart S, Wang T, Taylor JA, Duijts L, Sharp GC, Jankipersadsing SA, Nilsen RM, Vaez A, Fallin MD, Hu D, Litonjua AA, Fuemmeler BF, Huen K, Kere J, Kull I, Munthe-Kaas MC, Gehring U, Bustamante M, Saurel-Coubizolles MJ, Quraishi BM, Ren J, Tost J, Gonzalez JR, Peters MJ, Haberg SE, Xu Z, van Meurs JB, Gaunt TR, Kerkhof M, Corpeleijn E, Feinberg AP, Eng C, Baccarelli AA, Benjamin Neelon SE, Bradman A, Merid SK, Bergstrom A, Herceg Z, Hernandez-Vargas H, Brunekreef B, Pinart M, Heude B, Ewart S, Yao J, Lemonnier N, Franco OH, Wu MC, Hofman A, McArdle W, Van der Vlies P, Falahi F, Gillman MW, Barcellos LF, Kumar A, Wickman M, Guerra S, Charles MA, Holloway J, Auffray C, Tiemeier HW, Smith GD, Postma D, Hivert MF, Eskenazi B, Vrijheid M, Arshad H, Anto JM, Dehghan A, Karmaus W, Annesi-Maesano I, Sunyer J, Ghantous A, Pershagen G, Holland N, Murphy SK, DeMeo DL, Burchard EG, Ladd-Acosta C, Snieder H, Nystad W, Koppelman GH, Relton CL, Jaddoe VW, Wilcox A, Melen E, London SJ. DNA Methylation in Newborns and Maternal Smoking in Pregnancy: Genome-wide Consortium Meta-analysis. Am J Hum Genet. 2016;98(4):680-96. Epub 2016/04/05. doi: 10.1016/j.ajhg.2016.02.019. PubMed PMID: 27040690; PMCID: PMC4833289.

37.     Sharp GC, Ho K, Davies A, Stergiakouli E, Humphries K, McArdle W, Sandy J, Davey Smith G, Lewis SJ, Relton CL. Distinct DNA methylation profiles in subtypes of orofacial cleft. Clin Epigenetics. 2017;9:63. Epub 2017/06/13. doi: 10.1186/s13148-017-0362-2. PubMed PMID: 28603561; PMCID: PMC5465456.

38.     Shu X, Shu S, Zhai Y, Zhu L, Ouyang Z. Genome-Wide DNA Methylation Profile of Gene cis-Acting Element Methylations in All-trans Retinoic Acid-Induced Mouse Cleft Palate. DNA and cell biology. 2018. Epub 2018/10/03. doi: 10.1089/dna.2018.4369. PubMed PMID: 30277813.

39.     Xu Z, Lie RT, Wilcox AJ, Saugstad OD, Taylor JA. A comparison of DNA methylation in newborn blood samples from infants with and without orofacial clefts. Clinical epigenetics. 2019;11(1):40. doi: 10.1186/s13148-019-0638-9.

40.     Young JI, Slifer S, Hecht JT, Blanton SH. DNA Methylation Variation Is Identified in Monozygotic Twins Discordant for Non-syndromic Cleft Lip and Palate. Frontiers in Cell and Developmental Biology. 2021;9(860). doi: 10.3389/fcell.2021.656865.

41.     Zhang B, Zhang Y, Wu S, Ma D, Ma J. DNA methylation profile of lip tissue from congenital nonsyndromic cleft lip and palate patients by whole-genome bisulfite sequencing. Birth defects research. 2023;115(2):205-17. Epub 20221009. doi: 10.1002/bdr2.2102. PubMed PMID: 36210532; PMCID: PMC10092010.

42.     Alvizi L, Brito LA, Kobayashi GS, Bischain B, da Silva CBF, Ramos SLG, Wang J, Passos-Bueno MR. mir152 hypomethylation as a mechanism for non-syndromic cleft lip and palate. Epigenetics. 2022;17(13):2278-95. Epub 20220901. doi: 10.1080/15592294.2022.2115606. PubMed PMID: 36047706; PMCID: PMC9665146.

43.     Charoenvicha C, Sirimaharaj W, Khwanngern K, Chattipakorn N, Chattipakorn SC. Alterations in DNA Methylation in Orofacial Clefts. Int J Mol Sci. 2022;23(21). Epub 20221022. doi: 10.3390/ijms232112727. PubMed PMID: 36361518; PMCID: PMC9654384.

44.     Howe LJ, Richardson TG, Arathimos R, Alvizi L, Passos-Bueno M-R, Stanier P, Nohr E, Ludwig KU, Mangold E, Knapp M, Stergiakouli E, St Pourcain B, Davey Smith G, Sandy J, Relton CL, Lewis SJ, Hemani G, Sharp GC. DNA methylation mediates genetic liability to non-syndromic cleft lip/palate. bioRxiv. 2018.

45.     Howe LJ, Richardson TG, Arathimos R, Alvizi L, Passos-Bueno MR, Stanier P, Nohr E, Ludwig KU, Mangold E, Knapp M, Stergiakouli E, Pourcain BS, Smith GD, Sandy J, Relton CL, Lewis SJ, Hemani G, Sharp GC. Evidence for DNA methylation mediating genetic liability to non-syndromic cleft lip/palate. Epigenomics. 2019. Epub 2019/01/15. doi: 10.2217/epi-2018-0091. PubMed PMID: 30638414.

46.     Garland MA, Sun B, Zhang S, Reynolds K, Ji Y, Zhou CJ. Role of epigenetics and miRNAs in orofacial clefts. Birth defects research. 2020;112(19):1635-59. Epub 2020/09/15. doi: 10.1002/bdr2.1802. PubMed PMID: 32926553.

47.     Gibbs JR, van der Brug MP, Hernandez DG, Traynor BJ, Nalls MA, Lai SL, Arepalli S, Dillman A, Rafferty IP, Troncoso J, Johnson R, Zielke HR, Ferrucci L, Longo DL, Cookson MR, Singleton AB. Abundant quantitative trait loci exist for DNA methylation and gene expression in human brain. PLoS genetics. 2010;6(5):e1000952. Epub 2010/05/21. doi: 10.1371/journal.pgen.1000952. PubMed PMID: 20485568; PMCID: PMC2869317.

48.     Hannon E, Gorrie-Stone TJ, Smart MC, Burrage J, Hughes A, Bao Y, Kumari M, Schalkwyk LC, Mill J. Leveraging DNA-Methylation Quantitative-Trait Loci to Characterize the Relationship between Methylomic Variation, Gene Expression, and Complex Traits. Am J Hum Genet. 2018;103(5):654-65. Epub 20181025. doi: 10.1016/j.ajhg.2018.09.007. PubMed PMID: 30401456; PMCID: PMC6217758.

49.     Hannon E, Spiers H, Viana J, Pidsley R, Burrage J, Murphy TM, Troakes C, Turecki G, O'Donovan MC, Schalkwyk LC, Bray NJ, Mill J. Methylation QTLs in the developing brain and their enrichment in schizophrenia risk loci. Nature neuroscience. 2016;19(1):48-54. Epub 2015/12/01. doi: 10.1038/nn.4182. PubMed PMID: 26619357; PMCID: PMC4714325.

50.     Do C, Lang CF, Lin J, Darbary H, Krupska I, Gaba A, Petukhova L, Vonsattel JP, Gallagher MP, Goland RS, Clynes RA, Dwork A, Kral JG, Monk C, Christiano AM, Tycko B. Mechanisms and Disease Associations of Haplotype-Dependent Allele-Specific DNA Methylation. Am J Hum Genet. 2016;98(5):934-55. doi: 10.1016/j.ajhg.2016.03.027. PubMed PMID: 27153397; PMCID: PMC4863666.

51.     Gutierrez-Arcelus M, Lappalainen T, Montgomery SB, Buil A, Ongen H, Yurovsky A, Bryois J, Giger T, Romano L, Planchon A, Falconnet E, Bielser D, Gagnebin M, Padioleau I, Borel C, Letourneau A, Makrythanasis P, Guipponi M, Gehrig C, Antonarakis SE, Dermitzakis ET. Passive and active DNA methylation and the interplay with genetic variation in gene regulation. eLife. 2013;2:e00523. Epub 20130604. doi: 10.7554/eLife.00523. PubMed PMID: 23755361; PMCID: PMC3673336.

52.     Pierce BL, Tong L, Argos M, Demanelis K, Jasmine F, Rakibuz-Zaman M, Sarwar G, Islam MT, Shahriar H, Islam T, Rahman M, Yunus M, Kibriya MG, Chen LS, Ahsan H. Co-occurring expression and methylation QTLs allow detection of common causal variants and shared biological mechanisms. Nat Commun. 2018;9(1):804. Epub 20180223. doi: 10.1038/s41467-018-03209-9. PubMed PMID: 29476079; PMCID: PMC5824840.

53.     Wu Y, Zeng J, Zhang F, Zhu Z, Qi T, Zheng Z, Lloyd-Jones LR, Marioni RE, Martin NG, Montgomery GW, Deary IJ, Wray NR, Visscher PM, McRae AF, Yang J. Integrative analysis of omics summary data reveals putative mechanisms underlying complex traits. Nat Commun. 2018;9(1):918. Epub 20180302. doi: 10.1038/s41467-018-03371-0. PubMed PMID: 29500431; PMCID: PMC5834629.

54.     McRae AF, Powell JE, Henders AK, Bowdler L, Hemani G, Shah S, Painter JN, Martin NG, Visscher PM, Montgomery GW. Contribution of genetic variation to transgenerational inheritance of DNA methylation. Genome biology. 2014;15(5):R73. doi: 10.1186/gb-2014-15-5-r73.

55.     Min JL, Hemani G, Hannon E, Dekkers KF, Castillo-Fernandez J, Luijk R, Carnero-Montoro E, Lawson DJ, Burrows K, Suderman M, Bretherick AD, Richardson TG, Klughammer J, Iotchkova V, Sharp G, Al Khleifat A, Shatunov A, Iacoangeli A, McArdle WL, Ho KM, Kumar A, Söderhäll C, Soriano-Tárraga C, Giralt-Steinhauer E, Kazmi N, Mason D, McRae AF, Corcoran DL, Sugden K, Kasela S, Cardona A, Day FR, Cugliari G, Viberti C, Guarrera S, Lerro M, Gupta R, Bollepalli S, Mandaviya P, Zeng Y, Clarke T-K, Walker RM, Schmoll V, Czamara D, Ruiz-Arenas C, Rezwan FI, Marioni RE, Lin T, Awaloff Y, Germain M, Aïssi D, Zwamborn R, van Eijk K, Dekker A, van Dongen J, Hottenga J-J, Willemsen G, Xu C-J, Barturen G, Català-Moll F, Kerick M, Wang C, Melton P, Elliott HR, Shin J, Bernard M, Yet I, Smart M, Gorrie-Stone T, Shaw C, Al Chalabi A, Ring SM, Pershagen G, Melén E, Jiménez-Conde J, Roquer J, Lawlor DA, Wright J, Martin NG, Montgomery GW, Moffitt TE, Poulton R, Esko T, Milani L, Metspalu A, Perry JRB, Ong KK, Wareham NJ, Matullo G, Sacerdote C, Panico S, Caspi A, Arseneault L, Gagnon F, Ollikainen M, Kaprio J, Felix JF, Rivadeneira F, Tiemeier H, van Ijzendoorn MH, Uitterlinden AG, Jaddoe VWV, Haley C, McIntosh AM, Evans KL, Murray A, Räikkönen K, Lahti J, Nohr EA, Sørensen TIA, Hansen T, Morgen CS, Binder EB, Lucae S, Gonzalez JR, Bustamante M, Sunyer J, Holloway JW, Karmaus W, Zhang H, Deary IJ, Wray NR, Starr JM, Beekman M, van Heemst D, Slagboom PE, Morange P-E, Trégouët D-A, Veldink JH, Davies GE, de Geus EJC, Boomsma DI, Vonk JM, Brunekreef B, Koppelman GH, Alarcón-Riquelme ME, Huang R-C, Pennell CE, van Meurs J, Ikram MA, Hughes AD, Tillin T, Chaturvedi N, Pausova Z, Paus T, Spector TD, Kumari M, Schalkwyk LC, Visscher PM, Davey Smith G, Bock C, Gaunt TR, Bell JT, Heijmans BT, Mill J, Relton CL, Consortium B. Genomic and phenotypic insights from an atlas of genetic effects on DNA methylation. Nat Genet. 2021;53(9):1311-21. doi: 10.1038/s41588-021-00923-x.

56.     van Dongen J, Nivard MG, Willemsen G, Hottenga JJ, Helmer Q, Dolan CV, Ehli EA, Davies GE, van Iterson M, Breeze CE, Beck S, Suchiman HE, Jansen R, van Meurs JB, Heijmans BT, Slagboom PE, Boomsma DI. Genetic and environmental influences interact with age and sex in shaping the human methylome. Nat Commun. 2016;7:11115. Epub 2016/04/08. doi: 10.1038/ncomms11115. PubMed PMID: 27051996; PMCID: PMC4820961.

57.     Ng B, White CC, Klein H-U, Sieberts SK, McCabe C, Patrick E, Xu J, Yu L, Gaiteri C, Bennett DA, Mostafavi S, De Jager PL. An xQTL map integrates the genetic architecture of the human brain's transcriptome and epigenome. Nat Neurosci. 2017;20(10):1418-26. doi: 10.1038/nn.4632.

58.     Schulz H, Ruppert A-K, Herms S, Wolf C, Mirza-Schreiber N, Stegle O, Czamara D, Forstner AJ, Sivalingam S, Schoch S, Moebus S, Pütz B, Hillmer A, Fricker N, Vatter H, Müller-Myhsok B, Nöthen MM, Becker AJ, Hoffmann P, Sander T, Cichon S. Genome-wide mapping of genetic determinants influencing DNA methylation and gene expression in human hippocampus. Nature communications. 2017;8(1):1511-. doi: 10.1038/s41467-017-01818-4. PubMed PMID: 29142228.

59.     Grundberg E, Meduri E, Sandling JK, Hedman AK, Keildson S, Buil A, Busche S, Yuan W, Nisbet J, Sekowska M, Wilk A, Barrett A, Small KS, Ge B, Caron M, Shin SY, Lathrop M, Dermitzakis ET, McCarthy MI, Spector TD, Bell JT, Deloukas P. Global analysis of DNA methylation variation in adipose tissue from twins reveals links to disease-associated variants in distal regulatory elements. Am J Hum Genet. 2013;93(5):876-90. Epub 2013/11/05. doi: 10.1016/j.ajhg.2013.10.004. PubMed PMID: 24183450; PMCID: PMC3824131.

60.     van Dongen J, Ehli EA, Jansen R, van Beijsterveldt CEM, Willemsen G, Hottenga JJ, Kallsen NA, Peyton SA, Breeze CE, Kluft C, Heijmans BT, Bartels M, Davies GE, Boomsma DI. Genome-wide analysis of DNA methylation in buccal cells: a study of monozygotic twins and mQTLs. Epigenetics & chromatin. 2018;11(1):54-. doi: 10.1186/s13072-018-0225-x. PubMed PMID: 30253792.

61.     Smith AK, Kilaru V, Kocak M, Almli LM, Mercer KB, Ressler KJ, Tylavsky FA, Conneely KN. Methylation quantitative trait loci (meQTLs) are consistently detected across ancestry, developmental stage, and tissue type. BMC genomics. 2014;15:145. Epub 20140221. doi: 10.1186/1471-2164-15-145. PubMed PMID: 24555763; PMCID: PMC4028873.

62.     Kim E, Kwak SH, Chung HR, Ohn JH, Bae JH, Choi SH, Park KS, Hong JS, Sung J, Jang HC. DNA methylation profiles in sibling pairs discordant for intrauterine exposure to maternal gestational diabetes. Epigenetics. 2017;12(10):825-32. Epub 20171127. doi: 10.1080/15592294.2017.1370172. PubMed PMID: 29099273; PMCID: PMC5788413.

63.     Romanowska J, Haaland ØA, Jugessur A, Gjerdevik M, Xu Z, Taylor J, Wilcox AJ, Jonassen I, Lie RT, Gjessing HK. Gene–methylation interactions: discovering region-wise DNA methylation levels that modify SNP-associated disease risk. Clinical epigenetics. 2020;12(1):109. doi: 10.1186/s13148-020-00881-x.

64.     Moreno Uribe LM, Fomina T, Munger RG, Romitti PA, Jenkins MM, Gjessing HK, Gjerdevik M, Christensen K, Wilcox AJ, Murray JC, Lie RT, Wehby GL. A Population-Based Study of Effects of Genetic Loci on Orofacial Clefts. Journal of dental research. 2017;96(11):1322-9. doi: 10.1177/0022034517716914. PubMed PMID: PMC5607104.

65.     Eads CA, Danenberg KD, Kawakami K, Saltz LB, Blake C, Shibata D, Danenberg PV, Laird PW. MethyLight: a high-throughput assay to measure DNA methylation. Nucleic acids research. 2000;28(8):e32-00. doi: 10.1093/nar/28.8.e32.

66.     Cottrell S, Jung K, Kristiansen G, Eltze E, Semjonow A, Ittmann M, Hartmann A, Stamey T, Haefliger C, Weiss G. Discovery and Validation of 3 Novel DNA Methylation Markers of Prostate Cancer Prognosis. J Urol. 2007;177(5):1753-8. doi: 10.1016/j.juro.2007.01.010.

67.     Shabalin AA. Matrix eQTL: ultra fast eQTL analysis via large matrix operations. Bioinformatics. 2012;28(10):1353-8. Epub 2012/04/12. doi: 10.1093/bioinformatics/bts163. PubMed PMID: 22492648; PMCID: PMC3348564.

68.     Huan T, Joehanes R, Song C, Peng F, Guo Y, Mendelson M, Yao C, Liu C, Ma J, Richard M, Agha G, Guan W, Almli LM, Conneely KN, Keefe J, Hwang S-J, Johnson AD, Fornage M, Liang L, Levy

D. Genome-wide identification of DNA methylation QTLs in whole blood highlights pathways for cardiovascular disease. Nature Communications. 2019;10(1):4267. doi: 10.1038/s41467-019-12228-z.

69.     Pan H, Holbrook JD, Karnani N, Kwoh CK. Gene, Environment and Methylation (GEM): a tool suite to efficiently navigate large scale epigenome wide association studies and integrate genotype and interaction between genotype and environment. BMC Bioinformatics. 2016;17:299. Epub 2016/08/03. doi: 10.1186/s12859-016-1161-z. PubMed PMID: 27480116; PMCID: PMC4970299.

70.     Ruiz-Arenas C, Hernandez-Ferrer C, Vives-Usano M, Marí S, Quintela I, Mason D, Cadiou S, Casas M, Andrusaityte S, Gutzkow KB, Vafeiadi M, Wright J, Lepeule J, Grazuleviciene R, Chatzi L, Carracedo Á, Estivill X, Marti E, Escaramís G, Vrijheid M, González JR, Bustamante M. Identification of autosomal cis expression quantitative trait methylation (cis eQTMs) in children's blood. eLife. 2022;11:e65310. doi: 10.7554/eLife.65310.

71.     Clark AD, Nair N, Anderson AE, Thalayasingam N, Naamane N, Skelton AJ, Diboll J, Barton A, Eyre S, Isaacs JD, Pratt AG, Reynard LN. Lymphocyte DNA methylation mediates genetic risk at shared immune-mediated disease loci. Journal of Allergy and Clinical Immunology. 2020;145(5):1438-51. doi: https://doi.org/10.1016/j.jaci.2019.12.910.

72.     Cornelis JV, Cheng-Jian X, Judith MV, Nick HTtH, Wim T, Irene HH, Martijn CN, Jeunard B, Antoon JvO, Karen A, Markus W, Gerard HK, Maarten van den B. Differential DNA methylation in bronchial biopsies between persistent asthma and asthma in remission. Eur Respir J. 2020;55(2):1901280. doi: 10.1183/13993003.01280-2019.

73.     Kehayova YS, Wilkinson JM, Rice SJ, Loughlin J. Mediation of the Same Epigenetic and Transcriptional Effect by Independent Osteoarthritis Risk–Conferring Alleles on a Shared Target Gene, COLGALT2. Arthritis & Rheumatology. 2023;75(6):910-22. doi: https://doi.org/10.1002/art.42427.

74.     Qin T, Li S, Henry LE, Chou E, Cavalcante RG, Garb BF, D'Silva NJ, Rozek LS, Sartor MA. Whole-genome CpG-resolution DNA Methylation Profiling of HNSCC Reveals Distinct Mechanisms of Carcinogenesis for Fine-scale HPV+ Cancer Subtypes. Cancer Research Communications. 2023;3(8):1701-15. doi: 10.1158/2767-9764.CRC-23-0009.

75.     Fishilevich S, Nudel R, Rappaport N, Hadar R, Plaschkes I, Iny Stein T, Rosen N, Kohn A, Twik M, Safran M, Lancet D, Cohen D. GeneHancer: genome-wide integration of enhancers and target genes in GeneCards. Database : the journal of biological databases and curation. 2017;2017:bax028. doi: 10.1093/database/bax028. PubMed PMID: 28605766.

76.     Wilderman A, VanOudenhove J, Kron J, Noonan JP, Cotney J. High-Resolution Epigenomic Atlas of Human Embryonic Craniofacial Development. Cell reports. 2018;23(5):1581-97. Epub 2018/05/03. doi: 10.1016/j.celrep.2018.03.129. PubMed PMID: 29719267; PMCID: PMC5965702.

77.     Yankee TN, Oh S, Winchester EW, Wilderman A, Robinson K, Gordon T, Rosenfeld JA, VanOudenhove J, Scott DA, Leslie EJ, Cotney J. Integrative analysis of transcriptome dynamics during human craniofacial development identifies candidate disease genes. Nature Communications. 2023;14(1):4623. doi: 10.1038/s41467-023-40363-1.

78.     Salagovic J, Klimcakova L, Zabavnikova M, Behunova J, Hudakova T, Fedeles J, Molnarova A, Podracka L. Polymorphisms at 1q32, 8q24, and 17q22 loci are associated with nonsyndromic cleft lip with or without cleft palate risk in the Slovak population. Biomed Pap Med Fac Univ Palacky Olomouc Czech Repub. 2017;161(2):152-7. Epub 2017/04/20. doi: 10.5507/bp.2017.009. PubMed PMID: 28422192.

79.     Wang F, Jiang Y, Yang S, Liu Q, Lin J, Zhang H. Associations between the NOGGIN rs227731 polymorphism and NSCL/P risk may be associated with ethnicities: A meta-analysis. Birth defects research. 2017;109(6):445-51. Epub 2017/04/12. doi: 10.1002/bdra.23612. PubMed PMID: 28398705.

80.    de Aquino SN, Messetti AC, Hoshi R, Borges A, Viena CS, Reis SR, Oliveira Swerts MS, Graner E, Martelli-Junior H, Coletta RD. Analysis of susceptibility polymorphisms for nonsyndromic cleft lip with or without cleft palate in the Brazilian population. Birth Defects Res A Clin Mol Teratol. 2014;100(1):36-42. Epub 2014/01/22. doi: 10.1002/bdra.23204. PubMed PMID: 24446087.

81.    Mostowska A, Hozyasz KK, Wojcicka K, Biedziak B, Jagodzinski PP. Polymorphic variants at 10q25.3 and 17q22 loci and the risk of non-syndromic cleft lip and palate in the Polish population. Birth Defects Res A Clin Mol Teratol. 2012;94(1):42-6. Epub 2011/10/25. doi: 10.1002/bdra.22862. PubMed PMID: 22021054.

82.    Leslie EJ, Taub MA, Liu H, Steinberg KM, Koboldt DC, Zhang Q, Carlson JC, Hetmanski JB, Wang H, Larson DE, Fulton RS, Kousa YA, Fakhouri WD, Naji A, Ruczinski I, Begum F, Parker MM, Busch T, Standley J, Rigdon J, Hecht JT, Scott AF, Wehby GL, Christensen K, Czeizel AE, Deleyiannis FW, Schutte BC, Wilson RK, Cornell RA, Lidral AC, Weinstock GM, Beaty TH, Marazita ML, Murray JC. Identification of functional variants for cleft lip with or without cleft palate in or near PAX7, FGFR2, and NOG by targeted sequencing of GWAS loci. Am J Hum Genet. 2015;96(3):397-411. Epub 2015/02/24. doi: 10.1016/j.ajhg.2015.01.004. PubMed PMID: 25704602; PMCID: PMC4375420.

83.    Gaunt TR, Shihab HA, Hemani G, Min JL, Woodward G, Lyttleton O, Zheng J, Duggirala A, McArdle WL, Ho K, Ring SM, Evans DM, Davey Smith G, Relton CL. Systematic identification of genetic influences on methylation across the human life course. Genome biology. 2016;17(1):61. doi: 10.1186/s13059-016-0926-z.

84.    Butali A, Mossey PA, Adeyemo WL, Eshete MA, Gowans LJJ, Busch TD, Jain D, Yu W, Huan L, Laurie CA, Laurie CC, Nelson S, Li M, Sanchez-Lara PA, Magee WP, Magee KS, Auslander A, Brindopke F, Kay DM, Caggana M, Romitti PA, Mills JL, Audu R, Onwuamah C, Oseni GO, Owais A, James O, Olaitan PB, Aregbesola BS, Braimah RO, Oginni FO, Oladele AO, Bello SA, Rhodes J, Shiang R, Donkor P, Obiri-Yeboah S, Arthur FKN, Twumasi P, Agbenorku P, Plange-Rhule G, Oti AA, Ogunlewe OM, Oladega AA, Adekunle AA, Erinoso AO, Adamson OO, Elufowoju AA, Ayelomi OI, Hailu T, Hailu A, Demissie Y, Derebew M, Eliason S, Romero-Bustillous M, Lo C, Park J, Desai S, Mohammed M, Abate F, Abdur-Rahman LO, Anand D, Saadi I, Oladugba AV, Lachke SA, Amendt BA, Rotimi CN, Marazita ML, Cornell RA, Murray JC, Adeyemo AA. Genomic analyses in African populations identify novel risk loci for cleft palate. Hum Mol Genet. 2019;28(6):1038-51. Epub 2018/11/20. doi: 10.1093/hmg/ddy402. PubMed PMID: 30452639; PMCID: PMC6400042.

85.    Carlson JC, Taub MA, Feingold E, Beaty TH, Murray JC, Marazita ML, Leslie EJ. Identifying Genetic Sources of Phenotypic Heterogeneity in Orofacial Clefts by Targeted Sequencing. Birth defects research. 2017. Epub 2017/04/23. doi: 10.1002/bdra.23605. PubMed PMID: 28432837.

86.    Gowans LJ, Adeyemo WL, Eshete M, Mossey PA, Busch T, Aregbesola B, Donkor P, Arthur FK, Bello SA, Martinez A, Li M, Augustine-Akpan EA, Deressa W, Twumasi P, Olutayo J, Deribew M, Agbenorku P, Oti AA, Braimah R, Plange-Rhule G, Gesses M, Obiri-Yeboah S, Oseni GO, Olaitan PB, Abdur-Rahman L, Abate F, Hailu T, Gravem P, Ogunlewe MO, Buxó CJ, Marazita ML, Adeyemo AA, Murray JC, Butali A. Association Studies and Direct DNA Sequencing Implicate Genetic Susceptibility Loci in the Etiology of Nonsyndromic Orofacial Clefts in Sub-Saharan African Populations. Journal of dental research. 2016;95(11):1245-56. Epub 20160701. doi: 10.1177/0022034516657003. PubMed PMID: 27369588; PMCID: PMC5076758.

87.    Murray T, Taub MA, Ruczinski I, Scott AF, Hetmanski JB, Schwender H, Patel P, Zhang TX, Munger RG, Wilcox AJ, Ye X, Wang H, Wu T, Wu-Chou YH, Shi B, Jee SH, Chong S, Yeow V, Murray JC, Marazita ML, Beaty TH. Examining markers in 8q24 to explain differences in evidence for association with cleft lip with/without cleft palate between Asians and Europeans. Genetic epidemiology. 2012;36(4):392-9. Epub 2012/04/18. doi: 10.1002/gepi.21633. PubMed PMID: 22508319; PMCID: PMC3615645.

88.     Mukhopadhyay N, Feingold E, Moreno-Uribe L, Wehby G, Valencia-Ramirez LC, Muñeton CPR, Padilla C, Deleyiannis F, Christensen K, Poletta FA, Orioli IM, Hecht JT, Buxó CJ, Butali A, Adeyemo WL, Vieira AR, Shaffer JR, Murray JC, Weinberg SM, Leslie EJ, Marazita ML. Genome-Wide Association Study of Non-syndromic Orofacial Clefts in a Multiethnic Sample of Families and Controls Identifies Novel Regions. Front Cell Dev Biol. 2021;9:621482. Epub 20210409. doi: 10.3389/fcell.2021.621482. PubMed PMID: 33898419; PMCID: PMC8062975.

89.     Mukhopadhyay N, Feingold E, Moreno-Uribe L, Wehby G, Valencia-Ramirez LC, Restrepo Muñeton CP, Padilla C, Deleyiannis F, Christensen K, Poletta FA, Orioli IM, Hecht JT, Buxó CJ, Butali A, Adeyemo WL, Vieira AR, Shaffer JR, Murray JC, Weinberg SM, Leslie EJ, Marazita ML. Genome-wide association study of multiethnic nonsyndromic orofacial cleft families identifies novel loci specific to family and phenotypic subtypes. Genetic epidemiology. 2022;46(3-4):182-98. Epub 20220222. doi: 10.1002/gepi.22447. PubMed PMID: 35191549; PMCID: PMC9086172.

90.     Leslie EJ, Carlson JC, Shaffer JR, Feingold E, Wehby G, Laurie CA, Jain D, Laurie CC, Doheny KF, McHenry T, Resick J, Sanchez C, Jacobs J, Emanuele B, Vieira AR, Neiswanger K, Lidral AC, Valencia-Ramirez LC, Lopez-Palacio AM, Valencia DR, Arcos-Burgos M, Czeizel AE, Field LL, Padilla CD, Cutiongco-de la Paz EM, Deleyiannis F, Christensen K, Munger RG, Lie RT, Wilcox A, Romitti PA, Castilla EE, Mereb JC, Poletta FA, Orioli IM, Carvalho FM, Hecht JT, Blanton SH, Buxo CJ, Butali A, Mossey PA, Adeyemo WL, James O, Braimah RO, Aregbesola BS, Eshete MA, Abate F, Koruyucu M, Seymen F, Ma L, de Salamanca JE, Weinberg SM, Moreno L, Murray JC, Marazita ML. A multi-ethnic genome-wide association study identifies novel loci for non-syndromic cleft lip with or without cleft palate on 2p24.2, 17q23 and 19q13. Hum Mol Genet. 2016;25(13):2862-72. Epub 2016/04/02. doi: 10.1093/hmg/ddw104. PubMed PMID: 27033726; PMCID: PMC5181632.

91.     Kanazawa I, Takeno A, Tanaka KI, Notsu M, Sugimoto T. Osteoblast AMP-Activated Protein Kinase Regulates Postnatal Skeletal Development in Male Mice. Endocrinology. 2018;159(2):597-608. Epub 2017/11/11. doi: 10.1210/en.2017-00357. PubMed PMID: 29126229.

92.     Li ZW, Zhu YR, Zhou XZ, Zhuo BB, Wang XD. microRNA-135b expression silences Ppm1e to provoke AMPK activation and inhibit osteoblastoma cell proliferation. Oncotarget. 2017;8(16):26424-33. doi: 10.18632/oncotarget.15477. PubMed PMID: 28460435; PMCID: PMC5432269.

93.     Chen M-B, Liu Y-Y, Cheng L-B, Lu J-W, Zeng P, Lu P-H. AMPKα phosphatase Ppm1E upregulation in human gastric cancer is required for cell proliferation. Oncotarget. 2017;8(19):31288-96. doi: 10.18632/oncotarget.16126. PubMed PMID: 28423719.

94.     Li P, Fan JB, Gao Y, Zhang M, Zhang L, Yang N, Zhao X. miR-135b-5p inhibits LPS-induced TNFα production via silencing AMPK phosphatase Ppm1e. Oncotarget. 2016;7(47):77978-86. Epub 2016/10/30. doi: 10.18632/oncotarget.12866. PubMed PMID: 27793001; PMCID: PMC5363637.

95.     Fan JB, Ruan JW, Liu W, Zhu LQ, Zhu XH, Yi H, Cui SY, Zhao JN, Cui ZM. miR-135b expression downregulates Ppm1e to activate AMPK signaling and protect osteoblastic cells from dexamethasone. Oncotarget. 2016;7(43):70613-22. Epub 2016/09/24. doi: 10.18632/oncotarget.12138. PubMed PMID: 27661114; PMCID: PMC5342578.

96.     Zhang BH, Shi JY, Lin YS, Shi B, Jia ZL. VAX1 gene associated non-syndromic cleft lip with or without palate in Western Han Chinese. Arch Oral Biol. 2018;95:40-3. Epub 20180718. doi: 10.1016/j.archoralbio.2018.07.014. PubMed PMID: 30048854.

97.     Dixon MJ, Marazita ML, Beaty TH, Murray JC. Cleft lip and palate: understanding genetic and environmental influences. Nat Rev Genet. 2011;12(3):167-78. doi: 10.1038/nrg2933. PubMed PMID: 21331089; PMCID: PMC3086810.

98.     Hallonet M, Hollemann T, Pieler T, Gruss P. Vax1, a novel homeobox-containing gene, directs development of the basal forebrain and visual system. Genes Dev. 1999;13(23):3106-14. doi: 10.1101/gad.13.23.3106. PubMed PMID: 10601036; PMCID: PMC317183.

99.     Nikopensius T, Birnbaum S, Ludwig KU, Jagomägi T, Saag M, Herms S, Knapp M, Hoffmann P, Nöthen MM, Metspalu A, Mangold E. Susceptibility locus for non-syndromic cleft lip with or without cleft palate on chromosome 10q25 confers risk in Estonian patients. Eur J Oral Sci. 2010;118(3):317-9. doi: 10.1111/j.1600-0722.2010.00741.x. PubMed PMID: 20572868.

100.    Butali A, Suzuki S, Cooper ME, Mansilla AM, Cuenco K, Leslie EJ, Suzuki Y, Niimi T, Yamamoto M, Ayanga G, Erkhembaatar T, Furukawa H, Fujiwawa K, Imura H, Petrin AL, Natsume N, Beaty TH, Marazita ML, Murray JC. Replication of genome wide association identified candidate genes confirm the role of common and rare variants in PAX7 and VAX1 in the etiology of nonsyndromic CL(P). American journal of medical genetics Part A. 2013;161a(5):965-72. Epub 2013/03/07. doi: 10.1002/ajmg.a.35749. PubMed PMID: 23463464; PMCID: PMC3634899.

101.    de Aquino SN, Messetti AC, Bagordakis E, Martelli-Júnior H, Swerts MS, Graner E, Coletta RD. Polymorphisms in FGF12, VCL, CX43 and VAX1 in Brazilian patients with nonsyndromic cleft lip with or without cleft palate. BMC Med Genet. 2013;14:53. Epub 20130516. doi: 10.1186/1471-2350-14-53. PubMed PMID: 23679094; PMCID: PMC3660181.

102.    Wang J, Roy SK, Richard SA, Xu Y. Genome-wide Association Studies of REST Gene Associated Neurological Diseases/traits with Related Single Nucleotide Polymorphisms. Curr Neurovasc Res. 2023;20(3):410-22. doi: 10.2174/1567202620666230727153306. PubMed PMID: 37518996; PMCID: PMC10556398.

103.    Park JG, Tischfield MA, Nugent AA, Cheng L, Di Gioia SA, Chan WM, Maconachie G, Bosley TM, Summers CG, Hunter DG, Robson CD, Gottlob I, Engle EC. Loss of MAFB Function in Humans and Mice Causes Duane Syndrome, Aberrant Extraocular Muscle Innervation, and Inner-Ear Defects. Am J Hum Genet. 2016;98(6):1220-7. Epub 20160512. doi: 10.1016/j.ajhg.2016.03.023. PubMed PMID: 27181683; PMCID: PMC4908193.

104.    Gemelli C, Montanari M, Tenedini E, Zanocco Marani T, Vignudelli T, Siena M, Zini R, Salati S, Tagliafico E, Manfredini R, Grande A, Ferrari S. Virally mediated MafB transduction induces the monocyte commitment of human CD34+ hematopoietic stem/progenitor cells. Cell Death Differ. 2006;13(10):1686-96. Epub 20060203. doi: 10.1038/sj.cdd.4401860. PubMed PMID: 16456583.

105.    Kim H. The transcription factor MafB promotes anti-inflammatory M2 polarization and cholesterol efflux in macrophages. Sci Rep. 2017;7(1):7591. Epub 20170808. doi: 10.1038/s41598-017-07381-8. PubMed PMID: 28790455; PMCID: PMC5548719.

106.    Meng Y, Li J, Ye Z, Yin Z, Sun Q, Liao Z, Li G, Deng J, Liu L, Yu Y, Wu L, Zhou H, Shen N. MicroRNA-148a facilitates inflammatory dendritic cell differentiation and autoimmunity by targeting MAFB. JCI Insight. 2020;5(8). Epub 20200423. doi: 10.1172/jci.insight.133721. PubMed PMID: 32213710; PMCID: PMC7205423.

107.    Russell R, Carnese PP, Hennings TG, Walker EM, Russ HA, Liu JS, Giacometti S, Stein R, Hebrok M. Loss of the transcription factor MAFB limits β-cell derivation from human PSCs. Nat Commun. 2020;11(1):2742. Epub 20200602. doi: 10.1038/s41467-020-16550-9. PubMed PMID: 32488111; PMCID: PMC7265500.

108.    Fontoura C, Silva RM, Granjeiro JM, Letra A. Further evidence of association of the ABCA4 gene with cleft lip/palate. Eur J Oral Sci. 2012;120(6):553-7. Epub 20121015. doi: 10.1111/eos.12001. PubMed PMID: 23167473; PMCID: PMC4438764.

109.    Lennon CJ, Birkeland AC, Nuñez JA, Su GH, Lanzano P, Guzman E, Celis K, Eisig SB, Hoffman D, Rendon MT, Ostos H, Chung WK, Haddad J, Jr. Association of candidate genes with

nonsyndromic clefts in Honduran and Colombian populations. Laryngoscope. 2012;122(9):2082-7. Epub 20120702. doi: 10.1002/lary.23394. PubMed PMID: 22753311.

110.    Pan Y, Zhang W, Du Y, Tong N, Han Y, Zhang H, Wang M, Ma J, Wan L, Wang L. Different roles of two novel susceptibility loci for nonsyndromic orofacial clefts in a Chinese Han population. Am J Med Genet A. 2011;155a(9):2180-5. Epub 20110810. doi: 10.1002/ajmg.a.34170. PubMed PMID: 21834038.

111.    Yuan Q, Blanton SH, Hecht JT. Association of ABCA4 and MAFB with non-syndromic cleft lip with or without cleft palate. Am J Med Genet A. 2011;155a(6):1469-71. Epub 20110512. doi: 10.1002/ajmg.a.33940. PubMed PMID: 21567910; PMCID: PMC3711398.

112.    Koss H, Bunney TD, Behjati S, Katan M. Dysfunction of phospholipase Cγ in immune disorders and cancer. Trends in biochemical sciences. 2014;39(12):603-11. Epub 20141030. doi: 10.1016/j.tibs.2014.09.004. PubMed PMID: 25456276.

113.    Cockcroft S, Thomas GM. Inositol-lipid-specific phospholipase C isoenzymes and their differential regulation by receptors. Biochem J. 1992;288 ( Pt 1)(Pt 1):1-14. doi: 10.1042/bj2880001. PubMed PMID: 1332691; PMCID: PMC1132071.

114.    Tao P, Han X, Wang Q, Wang S, Zhang J, Liu L, Fan X, Liu C, Liu M, Guo L, Lee PY, Aksentijevich I, Zhou Q. A gain-of-function variation in PLCG1 causes a new immune dysregulation disease. J Allergy Clin Immunol. 2023. Epub 20230706. doi: 10.1016/j.jaci.2023.06.020. PubMed PMID: 37422272.

115.    Moreno Uribe LM, Fomina T, Munger RG, Romitti PA, Jenkins MM, Gjessing HK, Gjerdevik M, Christensen K, Wilcox AJ, Murray JC, Lie RT, Wehby GL. A Population-Based Study of Effects of Genetic Loci on Orofacial Clefts. J Dent Res. 2017;96(11):1322-9. Epub 20170629. doi: 10.1177/0022034517716914. PubMed PMID: 28662356; PMCID: PMC5607104.

116.    Diaz Perez KK, Curtis SW, Sanchis-Juan A, Zhao X, Head T, Ho S, Carter B, McHenry T, Bishop MR, Valencia-Ramirez LC, Restrepo C, Hecht JT, Uribe LM, Wehby G, Weinberg SM, Beaty TH, Murray JC, Feingold E, Marazita ML, Cutler DJ, Epstein MP, Brand H, Leslie EJ. Rare variants found in clinical gene panels illuminate the genetic and allelic architecture of orofacial clefting. Genet Med. 2023;25(10):100918. Epub 20230615. doi: 10.1016/j.gim.2023.100918. PubMed PMID: 37330696; PMCID: PMC10592535.

117.    Leslie EJ, Mansilla MA, Biggs LC, Schuette K, Bullard S, Cooper M, Dunnwald M, Lidral AC, Marazita ML, Beaty TH, Murray JC. Expression and mutation analyses implicate ARHGAP29 as the etiologic gene for the cleft lip with or without cleft palate locus identified by genome-wide association on chromosome 1p22. Birth Defects Res A Clin Mol Teratol. 2012;94(11):934-42. Epub 2012/09/26. doi: 10.1002/bdra.23076. PubMed PMID: 23008150; PMCID: PMC3501616.

118.    Suzuki M, Mizuno A. A novel human Cl(-) channel family related to Drosophila flightless locus. J Biol Chem. 2004;279(21):22461-8. Epub 20040308. doi: 10.1074/jbc.M313813200. PubMed PMID: 15010458.

119.    Halleran AD, Sehdev M, Rabe BA, Huyck RW, Williams CC, Saha MS. Characterization of tweety gene (ttyh1-3) expression in Xenopus laevis during embryonic development. Gene Expr Patterns. 2015;17(1):38-44. Epub 20141223. doi: 10.1016/j.gep.2014.12.002. PubMed PMID: 25541457; PMCID: PMC4402287.

120.    Yu AC, Zambrano RM, Cristian I, Price S, Bernhard B, Zucker M, Venkateswaran S, McGowan-Jordan J, Armour CM. Variable developmental delays and characteristic facial features-A novel 7p22.3p22.2 microdeletion syndrome? Am J Med Genet A. 2017;173(6):1593-600. Epub 20170425. doi: 10.1002/ajmg.a.38241. PubMed PMID: 28440577.

121.     Huang X, Li Q, Zheng X, Jiang C. TTYH3 Promotes Cervical Cancer Progression by Activating the Wnt/β-Catenin Signaling Pathway. Cancer Invest. 2024;42(8):726-39. Epub 20240827. doi: 10.1080/07357907.2024.2395014. PubMed PMID: 39189652.

122.     Liu H, Leslie EJ, Carlson JC, Beaty TH, Marazita ML, Lidral AC, Cornell RA. Identification of common non-coding variants at 1p22 that are functional for non-syndromic orofacial clefting. Nat Commun. 2017;8:14759. Epub 2017/03/14. doi: 10.1038/ncomms14759. PubMed PMID: 28287101; PMCID: PMC5355807.

123.     Ranji P, Pairet E, Helaers R, Bayet B, Gerdom A, Gil-da-Silva-Lopes VL, Revencu N, Vikkula M. Four putative pathogenic ARHGAP29 variants in patients with non-syndromic orofacial clefts (NsOFC). Eur J Hum Genet. 2024. Epub 20241106. doi: 10.1038/s41431-024-01727-3. PubMed PMID: 39506048.

124.     Conte F, Oti M, Dixon J, Carels CE, Rubini M, Zhou H. Systematic analysis of copy number variants of a large cohort of orofacial cleft patients identifies candidate genes for orofacial clefts. Hum Genet. 2016;135(1):41-59. Epub 2015/11/13. doi: 10.1007/s00439-015-1606-x. PubMed PMID: 26561393; PMCID: PMC4698300.

125.     Dickinson AJ, Sive HL. The Wnt antagonists Frzb-1 and Crescent locally regulate basement membrane dissolution in the developing primary mouth. Development. 2009;136(7):1071-81. Epub 20090218. doi: 10.1242/dev.032912. PubMed PMID: 19224982; PMCID: PMC2685928.

126.     Hoang B, Moos M, Jr., Vukicevic S, Luyten FP. Primary structure and tissue distribution of FRZB, a novel protein related to Drosophila frizzled, suggest a role in skeletal morphogenesis. J Biol Chem. 1996;271(42):26131-7. doi: 10.1074/jbc.271.42.26131. PubMed PMID: 8824257.

127.     Hoang BH, Thomas JT, Abdul-Karim FW, Correia KM, Conlon RA, Luyten FP, Ballock RT. Expression pattern of two Frizzled-related genes, Frzb-1 and Sfrp-1, during mouse embryogenesis suggests a role for modulating action of Wnt family members. Dev Dyn. 1998;212(3):364-72. doi: 10.1002/(sici)1097-0177(199807)212:3<364::Aid-aja4>3.0.Co;2-f. PubMed PMID: 9671940.

128.     Kamel G, Hoyos T, Rochard L, Dougherty M, Kong Y, Tse W, Shubinets V, Grimaldi M, Liao EC. Requirement for frzb and fzd7a in cranial neural crest convergence and extension mechanisms during zebrafish palate and jaw morphogenesis. Dev Biol. 2013;381(2):423-33. Epub 20130625. doi: 10.1016/j.ydbio.2013.06.012. PubMed PMID: 23806211.

129.     Lin K, Wang S, Julius MA, Kitajewski J, Moos M, Jr., Luyten FP. The cysteine-rich frizzled domain of Frzb-1 is required and sufficient for modulation of Wnt signaling. Proc Natl Acad Sci U S A. 1997;94(21):11196-200. doi: 10.1073/pnas.94.21.11196. PubMed PMID: 9326585; PMCID: PMC23413.

130.     Janečková E, Feng J, Guo T, Han X, Ghobadi A, Araujo-Villalba A, Rahman MS, Ziaei H, Ho TV, Pareek S, Alvarez J, Chai Y. Canonical Wnt signaling regulates soft palate development by mediating ciliary homeostasis. Development. 2023;150(5). Epub 20230309. doi: 10.1242/dev.201189. PubMed PMID: 36825984; PMCID: PMC10108707.