FedWeight: Mitigating Covariate Shift of Federated Learning on Electronic Health Records Data through Patients Re-weighting

He Zhu[1,5], Na Li[6], Xiaoxiao Li[7], Dianbo Liu[3,4*], David Buckeridge[2*], and Yue Li[1,5*]

[1]*School of Computer Science, McGill University*
[2]*School of Population and Global Health, McGill University*
[3]*School of Medicine, National University of Singapore*
[4]*College of Design and Engineering, National University of Singapore*
[5]*Mila - Quebec AI Institute*
[6]*Community Health Sciences, University of Calgary*
[7]*Electrical and Computer Engineering, University of British Columbia*
[*]*Correspondence to dianbo@nus.edu.sg, david.buckeridge@mcgill.ca, yueli@cs.mcgill.ca*

Abstract

Federated Learning (FL) has emerged as a promising approach for research on real-world medical data distributed across different organizations, as it allows analysis of distributed data while preserving patient privacy. However, one of the prominent challenges in FL is covariate shift, where data distributions differ significantly across different clinical sites, like hospitals and outpatient clinics. These differences in demographics, clinical practices, and data collection processes may lead to significant performance degradation of the shared model when deployed for a target population. In this study, we propose a Federatively Weighted (FedWeight) framework to mitigate the effect of covariate shift on Federated Learning. Leveraging the data distribution estimated by density estimator models, we re-weight the patients from the source clinical sites, making the trained model aligned with the data distribution of the target site, thus mitigating the covariate shift between source and target sites. To make our approach also applicable to unsupervised learning, we integrate FedWeight into a novel federated embedded topic model (ETM), namely FedWeight-ETM. We evaluated FedWeight in cross-site FL within the eICU dataset and also cross-dataset FL between eICU and MIMIC-III data. Compared with the baseline, FedWeight-corrected FL models demonstrate superior performance for predicting patient mortality, ventilator use, sepsis diagnosis, and length of stay in the intensive care unit (ICU). Moreover, FedWeight outperforms FedAvg in identifying important features relevant to the clinical outcomes. Leveraging

Shapley Additive Explanations (SHAP), the FedWeight-corrected classifiers reveal subtle yet significant associations between drugs, lab tests, and patient outcomes. Using FedWeight-ETM, we identified known disease topics involving renal or heart failure predictive of future mortality at the ICU readmission. Together, FedWeight provides a robust FL framework to address the challenge of covariate shift from clinical silos in predicting critical patient outcomes and providing meaningful clinical features.

1   Introduction

Training machine learning models on large-scale electronic health record (EHR) data is promising for advancing medical research and improving patient outcomes [1]. However, EHR data are from different healthcare institutions are not easily pooled, because it is difficult to move these data out of institutions due to laws and regulations about data privacy and data governance, and transmission costs [2]. To enable machine learning in the context of these challenges, Federated Learning (FL) trains the model on local datasets that remain in each healthcare institution and shares only the model parameters with the central server for model averaging [3]. Although FL has been widely used in the clinical settings [4–6], traditional FL assumes the same data distributions for each silo [7–12]. This is an unrealistic assumption because of covariate shifts due to differences in patient demographics, clinical practices, and data collection methods between institutions [13, 14]. Such disparities may lead to poor performance on out-of-distribution (OOD) clinical data, resulting in inaccurate predictions and uninterpretable clinical outcomes. For instance, a federated model trained on data from specialized hospitals may be unable to accurately predict patient outcomes at a community hospital. Therefore, given the significance of AI safety and quality [15–18], it is essential to develop a framework to mitigate the effect of covariate shifts on FL, thus providing more robust inference to enhance disease prevention strategies and promote fairness in healthcare decisions and outcomes [19–21].

Recently, Shimodaira *et al.* introduce the weighted log-likelihood method to address covariate shift using importance sampling, assigning weights to the training loss based on the ratio of test to training input densities [13]. Building upon their work, methods have been proposed to estimate the reweighting ratios using Kernel Mean Matching [22, 23], as well as Kullback-Leibler Importance Estimation Procedure (KLIEP) [24], which have been extensively employed to alleviate covariate shifts in centralized learning environments. However, all these methods rely on access to both training and test samples of all patients to estimate the reweighting ratios, which is impractical in FL settings due to privacy constraints. To mitigate this challenge, FedCL and FedDNA employ a model-level reweighting approach by sharing statistical model parameters across clients [25, 26], but their improvements are limited as the reweighting is estimated solely at the model level. In contrast, several existing methods estimate the sample-level reweights by sharing unlabelled data across clients [27, 28], which may increase communication costs and compromise data privacy, thus violating FL principles. To address this issue, FedDisk trains a global density estimator along with multiple local density estimators to compute reweighting ratios [29], but their experiments demonstrate only minor improvements on real-world image

data. In conclusion, existing methods either provide limited performance improvements or compromise privacy, making them unsuitable for healthcare, where patient confidentiality is paramount. Given that FL in healthcare is still nascent, with limited algorithms addressing co-variate shifts [30, 31], there is a pressing need to develop a privacy-preserving solution with improved performance in clinical settings.

In this study, we first describe covariate shifts in two widely used public EHR datasets, namely the eICU Collaborative Research Database (eICU) [32] and the Medical Information Mart for Intensive Care III (MIMIC-III) [33]. We then present a novel FL framework — Federated Weighted Log-likelihood (FedWeight). Specifically, FedWeight incorporates the weighted log-likelihood method within the federated framework [13], which probabilistically re-weights the patients from the source clinical sites, aligning the trained model with the data distribution of the target site. FedWeight can be applied to both supervised and unsupervised tasks. We evaluate our framework using eICU and MIMIC-III datasets. Within the eICU, we perform cross-hospital FL. We then conduct federated training between eICU and MIMIC-III, while addressing their distribution differences. The experiments demonstrate that compared with FedAvg, our approach can provide more accurate predictions of patient mortality, ventilator use, sepsis diagnosis, and ICU length of stay.

2   Results

2.1   ML-based data harmonization reduces administrative differences between eICU hospitals

Different hospitals may adopt distinct administration practices, leading to variations in naming the same drug across clinical sites. This motivates us to harmonize the EHR data before model training. Specifically, in the eICU dataset, different hospitals may use distinct encoding of drug administration. Some may use the generic name (e.g. "acetylsalicylic acid"), while others use the trade name (e.g. "aspirin"), although both refer to the same drug. Moreover, the dosage information is included in some drug names (e.g. "aspirin 10 mg") but not all. Consequently, there is almost no overlap in drug encodings across hospitals (Figure 2c). Further, over 40% of drug names are not recorded (Figure 2a), although some of them have Hierarchical Ingredient Code List (HICL) codes. However, these codes are not widely used in other hospitals, such as those in MIMIC-III, which impedes the training of FL models across hospitals and datasets. As a result, we observed distinct clusters of patients by hospitals (Figure 1a). To address this, we developed a method to impute missing drug names. We also proposed a drug harmonization framework to combine drugs with similar identities and exclude dosage information (**Methods** Section 4.3). This preprocessing step successfully decreased the unrecorded drug proportions to approximately 20% (Figure 2b) and increased the number of common drugs to over 90%. The clustering of patients from different hospitals shows better mixing, although some hospitals still exhibit distinct clusters. (Figure 1b). To prepare for model training across datasets, we also

Figure 1: Visualization of covariate shift in the eICU and MIMIC-III dataset. **a** - **b.** UMAP projection of drug data across hospitals on eICU before and after drug harmonization. The UMAP-reduced drug data is visualized in a scatter plot, with each data point coloured by its respective hospital. **c.** UMAP projection of harmonized medication data from eICU and MIMIC-III. **d** - **g.** Distribution differences in patient demographics, calculated as the proportion of patients in each demographic group within each hospital. **h** - **j.** In-hospital and out-hospital distribution by ML-based density estimators. For each hospital, the in-hospital estimate was calculated by applying its own density estimator to its patient data. The out-hospital estimate was computed by applying the same density estimator to patient data from a different hospital.

harmonized data between two data domains namely eICU and MIMIC-III (Figure 1c). Interestingly, we observed two distinct clusters from the MIMIC-III drug data (Figure S1a). Enrichment

Figure 2: Effect of drug imputation and drug harmonization in the eICU dataset. **a** - **b.** The drug rates before and after imputation. Since some drugs may lack recorded names, drug rates indicate the proportion of administered drugs with recorded names over all unique drugs. **c** - **d.** Percentage of overlapped drugs across hospitals before and after harmonization. The percentage was computed as the proportion of drugs with recorded names present in both Hospital A and B relative to all drugs in Hospital B.

analysis revealed that one cluster is significantly associated with planned hospital admissions (elective ICU admission) and cardiovascular surgery patients (Figure S1b-d). These covariate shifts due to the heterogeneous distributions between hospitals and between datasets motivate us to develop FedWeight as experimented next.

2.2 Heterogeneous patient demographics across eICU hospitals

In addition to the drug data, we also observed demographic differences among the hospitals in the eICU dataset. Specifically, we analyzed the distribution of patients by age, sex, BMI, and ethnicity within each hospital (Figure 1d-g). Although most patients were aged from 50 to 89,

Hospital 148 had a higher proportion of younger patients (< 30) (Figure 1d). In addition, Hospital 458 had a higher proportion of ethnically African patients, whereas Hospitals 167 and 165 had more Native American patients (Figure 1e). Despite the overall low proportion of underweight patients across all hospitals, Hospital 199 manifested a slightly lower proportion of underweight patients and a higher proportion of obese patients (Figure 1f). Furthermore, while most hospitals had more male patients than female patients, Hospital 283 had an almost equal proportion of male and female patients (Figure 1g). These discrepancies in patient demographics across hospitals may hinder the effective generalization of models in FL settings.

2.3   Quantifying data likelihood by ML-based density estimators

To address the problem of covariate shift, we first measured data distributions using model-based approach. To this end, we experimented with 3 deep learning density estimators, namely Masked Autoencoder for Density Estimation (MADE) [34], Variational Autoencoder (VAE) [35], and Vector Quantized Variational Autoencoder (VQ-VAE) [36]. Through each model, we can compare the data likelihoods between the training data from one hospital and test data from another hospital. In most hospitals, the in-hospital likelihood is significantly larger than the out-hospital likelihood (Figure 1h-j), indicating that each hospital's model aligns more closely with its own data than other hospitals' data.

2.4   Proposed FedWeight to address covariate shift

Putting the above together, we developed **FedWeight**, a machine learning method to re-weight the patients from source hospitals, aligning the trained model with the data distribution observed in the target hospital (Figure 3a-d). Specifically, the target hospital shares its density estimator with all source hospitals, which use it to calculate the reweighting ratio to train their local models. Intuitively, we assign larger weights to the patients from the source hospitals whose data distribution is similar to the target population and smaller weights to those with dissimilar data distribution. Consequently, the trained model is better aligned with the target hospital data distribution, thereby effectively addressing the covariate shift problem in the FL settings. During training, only model parameters are shared between the target hospital and the source hospitals, thus effectively safeguarding patient privacy.

This method facilitates adapting source hospital data to the target. Additionally, we could also allow hospitals to mutually benefit from each other. To achieve this, we developed **Symmetric FedWeight** (Figure 3e-h). For simplicity, we assumed a federated network with two hospitals, which can be easily extended to multiple hospitals. These hospitals will treat each other as the target and themselves as the source to calculate the reweighting ratio (Figure 3g). The symmetric strategy enables both hospitals to adapt to each other's data distributions, ensuring mutual benefit.

Figure 3: FedWeight training process. **a.** The target and source hospitals independently train their density estimators. **b.** The target hospital shares its density estimator with all source hospitals. **c.** Each source hospital calculates a patient-specific re-weight, which is used to train its local model. **d.** The target hospital aggregates the parameters from the source hospitals using the FL algorithm, which better generalizes the target hospital's data. Symmetric FedWeight training process. **e.** Hospital A and B independently train their density estimators. **f.** Hospital A and B share their density estimators to each other. **g.** Hospital A and B regard each other as targets, estimating the re-weight, which is used to train its local model. **h.** Aggregate the local models from Hospital A and B using FL algorithm.

2.5 Simulation study

To evaluate the effectiveness of our method, we first undertook simulation experiments on patients' data. We trained the FedWeight model on the simulation dataset by employing density estimators, namely MADE, VAE, and VQ-VAE. Given our imbalanced labels, the Area Under the Precision-Recall Curve (AUPRC) was selected as the evaluation metric for model prediction. We observed that all FedWeight models achieve an average AUPRC of 0.923, which significantly surpassed that of FedAvg, which had an average AUPRC of 0.917 (Wilcoxon test p-value < 0.05) (Figure S2a). This result demonstrates that our proposed model has enhanced predictive capability over this baseline with the datasets examined.

We also examined FedWeight's capability in identifying influential features. Specifically, we evaluated the trained model by comparing its weights to a sparse reference model that simulates true influential features, using AUPRC as the evaluation metric. FedWeight models, when trained with density estimators such as VAE, VQ-VAE, and MADE, achieve comparable AUPRC results for influential features. However, they all significantly outperform the baseline. This demonstrates FedWeight has enhanced capacity to identify influential features.

Additionally, we computed the Pearson correlation between the weights of the reference model and the trained model. We observed that FedWeight exhibits a higher correlation with the reference model's weights (Figure S2c). Therefore, FedWeight is more effective in capturing influential features, thus having better model interpretability.

2.6 Predicting critical outcomes from eICU data

Accurately predicting critical clinical events can drastically improve patient outcomes, especially in the ICU. We conducted experiments to predict four outcomes, namely mortality, ventilator use, sepsis, and ICU length of stay, using the first 48 hours of data from the eICU dataset (Section 4.3). FedWeight outperformed FedAvg for most hospitals (Figure 4a-d). Moreover, it also demonstrated comparable performance to the centralized model.

For mortality prediction, Hospital 458 and 420 had the best results for all methods (Figure 4a). FedWeight with the VQ-VAE density estimator provided superior results in all target hospitals (Wilcoxon test p-value < 0.05). Moreover, all FedWeight variants surpassed FedAvg in performance. They also show comparable performance to the centralized model.

For ventilator prediction, Hospital 167 and 458 demonstrate the highest AUPRC for all methods (Figure 4b), as they had the most patients and fewer imbalanced labels. Moreover, FedWeight exceeded the AUPRC of FedAvg, especially in Hospital 252, 420, 458. Furthermore, the overall performance was close to that of the centralized model. FedWeight with the VAE density estimator demonstrated the best performance in all target hospitals.

For sepsis prediction, Hospital 420, with fewer imbalanced labels, yielded the most favourable outcomes for all methods (Figure 4c). Again, the FedWeight strategies also significantly outperformed FedAvg. FedWeight using MADE and VAE provideed the most accurate results in all target hospitals.

Figure 4: Performance comparison of federated models across clinical outcome predictions, including mortality, ventilator, sepsis, and ICU length of stay prediction. FedWeight methods were compared with FedAvg and a centralized model (i.e., model trained on the pooled eICU data and corrected by the covariate indicator variable for each hospital). **a** - **d.** Performance of clinical outcome predictions on eICU, which was evaluated on a bootstrap-sampled dataset from five target hospitals (167, 199, 252, 420, and 458), with mean and standard deviation computed from the bootstrap samples. **e.** Performance of cross-dataset federated model trained on eICU and evaluated on a bootstrapped test set of MIMIC-III. One-sided Wilcoxon test p-values were calculated to compare FedWeight with FedAvg. * denotes the p-values < 0.05, and ** represents the p-values < 0.01.

For predicting ICU length of stay, most FedWeight variants demonstrated lower loss com-

pared to FedAvg and exhibited performance close to that of the benchmark centralized model, especially in Hospital 199 (Figure 4d).

Together, FedWeight surpassesed the baseline in predicting mortality, ventilator use, sepsis diagnosis, and ICU length of stay across various hospitals, highlighting its effectiveness in clinical predictions.

2.7 Cross-dataset federated learning

In practice, each EHR dataset often requires separate access approval, making it difficult to share or centralize the data. In this scenario, models trained on each dataset can be pooled via the FL framework. To this end, we performed cross-dataset analysis by training our model on all hospitals from the eICU dataset and making predictions on MIMIC-III (Figure 4e). As expected, the models trained on eICU hospitals demonstrate worse performance on MIMIC-III (Figure 4e), compared to the performance on the held-out patients from eICU (Figure 4a-d). Even so, FedWeight outperforms FedAvg, especially in ventilator use and ICU length of stay predictions, with all FedWeight variants significantly surpassing FedAvg (Wilcoxon test p-value < 0.01) and achieving results comparable to the centralized model (i.e., model trained on the pooled eICU data and corrected by the covariate indicator variable for each hospital). For mortality prediction, FedWeight using MADE and VAE as density estimators achieve significantly higher AUPRC (Wilcoxon test p-value < 0.05) (Figure 4e). In sepsis prediction, FedWeight models with VAE and VQ-VAE demonstrate significantly superior performance (Wilcoxon test p-value < 0.05). In addition, all FedWeight variants show statistical significance in ventilator and ICU length of stay prediction (Wilcoxon test p-value < 0.01). These findings indicate that FedWeight is capable of addressing the distribution differences across different datasets.

2.8 Detecting clinical features by FedWeight+SHAP analysis

We sought to assess whether FedWeight improves detecting relevant features for the clinical outcome predictions using Shapley Additive Explanations (SHAP) [37]. As a reference, we computed the SHAP values of the centralized model. We used the Pearson correlation of the SHAP values between the federated model and the centralized models as the evaluation metric. We first performed the experiments using hospitals within the eICU dataset. FedWeight demonstrated a superior correlation of SHAP values compared to FedAvg across almost all hospitals, especially in predicting ventilator use. Specifically, for mortality prediction, all the FedWeight methods significantly outperform the baseline method (Wilcoxon test p-value < 0.01), except for Hospital 458 (Figure 5a). Moreover, the FedWeight model using MADE and VAE density estimator demonstrated significantly higher correlation of SHAP values in all hospitals (Wilcoxon test p-value < 0.01). For ventilator prediction, FedWeight using VAE and VQ-VAE density estimator significantly outperformed FedAvg in all target hospitals (Wilcoxon test p-value < 0.01) (Figure 5b). Regarding sepsis prediction, FedWeightVAE demonstrated the most significant results in all target hospitals (Wilcoxon test p-value < 0.05) (Figure 5c). For the length of stay

Figure 5: Comparison of FedWeight and FedAvg in capturing influential features. SHAP values were leveraged to identify feature importance, calculated from test data fed into the trained model. For mortality and ICU length of stay prediction, SHAP values were summed across all samples. For ventilator and sepsis prediction, SHAP values were aggregated across all time windows and samples, resulting in a one-dimensional vector of feature importance. We then calculated the Pearson correlation of feature importance between the federated and centralized model. **a** - **d**. Pearson correlation of SHAP-based feature importance for clinical outcome predictions in eICU. The models were validated on the bootstrapped test sets of five target hospitals (167, 199, 252, 420, 458). **e.** Pearson correlation of SHAP-based feature importance for cross-dataset federated learning, where models were trained on eICU, and the correlation was computed on the bootstrapped test set of MIMIC-III. One-sided Wilcoxon test p-values were calculated against the baseline. * denotes the p-values < 0.05, and ** represents the p-values < 0.01.

prediction, FedWeightMADE demonstrated significantly stronger correlations with the benchmark across all target hospitals, whilst FedWeightVAE showed significantly higher correlations in Hospital 167, 252, 420, 458 (Wilcoxon test p-value < 0.01) (Figure 4d). To conclude, our experiments on the eICU dataset demonstrated that FedWeight SHAP values are more consistent with the benchmark SHAP values. As SHAP values quantify feature importance, this suggests that FedWeight is more effective in capturing influential features.

We further performed a cross-dataset analysis, using eICU hospitals as the source and MIMIC-III as the target. FedWeight demonstrated significantly higher correlations in predicting mortality, sepsis, and ICU length of stay (Wilcoxon test p-value < 0.01) (Figure 5e). Notably, FedWeight utilizing the VAE as the density estimator exhibited the best performance in mortality and sepsis prediction, underscoring its capacity to capture features highly associated with a patients' length of stay in the ICU.

2.9   Top drugs and lab tests attributed to the clinical outcomes

Given the strong quantitative results for the feature attributions, we turn to individual drugs and lab tests that exhibit high SHAP values based on the best FedWeight model, namely FedWeightVAE.

Mortality   We leveraged drug administration and lab tests from the initial 48 hours of ICU admission to predict patient mortality beyond this period. The drugs of the highest correlation with patient mortality are predominantly vasopressors or anaesthetics (Figure 6a). Specifically, glycopyrrolate is the foremost drug in mortality prediction, which reflects its role as an anticholinergic agent to manage respiratory secretions and mitigate vagal reflexes in critically ill patients during surgery [38]. Its association with mortality may serve as a proxy marker for high-acuity clinical scenarios, highlighting its relevance as a potential marker of severe physiological compromise in ICU settings. Following glycopyrrolate, vasopressors like vasopressin epinephrine, and phenylephrine emerge as critical prior to patients mortality, highlighting their role as primary stress hormones typically administered in the context of life-threatening conditions such as septic shock, cardiogenic shock, or profound hypotension [39–42]. Therefore, this correlation suggests that the necessity for vasopressor support often signals an advanced stage of critical illness, marked by high acuity and poor prognostic outcomes. After vasopressors, we also observed the administration of morphine prior to patient mortality in the ICU, which is likely attributable to its role in palliative care and the management of refractory pain and dyspnea in critically ill patients [43].

We identified a significant correlation between elevated blood urea nitrogen (BUN) levels and subsequent patient mortality, highlighting its utility as a pivotal biomarker for mortality risk in critical care (Figure 6b). Elevated BUN reflects underlying pathophysiological processes, including renal insufficiency, systemic hypoperfusion, and heightened protein catabolism, which are key indicators of severe illness [44–48]. Therefore, this association underscores the prognostic significance of BUN in stratifying patient risk and guiding therapeutic interventions. Additionally,

Figure 6: Feature importance of drug and lab tests for clinical outcome predictions identified by FedWeight+SHAP. **a.** SHAP values of the top 5 most important drugs identified by FedWeight for each clinical outcome. We computed SHAP values for each drugs present on target hospital. For mortality and ICU length of stay prediction, SHAP values were summed across all patients, while for ventilator and sepsis prediction, they were summed across both time windows and samples, resulting in a one-dimensional feature importance vector. For each clinical outcome, the top 5 most important drugs were selected, visualized with colour intensities indicating their feature importance. **b** - **e.** SHAP values of the top 12 most important lab tests for each clinical outcome, identified by FedWeightVAE. Each dot represents a sample from the target hospital. Features are ordered by their mean absolute SHAP value.

increased lactate levels are another profound indicators of mortality. This verifies the existing medical knowledge, as elevated lactate is a key clinical criterion for tissue hypoxia in critically ill patients, which is strongly associated with ICU mortality [49–52].

Ventilator use Since most patients initiate ventilation within 72 hours of ICU admission, we established a 72-hour observation window, further segmented into six 12-hour intervals. We aim to utilise drug administration and lab test results from each interval to predict ventilator use in the next. Additionally, we observed that 65.5% of patients in the eICU dataset experience multiple episodes of mechanical ventilation during their ICU stay. As a result, ventilator treatment may appear in multiple intervals for a single patient. Among all the drugs, propofol stands out for ventilator prediction, which is commonly administered in critical care to facilitate patient-ventilator synchrony, minimize agitation, and ensure tolerance of invasive respiratory support [53, 54]. However, since propofol is commonly administered during intubation, its association with subsequent ventilator use likely stems from multiple ventilation episodes throughout the ICU stay. Following propofol, our model identified chlorhexidine as a drug administered to ICU patients who subsequently require mechanical ventilation, which facilitates the identification of patients at high risk for ventilation, while enabling medical practitioners to proactively prepare for ventilatory support. This identification verifies the established clinical knowledge. As a broad-spectrum antiseptic with bactericidal properties, chlorhexidine is widely utilized in oral care protocols within the ICU to mitigate the microbial colonization of the oropharynx and trachea, which are primary precursors to ventilator-associated pneumonia (VAP) [55–57]. Our model also identified that etomidate is the third most commonly administered drug prior to ventilator use. Specifically, etomidate is a rapid-onset, short-acting intravenous anaesthetic commonly employed before endotracheal intubation. Although its administration might be associated with adrenal suppression, etomidate is particularly suitable for hemodynamically unstable patients due to its minimal impact on blood pressure [58, 59].

For ventilator prediction, elevated arterial blood gas (ABG) parameters, including oxygen saturation (O2sat), demonstrate a strong correlation with the subsequent initiation of ventilatory support (Figure 6c). While this may appear counterintuitive, as oxygen saturation measurements are recorded before the initiation of ventilation, one would anticipate these parameters to be elevated only after ventilator treatment begins. Multiple episodes of mechanical ventilation could explain this phenomenon. This pattern could also arise from the initial use of high-flow oxygen or non-invasive ventilation (e.g., CPAP or BiPAP) as a preparatory measure before transitioning to invasive mechanical ventilation. Therefore, it is conceivable that patients undergo an initial period of ventilatory support, exhibit elevated oxygen saturation values, and subsequently require re-initiation of mechanical ventilation. Moreover, we also observed the elevated blood urea nitrogen (BUN) levels are highly correlated with the subsequent ventilatory support. Such elevations often stem from renal insufficiency, hypovolemia, or heightened protein catabolism associated with conditions like sepsis or multiorgan dysfunction. These pathophysiological states frequently precede respiratory failure, necessitating the commencement of mechanical ventilation. Therefore, markedly elevated BUN levels may serve as an ICU

risk marker which often guides decisions on fluid management, dialysis, and ventilation timing.

Sepsis   Similar to ventilation use, we designed a 72-hour observation window comprising six 12-hour intervals. We aim to employ drug administration and lab tests from each interval to predict sepsis diagnosis in the subsequent interval. Confirming the existing medical knowledge and literature, antibiotics such as vancomycin, exhibit the highest overall feature attribution [60, 61], followed by piperacillin [62], cefepime [63], and metronidazole [64]. Notably, our model make prediction based on drugs administered prior to the sepsis diagnosis. Given that approximately 90% of sepsis cases are community-onset [65], with most patients presenting infection symptoms prior to sepsis diagnosis, it is routine for patients to receive these antibiotics before a formal diagnosis of sepsis.

For sepsis prediction, the white blood cell (WBC) count holds the highest SHAP value (Figure 6d), underscoring its strong association with septicemia and septic shock. Leukocytosis generally reflects the immune system's activation in response to bacterial infection, driven by pro-inflammatory cytokines [66]. Therefore, clinicians may leverage WBC with other laboratory markers, including creatinine, platelet count, and lactate, for early diagnosis and monitoring of the condition.

ICU length of stay   Using drug administration and lab tests from the first 48 hours of ICU admission, we aim to predict the remaining ICU stay duration. Patients receiving cardiovascular drugs, including midodrine, alteplase, and nicardipine, tend to have prolonged ICU stays (Figure 6a). This indicates that patients with cardiac surgery often require prolonged monitoring and extended care, as supported by existing research [67].

We observed an elevated platelet count as a strong indicator of prolonged ICU length of stay (Figure 6e). Specifically, thrombocytosis serves as a surrogate marker of heightened inflammatory activity and immune system activation. This hyperactive platelet response may signify the severity of the underlying pathology, such as infection, malignancy, or surgical recovery, all of which demand intensive and sustained care. Additionally, elevated platelet levels are indicative of a hypercoagulable state, predisposing patients to thrombotic complications, such as deep vein thrombosis or pulmonary embolism, which necessitates prolonged ICU stay for vigilant monitoring and therapeutic intervention [68]. Therefore, the association of thrombocytosis with prolonged ICU duration underscores its role as a biomarker of systemic stress and disease severity, providing insights into patient prognosis and resource allocation in the ICU.

Together, our combined FedWeight+SHAP analysis detected known drugs and lab tests associated with the clinical outcome predictions. This identification of influential variables warrants future investigation to explore causal mechanisms for the nuanced clinical associations identified in this analysis.

Figure 7: Federated latent topic modelling. **a.** FedAvg-ETM. In each local hospital $k \in K$, $\mathbf{X}_k$ is input into the encoder, whose output goes into two separate linear layers and produces $\boldsymbol{\mu}_k$ and $\boldsymbol{\sigma}_k$. Then, through the re-parameterization trick, we obtain the latent representation $\mathbf{Z}_k$. After applying the softmax function, $\mathbf{Z}_k$ gives the patient-topic distribution $\boldsymbol{\theta}_k$. The learnable topic embedding $\boldsymbol{\alpha}_k$ and ICD embedding $\boldsymbol{\rho}_k$ generate the topic-ICD mixture $\boldsymbol{\beta}_k$. Then, $\boldsymbol{\beta}_k$ is multiplied with $\boldsymbol{\theta}_k$ to reconstruct the input. During federated averaging, only the encoder network and the ICD embedding are uploaded to the target hospital for aggregation, whilst all other model parameters are kept locally updated. **b.** FedWeight-ETM. FedWeight-ETM builds on FedAvg-ETM by applying a re-weight to the log-likelihood term of the ELBO function for each patient.

2.10 Mortality-associated latent topics captured by FedWeight-ETM

In the aforementioned experiments, FedWeight demonstrated exceptional performance in supervised prediction tasks. To identify sets of correlated clinical features in an unsupervised setting, a topic model is a natural choice. To achieve this, we incorporated FedWeight into Embedded Topic Model (ETM) (**Methods** Section 4.2). First, we developed FedAvg-ETM, which shares the encoder weights and clinical feature embeddings between silos (i.e., eICU and MIMIC-III) to infer robust latent topics from distributed EHR data (Figure 7a). To address covariate shifts, we further extended FedAvg-ETM to FedWeight-ETM (Figure 7b). Quantitatively, we benchmarked the performances of non-federated ETM, FedAvg-ETM, and FedWeight-ETM by predicting readmission mortality using the corresponding topic mixtures as inputs to a logistic regression classifier. Overall, FedAvg-ETM and FedWeight-ETM demonstrated comparable performance and outperformed the baseline non-federated ETM across different topic numbers. Details were provided in **Appendix** Section S2.

We evaluated FedWeight-ETM's ability to capture topics related to patient mortality. We identified the top 5 topics that are significantly associated with patient mortality. Figure 8a and b display the top International Classification of Diseases (ICD) codes under each topic supported by the Fisher-exact test p-values for eICU and MIMIC-III data, respectively. For eICU, Topic 16 is the most significant topic with the top ICD code being chronic renal failure (ICD-585). Indeed, renal failure disrupts fluid, electrolyte, and metabolic equilibrium and ultimately systemic homeostasis. For critically ill patients, it exacerbates comorbidities such as sepsis and is a hallmark of multiorgan dysfunction syndrome (MODS), perpetuating systemic inflammation and hemodynamic instability, which increases the risks of patient mortality [69, 70]. As the second significant topic, Topic 31 involves cardiac dysrhythmias (ICD-427), which has direct impact on hemodynamic stability and causes critical conditions such as myocardial ischemia, heart failure, or sepsis. Moreover, ventricular arrhythmias and atrial fibrillation can lead to reduced cardiac output, tissue hypoperfusion, and multiorgan dysfunction in critically ill patients [71]. As the third focus, Topic 28 involves acute renal failure (ICD-584). We observed that while acute renal failure is strongly associated with mortality, its correlation is weaker than that of chronic renal failure. This may be due to its potential reversibility with timely medical intervention [72].

We also observed clinically meaningful topics inferred from the MIMIC-III dataset (Figure 8b). Namely, Topic 17 is characterized by septicemia (ICD-38), which triggers widespread endothelial damage, capillary leakage, and coagulopathy, culminating in septic shock and multi-organ failure [73]. Furthermore, for Topic 10, the top disease identified by our model is abnormal findings on examination of blood (ICD-790). Severe abnormalities in blood parameters—such as metabolic acidosis, electrolyte imbalances, coagulation abnormalities, and hematologic dyscrasias—can precipitate multi-organ failure, hemodynamic instability, and increased susceptibility to infections [74]. These abnormalities often reflect underlying pathophysiological insults, such as sepsis, acute kidney injury, or haematological malignancies, which all increase the risks of patient mortality. As the third significant topic, Topic 25 is identified as complications peculiar to certain specified procedures (ICD-996). These complications involve device-related

Figure 8: Relationship between the most and least mortality-associated topics and ICD codes identified by FedWeight-ETM. We identified the top 5 mortality-associated topics based on Wilcoxon test on topic proportions between deceased and surviving patients. The colour intensity of the heatmap indicates the probability of an ICD codes within a given topic. We selected the top ICD codes whose probability is greater than 0.08. These ICD codes were validated by Fisher's exact test p-values on the eICU and MIMIC-III data, visualized in a scatter plot for the most and least mortality-associated ICD codes. **a.** Mortality-associated topics and ICD codes on eICU. **b.** Mortality-associated topics and ICD codes on MIMIC-III.

infections, graft failures, prosthetic malfunctions, and post-surgical hemorrhage, which can pre-

cipitate systemic instability, multi-organ failure, or sepsis. This indicates that FedWeight-ETM is able to efficaciously capture semantically meaningful topics, thus helping to uncovering clinically relevant patterns in healthcare data. The association of these medical conditions with mortality underscores the necessity of rigorous monitoring, prompt identification, and timely intervention to mitigate these risks and improve patient survival.

3 Discussion

FL is a promising approach for leveraging decentralized EHR data. However, FL notoriously suffers from the covariate shift issue, where data distributions differ significantly across clinical sites. These differences in demographics, clinical practices, and data collection processes may lead to significant performance degradation of the shared model when deployed for a target population. To mitigate this issue, we propose FedWeight, where we probabilistically re-weight the patients from the source hospitals. Intuitively, samples more similar to the target distribution receive higher weights, thus contributing more during training, whereas those less similar are assigned lower weights, thereby contributing less during training. This approach ensures that the data more relevant to the target distribution has a more significant impact on model training, thus aligning the trained model with the data distribution of the target hospital and effectively addressing covariate shifts in FL environments.

We conducted extensive experiments by FL across hospitals within the eICU dataset and between the eICU and MIMIC-III datasets. Our approach demonstrates the following strengths: (1) enhances the generalization of FL classifiers for predicting clinical outcomes such as mortality, sepsis, ventilator usage, and ICU length of stay; (2) uncovers subtle yet significant drugs and lab tests associated with clinical outcomes; and (3) identifies relationships between diseases, involving renal and heart failure, and future mortality at ICU readmission. In future work, we aim to theoretically investigate why certain density estimators (e.g. VAE) perform better in specific prediction tasks. We also plan to develop density estimators and federated models that perform well even on small datasets. Furthermore, we now aggregated patient clinical data across all time points to estimate density. We will also explore density estimators designed for time series data. Moreover, although our method enhances performance on the target hospital, its ability to generalize to entirely new or highly heterogeneous sites remains uncertain. A potential future direction could involve systematically analyzing the degree of data heterogeneity under which FedWeight outperforms other methods. We will establish quantitative benchmarks to rigorously assess its effectiveness across different levels of distribution shifts. Besides, since some patients experience multiple ventilation episodes, medications administered during treatment may be mistakenly identified as predictors of future treatments. We aim to mitigate this spurious correlation in our future work. Finally, we will integrate our method into existing open-source Software Development Kit (SDK) such as Flower [75], Federated AI Technology Enabler (FATE) [76], FedScale [77], as well as NVIDIA Federated Learning Application Runtime Environment (NVIDIA FLARE) [78], for practical application in real-world health institutions.

4 Methods

4.1 FedWeight

Problem formulation   Assuming there are K source hospitals and one target hospital τ in the federated network. For each source hospital $k \in K$, we have input data $\mathbf{X}_k \in \mathbb{R}^{N_k \times D}$ and labels $\mathbf{y}_k \in \mathbb{R}^{N_k}$, where N_k is the number of patients in hospital k and D is the feature size, and $N = \sum_{k=1}^{K} N_k$ is the total number of patients in the federated network. The notations and their corresponding descriptions are outlined in Table S1.

Federated Learning   Before delving into FedWeight, we first described the baseline Federated Average (FedAvg) [3], whose training process is as follows:

- **Local computation:** Each source hospital $k \in K$ updates its model parameters $\mathbf{w}_k$ on its local data.

- **Parameter sharing:** Then, the source hospital k sends its updated parameters $\mathbf{w}_k$ to the target hospital τ.

- **Aggregation at the target hospital:** The target hospital τ receives these updated parameters and aggregates them to update the global model, which is then sent back to the source hospitals for the next round training.

$$\mathbf{w} = \sum_{k=1}^{K} \frac{N_k}{N} \mathbf{w}_k$$

Weighted log-likelihood   However, FedAvg does not account for the covariate shift issue. Specifically, covariate shift can happen when the source and target hospitals may have different local resources / clinical practices: $p_k(\mathbf{X}) \neq p_\tau(\mathbf{X})$ while clinical outcomes given certain drugs remain similar across hospitals: $p_k(\mathbf{y}|\mathbf{X}) = p_\tau(\mathbf{y}|\mathbf{X})$. To mitigate this issue, we may employ the weighted log-likelihood algorithm [13], which modifies the standard log-likelihood calculation by assigning weights to different data points. Samples more similar to the target distribution are given higher weights, thus contributing more during training, while those less similar are assigned lower weights, thereby contributing less during training. Consequently, this approach ensures that the trained model gives preference to data which are more relevant to the target distribution, thus addressing covariate shift in FL environments.

The weighted log-likelihood for hospital k can be expressed mathematically as follows:

$$\mathcal{L}(\mathbf{w}_k) = \frac{1}{N_k} \sum_{n=1}^{N_k} \varphi(\mathbf{x}_n) \log p(y_n|\mathbf{x}_n, \mathbf{w}_k)$$

where $\varphi(\mathbf{x}_n) = \left(\frac{p_\tau(\mathbf{x}_n)}{p_k(\mathbf{x}_n)} \right)^\lambda$ denotes the weight assigned to the n-th data point, p_τ represents the density estimator trained on target hospital τ, and p_k denotes the density estimator trained on

source hospital k. Furthermore, λ is a hyper-parameter that controls the degree of re-weighting. Moreover, we employed various density estimators such as MADE, VAE, and VQ-VAE on account of their suitability in estimating the underlying probability distributions.

We developed FedWeight by incorporating the weighted log-likelihood algorithm into FL settings. Therefore, the log-likelihood for the entire federated network can be expressed as follows:

$$
\begin{aligned}
\mathcal{L}(\mathbf{w}) &= \sum_{k=1}^{K} \frac{N_k}{N} \mathcal{L}(\mathbf{w}_k) \\
&= \sum_{k=1}^{K} \frac{N_k}{N} \frac{1}{N_k} \sum_{n=1}^{N_k} \varphi(\mathbf{x}_n) \log p(y_n | \mathbf{x}_n, \mathbf{w}_k) \\
&= \frac{1}{N} \sum_{k=1}^{K} \sum_{n=1}^{N_k} \varphi(\mathbf{x}_n) \log p(y_n | \mathbf{x}_n, \mathbf{w}_k)
\end{aligned}
$$

Density estimation To calculate the re-weight $\varphi(\mathbf{x}_n)$, we need to effectively train the density estimator p_k and p_τ, which is described below.

(1) Masked Autoencoder for Density Estimation (MADE) The Masked Autoencoder for Density Estimation (MADE) is a neural network model for efficient density estimation in high-dimensional data [34]. It modifies the traditional autoencoder architecture by applying masks to its connections, ensuring that the output at each unit satisfies the autoregressive property, thus effectively modelling the input data's joint distribution.

The density estimation using the autoregressive property is calculated as follows:

$$
p(\mathbf{x}) = \prod_{d=1}^{D} p(x_d | \mathbf{x}_{<d})
$$

To achieve the autoregressive property, the conventional autoencoder architecture is modified by masking some of the model weights. Therefore, the MADE training algorithm consists of the following three steps:

- **Number Assignment:** Initially, a unique integer ranging from 1 to D is sequentially assigned to each input and output unit of the autoencoder model. Moreover, every hidden unit is randomly allocated an integer within the range of 1 to $D-1$, inclusive.

- **Mask Construction:** Assuming the autoencoder model consists of L layers, let $\mathbf{M}^{\mathbf{W}^l}$ represent the mask matrix between layer l and its preceding layer $l-1$, except for the output layer. We connect the unit k' in layer l and the unit k in its preceding layer $l-1$ only if the assigned integer $m^l(k')$ is greater than or equal to $m^{l-1}(k)$; in all other cases, we

apply masks. Therefore, this mask matrix is calculated as follows:

$$M_{k',k}^{\mathbf{W}^l} = \mathbb{I}[m^l(k') \geq m^{l-1}(k)]$$

Let $\mathbf{M}^{\mathbf{V}}$ denote the mask matrix between the output layer L and the last hidden layer $L-1$. Specifically, we connect the output unit d' and the unit d in the last hidden layer only if the assigned integer $m^L(d')$ is **strictly** greater than $m^{L-1}(d)$; in all other cases, we apply masks. Therefore, this mask matrix is calculated as follows:

$$M_{d',d}^{\mathbf{V}} = \mathbb{I}[m^L(d') > m^{L-1}(d))]$$

- **Output Calculation:** Then the autoencoder output $\hat{\mathbf{x}}$ is computed as follows:

$$h^l(\mathbf{x}) = \alpha(\mathbf{b}^l + (\mathbf{W}^l \odot \mathbf{M}^{\mathbf{W}^l})h^{l-1}(\mathbf{x})),$$
$$\hat{\mathbf{x}} = \sigma(\mathbf{b}^L + (\mathbf{W}^L \odot \mathbf{M}^{\mathbf{V}})h^{L-1}(\mathbf{x}))$$

where:

  - The masks $\mathbf{M}^{\mathbf{W}^l}$ and $\mathbf{M}^{\mathbf{V}}$ are applied through element-wise multiplication with their respective model weights $\mathbf{W}^l$ and $\mathbf{W}^L$.
  - $\mathbf{b}^l$ and $\mathbf{b}^L$ are the bias for the model hidden and output layers.
  - α is the activation function in the hidden layer, while σ denotes the sigmoid function.

Given the output $\hat{\mathbf{x}}$, we find the model parameters to maximize the reconstruction likelihood.

(2) Variational Autoencoder (VAE) Variational Autoencoder (VAE) is a generative machine learning model that uses deep neural networks to encode data into a latent space and then decode it back, enabling tasks like data generation and density estimation [35]. Specifically, a VAE model consists of two coupled components: an encoder model $q_\varphi(\mathbf{z}|\mathbf{x})$, and a decoder model $p_\theta(\mathbf{x}|\mathbf{z})$. During training, the encoder maps the observations $\mathbf{x}$ to an approximate posterior over the latent variables, parameterized by a mean $\boldsymbol{\mu}$ and variance $\boldsymbol{\sigma}^2$. Instead of directly sampling the latent representation $\mathbf{z}$ from the encoded latent distribution $q_\varphi(\mathbf{z}|\mathbf{x})$, which is non-differentiable, $\mathbf{z}$ can be expressed as $\mathbf{z} = \boldsymbol{\mu} + \boldsymbol{\sigma} \odot \boldsymbol{\epsilon}$, where $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$. Finally, the decoder ingests the latent representations $\mathbf{z}$ to reconstruct the initial data $\hat{\mathbf{x}}$.

Furthermore, the decoder parameters θ and the variational parameters φ are learned by maximizing the Evidence Lower Bound (ELBO), where $p(\mathbf{z})$ denotes a standard normal distribution.

$$ELBO = \underbrace{\mathbb{E}_q\left[\log p_\theta(\mathbf{x}|\mathbf{z})\right]}_{\text{Reconstruction log-likelihood}} - KL\left[q_\varphi(\mathbf{z}|\mathbf{x})||p(\mathbf{z})\right]$$

Based on $\log p(\mathbf{x}) = KL[q_\varphi(\mathbf{z}|\mathbf{x})||p(\mathbf{z}|\mathbf{x})] + ELBO$, when the model is converged, we expect the variational distribution $q_\varphi(\mathbf{z}|\mathbf{x})$ approaches the true posterior distribution $p(\mathbf{z}|\mathbf{x})$, leading to their KL divergence converges to zero. Consequently, we may leverage ELBO in density estimation, as $\log p(\mathbf{x})$ approaches to ELBO when the model is converged.

(3) Vector Quantized Variational Autoencoder (VQ-VAE) In addition to the aforementioned density estimators, we may also utilize the Vector Quantized Variational Autoencoder (VQ-VAE) to estimate density [36]. Compared with traditional VAE that utilizes continuous latent representations, VQ-VAE employs a discretized latent space. Moreover, VQ-VAE uses categorical distributions for both posteriors $q(\mathbf{z}|\mathbf{x})$ and priors $p(\mathbf{z})$.

Similar to the VAE architecture, VQ-VAE also contains an encoder and decoder. Further, VQ-VAE maintains a codebook $\mathbf{E} = \{\mathbf{e}_1, \mathbf{e}_2, ..., \mathbf{e}_C\}$, where C is the total codewords. For each codeword $\mathbf{e}_c \in \mathbb{R}^H$, H denotes the codeword dimension.

When training VQ-VAE, the encoder takes $\mathbf{x}$ as input and generates $\mathbf{z}_e$. Then $\mathbf{z}_e$ goes into the codebook $\mathbf{E}$ to find the index of the **nearest** codeword $\hat{c} = \mathrm{argmin}_{c \in C}||\mathbf{z}_e - \mathbf{e}_c||^2$. Then we use this index to construct the one-hot encoded latent representation $\mathbf{z} = [z_c]_{\times C}$, where $z_c = \mathbb{I}[c = \hat{c}]$. Subsequently, the decoder takes $\mathbf{e}_{\hat{c}}$ as input and reconstructs the original input data $\hat{\mathbf{x}}$. Finally, we may leverage ELBO in density estimation.

FedWeight algorithm design The FedWeight training process is as follows:

- **Density estimator training:** At the beginning of the training process, the target hospital τ trains a density estimator p_τ. Meanwhile, each source hospital $k \in K$ also trains its own density estimator p_k. We selected specific density estimator models, including MADE, VAE, and VQ-VAE on account of their suitability in estimating the underlying probability distributions (Figure 3a).

- **Sharing of target density estimator:** Then the target hospital distributes its density estimator p_τ to all the source hospitals (Figure 3b).

- **Re-weighted local model training:** Upon receipt p_τ, each source hospital $k \in K$ computes the re-weight $\varphi(\mathbf{x}_n) = \left(\frac{p_\tau(\mathbf{x}_n)}{p_k(\mathbf{x}_n)}\right)^\lambda$ for patient $n \in N_k$, where λ is a hyper-parameter that controls the degree of re-weighting. Leveraging such re-weight, the source hospital k trains its own model locally. Then, the parameters of such model $\mathbf{w}_k$ is further sent to the target hospital τ (Figure 3c).

- **Aggregation at the target hospital:** The target hospital τ receives these updated parameters and aggregates them to update the global model, which has better generalization capabilities when applied to the target hospital's data (Figure 3d).

$$\mathbf{w} = \sum_{k=1}^{K} \frac{N_k}{N} \mathbf{w}_k$$

FedWeight effectively adapts the model trained on source hospital data to a target hospital. However, there may be scenarios in which hospitals must mutually benefit from each other's data distributions. To address the collaboration among these hospitals, we developed a **Symmetric FedWeight** training paradigm. For simplicity, we assumed a federated network with two hospitals, namely A and B. During local training at Hospital A, Hospital B is treated as the target to estimate the reweighting ratios. Conversely, when training at Hospital B, Hospital A serves as the target for reweighting. This symmetric approach allows both hospitals to benefit from each other by adapting to their respective data distributions (Figure 3e-h).

Moreover, to protect patients' privacy, only the parameters of the models are exchanged within the federated networks, ensuring no transmission of raw data.

4.2 Federated embedded topic model

ETM   ETM is a generative model of documents that learns interpretable topics and word embeddings and is robust to large vocabularies. For each patient diagnosis data point $\mathbf{x}_n$, the encoder, parameterized by $\mathbf{w}_\theta$, maps it to $\boldsymbol{\mu}_n$ and $\log \boldsymbol{\sigma}_n^2$ through two distinct linear layers, parameterized by $\mathbf{w}_\mu$ and $\mathbf{w}_\sigma$.

$$\boldsymbol{\mu}_n, \log \boldsymbol{\sigma}_n^2 = f(\mathbf{x}_n; \mathbf{w}_\theta, \mathbf{w}_\mu, \mathbf{w}_\sigma)$$

Through the re-parameterization trick, we obtain the latent represatation $\mathbf{z}_n$, which eventually outputs the patient-topic mixture $\boldsymbol{\theta}_n$ via a softmax operation across all latent topics.

$$\mathbf{z}_n = \boldsymbol{\mu}_n + \boldsymbol{\sigma}_n \circ \boldsymbol{\epsilon}_n, \ \text{ where } \ \boldsymbol{\epsilon}_n \sim \mathcal{N}(0, \boldsymbol{I})$$
$$\boldsymbol{\theta}_n = \text{softmax}(\mathbf{z}_n)$$

The decoding process involves the multiplication of two learnable matrices: the topic embedding matrix $\boldsymbol{\alpha}$ and the ICD embedding matrix $\boldsymbol{\rho}$. The operation yields a topic-ICD mixture $\boldsymbol{\beta}$, which represents the probabilistic association between topics and ICD codes. Eventually, $\boldsymbol{\beta}$ is multiplied by $\boldsymbol{\theta}_n$, resulting in the reconstruction of the input $\hat{\mathbf{x}}_n$.

$$\boldsymbol{\beta} = \boldsymbol{\alpha}\boldsymbol{\rho}$$
$$\hat{\mathbf{x}}_n = \boldsymbol{\theta}_n\boldsymbol{\beta}$$

ETM is trained by maximizing the following ELBO function:

$$ELBO_n = \mathbb{E}_q[\log p(\mathbf{x}_n|\mathbf{z}_n)] - KL[q(\mathbf{z}_n|\mathbf{x}_n) \ \| \ p(\mathbf{z}_n)]$$

FedAvg-ETM   The topics are not identifiable between hospitals, preventing directly model averaging. Instead, we aggregate only $\mathbf{w}_\theta$ and $\boldsymbol{\rho}$, while $\mathbf{w}_\mu$, $\mathbf{w}_\sigma$, and $\boldsymbol{\alpha}$ are kept locally updated (Figure 7a). Specifically, the FedAvg-ETM training process is as follows:

1. **Local computation:** Each source hospital $k \in K$ updates its local model parameters $\mathbf{w}^k = \{\mathbf{w}_\theta^k, \mathbf{w}_\mu^k, \mathbf{w}_\sigma^k, \boldsymbol{\alpha}^k, \boldsymbol{\rho}^k\}$.

2. **Parameter sharing:** Then, the source hospital k sends the **non-topic-associated** model parameters $\mathbf{w}_\theta^k$ and $\boldsymbol{\rho}^k$ to the target hospital τ.

3. **Aggregation at the target hospital:** The target hospital τ receives these updated parameters and aggregates them to update the global model, which is then sent back to the source hospitals for the next round of training.

$$\mathbf{w}_\theta = \sum_{k=1}^{K} \frac{N^k}{N} \mathbf{w}_\theta^k \quad \boldsymbol{\rho} = \sum_{k=1}^{K} \frac{N^k}{N} \boldsymbol{\rho}^k$$

FedWeight-ETM Variations in clinical practices, patient demographics, and data collection methods, may engender covariate shifts in clinical data, which hinders FedAvg-ETM from effectively uncovering semantically meaningful latent topics. To address this challenge, we proposed FedWeight-ETM, which integrates the FedWeight framework with the ETM model. Specifically, we assign the reweighting ratio $\varphi(\mathbf{x}_n) = \left(\frac{p_\tau(\mathbf{x}_n)}{p_k(\mathbf{x}_n)}\right)^\lambda$ to the likelihood term of the ELBO function of the n-th data point (Figure 7b), leading to the weighted ELBO function:

$$ELBO_n = \varphi(\mathbf{x}_n)\mathbb{E}_q[\log p(\mathbf{x}_n|\mathbf{z}_n)] - KL[q(\mathbf{z}_n|\mathbf{x}_n) \parallel p(\mathbf{z}_n)]$$

4.3 Data preprocessing

We applied our method to the eICU Collaborative Research Database with data from 208 hospitals and 200,859 patients, and the MIMIC-III dataset. For the eICU dataset, we selected 10 hospitals with the most patients: Hospitals 167, 420, 199, 458, 252, 165, 148, 281, 449, and 283. The dataset includes 1,399 distinct drugs, which were binarized for model analyses; a value of "1" indicates the patient received the drug, whereas "0" signifies no administration. Since patients typically receive only 1% to 2% of the total drug catalogue, this binarization results in a sparse matrix with a predominance of "0" values in the dataset. In addition to drug data, our model incorporates patient demographics (age, sex, BMI, and ethnicity) to enhance predictive accuracy. Age was encoded into eight categories: <30, 30-39, 40-49, 50-59, 60-69, 70-79, 80-89, and >89 years. Sex was dichotomized, with "1" representing males and "0" indicating females. BMI was categorized into four groups: underweight (<18.5), normal (18.5-24.9), overweight (25-29.9), and obesity (≥ 30) [79]. Ethnicity was classified into five categories: Caucasian, African American, Hispanic, Asian, and Native American, reflecting the diversity within the patient population. For MIMIC-III dataset, we removed the newborn patients. Moreover, given the small sample size, we excluded the MIMIC patients in Cluster 2 for simplicity (Figure S1a).

Drug imputation In the eICU dataset, approximately one-third of drug names are not recorded across all the 208 hospitals. If these unrecorded drugs are represented as "0", we tend to have an extremely sparse input, which hampers the training of density estimators due to the lack of informative features. For instance, if density estimators like MADE, VAE, or VQ-VAE are trained under such conditions, these auto-encoders tend to reconstruct more zeros than ones. To address this challenge, we implemented a strategy for imputing the missing drug names. While some drugs lack recorded names, we observed that they possess Hierarchical Ingredient Code List (HICL) codes —— a standardized coding system for identifying healthcare products. Consequently, we could match these unrecorded drugs with their counterparts with both HICL codes and recorded names in the database. This imputation method allows us to fill in missing drug names based on their HICL codes. As a result, the proportion of unrecorded drugs in our dataset decreases to around 20%, which enhances the accuracy and reliability of our models by providing a richer set of features (Figure 2a-b).

Drug harmonization We discovered that different hospitals may use different drug names, although they share the same identity (e.g. "aspirin" and "acetylsalicylic acid"). We also found that the dosage information is included in some drug names (e.g. "aspirin 10 mg"). Consequently, we have minimal overlap in drug names between hospitals, further exacerbating the challenge of sparse input data and hindering the training process. To address this, we implemented a harmonization process to standardize drug names. We merged those with shared identities while disregarding dosage details. Our algorithm comprises two main steps:

- **Initial drug mapping:** We maintained a reference panel of drug names on the server, containing the 237 most common drugs [80], which is periodically updated and shared with client hospitals. For each drug name from a client hospital (e.g., "aspirin 10mg"), the algorithm checks against each name in the reference panel to find if the reference name (e.g., "aspirin") is included in the client's drug name ("aspirin 10mg"). If a match is detected, the client's drug name is mapped to the reference drug name (e.g., "aspirin"). This approach enables accurate mapping of most drug names without dosage information.

- **Use of BioWordVec for mapping:** If no direct match is found, the algorithm initiates a similarity analysis. First, the client's drug name and the reference drug name are converted into their respective word embeddings using BioWordVec [81]. Then, we compute the cosine similarity between these two word embeddings. After comparing with all reference drug names, the reference name with the highest cosine similarity is selected. This approach effectively matches drugs with similar identities.

After harmonization, the proportion of overlapping drugs across hospitals significantly increases (Figure 2c-d).

Lab tests data preprocessing We also included lab tests as model input. However, we encountered several challenges due to discrepancies between the eICU and MIMIC-III datasets.

The eICU dataset contains 158 unique lab tests, whilst MIMIC-III includes 590 unique lab tests, with differences in test names, abbreviations, and units across the two datasets. Referring to a previous study that focused on the 29 most influential lab tests for model training [80], we identified 12 lab tests common to both datasets (Table S2). Due to variations in the scale of lab results, we normalized the data to ensure consistency and comparability.

Time series data preprocessing   We used the eICU dataset to construct predictive models for patients' mortality, ventilator use, and sepsis diagnosis, and ICU length of stay. To prepare data for mortality and length of stay prediction, we segmented the dataset into two intervals: drug data within the initial 48 hours of ICU admission and mortality outcomes and length of stay after 48 hours. The models were trained using patient demographics and drug administration data from the first 48 hours to predict mortality and length of stay beyond this period. For ventilator use and sepsis diagnosis, we designed a 72-hour observation period after ICU entry, further divided into six 12-hour intervals. The objective was to use drug and demographic information from each interval to predict ventilator use and sepsis occurrence in the next interval.

4.4   Simulation Study

Generating inputs   We created the simulation dataset input from the preprocessed eICU dataset, which included imputed and harmonized drug data alongside age, sex, BMI, and ethnicity information. For simplicity, we assumed one source hospital and one target hospital in the network. To generate the simulation dataset, we utilized the input from Hospital 167, which has the most patients, as the target input $\mathbf{X}_\tau$. We further combined the remaining 9 hospitals as the source input $\mathbf{X}_k$.

Generating labels   We simulated labels by first creating a linear model $f(\mathbf{X}) = \mathbf{Xw} + \mathbf{b}$, where the weight was sampled from Gaussian distribution: $\mathbf{w} \sim \mathcal{N}(0, \mathbf{I})$. Meanwhile, we created a binary mask sampled from Bernoulli distribution: $\mathbf{m} \sim Bernoulli(0.15)$. Then we applied the binary mask $\mathbf{m}$ to the weight: $\mathbf{w} = \mathbf{w} \odot \mathbf{m}$. As a result, 15% of the weight values were sampled from Gaussian distribution while setting the rest of weight values as 0, which simulated around 15% of features are causal.

In terms of the bias of the linear model $\mathbf{b} \in \mathbb{R}^N$, we set it as: $\mathbf{b} = -\log \frac{1-\boldsymbol{\pi}}{\boldsymbol{\pi}}$ where $\boldsymbol{\pi} = [0.15]_{\times N}$. Then we created imbalanced dataset with most labels were 0, which simulated the real-world scenario.

Finally, we generated simulated labels for source hospital: $\mathbf{y}_k = \mathbb{I}[f(\mathbf{X}_k) > 0]$, and target hospital $\mathbf{y}_\tau = \mathbb{I}[f(\mathbf{X}_\tau) > 0]$. We used the same linear model f to generate source and target labels, ensuring consistent conditional probabilities and simulating the covariate-shift problem.

4.5 Experimental design for reliable model evaluation

To enable early stopping and hyperparameter selection, we created a validation set, represented by $(\mathbf{X}_\tau^{(val)}, \mathbf{y}_\tau^{(val)})$, assuming part of the labels are available on the target hospital. The remaining data from the target hospital, without accessible labels, formed the test set, $(\mathbf{X}_\tau^{(test)}, \mathbf{y}_\tau^{(test)})$.

For a stable assessment of the model performance, our experimental design employed multiple seeds. We selected the 5 hospitals (Hospital 167, 199, 252, 420, 458) from the 10 hospitals with the most patients as the target hospitals. We looped these 5 hospitals, and each hospital under the loop was alternately designated as the target, while the remaining 9 were the source hospitals. Models were trained on the source hospitals and evaluated on the target. We employed bootstrap sampling to generate 100 distinct test sets based on the existing test set $(\mathbf{X}_\tau^{(test)}, \mathbf{y}_\tau^{(test)})$. We then evaluated our model on these 100 test sets, obtaining 100 individual results. By calculating the mean and standard deviation of these results, we ensured a more reliable assessment of the model's performance.

5 Data availability

The eICU dataset analyzed in this paper are publicly available through eICU Collaborative Research Database (`https://eicu-crd.mit.edu/about/eicu/`). The MIMIC-III dataset are publicly available through PhysioNet (`http://mimic.physionet.org`).

6 Code availability

We implemented FedWeight in Python3.9. The software is available at: `https://github.com/li-lab-mcgill/FedWeight`.

7 Acknowledgments

Y.L. is supported by NOVA-FRQNT-NSERC grant (FRQ-NT 2023-NOVA-328677), Canada Research Chair (Tier 2) in Machine Learning for Genomics and Healthcare (CRC-2021-00547) and Natural Sciences and Engineering Research Council (NSERC) Discovery Grant (RGPIN-2016-05174). H.Z. is supported by Artificial Intelligence for Public Health and Health Equity Trainee Scholarship.

8 Author contributions

Y.L., D.L., and H.Z. coined the idea. Y.L., D.L., and D.B. supervised the work. H.Z. implemented the software and conducted the experiments. Y.L., D.L., D.B., and H.Z. analyzed the data with the help from N.L.. All authors contributed to the final writing of the manuscript.

Declaration of Interests

The authors declare no competing interests.

References

1. Muralidharan, V. *et al.* A scoping review of reporting gaps in fda-approved ai medical devices. *npj Digital Medicine* **7**, 273 (2024).

2. Miotto, R., Wang, F., Wang, S., Jiang, X. & Dudley, J. T. Deep learning for healthcare: review, opportunities and challenges. *Briefings in Bioinformatics* (2017).

3. McMahan, H. B., Moore, E., Ramage, D., Hampson, S. & Aguera y Arcas, B. Communication-efficient learning of deep networks from decentralized data. *Proceedings of the 20th International Conference on Artificial Intelligence and Statistics (AISTATS)* (2017).

4. Sheller, M. J. *et al.* Federated learning in medicine: facilitating multi-institutional collaborations without sharing patient data. *Scientific Reports* **10** (2020).

5. Huang, L. *et al.* Patient clustering improves efficiency of federated machine learning to predict mortality and hospital stay time using distributed electronic medical records. *Journal of Biomedical Informatics* **99** (2019).

6. Dayan, I. *et al.* Federated learning for predicting clinical outcomes in patients with COVID-19. *Nature Medicine* **27** (2021).

7. Wahba, G. *SPLine models for observational data* (1990). URL `https://doi.org/10.1137/1.9781611970128`.

8. Bishop, C. M. Neural networks for pattern recognition. *Choice Reviews Online* (1994).

9. Vapnik, V. *The nature of statistical learning theory* (1995). URL `https://doi.org/10.1007/978-1-4757-2440-0`.

10. Hart, P. D., Duda, R. O. & Stork, D. G. *Pattern classification* (2012).

11. Hastie, T., Tibshirani, R. & Friedman, J. H. *The elements of statistical learning: data mining, inference, and prediction* (2013).

12. Schölkopf, B. & Smola, A. J. Learning with Kernels. *The MIT Press eBooks* (2018). URL `https://doi.org/10.7551/mitpress/4175.001.0001`.

13. Shimodaira, H. Improving predictive inference under covariate shift by weighting the log-likelihood function. *Journal of Statistical Planning and Inference* **90** (2000).

14. Pan, S. J. & Yang, Q. A survey on transfer learning. *IEEE Transactions on Knowledge and Data Engineering* (2010).

15. Amodei, D. *et al.* Concrete problems in AI safety. *arXiv:1606.06565* (2016).

16. Ross, P. & Spates, K. Considering the safety and quality of artificial intelligence in health care. *The Joint Commission Journal on Quality and Patient Safety* (2020).

17. Macrae, C. Governing the safety of artificial intelligence in healthcare. *BMJ Quality and Safety* (2019).

18. Habli, I., Lawton, T. & Porter, Z. Artificial intelligence in health care: accountability and safety. *Bulletin of The World Health Organization* (2020).

19. Schrouff, J. *et al.* Diagnosing failures of fairness transfer across distribution shift in real-world medical settings. *36th Conference on Neural Information Processing Systems* (2022).

20. Singh, H., Mhasawade, V. & Chunara, R. Generalizability challenges of mortality risk prediction models: A retrospective analysis on a multi-center database. *PLOS digital health* (2022).

21. Schumann, C. *et al.* Transfer of Machine Learning Fairness across Domains. *arXiv:1906.09688* (2019). URL `http://export.arxiv.org/pdf/1906.09688`.

22. Huang, J., Smola, A. J., Gretton, A., Borgwardt, K. M. & Schölkopf, B. *Correcting sample selection bias by unlabeled data* (2007).

23. Fang, T., Lu, N., Niu, G. & Sugiyama, M. Rethinking Importance Weighting for Deep Learning under Distribution Shift. *Advances in Neural Information Processing Systems 33 (NeurIPS 2020)* (2020).

24. Sugiyama, M., Nakajima, S., Kashima, H., Buenau, P. V. & Kawanabe, M. Direct Importance Estimation with Model Selection and Its Application to Covariate Shift Adaptation. *Advances in Neural Information Processing Systems 20 (NIPS 2007)* (2007).

25. Wang, M., Guo, J. & Jia, W. Federated Multi-Phase Curriculum learning to synchronously correlate user heterogeneity. *IEEE Transactions on Artificial Intelligence* (2023).

26. Duan, J.-H., Li, W. & Lu, S. *FedDNA: Federated Learning with Decoupled Normalization-Layer Aggregation for Non-IID Data* (2021).

27. Ramezani-Kebrya, A., Liu, F., Pethick, T., Chrysos, G. & Cevher, V. Federated learning under covariate shifts with generalization guarantees. *Transactions on Machine Learning Research* (2023). URL `https://openreview.net/forum?id=N7lCDaeNiS`.

28. Xu, Y., Cui, W., Xu, J. & Cheng, H. Federated covariate shift adaptation for missing target output values. *International Journal of Wavelets, Multiresolution and Information Processing* (2023).

29. Nguyen, H., Wu, P. & Chang, J. M. Federated Learning for distribution skewed data using sample weights. *IEEE Transactions on Artificial Intelligence* (2023).

30. Chaddad, A. *et al.* Explainable, Domain-Adaptive, and federated artificial intelligence in medicine. *IEEE/CAA Journal of Automatica Sinica* (2023).

31. Zhang, M., Wang, Y. & Luo, T. Federated Learning for Arrhythmia Detection of Non-IID ECG. *International Conference on Computer and Communications (ICCC)* (2020).

32. Pollard, T. J. *et al.* The eICU collaborative research database, a freely available multi-center database for critical care research. *Scientific Data* **5** (2018).

33. Johnson, A. E. *et al.* MIMIC-III, a freely accessible critical care database. *Scientific Data* (2016).

34. Germain, M., Gregor, K., Murray, I. & Larochelle, H. MADE: Masked autoencoder for distribution estimation. *Proceedings of the 32nd International Conference on Machine Learning* (2015).

35. Kingma, D. P. & Welling, M. Auto-encoding variational bayes. *International Conference on Learning Representations* (2014).

36. van den Oord, A., Vinyals, O. & Kavukcuoglu, K. Neural discrete representation learning. *31st Conference on Neural Information Processing Systems* **30** (2017).

37. Lundberg, S. M. & Lee, S.-I. A unified approach to interpreting model predictions. *31st Conference on Neural Information Processing Systems* **30** (2017).

38. Gallanosa, A., Stevens, J. & Quick, J. Glycopyrrolate (2023). URL `https://www.ncbi.nlm.nih.gov/books/NBK526035/`.

39. Holmes, C. L. & Walley, K. R. Vasopressin in the ICU. *Current Opinion in Critical Care* (2004).

40. Sharman, A. & Low, J. A. Vasopressin and its role in critical care. *Continuing Education in Anaesthesia, Critical Care and Pain* (2008).

41. Belletti, A. *et al.* Effect of continuous epinephrine infusion on survival in Critically Ill patients: A Meta-Analysis of Randomized Trials*. *Critical Care Medicine* (2020).

42. Patel, V. V., Sullivan, J. B. & Cavanaugh, J. Analysis of mortality in patients treated with phenylephrine in septic shock. *Journal of Pharmacy Practice* (2021).

43. Ostermann, M. E., Keenan, S. P., Seiferling, R. A. & Sibbald, W. J. Sedation in the intensive care unit: A systematic review. *JAMA* (2000).

44. Wernly, B. *et al.* Blood urea nitrogen (BUN) independently predicts mortality in critically ill patients admitted to ICU: A multicenter study. *Clinical Hemorheology and Microcirculation* (2018).

45. Arihan, O. *et al.* Blood Urea Nitrogen (BUN) is independently associated with mortality in critically ill patients admitted to ICU. *PLoS ONE* (2018).

46. Saygitov, R. T., Glezer, M. G. & Semakina, S. V. Blood urea nitrogen and creatinine levels at admission for mortality risk assessment in patients with acute coronary syndromes. *Emergency Medicine Journal* (2010).

47. Cauthen, C. A. *et al.* Relation of blood urea nitrogen to Long-Term mortality in Patients with Heart Failure. *The American Journal of Cardiology* (2008).

48. Yin, J. *et al.* Blood urea nitrogen and clinical prognosis in patients with COVID-19: A retrospective study. *Medicine* (2024).

49. Bernhard, M. *et al.* Elevated admission lactate levels in the emergency department are associated with increased 30-day mortality in non-trauma critically ill patients. *Scandinavian Journal of Trauma Resuscitation and Emergency Medicine* (2020).

50. Bou Chebl, R., El Khuri, C., Shami, A. *et al.* Serum lactate is an independent predictor of hospital mortality in critically ill patients in the emergency department: a retrospective study. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine* (2017).

51. Chen, Y.-X. & Li, C.-S. Lactate on emergency department arrival as a predictor of mortality and site-of-care in pneumonia patients: a cohort study. *Thorax* (2015).

52. Kruse, O., Grunnet, N. & Barfod, C. Blood lactate as a predictor for in-hospital mortality in patients admitted acutely to hospital: a systematic review. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine* (2011).

53. Barr, J. Propofol: a new drug for sedation in the intensive care unit. *International Anesthesiology Clinics* (1995).

54. Marinella, M. A. Propofol for sedation in the intensive care unit: essentials for the clinician. *Respiratory Medicine* (1997).

55. Klompas, M., Speck, K., Howell, M. D., Greene, L. & Berenholtz, S. M. Reappraisal of routine oral care with chlorhexidine gluconate for patients receiving mechanical ventilation. *JAMA Internal Medicine* (2014).

56. Dai, W.-Y. *et al.* Meta-Analysis of the efficacy and safety of chlorhexidine for Ventilator-Associated Pneumonia Prevention in mechanically Ventilated patients. *Evidence-based Complementary and Alternative Medicine* (2022).

57. Zuckerman, L. M. Oral Chlorhexidine Use to Prevent Ventilator-Associated Pneumonia in Adults. *Dimensions of Critical Care Nursing* (2016).

58. Park, H. Y. *et al.* Effects of etomidate use in ICU patients on ventilator therapy: a study of 12,526 patients in an open database from a single center. *Korean journal of anesthesiology* (2021).

59. Bruder, E. A., Ball, I. M., Ridi, S., Pickett, W. & Hohl, C. Etomidate for sedating critically ill people during emergency endotracheal intubation. *Cochrane Database of Systematic Reviews* (2015).

60. Febrinasari, R. P., Benedictus, B. & Azmiardi, A. Systematic Review: A Comparison between Vancomycin and Daptomycin for Sepsis Infection Antibiotic Therapy. *Open Access Macedonian Journal of Medical Sciences* (2021).

61. Vazquez, M., Fagiolino, P., Boronat, A., Buroni, M. & Maldonado, C. Therapeutic drug monitoring of vancomycin in severe sepsis and septic shock. *International Journal of Clinical Pharmacology and Therapeutics* (2008).

62. Hagel, S. *et al.* Therapeutic drug monitoring-based dose optimisation of piperacillin/tazobactam to improve outcome in patients with sepsis (TARGET): a prospective, multi-centre, randomised controlled trial. *Trials* (2019).

63. Shliapnikov, C. & Rybkin, A. The role of cefepime, a 4th-generation cephalosporin, in treating patients with surgical sepsis. *PubMed* (1999).

64. Eykyn, S. J. & Phillips, I. Metronidazole and anaerobic sepsis. *The British Medical Journal* (1976).

65. Tonai, M. *et al.* Hospital-onset sepsis and community-onset sepsis in critical care units in Japan: a retrospective cohort study based on a Japanese administrative claims database. *Critical Care* (2022).

66. Rimmer, E. *et al.* White blood cell count trajectory and mortality in septic shock: a historical cohort study. *Canadian Journal of Anesthesia/Journal canadien d anesthésie* (2022).

67. Curran, T. F. Outcomes after prolonged ICU stays in postoperative cardiac surgery patients. *Federal practitioner* (2022).

68. Malato, A. *et al.* The impact of deep vein thrombosis in critically ill patients: a meta-analysis of major clinical outcomes. *Blood Transfusion* (2015).

69. Levy, E. M. The effect of acute renal failure on mortality. A cohort analysis. *JAMA* (1996).

70. Go, A. S., Chertow, G. M., Fan, D., McCulloch, C. E. & Hsu, C.-Y. Chronic kidney disease and the risks of death, cardiovascular events, and hospitalization. *New England Journal of Medicine* (2004).

71. Artucio, H. & Pereira, M. Cardiac arrhythmias in critically ill patients: epidemiologic study. *Critical Care Medicine* (1990).

72. Gross, C., Jonasson, J. M., Buchebner, D. & Agvall, B. Prognosis and mortality within 90 days in community-acquired acute kidney injury in the Southwest of Sweden. *BMC Nephrology* (2023).

73. Bauer, M. *et al.* Mortality in sepsis and septic shock in europe, north america and australia between 2009 and 2019—results from a systematic review and meta-analysis. *Critical Care* (2020).

74. Roenhoej, R. *et al.* Abnormal routine blood tests as predictors of mortality in acutely admitted patients. *Clinical Biochemistry* (2019).

75. Beutel, D. J. *et al.* Flower: A friendly federated learning research framework. *arXiv preprint arXiv:2007.14390* (2020). URL https://arxiv.org/abs/2007.14390.

76. Liu, Y., Fan, T., Chen, T., Xu, Q. & Yang, Q. FATE: an industrial grade platform for collaborative learning with data protection. *Journal of Machine Learning Research* (2021).

77. Lai, F. *et al.* FedScale: Benchmarking model and system performance of federated learning at scale. *International Conference on Machine Learning (ICML)* (2022).

78. Roth, H. R. *et al.* Nvidia flare: Federated learning from simulation to real-world. *arXiv:2210.13291* (2022). URL https://arxiv.org/abs/2210.13291.

79. Weir, C. B. & Jan, A. BMI Classification Percentile And Cut Off Points. *StatPearls* (2019). URL https://www.ncbi.nlm.nih.gov/books/NBK541070/.

80. Pan, W., Wu, Z., Rajendran, S. & Wang, F. An Adaptive Federated Learning Framework for Clinical Risk Prediction with Electronic Health Records from Multiple Hospitals. *Patterns* (2024).

81. Zhang, Y., Chen, Q., Yang, Z., Lin, H. & Lu, Z. BioWordVec, improving biomedical word embeddings with subword information and MeSH. *Scientific Data* (2019).

82. Hochreiter, S. & Schmidhuber, J. Long Short-Term memory. *Neural Computation* (1997).

83. Higgins, I. *et al.* beta-vae: Learning basic visual concepts with a constrained variational framework. *International Conference on Learning Representations* (2017).

84. Bowman, S. R. *et al.* Generating sentences from a continuous space. *SIGNLL Conference on Computational Natural Language Learning (CONLL)* (2016).

Supplementary information

S1 Model Architecture

For the simulation dataset, our objectives are to evaluate the model's predictive performance and interpretability. Due to the ease of linear models in revealing influential features, we opted for a linear model in our simulation studies. In contrast, the real-world mortality and ICU length of stay prediction using the eICU dataset involves leveraging patient demographics and drug data from the initial 48 hours of ICU admission to predict mortality after this period. Therefore, we selected a Multi-Layer Perceptron (MLP) architecture, reflecting its capacity for handling complex, non-linear relationships in high-dimensional data. For predicting ventilator use and sepsis diagnosis, which requires analyzing current interval data to predict outcomes in the next interval, we employed a Long Short-Term Memory (LSTM) architecture [82], owing to its proficiency in capturing temporal dependencies within sequence data.

Regarding our FedWeight density estimators, we utilized a standard MADE architecture. For density estimation with VAE, we employed a beta-VAE with KL divergence annealing to mitigate posterior collapse [83, 84]. In employing VQ-VAE as a density estimator, we adopted an Exponential Moving Average (EMA) method for stable codebook updates [36].

S2 Readmission mortality prediction using logistic regression with FedWeight-ETM topics as input features

We conducted experiments on patient ICD diagnostic records benchmarking non-federated ETM trained within each silo, FedAvg-ETM, and FedWeight-ETM.

We benchmarked the prediction performance on eICU (i.e., trained eICU (+ MIMIC-III) and tested on eICU held-out patients). In this setting, the ETM model was trained solely on the eICU dataset, whilst FedAvg- and FedWeight-ETM models were trained federatively on both eICU and MIMIC-III. For FedWeight, we used eICU as the source and MIMIC-III as the target to compute the reweighting ratios. We then trained a logistic regression model using patient-topic mixtures and mortality labels to predict readmission mortality in the eICU test dataset. We observed that all methods exhibit low AUPRC when the number of topics is small, followed by a peak in AUPRC at 32 topics (Figure S3a).

Beyond this point, the AUPRC declines as the number of topics increases. Specifically, ETM performs better with fewer topics, whilst FedAvg-ETM and FedWeight-ETM outperform ETM as the number of topics increases. Among these, FedWeight-ETM slightly surpasses FedAvg-ETM, particularly when the number of topics is 32, demonstrating its advantage in capturing meaningful patterns at this optimal topic number.

In the second experiment (MIMIC on MIMIC) (Figure S3b), we used MIMIC-III as the source and eICU as the target to estimate the FedWeight reweighting ratios. We then predicted mortality on the MIMIC-III test dataset. We discovered that across all methods, the AUPRC is lowest when the number of topics is small, which increases significantly as the number of topics rises, and peaks at 16 or 32 topics, then gradually declines with further increases of topics. Amongst all the methods, FedAvg-ETM achieves the highest AUPRC, particularly at 16 topics, with FedWeight-ETM exhibiting similar performance. In contrast, ETM performs notably worse than the federated topic models. This demonstrates that federated models, by leveraging data from multiple sources, can better uncover latent topic patterns.

Figure S1: Intra-dataset covariate shifts in the MIMIC-III dataset. **a.** K-mean clustering MIMIC-III drugs. **b.** Negative log of p-values from Fisher-exact test for admission types. "NEWBORN" indicates admissions pertaining to patient's birth. "EMERGENCY" indicates unplanned medical care. "ELECTIVE" indicates unplanned admission. **c.** Negative log of p-values from Fisher-exact test for surgery procedures.

Figure S2: Performance of FedWeight on the simulation dataset. **a.** AUPRC for model predictions. **b.** AUPRC compared model weights and influential features. **c.** Pearson correlation calculated between the weights of the trained and reference models. We evaluated their statistical significance using the Wilcoxon test against the baseline, denoting * for test p-values < 0.05 and ** for p-values < 0.01.

Figure S3: Performance comparison of ETM-based models in predicting readmission mortality. We used first-admission ICD codes as input to ETM-based models to generate patient-topic mixtures. A logistic regression model was then used to predict readmission mortality based on these mixtures. **a.** Performance on eICU. **b.** Performance on MIMIC-III.

Table S1: Notations in FedWeight

Notation	Description
K	Total number of source hospitals
τ	Target hospital in the federated network
k	Source hospital in the federated network
D	Feature size across the federated network
N_k	Number of patients in source hospital k
N	Total number of patients across all source hospitals
$\mathbf{X}$	Input data for hospital
$\mathbf{y}$	Labels for hospital
p	Density estimator for hospital
f	The linear model used in generating simulation labels
$\mathbf{w}_k$	Local model parameters for source hospital k
$\mathbf{w}$	Aggregated model parameters for target hospital τ
$\mathbf{q}$	Model parameters for the ETM encoder
$\mathbf{u}$	Model parameters for the ETM linear layer which outputs $\boldsymbol{\mu}$
$\mathbf{s}$	Model parameters for the ETM linear layer which outputs $\boldsymbol{\sigma}$
$\boldsymbol{\rho}$	ETM ICD embedding
$\boldsymbol{\alpha}$	ETM topic embedding
$\boldsymbol{\theta}$	ETM patient-topic mixture
$\boldsymbol{\beta}$	ETM topic-ICD mixture
$\mathbf{Z}$	ETM latent representation
φ	The re-weighting ratio used in the weighted log-likelihood algorithm
λ	Hyper-parameter that controls the degree of re-weighting

Table S2: Lab tests, abbreviations, and units

Lab Tests	Abbreviations	Units
Oxygen Saturation	o2sat	%
Partial pressure of oxygen	pao2	mm Hg
Partial pressure of carbon dioxide	paco2	mm Hg
pH value of blood	ph	Units
Albumin	albumin	g/dL
Bands	bands	%
Blood urea nitrogen	bun	mg/dL
Hematocrit	hct	%
International Normalized Ratio (PT test)	inr	ratio
Lactate	lactate	mmol/L
Platelets	platelets	K/mcL
White blood cell	wbc	K/mcL