

Change targets, messaging, and content delivery for a community-engaged social media campaign addressing HIV-related stigma in Peru: a qualitative study

Alyson Nunez^{1*}, Marguerite Curtis^{1*}, Milagros Wong², Kristin A. Kosyluk³, Jerome T. Galea⁴, Molly F. Franke^{1§#}, Renato A. Errea^{2#}

*These authors have contributed equally to the work.

#These authors have contributed equally to the work.

1 Department of Global Health and Social Medicine, Harvard Medical School, Boston, Massachusetts, USA.

2 Socios En Salud Sucursal Perú, Lima, Peru.

3 Department of Mental Health Law and Policy, University of South Florida, Tampa, Florida, USA.

4 School of Social Work, University of South Florida, Florida, USA

§ Corresponding author: Molly Franke

Email: Molly_franke@hms.harvard.edu

641 Huntington Avenue

Boston, Massachusetts 02115

United States

ABSTRACT

Mitigating HIV-related stigma is critical to improving HIV outcomes and reaching global HIV targets. We conducted formative community-engaged research to identify key characteristics of a social media campaign to address HIV-related stigma among youth in Lima, Peru. Focus groups and in-depth interviews with young people living with HIV (men who have sex with men, transgender women, sex workers, persons with perinatally-acquired HIV, cisgender women, and Venezuelan migrants), HIV advocates, and healthcare providers were carried out from November 2022 to July 2023. Audio recordings were transcribed verbatim and analysed for emergent themes using framework analysis. Two change targets were identified for campaign messaging: 1) changing how HIV and people living with HIV (PLWH) are perceived, and 2) changing attitudes and actions towards PLWH. Messages aligning with the first change target included education to raise awareness that HIV does not discriminate; HIV is a chronic, treatable condition; and “undetectable equals untransmittable” (U=U). Messages aligning with the second target included normalizing open conversations about HIV, encouraging support versus pity for PLWH, fostering unity among HIV-affected communities, and promoting inclusion of PLWH. Participants provided recommendations to achieve these messaging objectives, including how, to whom and by whom messages are delivered.

Keywords: discrimination, youth, adolescents, men who have sex with men, transgender women

Word count

Abstract: 196

Main text: 3,892

1 INTRODUCTION

2 Stigma—defined as the social-cognitive process leading to the stereotyping and subsequent discrimination
3 of individuals or groups[1]—poses an enormous threat to global health and impedes the control and
4 treatment of infectious diseases such as human immunodeficiency virus (HIV). HIV-related stigma is
5 associated with harmful consequences for people living with HIV (PLWH), including delayed diagnosis,
6 treatment nonadherence, and nondisclosure to sexual partners[2-6]. Stigmatizing behaviours towards
7 PLWH can manifest as loss of employment or housing and rejection by friends, family members and
8 partners[7–9]. Additionally, HIV-related stigma is harmful to holders of stigmatizing beliefs, as it impedes
9 the acceptability of preventative practices and accurate assessment of one’s own HIV risk[10-12].

10 There is promising preliminary evidence on the positive effects of anti-stigma campaigns and the potential
11 for mass media communication strategies to normalize HIV[13-16]. Mass media campaigns (i.e., via
12 television, radio, and/or newspapers) have led to widespread health-related behaviour changes[16], and
13 those targeting HIV-related stigma have shown a modest impact on stigma reduction[15]. However, the
14 elevated cost and decreasing popularity of these media limits their utility for stigma reduction. The global
15 popularity of social media platforms, along with access to internet and mobile devices, allow for novel
16 approaches to stigma reduction, whereby content can be widely and inexpensively disseminated. Social
17 media is increasingly used as a platform for health communication campaigns, particularly targeting
18 sexually transmitted infections[17-21]. While some of these interventions aim to reduce stigma, such as the
19 *Get Yourself Tested* campaign in the United States[18], only one[19] included formal measures of stigma.
20 This intervention successfully reduced stigma, but did not report community engagement to inform
21 campaign content.

22 HIV-related stigma is common in Latin America[22], where there is additional pervasive stigma against
23 populations in which HIV is mainly concentrated: gay, bisexual, and other men who have sex with men
24 (GBMSM), transgender women (TGW), and cisgender female sex workers (FSWs)[23-29]. Recognizing
25 the critical need to reduce HIV-related stigma and the relatively unexplored potential of social media
26 platforms to reach both targets of stigmatizing behaviours and holders of stigmatizing beliefs, we conducted
27 formative qualitative research with PLWH, HIV advocates, and HIV healthcare providers to inform an HIV
28 stigma reduction social media campaign targeted to young people in Lima, Peru. We report findings on key
29 messages, content delivery strategies, and overarching change goals for the campaign.

30 METHODS

31 *Study setting*

32 This work was conducted in Lima, Peru, where the majority of PLWH in Peru, an estimated 110,000, reside
33 [29] and where a rising number of new HIV diagnoses in youth 18-29 has been reported[30]. HIV
34 prevalence in Peru is estimated to be 0.3-0.4%, but is notably higher in transgender women (32%), GBMSM
35 (10%), and FSWS (2.4%)[29],[31]. One recent study also found an elevated prevalence (1%) in Venezuelan
36 migrants[32].

37 38 *Participant recruitment and consent process*

39 From November 2022 to July 2023, we conducted focus groups and in-depth interviews with 110
40 participants living in Lima, Peru. Participants included a diverse group of young people living with HIV
41 (YPLWH) (N=75) ages 16-29, recruited at health facilities and through referrals by healthcare providers
42 and community partners. Participating YPLWH included GBMSM, TGW, FSWS, individuals with
43 perinatal transmission, cisgender women, and Venezuelan migrants (Table 1). Additional sessions were
44 conducted with healthcare providers (N=11), 73% of whom were women, and HIV advocates (N=24), 21%
45 of whom were assigned female sex at birth and 25% of whom were TGW.

46

47 **Table 1: Baseline characteristics of young people living with HIV (N=75)**

Characteristic	n (%)
Sex assigned at birth	
Female	26 (34.7)
Male	47 (62.7)
Intersex	2 (2.7)
Gender identity	
Woman	31 (41.3)
Man	33 (44.0)
Transgender woman	9 (12.0)
Genderfluid	1 (1.3)
Other	1 (1.3)
Sexual orientation	
Homosexual	29 (38.7)
Bisexual	10 (13.3)
Heterosexual	32 (42.7)
Other	4 (5.3)
Median age, years (IQR)	24 (21-27)
Sex worker	3 (4.0)
Venezuelan migrant	8 (10.7)
Perinatal HIV transmission	17 (22.7)

48

49 *Data collection*

50 MW, a Peruvian qualitative researcher and certified nurse with experience working with YPLWH,
51 conducted 13 focus groups and 16 in-depth interviews (the point of data saturation) using semi-structured
52 interview guides (Supplemental Table 1) tailored to each participant group. These guides, informed by the
53 Health Stigma and Discrimination Framework[33], were designed to facilitate discussion on experiences
54 with HIV-related stigma, common myths associated with HIV, messages to be portrayed, and desired
55 outcomes of the campaign. Sessions lasted between 60 and 120 minutes. In most sessions, only MW and
56 participants were present; however, a team member aided group facilitation in five. Participants were
57 compensated 50 soles (13 USD), plus any transportation costs. Focus groups with healthcare personnel and
58 all interviews were conducted virtually. Notes were not taken. No repeat interviews were conducted.
59 Sessions were audio-recorded, transcribed verbatim, and translated from Spanish to English. Transcripts
60 were not returned to participants for comment or correction.

61

62 *Data analysis*

63 Transcripts were uploaded into the qualitative research software Dedoose. Two qualitative analysts (AN,
64 MW) conducted a systematic, comparative, and inductive analysis of the transcripts guided by the
65 Framework Analysis[34]. Initial codebooks were guided by the interview guide and Health Stigma and
66 Discrimination Framework[33]; additional codes were created *de novo* as needed. Both analysts reviewed
67 each transcript to reach a consensus on codes, which were organized into emergent themes. Here, we report
68 results from excerpts related to change targets and messages for achieving these goals. The COREQ
69 checklist was used for rigor and transparency[35].

70

71 *Research ethics*

72 The Peruvian Vía Libre Bioethics Committee (Approval 8326) and Harvard Medical School Institutional
73 Review Board (Approval IRB22-0741) approved this study. Participants and guardians of participants <18
74 years old provided written informed consent or assent, as applicable. The study was granted a waiver of
75 consent for adolescents <18 years old who had not disclosed their HIV status to a guardian.

76

77 RESULTS

78
79 Among messages that participants identified as fundamental components of an anti-stigma campaign, two
80 categories emerged as targets (i.e., change goals): (1) messages that change perceptions of HIV and PLWH,
81 and (2) messages that change attitudes and behaviours towards PLWH. Within these, subcategories of
82 educational messaging were proposed to challenge the stereotypes and prejudices that lead to
83 discrimination.

84
85 *Messages that change perceptions of HIV and PLWH*

86
87 **HIV does not discriminate.** Participants highlighted the need to communicate that anyone is at risk of
88 contracting HIV regardless of sexual orientation, gender identity, age, or social class, as HIV “*doesn’t*
89 *distinguish social condition. It doesn’t matter if you’re a housewife or if you’re faithful, or if you’re not*
90 *trans, or not gay. All the same, you can [still have HIV].”* (YPLWH, cisgender woman) To combat the
91 widespread belief that HIV only affects specific populations, specifically the LGBTQIA+ community,
92 participants emphasized that messaging should clarify that risk is connected to behaviours (i.e. condomless
93 sex) and exposures rather than to groups. Participants agreed that an effective campaign would explain
94 these behaviours within a broader conversation about modes of transmission and address how HIV is not
95 transmitted to counter misconceptions that sharing utensils or having non-sexual physical contact with
96 PLWH implies a risk of acquisition. One healthcare provider emphasized:

97
98 *The majority of people think that because they use the same towel [...] or use the same soap, they’re*
99 *going to, or someone’s going to give them HIV. Obviously, we know it’s not like that, but it would be*
100 *great to make it clear that it’s through sex, or from needles, or transmission from mother to child.*
101 (Infectious Disease Specialist)

102
103 **HIV is a treatable chronic condition.** Participants recommended the campaign clearly present HIV as a
104 treatable chronic condition to change the narrative that an HIV diagnosis is a certain death sentence. One
105 participant explained, “*people should view it as if it was cancer or diabetes [...], something like that.*
106 *Because people see [HIV] as something so bad.”* (YPLWH, FSW) They emphasized that with proper
107 adherence to freely available treatment, PLWH can live normal, healthy lives, studying, working, and
108 having families. Participants expressed that these messages should also reach young people receiving a new
109 HIV diagnosis. One participant stated:

110
111 *For someone who’s just been recently diagnosed, one of the messages should be that your life’s going*
112 *to go on like always. Because a lot of people think their life is done, it’s over, or they’re not going to*
113 *be able to work, or they’re going to be rejected from jobs, things like that. The first important*
114 *message would be for them to understand that their life’s going to go on like it always has. It’s not*
115 *going to change. It’s just a condition that’s going to be there.* (YPLWH, GBMSM)

116
117 Participants advocated for messages describing the differences between HIV and AIDS to counter the
118 perception that the two are synonymous and associations of HIV with extreme sickness and death. Instead,
119 campaign messages should focus on living, vitality and the right for PLWH to participate in all aspects of
120 society without stigmatization.

121
122 **Undetectable = Untransmittable.** Participants spoke of the need to educate the general public that U = U,
123 meaning that PLWH who achieve an undetectable viral load through adherence to antiretroviral treatment
124 (ART), do not transmit HIV to others through sex. Many described their peers as not fully understanding
125 the implications of this statement, “*... they think there’s a but. So, I’m undetectable, but maybe there’s a*
126 *risk [...]*Even though there have been many campaigns, people are still scared of transmission, even if

127 *you're undetectable.*" (YPLWH, GBMSM) Participants felt messages should explain the effectiveness of
 128 ART and clarify that it is possible for PLWH to be sexually active without risk of transmitting HIV.

129
 130 Additional illustrative quotes are included for each subcategory in Table 2.

131
 132 **Table 2. Changing perceptions of HIV and PLWH: Subcategories, messages and illustrative quotes**

<i>Subcategory</i>	<i>Messages</i>	<i>Illustrative quotes</i>
HIV does not discriminate.	<ul style="list-style-type: none"> a. Anyone can contract HIV. b. HIV is a condition of behaviours and exposures rather than populations. c. How HIV is and is not transmitted 	<p><i>That people behind the screen watch this and say "wow [...] it's not just such and such people who acquire it, it could also happen to my aunt, to my dad who's a doctor, to my brother who's a student." (YPLWH, FSW)</i></p> <p><i>That HIV is a condition of behaviours, not of groups. That anyone can acquire HIV, not just those vulnerable groups, which is what we've been taught for so long. (Infectious Disease Specialist)</i></p> <p><i>Sharing, eating, drinking together, at a meet up, having fun together, we can't transmit [HIV] to them that way. That they know that transmission is only through blood, through sex, only that. (YPLWH, cisgender woman)</i></p>
HIV is a treatable chronic condition.	<ul style="list-style-type: none"> a. HIV is not a death sentence. b. HIV is not AIDS. c. PLWH can lead lives similar to those without HIV. 	<p><i>Make it clear that in this day and age, HIV isn't synonymous with death. What's more, people living with HIV have a normal life. Their life expectancy is the same as someone not living with HIV. Also, their quality of life is the same as someone not living with HIV. [Infectious Disease Specialist]</i></p> <p><i>And reinforce that we don't have AIDS, that AIDS is the last stage because a lot of times straight friends come and say, "Oh, you have AIDS, I'm sorry, you poor thing, how much time do you have left?" (HIV advocate)</i></p> <p><i>That people know that just because you're living with HIV doesn't mean your whole world ended [...] because someone living with HIV can live a normal life, you know, that would be the most important. (YPLWH, TGW)</i></p>
Undetectable = Untransmittable.	<ul style="list-style-type: none"> a. PLWH can have a fulfilling sex life without risk of transmission to their partners. b. PLWH can start a family and have children without putting them at risk of HIV. 	<p><i>People should have information and knowledge, and they should know that people who are undetectable don't transmit [HIV]. (YPLWH, TGW)</i></p> <p><i>Normalize the topic that if you have HIV you can still have sex. There are a lot of people I see who think that because they have HIV they can no longer have sex. (HIV advocate)</i></p> <p><i>I would like everyone to be clear that those of us who have HIV are taking medication, we have a life like anyone else's, with the same conditions. Basically that, you know? That yes, we are taking medication, that yes, we can be ok, we can start a family. (YPLWH, perinatal transmission)</i></p>

133
 134
 135
 136

137 *Messages that change attitudes and behaviours towards PLWH*

138

139 **Normalize talking about HIV.** Participants hoped that the anti-stigma campaign would lead to greater
140 freedom to discuss an HIV diagnosis without “... *that fear, that I don’t want to talk about HIV precisely*
141 *because I could be rejected, but rather that we’re free to talk about it*” (YPLWH, perinatal transmission)
142 and without “... *the fear that if someone finds out, what are they going to think.*” (YPLWH, perinatal
143 transmission) Participants felt that to achieve this, campaign content should present HIV as a normal topic
144 of conversation among family and friends and demonstrate how taboos around discussing HIV and resulting
145 discrimination negatively affect PLWH.

146

147 In participant experience, when HIV emerges in conversation, it is frequently accompanied by
148 discriminatory remarks or simple indifference. As one participant noted, to counter these reactions the
149 campaign should “*show what those words do to people, to us, how they affect our lives.*” (YPLWH,
150 GBMSM)

151

152 **Encourage support instead of pity for PLWH.** Participants wanted campaign messages to increase
153 awareness of what it means to live with HIV so as to remove feelings of pity. As one participant described,
154 “*I don’t want people to feel bad for me [...] and I also don’t want to hear ‘oh you poor thing’, because I’m*
155 *not a poor thing. I’m not a poor thing because I live with this.*” (YPLWH, FSW) Participants living with
156 HIV instead seek support, encouragement, and equal treatment from society. Thus, it was suggested that
157 messages educate audiences on how to approach HIV with empathy and also address feelings of self-stigma
158 and guilt among PLWH, particularly those with new diagnoses.

159

160 Though participants hope the campaign will lessen judgment of PLWH by society as a whole, special
161 mention was made of messages targeting stigma by healthcare providers. Participants suggested messages
162 foster reflection on how explicit and implicit acts of discrimination at health facilities could drive PLWH
163 away from medical care and on how all populations living with HIV deserve equal care, highlighting that
164 “*the health system has a responsibility to treat us with the same warmth and quality that everyone deserves.*”
165 (HIV advocate)

166

167 **Promote inclusion.** Participants living with HIV frequently cited a desire to be accepted and included in
168 society just as anyone else. They suggested campaign messages present acceptance of PLWH as a step
169 towards inclusion, as many times people “*know that there are people living with HIV, but that’s all, at a*
170 *distance, yeah I support them, I accept them no problem, but over there, leave them over there.*” (YPLWH,
171 GBMSM, Venezuelan migrant) However, they highlighted that true inclusion should go beyond acceptance
172 to the active invitation to participate in all aspects of society. Participants recommended that messages
173 challenge the public to include PLWH in their daily lives, especially in the workplace where participants
174 mentioned facing particular stigma and exclusion. One participant described, “*I was working in a*
175 *warehouse, they made me do a blood test and it came back positive [for HIV]. I had been working fine, but*
176 *it was really uncomfortable to have gone through the whole process and then they say ‘no you can’t work*
177 *here because you have this.*” (YPLWH, GBMSM)

178

179 **Foster unity among HIV-affected communities.** Several participants discussed their perceptions that the
180 LGBTQIA+ community itself is fractured and that for the general public to accept and include the
181 community, first “... *the community itself has to heal, because if we don’t make a change, how can we*
182 *expect other people to accept us and treat us differently?*” (YPLWH, GBMSM, Venezuelan migrant) They
183 recommend messages should strive to break down the stereotypes that different groups affected by HIV
184 have of one another to promote a united front against HIV-related stigma. As one participant noted, “*the*
185 *[LGBTQIA+] community has grown a lot [...] But it’s time we break down all these stereotypes because*
186 *we’re all one community and we’re all susceptible to HIV, to stigma, and to all the prejudice that exists.*”
187 (HIV advocate)

188
189 Additional illustrative quotes are included for each subcategory in Table 3

190
191 **Table 3. Changing attitudes and behaviours towards PLWH: Subcategories, messages and**
192 **illustrative quotes**

<i>Subcategory</i>	<i>Messages</i>	<i>Illustrative quotes</i>
Normalize talking about HIV	a. PLWH should be able to speak about HIV without fear of rejection.	<i>That we can talk to people about our problems without being afraid, talk about what we have, and that there's trust there so we can just be ourselves, knowing they won't discriminate. (YPLWH, perinatal transmission)</i>
	b. HIV should be able to be discussed just as any other topic.	<i>That people aren't afraid to talk about the issue. It should be just like talking about, I don't know, chocolate or vanilla ice cream, a normal thing. (YPLWH, GBMSM)</i>
	c. Discriminatory comments and reactions negatively affect PLWH.	<i>HIV isn't what's going to kill me, what's going to kill me is your disrespect, your discrimination, your distance. The disease won't kill me because it's a disease that with medication I'm going to control and then it'll be undetectable. But your indifference, how you look at me, you staying away from me, those are what kill me, stigmatize. Without saying anything you're discriminating against me. (HIV advocate, FSW)</i>
Encourage support instead of pity for PLWH	a. PLWH desire encouragement, motivation and empathy from the general population.	<i>Another thing I want people to know is that if they were living with HIV, they would want to be treated with respect. People should know that and think about how they would like to be treated. So they can put themselves in the other person's shoes, understand a bit, and reflect, because most people think this will never happen to them. (Healthcare provider)</i>
	b. Healthcare providers must provide empathetic, quality care to all.	<i>That people don't discriminate against us if we're living with HIV, that they support us, motivate us to keep living our lives just like anyone else. (YPLWH, TGW)</i> <i>Demonstrate that, or make [healthcare providers] aware that discriminating against people living with HIV pushes them away. (Healthcare provider)</i>
Promote inclusion	a. PLWH must be accepted and included in all aspects of society.	<i>... that we can participate, that people are more fair, that they include us, not exclude us. (YPLWH, TGW)</i>
	b. Discrimination against PLWH in the workplace is illegal and unacceptable.	<i>I'd love for you to get people to say, 'Wow, these [PLWH] are doing a service to society; they don't deserve to be discriminated against'. And then maybe people would see us being discriminated against and start to defend us, even people with status and power. And maybe people in the workplace would become aware and even say, 'I want to hire someone living with HIV because I know they might have more skills than others [...]' (YPLWH, GBMSM)</i>
Foster unity among HIV-affected communities	a. The LGBTQIA+ community should unite to counter HIV-related stigma and promote acceptance.	<i>I know that [stigma faced by trans women] is more intense, but we must break those stereotypes that we have, that discrimination against each other, the barriers between us. It's time we come together. We all have problems, some fewer than others of course, but society should understand that any of us could [have HIV]. No one is free from this. (HIV advocate)</i> <i>TGW 1: To start we should change our own mentality. We shouldn't attack each other, we should support each other. That's the most important.</i> <i>TGW 2: That would be the first step.</i> <i>TGW 1: Exactly, and then show society that we are united and that they should also accept us just as we are. (YPLWH, TGW)</i>

193
194

195 The underlying theme of all messages was that PLWH are no different from anyone else, with similar hopes,
196 fears, and aspirations, and should not be seen “... *as good or bad, but rather just as normal people, like*
197 *anyone else, with the same life expectancy, the same access, the same ability to work, the same difficulties*
198 *as anyone else.*” (HIV advocate)

199
200 *Content Delivery*

201
202 **Videos** Participants agreed that short videos (15-30 seconds) with captivating intros are the most powerful
203 means of conveying anti-stigma messages. The following types of videos were mentioned as potentially
204 most effective: storytimes (a TikTok trend in which someone films themselves telling a story in a casual
205 setting often while doing another activity), testimonials both positive and negative, and social experiments
206 in which PLWH interact with the public. Participants considered that spokespeople in these videos should
207 represent diverse identities in all aspects and especially gender, sexual orientation, race, socioeconomic
208 status, education, and religion. As one participant advised, “*Pay attention to the profiles you choose. They*
209 *should be varied in terms of orientation, identity, race, income, and everything else, because it’s*
210 *crosscutting, you know.*” (HIV advocate)

211
212 Generally, participants felt that PLWH would be more credible than actors sharing these messages and that
213 it would be essential that all actors and those creating content or disseminating the campaign receive training
214 on HIV and HIV-related stigma. Most agreed that “*the majority should be PLWH, but also show people*
215 *who aren’t living with HIV because maybe they’re also affected and want to talk about it.*” (HIV advocate)
216 According to participants, family members, friends, or colleagues of PLWH would be compelling
217 messengers of anti-stigma information.

218
219 *Focus group leader: Who should give these messages?*
220 *TGW 1: A trans woman, a gay man, an actor, or whoever you think is convenient and would do a*
221 *good job.*
222 *TGW 2: Exactly. To get a clear and direct message to the public.*
223 *[YPLWH, TGW]*

224
225 **Audience** Many participants, particularly those involved in HIV activism, noted that the majority of media
226 campaigns related to HIV are aimed at the LGBTQIA+ community. They agreed that this reinforces the
227 stereotype that HIV is only an issue for a single population and suggested that an anti-stigma campaign
228 should be aimed at all audiences equally.

229
230 *Also don’t just shut yourself in the community, I mean the gay and trans community, it would also be*
231 *good for the general population to be informed. Because the campaigns they do, it’s true, they’re*
232 *just for the gay and trans community, key populations. The general population always asks, always*
233 *has questions, so to break with that misinformation they have, it would be great to share this with*
234 *everyone.* (HIV advocate)

235
236 **Language** The importance of careful selection of language was another key area related to content delivery.
237 In Spanish, participants preferred ‘transmitir’ (to transmit) over ‘contagiar’ (to infect) when discussing
238 HIV-acquisition, and ‘vivir con VIH’ (to live with HIV) or ‘ser seropositivo’ (to be seropositive) to identify
239 PLWH. Participants described a feeling of coexisting with the virus as “[HIV] *is just one of the many*
240 *qualities one may have*” (HIV advocate) and as such agreed that HIV should be described as a health or life
241 condition rather than a disease.

242
243 **Images** The red ribbon symbolic of HIV was controversial when discussing imagery to associate with an
244 HIV stigma campaign. Many participants felt that the red ribbon is in itself stigmatizing as it labels the user
245 as having some relationship with AIDS. Others mentioned that the red ribbon has been over-used and that

246 ribbons, in general, are employed in sad or negative situations, contradicting the campaign's objective of
247 eliminating feelings of pity towards PLWH. However, a sub-group of activist participants noted that, for
248 them, the red ribbon is a symbol of solidarity.

249
250 Participants suggested not using images frequently associated with HIV-campaigns, such as condoms, pills,
251 or drops of blood, and were in favour of a unique logo, perhaps one incorporating a subtle reference to the
252 red ribbon (e.g. an animal mascot wearing a red scarf). In general, "[Imagery should be] as normal as
253 possible and not related to disease. If there are condoms or pills, then you are referencing the disease."
254 (YPLWH, GBMSM) The use of the LGBTQIA+ flag in the campaign was met with caution as it could
255 perpetuate the stereotype that HIV only affects members of this community.

256
257 Supplemental Table 2 includes additional illustrative quotes of campaign language and imagery.

258 259 DISCUSSION

260
261 In this study, a diverse group of YPLWH and other stakeholders, including healthcare providers and
262 advocates, identified two overarching targets (i.e., change goals) for a campaign to reduce HIV-related
263 stigma in Peru: to change how PLWH are perceived and to change attitudes and actions towards PLWH.
264 While these change targets emerged organically, they coincide with theoretical models for stigma and
265 stigma reduction that posit that a lack of correct information leads to prejudices and stereotypes and,
266 ultimately, discrimination (also known as enacted or experienced stigma), and that stigma change programs
267 should include specific change targets or goals(1)[33]. The messaging associated with the two change goals,
268 while initially targeted broadly toward youth living in Peru, also has the potential to positively impact
269 YPLWH by reducing self-stigma (i.e., stigma that results when a member of a stigmatized group believes
270 the stereotypes and prejudices associated with the stigmatized condition and applies them to the self). In
271 Peru, like many places in Latin America, PLWH frequently experience a high level of self-stigma[28].

272
273 Results suggest that messaging aligned with the first change target should convey factual information about
274 HIV and what it means to live with HIV. Three specific messages were identified as critical: (1) HIV does
275 not discriminate (i.e., no one is immune from potentially acquiring HIV); (2) HIV is a treatable chronic
276 condition; and, (3) if HIV is controlled (i.e., undetectable) in one's body, it will not be transmitted sexually
277 (i.e., U=U, or I=I in Spanish). Providing education to correct common misconceptions and erroneous beliefs
278 (e.g., "HIV is synonymous with death", "PLWH cannot have sex or start a family without risking HIV
279 transmission to their partner or baby") is a destigmatizing strategy that can reduce the prejudices and
280 stereotypes that lead to discriminatory behaviour[36]. The potential for community-derived U=U
281 messaging in combatting stigma reduction has emerged as a particularly promising educational messaging
282 strategy[37],[38]; however, less awareness of this concept has been reported in low- and middle-income
283 countries[38]. Messages aligning with the second change target aimed to address prejudice and
284 discrimination through actions, including normalizing open discussion about HIV, offering support versus
285 pity, ensuring inclusion in all aspects of societal life, and fostering unity within the different communities
286 most affected by HIV. Much attention has been paid to reducing stigma toward members of stigmatized
287 groups, but less to fostering affirming attitudes and building allyships, which is critical to improving
288 outcomes for stigmatized groups[39].

289
290 Translating key messages to content appealing to young people is critical to ensuring effectiveness. Our
291 findings offer practical suggestions for the creation of content to communicate these messages, some of
292 which broadly align with prior studies[39-41]. Among these is the importance of diversity in representation
293 to ensure relatability across various identities and because a focus on a single population, such as the
294 LGBTQIA+ community, may reinforce existing stigmas[40],[41]. We found that content imagery and tone
295 are also key considerations. Imagery that reinforces stereotypes or that is overly negative or aggressive may
296 be less effective than lighter messages, free of shame and judgment[42]. For example, several participants

297 referenced antiquated images from the pre-ART era that immortalize the misconception that HIV is
298 analogous to death. An upbeat tone with positive imagery may more appropriately reflect the lifesaving
299 advances in HIV treatment. Other research offers pragmatic guidance for social media strategies targeting
300 young people. For example, one study found that provocative images and wording to grab the user’s
301 attention, followed by fact-based content, were effective in motivating engagement with services
302 offline[42].

303
304 Language emerged as another important consideration. Although a desire to avoid stigmatizing words is
305 intuitive, language’s subjective and evolving nature may challenge this goal. Some participants noted that
306 they preferred the term “seropositive” to describe themselves and others living with HIV, while other
307 stakeholders in our study and beyond have identified the term as stigmatizing[43],[44]. Community
308 advisory committees, ideally comprised of people of similar ages as the target population, should play a
309 key role in validating campaign content and ensuring that the language is correct, understandable,
310 acceptable, and appealing without reinforcing stigma.

311
312 A strength of this work was the engagement of a diverse group of YPLWH representative of the groups
313 most impacted by HIV in Peru. Effective community engagement to inform campaign messaging ensures
314 that campaign priorities align with those of the most affected groups. While qualitative methodologies are
315 one strategy to elicit community input[41], other community-based participatory methods such as
316 crowdsourcing[37],[41],[44], and discrete choice experiments have also been used to engage communities
317 in public health campaigns. By focusing on the opinions of PLWH and incorporating those of health
318 providers and advocates, we aimed to elevate the voices of those with lived experience. We did not validate
319 our findings among the potential targets of the campaign (i.e., young people without a known HIV
320 diagnosis). If there were a misalignment between the messaging strategies identified by study participants
321 and those of the campaign targets, this could undermine the effectiveness of subsequent campaign content.
322 Finally, although we aimed to explore messages about stigmas that intersect with HIV (i.e., stigmas against
323 GBMSM or TGW), this proved challenging. While participants readily shared experiences of stigma due
324 to different parts of their identity, we could not generate in-depth discussion around lived experiences at
325 the intersection of these stigmas. Intersectional stigma is conceptually difficult to measure, perhaps even
326 more so in young people with fewer years of lived experience. Our qualitative guides likely required greater
327 depth and attention to the lived experiences of intersectional stigmas than we gave them in this multi-aim
328 study[45].

329 CONCLUSIONS

331 We employed a community-engaged approach to identify change goals, messages, and messaging
332 preferences to underpin a Spanish-language social media campaign targeting young people. While this
333 formative research was exclusive to Peru, the findings broadly align with the existing literature and likely
334 have broad applicability, expanding the evidence base regarding the use of social media to reduce stigma
335 related to HIV and, potentially, to other stigmatized conditions.

336 ACKNOWLEDGEMENTS

338 The authors are indebted to the study participants, and to the healthcare providers and community partners
339 who supported the study team in participant recruitment. We would also like to express our gratitude to
340 the Fogarty International Center for their financial support and to the Socios en Salud Youth Advisory
341 Board for their guidance throughout our project.

342 COMPETING INTERESTS

343 The authors report there are no competing interests to declare.

344 AUTHORS’ CONTRIBUTIONS

345 Study design: M.W., K.K., J.G., M.F., R.E.

346 Analysis design: K.K., J.G., M.F., R.E.
347 Study implementation: M.W.
348 Data management: A.N., M.C., M.W.
349 Data analyses: A.N., M.C., M.W., K.K., R.E.
350 Data interpretation: all authors
351 First draft: A.N., M.C., K.K., M.F., R.E.
352 Critical review of manuscript: all authors
353 Approved final version: all authors
354

355 DATA AVAILABILITY

356 The data that support the findings of this study are available upon reasonable request from the
357 corresponding author. The data are not publicly available due to privacy or ethical restrictions.

358 FUNDING

359 This work was supported by the Fogarty International Center of the National Institutes of Health under
360 award number R01TW012394.

361 SUPPORTING INFORMATION

362 **Supplemental Table 1:** Qualitative data collection guide developed for focus groups and interviews with
363 PLWH

364 **Supplemental Table 2:** Illustrative quotes for campaign language and imagery

365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380

381 REFERENCES

- 382 1. Pescosolido BA, Martin JK. The stigma complex. *Annu Rev Sociol.* 2015;41.
383 <https://doi.org/10.1146/annurev-soc-071312-145702>
- 384 2. Relf MV, Holzemer WL, Holt L, Nyblade L, Ellis Caiola C. A review of the state of the science of
385 HIV and stigma: context, conceptualization, measurement, interventions, gaps, and future
386 priorities. *J. Assoc Nurses AIDS Care.* 2021;32(3). <https://doi.org/10.1097/jnc.0000000000000237>
- 387 3. Katz IT, Ryu AE, Onuegbu AG, Psaros C, Weiser SD, Bangsberg DR, et al. Impact of HIV-related
388 stigma on treatment adherence: systematic review and meta-synthesis. *J Int AIDS Soc.* 2013;16(3
389 Suppl 2):18640. <https://doi.org/10.7448/ias.16.3.18640>
- 390 4. Gesesew HA, Gebremedhin AT, Demissie TD, Kerie MW, Sudhakar M, Mwanri L. Significant
391 association between perceived HIV related stigma and late presentation for HIV/AIDS care in low
392 and middle-income countries: A systematic review and metaanalysis. *PLoS ONE.* 2017;
393 12(3):e0173928. <https://doi.org/10.1371/journal.pone.0173928>
- 394 5. Fischer LS, Mansergh G, Lynch J, Santibanez S. Addressing disease-related stigma during
395 infectious disease outbreaks. *Disaster Med Public Health Prep.* 2019;13(5–6).
396 <https://doi.org/10.1017/dmp.2018.157>
- 397 6. Kerrigan D, Vazzano A, Bertoni N, Malta M, Bastos FI. Stigma, discrimination and HIV
398 outcomes among people living with HIV in Rio de Janeiro, Brazil: The intersection of multiple
399 social inequalities. *Glob Public Health.* 2017;12(2).
400 <https://doi.org/10.1080/17441692.2015.1064459>
- 401 7. Cloete A, Simbayi LC, Kalichman SC, Strebel A, Henda N. Stigma and discrimination
402 experiences of HIV-positive men who have sex with men in Cape Town, South Africa. *AIDS*
403 *Care.* 2008;20(9). <https://doi.org/10.1080/09540120701842720>
- 404 8. Oduenyi C, Ugwa E, Ojukwu Z, Ojukwu-Ajasigwe J. An exploratory study of stigma and
405 discrimination among people living with HIV/ AIDS in Abuja Municipal Area Council, Nigeria.
406 *Afr J Reprod Health.* 2019;23(1). <https://doi.org/10.29063/ajrh2019/v23i1.9>
- 407 9. Kalichman SC, Simbayi LC. HIV testing attitudes, AIDS stigma, and voluntary HIV counselling
408 and testing in a black township in Cape Town, South Africa. *Sex Transm Infect.* 2003;79(6).
409 <https://doi.org/10.1136/sti.79.6.442>
- 410 10. Gamarel KE, Nelson KM, Stephenson R, Santiago Rivera OJ, Chiamonte D, Miller RL.
411 Anticipated HIV stigma and delays in regular HIV testing behaviors among sexually-active young
412 gay, bisexual, and other men who have sex with men and transgender women. *AIDS Behav.*
413 2018;22(2). <https://doi.org/10.1007/s10461-017-2005-1>
- 414 11. Earnshaw VA, Chaudoir SR. From conceptualizing to measuring HIV stigma: A review of HIV
415 stigma mechanism measures. Vol. 13, *AIDS Behav.* 2009; 13(6). <https://doi.org/10.1007/s10461-009-9593-3>
416

- 417 12. Benavides JA, Caparrós C, da Silva RM, Lembo T, Tem Dia P, Hampson K, et al. The power of
418 music to prevent and control emerging infectious diseases. *Front Med (Lausanne)*. 2021;8.
419 <https://doi.org/10.3389/fmed.2021.756152>
- 420 13. Rivera A V., DeCuir J, Crawford ND, Amesty S, Harripersaud K, Lewis CF. Factors associated
421 with HIV stigma and the impact of a nonrandomized multi-component video aimed at reducing
422 HIV stigma among a high-risk population in New York City. *AIDS Care*. 2015;27(6).
423 <https://doi.org/10.1080/09540121.2014.998611>
- 424 14. Fakolade R, Adebayo SB, Anyanti J, Ankomah A. The impact of exposure to mass media
425 campaigns and social support on levels and trends of HIV-related stigma and discrimination in
426 Nigeria: Tools for enhancing effective HIV prevention programmes. *J Biosoc Sci*. 2010;42(3).
427 <https://doi.org/10.1017/s0021932009990538>
- 428 15. Aghaei A, Sakhaei A, Khalilimeybodi A, Qiao S, Li X. Impact of mass media on HIV/AIDS
429 stigma reduction: a systematic review and meta-analysis. *AIDS Behav*. 2023;27(10).
430 <https://doi.org/10.1007/s10461-023-04057-5>
- 431 16. Wakefield MA, Loken B, Hornik RC. Use of mass media campaigns to change health behaviour.
432 *Lancet*. 2010;376(9748). [https://doi.org/10.1016/s0140-6736\(10\)60809-4](https://doi.org/10.1016/s0140-6736(10)60809-4)
- 433 17. Kite J, Chan L, MacKay K, Corbett L, Reyes-Marcelino G, Nguyen B, et al. A model of social
434 media effects in public health communication campaigns: systematic review. *J Med Internet Res*.
435 2023;25:e46345. <https://doi.org/10.2196/46345>
- 436 18. Friedman AL, Brookmeyer KA, Kachur RE, Ford J, Hogben M, Habel MA, et al. An assessment
437 of the GYT: Get yourself tested campaign: An integrated approach to sexually transmitted disease
438 prevention communication. *Sex Transm Dis*. 2014;41(3).
439 <https://doi.org/10.1097/olq.000000000000100>
- 440 19. Vazquez Guillaumet LJ, Babey MM, Njah M, Blake H, Jasani A, Kyeng R, et al. Acceptability,
441 feasibility and potential of an intervention using secret Facebook groups to complement existing
442 HIV prevention strategies among female sex workers in Cameroon, a randomized pilot study.
443 *AIDS Care*. 2024;36(4). <https://doi.org/10.1080/09540121.2023.2275048>
- 444 20. Young SD, Cumberland WG, Lee SJ, Jaganath D, Szekeres G, Coates T. Social networking
445 technologies as an emerging tool for HIV prevention: a cluster randomized trial. *Ann Intern Med*.
446 2013;159(5). <https://doi.org/10.7326/0003-4819-159-5-201309030-00005>
- 447 21. Jalloh, C.; Beaudoin, C.; Kimani, J.; Rojas, C.; Castro, D.; Beti, B.; Wylie, J. “Online and on the
448 ground”: A pilot initiative to share electronic sexual health information with youth in Canada,
449 Colombia & Kenya. *Can J Infect Dis Med Microbiol*. 2014;25(82A)
- 450 22. Alianza Liderazgo en Positivo & Poblaciones Clave. Estudio multipaís índice de estigma y
451 discriminación en personas con VIH. 2023. Available at:
452 <https://www.slideshare.net/slideshow/informe-index-multipais-20pdf/259778924>

- 453 23. Piñeirúa A, Sierra-Madero J, Cahn P, Guevara Palmero RN, Martínez Buitrago E, Young B, et al.
454 The HIV care continuum in Latin America: challenges and opportunities. *Lancet Infect Dis*.
455 2015;15(7). [https://doi.org/10.1016/s1473-3099\(15\)00108-5](https://doi.org/10.1016/s1473-3099(15)00108-5)
- 456 24. Chow JY, Konda KA, Borquez A, Caballero P, Silva-Santisteban A, Klausner JD, et al. Peru's
457 HIV care continuum among men who have sex with men and transgender women: opportunities to
458 optimize treatment and prevention. *Int J STD AIDS*. 2016;27 (12).
459 <https://doi.org/10.1177/0956462416645727>
- 460 25. Shin S, Muñoz M, Caldas A, Wu Y, Zeladita J, Wong M, et al. Mental health burden among
461 impoverished HIV-positive patients in Peru. *J Int Assoc Physicians AIDS Care*. 2011;10(1).
462 <https://doi.org/10.1177/1545109710385120>
- 463 26. Nayar US, Stangl AL, De Zalduondo B, Brady LM. Reducing stigma and discrimination to
464 improve child health and survival in low-and middle-income countries: Promising approaches and
465 implications for future research. *J Health Commun*. 2014;19(sup1).
466 <https://doi.org/10.1080/10810730.2014.930213>
- 467 27. De Boni R, Veloso VG, Grinsztejn B. Epidemiology of HIV in Latin America and the Caribbean.
468 Vol. 9, *Curr Opin HIV AIDS*. 2014;9(2). <https://doi.org/10.1097/coh.0000000000000031>
- 469 28. Silva-Santisteban A, Bustamante MJ. Índice de estigma y discriminación hacia las personas con
470 VIH en Perú. 2018. Available at: [https://www.stigmaindex.org/wp-](https://www.stigmaindex.org/wp-content/uploads/2019/11/Peru_PLHIV-Stigma-Index-Report_2018.pdf)
471 [content/uploads/2019/11/Peru_PLHIV-Stigma-Index-Report_2018.pdf](https://www.stigmaindex.org/wp-content/uploads/2019/11/Peru_PLHIV-Stigma-Index-Report_2018.pdf)
- 472 29. UNAIDS. Country fact sheet, Peru. 2023. Available at:
473 <https://www.unaids.org/en/regionscountries/countries/peru>
- 474 30. Ministerio del Perú. Situación epidemiológica del VIH-Sida en el Perú. Ministerio del Perú.
475 2021;Febrero 20(WHO Library Cataloguing-in-Publication Data). Available at:
476 https://www.dge.gob.pe/epipublic/uploads/vih-sida/vih-sida_202112_11_111653.pdf
- 477 31. Perla ME, Ghee AE, Snchez S, McClelland RS, Fitzpatrick AL, Surez-Ognio L, et al. Genital tract
478 infections, bacterial vaginosis, HIV, and reproductive health issues among lima-based clandestine
479 female sex workers. *Infect Dis Obstet Gynecol*. 2012;2012. <https://doi.org/10.1155/2012/739624>
- 480 32. International Organization of Migration, Partners in Health, United States Centers for Disease
481 Control and Prevention. Biobehavioral Survey (BBS) among Venezuelan migrants living in
482 Lima/Callao and Trujillo. 2023. Available at: [https://migrationhealthresearch.iom.int/final-report-](https://migrationhealthresearch.iom.int/final-report-biobehavioral-survey-bbs-among-venezuelan-migrants-living-limacallao-and-trujillo)
483 [biobehavioral-survey-bbs-among-venezuelan-migrants-living-limacallao-and-trujillo](https://migrationhealthresearch.iom.int/final-report-biobehavioral-survey-bbs-among-venezuelan-migrants-living-limacallao-and-trujillo)
- 484 33. Stangl AL, Earnshaw VA, Logie CH, Van Brakel W, Simbayi LC, Barré I, et al. The Health
485 Stigma and Discrimination Framework: A global, crosscutting framework to inform research,
486 intervention development, and policy on health-related stigmas. *BMC Med*. 2019;17(1).
487 <https://doi.org/10.1186/s12916-019-1271-3>
- 488 34. Goldsmith LJ. Using framework analysis in applied qualitative research. *Qualitative Report*.
489 2021;26(6). <https://doi.org/10.46743/2160-3715/2021.5011>

- 490 35. Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research (COREQ): A
491 32-item checklist for interviews and focus groups. *Int J Qual Health Care*. 2007;19(6).
492 <https://doi.org/10.1093/intqhc/mzm042>
- 493 36. Corrigan PW, Morris SB, Michaels PJ, Rafacz JD, Rüsçh N. Challenging the public stigma of
494 mental illness: A meta-analysis of outcome studies. *Psychiatr Serv*. 2012;63(10).
495 <https://doi.org/10.1176/appi.ps.201100529>
- 496 37. Smith PJ, Joseph Davey DL, Schmucker L, Bruns C, Bekker LG, Medina-Marino A, et al.
497 Participatory prototyping of a tailored undetectable equals untransmittable message to increase
498 HIV testing among men in Western Cape, South Africa. *AIDS Patient Care STDS*. 2021;35(11).
499 <https://doi.org/10.1089/apc.2021.0101>
- 500 38. Bor J, Fischer C, Modi M, Richman B, Kinker C, King R, et al. Changing knowledge and attitudes
501 towards HIV treatment-as-prevention and “undetectable = untransmittable”: a systematic review.
502 *AIDS Behav*. 2021;25(12). <https://doi.org/10.1007/s10461-021-03296-8>
- 503 39. Gonzalez KA, Riggle EDB, Rostosky SS. Cultivating positive feelings and attitudes: A path to
504 prejudice reduction and ally behavior. *Transl Issues Psychol Sci*. 2015;1(4).
505 <https://doi.org/10.1037/tps0000049>
- 506 40. Fields EL, Long A, Bademosi K, Granderson R, Schumacher CM, Chandran A, et al. Identifying
507 community-informed language to promote HIV pre-exposure prophylaxis (Prep) in black LGBTQ
508 communities in Baltimore. *AIDS Educ Prev*. 2020;32(2).
509 <https://doi.org/10.1521/aeap.2020.32.2.152>
- 510 41. Kerr J, Ayangeakaa S, Combs R, Harris L, Sears J, Northington T, et al. Community-informed
511 development of a campaign to increase HIV pre-exposure prophylaxis (PrEP) awareness among
512 African-American young adults. *J Racial Ethn Health Disparities*. 2021;8(4).
513 <https://doi.org/10.1007/s40615-020-00848-x>
- 514 42. Goedel WC, Suttén Coats C, Sowemimo-Coker G, Moitra E, Murphy MJ, van den Berg JJ, et al.
515 Gay and bisexual men’s recommendations for effective digital social marketing campaigns to
516 enhance HIV prevention and care continuity. *AIDS Behav*. 2021;25(5).
517 <https://doi.org/10.1007/s10461-020-03078-8>
- 518 43. UNAIDS Terminology Guidelines. 2024. Available at:
519 https://www.unaids.org/en/resources/documents/2024/terminology_guidelines
- 520 44. Shah HS, Grieb SD, Flores-Miller A, Phillips KH, Page KR, Cervantes A, et al. A crowdsourcing
521 open contest to design a Latino-specific COVID-19 campaign: mixed methods analysis. *JMIR*
522 *Form Res*. 2022;6(5). <https://doi.org/10.2196/35764>
- 523 45. Turan JM, Elafros MA, Logie CH, Banik S, Turan B, Crockett KB, et al. Challenges and
524 opportunities in examining and addressing intersectional stigma and health. *BMC Med*.
525 2019;17(1). <http://dx.doi.org/10.1186/s12916-018-1246-9>