Women's experiences collecting and accessing water in Guatemala, Honduras, Kenya, and Zimbabwe:

A mixed-methods investigation

Bethany A. Caruso[1*], Thea Mink[1], Madeleine Patrick[1], Emily Ogutu[2], Cameron Dawkins[1], Olivia Bendit[1],

Mahnoor Fatima[1], Ingrid Lustig[1], Alicia Macler[1], Jera White[1], Alondra Zamora[1], Alberto Emanuel Santos

López[3], Héctor Salvador Peña Ramírez[3], Carlos Daniel Sic[3], Jorge Lemus Chávez[4], Sandra Antonio[5],

Jazmina Nohemí Irías[5], Gladys Ramos[6], Everlyne Atandi[7], Peter Mwangi[7], Peter Koome[8], Rohin Otieno

Onyango[8], Petronilla Andiba Otuya[8], Paul Ruto[8], Morris Chidavaenzi[9], Jammaine Jimu[9], Sithandekile

Maphosa[9], Makaita Maworera[9], Munyaradzi Damson[10], Sheela S. Sinharoy[1]

[1] Hubert Department of Global Health, Rollins School of Public Health, Emory University, Atlanta, GA,

USA

[2] Gangarosa Department of Environmental Health, Rollins School of Public Health, Emory University,

Atlanta, GA, USA

[3] World Vision Guatemala, Guatemala City, Guatemala

[4] Centro Universitario de Occidente, de la Universidad de San Carlos de Guatemala, Quetzaltenango,

Guatemala

[5] World Vision Honduras, Tegucigalpa, Honduras

[6] Universidad Nacional Autónoma de Honduras, Tegucigalpa, Honduras

[7] World Vision Kenya, Nairobi, Kenya

[8] St. Paul's University, Nairobi, Kenya

[9] World Vision Zimbabwe, Harare, Zimbabwe

[10] Datalyst Africa, Harare, Zimbabwe

* Corresponding author

Email: bcaruso@emory.edu (BAC)

ABSTRACT

1.8 billion people live in households that collect water from sources off household premises, creating burdens that disproportionately affect women. A current and comprehensive investigation of women's water collection experiences is needed to understand the gendered burden of this labor. This study used mixed methods, including (a) go-along, in-depth interviews (IDIs), (b) semi-structured observation, (c) activity-tracking smart watches, and (d) scales, to (1) understand women's practices, perspectives, and experiences going to water sources; (2) determine actual water collection time, distance, caloric expenditure, total elevation ascended, weight carried, and water volume collected; and (3) assess alignment of women's estimated and actual water journey times. Ninety-four women participated across four countries: Guatemala (n=22), Honduras (n=17), Kenya (n=22), Zimbabwe (n=33). Women reported accessing various sources depending on season and needs, faced risks due to terrain and animals, and experienced physical injury and mental burden. Experiences varied within and by country. The mean water journey time (including going to the source, activities at the source, and returning) was 82 minutes (range: 13 minutes (Guatemala) – 287 minutes (Kenya)). The mean distance traveled was 3.5km (range: 0.2km (Guatemala, Honduras, Zimbabwe) - 15.8km (Kenya)). Mean caloric expenditure was 231cal (range: 36 (Guatemala) – 952 (Zimbabwe)). The mean volume of water collected was 16.1L. (range: 3.7L (Kenya) - 38.2L (Zimbabwe). Women also carried children, wet laundry, and other items resulting in heavy loads. The mean total weight of loads brought from water sources was 19.3kg (range: 5.0kg (Honduras) - 50.1kg (Zimbabwe). Findings demonstrate how a lack of adequate and accessible water drains women of energy and time and poses risks to their well-being. Our findings reinforce the need to redouble efforts to improve water access in low-resource settings and rigorously measure the impacts of such efforts on women's lives.

1 INTRODUCTION

Water is critical for health(1) and sustaining life(2) and is a basic human right(3); yet a large proportion of the global population still lacks access, creating burdens that disproportionately affect women and girls(4). An estimated 1.8 billion people live in households where water is collected from sources located off household premises; in 70% of these households, women and girls are the primary water collectors (63% women and girls aged 15 and over; 7% girls younger than age 15)(4). This unpaid labor and drudgery are driven by sociocultural norms that conscript women to carry out this time-consuming, arduous, and risk prone work(5, 6). These quantitative estimates of water collection responsibility are vital for showing who bears this burden, but they do not capture what this burden entails. While thorough assessments of water collection time and energy requirements exist, they are outdated and primarily focused on East Africa(7)(8). Thus, a current, comprehensive, and more diverse investigation of water collection experiences, particularly among women, is needed to understand the gendered burden of this labor.

Research on how water fetching can impose disproportionate physical harms and cognitive demands on women is growing(9). Water fetching has been linked to poor musculoskeletal health, broken limbs, lacerations, general pain, fatigue, perinatal health problems, stress, chest pain, headaches, and hair loss for those carrying on the head(10-17). Research conducted in 21 low-income and middle-income countries found that 85% of respondents (6291/7401) reported experiencing a water-fetching-related injury. Among those, 72% were female(12). Further, research from Odisha, India illuminated the cognitive burden of water collection and management among women who described the myriad factors they needed to weigh when determining where to access water, including household needs (e.g., drinking, cooking), water characteristics (e.g., taste), source features (e.g., distance), and season, among others(18).

The imperative to improve water access to save women's time and energy was argued nearly 40

years ago(19), and while water access has improved to varying extents, water collection continues to

place substantial burdens on women's time and energy(20). Specific understanding of the time required

for water collection is improving, yet the impact of water collection on energy, beyond qualitative

reports of exhaustion, remains limited. Globally, women collectively spend an estimated 250 million

hours per day collecting water, over three times the amount of time spent by men.(21) The longer the

time needed to fetch water, the less likely that men are involved(22), further exacerbating the

disproportionate burden placed on women. As such, women are more likely to fetch water when the

work is intensified during dry seasons or climate-related droughts(23), which can require farther travel,

longer queues, and elevated water extraction effort(24-26). Moreover, as water collection times and

distances increase, the volume of water collected decreases(7, 19, 27, 28), meaning that women's

increasing efforts likely yield water quantities that are insufficient to meet their needs.

Water collection time has been associated with individuals' education and economic

opportunities and health (12, 29-34), reinforcing the need for more precise understanding of the

amount of time spent collecting water. Specifically, increases in time for water fetching have been

associated with increased risk of injury(12) and intimate partner violence(29). Research in India found

higher water-fetching times among children to be associated with lower school math, reading, and

writing scores(30). Conversely, improved water access has resulted in time savings, enabling leisure,

social, and economic activities(31, 32). For example, the provision of piped water into households in

Malawi resulted in a 77% reduction in time spent collecting water among women, 69% among girls, and

72% among boys, amounting to an estimated 3.8 hours of time savings per household per week, mostly

for women and girls. Further, water use increased by 32%, including for domestic and productive

uses(32). Additionally, a decrease in roundtrip water collection time by 15 minutes in Sub-Saharan Africa

was associated with improved child health outcomes(33), which may further reduce time women spend

caring for sick children(34).

Better data on time, distance, and energy related to water collection are needed. Most data on

water collection time are self-reported; other methods to measure time are sparse(17, 35-37). While

self-report time data are critical for large-scale global monitoring efforts, more precise measures are

needed for research(38). A limited body of research has compared self-reported and measured water

collection time with mixed results, self-reported time to be underestimated in some settings and

overestimated in others(35-37). Data is also limited on distances traveled to water sources(39); among

studies that report distance, most involved participants who traveled less than 1 kilometer(35, 40-42).

Distance may be understudied because time is often used as a proxy(17, 33, 43). However, time is an

inaccurate substitute for measuring distance(37, 44); variability in terrain, elevation, weather, physical

ability, health status and the load carried all influence the time it takes to traverse the same distance(45).

Finally, women likely expend considerable energy collecting water and doing water work, like laundry

and watering animals and garden crops, at sources, yet little is known about this energy expenditure.

Existing studies are limited by small sample sizes and use simulations to generate estimates(46, 47).

Research that improves understanding of the energy and time required and distance traveled for water

work is needed.

This study sought to comprehensively understand women's burden traveling to, performing

work at, and returning from water sources in four countries: Guatemala, Honduras, Kenya, and

Zimbabwe. It aimed to: (1) understand women's practices, perspectives, and experiences going to water

sources; (2) determine actual water collection time, distance, caloric expenditure, total elevation

ascended, work performed at source, weight carried, and water volume collected and transported; (3)

assess alignment of women's estimated and actual water journey times.

2 METHODS

2.1 Study Design and Motivation

This mixed methods study used (a) go-along, in-depth interviews (IDIs), (b) semi-structured observation, (c) activity-tracking smart watches, and (d) scales to elucidate women's experiences going to water sources (**Figure 1**). Research took place across four countries (Guatemala, Honduras, Kenya, and Zimbabwe) in communities where World Vision is delivering its *Strong Women, Strong World: Beyond Access* (SWSW) programming. The SWSW program seeks to increase women's empowerment and improve women's well-being through an integrated intervention that includes water improvements and economic empowerment training and support. The SWSW program theorizes that by providing better water services, women will have increased access to water, saving them time and energy and enabling them to use water, time, and energy for other purposes, including economic activities. World Vision hypothesized that women in the SWSW program areas expend substantial time and energy to retrieve limited quantities of water. To test this hypothesis, Emory University worked with World Vision USA and local World Vision offices in Guatemala, Honduras, Kenya, and Zimbabwe to design the present study. Each World Vision country office identified a local learning partner to contribute research expertise and lead data collection activities: Centro Universitario de Occidente de la Universidad de San Carlos de Guatemala, the National Autonomous University of Honduras, St. Paul's University in Kenya, and Datalyst Africa in Zimbabwe.

Fig 1. Mixed-methods data collection activities to understand women's water collection experiences

2.2 Community and Participant Sample Size and Eligibility, and Setting Specific Details

We aimed to engage at least six communities in each country to capture diversity in geography

and population. Communities in Guatemala, Honduras, Kenya, and Zimbabwe were purposively selected

by World Vision country offices for their engagement in World Vision SWSW programming. Almost all

communities selected had water interventions planned, but not yet completed. In Kenya, World Vision

had just completed construction of a borehole in one community, which was used by participants in that

community and from two neighboring communities. As such, participants from three of six engaged

communities in Kenya already had access to the new source.

We aimed to engage at least three women per community for a total minimum sample of

approximately 18 women per country. Samples of 9-17 participants have been deemed to be sufficient

for qualitative research(48), and we used this as a guideline per country sample size. Where resources

allowed, we sought to increase the number per community to increase variability of data captured with

the smart watches. In all settings, we recruited and enrolled a convenience sample of individuals who

were eligible to participate if they were: women, bore responsibility for going to water sources for

household needs, lived in a community selected as eligible, and were age 18 or older. Specific details

around participant recruitment varied slightly in each country, as described below.

Research in Guatemala took place in seven rural communities in the departments of

Quetzaltenango and Totonicapán, mountainous regions in the southern part of the country. In

Guatemala, 95% of the population (91% rural; 98% urban) has at least basic drinking water service (i.e.,

from an improved source requiring no more than 30 minutes roundtrip for collection)(4). Guatemala is a

multicultural, multiethnic, and multilingual country; the majority of the population (56%) is of mixed

indigenous-Spanish descent and 42% are Mayan. The research team deliberately recruited members of

the Mayan population; women from two different Mayan linguistic groups (K'iche' and Mam)

participated. Spanish is the official language of Guatemala and is spoken by 70% of the population; 8% of

the Guatemalan population speak K'iche' and 4% speak Mam(49). In Guatemala, individuals were invited

to voluntarily participate through local community authorities, with whom the research team discussed

the research. Some women were also recruited by the recommendation of other participating women.

World Vision staff did not participate in recruitment.

Research in Honduras took place in six communities across two areas—one rural and one peri-

urban—in the department of El Paraíso in eastern Honduras. The urban area is hilly and forested while

the rural area is mountainous. In Honduras, 96% of the population (91% rural; >99% urban) has at least

basic drinking water service(4). Most (90%) of the population of Honduras is of mixed indigenous and

European descent; 7% is indigenous. Spanish is the official language(50). In Honduras, the local World

Vision team introduced local community leaders (members of boards of trustees, water boards,

churches) to the local research team members who explained the research project. The local leaders

brought the research team to neighborhoods within the community, in which the research team went

door to door to explain the research and recruit eligible and interested participants. Neither World

Vision staff nor the local leaders participated in recruitment.

Research in Kenya took place in six rural communities across Isiolo and Samburu Counties. The

areas are arid and semi-arid and characterized by drought and water scarcity. In Kenya, 63% of the

population (53% rural; 86% urban) has at least a basic drinking water service(4). Kenya is also a

multicultural, multiethnic, and multilingual country. English and Kiswahili are both official languages and

there are numerous indigenous languages(51). In Kenya, the local research team approached village

leaders in the study area to explain the study and seek their help identifying potential participants. The

village leaders approached potential participants to gauge their interest in participating and relayed this

information back to the research team. The team then visited interested individuals in person to confirm

their eligibility, provide a detailed explanation of the study, and initiate the consent process. World

Vision staff did not participate in recruitment.

Research in Zimbabwe took place in six rural communities in   Gwanda and Nyanga districts. The

districts are drought-prone, and data was collected during the dry season. Gwanda has flat, agricultural

lands whereas Nyanga is forested, mountainous and has flat agricultural lands in the low-lying areas. In

Zimbabwe, 62% of the population (48% rural; 93% urban) has at least basic drinking water service(4).

Shona (80.9%), Ndebele (11.5%), and English (0.3%) are the official languages(52). Members of the

Gwanda and Nyanga World Vision offices introduced the local research team to local village leaders,

who provided lists of households from which women were selected, ensuring that they lived far from

each other to increase variability. The research team visited women to explain the research and invite

participation. Neither World Vision staff nor village leaders participated in recruitment.

Additional information regarding the ethical, cultural, and scientific considerations specific to

inclusivity in global research is included in the **S1 Checklist**.

2.3 Data Collection Tools and Procedures

Data collection took place from June-September 2023 (Guatemala: July 11-August 2; Honduras:

July 5-August 5; Kenya: June 22-July 24; Zimbabwe: July 15-July 22 and September 22-September 30.

All participants provided answers to a short survey on demographic characteristics and water

access (e.g., primary water source). Women were also asked two questions informed by the

Demographic and Health Surveys (DHS) and Multiple Indicator Cluster Surveys (MICS) to get their

estimates of water collection time: (1) "How long does it take for members of your household to go to

the drinking water source, get water, and come back?" and (2) "How long does it take for members of

your household to go to the water source for other uses, get water, and come back?"

To understand women's own perspectives about water collection and water work at sources, we

carried out go-along in-depth interviews (See 'a' in Figure 1). Go-along interviews enable the interviewer

and the participant to inhabit the space about which they are talking(53) and give participants active

roles as guides through the space(54). Because the environment can prompt discussion about what is

observed and discussed, go-along interviews can produce richer insights than sedentary interviews(55).

This method has been used to examine women's water fetching experiences and environmental risks in

an informal settlement in urban Malawi(15) and to explore experiences of water use, access, and

insecurity in Western Kenya(56). In our study, go-along interviews started in participant's homes,

continued while going to and from the water source (during pauses in walking or at the source), and

concluded back at participant's homes.

To capture additional information about experiences not gained through other methods, semi-

structured observations were carried out during the journeys (See 'b' in Figure 1). Observation guides

prompted team members to record information about, for example, the terrain, if women were

accompanied, how water was extracted and transported home, and if women engaged in other activities

on their journeys.

To determine total time, distance, elevation ascent, and caloric expenditure, participants wore

Garmin Forerunner 255 activity-tracking watches (See 'c' in Figure 1). Watches were calibrated prior to

the start of water collection using each participant's weight, height, and age to estimate caloric

expenditure. A data collection team member also wore a watch as a back-up, which was calibrated using

the participant's weight, height, and age. Watches started recording when the participant left their

home and finished upon return.

To determine the weight of items carried, Seca 876 scales were used (See 'd' in Figure 1). A data

collection team member determined the weight of any items, such as a water container or a child, that

each woman carried when she left home. The research team carried the scales to weigh anything

acquired along the way. Items were re-weighed when women left the water source before beginning the

return journey, such as water containers with water or wet laundry. If anything was difficult to place on

the scale alone (e.g., clothing), team members weighed the woman while holding the item(s) and later

subtracted her weight.

2.4 Data Collection Training

Emory team members trained the local research teams to carry out and supervise data collection. The five-day trainings provided background information about the SWSW program and guidance on ethical research conduct, principles of qualitative and quantitative research, data management, and how to use data collection tools and instruments (watches, scales, recorders). The local research teams also reviewed all tool translations to ensure consistent meaning and understanding across languages. Tools and procedures were piloted prior to data collection.

2.5 Data Collection and Management

Two trained local research team members collected data from each participant. Typically, one team member explained and provided the watch to the participant, collected demographic data, and led the go-along, in-depth interview while the other team member took notes, led the semi-structured observation, and recorded data on all weighed items. Working in teams of two lessened the work burden and provided team members with a greater sense of security, particularly when journeying far in new locales.

Go-along, in-depth interview, observation, and demographic survey tools were translated into Spanish (Honduras and Guatemala), Samburu (Kenya), and Ndebele and Shona (Zimbabwe). Data collection team members in Guatemala who were native K'iche' or Mam speakers verbally translated tools from Spanish into K'iche' and Mam, as these languages are not typically written.

Digital voice recorders were used to record go-along interviews. Audio recordings in K'iche', Mam, and Spanish were first transcribed into Spanish by trained members of the local research teams and then translated into English by a hired translator. Audio recordings in Samburu, Ndebele, and Shona were simultaneously transcribed and translated into English by trained team members.

Demographic and observation data were recorded on paper forms. Trained team members in

Kenya and Zimbabwe typed up observation notes in English. In Guatemala and Honduras, observation

notes were typed up in Spanish and then translated into English by a hired translator. Demographic and

select numeric or short answer data from observation sheets, which included weight data, were double

entered into a database (by different members of the Emory team) and then compared for accuracy.

Discordant data were checked against scanned files and corrected.

Data from the watches were synced to the 'Garmin Connect' web platform and renamed with

participant IDs. Data were then downloaded and exported from the Garmin web platform as CSV and

TCX files and uploaded onto a secure OneDrive folder.

All data, including recordings and scanned documents (e.g., surveys, notes), were de-identified,

uploaded to secure OneDrive folders, and stored using unique identifiers that linked all data for each

participant.

2.6 Data Analysis

To learn about women's practices, experiences, and perceptions of water collection, and the

environments and factors that influence them, (aim 1; activities 'a' and 'b' in Figure 1), we conducted a

rapid thematic analysis of all transcripts from go-along interviews and observation notes. Using Excel to

organize data, we created a matrix of key themes of interest identified *a priori* (e.g., methods of

collection and carriage, terrain, perceived risks). Each transcript and all observation notes were read

line-by-line, and relevant information was extracted, organized by the community, and then summarized

into profiles for each community based on individual data. Community profiles were further summarized

into country profiles.

To determine actual water collection time, distance, caloric expenditure, total elevation ascent,

and water volume and weight carried (aim 2; activities 'c' and 'd' in Figure 1), we calculated descriptive

statistics of watch and weight data. The volume of water collected was derived from the weight data;

the weight of containers was subtracted from the total weight of the water-filled containers to

determine weight of the water alone. The weight of water was then converted to volume (1kg=1L). We

generated graphs to visualize trends related to participant journeys, with attention to time, distance,

caloric expenditure, water volume, and the source visited (improved or unimproved).

Time, distance, total elevation ascent, and caloric expenditure data were manipulated for select

participants in each country based on identified data collection challenges, enabling us to retain and use

data collected. Specifically, in Honduras, one water journey was only recorded one-way. For this

observation, time and distance were doubled to estimate a round-trip. In Kenya, data for two water

journeys are the mean of the participant and enumerator watches as these were not properly

distinguished during data collection (individual observations were similar). Watch data from data

collectors were used as proxies for two participants (Guatemala, Zimbabwe) because participant water

journeys were accidentally not recorded.

To determine if women's estimated and actual water collection times were aligned (aim 3), we

calculated descriptive statistics (mean, range, SD) of the watch-measured times and women's estimates

of water collection time, specifically among the subset that collected water. We calculated the

difference between estimated and actual times and identified the number and proportion of

observations by country and type (drinking, other uses) that were aligned and that were discordant. We

leveraged the go-along interview and observation data to determine and classify women's intended use

of the water they were collecting. Water was classified as for 'drinking' if a participant indicated that any

of the water would be used for drinking, even if they also would use some of the water for other

purposes. Water was classified as for 'other uses' (e.g., bathing, washing dishes) if women did not

explicitly mention using any of it for drinking. (We acknowledge that the WHO/UNICEF Joint Monitoring

Programme for Water Supply, Sanitation, and Hygiene (JMP) considers 'drinking water' to include

'drinking, cooking, personal hygiene and other domestic uses.(57) However, we separated as described

because we expect participants responded to what was asked of them, not to otherwise determined

categories (See section 2.3)).

Quantitative analyses were done in Stata (version 18.0), Microsoft Excel, and R (v4.2.1);

qualitative analyses were facilitated using Microsoft Excel.

The study protocol was reviewed by the Institutional Review Board committee of Emory

University, Atlanta, Georgia, USA, which determined that the research was exempt from further review

and approval (IRB00005955). It was approved by the Comité de Ética del Centro Universitario de

Occidente, de la Universidad de San Carlos de Guatemala in Guatemala (Acta 1.23 C.E. DICUNOC);

Universidad Nacional Autónoma de Honduras (CEIFCS-2023-P18) in Honduras; St. Paul's University -

Institutional Scientific Ethics Committee (ERB No. 38), the National Commission for Science, Technology

and Innovation (NACOSTI/P/23/27117) in Kenya; and the Medical Research Council of Zimbabwe

(MRCZ/A/3054) in Zimbabwe. All participants provided informed consent. In Guatemala, Honduras, and

Kenya, participants consented verbally, and enumerators signed the consent form to affirm. In

Zimbabwe, participants provided signed consents.

3 RESULTS

3.1 Participant Information

Ninety-four women from 25 communities in four countries participated in data collection (**Table**

1). For 93 women, a single journey to a single water source was recorded. One woman in Guatemala

participated in two journeys to capture that participant's different water sources.

Across all settings, the mean participant age was 37.1 years (SD = 13.7). The largest proportions

of participants in Guatemala (12; 55%) and Honduras (11; 65%) were unmarried with a partner or single;

the majority in Kenya (17; 77%) and Zimbabwe (23; 70%) were married. The average household size was

5.8 members (SD = 2.9). The greatest proportion to never attend school was in Kenya (20; 91%) and the

greatest proportion to engage in economic activities was in Guatemala (20; 91%).

331 *3.2 Reported and Observed Water Collection Practices and*

332 *Experiences*

Sources. Based on the demographic survey, women reported a variety of sources from which

they would primarily collect water for drinking and 'other uses.' Overall, most participants, including all

from Kenya and Zimbabwe, reported that their primary drinking water source was located outside of

their dwelling, yard, or plot (83; 89%) and almost half (42; 46%) indicated that they collected drinking

water from a source considered unimproved. However, there was variation between countries. Over

half of the participants in Guatemala (15; 71%), Honduras (11; 65%), and Zimbabwe (19; 58%) reported

collecting drinking water from a source considered improved, compared to just 24% (5) in Kenya.

Surface water was the most reported unimproved drinking water source in Guatemala (5; 24%),

Honduras (4; 24%) and Zimbabwe (13; 39%). In Kenya, most participants reported that they typically

collect drinking water from unprotected dug wells (15; 71%). On average, women reported collecting

water almost daily (5.6 days/week); though the mean number of reported collection days was lower in

Honduras compared to the other countries (Guatemala 6.4 days; Honduras 4.1 days; Kenya 5.7 days;

Zimbabwe 5.9 days).

While all women in Kenya reported that they would use the same water sources for drinking and

'other uses', sizable proportions of women in Guatemala (91%), Honduras (41%), and Zimbabwe (46%)

said they visited different sources for drinking and 'other uses.' The proportions of women using surface

water for 'other uses' were greater than the proportions using it for drinking in both Guatemala (+21%)

and Zimbabwe (+23%). No women reported using rainwater for drinking, but approximately one third

reported using it for 'other uses' in both Guatemala (6; 33%) and Honduras (5; 29%).

On the days they were followed, women visited improved and unimproved sources. In each of

the three communities in Kenya where participants had access to a recently installed borehole, water

was not always available. In each of these communities, two participants went to the new borehole and

two went to an existing unprotected source. Participants also described other sources they would

frequent depending on need or season. For example, participants from all countries reported collecting

rainwater during the rainy season. Women from some communities discussed how they sometimes got

water from family or neighbors (Honduras, Guatemala) or by paying someone, typically men (Zimbabwe).

Table 2 provides thematic summaries of the qualitative data by country; **S1 Table** provides country-

specific summaries by community.

Activities at the source. The primary reason for most (83; 87%) participants to visit water sources

was to collect water. Among those, four women (2 Guatemala, 2 Kenya) also brought clothing to wash

(**Table 3**). Ten women (9 Guatemala, 1 Kenya) did not collect water; they visited sources to wash clothes,

which they brought with them.

Table 1. Demographic and water characteristics of water journey participants, by country

	Country									
	Guatemala (n=22; 7 communities)		Honduras (n=17; 6 communities)		Kenya (n=22; 6 communities)		Zimbabwe (n=33; 6 communities)		Total (n=94; 25 communities)	
Age (mean, SD)	36.2	15.4	38.9	14.5	30.2	10.1	41.3	12.8	37.1	13.7
Household size[1] (mean, SD)	6.5	3.5	4.2	2.2	6	2.3	5.9	2.8	5.8	2.9
Marital status (n, %)										
Single, never married	3	13.6	4	23.5	1	4.5	1	3.0	9	9.6
Unmarried, but have partner	9	40.9	7	41.2	1	4.5	0	0.0	17	18.1
Married	8	36.4	3	17.6	17	77.3	23	69.7	51	54.3
Separated	2	9.1	1	5.9	1	4.5	3	9.1	7	7.4
Divorced	0	0.0	0	0.0	0	0.0	2	6.1	2	2.1
Widowed	0	0.0	2	11.8	2	9.1	4	12.1	8	8.5
Completed schooling[2] (n, %)										
Never attended school	4	18.2	4	25.0	20	90.9	0	0.0	28	30.4
Some primary	9	40.9	6	37.5	0	0.0	8	25.0	23	25.0
Primary	7	31.8	5	31.3	1	4.5	11	34.4	24	26.1
Secondary	2	9.1	1	6.3	0	0.0	13	40.6	16	17.4
Above secondary	0	0.0	0	0.0	1	4.5	0	0.0	1	1.1
Engaged in economic activities (n, %)	20	90.9	6	35.3	6	27.3	25	75.8	57	60.6
Primary source of drinking water[3] (n, %)										
Improved Sources	*15*	*71.4*	*11*	*64.7*	*5*	*23.8*	*19*	*57.6*	*50*	*54.3*
Piped water	8	38.1	7	41.2	0	0.0	4	12.1	19	20.7
Tube well/borehole	0	0.0	0	0.0	3	14.3	14	42.4	17	18.5
Protected dug well	7	33.3	1	5.9	2	9.5	0	0.0	10	10.9
Protected spring	0	0.0	0	0.0	0	0.0	1	3.0	1	1.1
Bottled water	0	0.0	3	17.7	0	0.0	0	0.0	3	3.3
Unimproved Sources	*6*	*28.6*	*6*	*35.3*	*16*	*76.2*	*14*	*42.4*	*42*	*45.6*
Unprotected dug well	0	0.0	2	11.8	15	71.4	1	3.0	18	19.6
Unprotected spring	1	4.8	0	0.0	1	4.8	0	0.0	2	2.2
Surface water	5	23.8	4	23.5	0	0.0	13	39.4	22	23.9
Other (unspecified)	1	4.5	0	0.0	0	0.0	0	0.0	1	1.1
Location of drinking water source[4] (n, %)										
In dwelling	3	13.6	1	5.9	0	0.0	0	0.0	4	4.3
In yard/plot	4	18.2	2	11.8	0	0.0	0	0.0	6	6.5
Elsewhere (beyond yard/plot)	15	68.2	14	82.4	21	100.0	33	100.0	83	89.2
Days /week needed to collect water[5]										
Mean, SD	6.4	1.5	4.1	2.5	5.7	1.9	5.9	2.0	5.6	2.1
Range		(2-7)		(1-7)		(2-7)		(1-7)		(1-7)
Primary source of water for other uses[6] (n, %)										
Improved Sources	*10*	*55.6*	*12*	*70.6*	*5*	*23.8*	*10*	*31.2*	*37*	*42.1*
Piped water	2	11.1	6	35.3	0	0.0	1	3.1	9	10.2
Tube well/borehole	0	0.0	0	0.0	3	14.3	9	28.1	12	13.6
Protected dug well	2	11.1	1	5.9	2	9.5	0	0.0	5	5.7
Rainwater	6	33.3	5	29.4	0	0.0	0	0.0	11	12.5
Unimproved Sources	*8*	*44.4*	*5*	*29.4*	*16*	*76.2*	*22*	*68.8*	*51*	*58.0*

	n	%	n	%	n	%	n	%	n	%
Unprotected dug well	0	0.0	2	11.8	15	71.4	2	6.3	19	21.6
Unprotected spring	0	0.0	0	0.0	1	4.8	0	0.0	1	1.1
Surface water	8	44.4	3	17.6	0	0.0	20	62.5	31	35.2
Other (unspecified)	4	18.2	0	0.0	0	0.0	1	3.0	5	5.4

Respondents with source for 'other uses' that is different than primary drinking water source[7]

	n	%	n	%	n	%	n	%	n	%
(n, %)	20	90.9	7	41.2	0	0.0	15	45.5	42	45.2

Household size: 1 missing (Zimbabwe)

Completed schooling: 1 missing (Honduras); 1 missing (Zimbabwe)

Primary source of drinking water: 1 missing (Kenya); Sources noted as 'other' were not classified as either improved or unimproved and were excluded from percentage totals

Location of primary drinking water source: 1 missing (Kenya)

Days in a week needed to collect water: 4 do not know (Guatemala); 1 missing (Kenya)

Primary source of water for other uses: 1 missing (Kenya); Sources noted as 'other' were not classified as either improved or unimproved and were excluded from percentage totals

Different primary source types for drinking and 'other uses': 1 missing (Kenya)

The methods women used to extract water varied depending on the source. In Guatemala

women used buckets or 'tinajas' (clay pots) to pull water from a well or filled containers by dipping them

into rivers. Some women in both Guatemala and Honduras used hoses to fill containers. Other methods

for extracting water were more arduous. Women in both Honduras and Zimbabwe used hand pumps,

which women said required strength or as noted in Zimbabwe, assistance from other women also

present at the sources. In Kenya and Zimbabwe, women dug for water at riverbeds. The process

required them to dig deep enough to reach water, wait for water to pool, and then remove it with small

vessels until it ran clear enough to be deemed suitable for collection. Women in Kenya said they

sometimes needed to dig so deep during the dry season that they would be fully inside the holes.

In addition to water collection or washing clothing, many women carry out other activities at or

on the way to and/or from water points. For example, at least one woman in each country was

simultaneously caring for a child, carrying them on their back or front; in Guatemala and Honduras, at

least one woman in each community reported that children regularly accompany them to water sources.

Women noted older children occasionally accompany them and help carry water if strong enough. While

at water points women also washed the clothing they were wearing (Guatemala, Kenya, Zimbabwe) and

eating utensils (Guatemala); bathed themselves or children (Honduras, Kenya); and tended to and

watered small vegetable gardens near the source (Zimbabwe). While journeying to or from water points,

women discussed or were observed collecting firewood (Honduras, Zimbabwe, Kenya), going to the

store (Honduras), harvesting mopane worms (Zimbabwe), and watering goats (Kenya), all of which can

lengthen the journey time.

Methods of carriage. Most women transported filled water containers without assistance

(carried or rolled: 97%), while a few used assistance (wheelbarrow or vehicle: 3%). Women in Honduras,

Guatemala, and Zimbabwe carried water on their heads and women in Kenya carried on their back,

often assisted by a rope on their head, or by dragging or rolling the water container behind them.

Carrying strategies depended on the size of the vessel and what else was being carried, like children. In

Kenya, women said they once had donkeys to help carry water, but many donkeys had died due to the

extended drought. One woman followed in Guatemala paid to travel by bus part of the way. Some in

Zimbabwe used carts or wheelbarrows to collect larger quantities. One Zimbabwean woman described

the work of using wheelbarrows: "When I am using a wheelbarrow, I make sure that I rest at least two

times on my way back home from the dam, because it is very hard to push a wheelbarrow along the way.

Plus, I will be carrying three buckets of water that are heavy. When carrying on my head, I rest for about

two   to   three   times   on   my   way   back."

Table 2a. Thematic descriptions of women's water collection experiences and practices, by country

Country & Theme	Guatemala (7 communities, 22 Participants)	Honduras (6 communities, 17 Participants)	Kenya (6 communities, 21 Participants)	Zimbabwe (6 communities, 33 Participants)
Water Sources Visited During Data Collection	Across communities, participants went to a variety of different sources, including rivers, wells, springs, and public tanks.	Across communities, participants went to a variety of different sources, including springs, wells, public and private taps, and rivers.	Across all communities, women collected water from an unprotected well, which they would dig in the riverbed. Some women in 3 communities (4, 5, & 6) collected from a new borehole. Women in communities 1 & 4 reported some water pooling in rocks at the top of a hill as another source.	Across 5 of 6 communities (1, 2, 3, 5, & 6), women visited unprotected dug wells at water sources. In 2 communities (1 & 3), participants visited dams. In 4 communities (3, 4, 5, & 6), women also visited improved water sources such as taps (4 & 6) or boreholes (3, 4, 5, & 6).
Other Water Sources Used by Participants, But Not During Data Collection	Women said rivers or springs are available in all communities. In 4 communities (2, 3, 4, & 6), women reported that some of their drinking water was provided for them for free by family members or neighbors who had household access to water. In 5 communities (3, 4, 5, 6, & 7), there is a community water source such as a tank or a well. In all but 1 community (6), women discussed rainwater collection during the winter rainy season.	Women in all communities reported accessing a natural water source, such as a river or spring. Rainwater harvesting was reported in 3 communities (1, 3, & 5) to collect more water during the winter months. In 3 communities (2, 3, & 5), women report that they may access drinking water from family members or neighbors with household water connections.	Women reported using alternative water sources during the rainy season, specifically surface water collection (e.g., small dams, gullies, small streams). Women in community 2 noted collecting rainwater, and women in community 1 described using an unprotected spring. Most women in communities with borehole access reported that they would access water via dug well at the riverbed for some purposes. In one community (4) a new borehole was noted as an alternative source that was a great help, but the participant chose to collect water at the riverbed for an undisclosed reason.	Participants in 2 communities (3, 4 & 6) reported access to improved water sources: a borehole (in 4 & 6) and a tap at a school (3). A participant in community 3 chose to collect water from the dam because the water in the borehole was running dry. Women in 5 of 6 communities (1, 2, 3, 5, & 6) reported using additional water sources in rainy season, such as rainwater harvesting from rooftops (1, 3, 5, & 6), or surface waters such as streams or "paddles" (1, 2, 3, 5, 6). One participant in community 3 occasionally paid someone for water.
Method of collection	Limited information was recorded about methods of collection at the source. In community 1, the woman who was collecting water (others were washing clothing), drew water from a well using a bucket and in community 2, a woman filled her containers directly from the river. In community 3, two participants extracted water from the source using a hose. (In communities 4, 5, & 6, the methods of collection were not recorded.)	Limited information was recorded about methods of collection at the source. In 2 communities (2 & 3), water was piped from a source and women collected water through the hose. In community 6, one participant had to manually pump to get water.	Across all communities, women who collected water at the riverbed dug the well at the river, scooping out and throwing away the brown, sandy water initially seeping in. They then collected the clearer water after the well re-filled. In community 6, women said that three people needed to get inside the dug well to get the water to clarify the depth of the hole. In 2 communities (1 & 2), women described using water pans to collect rainwater in the rainy season.	In 5 of 6 communities (1, 2, 3, 5, & 6), women dug a hole in or near a riverbed (or used a hole that had been dug by others) to collect water. In 3 communities (3, 5, & 6), participants that used the borehole reported needing the help of others to pump water. (In community 4, the method of water collection was not noted.)

Community names are not shared. We assigned ID numbers for readers to see where information is from across themes.

400   **Table 2b. Thematic descriptions of women's water collection experiences and practices, by country** *(continued)*

Country & Theme	Guatemala (7 communities, 22 Participants)	Honduras (6 communities, 17 Participants)	Kenya (6 communities, 21 Participants)	Zimbabwe (6 communities, 33 Participants)
Number and timing of trips	Participants reported collecting water between 1 and 20 times a day, depending on how much water was needed and the availability of water at the source. In all communities, women reported leaving the house to fetch water less often in the rainy season. In community 4, one participant was careful to only collect once a day because she pays to travel to a water source.	The number of times water is collected varies across participants, communities, and seasons, from water collection every two weeks (community 6) to water collection over 10 times in a day (community 4). Women in 5 communities (1,2,3,4,6) reported going more often in the summer dry season than the winter rainy season. In one community (5), a participant reported going half as often in the summer because there is less water available to collect. When women go during the day depends on weather, needs, and other responsibilities. In 3 communities (4,5,6), women said they time their water collection to avoid the heat of the sun. In 1 community (3), a participant reported avoiding the water source in the afternoon because of mosquitoes.	Most participants who go to the riverbed go only once a day. Participants in 3 communities (3,5,6) noted only going once per day because of the distance. Those in 3 different communities (1,2,4) said they may go twice to the riverbed if needed, but those in community 4 clarified that it is not possible to go twice in the dry season. Women in two communities (5,6) said they sometimes go to the borehole twice in a day, but said going twice is not always possible because the borehole is not open every day, there are long lines, and the hours are limited. Women in two communities (2,5) said they may go more often in the rainy season to smaller, closer sources. Women across all communities discussed collecting water in the morning because of the heat in the afternoon and, as those in community 5 noted, it could be dark by the time they get home if they start later in the day.	The number of times women would collect water per day varied. In 2 communities (1,2), women who got water via sand abstraction reported collecting 1-3 times a day; one woman collected up to 5 times. Participants in 3 communities (4,5,6) collected water from dug wells 2-3 times per day and made trips to a borehole or tap. The number of trips depended on the quantity of water they could collect per trip, which would increase if they had use of a wheelbarrow or scotch cart (1,2,3,6). In 4 communities (1,2,5,6), women said they decreased the number of trips in the rainy season because they had closer sources then. In 4 communities, women said they would collect early in the morning to avoid long queues at the river (1 & 6) or borehole (3,4). Weather (2), borehole hours (3), events (e.g., funerals) (3), and visitors, who participants did not want to leave at their house (6) also influenced timing.
Method of carriage	A variety of tactics for carrying water were observed or described. In 3 communities (4,5,6), at least one participant reported transporting water or laundry in a vehicle, for which participants occasionally paid (4,6). In 2 communities (2,4), women balanced water containers ('tinajas') on their heads, cushioned by a rolled-up blanket ('yagual'). A participant from community 1 noted that her husband would help her carry water, and a participant in community 5 said that if she needs to carry more than one container, she ties them together with a rope and pulls them. A participant from community 1 who did laundry carried the laundry in a large sheet.	Women across all communities carried water in buckets by their handles. In 2 communities (2,4), some women carried water on their heads.	Women across all communities collected water in jerricans that hold 20 liters, 10 liters, or, in 2 communities (4,5), 5 liters of water. Women would carry these on their backs or drag or roll them behind them. Decisions around which jerrican to use were based on women's strength or exhaustion, and if children were being carried. If older children were accompanying their mothers, they might assist by carrying small (5L) jerricans, or by rolling a jerrican. In 3 communities (1,2,4), women said donkeys used to carry water, but that they died in the drought.	Participants in all communities carry water in buckets on their heads. Some participants in 4 communities (1,2,3,6) reported occasionally having access to wheelbarrows or scotch carts to facilitate water carriage, which enabled them to collect more water.

Community names are not shared. We assigned ID numbers for readers to see where information is from across themes.

401   **Table 2c. Thematic descriptions of women's water collection experiences and practices, by country** *(continued)*

Country & Theme	Guatemala (7 communities, 22 Participants)	Honduras (6 communities, 17 Participants)	Kenya (6 communities, 21 Participants)	Zimbabwe (6 communities, 33 Participants)
Terrain and risks	In 6 of 7 communities (1,2,3,4,6,7), women described their path to water sources as steep and slippery, especially during the rainy season. Women in 3 communities (1,2,7), reported that either they or a family member had slipped and fallen while collecting water. In 2 communities (2,6), women stated that in the rain, rivers would overflow and become dangerous. In 3 communities (1,2,3), participants said that there were snakes on the pathway to the water source; dogs and people were also noted risks (5).	Women in all communities reported routes to the water source to be slippery or become more treacherous in the rain. Participants in 2 communities (2,3) reported snakes along the path and those in another (3) reported coyotes and mosquitoes.	The terrain was mostly flat in several communities but was hilly in 2 (4,6). Women across all communities traversed gullies on the way to and from water collection points, which could pose challenges. Women from all communities also reported wild elephants to be a risk during water collection, and some also reported risks from leopards and other people.	In all communities, the routes to the water sources are hilly or mountainous, with loose stones. In 5 of 6 communities (1,3,4,5,6), there are sandy stretches that can be hot and difficult to walk on while carrying water or pushing wheelbarrows. In 4 communities (2,3,5,6), there are prickly or thorny trees and bushes on the pathway. Participants in 2 communities (4,5) said motorbikes go by quickly making it dangerous. In 3 communities (2,3,6), women reported there to be dangerous people, including those who sexually harassed or assaulted women (2,6) or children (3), or traditional healers performing rituals at the dam (3). Participants in all communities reported risks from animals (e.g., snakes (1,3,6), hyenas or jackals (2,6), baboons (5)).
Physical effects of water collection	In 2 communities (1,3), women reported discomfort and pain in in various parts of the body from water collection. A participant in community 4, reported that she has a hard time carrying heavy things, including water, since a caesarian section. A pregnant participant in community 6 reported that she had to reduce the amount of water she carried and now needs help from her husband. No data were reported from the remaining 3 communities.	Participants across communities reported fatigue, exhaustion, and physical discomfort of some type related to water collection. In one community (4), a participant reported having fallen while going to the source, and in another (6), a participant had sprained an ankle.	Women in all communities mentioned exhaustion and pain in the body. In one community (2), a participant reported possible miscarriages because of water collection. Women in community 6 reported having more energy because they accessed water from the new borehole, which made their water collection easier.	Across communities, women reported numbness, exhaustion, and pain in the neck, back, chest, and other body parts. A participant in community 1 noted that the journey is harder on her physically when she uses a wheelbarrow because it is heavy and hard to push. In communities with boreholes, one participant (3) noted that pumping at the borehole requires a lot of effort and energy and another reported shoulder pain from the pumping (6).

Community names are not shared. We assigned ID numbers for readers to see where information is from across themes.

404   **Table 2d. Thematic descriptions of women's water collection experiences and practices, by country** *(continued)*

Country & Theme	Guatemala (7 communities, 22 Participants)	Honduras (6 communities, 17 Participants)	Kenya (6 communities, 21 Participants)	Zimbabwe (6 communities, 33 Participants)

Children	Across all communities, some participants carried children when collecting water and some reported that older children may assist with water collection.	Across all communities, women reported being accompanied by children. In 2 communities (1,2), participants were observed with children. In one community (2), one participant carried a toddler when she did not want to walk, and another had two children with her (the older child carried the younger child). Participants in 3 communities (1,2, 3) reported that children may help with carrying if they are old enough.	Women across all communities reported that children occasionally collected water to help their mothers but are not the primary water collectors. Children are often left at home to be watched by other women (4,5,6) or older siblings (2,3) while the mother collects water. In only one community (5) was a participant observed to carry a child on a water journey. In 3 communities (2,4,5), mothers with small children reported that they sometimes carry them while collecting water. One woman (community 2) said they carry water on their backs and babies in front.	In 4 of 6 communities (1 2,5,6), at least one participant was observed carrying a child while collecting water; women in all communities said they sometimes carried a child while collecting water. In 5 communities (1-4,6), older children were reported to help with water collection or occasionally collected by themselves. Children under 18 were reported to not be allowed to collect water at the river (community 3). One pregnant participant (community 5) with a small child had someone mind the child while she collected water.
Other work done during water collection	Women across all communities reported washing clothing at water points. For many participants, trips to water points for washing clothes were said to be separate from trips for water collection. Women from 4 communities (2,4,5,6) said they would wash utensils or dishes at sources as well; other women carried water home to wash dishes or utensils there.	Participants across all communities reported washing clothing at the water point. Participants in 3 communities (1,2,4) also reported collecting firewood on their way to or from water points. Participants in 5 communities (1-5) commonly bathed themselves or children at water sources. In one community (6), a participant also reported going to the grocery store on her way to the water point.	Women across all communities reported collecting firewood on the way to or from water sources and washing clothes while at water collection points. Women in 4 communities (2,3,4,5) also took goats to get water at collection sites.	Women reported participating in various activities when collecting water. In all communities, women reported collecting firewood on the journey (except community 6) and washed clothes at the water source. In 4 communities (1,2,3,6), participants reported picking fruit when in season. Some women also watered gardens by the water source (5,6) and one woman said she harvested mopane worms (2).
Perceived water quality and water treatment	Perceptions of water quality varied across communities and by sources. Women in some communities discussed treating water by boiling (2,4) or filtering (4,5). Participants in all communities reported work to maintain water sources to preserve the cleanliness of the water (e.g., cleaning the inside of wells (1,2,3) or tanks (6,7), or cleaning mud and debris out of areas where spring or surface water is collected (4,5,6, 7).	Perceptions of water quality varied widely between communities and by sources. Women in five communities (1,3,4,5,6) said that river or stream water was contaminated. In three communities (1,2,4), women reported that the river gets dirty in the rain. One woman (3) perceived water from her primary well to be safe for consumption. Participants were not asked about treating water.	Perceptions of water quality varied by community and source. Women in 3 communities (2,3,4) perceived dam water to be the most contaminated because the dams are used by animals. Women in all communities poured out the initial water from dug wells and then collected clearer water eventually seeping through. Women in 5 of 6 communities (1,2,4,5,6) reported that they had at some point treated water with chlorine but do not regularly or currently. A woman in community 3 said she would boil water if children had diarrhea.	Across communities, perceptions and practices related to water quality varied. Dam water was perceived to be unsafe because animals drink the water (1,3). In 3 communities (2,3,6), participants reported treating water (sieves, tablets, or boiling) though some said boiling and tablets negatively affected the taste of water (6). Water safety techniques (e.g., putting lids on buckets, using one utensil to extract drinking water, separating drinking from other water) were reported (5,6). Women typically considered the borehole water to be clean, though one woman (5) reported borehole water to be rusty. Neither water quality nor treatment activities were mentioned in community 4.

Community names are not shared. We assigned ID numbers for readers to see where information is from across themes.

Frequency of collection. Women may go to different sources throughout the course of the day or

week and the season. Factors such as the weather, family needs, the volume they can carry, other

responsibilities (e.g., childcare), available support for carrying water (e.g., other people, wheelbarrow),

queues at the water source, and distance to sources influence frequency and source selected. In

Honduras and Guatemala, the frequency ranges were the widest. In Honduras, one participant said she

collected water once every two weeks, economizing the water to make it last; another said she would

go at least ten times per day in the summer. In Guatemala, one participant collected water once per day

because she paid for transport to carry the heavy loads; another said she would go up to 20 times in a

day if she needed to collect water for bathing. In Zimbabwe, most women reported going to (often

varied) water sources several times a day, though the number of trips could depend on their access to a

scotch cart (a wooden cart typically pulled by donkeys) or wheelbarrow to transport greater volumes of

water. In Kenya, women typically collected water only once per day because of the heat and the

distance to water sources. Some women in Kenya would go more than once, but not to riverbeds in the

dry season. Those with borehole access would sometimes go twice, but long lines and limited operating

hours were barriers.

421   *3.3 The Burden of Collection*

The time and distance traveled, the elevation gained, the terrain and risks encountered, the

energy expended, the weights carried, and the volumes of water collected varied within and across

countries.

Time. Across countries, the mean water journey time—including going to the source, completing

activities at the source (e.g., extracting water, washing clothes), and returning was 82min (SD = 73.3)

(**Table 3**). Mean times were shortest in Honduras (38min), comparable in Zimbabwe (58min) and

Guatemala (68min), and markedly greater in Kenya (164min). Within each country, the shortest journeys

were less than 20 minutes (Guatemala: 13min; Honduras: 18min; Kenya: 19min; Zimbabwe: 11min). In

three countries, the longest journeys lasted over three hours, and over 4.5 hours in two (Guatemala:

195min; Honduras: 100min; Kenya: 287min; Zimbabwe: 282min).

Distance. The mean distance for all water journeys was 3.5km (SD = 3.9) (**Table 3**). Mean

distances were shortest in Guatemala (1.1km) and Honduras (1.1km), greater in Zimbabwe (2.8km), and

greatest in Kenya (8.5km). Within each country, the shortest journeys were all less than 1km

(Guatemala: 0.2km; Honduras: 0.2km; Kenya: 0.7km; Zimbabwe: 0.2km) and the farthest were all over

2km, with the longest in Kenya (Guatemala: 3.7km; Honduras: 2.7km; Kenya: 15.8km; Zimbabwe: 8.7km).

As **Figure 2** shows, those walking the greatest distances and taking the greatest amount of time to

collect water were largely doing so to get water from unimproved sources.

Fig 2. Time, distance, and (a) caloric expenditure, (b) volume of water collected, and (c) source

type among participants followed in Guatemala, Honduras, Kenya, and Zimbabwe. The graphs

show a woman from Zimbabwe who spent more time, went farther, expended more calories

and collected more water than other women in Zimbabwe. She tended a garden by the source;

her work hauling water to it resulted in these increased, outlier values.)

Total elevation ascent. The overall mean total elevation ascent across countries was 57m (SD =

49.8), with the smallest mean values in Zimbabwe (40m) and Honduras (44m), larger values in

Guatemala (57m), and the largest in Kenya (89m) (**Table 3**). For each country, the smallest total

elevation ascent values were all under 15m (Guatemala: 6m; Honduras: 12m; Kenya: 7m; Zimbabwe:

1m). Three of the largest total elevation ascent values were at or over 200m (Guatemala: 299m;

Honduras: 74m; Kenya: 214m; Zimbabwe: 200m).

Environmental risks. Women traversed challenging terrain and encountered risks while going to

water points. Paths were particularly challenging in Guatemala and Zimbabwe. Women in Guatemala

went down steep and fragile hillsides. In Zimbabwe, women described hilly and mountainous terrain

with loose stones and portions with sand that could burn feet and were hard to walk on or push

wheelbarrows through, particularly when carrying heavy loads. The rains made terrain treacherous

(Honduras) or even more challenging to traverse (Guatemala) because of overflowing rivers and slippery

slopes. Women in Guatemala, Kenya, and Zimbabwe said other people could pose risks. In Zimbabwe,

women were concerned about fast motorbikes and experiencing sexual harassment and assault.

Women reported animals to be risks in Guatemala (snakes, dogs), Honduras (snakes, coyotes), Kenya

(elephants, leopards) and Zimbabwe (bees, snakes, hyenas, jackals, and baboons). Women navigated

challenging terrain and animal encounters while carrying cumbersome loads:

"There are many stones, big and small, along the way, making it difficult to walk even when

carrying an empty bucket. I once fell and broke my bucket that was full of water. The path is also

infested with snakes as they come to seek shelter from the thorny trees." (Zimbabwe)

Caloric expenditure. The overall mean caloric expenditure was 231cal (SD = 193.5) (See **Table 3**).

Mean expenditure was the least in Guatemala (118cal) and Honduras (144cal), greater in Zimbabwe

(223cal) and greatest in Kenya (421cal). Within each country, the smallest caloric expenditure values

were all under 100cal (Guatemala: 36cal; Honduras: 42cal; Kenya: 65cal; Zimbabwe: 37cal) and the

greatest were all over 350cal, (Guatemala: 393cal; Honduras: 354cal; Kenya: 736cal; Zimbabwe: 952cal).

There is a trend between the water journey time, distance, and calories expended. Most participants

from Kenya spent more time, traveled farther, and expended more calories than all other participants

(**Figure   2**).

Table 3. Water journey activity, source, time, distance, caloric expenditure, and elevation ascent, by country

	Country									
	Guatemala (n=22; 7 communities)		Honduras (n=17; 6 communities)		Kenya (n=22; 6 communities)		Zimbabwe (n=33; 6 communities)		Total (n=94; 25 communities)	
Number of water journeys[1]		23		17		22		33		95
Primary water journey activity[1] (n, %)										
Water collection	12	52.2	17	100.0	19	86.36	33	100.0	81	85.26
Washing clothes/ laundry	9	39.1	0	0.0	1	4.55	0	0.0	10	10.53
Water collection & washing clothes	2	8.7	0	0.0	2	9.09	0	0.0	4	4.21
Time in minutes[1-3]										
n		22		16		22		33		93
Mean, SD	67.7	52.1	38.1	19.9	164.4	87.3	57.7	45.7	81.9	73.3
Median		58.5		31.7		189.3		47.6		53.4
Range (min, max)		(13.3 - 194.5)		(18.3 - 99.5)		(19.2 - 287.2)		(10.8 - 282.0)		(10.8 - 287.2)
Difference (shortest & longest)		181.2		81.2		268.0		271.2		276.4
Distance (km)[1-3]										
n		22		16		22		33		93
Mean, SD	1.1	0.8	1.1	0.7	8.5	4.8	2.8	1.8	3.5	3.9
Median		0.9		0.9		9.0		2.4		1.9
Range (min, max)		(0.2 - 3.7)		(0.2 - 2.7)		(0.7 - 15.8)		(0.2 - 8.7)		(0.2 - 15.8)
Difference (shortest & longest)		3.5		2.5		15.1		8.5		15.6
Caloric expenditure (cal)[1-3]										
n	22		16		22		33		93	
Mean, SD	117.9	87.6	143.9	94.2	421.3	218.8	223	171.6	231.4	193.5
Median		97.5		103.5		489.5		175.0		161.0
Range (min, max)		(36 - 393)		(42 - 354)		(65 - 736)		(37 - 952)		(36 - 952)
Difference (least & greatest)		357		312		671		915		916
Total elevation ascent (m)[1,2,4]										
n	22		15		22		33		92	
Mean, SD	57.0	61.7	44.0	18.8	89.2	56.9	40.0	34.4	56.5	49.8
Median		35.5		45.0		86.4		36.0		42.0
Range (min, max)		(6.1 - 299.0)		(12.0 - 74.0)		(7.0 - 214.9)		(1.0 - 200.0)		(1.0 - 299.0)
Difference (least & greatest)		292.9		62.0		207.9		199.0		298.0

[1] 1 participant completed two separate water journeys to different water sources (Guatemala)

[2] 1 participant was excluded from aggregate analyses for time, distance, caloric expenditure, and ascent because she took a bus for part of her water journey and was only on foot for part of the time. Exclusion ensures that the values noted here reflect journeys taken completely on foot. Her specific journey lasted 32.6 minutes, covered 5.4 km, burned 103 calories, and resulted in a gain of 61.3 meters. The distance and ascent values are greater than 1 standard deviation higher than the country mean for these variables. The weight of what she carried is included in all analyses as she did have to carry items to and from the source, even if on the bus part of the way. The total weight of water she collected was 11.3 kg (11.3 L) (Guatemala)

[3] 1 missing & 1 one-way observation was doubled to estimate roundtrip metrics (Honduras)

[4] 2 missing (Honduras)

474   *Weight of loads carried and volumes of water collected.* The weight and volumes of water

475  women carried varied depending on why they were going to the source (e.g., water collection, laundry),

476  if they were carrying children, and whether they were traveling to or returning from sources (**Table 4**).

477   The mean weights of what was carried to water points were higher in Guatemala (6.2kg) and

478  Zimbabwe (3.4kg) compared to Honduras (1.4kg) and Kenya (2.0kg). Empty water containers, which

479  were carried by over half of women followed in Guatemala (12/22) and by most women followed in

480  Honduras, Kenya, and Zimbabwe weighed less than 1kg on average (range: 0.2kg in Honduras to 3.3kg in

481  Kenya). Children were carried in all countries, contributing to the loads carried to and from the sources.

482  More women were observed carrying children in Zimbabwe (7) and Guatemala (6) than in Honduras (1)

483  and Kenya (1). Among these women, the mean child weight was 9.3kg (range: 5.6-12.5kg, both in

484  Guatemala). Dry laundry was carried to water sources by 13 women, all from Guatemala (11) and Kenya

485  (2). The two in Kenya carried less than 2kgs of dry laundry weight. In Guatemala, the mean dry laundry

486  weight carried was 6.5kgs (range: 2.5-14.3kg). Sixteen women (from Guatemala, Kenya, and Zimbabwe)

487  carried other items [range: 0.2kg for a rope (Kenya) or item to scoop water from dug wells (Zimbabwe)

488  to 11kg for a bucket of manure carried one-way for gardening (Zimbabwe)].

489   The mean total weight of loads carried or transported from water sources was 19.3kg. The

490  greatest loads were observed in Guatemala (43.7kg) and Zimbabwe (50.1kg) compared to Honduras

491  (31.7kg) and Kenya (33.6kg). The mean proportional increase in weight carried from the source

492  compared to the journey to the source was 1512%. The increase in weights carried for the return

493  journey is attributable to water, whether women brought back water for later use or wet laundry. The

494  mean weight of containers filled with water during return trips was 17.0kg. The smallest weight of a

495  filled container was observed in Honduras (5.0kg) and the greatest was observed in Zimbabwe (40.1kg).

496  Women in Guatemala who brought wet laundry back from the source carried a mean weight of 18.3kg

497  (range: 8.8-36.5kg), representing a mean increase 11.8kg over the dry laundry (range: 5.6-26.6kg).

The overall mean volume of water collected was 16.1L. The smallest volume of water collected

was observed in Kenya (3.7L) and the most was in Zimbabwe (38.2L). Across countries, the mean

volumes of water collected and carried were greatest in Zimbabwe (19.6L) followed by Guatemala

(15.3L), Honduras (14.8L) and Kenya (12.0L). One women from Zimbabwe reflected on the limited

quantity of water she collects and how she manages: "[If] we had a lot of water, we would be using

water for many different reasons; but due to unavailability of water we are minimizing the water so that

it can accommodate all of the activities that we are undertaking. Therefore, we normally get water from

the far sources; and yet we are able to carry a small jerrican." (Zimbabwe)

On average, women expended 17.4 calories per liter collected; the mean number of calories

expended per liter of water carried were comparable in Guatemala (7.8cal/L), Honduras (11.1cal/L), and

Zimbabwe (11.8cal/L), and were much greater in Kenya (39.3cal/L). The least amount of calories exerted

for one liter of water collected was in Zimbabwe (1.9cal/L), and the greatest was in Kenya (79.5cal/L).

The water weight that women carried or transported was equivalent to substantial proportions

of their body weight. The mean proportion of collected water weight to participant body weight was

28%. Across countries, values were comparable (Guatemala: 29%; Honduras: 23%; Kenya: 25%;

Zimbabwe: 32%). The smallest proportion was recorded in Kenya (9%) and the largest proportion in

Guatemala (72%). (Note, the Guatemalan woman bearing this weight was carrying it unassisted (see

Table 4)). As **Figure 2** shows**,** there is a trend between the journey time, distance, and volume (or

weight) carried. Specifically, women in Kenya, who were experiencing a drought, walked the greatest

distances, took the greatest time fetching water and collected smaller volumes of water. Conversely,

higher volumes of water were collected by those traveling shorter distances for less time.

Physical and Mental Impacts. Women reported myriad physical impacts of water collection from

carrying heavy loads of water, traversing challenging terrain, and expending effort going to and working

at sources. Women discussed general discomfort, pain (neck, back, chest, shoulders, arms, knees, legs,

stomach, ribs), headache, shortness of breath, blisters, weakness, numbness, fatigue, poor sleep, and

exhaustion. As one woman in Kenya commented, "You feel a headache after you finish all the tasks; you

feel very tired. You cannot even sleep well due to the tiredness. You feel pain in your ribs, on your back,

headache; so you feel pain all over your body."

Women reported specific injuries related to animal encounters or falling, whether experienced

by them or those they know. Pregnancy was said to exacerbate discomfort and pain (Kenya) and require

women to carry smaller quantities of water (Guatemala). Women in all countries reported falling and

slipping, and those in Guatemala specifically reported sprained ankles and broken legs because of the

mountainous terrain, made riskier if barefoot. As one woman shared:

"Once, my daughter broke her leg while fetching water here. She didn't like going to get water,

so she carried a larger jar on her side. During one of these trips, she slipped, and her foot got

trapped, leading to the accident. I often advised her to make multiple trips to ease the burden,

but she preferred carrying it all at once." (Guatemala)

Finally, women commented on the mental burden of water collection, the need to ration, and

the implications of not having water, which can be stressful. One woman from Honduras noted the

deliberation involved in water use given the effort to get water:

"In fact, I ration its use because it really represents an effort for me. I need to evaluate when it

will be possible to use it, since bringing it involves a long journey. Therefore, I use it in a

controlled manner, treating it with great care due to the effort involved in obtaining it.

Furthermore, I am very clear for what purposes I will use it. I don't waste it unnecessarily

because I know that getting it is difficult." (Honduras)

3.4 Alignment of Estimated and Measured Water Collection Time

On average, those who indicated they would use the water for drinking (n=30) over-estimated

by 39 minutes and those who collected for 'other uses' (n=6) under-estimated by 11 minutes (**Table 5**).

Average estimates from participants in Honduras, Zimbabwe, and Guatemala were not far from the

average measured times, and times were greatly over-estimated by Kenyan participants. Specifically,

participants underestimated across all water collection activities by 19 minutes in Guatemala and by 2

minutes in Honduras and overestimated by 11 minutes in Zimbabwe. In Kenya, participants

overestimated by an average of 85 minutes. About half (47%) of participants estimated their water

collection time to be within 15 minutes of the measure time (**S2 Table**). Among those whose estimates

were greater than 15 minutes off, a greater proportion overestimated (42%) than underestimated (11%).

554
555
556

Table 4. Water journey transport method and item weight, by country

	Country									
	Guatemala[1] (n=23)		Honduras (n=17)		Kenya (n=22)		Zimbabwe (n=33)		Total (n=95)	
Water container transport method[2] (n, %)										
Carried (arms, head, back with rope) or rolled	22	95.7	16	94.1	22	100.0	30	96.8	90	96.8
Pushed wheelbarrow	0	0.0	1	5.9	0	0.0	1	3.2	2	2.2
Used vehicle	1	4.4	0	0.0	0	0.0	0	0.0	1	1.1

Weight carried (kg)

Total Weight (kg) Carried to Water Point[3]

n		23		17		19		33		92
Mean, SD	6.2	5.2	1.4	3.1	2.0	1.8	3.4	5.2	3.4	4.6
Median		5.2		0.6		1.8		0.9		1.0
Range (min, max)		(0.4-17.1)		(0.2-13.4)		(0.4-8.4)		(0.7-22.8)		(0.2-22.8)

Total Weight (kg) Carried from Water Point[4]

n		21		17		18		31		87
Mean, SD	21.6	10.5	16.2	6.0	14.0	6.4	22.5	6.8	19.3	8.3
Median		20.8		16		11.6		20.6		19.4
Range (min, max)		(8.8-43.7)		(5.0-31.7)		(6.1-33.6)		(14.9-50.1)		(5.0-50.1)

Total Weight (kg) Increase on Return Journey[5]

n		21		17		17		31		86
Mean kg increase, SD	14.9	8.6	14.8	4.5	12.1	6.4	19.4	4.0	15.9	6.5
Median		11.3		15.5		10.5		19.2		16.5
Range - kg (min, max)		(5.6-35.3)		(4.6-21.1)		(3.7-31.3)		(13.4-38.2)		(3.7-38.2)
Mean % increase compared to weight carried to point		706%		2574%		1004%		1740%		1512%
Range - % (min, max)		(41%-2475%)		(137%-9150%)		(125%-3025%)		(59%-2886%)		(41%-9150%)

Weight (kg) of Water Containers, Empty[6]

n		12		17		19		33		81
Mean, SD	0.7	0.4	0.7	0.3	1.4	0.8	0.9	0.2	0.9	0.5
Median		0.7		0.6		1.1		0.9		0.8
Range (min, max)		(0.4-2.0)		(0.2-1.5)		(0.4-3.3)		(0.7-1.9)		(0.2-3.3)

Weight (kg) of Water Containers, With Water[7]

n		12		17		18		31		78
Mean, SD	15.9	8.0	15.4	4.6	13.4	6.4	20.5	4.0	17.0	6.1
Median		13.2		16.0		11.6		20.1		18.3
Range (min, max)		(8.8-36.0)		(5.0-21.7)		(6.1-33.6)		(14.9-40.1)		(5.0-40.1)

Weight (kg) and Volume (L) of Water (1kg water=1L)[8]

n		12		17		18		31		78
Mean, SD	15.3	7.9	14.8	4.5	12.0	6.2	19.6	3.8	16.1	6.1
Median		12.9		15.5		10.5		19.3		17.2
Range (min, max)		(8.4-35.3)		(4.6-21.1)		(3.7-31.1)		(14.0-38.2)		(3.7-38.2)

Calories Expended per Liter of Water Carried[9]

n		12		16		18		31		77
Mean, SD	7.8	4.5	11.1	7.9	39.3	24.5	11.8	9.4	17.4	18.3
Median		6.2		8.9		41.0		8.9		9.4
Range (min, max)		(2.9-16.6)		(2.7-27.4)		(2.1-79.5)		(1.9-49.1)		(1.9-79.5)

Weight (kg) of Carried Children

n		6		1		1		7		15
Mean, SD	8.5	2.4	12.8	NA	7.1	NA	9.7	1.8	9.3	2.3
Median		8.2		12.8		7.1		10		9.8
Range (min, max)	(5.6-12.5)		NA		NA		(6.7-12.2)		(5.6-12.5)	

Weight (kg) of Other Items Carried[10]

n		6		0		2		8		16
Mean, SD	1.8	1.7			0.7	0.6	1.9	3.8	1.7	2.8
Median		1.1				0.7		0.2		0.7
Range (min, max)	(0.6-5.2)				(0.2-1.1)		(0.2-11.0)		(0.2-11.0)	

Weight (kg) of Dry Laundry[11]

n		11		0		2		0		13
Mean, SD	6.5	3.6			1.8	0.1			5.8	3.7
Median		4.7				1.8				4.5
Range (min, max)	(2.5-14.3)				(1.7-1.8)				(1.7-14.3)	

Weight (kg) of Wet Laundry[12]

n		11		0		2		0		13
Mean, SD	18.3	9.0			1.9	0.8			15.8	10.3
Median		15.6				1.9				14.3
Range (min, max)	(8.8-36.5)				(1.3-2.5)				(1.3-36.5)	

Laundry Weight Increase (Dry to Wet)[13]

n		11		0		1		0		12
Mean kg increase, SD	11.8	6.3			0.7	NA			10.9	6.8
Median		10.9				0.7				10.0
Range kg increase (min, max)	(5.6-26.6)				NA				(0.7-26.6)	
Mean % increase		200%				NA				187%
Range % increase (min, max)	(96%-362%)				NA				(39%-363%)	

Participant Body Weight (kg)

n		22		17		22		33		94
Mean, SD	56.9	8.5	65.4	14.5	48.2	7.9	64.4	12.1	59.0	12.8
Median		56.8		65.0		49.7		62.7		56.8
Range (min, max)	(43.8-84.0)		(44.4-94.6)		(35.8-64.2)		(48.9-98.0)		(35.8-98.0)	

Weight (kg) of Containers with Water as Proportion of Participant Body Weight[14]

n		12		17		18		31		78
Mean		29%		23%		25%		32%		28%
Range (min, max)	(15%-72%)		(6%-39%)		(9%-61%)		(16%-63%)		(6%-72%)	

[1] 1 participant completed two separate water journeys to different water sources (Guatemala)

[2] 2 missing (Zimbabwe)

[3] 3 missing (Kenya)

[4] 2 missing (Guatemala); 4 missing (Kenya); 2 missing (Zimbabwe)

[5] 2 missing (Guatemala); 5 missing (Kenya); 2 missing (Zimbabwe)

[6] 11 missing (Guatemala); 3 missing (Kenya)

[7] 11 missing (Guatemala); 4 missing (Kenya); 2 missing (Zimbabwe)

[8] 11 missing & 2 include weight of water container in calculation (Guatemala); 4 missing & 1 includes weight of water container in calculation (Kenya); 2 missing (Zimbabwe)

[9] 11 missing (Guatemala); 1 missing (Honduras); 4 missing (Kenya); 2 missing (Zimbabwe)

[10] *Other items:* soap, basins/washbowls, dishes, brooms (Guatemala); rope (Kenya); scooping plates, manure, vegetables (Zimbabwe)

[11] 1 missing (Kenya)

[12] 1 missing (Kenya)

[13] 2 missing (Kenya)

[14] 10 missing & 1 participant was excluded from aggregate analyses for time, distance, caloric expenditure, and ascent because she took a bus for part of her water journey and was only on foot for part of the time. The weight of what she carried is included in all analyses as she did have to carry items to and from the source, even if on the bus part of the way (Guatemala); 4 missing (Kenya); 2 missing (Zimbabwe)

Table 5. Estimated and measured times to collect water overall (n=36), stratified by if for drinking (n=30) or other uses (n=6)

	n	Estimated time (min)			Measured time (min)			Difference (estimated - measured min)		
		Mean	Range	SD	Mean	Range	SD	Mean	Range	S
All water collection[1]										
All	36	115.4	(10.0 - 520.0)	141.2	85.0	(18.3 - 264.4)	77.7	30.4	(-121.1 - 303.3)	81
Guatemala	4	18.8	(10.0 - 30.0)	8.5	38.2	(20.4 - 70.40)	22.1	-19.4	(-60.4 - (-0.1))	27
Honduras	5	28.0	(20.0 - 30.0)	4.5	29.6	(18.3 - 41.1)	8.1	-1.6	(-11.1 - 11.7)	9
Kenya	12	258.3	(60.0 - 520.0)	169.4	173.1	(20.9 - 264.4)	76.6	85.3	(-121.1 - 303.2)	123
Zimbabwe	15	56.0	(20.0 - 105.0)	24.7	45.4	(18.7 - 83.4)	17.8	10.6	(-20.3 - 65.5)	22
Drinking water										
All	30	132.5	(15.0 - 520.0)	149.0	93.8	(18.7 - 264.4)	82.1	38.7	(-121.1 - 303.2)	86
Guatemala	3	21.7	(15.0 - 30.0)	7.6	27.4	(20.4 - 31.8)	6.2	-5.7	(-11.8 - (-0.1))	5
Honduras	3	26.7	(20.0 - 30.0)	5.8	33.8	(28.8 - 41.1)	6.4	-7.1	(-11.1 - (-1.6))	5
Kenya	12	258.3	(60.0 - 520.0)	169.4	173.1	(20.9 - 264.4)	76.6	85.3	(-121.1 - 303.2)	123
Zimbabwe	12	60.8	(20.0 - 105.0)	23.8	46.2	(18.7 - 83.4)	18.8	14.6	(-20.3 - 65.5)	23
Water for other uses[2]										
All	6	30.0	(10.0 - 60.0)	16.7	40.7	(18.3 - 70.4)	20.3	-10.7	(-60.4 - 11.7)	25
Guatemala	1	10.0	(10.0 - 10.0)		70.4	(70.4 - 70.4)		-60.4	(-60.4 - (-60.4))	
Honduras	2	30.0	(30.0 - 30.0)	0.0	23.2	(18.3 - 28.2)	7.0	6.8	(1.8 - 11.7)	7
Kenya	0									
Zimbabwe	3	36.7	(20.0 - 60.0)	20.8	42.4	(33.1 - 60.9)	16.0	-5.7	(-13.2 - (-0.9))	6

[1] Data from 10 women were excluded because they did not collect water to bring home (9 Guatemala, 1 Kenya) and from 10 women because they did not provide estimated time to collect water (5 Guatemala, 1 Honduras, 4 Zimbabwe). Data from 38 women were excluded because their water source type was not the same for their estimated and measur water journey times (5 Guatemala, 10 Honduras, 9 Kenya, 14 Zimbabwe). For the one Guatemala participant who completed two water journeys, only her water journey to colle water was included.

[2] Water was classified as for 'other uses' if the respondent did not explicitly say it would be used for drinking.

4 DISCUSSION

Our mixed-methods investigation applied novel approaches to comprehensively assess women's lived experiences while traveling to, carrying out work at, and returning from water sources across Guatemala, Honduras, Kenya, and Zimbabwe. Findings illuminate how this unpaid, gendered task demands women's energy, consumes their time, exposes them to risks, and impacts their well-being—often for water from unimproved sources that is of insufficient quantity to ensure needs are met. Our results confirm the hypothesis made by our World Vision SWSW program partners regarding substantial water burden in these areas and support previous calls for water service improvement efforts to prioritize initiatives that reduce the disproportionate time, energy, and physical and mental health burdens of women and ameliorate this gender inequitable burden(19, 33).

In our study, the time women spent, the energy they expended, the distances they traveled, and the elevation they gained going to water sources diverge widely from the results of previous research, revealing how limited the understanding of women's experiences has been. In the more extreme examples, women in our study spent over 4.5 hours (Kenya and Zimbabwe), burned over 700 calories (Kenya and Zimbabwe), walked almost 16 kilometers (Kenya), and ascended nearly 300 meters in elevation (Guatemala) to access and use water sources. Participants in previous research typically traveled less than 1km with measured times of less than one hour(33, 35, 40-42) (exception is a 2014 study in Mozambique)(44); data tracking caloric expenditure is dated(7) and elevation ascent data could not be found.

Journeys that were not as extreme should not be assumed to be of inconsequential impact. Even for women walking shorter distances for less time, our data suggests that this persistent labor amounts to a tremendous inequity of lost time and energy and bodily strain over the course of a lifetime. Moreover, we recognize that impacts are likely greater than what we found. Firstly, we assessed the

burden of a single journey, not the burden of a single day. Thus, efforts may be far greater than what we

recorded. Secondly, we did not assess financial or opportunity costs. Even those spending less time and

energy were still depleting important resources that could have been channeled for other uses. Thirdly,

it is possible that all participants felt the effect of a 'bandwidth tax', which is the mental toll or cognitive

exhaustion experienced among those living in scarcity due to constant thinking about how to manage

limited and tenuous resources(58). As Ray and Smith (2021) have discussed, insufficient, unreliable, and

inadequate water resources can impose such a 'bandwidth tax'(59) Consistent with other studies(11, 12),

many of our participants described water journeys as physically and mentally exhausting. Adapting a

point by Ray and Smith(59), when people are consumed with managing their most immediate needs—

like water needs—they may lack the cognitive energy to plan for or even recognize other current or

future needs for themselves or their families. Thus, while our work provides critical and novel data on

the water access burden, we expect the burden among all participants is likely greater than what we

captured.

Those expending the most energy and time collecting water were getting small quantities and

from unsafe sources, raising questions about additional unmeasured impacts on health, particularly for

proper hydration, nutrition, and hygiene. We found that on average, women in Kenya fetched once per

day as the sole collector in the household, took over 2.5 hours to collect water, expended 40 calories per

liter of water, and collected only 12 liters for a household of six. Assuming water is distributed equally,

all household members would have just two liters of water per day. Based on the WHO publication on

'Domestic water quantity, service level and health', quantities below 5.3 liters/person/day from sources

greater than 1 kilometer away and requiring over 30 minutes of collection time are considered a 'very

high' level of health concern because adequate amounts needed for drinking, cooking, and hygiene

cannot be assured(60). Further, the average adult woman needs at least 2.7L of water per day for

drinking alone, and much more with heavy physical activity and in hot, dry environments (61). Thus, the

women collecting water could be spending considerably more energy per liter they are allotted

compared to other household members. Further, because data on intrahousehold use is limited and

equitable distribution is not certain(9) women could be getting even less water than others, as has been

shown with food (62). Compounding the limited water available and the energy expense, in many

countries women eat 'last and least' in the family, potentially leading to an 'energy output-input'

imbalance(62). Thus, women may not be adequately replacing both the water and the calories they

expend for water collection. While we did not investigate hydration, diet, body composition (e.g., fat

mass), or hygiene, some women in our study did discuss these issues and further work on caloric and

water adequacy is warranted.

Consistent with other studies,(11, 12, 16, 58) participants described how water journeys could

cause injuries, and our assessment of the amount of weight women carried amplifies specific concerns

about musculoskeletal health impacts. A recent scoping review(16) on water carriage suggests that

impacts to musculoskeletal health may be substantial, warranting further research. The authors note

that musculoskeletal health is not widely assessed, and pain associated with it may be under-reported

as women may accept musculoskeletal pain as a consequence of their role as water collectors(16). Other

research argues that, because cervical spinal conditions are a common cause of disability, carrying water,

specifically on the head, may be a major contributor to musculoskeletal disease burden in low-income

countries(11, 63). Our novel data detailing the extensive weights women carried provide further

justification for water infrastructure improvement efforts that center women's health and aspire to

eliminate or at least reduce the physical burden of water carriage. Findings also support the need for

more research on how water carriage may contribute to musculoskeletal health and disability. However,

we emphasize that such research should extend beyond water carriage to encompass all items carried

for water-related work given the tremendous non-water weights we observed women to shoulder (e.g.,

clothing). Based on context, such research could also incorporate firewood collection, a similarly

gendered and physically demanding activity(17, 64, 65).

Our research provides further evidence of multiple water source use and the need for research

and monitoring to look beyond collection from primary drinking water sources alone. Water collection

from multiple sources has been found to be a global phenomenon—even among those with improved

sources(66)—with source selection determined by need, season, and source characteristics, among

other factors(18) In our study, large proportions of women in Guatemala, Honduras, and Zimbabwe

reported visiting different sources for drinking and 'other uses.' For some, as we saw in Guatemala,

women did not collect water but carried out strenuous laundering activities at the source. The recently

published 'Priority Gender-Specific Indicators for WASH Monitoring Under SDG Targets 6.1 and 6.2'

recommends new indicators that could shed light on the burden of collecting from multiple sources (i.e.,

'Average time primary water collector spends per day collecting drinking water, by sex and age' and

'Average time primary water collector spends per day collecting water for all household needs, by sex

and age')(67). These indicators do not encompass water work at the sources, which, based on our

findings, deserves more attention.

Our findings show that self-reported time estimates for water collection are imperfect, but still

valuable for monitoring. Half of respondents' estimates were within 15 minutes of the measured time.

Most of the participants who over-estimated by more than 15 minutes were from Kenya where water

journeys were very long. Field team members remarked that these women often take animals with

them to water at the sources, which can lengthen the process substantially. However, they did not bring

animals on the days they were followed. If they had, estimates could have been closer to actual water

journey times. Those who were closer to sources had more accurate estimations, thus we do not expect

that time estimates would interfere with how JMP classifies access for global monitoring, though self-

reported time burden assessments may need caution when including those who are particularly far from

sources.

Programs that aim to improve water infrastructure should leverage the methodologies we

employed to objectively evaluate if programs positively impact the lives of women and girls.

Unfortunately, while the myriad benefits of how improved water services can improve the lives of

women and girls have been long proclaimed(19, 68), rigorous assessments to determine impact are

scant. Benefits are not guaranteed and may be (erroneously) assumed, instead of tested. As one study in

Zambia demonstrates, water improvements may ease burden for some but shift it to others. Specifically,

while researchers found that a new, closer, improved water supply (borehole) significantly decreased

the amount of time adult women spent on water-related work, girls who lived close to the new borehole

significantly increased the time they spent on water-related work and significantly decreased the

amount of time they spent in school and on homework (with no changes observed among boys)(69).

Alternatively, even if closer sources are made available, women may just frequent them more often.

They may get more water but spend no less time or energy in the process. Or, women may still use

farther sources if a newer one is not always available (as observed in our study among Kenyan women)

or does not meet their specific preferences or needs (as noted in India(18), Ghana,(70) and observed in

our study among women from Guatemala who carried heavy clothing loads to and from the river for

washing despite access to other sources). Thus, evaluations, using approaches like ours, should be used

to assess quantitative and qualitative impacts on women's lives.

5 STRENGTHS AND LIMITATIONS

Our mixed-methods study provides a comprehensive understanding of women's experiences

accessing water sources using novel and state-of-the-art approaches, though limitations exist.

Community sampling was purposive, and individual sampling was by convenience, thus data may not be

representative, and generalizability is limited. We only assessed a single journey to a water source per

enrolled household due to time constraints. Future research could collect data throughout the day and

from all water collectors within a household to capture daily burden for individuals and the household

more accurately. Journeys were recorded as a complete experience, preventing assessment of the time

and energy burden of various activities within the journey (e.g., the walk to source vs. the walk from

source; water extraction). Caloric expenditure is modeled by watches, which become more precise over

time as the watch learns about the individual wearer's behavior. However, watches were worn only

once by each participant, so estimates may be imperfect. Transcriptions of some IDIs were not verbatim,

but summaries and descriptions of conversations still contributed to the analysis.

691 6 CONCLUSION

While water is necessary to sustain life and thus should be considered a source of nourishment,

we show how a lack of adequate and accessible water was actually depleting, draining women of energy

and time, and posing risks to their well-being. Crider and Ray (2022) argue that research in the sector

often focuses on women for their instrumental value or how they can support programming, instead of

their intrinsic worth(9). Through our use of quantitative and qualitative research methods, we centered

women's lived experiences, which have intrinsic value. The results of our study further reinforce the

need to not only redouble efforts to improve water access in low-resource settings but to rigorously

measure the impacts of such efforts on women's lives.

700 ACKNOWLEDGEMENTS

We are grateful to the participants of this study for their time and for sharing their experiences.

We would like to thank the following partners and enumerators for their research collaboration. From

Centro Universitario de Occidente, de la Universidad de San Carlos de Guatemala: Raul Bethancourt, Luis

Alfredo López Cortez, Leonardo Cabrera López, Facundo Zacarías López Sánchez, Alex Fernando López

Cortez, Zaida Anaí Vasquez García, Gladys Judith Vásquez García López, Vilma Rosario López Sánchez,

Any Gabriela León Ramon, Heidy Maribel Baten Velásquez, Lucy Antonieta Ixmucané Capriel Herrera,

Lesbia Beatriz Xiloj Hernandez, Ana Rosa León, José Joél Nolasco Yacabalquiej, Ingrid Johana Guachiac

Gómez; from World Vision Guatemala: Enri Maldonado, Erick Calderón, Estuardo Gonzales; from

Universidad Nacional Autónoma de Honduras: Rosaura Suyapa Rodriguez Funez, Ashly Michelle Méndez,

Nastin Nahomy Tilguant Ávila, Brenda Paola Cruz, Yesi Karolina Murillo Vega, Jorge Luis Cálix Barahona,

Dulce Milagro Sevilla Sosa, Wilber Adonay Ávila Valladares, Néstor Fabricio Dávila González, Pedro

Alejandro Hernández; from St. Paul's University, Kenya: Lesoito Jeremiah Christopher, Leshaule

Benedicto Alois, Patricia Lekiliyo, Lechalote Ben Ljania, Nasieku Lolchuragi, Lekalantula Raina Deborah,

Glory Karimi Kiriinya, Lekonte Salome Mary, Lelikoo Naanyu Sarah, John Kalasinga Lerosion; from World

Vision Kenya: Kevin Muche; from Datalyst Africa: Molly Manyonganise, Eve Nyemba, Sunga Mzeche,

Nyasha Kamwire, Luckness Zimanyiwa, Sibonisiwe Mpofu, Tapiwa Mapfumo, Kudzai Zibgwi, Pricilla

Moyo, Gladys Mhaka, Sithokozile Sibanda, Gugulethu Mnkandhla; and from World Vision Zimbabwe:

Nobuhle Mlotshwa. Lastly, we would like to thank Jorge Beteta for his translation services.

REFERENCES

1.   Wolf J, Johnston RB, Ambelu A, Arnold BF, Bain R, Brauer M, et al. Burden of disease attributable

to unsafe drinking water, sanitation, and hygiene in domestic settings: a global analysis for selected

adverse health outcomes. Lancet. 2023;401(10393):2060-71.

2.   ICWE. The Dublin Statement on Water and Sustainable Development International Conference

on Water and the Environment; Dublin, Ireland1992.

3.   The human right to water and sanitation. , (28 July 2010, 2010).

4.   United Nations Children's Fund (UNICEF) and World Health Organization (WHO). Progress on

household drinking water, sanitation and hygiene 2000–2022: special focus on gender. New York; 2023.

5.   Caruso BA, Chipungu J, Kulkarni S, Ray I. Women, work, and water. Lancet. 2023.

6.   Caruso BA. Water is life, particularly for women. Nature Water. 2023;1(2):124-.

7.   White GF, Bradley DJ, White AU. Drawers of water: domestic water use in East Africa. Chicago:

University of Chicago Press; 1972.

8.   Thompson J. Drawers of water II: 30 years of change in domestic water use & environmental

health in east Africa. Summary: Iied; 2001.

9.   Crider YS, Ray I. Water and Development: A Gender Perspective. Oxford Research Encyclopedia

of Environmental Science2022.

10.   Gandhi S, Dilbaghi M, Raina K. Rural women carry the load of fetching water. Indian Journal of

Social Work. 2003;64:65-75.

11.   Geere J-AL, Cortobius M, Geere JH, Hammer CC, Hunter PR. Is water carriage associated with the

water carrier's health? A systematic review of quantitative and qualitative evidence. BMJ global health.

2018;3(3):e000764.

12.   Venkataramanan V, Geere J-AL, Thomae B, Stoler J, Hunter PR, Young SL. In pursuit of

'safe'water: the burden of personal injury from water fetching in 21 low-income and middle-income

countries. BMJ global health. 2020;5(10):e003328.

13.   Asaba RB, Fagan H, Kabonesa C, Mugumya F. Beyond distance and time: gender and the burden

of water collection in rural Uganda. WH2O: Journal of Gender and Water. 2013;2(1):31-8.

14.   Mukiese J-MN, Sangana CB, Kiyombo GM. The Burden of Musculoskeletal Pains and Injuries

from Water-Carrying Among School-Going Adolescent Girls in Kinshasa's Peri-Urban Settings,

Democratic Republic of Congo. European Journal of Development Studies. 2024;4(3):52-61.

15.   Adams EA, Byrns S, Kumwenda S, Quilliam R, Mkandawire T, Price H. Water journeys: Household

water insecurity, health risks, and embodiment in slums and informal settlements. Soc Sci Med.

2022;313:115394.

16.   Meeuwisse S, Elliott SJ, Bennett A, Kapoor V. Water fetching and musculoskeletal health across

the life-course in Sub-Saharan Africa: A scoping review. PLOS Global Public Health. 2024;4(9):e0003630.

17.   Ho EW, Strohmeier-Breuning S, Rossanese M, Charron D, Pennise D, Graham JP. Diverse health,

gender and economic impacts from domestic transport of water and solid fuel: A systematic review.

International Journal of Environmental Research and Public Health. 2021;18(19):10355.

18.   Patrick M, Hennink M, Dasmohapatra M, Rout M, Clasen T, Caruso BA. Influences on water

source selection and use among women in rural Odisha, India. Frontiers in Water. 2023;5:7.

19.   Cairncross S. The benefits of water supply. Developing world water. 1987;2:30-4.

20.   Caruso BA, Conrad A, Patrick M, Owens A, Kviten K, Zarella O, et al. Water, Sanitation, and

Women's Empowerment: A systematic review and qualitative metasynthesis. PLOS Water.

2022;1(6):e0000026.

21.   UN-Women, DESA. Progress on the Sustainable Development Goals: The Gender Snapshot 2024.

New York: UN-Women and DESA; 2024.

22.   Sorenson SB, Morssink C, Campos PA. Safe access to safe water in low income countries: water

fetching in current times. Soc Sci Med. 2011;72(9):1522-6.

23.   Sinharoy SS, Caruso BA. On World Water Day, gender equality and empowerment require

attention. Lancet Planet Health. 2019;3(5):e202-e3.

24.   Van Houweling E. "A good wife brings her husband bath water": Gender roles and water

practices in Nampula, Mozambique. Society & natural resources. 2016;29(9):1065-78.

25.   Thai NV, Guevara JR. Women and water management: A policy assessment—A case study in An

Giang Province, Mekong Delta, Vietnam. Asia-Pacific Journal of Rural Development. 2019;29(1):77-97.

26.   Chew JF, Corlin L, Ona F, Pinto S, Fenyi-Baah E, Osei BG, et al. Water source preferences and

water quality perceptions among women in the eastern region, Ghana: a grounded theory study.

International Journal of Environmental Research and Public Health. 2019;16(20):3835.

27.   Zozmann H, Klassert C, Klauer B, Gawel E. Water procurement time and its implications for

household water demand—insights from a water diary study in five informal settlements of pune, India.

Water. 2022;14(7):1009.

28.   Cassivi A, Guilherme S, Bain R, Tilley E, Waygood EOD, Dorea C. Drinking water accessibility and

quantity in low and middle-income countries: A systematic review. International Journal of Hygiene and

Environmental Health. 2019;222(7):1011-20.

29.   Bachwenkizi J, Mohamed H, Funsan P, Rweyemamu D, Nelson W, Shao M, et al. Access to water

sources and intimate partner violence against women in 26 Sub-Saharan African countries. Hygiene and

Environmental Health Advances. 2023;7:100063.

30.   Hamlet LC, Chakrabarti S, Kaminsky J. Reduced water collection time improves learning

achievement among primary school children in India. Water Research. 2021;203:117527.

31.   Bisung E, Elliott SJ. Improvement in access to safe water, household water insecurity, and time

savings: A cross-sectional retrospective study in Kenya. Soc Sci Med. 2018;200:1-8.

32.   Winter JC, Darmstadt GL, Davis J. The role of piped water supplies in advancing health, economic

development, and gender equality in rural communities. Soc Sci Med. 2021;270:113599.

33.   Pickering AJ, Davis J. Freshwater availability and water fetching distance affect child health in

sub-Saharan Africa. Environmental science & technology. 2012;46(4):2391-7.

34.   Caruso BA, Sinharoy SS. Gender data gaps represent missed opportunities in WASH. The Lancet

Global Health. 2019;7(12):e1617.

35.   Crow B, Davies J, Paterson S, Miles J. Using GPS and recall to understand water collection in

Kenyan informal settlements. Water international. 2013;38(1):43-60.

36.   Dongzagla A, Nunbogu AM, Fielmua N. Does self-reported water collection time differ from

observed water collection time? Evidence from the Upper West Region of Ghana. Journal of Water,

Sanitation and Hygiene for Development. 2020;10(2):357-65.

37.   Cassivi A, Tilley E, Waygood EOD, Dorea C. Evaluating self-reported measures and alternatives to

monitor access to drinking water: a case study in Malawi. Science of the Total Environment.

2021;750:141516.

38.   Seymour G, Malapit HJ, Quisumbing AR. Measuring time use in development settings. World

Bank Policy Research Working Paper. 2017(8147).

39.   Nygren BL, O'Reilly CE, Rajasingham A, Omore R, Ombok M, Awuor AO, et al. The relationship

between distance to water source and moderate-to-severe diarrhea in the global enterics multi-center

study in Kenya, 2008–2011. The American journal of tropical medicine and hygiene. 2016;94(5):1143.

40.   Gorter AC, Sandiford P, Smith GD, Pauw JP. Water supply, sanitation and diarrhoeal disease in

Nicaragua: results from a case-control study. International journal of epidemiology. 1991;20(2):527-33.

41.   Gorter AC, Sandiford P, Pauw J, Morales P, Pérez RM, Alberts H. Hygiene behaviour in rural

Nicaragua in relation to diarrhoea. International journal of epidemiology. 1998;27(6):1090-100.

42.   Semenza JC, Roberts L, Henderson A, Bogan J, Rubin C. Water distribution system and diarrheal

disease transmission: a case study in Uzbekistan. The American journal of tropical medicine and hygiene.

1998;59(6):941-6.

43.   Wang X, Hunter PR. A systematic review and meta-analysis of the association between self-

reported diarrheal disease and distance from home to water source. The American journal of tropical

medicine and hygiene. 2010;83(3):582.

44.   Ho JC, Russel KC, Davis J. The challenge of global water access monitoring: evaluating straight-

line distance versus self-reported travel time among rural households in Mozambique. Journal of water

and health. 2014;12(1):173-83.

45.   Smiley SL, Curtis A, Kiwango JP. Using spatial video to analyze and map the water-fetching path

in challenging environments: a case study of Dar es Salaam, Tanzania. Tropical Medicine and Infectious

Disease. 2017;2(2):8.

46.   La Frenierre J. The burden of fetching water: Using caloric expenditure as an indicator of access

to safe drinking water—A case study from Xieng Khouang Province, Lao PDR: University of Denver; 2009.

47.   Russel KC, Buman MP, Davis J, Haskell WL. Estimating energy expenditure of head-hauling water

and grain grinding from heart rate monitor measurements in northern Mozambique. Public Health Nutr.

2020;23(16):2886-97.

48.   Hennink M, Kaiser BN. Sample sizes for saturation in qualitative research: A systematic review of

empirical tests. Soc Sci Med. 2022;292:114523.

49.   CIA. The World Factbook: Guatemala Country Factsheet [updated 29 October 2024; cited 2024

5 November]. Available from: https://www.cia.gov/the-world-

factbook/countries/guatemala/factsheets/.

50.   CIA. The World Factbook: Honduras Country Factsheet [updated 28 October 2024; cited 2024

5 November]. Available from: https://www.cia.gov/the-world-factbook/countries/honduras/.

51.   CIA. The World Factobook: Kenya Country Factsheet [updated 28 October 2024; cited 2024 5

November]. Available from: https://www.cia.gov/the-world-

factbook/countries/kenya/factsheets/.

52.   CIA. The World Factobook: Zimbabwe Country Factsheet [updated 29 October 2024; cited 2024

5 November]. Available from: https://www.cia.gov/the-world-

factbook/countries/zimbabwe/factsheets/.

53.   Carpiano RM. Come take a walk with me: The "Go-Along" interview as a novel method for

studying the implications of place for health and well-being. Health & place. 2009;15(1):263-72.

54.   Garcia CM, Eisenberg ME, Frerich EA, Lechner KE, Lust K. Conducting go-along interviews to

understand context and promote health. Qualitative health research. 2012;22(10):1395-403.

55.   Evans J, Jones P. The walking interview: Methodology, mobility and place. Applied geography.

2011;31(2):849-58.

56.   Boateng GO, Collins SM, Mbullo P, Wekesa P, Onono M, Neilands TB, et al. A novel household

water insecurity scale: Procedures and psychometric analysis among postpartum women in western

Kenya. PLoS One. 2018;13(6):e0198591.

57.   WHO/UNICEF Joint Monitoring Programme for Water Supply S, and Hygiene. Drinking Water

[Available from: https://washdata.org/monitoring/drinking-water.

58.   Mullainathan S, Shafir E. Scarcity: Why having too little means so much. 2013. London: Allen

Lane.

59.   Ray I, Smith KR. Towards safe drinking water and clean cooking for all. The Lancet Global Health.

2021;9(3):e361-e5.

60.   Howard G, Bartram J, Williams A, Overbo A, Geere J-A, Organization WH. Domestic water

quantity, service level and health: World Health Organization; 2020.

61.   Sawka MN, Cheuvront SN, Carter R. Human water needs. Nutrition reviews.

2005;63(suppl_1):S30-S9.

62.   Batliwala S, Reddy AK. Energy for women and women for energy (engendering energy and

empowering women). Energy for Sustainable Development. 2003;7(3):33-43.

63.   Geere J-A, Bartram J, Bates L, Danquah L, Evans B, Fisher MB, et al. Carrying water may be a

major contributor to disability from musculoskeletal disorders in low income countries: a cross-sectional

survey in South Africa, Ghana and Vietnam. Journal of global health. 2018;8(1).

64.   Njenga M, Gitau JK, Mendum R. Women's work is never done: Lifting the gendered burden of

firewood collection and household energy use in Kenya. Energy Research & Social Science.

2021;77:102071.

65.   Clancy JS, Skutsch M, Batchelor S. The Gender-Energy-Poverty Nexus: Finding the energy to

address gender concerns in development. 2002.

66.   Daly SW, Lowe J, Hornsby GM, Harris AR. Multiple water source use in low-and middle-income

countries: a systematic review. Journal of Water and Health. 2021;19(3):370-92.

67.   Bethany A. Caruso, Jenala Chipungu, Julie Hennegan, Albert Motivans, Lauren Pandolfelli,

Madeleine Patrick, et al. Priority Gender-Specific Indicators for WASH Monitoring under SDG Targets 6.1

and 6.2: Recommendations for National and Global Monitoring. New York; 2024.

68.   Brocklehurst C, Bartram J. Swimming upstream: why sanitation, hygiene and water are so

important to mothers and their daughters. Bull World Health Organ. 2010;88(7):482-.

69.   Shimamura Y, Shimizutani S, Taguchi S, Yamada H. The impact of better access to improved

water sources on health, schooling, and water collection of girls and boys in rural Zambia. The Journal of

Development Studies. 2022;58(9):1750-71.

70.   Kulinkina AV, Sodipo MO, Schultes OL, Osei BG, Agyapong EA, Egorov AI, et al. Rural Ghanaian

households are more likely to use alternative unimproved water sources when water from boreholes

has undesirable organoleptic characteristics. International journal of hygiene and environmental health.

2020;227:113514.

Supporting Information

S1 Checklist. PLOS Global Public Health inclusivity in global research

S1 Table. Thematic descriptions of women's water collection experiences and practices, by community

S2 Table. Water Journey estimated time comparison and discordance, by country

S1 Protocol. Protocol for go-along interviews, semi-structured observations, and research equipment

S1 Tools. Data collection tools for participant demographic information, go-along interviews, and semi-

structured observations