Integrative Genome-Wide Association Studies of COVID-19 Susceptibility and Hospitalization Reveal Risk Loci for Long COVID

Zhongshan Cheng[1*]

1. Center for Applied Bioinformatics, St. Jude Children's Research Hospital, 262 Danny Thomas Hospital, Memphis, TN 38105, USA

   *Corresponding authors:

   Zhongshan Cheng: Senior Bioinformatics Research Scientist, Center for applied bioinformatics, St. Jude Children's Research Hospital, 262 Danny Thomas Place, Memphis, TN 38105, United States; email: cheng.zhong.shan@gmail.com

Keywords: COVID-19, SARS-CoV2, Long COVID, GWAS, TWAS, Hospitalization, Susceptibility

Abstract

Long COVID presents a significant public health challenge, characterized by over 200 reported symptoms spanning multiple organ systems, and genome-wide association studies (GWAS) has been hindered by the disorder's symptom heterogeneity and the limited power of available datasets. To overcome the challenges of limited sample size or heterogeneity of long COVID cohorts, an integrative analysis using COVID-19 GWAS data from the Host Genetics Initiative (HGI, release 7) is performed by leveraging proxy phenotypes, resulting in 62 independent single nucleotide polymorphisms (SNPs) prioritized as potentially associated with long COVID. These SNPs are categorized into three groups: (1) severe COVID-19-specific SNPs, such as SNPs mapped to two well-known loci involved in SARS-CoV2 entry (*ACE2* [rs190509934] and *TMPRSS2* [rs12329760]), as well as variants of *DPP9* (rs7251000), *FOXP4* (rs12660421) and *HLA-DQA1* (rs17219281), exhibiting associations with severe COVID-19 but displaying weaker signals in non-hospitalized COVID-19 cases; (2) SNPs associated with both severe and mild COVID-19, including the SNP close to *ABO* (rs505922), representing a catalog of SNPs predispose to both acute COVID-19 and long COVID; and (3) non-hospitalization-specific SNPs, such as variants in *KCTD16* (rs62401842) and *WASF3* (rs56143829), highlighting genetic contributors specific to mild COVID-19 cases that might also contribute to long COVID. Further transcriptome-wide association studies (TWASs) across 48 GTEx tissues, reveal that long COVID exhibits weaker association signals across heart, brain, and muscle-related tissues, as determined by correlations between gene expression of adjacent genes of candidate SNPs (43 out of 62 SNPs) and different COVID phenotypes, with notable TWAS hits including *DPP9* (rs7251000), *CCR1* (rs17078348), and *THBS3* (rs41264915). Phenome-wide TWASs also link *HLA-DQA1* (rs17219281), *HLA-A* (rs9260038), and *HLA-C* (rs1634761), *GSDMB* (rs9916158) and *FOXP4* (rs12660421) with long COVID. Taken together, current integrative approach offers a scalable framework for long COVID research by maximizing the statistical power of existing large-scale COVID-19 GWASs and provides novel insights into the genetic underpinnings of long COVID.

Introduction

Long COVID remains a critical and complex public health issue, with a diverse array of over 200 reported symptoms affecting multiple organ systems[1]. These symptoms, which include fatigue, cognitive impairment, respiratory issues, and cardiovascular problems, can persist well beyond the acute phase of COVID-19 infection, revealing a unique challenge for researchers to identify genetic risk factors using genome-wide association studies (GWAS), as the wide variation in symptom clusters may arise from different biological mechanisms of long COVID and require extensive sample sizes to detect meaningful associations. To address this complexity, long COVID GWAS requires not only large and diverse patient datasets but also refined approaches to phenotype classification and statistical analyses.

Among the most significant efforts to understand the genetic basis of long COVID is the large GWAS conducted by Lama et al.[2], which analyzed approximately 3,500 long COVID cases but not powerful enough to reveal the full genetic landscape due to several key limitations. Primarily, the study drew cases from early phases of the COVID-19 pandemic based on more immediate post-acute symptoms rather than on late-onset long COVID, which can emerge years after initial infection. As a result, potential long COVID cases with delayed symptom onset may be classified as controls, resulting in dilution of statistical power needed to detect true genetic associations. Despite these limitations, Lama et al.[2] identified a single gene, *FOXP4*, on chromosome 17, reaching genome-wide significance for association with long COVID. *FOXP4* variants were found to be strongly associated with both long COVID and severe COVID-19, in line with a shared genetic predisposition that may drive both conditions. However, a stronger association of *FOXP4* with severe COVID-19 suggests that *FOXP4* may contribute primarily to risk factors influencing severe acute COVID-19 cases that later progress to long COVID. Given that long COVID can also follow mild COVID-19, additional genetic contributors associated specifically with mild cases likely remain undiscovered.

Another long COVID GWAS conducted by Taylor et al.[3] provided further insights into genetic factors specific to a subtype of long COVID by investigating genetic overlap between long COVID and Myalgic Encephalomyelitis/Chronic Fatigue Syndrome (ME/CFS), a condition with substantial symptom overlap, including chronic fatigue and cognitive impairments. They suggested that it is beneficial to analyze long COVID subtypes separately, as different symptom profiles could be associated with distinct genetic and biological pathways. However, their GWAS faced limitations due to its relatively small sample size (n < 1,000 long COVID cases) with restricted statistical power. Additionally, the study only reported SNPs derived by network analysis and did not identify any SNPs that reached genome-wide significance for long COVID, emphasizing the need for larger, more statistically robust studies. The absence of genome-

wide significant SNPs underscores a major challenge in long COVID research: even well-defined

subtypes require substantial sample sizes to detect genetic variants with confidence.

More recently, a multi-ancestry GWAS was deposited at the preprint server medRxiv by Chaudhary et

al.[4], which expanded the scope of long COVID GWAS by including participants from diverse ancestral

backgrounds. Notably, several immune-related loci, such as *ABO*, *BPTF*, and *HLA-DQA1*, were reported

to be linked with chronic fatigue syndrome (CFS), fibromyalgia, and depression, suggesting potential

immune dysregulation in long COVID. Despite this, this multi-ancestry long COVID GWAS still has

limited sample size in non-European samples regarding the heterogeneity of long COVID symptoms,

leading to the lack of sufficient power to detect associations for less common symptom clusters or subtle

genetic variants. Furthermore, the study relied on self-reported cases, which may introduce

misclassification biases, particularly given that long COVID lacks a single diagnostic marker. Self-

reported symptoms, though valuable for capturing diverse experiences, can introduce noise that

complicates statistical analysis and gene identification.

To advance insights into long COVID genetics, additional large-scale GWAS or alternative analyses that

utilize existing COVID-19 GWAS summary statistics are essential to uncover genetic factors masked by

current sample size limitations. Integrative approaches leveraging large publicly available GWAS of

COVID-19 hospitalization and SARS-CoV-2 susceptibility may provide the statistical power required to

detect SNPs linked to long COVID.

In this study, an integrative analysis of COVID-19 GWASs from the Host Genetics Initiative (HGI,

release 7) was performed with the assumption that these phenotypes of COVID-19 hospitalization and

susceptibility considered together to be a proxy phenotype of long COVID. Hypothesizing that long

COVID may develop after either mild or severe COVID-19, this study was aimed to identify potential

shared or hospitalization-specific genetic elements, as well as non-hospitalization-specific genetic

elements. The current analysis prioritized 62 top independent SNPs potentially associated with long

COVID into three SNP categories: (1) "Severe COVID-19-Specific SNPs", such as these previously

reported long COVID risk SNPs rs12660421 (*FOXP4*), rs10774679 (*OAS3*), and rs17219281 (*HLA-*

DQA1), which show significant associations in both GWASs of hospitalized COVID-19 vs. general

population and SARS-CoV-2 infection vs. general population with relatively weaker associations in

hospitalized vs. non-hospitalized COVID-19; (2) "SNPs Associated with Both Severe and Mild COVID-

19", including the well-known long COVID risk SNPs, rs505922 (*ABO*), covering SNPs that may

contribute to long COVID risk across a range of COVID-19 severities; (3) an extra SNP category

specifically emerged from non-hospitalized COVID-19 patients ("Non-Hospitalization-Specific Long

COVID SNPs") was further explored to prioritize 20 potential long COVID SNPs, such as the reported

long COVID SNP rs62401842 (*KCTD16*), as well as rs56143829 (*WASF3*). These prioritized SNPs were

evaluated in association with long COVID based on published long COVID GWASs, and transcriptome-

wide association studies (TWAS) were performed for these HGI COVID-19

hospitalization/susceptibility/long COVID GWASs to determine the expression level association of genes

close to these candidate SNPs with these phenotypes, resulting in the conclusion that long COVID is

likely to have different tissue etiology, particularly occurring in heart, brain, and muscle related tissues.

Further phenome-wide TWASs were conducted to evaluate additional candidate genes potentially

involved in long COVID. This integrative approach prioritizes SNPs as well as genes close to them with

potential relevance to long COVID, offering a scalable alternative to screen long COVID risk loci by

maximizing the statistical power of existing large-sample COVID-19 GWAS data from HGI.

Methods

Genome-wide Screening of Genetic SNPs Potentially Associated with Long COVID

Given the challenge of identifying genetic associations with long COVID due to its diverse

symptomatology and the limited cohort sizes in GWAS, proxy phenotypes such as SARS-CoV-2

susceptibility (also refer to COVID-19 susceptibility in this study) and COVID-19 hospitalization were

leveraged to prioritize relevant SNPs. This strategy is supported by evidence that long COVID can

develop following both mild and severe COVID-19, suggesting shared or distinct genetic risk factors may

influence acute-phase outcomes of COVID-19 (e.g., hospitalization, non-hospitalization, or susceptibility

to SARS-CoV2 infection) and subsequent long COVID susceptibility.

Publicly available COVID-19 GWAS datasets were downloaded from the HGI (release 7), including HGI-

C2 (COVID-19 [n=159,840] vs. the general population [n=2,782,977]), HGI-B1 (hospitalized [n=16,512]

vs. non-hospitalized COVID-19 [n=71,321]), and HGI-B2 (hospitalized COVID-19 [n=44,986] vs. the

general population [n=2,356,386]), and analyzed using an integrative analysis pipeline to prioritize

potential long COVID SNPs. Although HGI also shared the long COVID GWASs with 4 slightly different

case-control designs that were published by Lammi et al[2], these data were not initially utilized to

prioritize long COVID SNPs due to the fact that these long COVID GWASs contain potential bias to

severe COVID-19, which may lead to the missing of mild COVID-19 related long COVID risk SNPs.

Again, the previously published HGI long COVID GWASs only identified one genome-wide significant

locus represented by *FOXP4*. Therefore, in the current study the SNP prioritization workflow primarily

focused on two key long COVID risk SNP categories related to severe COVID-19 that are potentially

relevant to long COVID. The first SNP categories is "Severe COVID-19-Specific SNPs", which includes

SNPs associated with both SARS-CoV-2 susceptibility and COVID-19 hospitalization but tends to be

more prevalent within hospitalized than in non-hospitalized cases. These SNPs are hypothesized to influence long COVID risk based on the epidemiological observation that multiple SARS-CoV2 infections and COVID-19 hospitalizations elevate the likelihood of developing the long COVID condition[5]. The following selection criteria was applied, including (a) $P \leq 0.05$ in HGI-C2, capturing SNPs linked to COVID-19 susceptibility; (b) $P \leq 0.05$ in HGI-B1, identifying SNPs that predispose individuals to COVID-19 hospitalization rather than mild COVID-19; (c) $P \leq 0.05$ in HGI-B2, identifying SNPs distinguishing hospitalized COVID-19 individuals from the general population; (d) $P > 0.05$ in the differential GWAS of HGI-B1 vs. HGI-B2, excluding SNPs with differential effect sizes between these comparisons; (e) all SNPs required to display consistent effect directions across HGI-C2, HGI-B1, and HGI-B2. Meanwhile, the second SNP categories is "SNPs Associated with Both Severe and Mild COVID-19", which influence both mild and severe COVID-19 outcomes that are plausible candidates for long COVID risk, given that non-hospitalized cases can also develop long COVID. These SNPs are unlikely to show significant differentiation between hospitalized and non-hospitalized cases but may exhibit associations with SARS-CoV-2 susceptibility and COVID-19 hospitalization when compared to the general population. The selection criteria for this group were performed as (a) $P \leq 0.05$ in HGI-C2, identifying COVID-19 susceptibility-related SNPs; (b) $P \leq 0.05$ in HGI-B2, identifying SNPs associated with COVID-19 hospitalization; (c) $P > 0.05$ in HGI-B1, indicating no significant difference in SNP effects between severe and mild COVID-19; (d) $P \leq 0.05$ in the differential GWAS of HGI-B1 vs. HGI-B2, capturing SNPs with distinct effects in these datasets; (e) all prioritized SNPs need to fulfill consistent effect size directions across HGI-C2 and HGI-B2. Finally, SNPs met either of the above two sets of conditions were further evaluated based on the criterion of passing the $P < 1E-7$ in any of the three GWAS datasets (HGI-B1, HGI-B2, or HGI-C2). This prioritization framework provides a systematic approach to identifying genetic variants potentially predisposing individuals to long COVID by leveraging large-scale COVID-19 GWAS data, resulting in 42 top independent SNPs that were subjected to further evaluation on its potential involvement in long COVID by COVID19_GWAS_Analyzer[6,7] (https://github.com/chengzhongshan/COVID19_GWAS_Analyzer).

Identification of SNPs Potentially Associated with Long COVID in Non-Hospitalized Cases
Current COVID-19 GWAS analyses from the HGI are predominantly biased toward hospitalized cases, potentially overlooking SNPs associated with long COVID that emerge specifically in non-hospitalized individuals. To address this gap, SNPs were filtered using the following criteria: (1) $P > 0.05$ in the GWAS comparing hospitalized cases with the general population (HGI-B2), which excludes SNPs associated with COVID-19 hospitalization; (2) $P \leq 0.05$ in the GWAS comparing hospitalized cases with non-hospitalized cases (HGI-B1), which captures SNPs significantly differentiating these two groups; (3)

$P \leq 0.05$ in the differential GWAS between HGI-B1 and HGI-B2, retaining SNPs with differential

association signals between these datasets. To further refine the selection, independent SNPs met a

suggestive significance threshold ($P<1E-5$) were prioritized and ensured no other genome-wide

significant SNPs located within a 500-kb window in either HGI-B2 or HGI-C2. The use of a relaxed

significance threshold aimed to account for the higher heterogeneity among non-hospitalized cases in

HGI-B1 and the inherent bias of HGI GWASs toward identifying SNPs associated with hospitalization.

This approach enables the identification of SNPs that might represent unique genetic contributors to long

COVID risk in non-hospitalized individuals.

Comparing Two COVID-19 Hospitalization GWASs Using the Differential Z-Score Method

To assess differences in effect sizes of SNPs between two COVID-19 GWAS datasets, such as one GWAS

comparing hospitalized cases with non-hospitalized cases (HGI-B1) and the other GWAS comparing

hospitalized cases with the general population (HGI-B2), as well as comparison SNP effect size between

sex-stratified COVID-19 hospitalization GWASs, differential z-score method[7] was applied in current

study. This approach quantifies the disparity in effect sizes (β) of each SNP between two GWASs by

calculating a differential z-score, which incorporates adjustments for potential sample overlap between

studies. The corresponding p-value was derived from the normalized differential z-scores. The differential

z-score is defined as:

$$\Delta Z - score = \frac{gwas1.\beta - gwas2.\beta}{\sqrt{(gwas1.se^2 + gwas2.se^2)}}$$

$$diff.zscore = Normalization(\Delta Z - score)$$

$$P = pnorm(-|diff.zscore|) * 2$$

Here, gwas1.β and gwas2.β represent the effect sizes for a given SNP from the two GWASs, while

gwas1.se and gwas2.se denote the corresponding standard errors. To account for genome-wide correlation

in effect sizes caused by overlapping samples, differential z-scores were normalized using the SAS

procedure "proc stdize" with the "std" method. Subsequently, differential p-values were computed based

on the normalized differential z-scores. The cumulative distribution function of the standard normal

distribution ("pnorm") was used to calculate p-values from the normalized scores.

Transcriptome-Wide Association Analysis Using S-MetaXcan

Transcriptome-wide association analysis (TWAS) was conducted using S-MetaXcan[8], which infers gene

expression association with a phenotype based on its corresponding GWAS summary statistics and GTEx

gene expression datasets across 48 tissues. Four COVID-19 GWAS datasets from the HGI were analyzed,

including HGI-B1, HGI-B2, HGI-C2, and HGI-B1-vs-B2, alongside four long COVID GWAS datasets

with subtly distinct case-control definitions from HGI[2] and an additional set of 221 GWASs from the UK

Biobank (UKB) (all sample sizes >5,000) shared by Benneal's lab (https://www.nealelab.is/uk-biobank).

Reference transcriptomic data from 48 tissues in GTEx v7[9] were utilized for the analyses.

In detail, S-MetaXcan[8] employs multivariate regression to integrate evidence across different GTEx

tissues (n=48), enhancing detection power. Default settings were applied, with the conditional number for

eigenvalue set at 30 during singular value decomposition (SVD) truncation and genes on autosomal

chromosomes were included in the analysis. First, TWAS results for COVID-19 (n=8, collectively

referred to as COVID TWASs) were evaluated for genes proximal to prioritized long COVID-associated

SNPs (±500kb to the candidate SNP termed as tag SNP). Genes showing nominal significance (*P*<0.05)

in any of the COVID TWASs across the 48 GTEx tissues were prioritized for further investigation.

Also, to expand these findings, phenome-wide association analyses were conducted for these prioritized

genes. Associations were examined across a TWAS database aggregating 193 previously published TWAS

results[8] and 229 TWASs generated in this study, yielding a total of 418 TWAS datasets.

Statistical analysis software and availability of codes and data

Unless otherwise specified, all analyses were performed using SAS OnDemand for Academics (freely

available at https://www.sas.com/en_us/software/on-demand-for-academics.html). The scripts and

datasets used in this study are available from the corresponding author upon reasonable request.

Local Manhattan plots for the top candidate long COVID-associated SNPs and all supplementary tables

are publicly accessible at the GitHub link

(https://github.com/chengzhongshan/COVID19_GWAS_Analyzer/tree/main/LongCOVID_data_and_scri

pts/LongCOVID_SNP_LocalManhattanPlots).

Results

Prioritizing Genetic Variants Associated with Long COVID Using COVID-19 GWAS Data

To uncover long COVID risk loci, an integrative workflow was developed (Fig. 1a) by leveraging

publicly available COVID-19 GWAS data. Acute COVID-19 phenotypes, such as SARS-CoV-2

susceptibility, COVID-19 hospitalization, and non-hospitalization, were used as proxy traits for the long

COVID phenotype to prioritize genetic variants potentially linked to long COVID. The analysis utilized

data from the HGI release 7, including GWAS of hospitalized vs. non-hospitalized cases (HGI-B1),

hospitalized cases vs. the general population (HGI-B2), and SARS-CoV-2 infection vs. the general

population (HGI-C2). These GWASs comprise large sample sizes (16,512, 44,986, and 159,840 COVID-

19 cases for HGI-B1, HGI-B2, and HGI-C2, respectively; Fig. 1A) and exhibit minimal genomic inflation

(λ close to 1), as shown in supplementary Fig. S1 and Fig. 1A, loci near *SLC6A20* and *ABO* displayed

notable significance in association signals. These regions, previously validated as COVID-19 risk loci[10],

are well-established contributors to COVID-19 susceptibility and severity.

Categorization of Prioritized Long COVID Risk SNPs Related to COVID-19 Severity

To explore the genetic basis of long COVID, we used the integrative workflow shown in Fig. 1A to

prioritize 42 independent SNPs (see Table 1), categorized into two groups: "Severe COVID-19-Specific

SNPs" (n=27) and "SNPs Associated with Both Severe and Mild COVID-19" (n=15). The first category

includes SNPs linked predominantly to SARS-CoV-2 susceptibility and severe disease outcomes, such as

hospitalization, and are more prevalent in hospitalized cases than in non-hospitalized cases. The second

category consists of SNPs associated with both severe and mild COVID-19, showing consistent

associations with SARS-CoV-2 susceptibility and hospitalization, but no significant differentiation

between hospitalized and non-hospitalized cases. Of the 42 SNPs, except two SNPs, rs150345524 (*SF3B1*,

P=5.6E-8 in HGI-B2) and rs11766643 (*SLC22A5*, *P*=7.5E-8 in HGI-C2), all SNPs passed the

conventional genome-wide significance threshold (*P*<5E-8) in at least one HGI COVID-19 GWAS.

To reveal biological insights from SNP-gene associations, each of the above COVID-19-associated SNPs

was linked with its corresponding nearest gene to create a SNP-gene pair, providing an intuitive

framework for assessing their biological relevance in long COVID. However, proximity-based

assignments may overlook distal regulatory effects, which will be further addressed in subsequent

analyses. Notable SNPs in the "Severe COVID-19-Specific SNPs" category include rs10774679 (*OAS3*),

rs17169628 (*SLC22A31*), rs12329760 (*TMPRSS2*), rs12660421 (*FOXP4*), rs17078348 (*SLC6A20*),

rs190509934 (*ACE2*), rs2834164 (*IFNAR2*), rs3785884 (*MAPT*), rs7251000 (*DPP9*), and rs76608815

(*SLC5A3*), among others. These SNPs (n=27) show stronger associations with severe COVID-19 than

with mild cases. Among "SNPs Associated with Both Severe and Mild COVID-19", five out 15 SNP-

gene pairs, including rs2290859 (*NXPE3*), rs505922 (*ABO*), rs149533170 (*IFNA21*), rs34536443 (*TYK2*),

and rs9916158 (*GSDMA/B*), demonstrates consistent associations across GWAS datasets without

differential effects between hospitalized and non-hospitalized groups. Notably, *ABO* (rs505922) has been

reported as a genetic locus associated with long COVID[4]. Several immune-related genes were found close

to these SNPs, including *IFNA21* (rs149533170), *TYK2* (rs34536443), *GSDMA/B* (rs9916158), *BCL11A*

(rs1123573), *ELF5* (rs2924480), and *JAK1* (rs11208552), highlighting the role of immune dysregulation

in long COVID. Additionally, *FUT2* is found close to rs492602, and the protein of FUT2 encodes

fucosyltransferase 2, a key enzyme in the ABO antigen synthesis pathway, further underscores the

involvement of *ABO* in COVID-19 outcomes[11]. Additionally, four genes—*TAC4*, *NXPE3*, *NSF*, and

FOXG1—linked to neuronal functions were close to rs75432325, rs2290859, rs1378358, and

rs150497803, respectively, while three genes—*ZKSCAN1*, *BCL11A*, and *FOXG1*—identified as

transcription factors were linked to rs2897075, rs1123573, and rs150497803, respectively.

Evaluation of Odds Ratios for 42 Prioritized Long COVID Risk SNPs

To characterize the 42 candidate long COVID risk SNPs, we evaluated their odds ratios (ORs) across
three GWAS datasets: SARS-CoV-2 susceptibility (HGI-C2), COVID-19 hospitalization (HGI-B1 and
HGI-B2, with the first used mild COVID-19 and the latter employed general population as controls) (Fig.
1B). Distinct differences in OR distributions were observed between the aforementioned two long SNP
categories. SNPs in the "Severe COVID-19-Specific SNPs" category exhibited greater deviation in ORs
from 1 compared to that of the other long COVID SNP category. To quantify these differences, absolute
effect sizes (β) corresponding to each OR were used for statistical comparisons, as β values account for
both protective (OR<1) and risk (OR>1) effects in a unified framework. As shown in Supplementary Fig.
S2, SNPs classified as "Severe COVID-19-Specific SNPs" displayed a higher frequency of statistical
significance (P<0.05) in absolute effect sizes across GWAS datasets. These SNPs also demonstrated
significantly larger absolute effect sizes in comparisons between HGI-B1/HGI-B2 and HGI-C2,
underscoring their stronger association with COVID-19 hospitalization. Notably, only in HGI-B1 did the
absolute effect sizes differ significantly between the two SNP categories, a result reflecting the selection
criteria: severe COVID-19-specific SNPs required P<0.05 in both HGI-B1 and HGI-B2, whereas SNPs
associated with both severe and mild COVID-19 needed to be P<0.05 in HGI-B2 but not in HGI-B1.
These findings emphasize the expected differences stemming from category-specific filtering criteria
while highlighting the relevance of severe COVID-19-specific SNPs to hospitalization outcomes and their
potential role in long COVID.

Expanded Screening of Potential Causal Genes Linked to 42 Prioritized Long COVID Risk SNPs

To identify potential causal genes associated with 42 prioritized long COVID risk SNPs, we employed
local Manhattan plots, centering each candidate SNP within a 500-kb genomic window (Fig. 2 & 3).
Given that 41 of these SNPs, except rs12329760 (*TMPRSS2*), are located in non-coding regions, manual
evaluation was employed to prioritize genes proximal to each SNP with robust association signals. In total,
18 SNPs were prioritized based on their local Manhattan plots, representing two categories of long
COVID risk loci.

The first category of long COVID SNPs (n=27), nine SNPs were prioritized (Fig. 2): rs17078348
(*SLC6A20*), rs12660421 (*FOXP4*), rs12329760 (*TMPRSS2*), rs190509934 (*ACE2*), rs61860402 (*SFTPD*),
rs10774679 (*OAS3*), rs7251000 (*DPP9*), rs3785884 (*MAPT*), and rs2834164 (*IFNAR2*). Strong
association signals (P <1E-30) were observed for rs17078348 (*SLCA20*), with the locus encompassing
several genes, including *LZTFL1*, *CCR9*, *FYCO1*, *CCR1* and *XCR1*, previously implicated in COVID-19

severity and susceptibility. Similarly, rs12660421 (*FOXP4*) showed significant association signals near *FOXP4*, a known long COVID risk gene, with the SNP residing within its promoter region and relatively distant from other nearby genes. For rs12329760 (*TMPRSS2*), a missense variant, nominally significant associations with COVID-19 hospitalization (HGI-B2) and SARS-CoV-2 susceptibility (HGI-C2) were detected. Notably, nearby SNPs exhibited weaker signals (*P*<0.05) in HGI-B1 and differential effect sizes between HGI-B1 and HGI-B2, suggesting a complex genetic architecture. Several antiviral genes, including *MX1* and *MX2*, were identified near this locus. Other prioritized loci included rs61860402 (*SFTPD*) and rs7251000 (*DPP9*), two genes relevant to lung function, and rs10774679 (*OAS3*) and rs2834164 (*IFNAR2*), which are located within interferon-related gene clusters. While rs3785884 (*MAPT*) was initially implicated, its association may be driven by *CRHR1*, a gene associated with age-related neurodegenerative processes[12], aligning with the increased risk of severe COVID-19 and long COVID in older populations.

In terms of the second category of SNPs (n=15), nine loci were prioritized (Fig. 3), including rs149533170 (*IFNA21*), rs34536443 (*TYK2*), rs75432325 (*TAC4*), rs505922 (*ABO*), rs1120591 (*RAB2A*), rs3897075 (*ZKSCAN1*), rs9916158 (*GSDMA/B*), rs1378358 (*NSF*), and rs2924480 (*ELF5*). Among these, rs505922 (*ABO*) and rs34536443 (*TYK2*) displayed genome-wide significant associations with both hospitalized and non-hospitalized COVID-19, as evidenced by consistent signals in HGI-B2 and HGI-C2 datasets. The remaining SNPs were located in gene-dense regions, complicating the identification of specific causal genes. For instance, rs149533170 (*IFNA21*) is located near a cluster of interferon genes, including *IFNB1* and multiple *IFNAs*, which are central to interferon-mediated antiviral responses[13]. Similarly, rs9916158 (*GSDMA/B*) is proximal to genes implicated in asthma, while rs75432325 (*TAC4*) resides near *DLX3* and *DLX4*, which are involved in developmental pathways[14]. Less complex loci, such as rs1120591 (*RAB2A*) and rs2924480 (*ELF5*), contained fewer neighboring genes, facilitating clearer prioritization. *RAB2A* is an insulin regulator[15], while *ELF5* encodes a transcription factor relevant to epithelial differentiation and innate immune response[16].

Sex-Biased Associations with COVID-19 Hospitalization among Top Candidate SNPs

Epidemiological evidence indicates that women are more likely to experience long COVID but less likely to develop severe COVID-19[17,18], potentially due to stronger immune responses to SARS-CoV-2 infection in women. To investigate whether the top 42 candidate SNPs associated with long COVID exhibit sex-biased associations with COVID-19 hospitalization, we analyzed publicly available sex-stratified GWAS summary statistics from the GRASP COVID-19 database[19] generated based on UK Biobank samples.

The selected sex-stratified COVID-19 hospitalization GWASs (cases=1,343 and controls=262,886 for

females; cases=1,917 and controls=221,174 for males) includes mixed ancestry populations comparable

to the four HGI COVID-19 GWAS datasets used in our study. Using the delta-z-score method to compare

male and female GWAS results, SNPs with differential effect sizes for COVID-19 hospitalization were

identified (Fig. 4A). Genomic inflation factors for both sex-stratified GWASs and the sex-biased

differential GWAS were close to 1, indicating no evidence of inflation (Fig. 4B).

For these sex-stratified COVID-19 hospitalization GWASs, compact local Manhattan plots (Fig. 4C) and

forest plots (Fig. 4D) were generated for these 42 candidate SNPs. Due to the smaller sample sizes in the

sex-stratified GWASs, only 15 SNPs reached nominal significance ($P<0.05$) in at least one of the three

analyses. Of these, rs17078348 (*SLC6A20*), a well-established severe COVID-19 risk SNP, showed

nominal significance in both male and female cohorts. Five SNPs displayed nominal significance

exclusively in females, including rs75432325 (*TAC4*), rs505922 (*ABO*), rs12660421 (*FOXP4*), rs7251000

(*DPP9*), and rs11766643 (*TFR2*). Eight SNPs were nominally significant only in males, including

rs897075 (*ZKSCAN1*), rs117169628 (*SLC22A31*), rs7515509 (*AK5*), rs76608815 (*SLC5A3*), rs12585036

(*ATP11A*), rs5023077 (*FBRSL1*), rs10774679 (*IFNAR21*), and rs12329760 (*HLA-C*). None of the SNPs

passed the stringent multiple testing correction threshold ($P<6E-4$), and local Manhattan plots for 12 of

the 42 SNPs revealed relatively weaker association signals in the sex-stratified GWAS compared to the

stronger signals observed in HGI-B2 (Fig. 5). This reduction in signal strength likely reflects the smaller

sample sizes in the sex-stratified datasets, with reduced statistical power to detect sex-biased associations.

Finally, aggregating the absolute effect sizes of SNPs across the four groups by sex and nominal

association strength ($P<0.05$) revealed no significant differences (ANOVA $P=0.15$) among these groups.

In summary, while the sex-biased GWAS analysis did not identify SNPs with statistically significant sex-

biased associations with COVID-19 hospitalization, it highlighted 15 SNPs with nominal significance,

including rs505922 (*ABO*) and rs12660421 (*FOXP4*), which have been previously implicated as risk

variants for both severe COVID-19 and long COVID. These findings warrant further investigation in

larger sex-stratified COVID-19 and long COVID GWAS.

357 Evaluation of Previously Reported Long COVID-Associated SNPs

To assess whether the 42 prioritized long COVID SNPs identified in our study overlap with or are

proximal to those reported in two independent long COVID GWASs by Taylor et al.[3] and Chaudhary et

al.[4], a set of 83 SNPs derived from SNP network analyses and 3 SNPs identified via traditional GWAS

methods specifically from the two studies was created. These SNPs were analyzed for their association

signals in three HGI GWAS datasets (supplementary Table S1). Notably, rs12660421 (*FOXP4*), identified

by the HGI long COVID GWAS, was excluded from this evaluation as it displayed stronger associations

with general COVID-19 phenotypes in the HGI datasets analyzed here. This SNP was already included

among the 42 prioritized long COVID risk SNPs.

Our evaluation was guided by the hypothesis that true long COVID associations reported by Taylor et al.[3]

and Chaudhary et al.[4]. would exhibit at least nominal significance in one of the HGI GWAS datasets. The

larger sample sizes and higher statistical power of the HGI datasets enhance their ability to detect

associations with traits closely related to long COVID, such as severe COVID-19 or susceptibility to

SARS-CoV-2 infection. Moreover, overlap between our prioritized SNPs and those reported in previous

studies would strengthen the hypothesis that the loci in question are implicated in long COVID

pathogenesis.

Among the three SNPs reported by Chaudhary et al.[4] (Fig. 6), rs643434 (*ABO*) and rs2080090 (*BPTF*)

demonstrated nominally significant associations (P<0.05) with COVID-19 hospitalization (HGI-B2) and

susceptibility to COVID-19 (HGI-C2). rs643434 (*ABO*) surpassed genome-wide significance thresholds

in both HGI-B2 (P= 6.65E-24) and HGI-C2 (P=8.69E-85) and showed non-significant association in

HGI-B2 (P=0.6) but demonstrated significantly differential effect sizes (P=6E-5) between HGI-B1

(hospitalized vs. non-hospitalized COVID-19) and HGI-B2. This supported its classification as a

"category 2" long COVID SNP (associated with both severe and mild COVID-19 phenotypes). Similarly,

rs2080090 (*BPTF*) showed stronger associations in HGI-B2 (P=1.9E-4) than in HGI-C2 (P=2.5E-2), with

no significant association in HGI-B1. This SNP was also categorized as a suggestive "category 2" long

COVID SNP. In contrast, rs9273363 (*HLA-DQB1*) did not exhibit significant associations across any of

the HGI GWAS datasets analyzed (P>0.2). Nevertheless, among these 42 candidate SNPs, rs17219281

(*HLA-DQA1*), a "category 1" long COVID SNP that tends to be more likely associated with hospitalized

rather than mild COVID-19 (see Table 1), is ~50kb downstream of rs9273363 (*HLA-DQB1*), strongly

implicating that rs17219281 may be the real SNP associated with long COVID. Taken together, the three

recently published long COVID loci, including *ABO*, *BPTF*, and *HLA-DQB1*, were completely covered

by these 42 prioritized long COVID SNPs.

Meanwhile, of the 80 SNPs reported by Taylor et al.[3], 12 displayed nominally significant associations in

at least one HGI GWAS dataset (Supplementary Table S2 and Fig. 6). These included four SNPs

associated with HGI-B2, five with HGI-B1, and seven with HGI-C2. Notably, rs4303401 (*FRMD6*)

surpassed the multiple-testing correction threshold (P<1.5E-4), representing a strong candidate SNP.

Expanding the analysis to include a ±500kb genomic window around the reported SNPs identified

additional suggestive genome-wide associations in the HGI datasets. For example, in Fig. 6, rs62401842

(proximal to *KCTD16*, which shows brain-specific expression) and rs7671107 and rs1017716 (adjacent to

ANAPC4, a gene implicated in aging and strongly supported as eQTLs accordingly to GTEx eQTL

database) emerged as promising candidates. Other noteworthy SNPs in Fig. 6 include rs12602210 (*BPTF*)

and rs17412601 and rs2290859 (both mapped to *NXPE3*), which demonstrate suggestive or genome-wide

significance in at least one HGI GWAS dataset. In summary, four loci, including *KCTD16*, *ANAPC4*,

BPTF, and *NXPE3*, revealed by Taylor et al[3]. derived from SNP network analysis are independently

confirmed to be nominally associated with COVID-19 related phenotypes in the current study; among

them *BPTF* is a loci confirmed to be associated with long COVID by Chaudhary et al., Taylor et al., and

current study. Therefore, these findings underscore the potential of integrating data from diverse GWAS

to elucidate the genetic architecture of long COVID and highlight candidate loci for further investigation.

Potential Long COVID SNPs Specifically Identified in Non-Hospitalized COVID-19 Cases

To identify potential long COVID SNPs specific to non-hospitalized COVID-19 cases, we screened SNPs

that showed suggestive genome-wide association signals in HGI-B1 (hospitalized vs. non-hospitalized,

$P<$ 1E-5) and HGI-B1-vs-B2 (differential GWAS between HGI-B1 and HGI-B2, $P<0.05$) but were not

significant in HGI-B2 (hospitalized vs. general population, $P>0.05$) and were not in proximity to any

genome-wide significant SNPs in HGI-B2. This selection identified 20 independent SNPs potentially

associated with long COVID that specifically emerged from mild (non-hospitalized) COVID-19 cases

(Table 2; local Manhattan plots accessible at GitHub link as pointed out in the Method section; Fig. 7). All

20 SNPs showed suggestive genome-wide significant differences in effect sizes between HGI-B1 and

HGI-B2 ($P<1E-5$). Although none met the genome-wide significance threshold ($P<5E-8$), notable

findings were observed. For example, in Fig. 7, rs62401842 (*KCTD16*) exhibited the strongest association

in HGI-B1 ($P=8.53E-7$), while showing no association in HGI-B2 ($P=0.38$) and a significant difference

between HGI-B1 and HGI-B2 ($P=3.18E-7$). Gene *KCTD16*, encoding potassium channel tetramerization

domain-containing 16[20], is highly expressed in brain tissues according to the GTEx database. In addition,

eight SNPs illustrated in Fig. 7 were mapped to genes with brain-related functions, including rs9799354

(*NLGN1*), rs112842080 (*CPLX2*), rs61858037 (*NRG3*), rs61939166 (*KIF21A*), rs367777 (*NAV3*),

rs56143829 (*WASF3*), rs11454577 (*AKAP6*), and rs6049828 (*SYNDIG1*). Among these, *WASF3* has been

implicated in mitochondrial dysfunction and may mediate exercise intolerance in myalgic

encephalomyelitis/chronic fatigue syndrome[21]. Additionally, among these 20 candidate SNPs identified,

only three (rs62401842 [*KCTD16*], rs9799354 [*NLGN1*], and rs112842080 [*CPLX2*]) exhibit male-biased

associations with COVID-19 hospitalization at nominal significance levels (Fig. 8A, B). Furthermore, six

candidate SNPs, including rs62401842 (*KCTD16*), rs9799354 (*NLGN1*), rs112842080 (*CPLX2*),

rs56143829 (*WASF3*), rs4737438 (*PENK*), and rs6049828 (*SYNDIG1*), demonstrate sex-biased

association patterns within a genomic window ranging from 500kb to 1000kb (Fig. 8C-H). However, the

limited sample sizes in sex-stratified COVID-19 GWAS preclude any candidate SNPs from achieving

statistical significance after correction for multiple testing in analyses of sex-biased associations with

COVID-19 hospitalization. In summary, while no genome-wide significant SNPs specific to non-hospitalized COVID-19 cases were identified, these 20 SNPs provide suggestive evidence of association with long COVID in mild COVID-19.

Transcriptome-Wide Association Study Identifies Candidate Genes near Prioritized SNPs for COVID-19 and Long COVID Phenotypes

To elucidate genetic mechanisms underlying COVID-19 and long COVID phenotypes, we performed a transcriptome-wide association study (TWAS) using S-MetaXcan across 48 GTEx tissues. This approach utilized COVID-19 GWAS summary statistics from HGI: HGI-B1, HGI-B2, HGI-C2, and HGI-B1-vs-B2. The analysis targeted 62 candidate SNPs previously prioritized for their potential association with long COVID and aimed to identify their adjacent genes whose expression levels may be associated with COVID-19 hospitalization or susceptibility phenotypes, as well as long COVID.

Of the 62 candidate SNPs, two SNPs (rs190509934 near *ACE2* and rs9570861 near *OR7E156P*) not included in the TWAS analysis. The former resides on chromosome X, which S-MetaXcan excludes due to the focus of S-MetaXcan only on autosomal genes, while the latter SNP had no protein-coding genes within a 500-kb window. The remaining 60 SNPs are corresponded to 753 adjacent genes tested by these COVID TWASs (Fig. S5) with detectable expression across the 48 GTEx tissues.

To simplify interpretation, we retained only the strongest association (smallest p-value) for each gene across 48 GTEx tissues. This approach was justified by the hypothesis that specific tissues, rather than all tissues, are likely to drive the pathogenicity of COVID-19. Applying varying significance thresholds ($P<0.05$, 0.01, 0.001, 0.0001, 0.00001), the analysis revealed 422, 152, 43, 19, and 11 adjacent genes meeting these criteria in at least one of the four HGI datasets (Fig. S3, Table S2). Using a genome-wide significance threshold ($P<3.5E-7$), three loci displayed significant associations: (1) *DPP9*, showing the strongest association signal ($P=4.6E-18$) in adipose visceral omentum tissue; (2) *CXCR6* gene cluster, which included multiple genes (*CXCR6*, *CCR3*, *CCR1*, and *LZTFL1*), displaying the most significant association ($P=1.3E-10$) in nerve tibial tissue; (3) *MUC1* gene cluster, comprising *MUC1*, *THBS3* and *FAM189B*, highlighted in multiple GTEx tissues, including lung and colon. Additional loci with suggestive TWAS significance ($P<3.5E-6$) included *FYCO1*, *SLC6A20*, and *XCR1*, with specific associations to HGI datasets B1, B2, or C2 in distinct GTEx tissues.

To investigate overlap with long COVID, TWAS was extended to four long COVID GWASs using the same S-MetaXcan framework. Genome-wide evaluation of TWAS signals across the eight HGI GWAS datasets (Fig. S4) confirmed that the top hits with association $P<1E-5$ originated exclusively from COVID-19 hospitalization and susceptibility GWAS, with no contributions from long COVID GWAS

(Fig. S4). Notably, the candidate genes proximal to these prioritized SNPs accounted for most of the top signals across the eight HGI GWAS (Fig. 9), with the exception of *CCR5*, *EXOSC7*, *FLT1P1*, *RLN1*, *GBAP1*, *ARHGEF2*, *ZSCAN31*, *ZKSCAN3*, *PGMSP2*, *PAQR5*, and *TMIE*. Further in-depth evaluation of these outlier genes in long COVID GWAS revealed most of them lack of significance (Fig. S5), reinforcing the utility of prioritizing top candidate SNPs and their adjacent genes for downstream analyses. Interestingly, although the strongest associations identified in COVID-19 hospitalization or susceptibility GWASs (P<1E-5) were not genome-wide significant in long COVID datasets, four genes (*CCR1*, *DPP9*, *MUC1*, and *THBS3*; Fig. S5) showed nominally significant associations (P<0.05) in long COVID GWASs across specific tissues, such as colon transverse, testis, and heart atrial appendage. These associations were generally weaker than those observed on acute COVID-19 phenotypes, suggesting tissue-specific contributions to long COVID pathophysiology. In details, *DPP9* demonstrated strong associations in adipose-related tissues for COVID-19 hospitalization but weaker signals in colon transverse for long COVID. *THBS3* showed widespread associations across 19 GTEx tissues for COVID-19 hospitalization but exhibited weaker signals in long COVID, particularly in colon sigmoid and spleen. *MUC1* (adjacent to *THBS3*) and *CCR1* exhibited similar patterns, with weaker long COVID associations in colon transverse and heart atrial appendage, respectively. Furthermore, a relaxed threshold (P<0.05) identified additional adjacent genes associated with long COVID (Fig. 10 & Fig. S6-7). These included *OAS3* (rs10774679), *TMPRSS2* (rs12329760), *HLA-A* (rs4s9260038), *HLA-C* (rs1634761), *HLA-DQA1* (rs17219281), *WASF3* (rs65143829), *IFNAR2* (rs2834164), *SFTPD* (rs61860402), and *GSDMA/B* (rs9916158), the associations of which enriched in tissues such as heart, brain, colon, adipose, and muscle. Notably, the well-established *ABO* locus, a COVID-19 susceptibility locus, did not show significant TWAS signals. This suggests that its effects may stem from non-expression-related mechanisms, such as genetic mutations influencing blood types.

In summary, TWAS analyses identified 52 SNP-gene pairs nominally associated with long COVID across tissues. While genes such as *DPP9, CCR1*, *THBS3*, and *MUC1* emerged as potential shared contributors to acute and long COVID phenotypes, their tissue-specific associations suggest distinct mechanisms driving long COVID. These findings highlight the importance of integrative approaches in uncovering the molecular underpinnings of long COVID.

Phenome-Wide Analysis of Gene Expression Associations with Candidate Genes

To investigate gene expression associations with long COVID-related phenotypes, we performed a phenome-wide analysis of gene expression associations (Pheno-TWAS) analysis targeting 918 tested TWAS genes located within a 1Mb window of 62 candidate SNPs previously linked to COVID-19. Recognizing the limitations of published long COVID GWASs, which often rely on imprecisely defined

phenotypes, this analysis extended beyond prior long COVID studies by utilizing a manually curated TWAS database encompassing 418 phenotypes across 48 GTEx tissues. Of the 918 tested TWAS genes, 400 genes (43.6%), including 87 associated with *HLA* loci (e.g., *HLA-DQA2*, *HLA-A*, and *HLA-C*; Fig. S8-10), passed a stringent bonferroni-adjusted significance threshold (P<2.7E-9). Genes located near *HLA* loci emerged as top hits, demonstrating significant associations with immune-related disorders such as asthma, rheumatoid arthritis, psoriasis, coeliac disease, and systemic lupus erythematosus—conditions that may contribute to long COVID susceptibility. In addition to *HLA*-associated loci, other 313 genes across 44 non-*HLA* loci surpassed the genome-wide significance threshold in TWASs. Key findings include: (1) *IFNAR2*: strongly associated with COVID-19 hospitalization (P=2.6E-10; Fig. S11); (2) *GSDMB*: linked to asthma (P=2.7E-57; Fig. S12); (3) *BPTF*: associated with trunk fat percentage (P=1E-18; Fig. S13); (4) *CCDC171*: associated with body fat percentage (P=2E-35; Fig. S14). Interestingly, *FOXP4* expression, which did not exhibit significant associations with long COVID phenotypes in prior TWASs, demonstrated a suggestive association with sleep duration (P=1.1E-5; Fig. S15). This finding highlights the potential for phenome-wide approaches to uncover previously uncharacterized associations relevant to long COVID. In total, pheno-TWAS identified 45 candidate SNPs associated with genome-wide significant expression-linked phenotypes. These include three major *HLA* loci (*HLA-DQA2*, *HLA-A*, and *HLA-C*) as well as non-*HLA* loci such as *IFNAR2*, *BPTF*, *CCDC171*, and *GSDMB*. The implicated phenotypes—ranging from COVID-19 hospitalization and immune disorders to body composition traits and sleep duration—provide a broad spectrum of potential biological pathways contributing to long COVID pathophysiology. Full results, including nominally significant associations, are provided in Supplementary Table S3.

Discussion

Long COVID, characterized by over 200 symptoms spanning multiple organ systems, presents significant challenges for GWAS due to difficulties in assembling large, representative cohorts. To date, only four genome-wide significant loci—*FOXP4*, *ABO*, *HLA-DQA1*, and *BPTF*—have been identified for long COVID[2,4]. Our integrative analysis identifies 62 candidate loci potentially associated with long COVID, leveraging large-scale COVID-19 GWAS datasets from the HGI. This approach addresses the inherent limitations of long COVID-specific GWAS, such as limited sample sizes and phenotype heterogeneity, by utilizing proxy phenotypes like SARS-CoV-2 susceptibility and COVID-19 hospitalization. By categorizing genetic variants into three groups—those associated with hospitalized COVID-19, those linked to both hospitalized and non-hospitalized cases, and those specific to non-hospitalized cases—our findings provide a scalable framework to investigate the genetic underpinnings of long COVID.

The analysis highlights 42 genome-wide or suggestive loci, including well-documented variants such as

rs12660421 (*FOXP4*) and rs505922 (*ABO*), which have been implicated in severe COVID-19 outcomes

and long COVID susceptibility. Notably, additional loci like rs17219281 (*HLA-DQA1*), rs9260038 (*HLA-*

A), rs1634761 (*HLA-C*), rs7251000 (*DPP9*), rs9916158 (*GSDMB*), rs12602210 (*BPTF*), rs17078348

(*CCR1*), and rs41264915 (*THBS3*), rs190509934 (*ACE2*) and rs12329760 (*TMPRSS2*) were identified.

Furthermore, a unique category comprising 20 SNPs emerged specifically in non-hospitalized COVID-19

cases. These include rs62401842 (*KCTD16*) and rs56143829 (*WASF3*), mapped to genes enriched in

brain-related functions, underscoring the complexity of long COVID's etiology.

While this approach provides critical insights, certain limitations warrant consideration. First, the reliance

on proxy phenotypes may obscure SNPs specific to mild COVID-19 cases, given the skewed

representation of hospitalized patients in HGI datasets. Indeed, non-hospitalized-specific SNPs did not

reach genome-wide significance, likely reflecting this bias. Nevertheless, 20 suggestive SNPs, enriched

near genes implicated in neurocognitive and age-related pathways, merit further validation in independent

cohorts. Second, the TWAS conducted using S-MetaXcan are constrained by the limited tissue specificity

of available GTEx datasets and COVID related GWASs. Despite identifying 7 loci—*HLA-DQA1*, *HLA-A*,

HLA-C, *DPP9*, *CCR1*, *THBS3*, *GSDMB*—that emerged as promising TWAS hits for long COVID, the

potential absence of COVID-19-specific tissues, such as nasopharyngeal or immune-relevant sites, may

hinder the detection of tissue-specific effects pertinent to long COVID's heterogeneity.

Our findings underscore the utility of integrative approaches to prioritize genetic risk loci for complex

traits like long COVID. The discovery of loci such as *FOXP4*, *ABO*, and *HLA-DQA1*, as well as *BPTF*,

suggests shared biological pathways between acute COVID-19 severity and its chronic manifestations.

Additionally, the identification of brain-enriched loci in non-hospitalized cases aligns with emerging

evidence linking long COVID to neuroinflammatory and neurodegenerative processes. Future studies,

particularly those incorporating diverse populations and long-term follow-up data, are essential to refine

these associations and unravel the intricate mechanisms underlying long COVID.

Ethics approval and consent to participate

All data used in this study were freely shared by other researchers. Since no patient-identifiable

information was involved, ethical approval and informed consent were not applicable.

Consent for publication

This study is a secondary analysis of publicly available COVID-19 and long COVID-related GWAS data.
No individual-identifiable information, such as personal identifiers or images, was used; therefore,
consent for publication is not applicable.

Availability of data and material

Declaration of interests

The authors declare no competing interests.

Funding

No funding is available for supporting current study.

Authors' contributions

Z.C. conceived the study, analyzed the data, and wrote the manuscript.

Acknowledgements

I am thankful to all researchers who participated and contributed to the HGI and GRASP project, as
without their generosities in freely sharing these COVID-19 hospitalization GWASs and other COVID-
19/long COVID related GWASs as these were essential for the current finding.

592   **TABLES**

593   **Table 1. Prioritized long COVID SNPs based on integrative analysis of published COVID-19**
594   **GWASs**

Variant information						HGI-B1			HGI-B2			HGI-C2			HGI-B1-vs-B2	Long COVID
chr	pos	rsid	allele	AF	gene	beta	se	p	beta	se	p	beta	se	p	DiffPval	SNP groups
1	47275770	rs10890422	C	0.402	CYP4B1	-0.05	0.02	3.28E-03	-0.05	0.01	4.43E-08	-0.02	0.00	1.10E-04	6.63E-01	Hosp
1	65412830	rs11208552	T	0.647	JAK1	-0.01	0.02	6.95E-01	-0.05	0.01	2.19E-09	-0.02	0.00	4.21E-04	5.14E-03	Both
1	77949123	rs7515509	A	0.366	AK5	0.08	0.02	5.71E-06	0.06	0.01	2.94E-12	0.02	0.00	2.16E-05	1.82E-01	Hosp
1	155167786	rs41264915	G	0.103	THBS3	-0.10	0.03	3.59E-04	-0.13	0.01	9.33E-23	-0.02	0.01	1.13E-03	2.64E-01	Hosp
2	26833309	rs1275969	T	0.560	CIB4	0.04	0.02	2.37E-02	0.05	0.01	2.03E-08	0.01	0.00	9.18E-03	8.15E-01	Hosp
2	60707588	rs1123573	G	0.367	BCL11A	-0.03	0.02	1.86E-01	-0.07	0.01	1.32E-14	-0.03	0.00	4.62E-09	1.84E-02	Both
2	198245297	rs150345524	A	0.413	SF3B1	0.05	0.02	1.69E-02	0.05	0.01	5.56E-08	0.01	0.00	4.66E-02	9.52E-01	Hosp
3	45847241	rs17078348	G	0.101	SLC6A20	0.29	0.03	1.15E-19	0.34	0.01	2.53E-130	0.08	0.01	2.01E-36	6.34E-02	Hosp
3	101525625	rs2290859	T	0.341	NXPE3	-0.03	0.02	1.55E-01	-0.07	0.01	2.34E-13	-0.05	0.00	4.43E-31	2.02E-02	Both
4	25435159	rs7684660	C	0.829	ANAPC4	-0.06	0.03	1.98E-02	-0.08	0.01	8.71E-12	-0.02	0.00	2.86E-05	4.57E-01	Hosp
5	131776967	rs10066378	C	0.131	SLC22A5	0.06	0.02	1.80E-02	0.07	0.01	1.63E-10	0.02	0.01	1.00E-03	4.94E-01	Hosp
6	29907544	rs9260038	G	0.622	HLA-A	-0.04	0.02	2.72E-02	-0.06	0.01	1.94E-10	-0.01	0.00	1.30E-02	3.40E-01	Hosp
6	31274027	rs1634761	T	0.500	HLA-C	-0.04	0.02	1.81E-02	-0.06	0.01	4.50E-12	-0.02	0.00	9.99E-07	2.95E-01	Hosp
6	32675645	rs17219281	A	0.072	HLA-DQA1	-0.10	0.04	1.01E-02	-0.12	0.02	5.53E-11	-0.03	0.01	1.81E-03	5.30E-01	Hosp
6	41488378	rs12660421	A	0.041	FOXP4	0.26	0.04	2.88E-10	0.24	0.02	2.82E-31	0.08	0.01	7.36E-15	6.55E-01	Hosp
7	99630342	rs2897075	T	0.364	ZKSCAN1	0.01	0.02	7.21E-01	0.05	0.01	8.92E-09	0.02	0.00	6.17E-06	7.17E-03	Both
7	100220850	rs11766643	G	0.204	TFR2	-0.05	0.02	2.45E-02	-0.04	0.01	4.71E-04	-0.03	0.01	7.46E-08	4.92E-01	Hosp
8	61537523	rs1120591	C	0.393	RAB2A	0.02	0.02	2.75E-01	0.05	0.01	8.64E-10	0.02	0.00	2.48E-05	4.32E-02	Both
9	15795833	rs79611697	T	0.068	CCDC171	0.08	0.04	3.75E-02	0.10	0.01	3.57E-08	0.03	0.01	3.26E-03	4.72E-01	Hosp
9	21172825	rs149533170	A	0.009	IFNA21	0.06	0.11	6.03E-01	0.28	0.05	1.38E-08	0.08	0.03	7.64E-04	2.21E-02	Both
9	136149229	rs505922	T	0.348	ABO	0.01	0.02	6.16E-01	0.09	0.01	4.14E-24	0.08	0.00	7.93E-90	2.33E-06	Both
10	81723695	rs61860402	T	0.112	SFTPD	0.08	0.03	2.65E-03	0.08	0.01	1.38E-10	0.03	0.01	2.76E-08	9.61E-01	Hosp
11	1241221	rs35705950	T	0.100	MUC5B	-0.09	0.03	3.84E-03	-0.10	0.01	3.41E-12	-0.02	0.01	3.31E-03	6.00E-01	Hosp
11	34529831	rs2924480	C	0.399	ELF5	-0.03	0.02	1.67E-01	-0.08	0.01	1.63E-16	-0.02	0.00	1.59E-06	1.31E-02	Both
12	113374748	rs10774679	T	0.653	OAS3	0.04	0.02	2.31E-02	0.07	0.01	3.25E-16	0.03	0.00	4.75E-10	7.42E-02	Hosp
12	133141973	rs5023077	C	0.513	FBRSL1	-0.05	0.02	1.03E-02	-0.07	0.01	2.84E-15	-0.02	0.00	3.14E-07	1.79E-01	Hosp
13	113535741	rs12585036	T	0.209	ATP11A	0.07	0.02	4.58E-04	0.10	0.01	3.15E-21	0.02	0.01	2.85E-06	2.22E-01	Hosp
14	29497258	rs150497803	T	0.043	FOXG1	0.00	0.05	9.79E-01	-0.09	0.02	6.68E-05	-0.07	0.01	2.79E-10	3.11E-02	Both
16	89262657	rs117169628	A	0.135	SLC22A31	0.09	0.03	4.68E-04	0.10	0.01	1.37E-14	0.03	0.01	2.78E-05	7.97E-01	Hosp
17	38182229	rs9916158	T	0.365	GSDMA/B	-0.01	0.02	4.85E-01	0.03	0.01	2.05E-04	0.02	0.00	1.11E-08	5.88E-03	Both
17	44057595	rs3785884	A	0.187	MAPT	-0.05	0.02	2.22E-02	-0.09	0.01	3.84E-16	-0.02	0.01	3.51E-04	6.69E-02	Hosp
17	44787312	rs1378358	T	0.167	NSF	-0.04	0.02	9.81E-02	-0.08	0.01	1.18E-13	-0.02	0.01	2.58E-03	2.75E-02	Both
17	47950329	rs75432325	C	0.042	TAC4	0.08	0.04	5.82E-02	0.18	0.02	2.24E-19	0.05	0.01	7.00E-07	2.04E-02	Both
19	4714468	rs7251000	T	0.432	DPP9	-0.07	0.02	3.98E-04	-0.08	0.01	1.17E-19	-0.03	0.00	1.18E-11	4.61E-01	Hosp
19	9009161	rs12976386	T	0.552	MUC16	0.04	0.03	1.14E-01	-0.03	0.01	9.37E-03	-0.03	0.01	3.75E-08	2.43E-03	Both
19	10463118	rs34536443	C	0.043	TYK2	0.05	0.06	3.58E-01	0.23	0.02	1.74E-21	0.05	0.01	1.29E-05	4.66E-04	Both
19	49206417	rs492602	G	0.460	FUT2	-0.02	0.02	1.69E-01	-0.06	0.01	1.71E-11	-0.04	0.00	3.44E-18	4.00E-02	Both
19	50882619	rs1405655	C	0.332	NR1H2	0.07	0.02	6.03E-05	0.08	0.01	5.92E-19	0.03	0.00	1.20E-09	7.95E-01	Hosp
21	34621948	rs2834164	C	0.480	IFNAR2	-0.10	0.02	1.51E-08	-0.11	0.01	4.14E-41	-0.03	0.00	6.54E-14	4.83E-01	Hosp
21	35353264	rs76608815	T	0.090	SLC5A3	0.11	0.03	9.64E-05	0.13	0.01	6.60E-21	0.04	0.01	1.49E-07	4.83E-01	Hosp
21	42852497	rs12329760	T	0.258	TMPRSS2	-0.07	0.02	8.60E-04	-0.06	0.01	2.16E-08	-0.01	0.00	5.32E-03	4.55E-01	Hosp
23	15620340	rs190509934	C	0.003	ACE2	-0.46	0.20	2.12E-02	-0.44	0.10	1.08E-05	-0.36	0.04	1.70E-18	9.38E-01	Hosp

Note: HGI: the COVID-19 host genetics initiative; chr: chromosome; pos: hg38 genomic position; allele:
alternative allele; AF: allele frequency for alternative allele; diffPval: statistical significance *P* for
differential effect size of each common SNP by comparing the two HGI GWASs, including HGI-B1 and
HGI-B2; Hosp: long COVID SNPs tend to be specifically emerged in hospitalized COVID-19 cases; Both:
long COVID SNPs tend to be emerged in both hospitalized and non-hospitalized COVID-19 cases.
Association Ps passed the suggestive genome-wide association significance threshold of P < 1E-7 are
highlighted in light red color, and differential effect size significance P < 0.05 is colored with light green.

Note: HGI-C2 (SASR-CoV2 susceptibility), HGI-B1 (hospitalized vs. non-hospitalized COVID-19), and

HGI-B2 (hospitalized vs. general population).

Table 2. Potential long COVID risk SNPs emerged specifically from non-hospitalized COVID-19

cases

Variant information						HGI-B1			HGI-B2			HGI-C2			HGI-B1-vs-B2
chr	pos	rsid	allele	AF	gene	beta	se	p	beta	se	p	beta	se	p	DiffPval
3	174147743	rs9799354	C	0.556	NLGN1	0.08	0.02	5.38E-06	0.01	0.01	2.71E-01	-0.01	0.00	1.79E-02	6.21E-06
5	128241667	rs201484359	G	0.304	SLC27A6	0.13	0.03	8.55E-06	0.02	0.01	1.25E-01	-0.01	0.01	8.91E-02	2.58E-05
5	143904835	rs62401842	A	0.042	KCTD16	0.27	0.06	8.53E-07	0.02	0.03	3.82E-01	0.00	0.01	8.72E-01	3.18E-07
5	175248251	rs112842080	T	0.021	CPLX2	-0.36	0.07	1.58E-06	-0.04	0.03	2.53E-01	0.03	0.02	6.69E-02	1.25E-06
6	37086436	rs147171940	A	0.038	PIM1	-0.19	0.04	5.96E-06	-0.04	0.02	1.33E-01	0.01	0.01	5.74E-01	5.92E-05
6	151444801	rs76392050	T	0.010	MTHFD1L	0.70	0.16	6.10E-06	0.10	0.06	6.16E-02	0.03	0.03	3.84E-01	5.88E-06
8	57438048	rs4737438	A	0.378	PENK	0.08	0.02	2.97E-06	0.01	0.01	2.79E-01	0.00	0.00	5.30E-01	2.98E-06
8	68035003	rs148340257	T	0.065	CSPP1	-0.22	0.05	8.05E-06	-0.03	0.02	1.77E-01	0.00	0.01	8.82E-01	7.21E-06
9	123344199	rs10760104	T	0.819	CDK5RAP2	-0.13	0.03	2.37E-06	-0.01	0.01	4.57E-01	0.00	0.01	8.77E-01	5.36E-07
10	54976003	rs55654117	C	0.272	MBL2	0.10	0.02	1.85E-06	0.01	0.01	2.61E-01	0.00	0.00	4.50E-01	1.13E-06
10	84981211	rs61858037	T	0.091	NRG3	0.15	0.03	9.21E-06	0.02	0.02	1.62E-01	0.00	0.01	9.58E-01	2.31E-05
12	39502966	rs61939166	G	0.011	KIF21A	-0.40	0.09	3.60E-06	-0.03	0.04	5.36E-01	-0.02	0.02	3.63E-01	1.10E-06
12	78552292	rs367777	T	0.358	NAV3	-0.10	0.02	3.13E-06	-0.01	0.01	2.77E-01	0.01	0.00	2.39E-01	1.89E-06
13	27062460	rs56143829	T	0.025	WASF3	0.24	0.05	1.26E-06	0.01	0.03	6.03E-01	0.00	0.01	7.56E-01	3.23E-07
13	63988259	rs9570861	A	0.552	PCDH20	0.08	0.02	5.51E-06	0.02	0.01	1.07E-01	-0.01	0.00	4.06E-02	4.83E-05
14	32937383	rs11454577	G	0.589	AKAP6	-0.09	0.02	5.54E-06	-0.01	0.01	3.22E-01	0.00	0.01	6.24E-01	7.71E-06
16	84887921	rs113067468	G	0.030	CRISPLD2	-0.30	0.07	4.48E-06	-0.05	0.03	1.20E-01	0.00	0.02	8.21E-01	2.18E-05
17	3454928	rs115744301	G	0.035	TRPV3	0.66	0.15	5.21E-06	0.02	0.05	6.89E-01	-0.03	0.03	2.52E-01	2.53E-07
17	13775336	rs11869231	T	0.310	COX10	-0.11	0.02	2.38E-06	0.00	0.01	6.52E-01	0.00	0.00	9.85E-01	2.26E-07
20	24627964	rs6049828	C	0.567	SYNDIG1	0.09	0.02	7.13E-06	0.01	0.01	2.65E-01	0.00	0.00	3.51E-01	5.43E-06

Note: abbreviations in the table headers are similar to that in Table 1.

SUPPLEMENTARY TABLES

Table S1. Association signals of published long COVID SNPs across 4 HGI COVID-19 GWASs

Table S2. Adjacent genes of these 62 candidate SNPs that pass the nominal significance threshold in

the transcriptome-wide association studies (TWAS) of COVID-19 hospitalization and susceptibility

Table S3. Adjacent genes of these 62 candidate SNPs that pass the nominal significance threshold in

the phenome-wide transcriptome association studies (pheno-TWASs)

FIGURES

Figure 1. Workflow to prioritize genetic single nucleotide polymorphisms (SNPs) potentially
associated with long COVID

(A) 42 top prioritized long COVID risk SNPs, with 27 SNPs associated specifically with hospitalization
risk and 15 showing common risk patterns across both hospitalized and non-hospitalized cases. These
SNPs were selected based on a significance threshold of $P \leq$ 1E-7 in at least one of the GWAS studies
(HGI-C2, HGI-B2, or HGI-B1). Each SNP is centered within a 1Mb window in local Manhattan plots to
highlight the genomic context, including known COVID-19 and long COVID risk loci. Key genes near
these SNPs, including *FOXP4*—a locus previously implicated in long COVID risk—and other notable
COVID-19-associated genes (*ACE2*, *OAS3*, *JAK1*, *IFNA2*, *DDP9*, and *ABO*), are annotated to emphasize
their potential relevance to SARS-CoV-2 pathology and long COVID risk. (B) Forest plot of 42 SNPs
identified from COVID-19 GWAS on susceptibility (HGI-C2) and hospitalization (HGI-B1 and HGI-B2),
showing suggestive (P<1E-7 or genome-wide significant associations (P<5E-8). Allele distributions are
compared between hospitalized and non-hospitalized cases (HGI-B1). SNPs are grouped as follows: (1)
hospitalization-specific, genome-wide significant (Hosp-specific, P<5E-8); (2) common across cases,
genome-wide significant (Common, P<5E-8); (3) hospitalization-specific, suggestive or non-significant
(hosp-specific, P = (5E-8, 1]); (4) common across cases, suggestive or non-significant (Common, P =
(5E-8, 1]). Colored dots represent odds ratios (ORs) with 95% confidence intervals (CIs) as error bars.
ORs with CIs overlapping the vertical line (OR = 1) are non-significant.

Figure 2. Local Manhattan plots for potential long COVID risk SNPs more likely associated with
hospitalized than non-hospitalized COVID-19

Figure 3. Local Manhattan plots for potential long COVID risk SNPs likely associated with both
hospitalized and non-hospitalized COVID-19

Figure 4. Evaluating top potential long COVID risk SNPs in sex-biased COVID-19 hospitalization
GWAS from UK Biobank

(A-B) Manhattan plots and QQ plots for sex-stratified COVID-19 hospitalization GWASs, as well as the
differential effect size GWAS between females and males. (C) Local Manhattan plot for these 42 SNPs
separated into two groups, including long COVID SNPs that tend to be specific in hospitalized cases and
that more frequent in both hospitalized and non-hospitalized cases. All association signals shown in the
figure are within a 100Mb genomic window for each potential long COVID risk SNP residing at the

center. Genes close to these 42 SNPs are added into the upper track. (C) Forest plots for these 42 potential

long COVID risk SNPs in sex-stratified COVID-19 hospitalization GWASs, as well as the differential

effect size GWAS between females and males. Odd Ratio (OR) and the corresponding 95% confidence

intervals (CI) of these SNPs are illustrated, with SNPs showing nominal significant associations among

female or male COVID-19 patients are colored in dark red and tagged by '*'. Note that there are 8 and 6

SNPs showing sex-biased associations with females and males, respectively, meanwhile, only one SNP

rs17078348 mapped to *SLC6A20* displaying nominally significant association signals in both sexes. The

genomic position of rs190509934 is used to collect all other SNPs around it, as it is not included in the

two UKB sex stratified COVID-19 GWASs.

Figure 5. Local Manhattan plots for potential long COVID SNPs displaying suggestively sex-biased
association with COVID-19 hospitalization
661

Figure 6. Evaluating published long COVID SNPs in the current HGI COVID-19 GWASs

(A) Forest plots for these 15 SNPs in currently studied COVID-19 GWASs (from left to right: HGI-B1,

HGI-B2, and HGI-C2), with genes closed to each of them listed on the right-y-axis. Note: beside 3 long

COVID risk SNPs reported by Chaudhary et al., other 12 long COVID SNPs were obtained by screening

83 previously reported long COVID SNPs by Taylor et al. (see Table S1) that show nominal significant

association in at least one of the 3 current HGI COVID-19 GWASs. (B-G) Local Manhattan plots

illustrate published long COVID SNPs approximate to these top candidate long COVID SNPs identified

in current study with each candidate SNP is located in the center of each genomic window; GWASs

labels from bottom to top in each panel are HGI-B1, HGI-B2, HGI-B1-vs-B2, and HGI-C2, respectively.

Figure 7. Local Manhattan plots for potential long COVID risk SNPs specifically associated with
non-hospitalized COVID-19
673
Figure 8. Evaluation of potential long COVID risk SNPs associated with non-hospitalized COVID-
19 in sex-biased COVID-19 hospitalization GWAS from UK Biobank

(A) Local Manhattan plots for candidate long COVID-associated SNPs identified exclusively in the non-

hospitalized COVID-19 group from sex-stratified COVID-19 hospitalization GWAS (the differential

effect size GWAS between females and males at the top, female GWAS in the middle, and male GWAS

shown at the bottom). (B) Forest plots displaying the odds ratios (OR) and corresponding 95% confidence

intervals (CI) for these SNPs in non-hospitalized COVID-19 patients from the sex-stratified GWAS (male

results at the top and female results at the bottom). SNPs with nominally significant associations in either

female or male groups are highlighted in dark red and marked with an asterisk (*). Notably, three SNPs

exhibit male-biased associations with COVID-19 hospitalization. (C-H) Detailed local Manhattan plots
for six prioritized SNPs identified specifically in the non-hospitalized COVID-19 group within the sex-
stratified GWAS.

Figure 9. Evaluation of top genes adjacent to prioritized long COVID risk SNPs (n=62) in
transcriptome-wide association studies (TWAS) of COVID-19 susceptibility, hospitalization, and
long COVID

(A) Bar plot showing the top genes adjacent to prioritized long COVID risk SNPs based on
transcriptome-wide association studies (TWAS) for COVID-19 susceptibility and hospitalization. Genes
are categorized by the GTEx tissue in which they display the most significant association. Besides
rs1078348, other two SNPs have multiple genes with the smallest TWAS *P* are highlighted. (B) Heatmap
depicting the significance (*P*) of gene-tissue associations for prioritized long COVID SNP-adjacent genes
across 48 GTEx tissues. Each cell represents the *P* for a specific gene in a given tissue, with a color
gradient indicating statistical significance: gray (*P*>0.05), yellow (*P*<0.05), and red (*P*<1E-6). Genes are
sorted by TWAS, and GTEx tissues are clustered to highlight patterns of tissue specificity.

Figure 10. Top genes adjacent to prioritized long COVID risk SNPs (n=23) in long COVID-19
transcriptome-wide association studies (TWASs)

Bar plots illustrating the best *P* for genes adjacent to prioritized long COVID risk SNPs (n=23) in 4 long
COVID TWASs across 48 GTEx tissues.

701
702
703

SUPPLEMENTARY FIGURES

Figure S1. QQ plots for four COVID-19 GWASs from HGI

Figure S2. Differential effect size analysis for candidate long COVID SNPs by SNP categories and GWAS

Figure S3. Total number of genes adjacent to prioritized long COVID risk SNPs (n=62) in COVID-19 transcriptome-wide association studies (TWASs)

Bar plots illustrating the number of genes adjacent to prioritized long COVID risk SNPs (n=62) as well as number of TWASs out of 4 COVID-19 related GWAS across 48 GTEx tissues passed different association P thresholds. Note that the 62 SNPs are colored according to its corresponding long COVID SNP category as depicted in Figure 1.

Figure S4. Transcriptome-wide association studies (TWAS) for COVID-19 susceptibility, hospitalization, and long COVID

(A) Top genome-wide significant TWAS-identified genes associated with COVID-19 susceptibility and hospitalization, annotated by their best significant tissue expression association (based on 48 GTEx tissues). No genes reached genome-wide significance in TWAS analyses of long COVID-related GWASs. (B) Manhattan plots depicting TWAS results for COVID-19 susceptibility and hospitalization (HGI datasets), as well as long COVID GWASs derived from four distinct case-control designs. Each dot represents a gene, with its genomic position along the x-axis and -log10(P) on the y-axis.

Figure S5. Evaluation of top genes identified from transcriptome-wide association studies (TWAS) of COVID-19 susceptibility and hospitalization in corresponding long COVID-related TWASs

Heatmap displaying the significance (P) of top genes identified in TWAS analyses of COVID-19 susceptibility and hospitalization across multiple tissues in long COVID-related TWASs. Each cell represents the P for the association of a gene with long COVID in a specific tissue from GTEx database. The color gradient denotes statistical significance: blue ($P>0.05$), yellow ($P<0.05$), and red ($P<1E-6$). Rows are sorted by genes, and columns represent tissues, allowing visualization of tissue-specific patterns of gene expression relevance.

Figure S6. Other top genes adjacent to prioritized long COVID risk SNPs (n=29) in long COVID-19 transcriptome-wide association studies (TWASs)

Bar plots illustrating the best *P* for genes adjacent to prioritized long COVID risk SNPs (n=29) in 4 long

COVID TWASs across 48 GTEx tissues.

Figure S7. Evaluation of top genes adjacent to prioritized long COVID risk SNPs (n=62) in

transcriptome-wide association studies (TWAS) of long COVID across 48 GTEx tissues

(A) Bar plot showing the number of SNP-gene pairs passed the nominal TWAS association threshold in

GTEx tissues. Note that SNP-gene pairs are defined by combining prioritized long COVID risk SNPs

with its adjacent genes. (B) Heatmap depicting the significance (*P*) of SNP-gene pairs across 48 GTEx

tissues in long COVID TWASs. Each cell represents the *P* for a specific gene in a given tissue, with a

color gradient indicating statistical significance: white (*P*>0.05), yellow (*P*<0.05), and red (*P*<1E-6).

Genes are sorted by SNP-gene pairs and TWAS names.

Figure S8. Evaluation of genes adjacent to prioritized long COVID risk SNP rs17219281 (*HLA-

DQA1) **in phenome-transcriptome-wide association studies (pheno-TWAS) across 48 GTEx tissues**

To the sake of clarity, heatmap only depicting the significance of SNP-gene pairs with *P*<1E-60 across 48

GTEx tissues in TWASs. Each cell represents the *P* for a specific gene in a given tissue, with blank cell

indicating the P not passed the threshold of *P*<1E-60.

Figure S9. Evaluation of genes adjacent to prioritized long COVID risk SNP rs9260038 (*HLA-A*) in

phenome-transcriptome-wide association studies (pheno-TWAS) across 48 GTEx tissues

To the sake of clarity, similar heatmap as that in Figure S8 only depicting the significance of SNP-gene

pairs with *P*<1E-60 across 48 GTEx tissues in TWASs.

Figure S10. Evaluation of genes adjacent to prioritized long COVID risk SNP rs1634761 (*HLA-C*)

in phenome-transcriptome-wide association studies (pheno-TWAS) across 48 GTEx tissues

To the sake of clarity, similar heatmap as that in Figure S8 only depicting the significance of SNP-gene

pairs with *P*<1E-60 across 48 GTEx tissues in TWASs.

Figure S11. Evaluation of genes adjacent to prioritized long COVID risk SNP rs2834164 (*IFNAR2*)

in phenome-transcriptome-wide association studies (pheno-TWAS) across 48 GTEx tissues

To the sake of clarity, the color gradient of heatmap denotes statistical significance: blue (P>0.05), yellow

(P<0.05), and red (P<1E-6). Rows are sorted by genes, and columns represent TWAS, allowing

visualization of tissue-specific patterns of gene expression relevance.across 48 GTEx tissues in TWASs.

Figure S12. Evaluation of genes adjacent to prioritized long COVID risk SNP rs9916158

(*GSDMA/B*) in phenome-transcriptome-wide association studies (pheno-TWAS) across 48 GTEx

tissues

Heatmap legend as that in Figure S11.

Figure S13. Evaluation of genes adjacent to prioritized long COVID risk SNP rs12602210 (*BPTF*)

in phenome-transcriptome-wide association studies (pheno-TWAS) across 48 GTEx tissues

Heatmap legend as that in Figure S11.

Figure S14. Evaluation of genes adjacent to prioritized long COVID risk SNP rs79611697

(*CCDC171*) in phenome-transcriptome-wide association studies (pheno-TWAS) across 48 GTEx

tissues

Heatmap legend as that in Figure S11.

Figure S15. Evaluation of genes adjacent to prioritized long COVID risk SNP rs12660421 (*FOXP4*)

in phenome-transcriptome-wide association studies (pheno-TWAS) across 48 GTEx tissues

Heatmap legend as that in Figure S11.

References

1. Ely, E.W., Brown, L.M., Fineberg, H.V., National Academies of Sciences, E., and Medicine Committee on Examining the Working Definition for Long, C. (2024). Long Covid Defined. N Engl J Med *391*, 1746-1753. 10.1056/NEJMsb2408466.

2. Lammi, V., Nakanishi, T., Jones, S.E., Andrews, S.J., Karjalainen, J., Cortés, B., O'Brien, H.E., Fulton-Howard, B.E., Haapaniemi, H.H., Schmidt, A., et al. (2023). Genome-wide Association Study of Long COVID. medRxiv, 2023.2006.2029.23292056. 10.1101/2023.06.29.23292056.

3. Taylor, K., Pearson, M., Das, S., Sardell, J., Chocian, K., and Gardner, S. (2023). Genetic risk factors for severe and fatigue dominant long COVID and commonalities with ME/CFS identified by combinatorial analysis. Journal of translational medicine *21*, 775. 10.1186/s12967-023-04588-4.

4. Chaudhary, N.S., Weldon, C.H., Nandakumar, P., Team, a.R., Holmes, M.V., and Aslibekyan, S. (2024). Multi-ancestry GWAS of Long COVID identifies immune-related loci and etiological links to chronic fatigue syndrome, fibromyalgia and depression. medRxiv, 2024.2010.2007.24315052. 10.1101/2024.10.07.24315052.

5. Cai, M., Xie, Y., Topol, E.J., and Al-Aly, Z. (2024). Three-year outcomes of post-acute sequelae of COVID-19. Nat Med *30*, 1564-1573. 10.1038/s41591-024-02987-8.

6. Zhang, K., Lin, S., Luo, Y.S., and Cheng, Z. (2024). Protocol to search for genetic factors related to severe COVID-19 by analyzing publicly available genome-wide association studies. STAR protocols *5*, 103028. 10.1016/j.xpro.2024.103028.

7. Cheng, Z., Cai, Y., Zhang, K., Zhang, J., Gui, H., Luo, Y.S., Zhou, J., and DeVeale, B. (2023). MAP3K19 regulatory variation in populations with African ancestry may increase COVID-19 severity. iScience *26*, 107555. 10.1016/j.isci.2023.107555.

8. Barbeira, A.N., Dickinson, S.P., Bonazzola, R., Zheng, J., Wheeler, H.E., Torres, J.M., Torstenson, E.S., Shah, K.P., Garcia, T., Edwards, T.L., et al. (2018). Exploring the phenotypic consequences of tissue specific gene expression variation inferred from GWAS summary statistics. Nat Commun *9*, 1825. 10.1038/s41467-018-03621-1.

9. GTEx-Consortium (2020). The GTEx Consortium atlas of genetic regulatory effects across human tissues. Science *369*, 1318-1330. 10.1126/science.aaz1776.

10. Severe-COVID-GWAS-Group, Ellinghaus, D., Degenhardt, F., Bujanda, L., Buti, M., Albillos, A., Invernizzi, P., Fernandez, J., Prati, D., Baselli, G., et al. (2020). Genomewide Association Study of Severe Covid-19 with Respiratory Failure. N Engl J Med *383*, 1522-1534. 10.1056/NEJMoa2020283.

11. Galeev, A., Suwandi, A., Cepic, A., Basu, M., Baines, J.F., and Grassl, G.A. (2021). The role of the blood group-related glycosyltransferases FUT2 and B4GALNT2 in susceptibility to infectious disease. International journal of medical microbiology : IJMM *311*, 151487. 10.1016/j.ijmm.2021.151487.

12. Polanczyk, G., Caspi, A., Williams, B., Price, T.S., Danese, A., Sugden, K., Uher, R., Poulton, R., and Moffitt, T.E. (2009). Protective effect of CRHR1 gene variants on the development of adult depression following childhood maltreatment: replication and extension. Archives of general psychiatry *66*, 978-985. 10.1001/archgenpsychiatry.2009.114.

13. Samuel, C.E. (2001). Antiviral actions of interferons. Clinical microbiology reviews *14*, 778-809, table of contents. 10.1128/CMR.14.4.778-809.2001.

14. Tamaoki, N., Takahashi, K., Aoki, H., Iida, K., Kawaguchi, T., Hatakeyama, D., Inden, M., Chosa, N., Ishisaki, A., Kunisada, T., et al. (2014). The homeobox gene DLX4 promotes generation of human induced pluripotent stem cells. Sci Rep *4*, 7283. 10.1038/srep07283.

15. Sugawara, T., Kano, F., and Murata, M. (2014). Rab2A is a pivotal switch protein that promotes either secretion or ER-associated degradation of (pro)insulin in insulin-secreting cells. Scientific Reports *4*, 6952. 10.1038/srep06952.

16. Pietzner, M., Chua, R.L., and Wheeler, E. (2022). ELF5 is a potential respiratory epithelial cell-specific risk gene for severe COVID-19. *13*, 4484. 10.1038/s41467-022-31999-6.

17. Bai, F., Tomasoni, D., Falcinella, C., Barbanotti, D., Castoldi, R., Mulè, G., Augello, M., Mondatore, D., Allegrini, M., Cona, A., et al. (2022). Female gender is associated with long COVID syndrome: a prospective

cohort study. Clinical microbiology and infection : the official publication of the European Society of
Clinical Microbiology and Infectious Diseases *28*, 611.e619-611.e616. 10.1016/j.cmi.2021.11.002.

18.   Neuhouser, M.L., Butt, H.I., Hu, C., Shadyab, A.H., Garcia, L., Follis, S., Mouton, C., Harris, H.R., Wactawski-
Wende, J., Gower, E.W., et al. (2024). Risk factors for long COVID syndrome in postmenopausal women
with previously reported diagnosis of COVID-19. Annals of epidemiology *98*, 36-43.
10.1016/j.annepidem.2024.08.003.

19.   Thibord, F., Chan, M.V., Chen, M.H., and Johnson, A.D. (2022). A year of Covid-19 GWAS results from the
GRASP portal reveals potential genetic risk factors. HGG Adv, 100095. 10.1016/j.xhgg.2022.100095.

20.   Zuo, H., Glaaser, I., Zhao, Y., Kurinov, I., Mosyak, L., Wang, H., Liu, J., Park, J., Frangaj, A., Sturchler, E., et
al. (2019). Structural basis for auxiliary subunit KCTD16 regulation of the GABA(B) receptor. Proc Natl Acad
Sci U S A *116*, 8370-8379. 10.1073/pnas.1903024116.

21.   Wang, P.Y., Ma, J., Kim, Y.C., Son, A.Y., Syed, A.M., Liu, C., Mori, M.P., Huffstutler, R.D., Stolinski, J.L.,
Talagala, S.L., et al. (2023). WASF3 disrupts mitochondrial respiration and may mediate exercise
intolerance in myalgic encephalomyelitis/chronic fatigue syndrome. Proc Natl Acad Sci U S A *120*,
e2302738120. 10.1073/pnas.2302738120.

Figure 1

SARS-CoV2 infected vs. general population
(HGI-C2: 159,840 vs. 2,782,977)

P ≤ 0.05

Differential GWAS (HGI-B1 vs. HGI-B2)

P > 0.05 *P ≤ 0.05* *P ≤ 0.05*

Hospitalized COVID-19 vs. general population
(HGI-B2: 44,986 vs. 2,356,386)

P ≤ 0.05 *P > 0.05* *P ≤ 0.05*

Hospitalized vs. non-hospitalized COVID-19
(HGI-B1: 16,512 vs. 71,321)

P ≤ 0.05 *P ≤ 0.05* *P > 0.05*

Independent SNPs with any $P <= 1\times10^{-7}$ with same effect size directions among HGI-C2 and HGI-B2

Independent SNPs (MAF>0.01) with any $P <= 1\times10^{-5}$ and no other adjacent genome-wide significant SNPs in either HGI-B2 or HGI-C2

Independent SNPs with any $P <= 1\times10^{-7}$ with consistent effect size directions between HGI-C2 and HGI-B2

Prioritize each top hit among 4 HGI-GWASs by manually evaluating local Manhattan plots

Category 1: Severe COVID-19-Specific SNPs (n=27)

Category 3: Mild COVID-19-Specific SNPs (n=20)

Category 2: SNPs Associated with Both Severe and Mild COVID-19 (n=15)

Determine association signals for these 62 candidate SNPs in sex-stratified COVID-19 GWASs and long COVID GWASs

Check genes close to these candidate SNPs in transcriptome-wide association studies (TWASs) of COVID-19 hospitalization/susceptibility and long COVID GWASs across 48 GTEx tissues

Evaluate genes adjacent to these candidate SNPs in phenotype-wide transcription association studies (Pheno-TWASs) across 48 GTEx tissues, covering a broad set of 420 phenotypes

Figure 2

Figure 3

Figure 4

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

Figure S6

Figure S7

Figure S8

Long COVID association signals for SNPs in the SNP category "Severe COVID-19-Specific SNPs"

Figure S9

Figure S10

Figure S11

Figure S12

Figure S13

Figure S14

Figure S15

Figure S16

Only combinations of genes and tissues with at least one TWAS association P < 0.01 are illustrated in the heatmap.

P<0.05   P<10⁻⁶

Figure S17

rs17219281..HLA-DQA2..BRD2......Brain Putamen basal ganglia..........
rs17219281..HLA-DQA2..BTNL2.....Nerve Tibial.......................
rs17219281..HLA-DQA2..C6orf10...Testis.............................
rs17219281..HLA-DQA2..COL11A2...Adrenal Gland......................
rs17219281..HLA-DQA2..COL11A2...Vagina.............................
rs17219281..HLA-DQA2..HLA-DMA...Heart Left Ventricle...............
rs17219281..HLA-DQA2..HLA-DMA...Liver..............................
rs17219281..HLA-DQA2..HLA-DMA...Skin Sun Exposed Lower leg..........
rs17219281..HLA-DQA2..HLA-DMB...Cells Transformed fibroblasts.......
rs17219281..HLA-DQA2..HLA-DMB...Pancreas...........................
rs17219281..HLA-DQA2..HLA-DPA1..Testis.............................
rs17219281..HLA-DQA2..HLA-DPA1..Thyroid............................
rs17219281..HLA-DQA2..HLA-DPA1..Whole Blood........................
rs17219281..HLA-DQA2..HLA-DPB1..Brain Cerebellum...................
rs17219281..HLA-DQA2..HLA-DQA1..Adipose Visceral Omentum...........
rs17219281..HLA-DQA2..HLA-DQA1..Colon Transverse...................
rs17219281..HLA-DQA2..HLA-DQA1..Liver..............................
rs17219281..HLA-DQA2..HLA-DQA1..Muscle Skeletal....................
rs17219281..HLA-DQA2..HLA-DQA2..Brain Hippocampus..................
rs17219281..HLA-DQA2..HLA-DQA2..Colon Sigmoid......................
rs17219281..HLA-DQA2..HLA-DQA2..Pancreas...........................
rs17219281..HLA-DQA2..HLA-DQA2..Prostate...........................
rs17219281..HLA-DQA2..HLA-DQB1..Adipose Subcutaneous...............
rs17219281..HLA-DQA2..HLA-DQB1..Brain Anterior cingulate cortex BA24.
rs17219281..HLA-DQA2..HLA-DQB1..Muscle Skeletal....................
rs17219281..HLA-DQA2..HLA-DQB1..Pancreas...........................
rs17219281..HLA-DQA2..HLA-DQB1..Prostate...........................
rs17219281..HLA-DQA2..HLA-DQB1..Small Intestine Terminal Ileum......
rs17219281..HLA-DQA2..HLA-DQB2..Artery Aorta.......................
rs17219281..HLA-DQA2..HLA-DQB2..Cells EBV-transformed lymphocytes...
rs17219281..HLA-DQA2..HLA-DQB2..Heart Atrial Appendage.............
rs17219281..HLA-DQA2..HLA-DQB2..Small Intestine Terminal Ileum......
rs17219281..HLA-DQA2..HLA-DQB2..Testis.............................
rs17219281..HLA-DQA2..HLA-DRA...Thyroid............................
rs17219281..HLA-DQA2..HLA-DRB1..Brain Anterior cingulate cortex BA24.
rs17219281..HLA-DQA2..HLA-DRB1..Muscle Skeletal....................
rs17219281..HLA-DQA2..HLA-DRB1..Ovary..............................
rs17219281..HLA-DQA2..HLA-DRB1..Whole Blood........................
rs17219281..HLA-DQA2..HLA-DRB5..Adipose Subcutaneous...............
rs17219281..HLA-DQA2..NOTCH4....Adipose Visceral Omentum...........
rs17219281..HLA-DQA2..PSMB8.....Ovary..............................
rs17219281..HLA-DQA2..PSMB8.....Stomach............................
rs17219281..HLA-DQA2..PSMB9.....Adipose Visceral Omentum...........
rs17219281..HLA-DQA2..TAP1......Cells EBV-transformed lymphocytes...
rs17219281..HLA-DQA2..TAP1......Testis.............................
rs17219281..HLA-DQA2..TAP2......Skin Not Sun Exposed Suprapubic.....
rs17219281..HLA-DQA2..TAP2......Skin Sun Exposed Lower leg..........
rs17219281..HLA-DQA2..TAP2......Testis.............................

Column labels:
UKB Malabsorption or coeliac disease
RA OKADA TRANS ETHNIC
RA OKADA EUR
IMMUNOBASE Systemic lupus erythematosus hg19
RA OKADA ASIAN
UKB Hypothyroidism or myxoedema
UKB Asthma
IMMUNOBASE Multiple sclerosis hg19
RA GWASmeta2 20090505

$P < 1\times10^{-60}$

Figure S18

rs9260038..HLA-A..GABBR1...Adipose Subcutaneous.................
rs9260038..HLA-A..GABBR1...Breast Mammary Tissue................
rs9260038..HLA-A..GABBR1...Nerve Tibial..........................
rs9260038..HLA-A..HLA-A....Esophagus Mucosa.....................
rs9260038..HLA-A..HLA-A....Ovary................................
rs9260038..HLA-A..HLA-A....Skin Not Sun Exposed Suprapubic.......
rs9260038..HLA-A..HLA-A....Skin Sun Exposed Lower leg............
rs9260038..HLA-A..HLA-A....Whole Blood...........................
rs9260038..HLA-A..HLA-F....Brain Cerebellar Hemisphere...........
rs9260038..HLA-A..HLA-F....Brain Cerebellum......................
rs9260038..HLA-A..HLA-F....Colon Sigmoid.........................
rs9260038..HLA-A..HLA-F....Esophagus Gastroesophageal Junction...
rs9260038..HLA-A..HLA-F....Liver.................................
rs9260038..HLA-A..HLA-F....Ovary.................................
rs9260038..HLA-A..HLA-F....Pancreas..............................
rs9260038..HLA-A..HLA-G....Brain Cerebellar Hemisphere...........
rs9260038..HLA-A..HLA-G....Brain Cortex..........................
rs9260038..HLA-A..HLA-G....Colon Transverse......................
rs9260038..HLA-A..HLA-G....Lung..................................
rs9260038..HLA-A..HLA-G....Testis................................
rs9260038..HLA-A..MAS1L....Stomach...............................
rs9260038..HLA-A..MOG......Artery Tibial.........................
rs9260038..HLA-A..MOG......Esophagus Gastroesophageal Junction...
rs9260038..HLA-A..MOG......Heart Left Ventricle..................
rs9260038..HLA-A..OR2H2....Adipose Subcutaneous..................
rs9260038..HLA-A..OR2H2....Heart Left Ventricle..................
rs9260038..HLA-A..OR2H2....Nerve Tibial..........................
rs9260038..HLA-A..OR2H2....Skin Sun Exposed Lower leg............
rs9260038..HLA-A..PPP1R11..Brain Cerebellum......................
rs9260038..HLA-A..PPP1R11..Cells Transformed fibroblasts.........
rs9260038..HLA-A..PPP1R11..Whole Blood...........................
rs9260038..HLA-A..RNF39....Brain Hippocampus.....................
rs9260038..HLA-A..RNF39....Brain Nucleus accumbens basal ganglia.
rs9260038..HLA-A..RNF39....Lung..................................
rs9260038..HLA-A..RNF39....Stomach...............................
rs9260038..HLA-A..TRIM10...Skin Sun Exposed Lower leg............
rs9260038..HLA-A..TRIM15...Colon Transverse......................
rs9260038..HLA-A..TRIM15...Skin Sun Exposed Lower leg............
rs9260038..HLA-A..TRIM26...Adipose Visceral Omentum..............
rs9260038..HLA-A..TRIM26...Artery Tibial.........................
rs9260038..HLA-A..TRIM26...Brain Caudate basal ganglia...........
rs9260038..HLA-A..TRIM26...Brain Cerebellum......................
rs9260038..HLA-A..TRIM26...Nerve Tibial..........................
rs9260038..HLA-A..TRIM26...Testis................................
rs9260038..HLA-A..TRIM31...Adrenal Gland.........................
rs9260038..HLA-A..TRIM31...Lung..................................
rs9260038..HLA-A..TRIM31...Pancreas..............................
rs9260038..HLA-A..TRIM31...Testis................................
rs9260038..HLA-A..TRIM39...Adipose Subcutaneous..................
rs9260038..HLA-A..TRIM39...Adrenal Gland.........................
rs9260038..HLA-A..TRIM39...Esophagus Mucosa......................
rs9260038..HLA-A..TRIM39...Nerve Tibial..........................
rs9260038..HLA-A..TRIM40...Testis................................
rs9260038..HLA-A..ZFP57....Heart Atrial Appendage................
rs9260038..HLA-A..ZFP57....Prostate..............................
rs9260038..HLA-A..ZFP57....Spleen................................
rs9260038..HLA-A..ZNRD1....Artery Tibial.........................
rs9260038..HLA-A..ZNRD1....Esophagus Gastroesophageal Junction...
rs9260038..HLA-A..ZNRD1....Thyroid...............................
rs9260038..HLA-A..ZNRD1....Whole Blood...........................

Column labels:
IMMUNOBASE Multiple sclerosis hg19
UKB Psoriasis
IMMUNOBASE Systemic lupus erythematosus hg19
RA OKADA ASIAN
UKB Diastolic blood pressure automated reading
MAGIC Manning et al FastingGlucose Interaction
UKB Waist circumference
UKB Sitting height
pgc.scz2
UKB Ankylosing spondylitis
UKB Sarcoidosis
UKB Hypothyroidism or myxoedema
UKB Hyperthyroidism aka thyrotoxicosis
UKB Standing height
RA GWASmeta2 20090505
RA OKADA EUR
RA OKADA TRANS ETHNIC
UKB Malabsorption or coeliac disease

$P < 1\times10^{-60}$

Figure S19

rs1634761..HLA-C..ABHD16A...Cells EBV-transformed lymphocytes.....
rs1634761..HLA-C..ABHD16A...Nerve Tibial.........................
rs1634761..HLA-C..AIF1......Breast Mammary Tissue.................
rs1634761..HLA-C..APOM......Prostate..............................
rs1634761..HLA-C..APOM......Stomach...............................
rs1634761..HLA-C..APOM......Testis................................
rs1634761..HLA-C..BAG6......Brain Cortex..........................
rs1634761..HLA-C..BAG6......Brain Frontal Cortex BA9..............
rs1634761..HLA-C..BAG6......Brain Hypothalamus....................
rs1634761..HLA-C..C6orf15...Skin Sun Exposed Lower leg............
rs1634761..HLA-C..C6orf25...Breast Mammary Tissue.................
rs1634761..HLA-C..C6orf47...Thyroid...............................
rs1634761..HLA-C..CCHCR1....Artery Aorta..........................
rs1634761..HLA-C..CCHCR1....Brain Cerebellar Hemisphere...........
rs1634761..HLA-C..CLIC1.....Brain Cortex..........................
rs1634761..HLA-C..CLIC1.....Liver.................................
rs1634761..HLA-C..CLIC1.....Spleen................................
rs1634761..HLA-C..CSNK2B....Brain Nucleus accumbens basal ganglia.
rs1634761..HLA-C..CSNK2B....Stomach...............................
rs1634761..HLA-C..DDAH2.....Brain Cerebellum......................
rs1634761..HLA-C..DDAH2.....Brain Frontal Cortex BA9..............
rs1634761..HLA-C..DDR1......Prostate..............................
rs1634761..HLA-C..DDX39B....Brain Anterior cingulate cortex BA24..
rs1634761..HLA-C..GPANK1....Adipose Subcutaneous..................
rs1634761..HLA-C..GPANK1....Muscle Skeletal.......................
rs1634761..HLA-C..HCG27.....Skin Sun Exposed Lower leg............
rs1634761..HLA-C..HCG27.....Testis................................
rs1634761..HLA-C..HLA-B.....Pancreas..............................
rs1634761..HLA-C..HLA-C.....Spleen................................
rs1634761..HLA-C..LSM2......Artery Aorta..........................
rs1634761..HLA-C..LST1......Brain Hypothalamus....................
rs1634761..HLA-C..LST1......Cells EBV-transformed lymphocytes.....
rs1634761..HLA-C..LTB.......Esophagus Gastroesophageal Junction...
rs1634761..HLA-C..LY6G5C....Adipose Visceral Omentum..............
rs1634761..HLA-C..LY6G6C....Brain Nucleus accumbens basal ganglia.
rs1634761..HLA-C..LY6G6C....Testis................................
rs1634761..HLA-C..LY6G6D....Brain Anterior cingulate cortex BA24..
rs1634761..HLA-C..LY6G6F....Esophagus Mucosa......................
rs1634761..HLA-C..LY6G6F....Testis................................
rs1634761..HLA-C..MICA......Brain Putamen basal ganglia...........
rs1634761..HLA-C..MICA......Prostate..............................
rs1634761..HLA-C..MICA......Stomach...............................
rs1634761..HLA-C..MICB......Artery Aorta..........................
rs1634761..HLA-C..MICB......Spleen................................
rs1634761..HLA-C..MSH5......Testis................................
rs1634761..HLA-C..MUC21.....Brain Cerebellum......................
rs1634761..HLA-C..MUC22.....Esophagus Mucosa......................
rs1634761..HLA-C..NFKBIL1...Brain Cerebellum......................
rs1634761..HLA-C..POU5F1....Stomach...............................
rs1634761..HLA-C..POU5F1....Whole Blood...........................
rs1634761..HLA-C..PRRC2A....Whole Blood...........................
rs1634761..HLA-C..PSORS1C1..Brain Cortex..........................
rs1634761..HLA-C..PSORS1C2..Vagina................................
rs1634761..HLA-C..TCF19.....Colon Transverse......................
rs1634761..HLA-C..TCF19.....Prostate..............................
rs1634761..HLA-C..TNF.......Brain Nucleus accumbens basal ganglia.
rs1634761..HLA-C..VARS2.....Brain Frontal Cortex BA9..............
rs1634761..HLA-C..VARS2.....Lung..................................
rs1634761..HLA-C..VARS2.....Pancreas..............................
rs1634761..HLA-C..VARS......Vagina................................
rs1634761..HLA-C..VWA7......Nerve Tibial..........................

Column labels:
IMMUNOBASE Multiple sclerosis hg19
RA GWASmeta2 20090505
UKB Basal metabolic rate
UKB Asthma
UKB Hypothyroidism or myxoedema
RA OKADA ASIAN
AdvancedAMD 2015
UKB Hyperthyroidism aka thyrotoxicosis
UKB Ankylosing spondylitis
RA OKADA EUR
RA OKADA TRANS ETHNIC
IMMUNOBASE Systemic lupus erythematosus hg19
UKB Malabsorption or coeliac disease

$P < 1 \times 10^{-60}$

Figure S20

Figure S21

Figure S22

Figure S23

rs79611697..CCDC171..C9orf92..Adipose Subcutaneous.................
rs79611697..CCDC171..C9orf92..Ovary.................................
rs79611697..CCDC171..CCDC171..Artery Aorta.........................
rs79611697..CCDC171..CCDC171..Brain Anterior cingulate cortex BA24.
rs79611697..CCDC171..CCDC171..Brain Caudate basal ganglia..........
rs79611697..CCDC171..CCDC171..Brain Cerebellar Hemisphere..........
rs79611697..CCDC171..CCDC171..Brain Cerebellum.....................
rs79611697..CCDC171..CCDC171..Brain Cortex.........................
rs79611697..CCDC171..CCDC171..Brain Putamen basal ganglia..........
rs79611697..CCDC171..CCDC171..Cells EBV-transformed lymphocytes.....
rs79611697..CCDC171..CCDC171..Colon Transverse.....................
rs79611697..CCDC171..CCDC171..Esophagus Mucosa.....................
rs79611697..CCDC171..CCDC171..Heart Atrial Appendage...............
rs79611697..CCDC171..CCDC171..Lung.................................
rs79611697..CCDC171..CCDC171..Skin Not Sun Exposed Suprapubic......
rs79611697..CCDC171..CCDC171..Skin Sun Exposed Lower leg...........
rs79611697..CCDC171..CCDC171..Thyroid..............................
rs79611697..CCDC171..PSIP1....Brain Anterior cingulate cortex BA24.
rs79611697..CCDC171..PSIP1....Breast Mammary Tissue................
rs79611697..CCDC171..PSIP1....Testis...............................
rs79611697..CCDC171..SNAPC3...Adipose Subcutaneous.................
rs79611697..CCDC171..SNAPC3...Artery Aorta.........................
rs79611697..CCDC171..SNAPC3...Brain Cerebellar Hemisphere..........
rs79611697..CCDC171..SNAPC3...Brain Hypothalamus...................
rs79611697..CCDC171..SNAPC3...Cells Transformed fibroblasts........
rs79611697..CCDC171..SNAPC3...Colon Sigmoid........................
rs79611697..CCDC171..SNAPC3...Heart Atrial Appendage...............
rs79611697..CCDC171..SNAPC3...Nerve Tibial.........................
rs79611697..CCDC171..SNAPC3...Skin Not Sun Exposed Suprapubic......
rs79611697..CCDC171..SNAPC3...Thyroid..............................

$P < 0.05$

$P > 0.05$ $P < 1 \times 10^{-6}$

Figure S24