Association of genetic risk and physical activity with incident type 2 diabetes

Xuan Zhou[1], Germán D. Carrasquilla[1], Malene R. Christiansen[1], Roelof A.J. Smit[1,2], Lars

Ängquist[1], Torben Hansen[1], Ruth J.F. Loos[1,2], Niels Grarup[1], Allan Linneberg[1,3,4], Tuomas O.

Kilpeläinen[1], Jordi Merino[1,5]

Affiliations:

[1] Novo Nordisk Foundation Center for Basic Metabolic Research, University of Copenhagen,

Copenhagen, Denmark,

[2] Charles Bronfman Institute for Personalized Medicine, Icahn School of Medicine at Mount

Sinai, New York, NY, USA

[3] Center for Clinical Research and Prevention, Copenhagen University Hospital - Bispebjerg and

Frederiksberg, The Capital Region, Copenhagen, Denmark

[4] Department of Clinical Medicine, Faculty of Health and Medical Sciences, University of

Copenhagen, Copenhagen, Denmark

[5] Diabetes Unit and Center for Genomic Medicine, Massachusetts General Hospital, Boston,

MA, USA.

Running title: Gene-physical activity interaction in diabetes

Word count: 3,648 words

Corresponding authors' contact information:

Jordi Merino, Ph.D. Novo Nordisk Foundation Center for Basic Metabolic Research, University

of Copenhagen, 665 Blegdamsvej 3B, DK-2200 Copenhagen, Denmark. Phone: +45 51 42 40 55

Email: jordi.merino@sund.ku.dk

Abstract

Objective: To assess whether daily step counts and genetic risk interact to influence the risk of developing type 2 diabetes.

Research Design and Methods: We analyzed data from 9,501 participants in the *All of Us* Research Program with both genetic and wearable device–derived physical activity data and without diabetes at baseline. Physical activity was quantified using daily step counts. Genetic risk was assessed using a global polygenic score. Incident type 2 diabetes was identified using electronic health record–linked diagnostic codes. Multivariable Cox proportional hazards models estimated hazard ratios (HRs) for type 2 diabetes across genetic risk and physical activity levels. We tested for additive interaction using the relative excess risk due to interaction (RERI). In secondary analyses we used physical-activity intensity measures using wearable-derived and self-reported intensity levels.

Results: Type 2 diabetes incidence rates ranged from 4.1 per 1,000 person-years (95% CI, 2.5–5.7) in individuals with high physical activity and low genetic risk to 18.4 (95% CI, 15.2–21.6) in those with low physical activity and high genetic risk (HR, 6.2 (95% CI: 3.97, 9.6)). A significant additive interaction was observed (RERI, 0.20; 95% CI, 0.04–0.36; P = .007), with 15% (95% CI, 2–27) of excess risk attributed to the interaction. Similar interaction patterns were found using device-based intensity metrics and self-reported physical activity measures.

Conclusions: These findings provide evidence of additive interactions between genetic risk and physical activity, underscoring the potential value of integrating genomic and device-derived data to identify individuals who would more likely benefit from increasing physical activity.

48   **Graphical abstract**

Article Highlights

Why did we undertake this study?

Despite longstanding interest in gene–environment interactions influencing the development of type 2 diabetes, identifying such interactions in human populations remains challenging, partially due to difficulties in accurately measuring environmental exposures. Step count is an objective, interpretable measure of daily ambulatory physical activity that offers a novel opportunity to evaluate these interactions. To date, no studies have specifically investigated how daily step counts may interact with genetic susceptibility to influence type 2 diabetes risk.

What is the specific question(s) we wanted to answer?

To what extent do genetic risk and physical activity, measured via daily step count, independently and jointly contribute to the risk of developing type 2 diabetes?

What did we find?

We observed a gradient of increasing type 2 diabetes risk associated with high genetic susceptibility and low step count. This gradient deviated from additivity, with the combination of high genetic risk and low physical activity accounting for substantial proportion of excess type 2 diabetes risk. The evidence of additive interactions between genetic risk and physical activity was replicated in an independent cohort using self-reported physical activity data and longer-term follow-up.

77 **What are the implications of our findings?**

78 Our results highlight the joint impact of genetic risk and physical inactivity on type 2 diabetes

79 development. These findings support the integration of genetic and behavioral risk factors in risk

80 stratification and prevention strategies and may inform targeted public health interventions aimed

81 at reducing the global burden of type 2 diabetes.

INTRODUCTION

The risk of developing type 2 diabetes varies considerably among individuals, reflecting differences in susceptibility to the often collectively called environmental factors (1,2). This variability underpins the significance of gene-environment interaction, wherein an individual's genetic profile, together with its downstream molecular effectors, interact with environmental exposures to uniquely modulate disease risk (3–6). While gene-environment interactions are pervasive in nature, shaping phenotypes such as plant growth (7), bird migration (8), and grasshopper adaptation to new environments (9), identifying such interactions in the context of human traits and diseases remains challenging (1), with often conflicting findings in the literature (10–13). This challenge is partly due to the limited precision and depth of environmental exposure assessments, particularly when self-reported (14,15).

Step count provides a simple and interpretable metric of daily ambulatory activity, offering advantages over self-reported physical activity measures or less intuitive accelerometer-derived intensity classifications (16). Prior studies have shown that higher step counts are associated with lower type 2 diabetes risk (17,18), and that accelerometer-measured intensity-specific physical activity may attenuate type 2 diabetes genetic risk (19). However, previous studies of gene–physical activity interactions have generally relied on single time-point physical activity assessments and have not accounted for the dynamic nature of physical activity over time. Moreover, no prior study has directly investigated how daily step counts interact with genetic risk on the development of type 2 diabetes.

In this study, we analyzed longitudinal data from a population-based cohort in the United States, comprising 9,501 adults with objectively measured physical activity over time, to test the

105   hypothesis that physical activity, measured via daily step count, interacts with genetic risk in the

106   development of type 2 diabetes. We further evaluated whether similar patterns were observed

107   using additional activity measures, including device-based intensity levels and self-reported

108   physical activity.

METHODS

Study design and population

We used data collected from participants in the *All of Us* Research Program. Details on the design and execution of the *All of Us* have been published previously (20,21). In this study, we used Curated Data Repository (CDR) version 7 for Controlled Tier (C2022Q4R9). This dataset included information on physical measurements and vital signs collected at enrollment, surveys, electronic health record (EHR), Fitbit data, and genetic information from participants enrolled between May 6, 2018, and July 1, 2022. Our analyses focused on participants who owned a Fitbit and agreed to share their Fitbit, EHR, and genetic information. We excluded participants who did not wear a Fitbit for at least 6 months. Only the authorized authors who completed *All of Us* Research Program Responsible Conduct of Research Training accessed the deidentified data from the Researcher Workbench (a secured cloud-based platform). Since the authors were not directly involved with the participants, institutional review board review was exempted in compliance with *All of Us* Research Program policy.

In a secondary analysis, we used data from Inter99 (21). Inter99 is a population-based pre-randomized lifestyle intervention study aiming to prevent ischemic heart disease and type 2 diabetes. Baseline data were collected from March 1999 until January 2001 and there were 6,784 participants. Follow-up was conducted after one, three, five, ten, and twenty years for a health examination, completion of questionnaires and risk assessment.

Physical activity assessment

Activity tracking data for this study came from the Bring Your Own Device program that

allowed individuals who already owned a tracking device (Fitbit, Inc) to consent to link their

activity data with other data in the *All of Us* cohort. By registering their personal device on the

patient portal, participants could share all activity data collected since the creation of their

personal device account. We used mean daily steps calculated on a monthly basis for each

participant. Consistent with prior physical activity data curation approaches (17), days with less

than 10 hours of wear time, less than 100 steps, or greater than 45 000 steps or for which the

participant was younger than 18 years were removed. In addition, we removed months with

fewer than 15 valid days of monitoring. To capture additional dimensions of physical activity,

we derived device-based measures of intensity, including daily lightly active minutes, moderate

active minutes, and very active minutes []. We also analyzed self-reported physical activity data

from the Inter99 study, assessed via validated questionnaires covering work and leisure-time

activity (22), at baseline and at one-, three-, and five-year follow-ups. Reported activity levels

were converted into minutes per week by summing responses based on a five-day working week.

Type 2 diabetes polygenic scores

We generated a global polygenic score for type 2 diabetes that captures overall genetic burden

using genetic variants and weights identified in a previous study that did not include participants

from this study (23). We followed standard procedures to filter genetic data to include only

variants that had passed initial quality control (24). We used PRS–continuous shrinkage (PRScs)

(25) to infer posterior SNP effect sizes under continuous shrinkage priors, with the scaling

parameter auto-estimated (25). Scores were computed with ancestry-specific variants and

weights.

Ascertainment of type 2 diabetes

Incident type diabetes cases were identified using any incident billing code in EHR. We

excluded any new diagnoses coded during the first six months of monitoring, assuming that such

conditions were likely prevalent, but not yet recognized clinically. The EHR data from different

participating sites were mapped and harmonized using the Observational Medical Outcomes

Partnership common data model (26). In Inter99, we used data from separate nationwide health

registries to define type 2 diabetes events.

Statistical analysis

A flow diagram describes how many participants were excluded based on the criteria used to

create the analytical dataset (Supplementary Figure 1). Descriptive statistics for participant's

demographic and clinical characteristics were presented by median and IQR for continuous

variables and frequency for categorical variables.

We used Cox proportional hazard models to examine associations of genetic risk and step counts

with type 2 diabetes risk. For these analyses, we censored cases that occurred 6 months after

completing enrolment to mitigate potential bias due to time-varying confounding. Person-time

for each participant was calculated from the six months after completing baseline assessment to

the diagnosis of type 2 diabetes, death, loss to follow-up, or the end of the follow-up period,

whichever came first. We modelled polygenic scores and step counts as continuous variables.

We averaged daily step counts within each month and subsequently treated step counts as a time-

varying covariate in the Cox models to account for month-to-month variations in physical

activity during the follow-up period. We also classified participants according to categories of

genetic risk and physical activity (nine categories based on thirds of genetic risk and step counts,

with low genetic risk and high physical activity as reference) and conducted stratified analyses.

We adjusted the multivariable models for age (in years, continuous), sex (male, female) and

genetic ancestry inferred from the first 10 principal components of genomic relationship matrix.

We verified the proportional hazards assumption of the Cox model by using the Schoenfeld

residuals technique (p values for the global test > 0.1). Analyses were conducted for complete

cases given the small number of people with missing covariates. In secondary analyses, we

further adjusted our models for BMI (quintiles of kg/m^2). The rationale for not including BMI in

the main model was because BMI can act as a collider or a mediator (27).

We tested for additive and multiplicative interactions of genetic risk and physical activity with

type 2 diabetes incidence. For additive interactions we assessed the relative excess risk due to

interaction (RERI) and further examined the decomposition of the joint effect—that is, the

population attributable proportions to genetic risk alone, physical activity alone, and to their

interaction(28). For these analyses, we used the following formula (RERI = RR11 - RR10 -

RR01 + 1) and (attributable proportion (AP) = RERI/ RR11) (28). CIs for each of the interaction

measures were calculated using the delta method described by Hosmer and Lemeshow. We

tested for multiplicative interactions using the log-likelihood ratio test to compare the goodness

of fit of a multivariable-adjusted model with and without the cross-product interaction term(29).

For these analyses, we considered genetic risk and physical activity as continuous variables and

modeled the effect per 1SD.

In secondary interaction analyses, we derived device-based measures of physical activity

intensity, including daily lightly active minutes, fairly active minutes, and very active minutes.

These intensity levels were defined using metabolic equivalents (METs): 1.5–3.0 METs for

lightly active, 3.0–6.0 METs for fairly active, and >6.0 METs for very active minutes (30,31).

We also incorporated total self-reported physical activity data from the Inter99 study,

categorizing participants as having high (7–12 h/week) or low (0–2 h/week) activity. Additional

analyses were conducted stratifying participants by ethnic group (European, African, East Asian,

and South Asian) and by sex.

Two-sided tests were used to assess whether polygenic risk for T2D and physical activity are

associated with T2D incidence. A one-sided test was used to assess whether RERI is greater than

0, i.e., testing the hypothesis that the combined effects of polygenic risk for T2D and physical

activity exceeds their individual effects. All statistical analyses were performed using R

software, version 4.0.3 (R Foundation). We followed the Better Precision-data Reporting of

Evidence from Clinical Intervention Studies & Epidemiology (BePRECISE) reporting guideline

(32).

RESULTS

Of 14,200 participants with available Fitbit data at the time of our analysis, 9,501 had valid

physical activity data, genetic information, and no diabetes at baseline (Supplementary Figure 1).

The median age was 56 years (interquartile range (IQR) 42–66), the median body mass index

(BMI) was 28.3 kg/m^2 (IQR 24.8–33.3), and most participants were women (72%), of European

ancestry (88%), and held a college degree (72%; Table 1). Participants contributed 11.6 million

person-days of Fitbit monitoring over a median of 5.7 years (IQR, 3.6–7.7), yielding 97.9 billion

recorded steps. Median daily step count was 7,891 (IQR, 5,748–10,471), and step count showed

a normal distribution (Supplementary Figure 2). Compared to non-include All of US

participants, included individuals were more likely to be younger, female, of European ancestry,

with lower BMI and higher educational attainment (Supplementary Table 1).

During 53,672 person-years of follow-up, 494 incident cases of type 2 diabetes were

documented. Crude type 2 diabetes incidence rates per 1,000 person-year according to daily step

counts tertiles were 13.5 (95% CI: 11.9-15.2), 7.5 (95% CI: 6.3-8.8), and 5.9 (95% CI: 4.7-7.0)

for the lowest, intermediate, and highest tertile, respectively (multivariable-adjusted HR of 0.40

(95% CI: 0.26, 0.63; Table 2). Each standard deviation (SD) increase in step count was

associated with lower risk of type 2 diabetes, with an adjusted HR of 0.65 (95% CI, 0.57–0.73).

Similarly, type 2 diabetes incidence increased across tertiles of polygenic risk, with incidence

rates per 1,000 person-years of 5.9 (95% CI: 4.8, 7.0) in the lowest, 7.9 (95% CI: 6.6, 9.2) in the

intermediate, and 14.0 (95% CI: 12.3, 15.8) in the highest tertile. In adjusted models, participants

in the highest tertile had 2.50 times the risk of type 2 diabetes compared with those in the lowest

(95% CI, 2.04–3.06; Table 2). When modelled in the continuous scale, each SD increase in

polygenic score was associated with an adjusted HR of 1.52 (95% CI, 1.38–1.67).

When examining joint associations, crude incidence rates ranged from 4.1 (95% CI, 2.5–5.7) per

1,000 person-years among participants with low genetic risk and high physical activity to 18.4

(95% CI, 15.2–21.6) among those with high genetic risk and low physical activity (Figure 1).

The corresponding multivariable-adjusted HR was 6.20 (95% CI, 3.97–9.60). Within the high

genetic risk group, individuals with low activity (~5,000 steps/day) had nearly double the

incidence of those with high activity (~13,000 steps/day; HR, 1.92; 95% CI, 1.36–2.69). The

joint association was higher than the sum of the risk associated with each factor alone, with a

relative excess risk due to interaction (RERI) of 0.20 (95% CI: 0.04, 0.36); p=0.007).

Decomposition of excess type 2 diabetes risk attributed 40% to genetic risk (95% CI: 31, 49),

45% to low physical activity (95% CI: 29, 62), and 15% to their interaction (95% CI: 3, 27;

Table 3). No evidence of multiplicative interaction was observed (p>0.05).

We conducted a series of sensitivity and subgroup analyses. When physical activity was

modeled by intensity, similar additive interactions were observed. The RERI was 0.21 (95% CI,

0.02–0.41; p=0.015) for moderate activity minutes, with 19% of excess risk attributable to

interaction, and 0.40 (95% CI, 0.21–0.59; p<0.001) for highly active minutes, with 31%

attributable to interaction (Supplementary Table 2). Adjustment for BMI did not materially

change these estimates (Supplementary Table 2). Using total physical activity assessed by self-

reported physical activity in Inter99, the RERI was 0.53 (95% CI, 0.30–0.76; p=0.007), with

44% of excess risk attributable to interaction (Supplementary Table 3). Adjustment for BMI

attenuated this proportion to 38% (95% CI, –6 to 82; Supplementary Table 3).

In ancestry-stratified analyses, no significant associations between genetics and physical activity

or interactions were observed in the African ancestry group (68 cases among 583 participants;

Supplementary Table 4). Due to small sample sizes, models for South Asian (5 cases among 85

participants) and East Asian (9 cases among 207 participants) did not converge. Sex-stratified

analyses showed additive interactions in both sexes. Among men, the RERI was 0.2 (95% CI:

0.01, 0.42; p=0.03), and 22% of excess risk was attributed to interaction. Among women, RERI

was 0.2 (95% CI: -0.04, 0.43; p=0.050), with an attributable proportion of 11% (Supplementary

Table 5).

DISCUSION

Our findings provide consistent evidence of an additive interaction between genetic risk and physical activity in the development of type 2 diabetes. Specifically, individuals with both high genetic risk and low step counts had higher risk than expected from the sum of their individual effects, with approximately one-fifth of excess risk attributable to this interaction. Similar patterns were observed when physical activity was assessed using device-based intensity metrics or self-reported total physical activity. Since additive interactions reflect differences in absolute risk across populations (28), these observations may have potential public health relevance, suggesting that reducing the overlap of these risk factors could contribute to lowering type 2 diabetes incidence. However, replication in other cohorts and settings will be important to better understand the robustness and generalizability of these results.

Our study builds on prior research investigating gene-environment interactions in type 2 diabetes, which has often yielded inconsistent findings (10–13). One likely contributor to these discrepancies is error in exposure assessment due to reliance on self-reported methods (14,15). Using accelerometer-based physical activity measures, a recent UK Biobank study demonstrated an additive effect between physical activity and polygenic scores on T2D risk, with greater absolute risk reductions observed among individuals at high genetic risk who engaged in moderate-to-vigorous physical activity (19). Our findings extend this evidence base in three important ways. First, unlike the UK Biobank study, which relied on single time-point physical activity assessment, we modelled physical activity as a time-varying exposure, capturing behavioural changes over a median follow-up of 5 years. This approach more accurately reflects behavioral dynamics leading up to diabetes onset (33). Second, we used daily step counts as the primary exposure. Compared with intensity-based classifications such as "moderate-to-

vigorous," which may vary in interpretation and are less intuitive, step counts provide a simple,

interpretable, and actionable metric for public health messaging (16). Third, our polygenic scores

were derived from an external discovery dataset that excluded participants from this study,

thereby mitigating potential inflation of type I error due to sample overlap, a methodological

limitation that may have influenced the UK Biobank analyses. Together, our study strengthens

the evidence that objectively measured physical activity interacts with genetic risk on the

development of type 2 diabetes.

Physical activity remains central to public health recommendations for type 2 diabetes

prevention (34), yet current guidelines do not account for heterogeneity in individual responses.

Factors such as genetic susceptibility, body composition, and diet can all influence the

effectiveness of physical activity (35). For example, in the ARIC cohort, the protective effects of

physical activity were attenuated in women with higher genetic predisposition to insulin

resistance (36), suggesting that greater physical activity may be needed to offset genetic risk.

Similarly, a prior analysis from the All of Us Research Program estimated that individuals at

high genetic risk for obesity would need to walk an additional ~2,300 steps per day to match the

obesity risk of those at lower risk (37). In our study, participants at high genetic risk who walked

~5,000 steps per day had nearly double the incidence of type 2 diabetes compared to those

walking ~13,000 steps per day. Our models estimated that 15% of type 2 diabetes cases could

have been prevented by eliminating one of the two risk factors, emphasizing the potential impact

of increasing activity levels, particularly among genetically susceptible individuals.

Although the proportion of risk attributable to gene–physical activity interactions was similar

when using step counts or device-based intensity measures, higher estimates were observed in

the Inter99 cohort for total physical activity. This discrepancy may reflect differences in follow-

up time, population characteristics such as age and obesity prevalence, or potential inflation due

to self-reported activity measures. Since population-attributable risk depends on both exposure

prevalence and effect size (38), exposure misclassification, especially with self-reported physical

activity, could partly explain these differences. While these excess risk estimates due to the

interaction may change over time with shifting diabetes incidence, our findings underscore the

ongoing relevance of gene–physical activity interactions in type 2 diabetes.

This study has several limitations. First, as an observational study, participants were not

randomized to high- or low-physical activity groups, limiting causal inference interpretations.

Recall-by-genotype studies could help test these hypotheses under experimental conditions.

Second, incident type 2 diabetes was defined solely based on any incident billing codes from the

electronic health record, which may introduce misclassification. However, this approach is

widely used in biobank research, including other All of Us analyses (17,39). Third, the number

of type 2 diabetes cases was relatively small, reflecting limited follow-up and the subset of

participants with linked wearable data. Fourth, while we used comprehensive polygenic scores

for type 2 diabetes, other genetic risk factors (i.e., for obesity) were not included. Although we

adjusted for BMI in secondary analyses, residual confounding, collider bias, or mediation effects

could still exist. Larger studies with longer follow-up and detailed modelling of BMI trajectories

may help address this. Finally, other molecular, clinical, or behavioral factors such as diet, body

composition, or gut microbiome were not explicitly included in our analyses, but could affect

risk or modify gene–environment interactions.

In conclusion, by quantifying the combined contributions of genetic risk and physical activity to

type 2 diabetes risk, our results underscore the potential value of integrating genomic data with

objective, device-derived measures of physical activity to better identify those who are most

likely to benefit from targeted interventions. Future studies with larger, diverse populations and

longer follow-up are needed to confirm these interactions and to evaluate how they can be

translated into effective public health and clinical interventions.

REFERENCES

1.	Franks PW, McCarthy MI. Exposing the exposures responsible for type 2 diabetes and obesity. Science. 2016 Oct 7;354(6308):69–73.

2.	McCarthy MI. Painting a new picture of personalised medicine for diabetes. Diabetologia. 2017 Feb 7.

3.	Franks PW, Pearson E, Florez JC. Gene-environment and gene-treatment interactions in type 2 diabetes: progress, pitfalls, and prospects. Diabetes Care. 2013 May 1;36(5):1413–21.

4.	Franks PW, Merino J. Gene-lifestyle interplay in type 2 diabetes. Curr Opin Genet Dev. 2018;50:35–40.

5.	DJ H. Gene-environment interactions in human diseases. Nat Rev Genet. 2005;6(4).

6.	Herrera-Luis E, Benke K, Volk H, Ladd-Acosta C, Wojcik GL. Gene–environment interactions in human health. Nat Rev Genet. 2024 Nov 1;

7.	Napier JD, Heckman RW, Juenger TE. Gene-by-environment interactions in plants: Molecular mechanisms, environmental drivers, and adaptive plasticity. Vol. 35, Plant Cell. 2023.

8.	Bourret A, Garant D. Candidate gene-environment interactions and their relationships with timing of breeding in a wild bird population. Ecol Evol. 2015;5(17).

9.	John B, Lewis KR. Chromosome variability and geographic distribution in insects. Science (1979). 1966;152(3723).

10.	Merino J, Guasch-Ferré M, Li J, Chung W, Hu Y, Ma B, et al. Polygenic scores, diet quality, and type 2 diabetes risk: An observational study among 35,759 adults from 3 US cohorts. PLoS Med. 2022;19(4).

11.   Langenberg C, Sharp SJ, Franks PW, Scott RA, Deloukas P, Forouhi NG, et al. Gene-
Lifestyle Interaction and Type 2 Diabetes: The EPIC InterAct Case-Cohort Study. PLoS
Med. 2014 May 20;11(5):e1001647.

12.   Wang T, Heianza Y, Sun D, Huang T, Ma W, Rimm EB, et al. Improving adherence to
healthy dietary patterns, genetic risk, and long term weight gain: gene-diet interaction
analysis in two prospective cohort studies. BMJ. 2018 Jan 10;360:j5644.

13.   Kilpeläinen TO, Qi L, Brage S, Sharp SJ, Sonestedt E, Demerath E, et al. Physical activity
attenuates the influence of FTO variants on obesity risk: a meta-analysis of 218,166 adults
and 19,268 children. Lewis C, editor. PLoS Med. 2011 Nov 1;8(11):e1001116.

14.   Hebert JR, Ebbeling CB, Matthews CE, Hurley TG, MA Y, Druker S, et al. Systematic
errors in middle-aged women's estimates of energy intake: Comparing three self-report
measures to total energy expenditure from doubly labeled water. Ann Epidemiol.
2002;12(8).

15.   Olds TS, Gomersall SR, Olds ST, Ridley K. A source of systematic bias in self-reported
physical activity: The cutpoint bias hypothesis. J Sci Med Sport. 2019 Aug 1;22(8):924–8.

16.   Bassett DR, Toth LP, LaMunion SR, Crouter SE. Step Counting: A Review of
Measurement Considerations and Health-Related Applications. Sports Medicine. 2017 Jul
1;47(7):1303–15.

17.   Master H, Annis J, Huang S, Beckman JA, Ratsimbazafy F, Marginean K, et al.
Association of step counts over time with the risk of chronic disease in the All of Us
Research Program. Nat Med. 2022;28(11).

18.   Ding D, Nguyen B, Nau T, Luo M, del Pozo Cruz B, Dempsey PC, et al. Daily steps and
health outcomes in adults: a systematic review and dose-response meta-analysis. Lancet
Public Health. 2025 Aug 1;10(8):e668–81.

19.   Luo M, Yu C, Del Pozo Cruz B, Chen L, Ding D. Accelerometer-measured intensity-
specific physical activity, genetic risk and incident type 2 diabetes: A prospective cohort
study. Br J Sports Med. 2023;57(19).

20. The "All of Us" Research Program. New England Journal of Medicine. 2019 Aug
15;381(7):668–76.

21. Jørgensen T, Borch-John sen K, Thomsen TF, Ibsen H, Glümer C, Pisinger C. A
randomized non-pharmacological intervention study for prevention of ischaemic heart
disease: Baseline results Inter99 (1). Eur J Prev Cardiol. 2003;10(5).

22. Aadahl M, Jørgensen T. Validation of a new self-report instrument for measuring physical
activity. Med Sci Sports Exerc. 2003;35(7).

23. Suzuki K, Hatzikotoulas K, Southam L, Taylor HJ, Yin X, Lorenz KM, et al. Genetic
drivers of heterogeneity in type 2 diabetes pathophysiology. Nature. 2024;627(8003).

24. All of Us Genomic Quality Report – User Support. Available from:
https://support.researchallofus.org/hc/en-us/articles/27633757470228-All-of-Us-
Genomic-Quality-Report

25. Ge T, Chen CY, Ni Y, Feng YCA, Smoller JW. Polygenic prediction via Bayesian
regression and continuous shrinkage priors. Nat Commun. 2019;10(1).

26. Observational Medical Outcomes Partnership (OMOP) common data model (CDM) –
User Support. Available from: https://support.researchallofus.org/hc/en-
us/articles/360039685271-Observational-Medical-Outcomes-Partnership-OMOP-
common-data-model-CDM

27. Munafò MR, Tilling K, Taylor AE, Evans DM, Davey Smith G. Collider scope: when
selection bias can substantially influence observed associations. Int J Epidemiol. 2018 Feb
1;47(1):226–35.

28. VanderWeele TJ, Tchetgen Tchetgen EJ. Attributing effects to interactions. Epidemiology
[Internet]. 2014 Sep [cited 2020 Jul 21];25(5):711–22.

29. Van Der Weele TJ, Knol MJ. A tutorial on interaction. Epidemiol Methods. 2014;3(1).

30. Fitbit Web API Data Dictionary.

31.   Bailey CP, Dodd KW, McClain JJ, Seo I, Wheeler W, Wolff-Hughes DL. Fitbit Physical
414   Activity and Sleep Data in the All of Us Research Program: Data Exploration and
415   Processing Considerations for Research. Med Sci Sports Exerc. 2025. In Press.

416   32.   Lim SS, Semnani-Azad Z, Morieri ML, Ng AH, Ahmad A, Fitipaldi H, et al. Reporting
417   guidelines for precision medicine research of clinical relevance: the BePRECISE
418   checklist. Nature Medicine 2024 30:7. 2024 Jul 19;30(7):1874–81.

419   33.   Hilden P, Schwartz JE, Pascual C, Diaz KM, Goldsmith J. How many days are needed?
420   Measurement reliability of wearable device data to assess physical activity. PLoS One.
421   2023 Feb 1;18(2):e0282162.

422   34.   3. Prevention or Delay of Diabetes and Associated Comorbidities: Standards of Care in
423   Diabetes—2024. Diabetes Care. 2024;47.

424   35.   Martiskainen T, Lamidi ML, Venojärvi M, Tikkanen H, Laatikainen T. Factors affecting
425   the effectiveness of the physical activity counselling intervention implemented in primary
426   health care in adults with type 2 diabetes. BMC Endocr Disord. 2023 Dec 1; 23(1).

427   36.   Yann C, Zhao K, Amit Arora C, Hsu CH. Association of physical activity with lower type
428   2 diabetes incidence is weaker among individuals at high genetic risk. Diabetologia.
429   2014;57(12).

430   37.   Brittain EL, Han L, Annis J, Master H, Hughes A, Roden DM, et al. Physical Activity and
431   Incident Obesity Across the Spectrum of Genetic Risk for Obesity. JAMA Netw Open.
432   2024 Mar 4;7(3):e243821–e243821.

433   38.   Rockhill B, Newman B, Weinberg C. Use and misuse of population attributable fractions.
434   Am J Public Health. 1998;88(1):15–9.

435   39.   Eastwood S V., Mathur R, Atkinson M, Brophy S, Sudlow C, Flaig R, et al. Algorithms
436   for the capture and adjudication of prevalent and incident diabetes in UK Biobank. PLoS
437   One. 2016 Sep 1;11(9).

Funding/Acknowledgements:

We gratefully acknowledge *All of Us* and Inter99 participants for their contributions, without

whom this research would not have been possible. The *All of Us* Research Program is supported

by grants 1 OT2 OD026549, 1 OT2 OD026554, 1 OT2 OD026557, 1 OT2 OD026556, 1 OT2

OD026550, 1 OT2 OD 026552, 1 OT2 OD026553, 1 OT2 OD026548, 1 OT2 OD026551, 1 OT2

OD026555, IAA AOD21037, AOD22003, AOD16037, and AOD21041 (regional medical

centers); grant HHSN 263201600085U (federally qualified health centers); grant U2C

OD023196 (data and research center); 1 U24 OD023121 (Biobank); U24 OD023176 (participant

center); U24 OD023163 (participant technology systems center); grants 3 OT2 OD023205 and 3

OT2 OD023206 (communications and engagement); and grants 1 OT2 OD025277, 3 OT2

OD025315, 1 OT2 OD025337, and 1 OT2 OD025276 (community partners) from the National

Institutes of Health (NIH).

This study was also supported by the Novo Nordisk Foundation grant NNF23SA0084103 and

the European Union. JM was additionally supported by grants from the European Commission

(HORIZON-EIC-2023-PATHFINDERCHALLENGES-01 - 101161509) and the EFSD/NNF

Future Leaders Award (#0094134). Views and opinions expressed are however those of the

author(s) only and do not necessarily reflect those of the European Union or European

Innovation Council and SMEs Executive Agency (EISMEA). Neither the European Union nor

the granting authority can be held responsible for them. RL was additionally supported by

personal grants from the Novo Nordisk Foundation (Laureate award no. NNF20OC0059313)

and the Danish National Research Fund (Chair DNRF161). TOK and MRC and were

additionally supported by grants from the Novo Nordisk Foundation (NNF22OC0074128) and

EFSD/Lilly European Diabetes Research Programme.

Ethics declarations:

The authors declare no competing interests. JM is the guarantor of this work, and, as such, had

full access to all the data in the study and takes responsibility for the integrity of the data and the

accuracy of the data analysis.

Figures and Tables

Figure 1: Risk of incident type 2 diabetes by global polygenic risk score for type 2 diabetes,

physical activity.

Figure Legend: Risk is indexed using crude incidence rate and hazard ratio adjusted for age,

sex and genetic ancestry. HR hazard ratio. Dots correspond to averages of daily step count

tertiles, which are ~ 5k, ~ 8k and ~13k. Histograms along the y-axis (to the right) show the

density of daily step counts. Incidence rates are computed for tertiles of daily step counts.

Table 1. Baseline characteristics of the included *All of US* research program participants.
481
482

Person-y$^\phi$	53,672	483 484
Age, years	56 (42-66)	485
Sex, Women, No. (%)	6,863 (72)	486 487
BMI, kg/m^2	28.3 (24.8-33.3)	488
Family history of diabetes, No. (%)	2,994 (32)	489 490 491
Smoker, No. (%)	3,136 (33)	492
Self-reported race/ethnicity$^\varepsilon$		493 494
White, No. (%)	8132 (86)	495
African, No. (%)	583 (6)	496
East Asian, No. (%)	207 (2)	497 498
South Asian, No. (%)	85 (1)	499
Education		500 501
College degree, No. (%)	6,856 (72)	502
Some/No college, No. (%)	2,587 (27)	503 504

505 **Table Legend:** Values are medians (interquartile range) for continuous variables; numbers and
506 (percentages) for categorical variables.
507

Table 2: Adjusted hazard ratios of T2D risk according to tertiles of daily step counts and genetic risk.

	Low step count	Intermediate step count	High step count	P for trend
No. of events/person-years	263 / 19415	136 / 18,046	95 / 16,210	—
Incidence rate (1,000 person-years; 95% CI)	13.5 (11.9-15.2)	7.5 (6.3-8.8)	5.9 (4.7-7.0)	—
Multivariable-adjusted model	1.00 (Ref)	0.68 (0.57-0.82)	0.40 (0.26-0.63)	<0.001

	Low genetic risk	Intermediate genetic risk	High genetic risk	P for trend
No. of events/person-years	107/ 18,180	142/ 17,991	245/ 17,500	—
Incidence rate (1,000 person-years; 95% CI)	5.9 (4.8-7)	7.9 (6.6-9.2)	14 (12.2-15.8)	—
Multivariable-adjusted model	1.00 (Ref)	1.59 (1.43-1.76)	2.50 (2.04-3.06)	<0.001

Table Legend: Hazards ratios and 95% CI for T2D risk. Daily step counts and genetic risk were categorized in three categories based on the distribution of physical activity and polygenic score. Cox proportional hazards models were adjusted for age (in years, continuous), sex (male, female), and the first 10 principal components of the genomic relationship matrix.

Table 3 Additive interactions between physical activity and global polygenic risk of T2D.

Main effects	
Physical activity[†]	1.53 (1.32, 1.74)
Polygenic score[‡]	1.61 (1.32, 1.74)
Joint effect	2.34 (1.87, 2.81)
Relative excess risk due to interaction	
Relative excess risk due to interaction	0.2 (0.04, 0.36)
P value	0.007
Attributable risk proportion, %	
Daily step count	39.8 (31.0, 48.5)
Polygenic score	45.4 (28.6, 62.1)
Additive interaction	14.8 (2.5, 27.2)

Table Legend: Multivariable adjusted risk of type 2 diabetes estimated from Cox proportional hazards models adjusted for age, sex, and the first ten principal components of genomic relationship matrix. The joint analysis was conducted to quantitatively estimate the combined association of physical activity and genetic risk simultaneously with the incidence of type 2 diabetes. The relative excess risk due to interaction (RERI) is based on joint effects of a one-standard-deviation increase in PRS and a one-standard-deviation decrease in daily step count using the following formula (RERI = RR11 - RR10 - RR01 + 1). The model's attributable risk proportion due to the interaction was calculated as AP = RERI/ RR11(28).

† per SD decrease in daily step counts.

‡ per SD increase in the global polygenic scores.