VOYAGER: an international consortium investigating the role of human papilloma virus and genetics in oral and oropharyngeal cancer risk and survival

Author list

Gormley M [1,2,3], Adhikari A [3], Dudding T [1,2], Pring M [2,3], Hurley K [3], Macfarlane GJ [4], Lagiou P [5], Lagiou A [6], Polesel J [7], Agudo A [8], Alemany L [9], Ahrens W [10], Healy CM [11], Conway DI [12], Canova C [13], Holcatova I [14], Richiardi L [15], Znaor A [16], Olshan AF [17], Hung RJ [18,19], Liu G [19,20], Bratman S [21], Zhao X [22], Holt J [22], Cortez R [23], Gaborieau V [23], McKay JD [23], Waterboer T [24], Brennan P [23], Hayes N [22], Diergaarde B [25], Virani S [23]

Affiliations

[1] MRC Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK.

[2] Bristol Dental School, University of Bristol, Bristol, UK.

[3] University Hospitals Bristol NHS Foundation Trust Bristol Dental Hospital, Bristol, UK

[4] School of Medicine, Medical Sciences and Nutrition, University of Aberdeen, UK

[5] School of Medicine, National and Kapodistrian University of Athens, Greece

[6] School of Public Health, University of West Attica, Greece

[7] Unit of Cancer Epidemiology, Centro di Riferimento Oncologico di Aviano (CRO) National Cancer Institute, IRCCS, Italy

[8] Nutrition and Cancer Unit, Cancer Epidemiology Research Program, Catalan Institute of Oncology/IDIBELL, Barcelona, Spain

[9] Infections and Cancer Unit, Cancer Epidemiology Research Program, Catalan Institute of Oncology/IDIBELL, Barcelona, Spain

[10] Epidemiological Methods and Etiological Research, Leibniz Institute for Prevention Research and Epidemiology – BIPS, Germany

[11] School of Dental Science, Dublin Dental University Hospital, Trinity College Dublin, Ireland

[12] School of Medicine, Dentistry, and Nursing, University of Glasgow, UK

[13] Department of Cardiac, Thoracic and Vascular Sciences University of Padova, Italy

[14] Institute of Hygiene and Epidemiology, Charles University Prague, Czech Republic

[15] Reference Centre for Epidemiology and Cancer Prevention in Piemonte, Italy

[16] Cancer Surveillance, International Agency for Research on Cancer, France

[17] Department of Epidemiology, Gillings School of Global Public Health, University of North Carolina,

US.

[18] Prosserman Centre for Population Health Research, Lunenfeld-Tanenbaum Research Institute,

Sinai Health System, Toronto, Canada.

[19] Dalla Lana School of Public Health, University of Toronto, Toronto, Canada.

[20] Computational Biology and Medicine Program, Princess Margaret Cancer Centre, Toronto Canada

[21] Department of Radiation Oncology, Princess Margaret Cancer Centre, University Health Network,

University of Toronto, Toronto, Canada.

[22] Department of Medicine, University of Tennessee, USA

[23] Genomic Epidemiology Group, World Health Organization, International Agency for Research on

Cancer, Lyon, France.

[24] Infections and Cancer Epidemiology, German Cancer Research Center (DKFZ), Heidelberg,

Germany.

[25] Department of Human Genetics, School of Public Health, University of Pittsburgh, and UPMC

Hillman Cancer Center, Pittsburgh, US.

Word count: 3,224

Abstract

Head and neck cancer (HNC) is the sixth most common cancer globally. Incidence and survival rates vary significantly across geographic regions and tumor subsites. This is partly due to differences in risk factor exposure, which includes tobacco smoking, alcohol consumption and human papillomavirus (HPV) infection, alongside detection and treatment strategies. The VOYAGER (human papillomaVirus, Oral and oropharYngeal cAncer GEnomic Research) consortium is a collaboration between five large North American and European studies which generated data on 10,530 participants (7,233 cases and 3,297 controls). The primary goal of the collaboration was to improve understanding of the role of HPV and genetic factors in oral cavity and oropharyngeal cancer risk and outcome. Demographic and clinical data collected by the five studies were harmonized, and HPV status was determined for the majority of cases. In addition, 999 tumors were sequenced to define somatic mutations. These activities generated a comprehensive biomedical resource that can be utilized to answer critical outstanding research questions to help improve HNC prevention, early detection, treatment, and surveillance.

Key words

Head and neck cancer, oral cancer, oropharyngeal cancer, human papilloma virus, risk factors, survival.

Background

Head and Neck Cancer Genomic Epidemiology

Head and neck cancer (HNC) which is primarily squamous cell carcinoma, includes cancers of the oral cavity, pharynx and larynx.(1, 2) Globally, the incidence of oral and oropharyngeal cancer is estimated at 8.0 and 2.0 per 100,000, respectively, and is predicted to increase by 30% by 2030.(3, 4) Five-year survival remains poor, averaging between 40 – 50% with hypopharynx cases experiencing the worst outcomes.(5) Incidence and survival rates vary significantly across geographic regions and tumor subsites, partly due to differences in risk factor exposure. Established HNC risk factors include tobacco smoking and alcohol intake, which together account for a similar population attributable risk for both oral (64%) and oropharyngeal (72%) cancer.(6) However, human papilloma virus (HPV) infection, particularly high-risk subtype 16, has emerged as another major risk factor for oropharyngeal cancer.(7-9) Worldwide, it is estimated that around 52,000 incident HNC cases are caused by a persistent HPV infection each year, with attributable fractions highest in high-income countries in North America and Europe.(10-13)

Given the decline of tobacco use in developed countries, the incidence rate of HPV driven [HPV(+)] oropharyngeal cancer is now surpassing that of oral cancer.(9-12, 14) HPV(+) oropharyngeal tumors are considered distinct entities, demonstrating more favorable treatment response and prognosis compared to non-HPV related oropharyngeal cancer [HPV(-)].(8, 12, 15-17) This is likely due to differences in etiology, patient and tumor characteristics, with HPV(+) oropharyngeal tumors presenting more frequently in younger individuals (<65 years), and in those reporting higher numbers of sexual partners with reduced cumulative tobacco exposure compared to HPV(-) cases.(12, 18) However, only a small proportion of those with an oral HPV infection will develop HNC and despite better long-term survival, up to 25% of patients still develop disease recurrence within 5 years after initial diagnosis.(19) To improve prevention, early detection and prognosis, a better understanding of the role of host genetics and interactions with modifiable risk factors, such

as tobacco and alcohol use in oral and oropharyngeal cancer risk and survival is

required.(20)

HPV driven carcinogenesis is characterized by increased expression of the viral oncogenes

E6 and E7, leading to increased degradation of tumor suppressor proteins p53 and Rb,

respectively and loss of cell cycle activation. This can result in genomic instability and

resistance to apoptosis.(21, 22) HPV(+) and HPV(-) head and neck tumors harbor a similar

burden of somatic variants. However, HPV(+) oropharyngeal tumors carry fewer copy-

number alterations, suggesting a higher degree of genomic stability.(23-26) Genome

profiling studies have provided a list of genes that are recurrently mutated in HNC, including

TP53, *CDKN2A* (which encodes for p16[INK4]), *NOTCH1* and *PIK3CA*.(23, 24, 27-29) Genes

recurrently mutated in HPV(+) oropharyngeal cancer are related to epithelial structure and

differentiation, in addition to *RB1* (encoding the Rb protein).(12, 23-26) The presence or

absence of particular somatic alterations in tumors may be good markers of cancer

prognosis and response to treatment, but there is still a need to identify novel somatic driver

alterations, particularly as relatively few HPV(+) oropharyngeal cancer cases have been

sequenced to date.(24, 30, 31) Identification of molecular markers associated with prognosis

could facilitate better monitoring and clinical decision making, including the use of de-

escalation treatment strategies among those at lower risk of recurrence or progression as a

means to improve quality of life and, conversely, more aggressive treatment in those

deemed at higher risk.

Compared to other major cancer sites such as breast, lung and colorectal, HNC is relatively

rare, hampering research efforts. Collaboration between studies to form extensive

biomedical databases and resources plays a crucial role in driving progress across various

domains of cancer research.

Construction and content

The VOYAGER Consortium: Design

In 2016, the VOYAGER (human papillomaVirus, Oral and oropharYngeal cAncer GEnomic

Research) consortium was established, bringing together five large North American and

European studies (**Figure 1**) with a focus on oral and oropharyngeal cancers. The primary

goal of the collaboration was to improve understanding of the role of HPV and genomic

factors in oral cavity and oropharyngeal cancer risk and outcome. The project was funded by

the US National Institute of Dental and Craniofacial Research (NIDCR; R01DE025712).

The VOYAGER consortium includes 10,530 participants in total (6,489 oral and

oropharyngeal cancer cases, 744 other head and neck cancer cases, and 3,297 controls),

with detailed demographic, risk factor and clinical data. The five studies comprising

VOYAGER have been previously described and are: (a) the Alcohol-related cancers and

genetic susceptibility in Europe (ARCAGE) study (32), (b) the Toronto Mount Sinai Hospital-

Princess Margaret (MSH-PMH) study (Toronto) (33), (c) the University of Pittsburgh case-

control study on head and neck cancer (Pittsburgh) (34), (d) the Carolina Head and Neck

Cancer Epidemiology (CHANCE) study (35), and (e) the Head and Neck 5000 study

(HN5000) (36). Ethical approval was obtained as described in the **Declarations** section

below.

Each study contributed oral cavity and oropharyngeal cancer cases as defined by the

following ICD-10 codes: oral cavity (C00.3-C00.6, C00.8-C00.9, C02.0-C02.3, C02.8, C02.9,

C03.0-C03.9, C04.0-C04.9, C05.0, C05.8, C05.9, C06.0-C06.9) oropharynx (C01-C01.9,

C02.4, C05.1-C05.2, C09.0-C10.9), head and neck not otherwise specified (NOS) (C76.0),

pharynx NOS (C14.0), or any cases with overlap of these sites. Demographic information,

including age, sex, ethnicity, geographic region and education level, and information on

established risk factors, e.g., smoking and alcohol history, were shared. Clinical variables

included ICD code, tumor, nodal and metastasis status, HPV status (defined by p16 or HPV

DNA), vital status, follow-up time, and treatment information. Given all cases were

diagnosed between 2002 – 2018, the American Joint Committee on Cancer (AJCC) 7th

edition was used, mirroring the staging used in clinical practice.(37)

Data Generation and Harmonization

The VOYAGER consortium generated several different types of data including clinical,

demographic and behavior variables. Data harmonization was conducted for variables that

were centralized across all studies, as illustrated in **Tables 1** and **2**. Upon receipt of data,

cleaning and validation checks were conducted to identify inconsistencies, outliers, and

missing values. Follow-up data was carefully collated and harmonized to capture follow-up

times, progression events, vital status and cause of death to facilitate high-quality prognostic

research. A comprehensive data dictionary was developed to specify final definitions,

formats, permissible values, coding schemes and classifications (i.e., for categorical

variables). Version control was implemented to ensure consistency of data use across

analyses. Further detail on the statistical analyses performed is available in the

Supplementary Methods.

HPV16 E6 serology was prioritized as a marker of HPV(+) oropharyngeal cancer as this has

been shown to be a highly sensitive and specific marker of HPV oncogenic infection in

oropharyngeal cancer and can be easily assayed from blood.(38, 39) Multiplex serology was

performed on 75% (n= 5,294) of all HNC cases and 92% (n= 3,250) of oropharyngeal cancer

cases using a previously developed Luminex assay.(40, 41) Multiplex serology generates

quantitative data expressed in median fluorescence intensity (MFI) units for each pathogen-

specific antigen and serum. Seropositivity for every antigen was based on previously

determined standardized cut-offs in order to optimize sensitivity and specificity.(42, 43)

When serology was missing, two concordant tumor markers, p16 immunohistochemistry

(IHC) and high-risk HPV DNA in-situ hybridization (ISH) were required to determine HPV

status. If p16 IHC and HPV DNA ISH were discordant or only one marker was available,

then HPV status was unknown. This algorithm was based on evaluation of biomarkers

performance compared to molecular reference method (serology) from known data, led by

consortium members.(39, 44)

Genotyping data was generated at the Center for Inherited Disease Research (CIDR) in

several rounds. The first round (X01HG007780) was performed using the Illumina

OncoArray, which was custom designed for cancer studies by the OncoArray Consortium

part of the Genetic Associations and Mechanisms in Oncology (GAME-ON) Network. All

samples (6,034 cases and 6,585 controls) were genotyped as part of the oral and pharynx

cancer OncoArray study, except for 1,023 controls from the Toronto study which were

genotyped as part of the Lung OncoArray. This genotyping data was used to conduct the

first genome-wide association study (GWAS) on head and neck cancer in 2017.(45) With the

confirmation of HPV status, genotyping data were also used to run a GWAS of oral and

oropharyngeal cancer, stratified by HPV status.(46) A second round of genotyping was

undertaken (X01HG010743) for an additional 1,491 samples in VOYAGER. This was

conducted on the All of Us Array, an Illumina array customized for the All of Us Consortium

and designed to include multiethnic context.(47) The genotyping data from OncoArray and

the All of Us Array, has contributed to the largest HNC GWAS to date including 19,073

cases and 38,357 controls identifying 29 independent genetic loci.(48)

DNA tumor sequencing data were also generated for 999 samples in the VOYAGER

consortium using a custom cancer gene panel that has been previously reported.(49-53)

Next generation sequencing was performed using the Agilent SureSelect protocol and

reagents according to manufacturer's specifications. The assay targets all genes of the

Cancer Gene Census, in addition to clinically relevant targets such as drug metabolizing

enzymes. The assay also performs whole genome sequencing of HPV16 and 18 using

methodology previously reported to offer clinical diagnostic accuracy comparable or better to

conventional approaches while at the same time capturing base level resolution across the

HPV genome.(50) The resulting libraries were sequenced on Illumina sequencers, primarily

NovaSeq according to manufacturer's specifications. Further details of the assay can be

found elsewhere.(54) Target depth was mean exon coverage of 500x coverage. All analytic

tools and the pipelines for integrating steps have been publicly reported and are available as

open source software including: BWA for sequence alignment (55), NGS Copy (54) for copy

number assessment, Strelka (56), ABRA for realignment and structural variant detection

(57), and UNMASC for variant prioritization and filtering.(49)

VOYAGER Data Resource: Clinical and Demographic Profile

The VOYAGER consortium comprises 10,530 participants in total (6,489 oral and

oropharyngeal cancer cases, 744 other head and neck cancer cases, and 3,297 controls).

The primary focus was on oral cavity and oropharyngeal cancers, with systematically

planned and standardized inclusion criteria using ICD-10 coding. However, additional cases

from other head and neck subsites were also included opportunistically. In total, 3,514

oropharynx cancers and 2,975 oral cavity cancers, alongside 744 cases from several other

head and neck cancer subsites were included. The controls included were used for the

genotyping studies. To facilitate the use of this resource, we present here a brief description

of the oral cavity and oropharyngeal cancers and controls, while noting that this resource

contains additional head and neck subsites, described in Error! Reference source not found.,

that may be valuable to the scientific community.

Cancer cases were contributed to by HN5000 (41%), followed by Toronto (23%), Pittsburgh

(13%), CHANCE (12%) and ARCAGE (11%). There were significant differences (p <0.0001)

in anatomical site and staging across studies, with HN5000 contributing the highest number

of oropharyngeal cancer cases (45%) and therefore higher numbers of late (stage III and IV)

disease (**Supplementary Table 1**). Differences were also detected between cases versus

controls across all clinical and demographic variables, except age (p= 0.308) (**Table 1**).

HPV16 E6 serology status was available for 3,250 (92%) oropharyngeal cancer cases, of

which 61% were HPV(+) (**Figure 1; Table 1**). When serology was missing, p16 and high-risk

HPV DNA ISH concordance determined HPV oropharyngeal cancer status in the remaining

~8% of cases. There was good concordance between p16 and HPV DNA ISH. The

proportion of HPV(+) and HPV(-) oropharyngeal tumors was similar across North American

and European regions (p= 0.156), but these cases varied in terms of all other demographic,

clinical and risk factor behavior variables (**Table 1; Table 2**).

There are in total 2,975 oral cavity, and 3,514 oropharyngeal cancer cases (2,138 HPV(+),

1,146 HPV(-) and 230 HPV status unknown oropharyngeal cancer cases included in

VOYAGER (**Table 1**). Most cases presented in males (73%), outnumbering females across

all subsites. The overall mean age at diagnosis in all cases was 60 years (SD= 10.7), with

the lowest mean age observed in the HPV(+) oropharyngeal cancer group (58 years (SD=

8.9)). Overall age at diagnosis for oropharyngeal cancer (59 (SD= 9.6)) was significantly

younger than that of oral cavity (61 (SD= 12.2)) (p< 0.0001) (**Table 1**). Only 3% of oral and

oropharyngeal squamous cell carcinoma cases (n= 224) presented under 40 years old, and

these were predominantly patients with cancer of the oral cavity (71%). The highest

frequency of postsecondary education was observed in the HPV(+) oropharyngeal cancer

group (42%), with the lowest in the HPV(-) oropharyngeal group (28%) (**Table 1**).

Most cases were current (33%) or former smokers (42%), except for the HPV(+)

oropharyngeal group which had the largest never smoker population (34%) with significant

differences in smoking found between subsites (p< 0.0001) (**Table 2**). Similarly, more than

half (63%) of cases reported current alcohol drinking, which was consistent across subsites.

The proportion of never drinkers was different in oral cavity (16%) compared to

oropharyngeal cancer (12%), and significantly lower in total cases (14%) versus controls

(22%) (p< 0.0001) (**Table 2**). Overweight BMI was most common in the HPV(+)

oropharyngeal cancer subsite (40%), with underweight BMI being the least frequently

recorded category across oral and oropharyngeal cancer cases. There were significant

differences between total cases and controls across all BMI categories (p<0.0001) (**Table 2**).

Overall, significantly more oropharyngeal cancer cases presented at stage IV (72%

compared to oral cavity cancers which presented earlier, at stage I (28%) and II (22%)

(p<0.0001) (**Table 1**). Non-surgical treatment using radiotherapy, with or without

chemotherapy, was the most common treatment modality for oropharyngeal cancer (66%).

Surgery was the most common treatment modality for cases of oral cavity cancer, with

almost half (47%) of these patients receiving surgery alone and another 40% undergoing

surgery plus adjuvant radiotherapy with or without chemotherapy. Collecting information on

disease outcome was a primary focus of the consortium given the high risk of recurrence

and poor survival associated with HNC. The median length of follow-up time for oral and

oropharyngeal cancer cases was 5.3 years. Overall median survival time was 9.8 years, and

the five-year survival rate was 66% across all oral cavity and oropharyngeal cancer sites

(**Table 3**).

Patients with HPV(+) oropharyngeal cancer had the best overall survival, with a median

survival time of 14.3 years and 5-year survival at 81% (**Table 3** and **Figure 2**). For patients

where HPV status was not available, probability of survival at 5 years fell between that of

HPV(+) and HPV(-) oropharyngeal cancer patients, at 63%, indicating these were likely a

mix of patients with and without HPV(+) oropharyngeal cancer. The probability of being

progression-free at 5 years after diagnosis was highest for HPV(+) oropharyngeal cancer

patients at 75%, and similar for all other subsites, ranging from 44% – 50%. Overall, the

probability of being progression-free at 5 years was 57%. Disease-specific survival reflected

similar trends across subsites (**Table 3** and **Figure 2**). HPV(+) oropharyngeal cancer

patients had the highest probability of disease-specific survival at 5 years (88%) and HPV(-)

oropharyngeal cancer had the lowest (67%). Across all cancers, the probability of disease-

specific survival was 78% (**Table 3** and **Figure 2**).

Median length of follow-up time across studies varied, but not significantly so. Median length

of follow-up time was 8.4 years for ARCAGE, 10.0 years for CHANCE, 4.9 years for

HN5000, 5.7 years for Pittsburgh and 5.2 years for Toronto (p= 0.4). Patients from the

ARCAGE study have lower overall, progression-free and disease-specific survival compared

to other studies for HPV(+) oropharyngeal cancers, however, there were only 63 HPV(+)

oropharyngeal cancer cases in this cohort (**Supplementary Figures 1 - 3**). In addition, the

smoking rates of patients in ARCAGE is much higher compared to other studies and this has

been shown to affect interpretation of its HPV(+) oropharyngeal cancer profiles(38).

The clinical, demographic and survival profile of the VOYAGER cohort are characteristic of

what we know of HNC patients, epidemiologically and clinically.(4, 58, 59) We show the

predominance of male patients, the established differences in age at diagnosis, education

and survival of HPV driven and non-HPV cancers, and the expected distributions of risk

factor behaviors.(4, 58) Therefore, VOYAGER provides a reliable representative resource

for further research into this patient population. Importantly, the overlap of data with follow-

up data, somatic tumor sequencing and germline genotyping is a strength of this resource

(**Figure 3**).

Utility and discussion

Several important findings have come from the VOYAGER consortium. Previous GWAS

studies of oral and oropharyngeal cancer conducted by our group highlighted the important

role of the human leukocyte antigen (HLA) region (6p21.3) in susceptibility to oropharyngeal

cancer.(45) In particular, a two-fold protective effect was observed for oropharyngeal cancer

and the HLA haplotype DRB1*1301-DQA1*0103-DQB1*0603. This haplotype was previously

reported to also be protective for cervical cancer, a cancer type that is primarily driven by

HPV infection. In VOYAGER, HPV status was determined in 92% of oropharyngeal tumors

via HPV16 E6 serology enabling the first GWAS focused on HPV driven HNCs using fine

mapping techniques.(46) Within the 6p21.3 locus, there were two specific loci (rs4713462

and rs9269942) independently associated with reduced risk of oropharyngeal cancer

(**Figure 4**). These loci were separately associated with antibodies against specific HPV16

proteins which implicates specific germline variants in the natural immune response against

HPV(+) oropharyngeal cancer, supporting the use of therapeutic vaccines to protect against

this disease.

A recent publication utilized VOYAGER data to develop a risk prediction model for HNC

including genetic markers, HPV serostatus, demographic and lifestyle risk factors in

populations of European ancestry. The addition of HPV serology provided substantial

predictive accuracy for oropharyngeal cancer (AUC= 0.94, 95%CI: 0.92 – 0.95 in men and

AUC= 0.92, 95%CI: 0.88 – 0.95 in women) above that of previously published models,

highlighting the need to consider primary prevention and intensive surveillance for

oropharyngeal cancer subgroups.(60) Importantly, however, while HPV serology is a marker

for HPV driven oropharyngeal cancer, the use of HPV serology needs to be carefully

evaluated among smokers as demonstrated in another VOYAGER publication.(38)

Diagnostic accuracy of HPV serology was evaluated utilizing VOYAGER data and found to

be highly sensitive and specific independent of age, sex, year of diagnosis, BMI at

diagnosis, current alcohol use and primary tumor size, but exhibited some variation in

diagnostic accuracy for heavy smokers and by lymph node involvement. This work provides

further evidence that this additional HPV biomarker can be used for early diagnosis.

Randomized clinical trials remain the 'gold standard' to ascertain the causal effect of

interventions or modifiable exposures. However, they are not always feasible in terms of

cost, time or ethics (61). Conversely, observational studies are subject to confounding, bias

and reverse causality. This has significantly limited study design in HNC research, given that

it is a relatively rare cancer with correlated risk factors, requiring long-term follow-up from the

point of exposure to disease onset. To overcome such limitations, Mendelian randomization

(MR) uses measured genetic variation to examine the causal effect of potentially modifiable

exposures on health outcomes in observational data.(62-64) The genotyping data available

in VOYAGER has contributed to multiple MR studies (65-70) investigating a wide range of

genetically proxied exposures. These studies have among other things demonstrated an

independent causal effect for both smoking and alcohol on both HPV(+) and HPV(-) HNC,

suggesting that the effect of alcohol may have been previously underestimated.(65, 69) This

work further strengthens the evidence to support public health messaging around prevention

in HNC.

Conclusion and future prospects

The value of HNC data generated as part of VOYAGER highlights its use for studies focused

on prognosis, particularly overall survival where data is most complete. The variation across

centers is an important consideration, as studies contributing to VOYAGER were conducted

across different geographical settings and health systems. The VOYAGER data resource is

particularly suited for genomic studies on risk factors and outcome.

References

1.   Zhou T, Huang W, Wang X, Zhang J, Zhou E, Tu Y, et al. Global burden of head and

neck cancers from 1990 to 2019. iScience. 2024;27(3):109282.

2.   Fitzmaurice C, Allen C, Barber RM, Barregard L, Bhutta ZA, Brenner H, et al. Global,

Regional, and National Cancer Incidence, Mortality, Years of Life Lost, Years Lived With

Disability, and Disability-Adjusted Life-years for 32 Cancer Groups, 1990 to 2015: A

Systematic Analysis for the Global Burden of Disease Study. JAMA Oncol. 2017;3(4):524-

48.

3.   Sung H, Ferlay J, Siegel RL, Laversanne M, Soerjomataram I, Jemal A, et al. Global

Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36

Cancers in 185 Countries. CA: A Cancer Journal for Clinicians. 2021;71(3):209-49.

4.   Johnson DE, Burtness B, Leemans CR, Lui VWY, Bauman JE, Grandis JR. Head

and neck squamous cell carcinoma. Nat Rev Dis Primers. 2020;6(1):92.

5.   Bosetti C, Carioli G, Santucci C, Bertuccio P, Gallus S, Garavello W, et al. Global

trends in oral and pharyngeal cancer incidence and mortality. International Journal of

Cancer. 2020;147(4):1040-9.

6.   Hashibe M, Brennan P, Chuang SC, Boccia S, Castellsague X, Chen C, et al.

Interaction between Tobacco and Alcohol Use and the Risk of Head and Neck Cancer:

Pooled Analysis in the International Head and Neck Cancer Epidemiology Consortium.

Cancer Epidem Biomar. 2009;18(2):541-50.

7.   Heck JE, Berthiller J, Vaccarella S, Winn DM, Smith EM, Shan'gina O, et al. Sexual

behaviours and the risk of head and neck cancers: a pooled analysis in the International

Head and Neck Cancer Epidemiology (INHANCE) consortium. Int J Epidemiol.

2010;39(1):166-81.

8.   Ang KK, Harris J, Wheeler R, Weber R, Rosenthal DI, Nguyen-Tân PF, et al. Human

papillomavirus and survival of patients with oropharyngeal cancer. N Engl J Med.

2010;363(1):24-35.

9.   Chaturvedi AK, Engels EA, Pfeiffer RM, Hernandez BY, Xiao W, Kim E, et al. Human

papillomavirus and rising oropharyngeal cancer incidence in the United States. J Clin Oncol.

2011;29(32):4294-301.

10.   Chaturvedi AK, Anderson WF, Lortet-Tieulent J, Curado MP, Ferlay J, Franceschi S,

et al. Worldwide trends in incidence rates for oral cavity and oropharyngeal cancers. J Clin

Oncol. 2013;31(36):4550-9.

11.   Mehanna H, Beech T, Nicholson T, El-Hariry I, McConkey C, Paleri V, et al.

Prevalence of human papillomavirus in oropharyngeal and nonoropharyngeal head and neck

cancer--systematic review and meta-analysis of trends by time and region. Head Neck.

2013;35(5):747-55.

12.   Lechner M, Liu J, Masterson L, Fenton TR. HPV-associated oropharyngeal cancer:

epidemiology, molecular biology and clinical management. Nature Reviews Clinical

Oncology. 2022;19(5):306-27.

13.   de Martel C, Georges D, Bray F, Ferlay J, Clifford GM. Global burden of cancer

attributable to infections in 2018: a worldwide incidence analysis. Lancet Glob Health.

2020;8(2):e180-e90.

14.   Lauritzen BB, Grønlund MW, Jakobsen KK, Justesen MM, Garset-Zamani M,

Carlander AF, et al. Epidemiological trends and survival of oropharyngeal cancer in a high

HPV-prevalent area: A Danish population-based study from 2000 to 2020. Int J Cancer.

2024;155(12):2169-79.

15.   Fakhry C, Westra WH, Li S, Cmelak A, Ridge JA, Pinto H, et al. Improved survival of

patients with human papillomavirus-positive head and neck squamous cell carcinoma in a

prospective clinical trial. J Natl Cancer Inst. 2008;100(4):261-9.

16.   Fakhry C, Zhang Q, Nguyen-Tan PF, Rosenthal D, El-Naggar A, Garden AS, et al.

Human papillomavirus and overall survival after progression of oropharyngeal squamous cell

carcinoma. J Clin Oncol. 2014;32(30):3365-73.

17.   Zanoni DK, Patel SG, Shah JP. Changes in the 8th Edition of the American Joint

Committee on Cancer (AJCC) Staging of Head and Neck Cancer: Rationale and

Implications. Curr Oncol Rep. 2019;21(6):52.

18.   Gillison ML, D'Souza G, Westra W, Sugar E, Xiao W, Begum S, et al. Distinct risk

factor profiles for human papillomavirus type 16-positive and human papillomavirus type 16-

negative head and neck cancers. J Natl Cancer Inst. 2008;100(6):407-20.

19.   INTEGRATE. Post-Treatment Head and Neck Cancer Care: National Audit and

Analysis of Current Practice in the United Kingdom. Clin Otolaryngol. 2021;46(1):284-94.

20.   Brenner N, Mentzer AJ, Hill M, Almond R, Allen N, Pawlita M, et al. Characterization

of human papillomavirus (HPV) 16 E6 seropositive individuals without HPV-associated

malignancies after 10 years of follow-up in the UK Biobank. EBioMedicine. 2020;62:103123.

21.   Huibregtse JM, Scheffner M, Howley PM. A cellular protein mediates association of

p53 with the E6 oncoprotein of human papillomavirus types 16 or 18. Embo j.

1991;10(13):4129-35.

22.   Scheffner M, Werness BA, Huibregtse JM, Levine AJ, Howley PM. The E6

oncoprotein encoded by human papillomavirus types 16 and 18 promotes the degradation of

p53. Cell. 1990;63(6):1129-36.

23.   Seiwert TY, Zuo Z, Keck MK, Khattri A, Pedamallu CS, Stricker T, et al. Integrative

and comparative genomic analysis of HPV-positive and HPV-negative head and neck

squamous cell carcinomas. Clin Cancer Res. 2015;21(3):632-41.

24.   The Cancer Genome Atlas Network. Comprehensive genomic characterization of

head and neck squamous cell carcinomas. Nature. 2015;517(7536):576-82.

25.   Gillison ML, Akagi K, Xiao W, Jiang B, Pickard RKL, Li J, et al. Human

papillomavirus and the landscape of secondary genetic alterations in oral cancers. Genome

Res. 2019;29(1):1-17.

26.   Dogan S, Xu B, Middha S, Vanderbilt CM, Bowman AS, Migliacci J, et al.

Identification of prognostic molecular biomarkers in 157 HPV-positive and HPV-negative

squamous cell carcinomas of the oropharynx. Int J Cancer. 2019;145(11):3152-62.

27.   Agrawal N, Frederick MJ, Pickering CR, Bettegowda C, Chang K, Li RJ, et al. Exome

sequencing of head and neck squamous cell carcinoma reveals inactivating mutations in

NOTCH1. Science. 2011;333(6046):1154-7.

28.   Pickering CR, Zhang J, Neskey DM, Zhao M, Jasser SA, Wang J, et al. Squamous

cell carcinoma of the oral tongue in young non-smokers is genomically similar to tumors in

older smokers. Clin Cancer Res. 2014;20(14):3842-8.

29.   Stransky N, Egloff AM, Tward AD, Kostic AD, Cibulskis K, Sivachenko A, et al. The

mutational landscape of head and neck squamous cell carcinoma. Science.

2011;333(6046):1157-60.

30.   Gauthier MA, Kadam A, Leveque G, Golabi N, Zeitouni A, Richardson K, et al. Long-

read sequencing of oropharyngeal squamous cell carcinoma tumors reveal diverse patterns

of high-risk Human Papillomavirus integration. Front Oncol. 2023;13:1264646.

31. Lechner M, Frampton GM, Fenton T, Feber A, Palmer G, Jay A, et al. Targeted next-

generation sequencing of head and neck squamous cell carcinoma identifies novel genetic

alterations in HPV+ and HPV- tumors. Genome Med. 2013;5(5):49.

32. Lagiou P, Georgila C, Minaki P, Ahrens W, Pohlabeln H, Benhamou S, et al. Alcohol-

related cancers and genetic susceptibility in Europe: the ARCAGE project: study samples

and data collection. Eur J Cancer Prev. 2009;18(1):76-84.

33. Thomas S, Carroll JC, Brown MC, Chen Z, Mirshams M, Patel D, et al. Nicotine

dependence as a risk factor for upper aerodigestive tract (UADT) cancers: A mediation

analysis. PLoS One. 2020;15(8):e0237723.

34. Troy JD, Grandis JR, Youk AO, Diergaarde B, Romkes M, Weissfeld JL. Childhood

passive smoke exposure is associated with adult head and neck cancer. Cancer Epidemiol.

2013;37(4):417-23.

35. Divaris K, Olshan AF, Smith J, Bell ME, Weissler MC, Funkhouser WK, et al. Oral

health and risk for head and neck squamous cell carcinoma: the Carolina Head and Neck

Cancer Study. Cancer Causes Control. 2010;21(4):567-75.

36. Ness AR, Waylen A, Hurley K, Jeffreys M, Penfold C, Pring M, et al. Establishing a

large prospective clinical cohort in people with head and neck cancer as a biomedical

resource: head and neck 5000. BMC Cancer. 2014;14(1):973.

37. Edge SB, Compton CC. The American Joint Committee on Cancer: the 7th edition of

the AJCC cancer staging manual and the future of TNM. Ann Surg Oncol. 2010;17(6):1471-

4.

38. Kusters JMA, Diergaarde B, Ness A, Schim van der Loeff MF, Heijne JCM,

Schroeder L, et al. Diagnostic accuracy of HPV16 early antigen serology for HPV-driven

oropharyngeal cancer is independent of age and sex. Int J Cancer. 2024;154(2):389-402.

39. Hibbert J, Halec G, Baaken D, Waterboer T, Brenner N. Sensitivity and Specificity of

Human Papillomavirus (HPV) 16 Early Antigen Serology for HPV-Driven Oropharyngeal

Cancer: A Systematic Literature Review and Meta-Analysis. Cancers (Basel). 2021;13(12).

40. Waterboer T, Sehr P, Pawlita M. Suppression of non-specific binding in serological

Luminex assays. Journal of Immunological Methods. 2006;309(1):200-4.

41. Waterboer T, Sehr P, Michael KM, Franceschi S, Nieland JD, Joos TO, et al.

Multiplex human papillomavirus serology based on in situ-purified glutathione s-transferase

fusion proteins. Clin Chem. 2005;51(10):1845-53.

42. Clifford GM, Shin H-R, Oh J-K, Waterboer T, Ju Y-H, Vaccarella S, et al. Serologic

Response to Oncogenic Human Papillomavirus Types in Male and Female University

Students in Busan, South Korea. Cancer Epidemiology, Biomarkers & Prevention.

2007;16(9):1874-9.

43. Kreimer AR, Johansson M, Waterboer T, Kaaks R, Chang-Claude J, Drogen D, et al.

Evaluation of Human Papillomavirus Antibodies and Risk of Subsequent Head and Neck

Cancer. Journal of Clinical Oncology. 2013;31(21):2708-15.

44. Morais E, Kothari S, Chen YT, Roberts CS, Gómez-Ulloa D, Fenoll R, et al. The

BROADEN study: The design of an observational study to assess the absolute burden of

HPV-related head and neck cancers. Contemp Clin Trials. 2022;115:106631.

45. Lesseur C, Diergaarde B, Olshan AF, Wünsch-Filho V, Ness AR, Liu G, et al.

Genome-wide association analyses identify new susceptibility loci for oral cavity and

pharyngeal cancer. Nat Genet. 2016;48(12):1544-50.

46. Ferreiro-Iglesias A, McKay JD, Brenner N, Virani S, Lesseur C, Gaborieau V, et al.

Germline determinants of humoral immune response to HPV-16 protect against

oropharyngeal cancer. Nat Commun. 2021;12(1):5945.

47.   Denny JC, Rutter JL, Goldstein DB, Philippakis A, Smoller JW, Jenkins G, et al. The

"All of Us" Research Program. N Engl J Med. 2019;381(7):668-76.

48.   Ebrahimi E, Sangphukieo A, Park H, Gaborieau V, Ferreiro-Iglesias A, Diergaarde B,

et al. Cross-ancestral GWAS identifies 29 novel variants across Head and Neck Cancer

subsites. medRxiv. 2024:2024.11.18.24317473.

49.   Little P, Jo H, Hoyle A, Mazul A, Zhao X, Salazar AH, et al. UNMASC: tumor-only

variant calling with unmatched normal controls. NAR Cancer. 2021;3(4):zcab040.

50.   Montgomery ND, Parker JS, Eberhard DA, Patel NM, Weck KE, Sharpless NE, et al.

Identification of Human Papillomavirus Infection in Cancer Tissue by Targeted Next-

generation Sequencing. Appl Immunohistochem Mol Morphol. 2016;24(7):490-5.

51.   Seifert BA, O'Daniel JM, Amin K, Marchuk DS, Patel NM, Parker JS, et al. Germline

Analysis from Tumor-Germline Sequencing Dyads to Identify Clinically Actionable

Secondary Findings. Clin Cancer Res. 2016;22(16):4087-94.

52.   Patel NM, Michelini VV, Snell JM, Balu S, Hoyle AP, Parker JS, et al. Enhancing

Next-Generation Sequencing-Guided Cancer Care Through Cognitive Computing.

Oncologist. 2018;23(2):179-85.

53.   Patel NM, Jo H, Eberhard DA, Yin X, Hayward MC, Stein MK, et al. Improved Tumor

Purity Metrics in Next-generation Sequencing for Clinical Practice: The Integrated

Interpretation of Neoplastic Cellularity and Sequencing Results (IINCaSe) Approach. Appl

Immunohistochem Mol Morphol. 2019;27(10):764-72.

54.   Zhao X, Wang A, Walter V, Patel NM, Eberhard DA, Hayward MC, et al. Combined

Targeted DNA Sequencing in Non-Small Cell Lung Cancer (NSCLC) Using UNCseq and

NGScopy, and RNA Sequencing Using UNCqeR for the Detection of Genetic Aberrations in

NSCLC. PLoS One. 2015;10(6):e0129280.

55. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient

alignment of short DNA sequences to the human genome. Genome Biol. 2009;10(3):R25.

56. Saunders CT, Wong WS, Swamy S, Becq J, Murray LJ, Cheetham RK. Strelka:

accurate somatic small-variant calling from sequenced tumor-normal sample pairs.

Bioinformatics. 2012;28(14):1811-7.

57. Mose LE, Wilkerson MD, Hayes DN, Perou CM, Parker JS. ABRA: improved coding

indel detection via assembly-based realignment. Bioinformatics. 2014;30(19):2813-5.

58. Rettig EM, D'Souza G. Epidemiology of head and neck cancer. Surg Oncol Clin N

Am. 2015;24(3):379-96.

59. Maxwell JH, Grandis JR, Ferris RL. HPV-Associated Head and Neck Cancer: Unique

Features of Epidemiology and Clinical Management. Annu Rev Med. 2016;67:91-101.

60. Budhathoki S, Diergaarde B, Liu G, Olshan A, Ness A, Waterboer T, et al. A risk

prediction model for head and neck cancers incorporating lifestyle factors, HPV serology and

genetic markers. Int J Cancer. 2023;152(10):2069-80.

61. Lawlor DA, Tilling K, Davey Smith G. Triangulation in aetiological epidemiology.

International Journal of Epidemiology. 2017;45(6):1866-86.

62. Smith GD, Ebrahim S. 'Mendelian randomization': can genetic epidemiology

contribute to understanding environmental determinants of disease? International Journal of

Epidemiology. 2003;32(1):1-22.

63. Davey Smith G, Hemani G. Mendelian randomization: genetic anchors for causal

inference in epidemiological studies. Hum Mol Genet. 2014;23(R1):R89-98.

64. Sanderson E, Glymour MM, Holmes MV, Kang H, Morrison J, Munafò MR, et al.

Mendelian randomization. Nature Reviews Methods Primers. 2022;2(1):6.

65. Gormley M, Dudding T, Sanderson E, Martin RM, Thomas S, Tyrrell J, et al. A

multivariable Mendelian randomization analysis investigating smoking and alcohol

consumption in oral and oropharyngeal cancer. Nature Communications. 2020;11(1).

66. Gormley M, Dudding T, Kachuri L, Burrows K, Chong AHW, Martin RM, et al.

Investigating the effect of sexual behaviour on oropharyngeal cancer risk: a methodological

assessment of Mendelian randomization. BMC Medicine. 2022;20(1).

67. Gormley M, Yarmolinsky J, Dudding T, Burrows K, Martin RM, Thomas S, et al.

Using genetic variants to evaluate the causal effect of cholesterol lowering on head and

neck cancer risk: A Mendelian randomization study. PLoS Genetics. 2021;17(4 April 2021).

68. Gormley M, Dudding T, Thomas SJ, Tyrrell J, Ness AR, Pring M, et al. Evaluating the

effect of metabolic traits on oral and oropharyngeal cancer risk using Mendelian

randomization. eLife. 2023;12:e82674.

69. Thakral A, Lee JJW, Hou T, Hueniken K, Dudding T, Gormley M, et al. Smoking and

alcohol by HPV status in head and neck cancer: a Mendelian randomization study. Nature

Communications. 2024;15(1):7835.

70. Kachuri L, Saarela O, Bojesen SE, Davey Smith G, Liu G, Landi MT, et al. Mendelian

Randomization and mediation analysis of leukocyte telomere length and risk of lung and

head and neck cancers. Int J Epidemiol. 2019;48(3):751-66.

Declarations

Ethics approval and consent to participate

The VOYAGER project was approved by the International Association for Research on Cancer (IARC) Ethics Committee (IEC) (Project No.: 16 – 34). The five parent studies were all approved by the respective institutional review boards (IRBs) at each of the participating center: University of North Carolina at Chapel Hill (Study No.: 01-0390), University of Pittsburgh (STUDY19120160), The University Health Network (Project No.: 07-0521), Sinai Health System (REB No.: 08-0191) and the National Health Service, Health Research Authority (Project ID: 24028).

Consent for publication

Not applicable.

Availability of data and materials

Information on the VOYAGER Consortium can be found at https://voyager.iarc.who.int/. The contact page can be used for data and collaboration requests. Non-commercial research projects are generally approved if the proposed research complies with the signed agreements between studies and their research participants.

Genotype data for the oral and pharynx cancer OncoArray study have been deposited at the database of Genotypes and Phenotypes (dbGaP) and are available under controlled access under accession phs001202.v1.p1. Genotype data for the All of Us study are also available via dbGaP under controlled access under accession phs003225.v1.p1.

The oral and pharyngeal GWAS summary statistics by cancer site and world region have been deposited in the IEU Open GWAS platform (https://gwas.mrcieu.ac.uk/) under the

GWAS IDs: ieu-b-89, ieu-b-90, ieu-b-94, ieu-b-96, ieu-b-93, ieu-b-97, ieu-b-91, ieu-b-95 and ieu-b-98.

Competing interests

Scott Bratman reports grants from AstraZeneca, personal fees and equity from Adela, patents licensed to Adela and Roche, and service on advisory board for EMD Serono. Where members are identified as personnel of the International Agency for Research on Cancer/ World Health Organization, the authors alone are responsible for the views expressed in this article and they do not necessarily represent the decisions, policy or views of the International Agency for Research on Cancer / World Health Organization.

Funding

VOYAGER was funded by US National Institute of Dental and Craniofacial Research (NIDCR) grant R01DE025712 (PIs: Brennan, Diergaarde, and Hayes: The role of germline and somatic DNA mutations in oral and oropharyngeal cancers). Genotyping using the OncoArray and the All of Us array was performed at the Center for Inherited Disease (CIDR) and funded by NIDCR 1X01HG007780-0 and NIDCR/NCI X01HG010743, respectively. The Alcohol-Related Cancers and Genetic Susceptibility Study in Europe (ARCAGE) was funded by the European Commission's fifth frame-work program (QLK1-2001-00182), the Italian Association for Cancer Research, Compagnia di San Paolo/FIRMS, Region Piemonte and Padova University (CPDA057222). The Carolina Head and Neck Cancer Epidemiology (CHANCE) study was supported in part by the National Cancer Institute (R01CA90731). The Head and Neck 5000 study was a component of independent research funded by the National Institute for Health Research (NIHR) under its Programme Grants for Applied Research scheme (RP-PG-0707-10034). The views expressed in this publication are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. Core funding was also provided through awards from Above and Beyond, University Hospitals Bristol and Weston Research Capability Funding and the NIHR Senior Investigator

award to Professor Andy Ness. Human papilloma virus (HPV) serology was supported by a Cancer Research UK Programme Grant, the Integrative Cancer Epidemiology Programme (grant number: C18281/A19169). The University of Pittsburgh head and neck cancer case-control study was supported by US National Institutes of Health grants P50CA097190 and P30CA047904. The MSH-PMH study was supported by the Canadian Cancer Society Research Institute and the Princess Margaret Head & Neck Translational Research Program, with philanthropic funds from Joe's Team and the Wharton, Elia, Riley, and Tozer families. G.L. is funded by the Alan B. Brown Chair in Molecular Genomics and the Lusi Wong Foundation Fund.

Author contributions

Analyses for this manuscript were conducted by M.G. and S.V. The manuscript was drafted by M.G., S.V. and A.A. All authors contributed to the interpretation of the results and critical revision of the manuscript. All authors read and approved the final manuscript.

Acknowledgements

The authors would like to thank all study participants.

MAIN TABLES

Table 1. Clinical and demographic characteristics of the oral and oropharyngeal cancer cases and controls included in VOYAGER, stratified by subsite and HPV status

		Total OPC	HPV(+) OPC	HPV(-) OPC	HPV(+) vs. HPV(-) OPC	OC	Total OPC vs. OC	Total OPC + OC cases	Controls	Total OPC + OC cases vs. controls
		N= 3,514	N= 2,138	N= 1,146	P-value	N= 2,975	P-value	N= 6,489	N= 3,297	P-value
Sex (%)										
	Female	705 (20)	370 (17)	285 (25)		1,071 (36)		1,776 (27)	1,213 (37)	
	Male	2,809 (80)	1,768 (83)	861 (75)	<0.0001	1,903 (64)	<0.0001	4,712 (73)	2,084 (63)	<0.0001
	Unknown	0	0	0		1		1	0	
Age at diagnosis (in yrs)	Mean (SD)	59 (9.6)	58 (8.9)	60 (9.8)		61 (12.2)		60 (10.8)	59 (11.9)	
	[min, max]	[33, 84]	[26, 85]	[21, 94]	<0.0001	[18, 94]	<0.0001	[18, 94]	[17, 91]	0.308
Ethnicity (%)										
	White	2,947 (94)	1,963 (97)	828 (88)		2,271 (91)		5,218 (93)	2,374 (91)	
	Black	93 (3)	14 (<1)	68 (7)		103 (4)		196 (3)	183 (7)	
	Asian	53 (1)	20 (1)	31 (3)		89 (4)		142 (3)	4 (<1)	
	Latino/ Hispanic	7 (<1)	5 (<1)	2 (<1)		2 (<1)		9 (<1)	0 (0)	
	American Indian	2 (<1)	1 (<1)	1 (<1)		1 (<1)		3 (<1)	2 (<1)	
	Other	28 (1)	14 (<1)	11 (<1)	<0.001	16 (<1)	<0.0001	44 (<1)	6 (<1)	<0.0001
	Unknown	384	121	205		493		877	728	
Geographic region (%)										
	Canada	903 (26)	542 (25)	263 (23)		567 (19)		1,470 (23)	975 (30)	
	USA	735 (21)	436 (21)	222 (19)		841 (28)		1,576 (24)	1,596 (48)	
	Europe	1,876 (53)	1,160 (54)	661 (58)	0.156	1,567 (53)	<0.0001	3,443 (53)	726 (22)	<0.0001
Education (%)										
	None/ some school	880 (30)	472 (26)	379 (42)		847 (34)		1,727 (32)	368 (12)	
	High school	925 (32)	566 (32)	275 (30)		827 (33)		1,752 (32)	753 (23)	
	Postsecondary	1,081 (38)	755 (42)	260 (28)	<0.0001	834 (33)	0.003	1,915 (36)	2,114 (65)	<0.0001
	Unknown	628	345	232		467		1,095	62	

Stage (AJCC 7th ed.) (%)	I	165 (5)	45 (2)	112 (10)		800 (28)		965 (15)	NA	NA
	II	292 (8)	115 (6)	159 (14)		636 (22)		928 (15)		
	III	504 (15)	280 (13)	191 (17)		326 (11)		830 (13)		
	IV	2,481 (72)	1,675 (79)	652 (59)	<0.0001	1,110 (39)	<0.0001	3,591 (57)		
	Unknown	72	23	32		103		175		
HPV status (%)	Negative	1,146 (35)	0 (0)	1,146 (100)		1,702 (96)		2,848 (56)	NA	NA
	Positive	2,138 (65)	2,138 (100)	0 (0)	<0.0001	66 (4)	<0.0001	2,204 (44)		
	Unknown	230	0 (0)	0 (0)		1,207		1,437		
Treatment (%)	Radio + chemo	1,716 (49)	1,155 (55)	462 (40)		168 (6)		1,884 (30)	NA	NA
	Surgery	209 (6)	64 (3)	124 (11)		1,148 (47)		1,357 (22)		
	Surgery + radio + chemo	562 (16)	401 (18)	130 (11)		449 (15)		1,011 (16)		
	Surgery + radio	356 (10)	195 (9)	142 (12)		742 (25)		1,098 (18)		
	Radiotherapy	594 (17)	298 (14)	250 (22)		116 (4)		710 (11)		
	Palliative care	28 (<1)	10 (<1)	12 (1)		12 (<1)		40 (<1)		
	Chemotherapy	12 (<1)	3 (<1)	8 (<1)		6 (<1)		18 (<1)		
	No treatment	37 (<1)	12 (<1)	18 (2)	<0.0001	64 (2)	<0.0001	101 (2)		
Vital status (%)	Alive	2,407 (68)	1,711 (80)	569 (50)		1,677 (56)		4,084 (63)	NA	NA
	Dead	1,107 (32)	427 (20)	577 (50)	<0.0001	1,298 (44)	<0.0001	2,405 (37)		
Cause of death (%)	HNC	620 (60)	239 (60)	325 (61)		711 (59)		1,331 (60)	NA	NA
	Other cancer	111 (11)	33 (8)	65 (12)		130 (11)		241 (11)		
	Other disease	296 (29)	129 (32)	143 (27)	0.057	359 (30)	0.843	655 (29)		
	Unknown	2,487	1,737	613		1,775		4,262		

Key: HPV, human papilloma virus; OPC, oropharyngeal cancer; OC, oral cancer; radio, radiotherapy; chemo, chemotherapy.

HPV16 E6 serology was prioritized as a marker of HPV(+) oropharyngeal cancer. When serology was missing, two concordant tumor markers, p16 immunohistochemistry (IHC) and high-risk HPV DNA in-situ hybridization (ISH) were required to determine HPV status.

Table 2. Information on established risk factor behavior in VOYAGER, stratified by subsite and HPV status

		Total OPC	HPV(+) OPC	HPV(-) OPC	HPV(+) vs. HPV(-) OPC P-value	OC	Total OPC vs. OC P-value	Total OPC + OC cases	Controls	Total OPC + OC cases vs. controls P-value
		N= 3,514	N= 2,138	N= 1,146		N= 2,975		N= 6,489	N= 3,297	
Smoking status (%)	Current	862 (28)	277 (15)	498 (51)		988 (37)		1,850 (32)	545 (16)	
	Former	1,425 (46)	949 (51)	381 (39)		1,001 (38)		2,426 (42)	1,298 (40)	
	Never	787 (26)	638 (34)	103 (10)	<0.0001	671 (25)	<0.0001	1,458 (26)	1,437 (44)	<0.0001
	Unknown	440	274	164		315		755	17	
Alcohol drinking status (%)	Current	1,984 (64)	1,212 (65)	643 (65)		1,618 (61)		3,602 (62)	1,888 (58)	
	Former	463 (15)	233 (12)	180 (18)		387 (14)		850 (15)	666 (20)	
	Never	376 (12)	230 (12)	95 (10)		437 (16)		813 (14)	722 (22)	
	Never/ Former *	278 (9)	211 (11)	67 (7)	<0.0001	228 (9)	<0.0001	506 (9)	0 (0)	<0.0001
	Unknown	413	252	161		305		718	21	
Body mass index, in kg/m² (%)	Healthy (18.5 to <25)	1,034 (35)	508 (28)	452 (47)		1,057 (42)		2,091 (38)	433 (27)	
	Underweight (<18.5)	91 (3)	18 (<1)	68 (7)		96 (4)		187 (3)	12 (<1)	
	Overweight (25.0 to <30)	1,080 (36)	744 (40)	280 (29)		820 (33)		1,900 (35)	623 (39)	
	Obese (>30)	782 (26)	579 (31)	159 (17)	<0.0001	523 (21)	<0.0001	1,305 (24)	528 (33)	<0.0001
	Unknown	527	289	187		479		1,006	1,701	

Key: HPV, human papilloma virus; OPC, oropharyngeal cancer; OC, oral cancer.

* Never/Former category used when participants were only asked if they were a current drinker (HN5000 study). For these participants there was no information available to determine whether they were a former or a never drinker if they were not a current drinker.

HPV16 E6 serology was prioritized as a marker of HPV(+) oropharyngeal cancer. When serology was missing, two concordant tumor markers, p16 immunohistochemistry (IHC) and high-risk HPV DNA in-situ hybridization (ISH) were required to determine HPV status.

Table 3. Survival outcomes for the oral and oropharyngeal cancer cases in VOYAGER, stratified by subsite and HPV status

	OPC HPV Unknown	HPV(+) OPC	HPV(-) OPC	OC	Total OPC + OC
Overall survival					
N (number of cases)	230	2,138	1,146	2,975	6,489
Median Survival time, yrs	10.3 years	14.3 years	5.6 years	7.8 years	9.8 years
Probability of survival at 5y (95%CI), %	62.8 (56.7, 69.5)	81.1 (79.4, 82.9)	53.1 (50.2, 56.2)	60.7 (58.9, 62.5)	66.1 (65.0, 67.4)
Log-rank p-value	<0.001				
Progression-free survival					
N (number of cases)	164	1,731	894	2,272	5,061
Median progression-free survival time, yr	3.8 years	10.5 years	3.5 years	5.1 years	6.3 years
Probability of progression-free survival at 5y (95%CI), %	47.2 (40.1, 55.7)	74.9 (72.7, 77.1)	43.7 (40.2, 47.3)	50.5 (48.3, 52.7)	57.4 (56.0, 58.9)
Log-rank p-value	<0.001				
Disease-specific survival*					
N (number of cases)	220	2,112	1,102	2,877	6,311
Disease-specific survival at 5y (95%CI), %	72.7 (66.8, 79.2)	88.4 (87.0, 89.8)	67.5 (64.6, 70.6)	74.6 (72.9, 76.3)	78.1 (77.0, 79.2)
Gray's p-value	<0.001				

Key: HPV, human papilloma virus; OPC, oropharyngeal cancer; OC, oral cancer; Overall, including oral and oropharyngeal cancers.

Progression-free survival data was not available for the CHANCE study.
*Median not reached for disease-specific survival.

HPV16 E6 serology was prioritized as a marker of HPV(+) oropharyngeal cancer. When serology was missing, two concordant tumor markers, p16 immunohistochemistry (IHC) and high-risk HPV DNA in-situ hybridization (ISH) were required to determine HPV status.

MAIN FIGURES

Figure 1. Overview of the five studies included in VOYAGER.

Key: HPV, human papilloma virus; OPC, oropharyngeal cancer (green); OC, oral cancer (blue); Other sites (black).
Created in BioRender. Gormley, M. (2025) https://BioRender.com/j07p735

Figure 2. a) Overall survival, b) progression-free survival, and c) disease-specific survival for oral and oropharyngeal cancer cases in VOYAGER.

Figure 3. Overlapping data available in VOYAGER. Genotyping data includes cases and controls, tumor sequencing samples are for oropharyngeal and oral cavity cases only and clinical data includes all available cases and controls

Created in BioRender. (2025) https://BioRender.com/h48b673

Figure 4. Viral host interactions suggest HLA loci that are specific for HPV16 viral proteins