Unbiased human genomic characterization of polyglutamine disorder genes to inform therapeutic strategies

Kevin Lucy Namuli[1,2], Britt I. Drögemöller[3], Galen E.B. Wright[1,2]

Author affiliations:

[1]Department of Pharmacology and Therapeutics, Rady Faculty of Health Sciences, University of Manitoba, Winnipeg, Canada.

[2]PrairieNeuro Research Centre, Kleysen Institute for Advanced Medicine, Health Sciences Centre and Rady Faculty of Health Sciences, University of Manitoba, Winnipeg, Canada

[3]Department of Biochemistry and Medical Genetics, Rady Faculty of Health Sciences, University of Manitoba, Winnipeg, Canada.

Correspondence to: Galen E.B. Wright

University of Manitoba,

Department of Pharmacology and Therapeutics

Winnipeg, MB, Canada

Phone: +1 (204) 272-3130

galen.wright@umanitoba.ca

Word count: 3,822

Running title: Genomic Insights Inform PolyQ Therapeutics

Abstract

Background: Polyglutamine (polyQ) disorders, such as Huntington disease (HD) and several spinocerebellar ataxias, are severe neurological disorders caused by repeat expansions of the glutamine codon. These conditions lack effective treatments, with therapeutic research focused on pathogenic gene knockdown. We aimed to profile these genes using diverse human genomic data to guide therapeutic strategies by identifying new biology and assessing off-target effects of knocking these genes down.

Methods: We conducted an unbiased phenome-wide study to identify human traits and diseases linked to polyQ disorder genes (Open Targets L2G>0.5). Network analyses explored shared trait associations and overlapping biological processes among these genes. Lastly, we assessed the theoretical druggability of polyQ disorder genes using recently identified features predictive of clinical trial success and compared them to repeat expansion (HD) modifier genes.

Results: We identified 215 human phenotype/polyQ disorder gene associations from 3,095 studies, indicating potential adverse effects from gene knockdown. Shared trait associations among polyQ disorder genes suggested overlapping biological processes despite distinct functions. Drug target profile analysis revealed an unfavorable risk profile for polyQ disorder genes, particularly *ATN1*, *ATXN1*, *ATXN7*, and *HTT*, due to genomic features such as constraint, molecular interactions, and tissue specificity. PolyQ disorder genes also showed significantly more safety-related risks than HD modifier genes ($P=7.03 \times 10^{-3}$).

Conclusion: Our analyses emphasize the pleiotropic nature of polyQ disorder genes, highlighting their potential risks as drug targets due to safety concerns. These findings reinforce the importance of exploring alternate therapeutic strategies, such as targeting genetic modifier genes, to mitigate these challenges.

Introduction

Polyglutamine (polyQ) disorders are a group of ten heritable neurodegenerative conditions characterized by dynamic repeat expansions of the glutamine codon in distinct genes.[1] These include Huntington disease (HD), spinal and bulbar muscular atrophy (SBMA), dentatorubral-pallidoluysian atrophy (DRPLA), and several spinocerebellar ataxias (i.e., SCAs: 1-3, 6, 7 and 17).[1] Collectively, they affect approximately 1-10 individuals per 100,000 people, contributing significantly to the increasing burden of neurological disorders globally.[2] Typically, the onset of these disorders occurs in the third or fourth decade of life, presenting with a combination of motor and behavioural impairments that progressively worsen and result in death in some conditions.[3]

To date, some progress has been made in elucidating the pathogenic mechanisms of polyQ disorders. Unfortunately, these disorders remain without a cure, with current pharmacological treatments focused on alleviating the symptoms of the disorders.[4,5] While gene knockdown strategies, such as antisense oligonucleotides (ASOs), have been the focus of current research and some of these ASOs have advanced to clinical trial stages, efficacy and safety issues remain substantial barriers to their implementation in the clinic.[4,6–8] For example, the ASO tominersen, which was designed to reduce *HTT* expression in HD, demonstrated adverse events and failed to achieve efficacy. As tominersen suppresses the production of wild-type *HTT*,[9,10] this disrupts the normal functions of the gene, highlighting the challenges associated with the therapeutic targeting of polyQ genes. This is particularly important given that all polyQ disorders present in an autosomal dominant manner, except for SBMA.

The role of adverse events in clinical trial failures underscores the importance of understanding the biological consequences of targeting specific genes when designing therapeutics.[11] In line with this, Plenge *et al*.[12] highlighted the importance of evaluating potential adverse clinical consequences *a priori* by identifying human phenotypes associated with genetic variation related to drug target genes.[12] The increasing availability of extensive genome-wide association studies (GWAS) and related repositories, assisted by fine-mapping techniques that link genetic associations to suspected causal genes,[13,14] allows for systematic and unbiased identification of these gene-trait links. For example, we have previously shown that common genetic variation related to *HTT* is associated with cognitive function in large human cohorts.[15,16]

Associations between polyQ disorder genes and human traits can also provide insights into the biological processes that these genes are involved in, aiding in the identification of additional potential therapeutic targets. Several studies have demonstrated that drugs with prior genetic evidence for the role that the gene targets play in the clinical phenotype being treated are more likely to succeed in clinical trials.[14,17–21] Notably, a recent study also identified additional genetic factors associated with clinical trial outcomes,[20] showing that genes displaying human genetic constraint, several interacting partners and broad gene expression were associated with clinical trial stoppage. These studies highlight the importance of including human genomic and functional information into drug development pipelines.

The challenges highlighted above emphasize the need to comprehensively evaluate polyQ disorder genes to identify potential off-target effects and study genomic features linked to therapeutic failures. Here, we aimed to characterize polyQ disorder genes using extensive human genomic data to inform therapeutic studies and disease biology. First, we analyzed and prioritized human trait and disease associations with common genetic variation in polyQ disorder genes using large repositories. We unbiasedly profiled important human genomic information through this approach to identify potential off-target effects. We also uncovered unique and shared biological features by examining the additional phenotypes associated with these genes outside of the polyQ disorders themselves. Finally, we evaluated the theoretical druggability of the polyQ disorder genes, ranking them according to features related to past increased clinical trial success and comparing them to the repeat expansion modifier genes that influence disease onset. Our comprehensive analysis enhances our understanding of the underlying biology of polyQ disorders and their pleiotropic nature and offers potential insights into the consequences of targeting the associated genes for polyQ disorder therapeutics. Findings from our study are of broad relevance to therapeutic strategies for other repeat expansion disorders, of which more than 50 have been identified to date.

Methods

Ethical approval and gene selection

This study obtained ethical approval from the University of Manitoba Bannatyne Research Ethics Board (H2022:354 HS25743). We profiled all known polyQ disorder genes (*n*=10, i.e., *AR*, *ATN1*, *ATXN1*, *ATXN2*, *ATXN3*, *ATXN7*, *CACNA1A*, *HTT*, *TBP* and *THAP11*). For GWAS-based analyses, we excluded *AR* since it is located on the X chromosome, and current GWAS datasets do not include this information.

Identification of human traits and diseases associated with polyQ disorder genes

To identify human traits and diseases associated with the polyQ disorder genes beyond their primary disorders, an unbiased, comprehensive phenome-wide study was conducted by analyzing data from the Open Targets Genetics database (OTG) (version 8).[13] This repository contains approximately 132,893 curated GWAS association signals from various studies and genomic biobanks such as FinnGen and the UK Biobank, including 8,317 human GWAS summary statistics (data extracted January 2024). To curate OTG data, we used the related application programming interface, implemented using GraphQL v2.0[13] and executed in R 4.3.3.

We accessed data specific to the polyQ disorder genes, using Ensembl identifiers[22] to fetch relevant genomic studies. We identified human traits where significant associations could be attributed to polyQ disorder genes using the locus-to-gene (L2G) fine-mapping tool.[23] L2G is designed to refine genome-wide significant GWAS signals ($P<5\text{x}10^{-8}$), identifying likely causal genes attributed to these signals.[13] This tool uses a scoring metric that quantifies the strength of evidence for gene-trait associations, ranging from 0 to 1.[13] Higher scores signify increased support for a particular gene-trait association. Therefore, traits overlapping with the polyQ disorder genes with a score greater than 0.5 were kept for further analysis as recommended by the developers of this tool.[23]

Finally, to examine other severe Mendelian disorders and phenotypes associated with polyQ disorder genes outside of those caused by the CAG repeat expansion, we profiled the Online Mendelian Inheritance in Man (OMIM) database.[24] We analyzed the OMIM genemap2 file (accessed June 2022) containing 5,779 phenotype entries.

Assessing shared traits among polyQ disorder genes

To investigate whether the polyQ disorder genes were linked to shared traits and the degree of this overlap, we performed a network analysis with R packages *ggraph* (version 2.1.0) and *arcdiagram* (version 0.1.12). All gene-trait associations with an L2G>0.5 were included in the network analysis. In the resulting network, each node represented a polyQ disorder gene, and each edge (link) represented a bidirectional relationship shared between two polyQ disorder genes. This network analysis aimed to identify networks of genes linked to particular traits, thereby providing insights into underlying mechanisms and potentially uncovering shared biology.

Prioritization of informative L2G-significant human trait associations for polyQ disorder genes

To prioritize individual traits and disease signals for the polyQ disorder genes, count-based traits (e.g., blood cell types) were removed. This was performed to minimize non-specific signals, as such traits are associated with numerous generic associations across the genome. Further, we restricted our analyses to include only gene-trait pairs derived from peer-reviewed PubMed studies to ensure the validity and robustness of the study findings.

Variant annotation and unique signal identification

We annotated the predicted functional effect of index genetic variants related to GWAS trait signals to understand their potential mechanistic contribution to the observed gene-trait associations. The Ensembl Variant Effect Predictor was used to annotate the L2G significant signals using the *most severe* consequence flag related to each polyQ disorder gene. SpliceAI was used to predict whether the variants alter splicing.[25] Genotype-Tissue Expression (GTEx) data was used to determine whether variants were linked to gene expression changes of the related polyQ disorder genes. Further, we attempted to identify independent signals at each gene by considering linkage disequilibrium (LD) between index variants. Independent signals, defined as haplotypes, were determined by pruning index variants with $r^2>0.5$ using the 1,000 Genomes European super-population with LDLink SNPClip.[26] Each haplotype was described as a set of variants in LD (i.e., $r^2>0.5$) with each other. A single tag variant was considered a proxy for all the variants in high LD and was used to capture these haplotypes. The Finnish population was excluded from this analysis to minimize potential confounding factors associated with extended haplotype blocks.[27] Resulting

120 haplotypes were assigned to prioritized polyQ disorder gene-trait pairs to determine unique signals

121 at each locus.

Assessment of the theoretical druggability and risk profiles of the polyQ disorder genes

To thoroughly evaluate polyQ disorder genes for theoretical druggability and drug target profiles

related to the risk of clinical trial failure, we assessed recently identified features associated with

increased clinical trial success.[20] Genes were categorized as favorable or unfavorable under each

category based on previously determined thresholds.[20] The following features were included in

our analyses:

(i) Genomic constraint: We examined genes' tolerance to loss-of-function mutations using

sequencing data from the Genome Aggregation Database, which contains 125,748

exomes and 15,708 genomes (gnomAD v2.1.1; accessed July 2022).[28] We used the

probability of loss-of-function intolerance (pLI) metric, which ranges from 0 to 1, with

more constrained genes having higher scores. Genes with a high degree of constraint

were categorized as unfavourable, indicated by a rank greater than 0.9 (corresponding

to *"extremely loss of function intolerant"* genes).[29]

(ii) Tissue expression: We accessed RNA tissue specificity data from the Human Protein

Atlas (HPA) database (accessed February 2023),[30] which uses data from the HPA and

GTEx to classify genes into five categories (i.e., *"tissue enriched"*, *"group enriched"*,

"tissue enhanced", *"low tissue specificity"* and *"not detected"*) based on fold change

in mRNA expression across 37 organ systems.[31] Genes whose RNA tissue specificity

was classified as *"low tissue specificity"* were ranked unfavorable as they are broadly

expressed.

Molecular interactions (MI): We used the Open Targets Platform to access data from

the IntAct database[32] related to MI evidence for the genes (accessed October 2024).

We used the MI score, which ranges from 0 to 1 (scores closer to 1 are higher

confidence) and is built using diverse data (i.e., publications, experimental detection

methods, and interaction types).[33] Genes with more than ten interacting partners with

MI scores exceeding 0.42 (considered high confidence[20]) were categorized as

unfavorable.

(iii) Druggability evidence: We assessed the Drug Gene Interaction Database (accessed July

2024)[34] to identify any known or potentially druggable genes among the polyQ disorder

genes. We assigned a favorable "*druggable any*" metric to polyQ disorder genes that
had any known or predicted evidence of interaction with drugs from this database.
153
Finally, we compared the number of unfavorable genomic features associated with increased
safety-related risk (i.e., constraint, expression and interactions) to these same metrics that have
been reported in our recent analysis of HD modifier genes[35] using the Wilcoxon test accompanied
by the calculation of the effect size (*r*) to quantify the magnitude of the observed difference. This
analysis aimed to assess whether targeting one of these gene sets presents a theoretically greater
safety risk than the other. *P*<0.05 was considered significant in these analyses.

Data sharing
The data used in the analyses described in this study are publicly accessible to researchers
through the consortia that generated them. Analysis scripts for data accession, analysis and
visualization are accessible at https://github.com/Wright-Lab-Neurogenomics-
Research/polyQ_genes_characterization_analyses

Results

Profiling polyQ disorder genes for associations with common human traits and diseases

We identified 3,095 association signals (1,467 unique traits) surrounding polyQ disorder genes
from the OTG database. These traits were distributed among 20 categories based on health
conditions or biological processes, with the broad *measurement* category containing the most
associations (*n*=2,521). Applying a filtering threshold of L2G>0.5 retained seven polyQ disorder
genes (i.e., *ATXN1*, *ATXN2*, *ATXN3*, *ATXN7*, *CACNA1A*, *HTT* and *TBP*) and identified 215 gene-
trait pairs (149 unique traits), signifying associations with strong evidence for polyQ disorder gene
involvement in the trait (**Figure 1**). *ATXN1* had the highest number of L2G>0.5 significant traits
reported (*n*=57), predominantly related to hematological (measurement) phenotypes. *ATXN3* and
TBP were exclusively linked to hematological phenotypes, while *ATXN7* was associated solely
with disease-related phenotypes. Network analysis of the 215 polyQ disorder gene-trait pairs with
L2G>0.5 revealed that five of the seven polyQ disorder genes shared phenotypic traits. *TBP* and
CACNA1A did not share traits with any polyQ disorder genes. *ATXN1* had the greatest number of
traits shared with the other polyQ disorder genes. *HTT* and *ATXN1* displayed the most shared traits,

including educational attainment and cognitive ability, in addition to neutrophil and white blood

cell count.

We applied further prioritization steps to obtain high confidence and informative traits linked to

polyQ disorder genes, including removing count-based terms and studies not found in PubMed.

This analysis retained 46 significant traits (L2G>0.5) and five polyQ disorder genes: *HTT, ATXN1,*

ATXN2, ATXN7, and *CACNA1A* (**Figure 2**). Furthermore, filtering out duplicate traits for each

gene retained 37 unique gene-trait associations. These informative trait associations represent

diverse disease phenotypes, including neuropsychiatric traits (e.g., *HTT*-depression, *CACNA1A*-

positive affect, *ATXN7*-schizophrenia), metabolic disorders (e.g., *ATXN2*-type 1 diabetes,

HTT/ATXN7-type 2 diabetes) and autoimmune disorders (e.g., *ATXN1*-multiple sclerosis, *ATXN1*-

systemic lupus erythematosus). Additionally, they include several other non-neurological traits or

diseases, such as *ATXN1*-atrial fibrillation, *ATXN1*-/*ATXN7*-breast cancer, and *ATXN7*-cataracts.

PolyQ disorder genes, such as *ATXN1* and *HTT,* were associated with non-pathogenic neurogenetic

traits, including educational attainment, cognitive ability, and aging. This collective observation

highlights the pleiotropic features related to these genes outside of polyQ pathobiology.

Investigation and annotation of the variants linked to unique polyQ disorder gene-common

trait associations

The index variants linked with the high-confidence unique traits associated with the polyQ disorder

genes fall into three consequence categories: intronic (*n*=26), splice (*n*=1), and those variants

located in untranslated regions (*n*=1) (**Figure 2**). Notably, all traits associated with the splice and

untranslated region variants were linked to *HTT*. In contrast, traits related to the remaining polyQ

disorder genes (i.e., *CACNA1A, ATXN1, ATXN2* and *ATXN7*) were exclusively associated with

intronic variants. The *HTT* splice variant showed associations with educational attainment and its

noncognitive aspects. *ATXN7* was associated with several pathogenic traits, including type 2

diabetes, schizophrenia, breast cancer and cataracts, each associated with a unique variant (total

n=4).

The 37 unique gene-trait associations arose from 30 distinct signals. A variant located in the

untranslated region in *HTT* was associated with the highest number of distinct traits (*n*=5),

including type 2 diabetes and walking pace (**Table 1**). The remaining 29 unique variants resulted

in 21 independent signals (i.e., *ATXN1, n*=8; *HTT, n*=6; *ATXN7, n*=3; *ATXN2, n*=2; *CACNA1A,*

n=2) referred to as haplotypes. The *HTT* haplotype two (index variant: intronic rs61348208), which

we have previously reported,[16] had the largest number of traits (*n*=6) associated with it, including

both pathogenic (e.g., depression) and non-pathogenic traits (e.g., aging traits and parental

lifespan). *ATXN1* exhibited the highest number of independent signals (*n*=8 haplotypes), with

ATXN1 haplotype three (index variant: intronic rs909788) (*n*=5) demonstrating the most trait

associations. However, these were all linked to highly correlated cognitive and socio-economic

traits.

Profiling polyQ disorder genes for severe Mendelian diseases beyond canonical polyQ
disorders

A subset of the polyQ disorder genes (i.e., *AR, ATN1, ATXN2, CACNA1A, HTT, TBP*) had

phenotype entries in the OMIM database beyond the primary polyQ disorders they cause (**Table**

2). Rare Mendelian disorder associations included androgen insensitivity and hypospadias related

to *AR*, and neurological conditions linked to *HTT* (i.e., neurodevelopmental disorder Lopes-

Maciel-Rodan syndrome), *CACNA1A* (i.e., developmental and epileptic encephalopathy) and

ATN1 (i.e., congenital hypotonia, epilepsy, developmental delay, and digital anomalies). The genes

ATXN2 and *TBP* were also associated with susceptibility to Parkinson's disease, a common

neurological disorder, while *ATXN2* was also associated with susceptibility to amyotrophic lateral

sclerosis (ALS), a neurodegenerative disease. Finally, *AR* was also associated with prostate cancer

susceptibility in OMIM. These findings illustrate that therapeutic polyQ disorder gene knockdown

may impair critical developmental or neurological pathways.

Analysis of human genomic safety profiles and the theoretical druggability of polyQ disorder
genes highlights potential safety concerns associated with therapeutic targeting

We evaluated genomic features that predict clinical trial success to better understand the theoretical

implications of therapeutically perturbing polyQ disorder genes. Most of these genes showed

unfavorable profiles based on multiple criteria (**Figure 3**). Notably, four genes (i.e., *ATN1, ATXN1*,

ATXN7 and *HTT*) ranked unfavorably for the three categories associated with clinical trial stoppage

due to safety concerns (i.e., genetic constraint, broad expression, and several interactors). Further,

all polyQ disorder genes displayed at least one unfavorable signal for one of these three features.

All polyQ disorder genes, except for *THAP11*, displayed many molecular interactors (i.e., more

than ten protein-protein interactions in the IntAct database). This was especially apparent for *HTT*,

ATXN1, *ATN1* and *ATXN3*, which showed over 100 high-confidence interactions (i.e., ten times above the threshold used for classification). While five polyQ disorder genes fell within the theoretical *'druggable any'* category, this metric indicates whether the gene can be theoretically perturbed, not an indicator of safety. Compared to HD age-of-onset modifier genes, polyQ disorder genes displayed significantly more unfavorable features associated with an increased safety risk (P=7.03x10^{-3}; mean HD modifiers=1.0, n=9; mean polyQ disorders=2.2, n=10; effect size r=0.628, large).

Discussion

Gene knockdown strategies targeting the primary disease-causing genes in polyQ disorders are being extensively explored as potential therapies in these repeat expansion disorders.[4,6,7,36] However, these strategies have been unsuccessful in polyQ therapeutic clinical trials, failing to achieve the desired therapeutic value and resulting in adverse safety profiles. To better understand these challenges, we systematically profiled polyQ disorder genes using large-scale genomic databases, highlighting their pleiotropic roles and identifying genomic features associated with increased clinical trial failure. We demonstrated that polyQ disorder genes might be unfavorable targets since they are associated with genomic features linked to an increased failure of drug targets in clinical trials. Notably, *ATN1* (DRPLA), *ATXN1* (SCA1), *ATXN7* (SCA7), and *HTT* (HD) had particularly deleterious risk profiles in our analyses.

Our analysis of human GWAS revealed that polyQ disorder genes display extensive pleiotropy beyond their canonical roles in polyQ-related neurodegeneration, indicating potential off-target effects of therapeutic knockdown.[37] For example, *HTT, ATXN1, ATXN7* and *CACNA1A* were associated with neuropsychiatric disorders such as depression and schizophrenia. Additionally, metabolic traits like type 1 and type 2 diabetes were linked to *ATXN2* and *ATXN7*, which is in line with the fact that metabolic dysfunction is observed in several polyQ disorders.[38] *ATXN7* was exclusively associated with pathogenic traits, including schizophrenia, diabetes, breast cancer, and cataracts, highlighting the potential risks of targeting this gene. Our OMIM analyses also revealed disease links to polyQ disorder genes outside canonical repeat-mediated disorders. For example, *ATN1* and *HTT* have been linked to neurodevelopmental disorders, while *CACNA1A* has been linked to channelopathies with diverse neurological manifestations. However, susceptibility to

Parkinson's disease and ALS are findings correlated with longer CAG repeats,[39] suggesting

possible benefits of mutation-specific knockdown related to these phenotypes.

In addition to disease associations, polyQ disorder genes were linked to critical non-pathogenic

phenotypes through GWAS. We have previously reported that common *HTT* genetic variation is

associated with important neurobiological traits such as educational attainment and longevity,[16]

which we replicate here. This study showed that *ATXN1* is also linked to similar traits, including

educational attainment and cognitive abilities. When comparing the variant effects between related

GWAS and GTEx expression, alleles linked to lower *ATXN1* expression were also associated with

lower educational attainment and decreased cognitive abilities. These findings suggest that

therapeutically altering these genes might inadvertently affect crucial cognitive functions, which

should be assessed during future clinical trials. In addition to individual gene-trait associations,

network analyses revealed that specific trait associations were shared between polyQ disorder

genes, with *ATXN1*, *ATXN2,* and *HTT* displaying the greatest number of shared interactions. For

example, associations with metabolic traits (type 2 diabetes and cholesterol levels) were shared by

HTT, ATXN2 and *ATXN7* and susceptibility to breast cancer was shared by *ATXN1* and *ATXN7.*

These analyses indicate that specific polyQ disorder genes may be involved in common biological

pathways, as previously reported by others.[40,41]

While half of the polyQ disorder genes were found to be potentially challenging to target

therapeutically with traditional approaches, RNA-targeted therapeutics provide a flexible solution

by enabling the targeting of any gene product. Nonetheless, our human genomic safety profile

analyses revealed that polyQ disorder genes consistently rank unfavorably for predictors of clinical

trial success,[20] which could provide some insight into the mechanisms driving gene-lowering trial

failures.[7,8,42] Specifically, *ATN1*, *ATXN1*, *ATXN7* and *HTT* ranked poorly across all safety-related

categories. These four genes are integral to transcriptional regulation and are, therefore, essential

for several critical physiological processes.[38] Therapeutic targeting of these genes could disrupt

several important processes, leading to adverse effects.

Most of the polyQ disorder genes (i.e., 80%) displayed broad expression patterns, with this pattern

of expression associated with increased trial stoppage risks [reported odds ratio (OR): 1.29 95%;

confidence interval (CI): 1.19-1.39].[20] Additionally, half of the polyQ disorder genes are highly

constrained, with LoF intolerance (pLI>0.9) also associated with an increased risk of trial stoppage

(reported OR: 1.37; 95% CI: 1.27-1.48).[20] Further, all polyQ disorder genes, except for *THAP11*,

displayed more than ten high-confidence protein-protein interactions, indicating their central

physiological roles – an effect that can, in part, be attributed to the polyQ repeats themselves.[43]

Specifically, genes with between 11 and 20 partners (e.g., *CACNA1A*) are at an increased risk of

clinical trial stoppage (reported OR: 1.31; 95% CI: 1.19-1.44),[20] while those with >20 partners

(i.e., 80% of polyQ disorder genes) are at an even greater risk (reported OR: 1.38; 95% CI: 1.26-

1.52).[20] Finally, when we compared the genomic safety predictions for polyQ disorder genes to

those of genetic modifiers of the age of HD onset, we demonstrated that the modifier genes showed

more favourable genomic safety scores. This provides additional support to prioritize these

modifiers for future therapeutic research. Beyond their favourable drug target profiles, a further

benefit of modifier-based therapeutic targeting is the potential for cross-repeat expansion disorder

relevance due to shared pathobiological mechanisms of these modifier genes (e.g., somatic repeat

instability).[35,44–47] This may also be useful for rarer forms of repeat expansion disorders, where

assembling large cohorts of individuals required for a disorder-specific clinical trial may be

difficult.[36]

In conclusion, our findings indicate that the pleiotropic roles and adverse genomic profiles of

polyQ disorder genes might diminish their potential as therapeutic targets for their respective

conditions. However, it should be noted that these genes do have human genetic evidence for

involvement in their primary diseases, a feature associated with increased clinical trial success.[14]

Therefore, polyQ disorder genes should not be entirely abandoned as therapeutic targets, especially

with some encouraging results from recent early clinical trials.[48] Instead, increased focus should

be placed on assessing age-dependent implications of knockdown, mutant allele-specific

perturbation and gene editing approaches. In parallel, building on our findings that genetic

modifiers of the age of HD onset have favourable genomic safety scores, future research should

investigate therapeutics that target repeat expansion disorder genetic modifiers. In line with this,

the promising role of therapeutic targets that prevent somatic expansions has been further

emphasized by recent reports that show that the impact of the expanded repeat only becomes

evident late in the lifetime of vulnerable cell types.[49] Taken together, this study has revealed the

importance of incorporating diverse human genomics data into drug development pipelines to
develop safer and more effective treatments.
339

Acknowledgements

We thank the Natural Sciences and Engineering Research Council of Canada through the Canada
Research Chairs Initiative and Discovery Grant Program, as well as the Health Sciences Centre
Foundation Winnipeg, for the financial support provided to GEBW. BID was also supported
through the Canadian Institutes of Health Research Canada Research Chairs Initiative. The
Research Manitoba Graduate Studentship and Rady Faculty of Health Sciences Student Top-Up
supported NKL.

Author roles

NKL performed the formal analyses and wrote the manuscript. GEBW and BD designed the study
and wrote and edited the manuscript. All authors agreed on the final version of the paper.
351

 FIGURES

Figure 1. PolyQ disorder genes are pleiotropic and associated with several human traits and diseases in diverse phenotypic categories. (A) polyQ-associated genes and their corresponding polyQ disorders. Colors represent different genes used across panels and figures. **(B)** Human traits and disorders linked to polyQ disorder genes (i.e., L2G>0.5) obtained from an analysis of genome wide association studies in the Open Targets Genetics Database. The highest number of gene-trait pairs were identified for *ATXN1*, *ATXN2*, and *HTT* and most traits were classified as measurements. **(C)** Heatmap of non-measurement-related traits (L2G>0.5) for each polyQ disorder gene, depicting several associations for disease including those related to cardiovascular system (e.g., atrial fibrillation), pancreas (e.g., type 2 and type 2 diabetes) and nervous system. **(D)** Network graph illustrating shared polyQ disorder gene-trait associations. Larger circles represent the number of traits associated with a particular gene, while thicker arcs indicate more shared trait associations between genes. Notable shared traits include *HTT* and *ATXN1* with educational attainment and cognitive ability; *ATXN1* and *ATXN7* with breast cancer; *HTT* and *ATXN2* with depression; and *ATXN1*, *ATXN3* and *ATXN2* with HDL cholesterol levels. This network highlights the pleiotropic effects of polyQ disorder genes across neurological, metabolic, and oncological traits. **Abbreviations:** L2G; locus-to-gene; *DRPLA*, Dentatorubral-Pallidoluysian Atrophy; *HD*, Huntington Disease; *SBMA*, Spinal and Bulbar Muscular Atrophy; *SCA*, Spinocerebellar Ataxia.

Figure 2. High-confidence and informative human gene-trait associations linked to the polyQ disorder genes. (A) A total of 46 traits linked to five polyQ disorder genes remained after applying a filtering threshold of L2G>0.5, while excluding count-based terms (e.g., blood cell counts) and studies lacking PubMed IDs. **(B)** L2G scores for individual high confidence trait associations (duplicate traits removed). PolyQ disorder genes exhibit associations with both pathogenic and non-pathogenic characteristics, across neurological, metabolic, and immune-related traits. **(C)** *HTT* and **(D)** *ATXN1* displayed the highest number of gene-trait pairs and are linked to neurological disorders and neurobiological phenotypes. In these panels, high confidence gene-trait associations are annotated and plotted by variant position and L2G score. Color indicates predicted variant consequence (*most severe*) of index variant for each trait association. Count based traits were plotted but not annotated. The grey-shaded segment represents polyQ disorder gene locus (GRCh38-genome build coding sequence start-stop position). **Abbreviations:** L2G; locus-to-gene, UTR; Untranslated region, GRCh38; Genome Reference Consortium Human Reference 38.

354
355

Figure 3. Genetic features associated with clinical trial success suggest that polyQ disorder genes may not be favorable therapeutic targets. (**A**) Genetic constraint features, gnomAD probability of loss-of-function intolerance (pLI) and negative observed vs expected loss-of-function variant ratio, indicate high levels of the constraint of the polyQ disorder genes. pLI allows non-constrained genes (pLI<0.1) to be differentiated from highly constrained ones (pLI>0.9). The observed/expected ratio is a quantitative measure of how tolerant the genes are to certain classes of variations. (**B**) Tissue specificity of the polyQ disorder genes, with *AR* and *CACNA1A* displaying tissue-enhanced and tissue-enriched, respectively, while others show low specificity. (**C**) PolyQ disorder genes interact with several other proteins and only *THAP11* displays less than 11 interacting partners (associated with decreased levels of clinical trial stoppage). (**D**) Aggregated summary of human genomic features used to rank the theoretical druggability and genomic safety of polyQ disorder genes. Genes ranked more unfavorable related to theoretical druggability and genomic safety, such as *HTT*, *ATN1*, *ATXN1* and *ATXN7*, are on the right of the plot. Individual favorable metrics are represented by the teal green tiles, while the deep purple tiles represent unfavorable features. (**E**) PolyQ disorder genes (yellow) display significantly more (*P*=0.007) unfavorable genomic safety scores based on constraint, tissue specificity and interactions (maximum score=3, increased risk of clinical trial stoppage) than Huntington disease modifier genes (blue). Mean scores are indicated by red line (mean polyQ=2.2, mean HD modifier genes=1.0).

TABLES

Table 1 High-confidence GWAS associations linked to polyQ disorder genes are associated with pathogenic and non-pathogenic human traits. L2G prioritization highlights traits linked to unique independent signals identified as haplotypes associated with polyQ disorder genes extending beyond the primary traits directly caused by the pathogenic CAG repeat expansions in these the genes.

Signal (Tag variant)[a]	Reported trait (PMID)	Variant ID	Annotation[b]	Effect AF EUR[c]	GWAS index P-value	GWAS index effect (95% CI)	L2G	CADD score
ATXN1: haplotype one (rs719316)	Multiple sclerosis (31604244)	rs719316	intronic	T: 0.529	2.00×10^{-13}	1.07 (1.05-1.09)[d]	0.79	5.92
ATXN1: haplotype two (rs17603856)	Systemic lupus erythematosus (27399966)	rs17603856	intronic	G: 0.346	3.00×10^{-12}	0.88 (0.85-0.91)[d]	0.79	10.94
ATXN1: haplotype three (rs909788)	Cognitive ability, years of educational attainment or schizophrenia (31374203)	rs6459480	intronic	A: 0.509	2.00×10^{-10}	NA	0.69	1.82
	Household income (31844048)	rs6459480	intronic	A: 0.509	5.00×10^{-10}	-0.01 (-0.02--0.01)	0.70	1.82
	Educational attainment (30038396)	rs9297016	intronic	G: 0.576	1.00×10^{-09}	-0.01 (-0.01--0.006)	0.68	2.57
	Educational attainment: years of education (27225129)	rs7772172	intronic	G: 0.576	1.00×10^{-08}	-0.01 (-0.02--0.01)	0.68	5.02

	Educational attainment (years of education) (30038396)	rs909788	intronic	C: 0.545	4.00×10^{-08}	-0.01 (-0.011--0.005)	0.77	5.75
ATXN1: haplotype four (rs6459472)	Central corneal thickness (32528159)	rs6459472	intronic	G: 0.573	3.00×10^{-09}	NA	0.72	2.97
ATXN1: haplotype five (rs9367926)	Male-pattern baldness (30573740)	rs6915310	intronic	T: 0.182	5.00×10^{-20}	0.04 (0.03 - 0.05)	0.72	4.27
	Balding type 1 (30595370)	rs9367926	intronic	G: 0.180	6.00×10^{-12}	NA	0.71	6.59
ATXN1: haplotype six (rs2237199)	Subcutaneous adipose tissue attenuation (27918534)	rs2237199	intronic	A: 0.112	1.00×10^{-08}	5.67 (3.73 - 7.61)	0.57	3.49
ATXN1: haplotype seven (rs7770062)	Atrial fibrillation (30061737)	rs73366713	intronic	A: 0.131	1.53×10^{-25}	0.90 (0.88-0.92)[d]	0.55	2.84
	Atrial fibrillation (29892015)	rs7770062	intronic	A: 0.131	9.00×10^{-21}	0.90 (0.88-0.92)[d]	0.54	2.33
ATXN1: haplotype eight (rs3819405)	Breast cancer (29059683)	rs3819405	intronic	T: 0.340	1.65×10^{-08}	0.96 (0.95-0.97)[d]	0.53	12.10
ATXN2: haplotype one (rs653178)	Celiac disease (20190752)	rs653178	intronic	T: 0.534	6.00×10^{-14}	0.83 (0.78-0.87)[d]	0.51	1.59
ATXN2: haplotype two (rs848130)	Type 1 diabetes (34012112)	rs848130	intronic	A: 0.192	1.60×10^{-14}	0.85 (0.82-0,89)[d]	0.67	2.02
ATXN7: haplotype one (rs832190)	Schizophrenia (31740837)	rs832190	intronic	T: 0.603	4.00×10^{-08}	NA	0.51	0.58

ATXN7: haplotype two (rs3821902)	Breast cancer (29059683)	rs3821902	intronic	G: 0.150	2.99×10^{-12}	1.06 (1.05-1.08)[d]	0.54	0.81
ATXN7: haplotype three (rs13434089)	Type 2 diabetes (32541925)	rs13434089	intronic	C: 0.141	4.00×10^{-31}	-0.06 (-0.07--0.05)	0.55	4.59
CACNA1A: haplotype one (rs5021328)	Appendicular lean mass (33097823)	rs5021328	intronic	C: 0.644	6.60×10^{-09}	-0.01 (-0.02--0.01)	0.86	0.63
CACNA1A: haplotype two (rs16003)	Positive affect (30643256)	rs16003	intronic	T: 0.463	3.00×10^{-08}	0.01 (0.00 - 0.01)	0.84	8.26
	Depressive symptoms (30643256)	rs16003	intronic	T: 0.463	3.00×10^{-08}	0.01 (0.00 - 0.01)	0.84	8.26
HTT: haplotype one (rs363096)	Noncognitive aspects of educational attainment (33414549)	rs363096	splice	C: 0.591	1.00×10^{-12}	0.05 (0.04 - 0.06)	0.72	6.29
	Educational attainment: years of education (30595370)	rs363096	splice	C: 0.591	1.00×10^{-10}	NA	0.74	6.29
HTT: haplotype two (rs61348208)	Depression (30718901)	rs7685686	intronic	G: 0.433	6.00×10^{-15}	0.983 (0.979-0.987)[d]	0.59	6.39
	Frailty index (34431594)	rs82334	intronic	C: 0.323	3.10×10^{-10}	-0.02 (-0.03--0.02)	0.54	4.13
	Parental lifespan (30642433)	rs61348208	intronic	T: 0.389	6.00×10^{-09}	0.23 (0.15 - 0.31)	0.63	5.46

	Trait (PMID)	rsID	Function	Allele: freq	P-value	Effect (CI)		
	Gastroesophageal reflux disease (34187846)	rs7685686	intronic	G: 0.433	1.10×10^{-08}	0.97 (0.96-0.98)[d]	0.54	6.39
	Aging traits: health span, parental lifespan or longevity (32678081)	rs61348208	intronic	T: 0.389	3.00×10^{-08}	NA	0.62	5.46
	Vertex-wise sulcal depth (34910505)	rs2071703	intronic	C: 0.325	3.00×10^{-08}	5.57 (3.60 - 7.54)	0.6	2.52
HTT: haplotype three (rs113928896)	Highest math class taken (30038396)	rs113928896	intronic	T: 0.143	2.00×10^{-09}	0.02 (0.01 - 0.02)	0.55	0.32
HTT: haplotype four (rs362307)	Automobile speeding propensity (30643258)	rs362307	UTR	T: 0.063	6.00×10^{-14}	-0.03 (-0.04--0.02)	0.51	3.23
	Worry/vulnerability: special factor of neuroticism (30867560)	rs362307	UTR	T: 0.063	1.00×10^{-09}	0.01 (0.005 - 0.011)	0.54	3.23
	Walking pace (33128006)	rs362307	UTR	T: 0.063	1.00×10^{-09}	-0.01 (-0.02--0.01)	0.54	3.23
	Type 2 diabetes (30297969)	rs362307	UTR	T: 0.063	1.00×10^{-09}	1.08 (1.05-1.11)[d]	0.54	3.23
	Predicted visceral adipose tissue (31501611)	rs362307	UTR	T: 0.063	2.00×10^{-09}	0.03 (0.02 - 0.04)	0.54	3.23
HTT: haplotype five (rs6828882)	Monobrow (27182965)	rs6828882	intronic	G: 0.101	1.00×10^{-22}	NA	0.53	8.54

| | | rs2798297 | intronic | A: 0.353 | 1.00×10^{-24} | NA | 0.52 | 2.08 |

HTT: haplotype six (rs2798297) — Metabolic biomarkers (33980691)

Haplotype, Group of variants in high LD with one another. Tag variant, Variant proxy for all variants in high LD at a haplotype. [a]LDlink SNPclip pruned signals using an r2 of 0.5 in the 1000 Genomes Project European superpopulations. [b]Ensembl VEP 'most severe' annotations. [c]1000 Genomes Project European superpopulation effect allele frequency. [d]Odds ratio (all other effects are beta estimates). **Abbreviations:** AF, Allele frequency; CADD, Combined Annotation Dependent Depletion; EUR, 1000 Genomes European super-population; L2G, Open Targets locus-to-gene scoring metric; NA, data Not available; PMID, PubMed Identifier; UTR, Untranslated region.

Table 2. Non-canonical disease associations of polyQ disorder genes potentially informs off target effects of therapeutic knockdown of pathogenic disease genes. Six polyQ disorder genes have OMIM phenotype entries outside of their main polyQ disorder. These phenotypes range from endocrine and metabolic conditions (e.g., androgen insensitivity), to neurological and neurodevelopmental disorders (e.g., developmental and epileptic encephalopathy, Lopes-Maciel-Rodan syndrome), and susceptibility to complex diseases.

Gene	OMIM phenotypes
AR	- Androgen insensitivity, partial, with or without breast cancer (MIM: 312300, X-linked recessive) - Androgen insensitivity (MIM: 300068, X-linked recessive) - Hypospadias 1, X-linked (MIM:300633, X-linked recessive) *Susceptibility to: Prostate cancer (MIM: 176807, autosomal dominant, somatic mutation)*
ATN1	Congenital hypotonia, epilepsy, developmental delay, and digital anomalies (MIM: 618494, autosomal dominant)
ATXN2	*Susceptibility to: Amyotrophic lateral sclerosis (MIM: 183090, autosomal dominant)*[*] *Susceptibility to: Parkinson disease, late-onset (MIM: 168600, autosomal dominant, multifactorial)*[*]
CACNA1A	- Developmental and epileptic encephalopathy 42 (MIM: 617106, autosomal dominant) - Episodic ataxia, type 2 (MIM: 108500, autosomal dominant)[*] - Migraine, familial hemiplegic, 1, with progressive cerebellar ataxia (MIM: 141500, autosomal dominant)
HTT	Lopes-Maciel-Rodan syndrome (MIM: 617435, autosomal recessive)
TBP	*Susceptibility to: Parkinson disease (MIM: 168600, autosomal dominant, multifactorial)*[*]

[*]CAG repeat mediated disorder/susceptibility. **Abbreviations: OMIM**, Online Mendelian Inheritance in Man

REFERENCES

1. Bunting, E.L., Hamilton, J., and Tabrizi, S.J. (2022). Polyglutamine diseases. Curr Opin Neurobiol *72*, 39–47. https://doi.org/10.1016/j.conb.2021.07.001.

2. Fan, H.-C., Ho, L.-I., Chi, C.-S., Chen, S.-J., Peng, G.-S., Chan, T.-M., Lin, S.-Z., and Harn, H.-J. (2014). Polyglutamine (PolyQ) diseases: genetics to treatments. Cell Transplant *23*, 441–458. https://doi.org/10.3727/096368914X678454.

3. Bonsor, M., Ammar, O., Schnoegl, S., Wanker, E.E., and Silva Ramos, E. (2024). Polyglutamine disease proteins: Commonalities and differences in interaction profiles and pathological effects. Proteomics *24*, e2300114. https://doi.org/10.1002/pmic.202300114.

4. Ghanekar, S.D., Kuo, S.-H., Staffetti, J.S., and Zesiewicz, T.A. (2022). Current and emerging treatment modalities for spinocerebellar ataxias. Expert Rev Neurother *22*, 101–114. https://doi.org/10.1080/14737175.2022.2029703.

5. Stoker, T.B., Mason, S.L., Greenland, J.C., Holden, S.T., Santini, H., and Barker, R.A. (2022). Huntington's disease: diagnosis and management. Pract Neurol *22*, 32–41. https://doi.org/10.1136/practneurol-2021-003074.

6. Perlman, S.L. (2020). Update on the Treatment of Ataxia: Medication and Emerging Therapies. Neurotherapeutics *17*, 1660–1664. https://doi.org/10.1007/s13311-020-00941-3.

7. Rook, M.E., and Southwell, A.L. (2022). Antisense Oligonucleotide Therapy: From Design to the Huntington Disease Clinic. BioDrugs *36*, 105–119. https://doi.org/10.1007/s40259-022-00519-9.

8. Esteves, S., Duarte-Silva, S., and Maciel, P. (2017). Discovery of Therapeutic Approaches for Polyglutamine Diseases: A Summary of Recent Efforts. Med Res Rev *37*, 860–906. https://doi.org/10.1002/med.21425.

9. Kwon, D. (2021). Failure of genetic therapies for Huntington's devastates community. Nature *593*, 180. https://doi.org/10.1038/d41586-021-01177-7.

10. McColgan, P., Thobhani, A., Boak, L., Schobel, S.A., Nicotra, A., Palermo, G., Trundell, D., Zhou, J., Schlegel, V., Sanwald Ducray, P., et al. (2023). Tominersen in Adults with Manifest Huntington's Disease. N Engl J Med *389*, 2203–2205. https://doi.org/10.1056/NEJMc2300400.

11. Plenge, R.M. (2016). Disciplined approach to drug discovery and early development. Sci Transl Med *8*, 349ps15. https://doi.org/10.1126/scitranslmed.aaf2608.

12. Plenge, R.M., Scolnick, E.M., and Altshuler, D. (2013). Validating therapeutic targets through human genetics. Nat Rev Drug Discov *12*, 581–594. https://doi.org/10.1038/nrd4051.

13. Ghoussaini, M., Mountjoy, E., Carmona, M., Peat, G., Schmidt, E.M., Hercules, A., Fumis,
L., Miranda, A., Carvalho-Silva, D., Buniello, A., et al. (2021). Open Targets Genetics:
systematic identification of trait-associated genes using large-scale genetics and functional
genomics. Nucleic Acids Res *49*, D1311–D1320. https://doi.org/10.1093/nar/gkaa840.

14. Minikel, E.V., Painter, J.L., Dong, C.C., and Nelson, M.R. (2024). Refining the impact of
genetic evidence on clinical success. Nature *629*, 624–629. https://doi.org/10.1038/s41586-
024-07316-0.

15. Schultz, J.L., Saft, C., and Nopoulos, P.C. (2021). Association of CAG Repeat Length in the
Huntington Gene With Cognitive Performance in Young Adults. Neurology *96*, e2407–
e2413. https://doi.org/10.1212/WNL.0000000000011823.

16. Slike, A.N., and Wright, G.E.B. (2022). Common huntingtin-related genetic variation is
associated with neurobiological and aging traits in humans. Cell Death Discov *8*, 311.
https://doi.org/10.1038/s41420-022-01114-1.

17. Ghoussaini, M., Nelson, M.R., and Dunham, I. (2023). Future prospects for human genetics
and genomics in drug discovery. Curr Opin Struct Biol *80*, 102568.
https://doi.org/10.1016/j.sbi.2023.102568.

18. Rusina, P.V., Falaguera, M.J., Romero, J.M.R., McDonagh, E.M., Dunham, I., and Ochoa,
D. (2023). Genetic support for FDA-approved drugs over the past decade. Nat Rev Drug
Discov *22*, 864. https://doi.org/10.1038/d41573-023-00158-x.

19. Zhang, X., Yu, W., Li, Y., Wang, A., Cao, H., and Fu, Y. (2024). Drug development
advances in human genetics-based targets. MedComm (2020) *5*, e481.
https://doi.org/10.1002/mco2.481.

20. Razuvayevskaya, O., Lopez, I., Dunham, I., and Ochoa, D. (2024). Genetic factors
associated with reasons for clinical trial stoppage. Nat Genet *56*, 1862–1867.
https://doi.org/10.1038/s41588-024-01854-z.

21. Nelson, M.R., Tipney, H., Painter, J.L., Shen, J., Nicoletti, P., Shen, Y., Floratos, A., Sham,
P.C., Li, M.J., Wang, J., et al. (2015). The support of human genetic evidence for approved
drug indications. Nat Genet *47*, 856–860. https://doi.org/10.1038/ng.3314.

22. Martin, F.J., Amode, M.R., Aneja, A., Austine-Orimoloye, O., Azov, A.G., Barnes, I.,
Becker, A., Bennett, R., Berry, A., Bhai, J., et al. (2023). Ensembl 2023. Nucleic Acids Res
51, D933–D941. https://doi.org/10.1093/nar/gkac958.

23. Mountjoy, E., Schmidt, E.M., Carmona, M., Schwartzentruber, J., Peat, G., Miranda, A.,
Fumis, L., Hayhurst, J., Buniello, A., Karim, M.A., et al. (2021). An open approach to
systematically prioritize causal variants and genes at all published human GWAS trait-
associated loci. Nat Genet *53*, 1527–1533. https://doi.org/10.1038/s41588-021-00945-5.

24. Amberger, J.S., Bocchini, C.A., Schiettecatte, F., Scott, A.F., and Hamosh, A. (2015).
OMIM.org: Online Mendelian Inheritance in Man (OMIM®), an online catalog of human

genes and genetic disorders. Nucleic Acids Res *43*, D789-798.
https://doi.org/10.1093/nar/gku1205.

25. de Sainte Agathe, J.-M., Filser, M., Isidor, B., Besnard, T., Gueguen, P., Perrin, A., Van
Goethem, C., Verebi, C., Masingue, M., Rendu, J., et al. (2023). SpliceAI-visual: a free
online tool to improve SpliceAI splicing variant interpretation. Hum Genomics *17*, 7.
https://doi.org/10.1186/s40246-023-00451-1.

26. Machiela, M.J., and Chanock, S.J. (2015). LDlink: a web-based application for exploring
population-specific haplotype structure and linking correlated alleles of possible functional
variants. Bioinformatics *31*, 3555–3557. https://doi.org/10.1093/bioinformatics/btv402.

27. Huyghe, J.R., Fransen, E., Hannula, S., Van Laer, L., Van Eyken, E., Mäki-Torkko, E.,
Aikio, P., Sorri, M., Huentelman, M.J., and Camp, G.V. (2011). A genome-wide analysis of
population structure in the Finnish Saami with implications for genetic association studies.
Eur J Hum Genet *19*, 347–352. https://doi.org/10.1038/ejhg.2010.179.

28. Karczewski, K.J., Francioli, L.C., Tiao, G., Cummings, B.B., Alföldi, J., Wang, Q., Collins,
R.L., Laricchia, K.M., Ganna, A., Birnbaum, D.P., et al. (2020). The mutational constraint
spectrum quantified from variation in 141,456 humans. Nature *581*, 434–443.
https://doi.org/10.1038/s41586-020-2308-7.

29. Lek, M., Karczewski, K.J., Minikel, E.V., Samocha, K.E., Banks, E., Fennell, T., O'Donnell-
Luria, A.H., Ware, J.S., Hill, A.J., Cummings, B.B., et al. (2016). Analysis of protein-coding
genetic variation in 60,706 humans. Nature *536*, 285–291.
https://doi.org/10.1038/nature19057.

30. Uhlén, M., Fagerberg, L., Hallström, B.M., Lindskog, C., Oksvold, P., Mardinoglu, A.,
Sivertsson, Å., Kampf, C., Sjöstedt, E., Asplund, A., et al. (2015). Proteomics. Tissue-based
map of the human proteome. Science *347*, 1260419.
https://doi.org/10.1126/science.1260419.

31. Digre, A., and Lindskog, C. (2021). The Human Protein Atlas—Spatial localization of the
human proteome in health and disease. Protein Science *30*, 218–233.
https://doi.org/10.1002/pro.3987.

32. Ochoa, D., Hercules, A., Carmona, M., Suveges, D., Baker, J., Malangone, C., Lopez, I.,
Miranda, A., Cruz-Castillo, C., Fumis, L., et al. (2023). The next-generation Open Targets
Platform: reimagined, redesigned, rebuilt. Nucleic Acids Res *51*, D1353–D1359.
https://doi.org/10.1093/nar/gkac1046.

33. Villaveces, J.M., Jiménez, R.C., Porras, P., Del-Toro, N., Duesbury, M., Dumousseau, M.,
Orchard, S., Choi, H., Ping, P., Zong, N.C., et al. (2015). Merging and scoring molecular
interactions utilising existing community standards: tools, use-cases and a case study.
Database (Oxford) *2015*, bau131. https://doi.org/10.1093/database/bau131.

34. Freshour, S.L., Kiwala, S., Cotto, K.C., Coffman, A.C., McMichael, J.F., Song, J.J., Griffith,
M., Griffith, O.L., and Wagner, A.H. (2021). Integration of the Drug-Gene Interaction

Database (DGIdb 4.0) with open crowdsource efforts. Nucleic Acids Res *49*, D1144–D1151.
https://doi.org/10.1093/nar/gkaa1084.

35. Namuli, K.L., Slike, A.N., Hollebeke, M.A., and Wright, G.E.B. (2025). Genomic
characterization of Huntington's disease genetic modifiers informs drug target tractability.
Brain Communications *7*, fcae418. https://doi.org/10.1093/braincomms/fcae418.

36. Coarelli, G., Coutelier, M., and Durr, A. (2023). Autosomal dominant cerebellar ataxias: new
genes and progress towards treatments. Lancet Neurol *22*, 735–749.
https://doi.org/10.1016/S1474-4422(23)00068-6.

37. Nguyen, P.A., Born, D.A., Deaton, A.M., Nioi, P., and Ward, L.D. (2019). Phenotypes
associated with genes encoding drug targets are predictive of clinical trial side effects. Nat
Commun *10*, 1579. https://doi.org/10.1038/s41467-019-09407-3.

38. Tandon, S., Aggarwal, P., and Sarkar, S. (2024). Polyglutamine disorders: Pathogenesis and
potential drug interventions. Life Sciences *344*, 122562.
https://doi.org/10.1016/j.lfs.2024.122562.

39. Rajan-Babu, I.-S., Dolzhenko, E., Eberle, M.A., and Friedman, J.M. (2024). Sequence
composition changes in short tandem repeats: heterogeneity, detection, mechanisms and
clinical implications. Nat Rev Genet *25*, 476–499. https://doi.org/10.1038/s41576-024-
00696-z.

40. Gui, H., Kwan, J.S., Sham, P.C., Cherny, S.S., and Li, M. (2017). Sharing of Genes and
Pathways Across Complex Phenotypes: A Multilevel Genome-Wide Analysis. Genetics *206*,
1601–1609. https://doi.org/10.1534/genetics.116.198150.

41. Peng, J., Zhou, Y., and Wang, K. (2021). Multiplex gene and phenotype network to
characterize shared genetic pathways of epilepsy and autism. Sci Rep *11*, 952.
https://doi.org/10.1038/s41598-020-78654-y.

42. Ward, J.M., Stoyas, C.A., Switonski, P.M., Ichou, F., Fan, W., Collins, B., Wall, C.E.,
Adanyeguh, I., Niu, C., Sopher, B.L., et al. (2019). Metabolic and Organelle Morphology
Defects in Mice and Human Patients Define Spinocerebellar Ataxia Type 7 as a
Mitochondrial Disease. Cell Rep *26*, 1189-1202.e6.
https://doi.org/10.1016/j.celrep.2019.01.028.

43. Klockgether, T., Mariotti, C., and Paulson, H.L. (2019). Spinocerebellar ataxia. Nat Rev Dis
Primers *5*, 24. https://doi.org/10.1038/s41572-019-0074-3.

44. Flower, M., Lomeikaite, V., Ciosi, M., Cumming, S., Morales, F., Lo, K., Hensman Moss,
D., Jones, L., Holmans, P., TRACK-HD Investigators, et al. (2019). MSH3 modifies somatic
instability and disease severity in Huntington's and myotonic dystrophy type 1. Brain *142*,
1876–1886. https://doi.org/10.1093/brain/awz115.

45. Laabs, B.-H., Klein, C., Pozojevic, J., Domingo, A., Brüggemann, N., Grütz, K., Rosales,
R.L., Jamora, R.D., Saranza, G., Diesta, C.C.E., et al. (2021). Identifying genetic modifiers

of age-associated penetrance in X-linked dystonia-parkinsonism. Nat Commun *12*, 3216. https://doi.org/10.1038/s41467-021-23491-4.

46. Bettencourt, C., Hensman-Moss, D., Flower, M., Wiethoff, S., Brice, A., Goizet, C., Stevanin, G., Koutsis, G., Karadima, G., Panas, M., et al. (2016). DNA repair pathways underlie a common genetic mechanism modulating onset in polyglutamine diseases. Ann Neurol *79*, 983–990. https://doi.org/10.1002/ana.24656.

47. Wright, G.E.B., Findlay Black, H., Collins, J.A., Gall-Duncan, T., Caron, N.S., Pearson, C.E., and Hayden, M.R. (2020). Interrupting sequence variants and age of onset in Huntington's disease: clinical implications and emerging therapies. Lancet Neurol *19*, 930–939. https://doi.org/10.1016/S1474-4422(20)30343-4.

48. Inc, uniQure (2024). uniQure Announces Positive Interim Data Update Demonstrating Slowing of Disease Progression in Phase I/II Trials of AMT-130 for Huntington's Disease. GlobeNewswire News Room. https://www.globenewswire.com/en/news-release/2024/07/09/2910220/0/en/uniQure-Announces-Positive-Interim-Data-Update-Demonstrating-Slowing-of-Disease-Progression-in-Phase-I-II-Trials-of-AMT-130-for-Huntington-s-Disease.html.

49. Handsaker, R.E., Kashin, S., Reed, N.M., Tan, S., Lee, W.-S., McDonald, T.M., Morris, K., Kamitaki, N., Mullally, C.D., Morakabati, N.R., et al. (2025). Long somatic DNA-repeat expansion drives neurodegeneration in Huntington's disease. Cell *0*. https://doi.org/10.1016/j.cell.2024.11.038.