The Prevalence, Prevention, and Treatment of Cardiovascular Diseases in Twelve African

Countries (2014-2019): An Analysis of the World Health Organisation STEPwise Approach to

Chronic Disease Risk Factor Surveillance

Authors and affiliations

Wingston Felix Ng'ambi[1], Janne Estill[1], Fatma Aziza Merzouki[1], Cosamas Zyambo[2],
Jonathan Chiwanda Banda[3], David Beran[4], Olivia Keiser[1]

[1]Institute of Global Health, University of Geneva, Geneva, Switzerland

[2]University of Zambia, Department of Community Health and Family Medicine, Lusaka, Zambia

[3]Division of Non-Communicable Diseases and Mental Health, Ministry of Health,
Lilongwe, Malawi

[4]Division of Tropical and Humanitarian Medicine, University of Geneva, Geneva, Switzerland

ABSTRACT

Introduction: Cardiovascular diseases (CVDs) are responsible for nearly a third of
deaths globally. We conducted this study to understand the prevalence of history of
CVDs, their prevention and treatment in twelve African countries using the World Health
Organization STEPwise Approach to Surveillance (WHO STEPS) data.

Methods: We used secondary STEPS data extracted from 12 African countries
between 2014 and 2019. CVD was defined as a self-reported history of heart attack,
angina, or stroke. Weighted percentages, counts, weighted odds ratios (OR), and the
corresponding 95% confidence intervals (95%CI) were computed using the R software.
We fitted logistic regression models to select the predictor variables from a regression
model for CVD prevalence, CVD prevention and CVD treatment binary endpoints.

Results: Amongst 60,294 individuals, the prevalence of CVD was 5%. The CVD
prevalence was higher in older individuals, females, individuals with hypertension,
smokers, people with high salt intake, and in certain countries. Eleven percent of the
23,630 individuals at high risk of CVD (≥40 years) but without a history of the disease
received CVD prevention treatment. Amongst the 2,895 persons with CVDs, 22%

received treatment and counselling for CVD: 34% (n=215) receiving aspirin, 32% (n=202) counselling for CVD risk factors, 11% (n=66) statins, and 24% (n=148) both statins and aspirin. The uptake of CVD treatment varied by hypertension status, sex, age and country.

Conclusion: The prevalence of CVD was relatively low and CVD treatment uptake was sub-optimal. Concerted efforts must be made to accelerate the diagnosis and expand treatment for CVDs in Africa if to curtail untimely deaths attributable to CVDs.

Running tittle: CVD prevalence, prevention and treatment in Africa

Key words: CVD, Cardiovascular diseases, Africa, NCD, WHO STEPS, Non-communicable diseases

INTRODUCTION

Despite the relatively low prevalence of cardiovascular diseases (CVDs) in Africa compared to developed countries, there is evidence of a steady increase, which poses a significant public health concern. One of the aims of the United Nations Sustainable Development Goals is to reduce premature mortality from non-communicable diseases (NCDs) by a third by the year 2030 [1]. The CVDs, like coronary heart disease and stroke, are the most common NCDs globally responsible for an estimated 9.6 million male deaths and 8.9 million female deaths worldwide in 2019, of which 6.1 million of were among people aged 30 to 70 years [2]. Of these deaths related to CVDs, more than 75% occurred in low- and middle-income countries [3]. Heart attacks and strokes accounted for 85% of these fatalities [4]. Globally between 1990 and 2020 the age standardized DALYs for CVDs equal those of the communicable, maternal, neonatal and nutritional (CMNN) diseases combined.

While infectious diseases have historically been the primary focus of public health efforts in the region, CVDs are becoming a significant and increasing problem [5]. The

prevalence of CVDs in Africa has been steadily increasing over the past few decades
[5]. Despite the growing prevalence of CVDs in Africa, there is a lack of multi-country
analyses assessing the full care continuum; from diagnosis to treatment; in this region.
According to the Global Burden of Disease (GBD) project, prevalence of CVDs and
mortality differ substantially between high-income and low-income regions. Particularly,
Africa is seeing rising rates of diabetes, obesity, and hypertension; all of which are
significant causes of CVD; but these patterns are not well-represented in international
research initiatives. The region is understudied in comparison to more resourced
locations because of issues with infrastructure, data availability, and healthcare systems
[6]. Furthermore, while global trends in prevalence of CVDs and their associated risk
factors have been extensively studied; Africa remains relatively understudied despite
the rising burden of CVDs [6]. A dearth of comprehensive and localised studies that are
suited to the region's particular socioeconomic and healthcare contexts is further
highlighted by recent reports from GBD collaborations that emphasise the need to shed
light on the CVD epidemiology in Africa. Through region-specific analysis, these
initiatives are starting to address modifiable risk factors; nevertheless, there are still
gaps in assessing the effectiveness and implementation of interventions [7]. Efforts to
address this growing health concern, like improved healthcare access, awareness
campaigns, and research to develop region-specific prevention and treatment
strategies, are critical [8] [9]. For example, the World Health Organization developed the
package of essential noncommunicable disease interventions (PEN) and related
strategies (PEN-PLUS) for resource limited settings covering also the care of severe
NCDs in these settings [2][10].

The prevalence of CVDs in Africa is influenced by a complex interplay of various
determinants such as hypertension, diet, physical activity, tobacco use, alcohol
consumption, age, gender, genetic factors, socio-economic deprivation (including
poverty and limited access to healthcare), environmental factors (pollution and limited
access to clean and safe water, and exposure to toxins), lack of awareness about CVD
risk factors and prevention, and limited availability and quality of healthcare services
and facilities in the region. These factors can significantly influence the diagnosis,
treatment, and prevention of CVD [11] [12]. In addition, the prevalence of CVDs in Africa

is influenced by multiple determinants that vary across the regions, making it important

to consider both spatial and temporal aspects when addressing the burden of CVDs in

the region. The study acknowledges the complex interplay of determinants influencing

the prevalence of CVDs in Africa, including socio-economic, environmental, and health

system factors. This approach highlights the importance of considering regional

variability, which is often overlooked in global analyses. By concentrating on Africa, the

study fills a crucial gap in the literature, particularly in the context of limited healthcare

resources and overlapping burdens of infectious diseases like HIV/AIDS and

tuberculosis. Understanding the CVD determinants in Africa is crucial for developing

effective strategies to reduce the prevalence of CVDs in Africa. For example, cessation

of tobacco use, reduction of salt in the diet, eating more fruit and vegetables, regular

physical activity, and avoiding harmful use of alcohol have been shown to be protective

of CVD [13] [14]. Furthermore, identifying the persons at highest risk of CVDs and

ensuring they receive appropriate treatment can prevent premature deaths [15]. While

the WHO STEPwise approach has provided a framework for monitoring NCDs, its data

remain underutilized, especially in Africa [16]. This study addresses a critical gap in

global CVD research by providing the first multi-country analysis of CVD prevalence,

prevention, and treatment in Africa using WHO STEPS data. Its findings are not only

relevant for Africa but also provide a template for utilizing similar data in other regions,

thus contributing to the global effort to reduce NCD-related premature mortality.

METHODS

Study design and setting

This is a secondary analysis of WHO STEPS data from countries with CVD data

collected between 2014 and 2019 in Africa (see Box 1). This analysis includes data

from 12 countries in Africa: Algeria, Benin, Botswana, Eswatini, Ethiopia, Kenya,

Malawi, Morocco, São Tomé and Príncipe, Sudan, Uganda, and Zambia. The WHO

STEPS assess risk factors for chronic non-communicable diseases and uses a multi-

stage cluster sampling of households. One individual within the age range of the survey

(18-69 years) was selected per household [17]. For example, Malawi's STEPS survey in

2017 and Zambia's STEPS survey in 2017 followed this methodology to capture critical

health data for analysis [18] [19]. The WHO STEPS use simple, standardized methods

for collecting, analysing and disseminating data on key NCD risk factors in the

countries. The survey covers key behavioural risk factors: tobacco use, alcohol use,

physical inactivity, and unhealthy diet; as well as key biological risk factors: overweight

and obesity, raised blood pressure, raised blood glucose, and abnormal blood lipids

[17]. The survey is conducted using a stepwise procedure that begins with a

questionnaire to collect data on risk factors (STEPS 1), progresses to basic physical

examinations (STEPS 2), and concludes with the more intricate collection of blood

samples for biochemical analysis (STEPS 3) [17]. The 2014-2019 period was selected

because data on CVD were unavailable before this timeframe.

WHO Package of Essential Noncommunicable intervention implementation

Across Africa, countries exhibit considerable heterogeneity in the timing and extent of
adopting national NCD strategies and implementing the WHO Package of Essential
Noncommunicable (PEN) disease interventions. The WHO PEN approach was first
introduced globally in 2010 and WHO/AFRO has supported member states in PEN
roll-out since at least 2008[20]. In North Africa, Algeria formalized its national NCD
strategic planning most recently (with an updated UNSDCF including NCDs from 2023)
and Morocco has integrated PEN elements into its primary care cancer prevention
efforts, with routine risk factor surveillance reported since 2018. In West Africa, Benin's
earlier NCD strategic plan (2014–2018) provides the backbone for current pilot PEN
activities, and Sao Tome and Principe incorporated PEN into its NCD action planning
upon joining WHO/AFRO's regional frameworks in 2023. In East Africa, Ethiopia's NCD
strategic planning dates back to at least 2014–2016, followed by updated strategies into
the early 2020s; PEN activities have been scaled in selected regions with quarterly
monitoring. Kenya's national NCD strategy (2015–2020) underpins systematic PEN
roll-out across counties, and Uganda's policy (published 2020) supports PEN training
for frontline staff. Southern African countries show similar evolutionary progress:
Botswana reports PEN guideline integration into primary care with service
improvements; Eswatini's strategic NCD plan dates from the 2012–2020 era; and
Zambia has embedded PEN within its NCD framework, recently launching a PEN-Plus
National Operational Plan in 2025 to expand care and training beyond primary care.
Sudan adopted an NCD strategy as early as 2010–2015, though political instability has
constrained consistent PEN implementation. In Malawi, PEN integration into the
National Health Strategic Plan has guided initial district hospital training since the early
2020s, with ongoing expansion to peripheral centres.

Variables and data management

The most recent WHO STEPS data on CVD risk factors were extracted from each country. The data were managed in R (see 1_**Create_dataset_for_analysis.R; Functions_rmph.R**).

Outcome variables: The primary outcome variable was whether an individual responded "yes" or "no" to the question "Have you ever had a heart attack, angina (chest discomfort caused by heart disease), or stroke (cerebrovascular accident or incident)?", which we considered as a proxy for having CVD. We excluded 141,833 patients with missing CVD information. Prevention of CVD or uptake of treatment for CVD was assessed with the questions "Are you currently taking aspirin regularly to prevent or treat heart disease?" (yes/no), "Are you currently taking statins (Lovostatin/Simvastatin/Atorvastatin or any other statin) regularly to prevent or treat heart disease?" (yes/no) or "During the past three years, has a doctor or other health worker advised you to do any of the following: quit using tobacco or not start, reduce alcohol consumption or not start, reduce salt in your diet, reduce refined sugar in your diet, or eat at least five servings of fruits and/or vegetables each day?" (yes/no).

Predictor variables: The predictor variables comprised the following known dichotomous risk and protective factors defined as follows: harmful alcohol use (defined as daily drinking and having at least 4 (for males) or 3 (for females) drinking occasions in the last 30 days) derived using the Alcohol Use Disorder Identification Test-C (AUDIT-C) [21]; low consumption of vegetables (i.e. "In a typical week, on how many days do you eat vegetables?" with less than five times a week being categorized as low consumption of vegetables); low consumption of fruits (i.e. "In a typical week, on how many days do you eat fruit?" with less than five times a week being categorized to have low fruit consumption); physical inactivity (less than 150 minutes of moderate-intensity physical activity per week) [22]; high salt intake (i.e. "How often do you add salt or a salty sauce such as soy sauce to your food right before you eat it or as you are eating it?" with those adding salt to food regularly or often eating processed food with high salt quantities are considered to have high salt intake); history of raised blood pressure (systolic blood pressure ≥160 and/or diastolic blood pressure ≥ 100 mmHg or currently on medication); and history of raised blood glucose (capillary whole blood value at least

6.1mmol/L) [23]. We also included smoking history (currently smoking, previously

smoking and never smoked); body mass index (<18.5, 18.5-24.0, 25.0-29.0, 30.0+

kg/m^2; excluding pregnant women), sex of the respondent (male, female), age (15-29,

30-39, 40-49, 50-59, 60+; in completed calendar years), highest education level (none,

primary, secondary, tertiary), occupation (government employee, non-government

employee, self-employed, retired, or not working (including e.g. students, unemployed)),

type of residence (rural, urban), country, and marital status (never married, married,

previously married).

The distribution of missingness in the analysed data is shown in Box 2. Overall, 26% of

the individuals had missing data. The variable with the greatest missing data was

diabetes with 8% missing data. Imputation was performed to handle the missing data

because it allows for the inclusion of all available data in the analysis, ensuring more

accurate and representative results. Different probability distributions were specified

depending on the type of variable. For binary variables, such as presence or absence of

a condition, we used a binomial distribution, which models the probability of a "success"

or "failure" outcome. For categorical variables with more than two groups, such as

education level or occupation, we applied a multinomial distribution that accounts for

multiple outcome categories and ensures that the probabilities across categories sum to

one. For continuous variables, including age, body mass index, and blood pressure, a

Gaussian distribution was used to generate values based on the mean and variance of

the observed data. These simulation models were implemented using random

assignment based on the observed distribution of each variable, ensuring that imputed

values reflected the empirical patterns in the data. This strategy allowed us to

approximate the underlying data-generating mechanism more realistically, reduce bias

from listwise deletion, and retain the full analytic sample.

We opted for simulation-based imputation instead of standard MICE approaches for

several reasons. First, the data exhibited complex patterns across multiple countries,

including high inter-variable correlations and heterogeneity in variable distributions,

which led to convergence issues and unstable chains when using MICE. Second, some

variables had non-standard distributions or rare categories that are not easily handled

by default MICE models. Third, simulation-based imputation allows direct specification

of the appropriate distribution for each variable type (binomial, multinomial, Gaussian),

ensuring that imputed values reflect the empirical distribution of the observed data.

Missing categorical data were imputed using the base R function sample(), drawing

from the observed distribution of non-missing values, while continuous variables such

as age were simulated using rnorm() based on the observed mean and standard

deviation, with results rounded to the nearest integer. Replacement values were

therefore drawn from overall empirical distributions, and we acknowledge that this

approach does not capture country-specific heterogeneity in risk factor distributions. To

support reproducibility and ensure plausibility, we set a random seed prior to simulation

and compared the distribution of imputed values with complete-case data (see **sample**

simulation code below or 2_who_steps_ncd_data_cleaning_code_2024.R).

Finally, this approach avoids the iterative dependency structure of MICE, which can

amplify biases in the presence of autocorrelation, while preserving sample size and

statistical power. Furthermore, imputation is a common practice in epidemiological

research to maintain data integrity and reduce bias due to incomplete datasets [24]. In

order to ascertain the uptake of CVD prevention treatment, all the individuals aged at

least 40 years were considered to be at risk of CVD and this formed the denominator for

this analysis [25], [26], [27], [28], [29]. Furthermore, the WHO STEPS data analysis

guide also considers the CVD naïve individuals aged at least 40 years as being at CVD

risk [30], [31].

Data analysis

We set up the survey design before fitting any models and used the same weight

variable, wstep1, throughout the analysis (see

3_**CVD_Analysis_SSA_2024_FINAL.Rmd**). For the CVD prevalence and prevention

datasets, we defined a full complex design using primary sampling unit (psu) as the

cluster, stratum as the stratification variable, and wstep1 as the sampling weight, with

nesting enabled. For the treatment dataset, only wstep1 was available, so we applied a

simple one stage design. All quasibinomial models for CVD prevalence were run on

these survey design objects, ensuring that the sampling structure was properly

accounted for. We also adjusted for inter country variation by including country fixed

effects, rather than treating countries as random clusters. This allowed us to capture

differences across countries in a clear and consistent way.

Our analysis is structured around the CVD care pathway; spanning diagnosis, treatment
initiation, and counselling/adherence support; which provides a systematic framework to
identify gaps in care and is particularly relevant in the African context, where resource
constraints and variations in health system capacity can lead to substantial drop-offs at
each stage of the cascade (see Box 3). Weighted percentages, counts, weighted odds
ratios (OR), and the corresponding 95% confidence intervals (95%CI) were computed
using the R software (version 4.3.2). The weighting variable from WHO STEPS was
based on demographic characteristics of the general population from which a sample
was taken (wstep1) [32]. We conducted a set of three weighted univariable and
multivariable logistic regression analyses, with quasibinomial distributions [33], of the
effects of the predictor variables on: CVD prevalence in the whole dataset, CVD
prevention amongst the persons at risk of CVD, and CVD treatment (and counselling)
among those with CVD (see 3_**CVD_Analysis_SSA_2024_FINAL.Rmd**).

The quasibinomial distribution, an extension of the standard binomial distribution, is
particularly suitable for survey data with binary outcomes, as it accounts for
overdispersion (see 3_**CVD_Analysis_SSA_2024_FINAL.Rmd**). For each survey-
weighted multivariable quasibinomial logistic regression model, survey weights were
incorporated directly into the model fitting to ensure that parameter estimates reflected
the complex sampling design. Weighted standard errors were calculated to account for
unequal selection probabilities, thereby producing valid confidence intervals and p-
values. We first fitted a full model including all candidate predictors and then applied
backward variable selection using the stepAIC function from the MASS R package,
retaining predictors that optimized model fit while respecting the survey weights [34].
The stepAIC function was used for stepwise selection, optimizing model fit by retaining
predictors that minimized the AIC. Notably, the clinically important variables; age, sex,
and hypertension status; were naturally retained in the final model based on the AIC
criterion, so no variables needed to be forced into the model. We calculated the P-
values for the bivariate logistic regression by comparing the model with the variable of
interest and the empty model. Weighted quasibinomial logistic regression models with
country fixed effects were used to account for country-level heterogeneity, as mixed-
effects models were not preferred due to heterogeneity in survey design and instability

with weighted data. Similarly, we calculated the P-values from adjusted models by comparing the final model with the model without the variable of interest. Wherever applicable the level of statistical significance was set at $P<0.05$.

Sensitivity analyses (see 3_**4_CVD sensitivity analysis 2025.R**) were conducted to assess the robustness of the imputation assumptions by comparing results from the simulation-based imputation to a complete case analysis as shown in Box 4. This approach evaluated whether the main findings were influenced by the distributional assumptions used for imputing missing values. Based on the results there was perfect agreement of the coefficients from the imputed and complete case analyses coefficients (see Box 4).

Ethical consideration

Individual written consent was obtained from all participants before data collection the during the original WHO STEPS surveys. We requested access for secondary use of the data from the World Health Organization, the funder of the WHO STEPS surveys. The WHO STEPS survey datasets were downloaded from https://extranet.who.int/ncdsmicrodata/index.php/catalog/629 [23]. This dataset is anonymized with no identifiable information on survey respondents.

RESULTS

Individuals included and excluded from the study

The distribution of individuals included in the study is shown in Figure 1. Of the 202,127 individuals included in the database, 60,294 (29.8%) were interviewed from 2014 onwards and included information on CVD.

Characteristics of the study population

The characteristics of individuals included in this study are shown in Table 1. Of the 60,294 individuals, 44% were aged 15-29 years while 6% were aged 60 years or above.

The proportions of males and females were similar. More individuals were from rural

areas (60%) than urban areas (40%). About a third of the individuals had no education

and almost half of the individuals were self-employed (see Table 1). The majority of

individuals were currently married. The countries with highest numbers of respondents

were Ethiopia (23%), Algeria (21%), and Sudan (16%).

Of the 60,294 individuals, 9% had hypertension, 6% had high cholesterol and 3% had

diabetes. Furthermore, 19,333 (33%) reported high salt uptake, 45,207 (81%) low fruit

uptake, and 28,905 (54%) low vegetable uptake (Table 1). The prevalence of low level

of physical activity was reported as 16% while the prevalence of reported high level of

physical activity was 65%. A total of 1,803 (3%) of the 60,294 individuals reported using

alcohol in a harmful way. Furthermore, 5,047 (10%) and 3348 (7%) of the 60,294

individuals reported being current and previous smokers, respectively. The prevalence

of reported current smoking was 10% while 6% of the individuals had previously

smoked or used tobacco. The prevalence of measured underweight was 10% and the

prevalence of overweight (i.e. BMI of at least 30 kg/m^2) was 14%. The highest number

of individuals were interviewed in 2015 or 2016 and the least in 2019.

Care cascade for the cardiovascular disease

Across the 12 study countries, a substantial gap was observed along the CVD care

cascade in Figure 2. While the total eligible population ranged widely, only a small

proportion had a documented diagnosis of CVD, and an even smaller fraction received

treatment. For example, in Algeria, out of 6,955 individuals, only 415 (6.0%) had a CVD

diagnosis and 178 (2.6%) were on treatment. Similarly, in Malawi, 312 of 4,186

participants (7.5%) were diagnosed, but just 64 (1.5%) received treatment. Diagnosis

rates were particularly low in Kenya (0.4%) and Eswatini (4.3%), while Morocco showed

the highest diagnosis proportion (3.3%). Treatment coverage among those diagnosed

also varied widely, with Algeria achieving 42.9%, Morocco 36.2%, and Zambia only

15.0%. Overall, the cascade illustrates that fewer than one in ten individuals are

diagnosed, and less than half of those diagnosed receive treatment, underscoring major

gaps in CVD detection and management across the region.

Prevalence of CVD and factors associated with CVD in twelve countries of Africa

The prevalence of CVD was 5% (2895/60,294) and increased with age (Table 1). The

prevalence of CVD was higher in females than males, in urban than rural areas, and in

individuals with hypertension, using alcohol harmfully, or being obese or overweight.

Uganda (9%), Malawi (7%), Botswana (6%), Algeria (6%) and Kenya (6%) had the

highest, and Sudan (1%), Zambia (3%), and Morocco (3%) the lowest prevalence of

CVD (Table 1).

Older age, male sex, certain countries of residence, having hypertension, high salt

uptake, and current or previous smoking were significantly associated with higher risk of

CVD. The adjusted estimates show that there was an increasing trend in CVD with age

(Figure 3). After adjusting for sex, area of residence (urban vs rural), highest education

level, marital status, country of residence, having hypertension, high salt intake,

physical activity level, and smoking status, the individuals aged 50-59 or at least 60

years were more likely to have CVD (aOR: 1.39, 95%CI: 1.17-1.66 and aOR: 1.67,

95%CI: 1.36-2.07, respectively) than those aged 15-29 years. Males were less likely to

have CVD than females (aOR: 0.65, 95%CI: 0.54-0.72, P<0.001). Country-specific

differences highlight the role of regional factors, with Sudan showing the lowest risk and

Uganda the highest (Table 2). Individuals with hypertension were almost three times

more likely to have CVD compared to those without hypertension (aOR: 2.68, 95%CI:

2.32-3.10, P<0.001). Similarly, individuals with high salt uptake were 23% more likely to

have CVD than those without high salt uptake (aOR: 1.23, 95%CI: 1.02-1.49, P=0.03).

Those who never smoked tobacco were less likely to have CVD compared to those

currently smoking (aOR: 0.69, 95%CI: 0.55-0.86, P<0.0001).

Uptake of CVD prevention amongst the individuals aged at least 40 years in twelve countries of Africa

375 The characteristics of the 23,630 individuals who were at risk of CVD and interviewed

376 on the uptake of CVD prevention treatment and counselling are shown in Table 2. Of

377 these 23,630 individuals, the majority (51%) were aged between 40 and 49 years, 53%

were males, 56% were from rural areas, 37% had no education, 27% were from Algeria while less than 0.1% were from Morocco, 3% had diabetes, 9% had high cholesterol level, 16% had hypertension, 29% had high salt uptake, 81% had low fruit uptake, 49% had low vegetable uptake, 20% had low physical activity, 12% were current tobacco smokers, and 4% were harmful alcohol users.

Only 11% received any form of prevention or counselling, highlighting a major prevention gap. The distribution of the uptake of CVD prevention by the characteristics of the respondents is shown in Figure 4. People aged at least 60 years, those with BMI of at least 30, those who had previously smoked tobacco, those with low physical activity and those with tertiary education had the highest uptake of CVD prevention. Similarly, higher uptake of CVD prevention was observed amongst the females, urban residents, those with diabetes, those with hypertension, those with low salt uptake, and those who were not harmful alcohol users.

The factors associated with the uptake of CVD prevention are shown in Figure 4. Individuals who were aged at least 50 years were more likely to get CVD prevention compared to those who were aged between 40 and 49 years. Males were less likely to get CVD prevention compared to females (aOR: 0.73, 95%CI: 0.58-0.932, P<0.0001). Urban residents are more likely to receive prevention (aOR: 1.65, 95%CI: 1.33-2.04, P<0.0001) than their rural counterparts. Educational attainment influences uptake, with individuals having a tertiary education showing the highest uptake (Figure 4). Previous smokers had higher (aOR: 1.17, 95% CI: 0.83-1.64) and non-smokers lower uptake (aOR: 0.88, 95% CI: 0.64-1.20) compared with current smokers. Individuals with hypertension were 17 times more likely to get CVD prevention compared to those without hypertension (aOR: 16.65, 95%CI:13.93-19.91, P<0.0001). There were also country variations in uptake of treatment for preventing CVDs (see Figure 4). These findings suggest that targeted interventions addressing these modifiable factors could improve CVD prevention uptake in Sub-Saharan Africa.

Uptake of treatment for CVDs in twelve countries of Africa

The treatment and counselling received by the respondents with CVD is shown in Figure 1. A total of 630 (22%) of the 2895 individuals with CVD received treatment and

counselling. Of these 630 patients, 32% received counselling, 34% received aspirin (but no statins), 11% received statins (but no aspirin), and 24% received both statins and aspirin. The distribution of the uptake of treatment amongst the persons who had CVD is shown in Table 1. Individuals aged below 40 years had lower treatment uptake than individuals aged at least 50 years. The highest treatment uptake was observed amongst those with hypertension (53%), previous smoking history (30%), and obesity (27%) or overweight (40%). The CVD cases in Sudan showed the highest treatment uptake (42%) while the CVD cases in Kenya showed the lowest treatment uptake (5%).

The factors associated with CVD treatment uptake are shown in Figure 5. Age has a major impact on treatment uptake; those in the 50–59 and 60+ age groups were far more likely to get therapy than people in the 15–29 age range. For instance, the adjusted model showed a robust correlation between age increased treatment uptake, with an aOR of 2.31 (95%CI:1.39-3.84) for those aged 50-59 and 2.90 (95%CI: 1.75-4.82) for those aged 60+ compared to the 15–29-year-old. In the adjusted model, sex did not significantly affect treatment uptake; males had an OR of 1.36 (0.97-1.89) compared to females. However, urban dwellers were more likely than their rural counterparts to receive therapy (OR 1.36, 0.97-1.90). Individuals with hypertension were 7 times more likely to take treatment for CVD than those without hypertension (aOR: 7.19, 95%CI: 5.15-9.78, P<0.001), and individuals with low vegetable uptake were less likely to take CVD treatment than those who had high uptake of vegetables (aOR: 0.58, 95%CI: 0.54-0.83, P=0.003). Kenya, Malawi and Zambia had lower CVD therapy uptake compared to Algeria while Sudan had a similar CVD uptake compared to Algeria. With ORs of 1.37 (0.69-2.74), 0.74 (0.50-1.10) and 1.09 (0.70-1.72), respectively, diabetes, physical inactivity and current smoking demonstrated a weaker or non-significant correlation.

DISCUSSION

This study examined the prevalence of CVD and the uptake of both preventive and treatment services across twelve African countries, drawing on nationally representative WHO STEPS data. The discussion is organized around three key themes: CVD

prevalence, prevention, and treatment uptake. This is followed by an outline of the
study's strengths and limitations, and finally the conclusions and recommendations.

Discussion for CVD prevalence

The overall CVD prevalence of 5% in this study was consistent with findings from a
meta-analysis that found the CVD prevalence to be 7% in SSA [35]. In contrast, other
regions exhibit higher CVD prevalence, for example, Europe, South and Southeast Asia
and Central Asia have some of the highest CVD mortality rates globally [36], [37].
Similarly, North America has a high prevalence of CVD, with projections indicating that
61% of U.S. adults will have some form of cardiovascular disease by 2050 [37]. The low
rates of CVDs in Africa may be attributed to limited diagnostic capabilities for
confirmation [38]. In addition, the age distribution of the sample, including the exclusion
of adults over 69 years, may have contributed to lower estimates, as CVD risk increases
with age. Both factors should be considered when interpreting these findings relative to
regional prevalence estimates. As people get older, their heart and blood vessels
naturally work less efficiently, which greatly increases the risk of heart disease and
stroke [39] and this trend was also observed in this study. The implication is to provide
the screening services to persons aged 40 years and above, as these are documented
to have a high risk of CVD [3] [40] [41].

Our study is also consistent with what has been reported in other settings that more
females had CVD than males [42]. Therefore, CVD prevention may be significantly
increased by offering sex-specific risk factor monitoring and intervention strategies [3]
[42]. High blood pressure is one of the risk factors for CVD that has a high prevalence
and is linked to strong evidence of causation [43]. Meta-analyses have shown that
lowering blood pressure can effectively prevent CVDs [42] [44] [45]. Further evidence
suggests that smoking is a key factor responsible for more than 30% of the CVD
mortality [46] [47, pp. 2011–2018]. Therefore, implementing interventions aimed at
smoking cessation or preventing exposure to passive smoking has the potential to
further reduce the risk of CVD.

Our study found that the prevalence of CVD varies geographically throughout Africa, which emphasises how crucial it is to take regional variances into account when creating focused healthcare interventions. In addition, contextual elements such as the varying stages of national NCD program implementation, differences in the rollout of WHO PEN interventions, and diverse cultural and socioeconomic environments likely contribute to the observed patterns [48]. Incorporating these broader contextual influences aligns with evidence from other regions, including Europe, where cultural practices, socioeconomic structures, and policy environments shape both CVD prevalence and management. Including these considerations provides a more comprehensive explanation for cross-country variation in our findings. For example, dietary practices, public health policies, and healthcare infrastructure have all been linked to differences in the burden of CVD between Western and Southern European countries [49]. These results highlight the necessity of investigating and addressing the fundamental causes of regional variations in CVD prevalence in order to create efficient, situation-specific management and prevention plans in Africa.

The increased prevalence of CVDs in Africa is largely caused by high salt intake, especially because it is linked to hypertension, a key risk factor for heart disease and stroke. According to a meta-analysis, the World Health Organization's recommended daily limit of 2 grammes of salt is frequently exceeded in Africa, raising the risk of hypertension and cardiovascular diseases [50]. Regions such as Europe, North America, and Australia have successfully reduced salt consumption through public health measures [51]. For example, in order to prevent hypertension, the American Heart Association recommends avoiding meals high in sodium and focusing on controlling blood pressure through a heart-healthy diet. In Africa, implementing food-based dietary guidelines (FBDGs) has been successful in reducing risk factors for CVD, including high blood pressure and cholesterol. According to studies, choosing a diet that complies with current dietary guidelines decreases blood pressure and cholesterol, which should reduce the risk of CVD by one-third in middle-aged and older people in good health [52]. It has been demonstrated that the Dietary Approaches to Stop Hypertension (DASH) diet, which places an emphasis on consuming less salt, lowers blood pressure, cholesterol, and saturated fats, all of which are risk factors for CVD[53].

In order to prevent premature mortality, lower the increasing burden of non-
communicable diseases in Africa, and limit CVD risk factors like hypertension and
hypercholesterolemia, it is imperative that high salt intake be addressed through multi-
sectoral policies and customised FBDGs.

The CVD were shown to be three times more common among people with hypertension
in Africa than in those with normal blood pressure. This pattern is similar to findings
from other locations, but the risk varies. Due in large part to improved blood pressure
control and easy access to antihypertensive medication, the relative risk of CVD among
hypertensive people in Asia, Latin America, North America and Europe is two to three
times higher compared with those without hypertension [54] [55] [56] [57]. This
emphasises the critical need for better blood pressure screening, reasonably priced
antihypertensive drugs, and strong primary healthcare interventions because
hypertension is a major modifiable risk factor for CVD. Ignoring this problem will
increase the already increasing burden of CVD, worsen health disparities, and put
further strain on the region's already vulnerable healthcare infrastructure. Effectively
managing hypertension could lower the risk of CVD considerably and help achieve
international goals such as the Sustainable Development Goal of lowering premature
mortality from non-communicable diseases by 2030.

Discussion for CVD prevention

Comparing Africa to other regions, the region's 11% adoption rate of CVD preventative
therapy is far lower. Due to strong healthcare systems, easily accessible drugs, and
effective follow-up procedures, the uptake of preventative medicines like aspirin, statins,
and antihypertensives approaches 50%–60% in high-income nations like Europe, North
America, and Australia [58] [59]. Although acceptance rates in Asia vary, they are
typically also high, ranging from 20% to 40%, especially in cities with more advanced
healthcare facilities [56] [60]. Similar to Africa, Latin America shows also uptake rates of
between 30% and 40%, however rural areas encounter similar difficulties [61]. Higher
CVD morbidity and mortality, higher healthcare costs as a result of acute care needs,
and growing disparities in healthcare access are only a few of the serious ramifications

of the poor uptake in Africa [62]. In order to improve CVD preventive care and lower the rising burden of CVD in the region, this finding emphasises the urgent need for targeted interventions like bolstering primary healthcare systems, guaranteeing affordable access to necessary medications, delegating tasks to community health workers, and implementing international initiatives like the WHO's HEARTS program.

According to our study, men were much less likely than women to obtain CVD prevention. This pattern is in accordance with research from other areas, such as Europe and Asia, where women are more likely to follow preventative care recommendations and have higher rates of healthcare utilisation [63]. Because of maternity and reproductive health services, women engage with healthcare systems more frequently, which opens up potential for early CVD risk assessment and intervention. Furthermore, because of cultural norms, a perceived lack of sensitivity to chronic diseases, and poorer health-seeking behavior, men may be less involved in preventive healthcare [64]. This gender gap is made worse in Africa by structural issues including inadequate health care. Systemic issues including low health literacy and resource shortages, especially in rural regions where men predominate, worsen the gender gap in Africa. Public health is significantly impacted by men's lower adoption of CVD prevention. Addressing these inequities is crucial because men in Africa are more likely to suffer from risk factors such smoking, excessive alcohol use, and hypertension [65]. This disparity might be closed and the burden of CVD considerably decreased with gender-specific programs that focus on male health behaviour, raise knowledge, and enhance access to CVD preventive treatment.

Compared to their rural counterparts, urban dwellers had almost double the likelihood of receiving CVD prevention. This gap between urban and rural areas is consistent with international findings showing that urban areas have better access to, infrastructure for, and awareness of, healthcare [66]. Preventive programs that encourage CVD screening and early intervention are more prevalent in urban locations, as are healthcare facilities and specialised physicians. On the other hand, access to preventative treatment is frequently restricted in rural Africa due to factors like lower socioeconomic status, understaffed healthcare facilities, and geographic inaccessibility [67]. Given that the

majority of people in Africa live in rural areas, the effects of this urban-rural divide are

crucial. Rural dwellers will continue to be disproportionately susceptible to untreated

cardiovascular disease risk factors in the absence of focused efforts, which will increase

morbidity and mortality. Increasing access to CVD prevention treatment in rural areas

through community-based initiatives, mobile health interventions, and decentralised

healthcare services must be a top priority for policymakers. For equal progress towards

reaching the Sustainable Development Goal of lowering premature mortality from non-

communicable diseases by 2030, this gap must be closed.

Discussion for CVD treatment uptake

There is not much evidence on the uptake of treatment for CVD in Africa. Just 22% of

people with CVD in Africa received treatment, indicating a significant gap in the

provision of necessary care for those impacted. This number is significantly lower than

in other continents like North America and Europe, where more than 70% of people with

CVD receive treatment because of their well-established healthcare systems, easy

access to drugs, and successful preventative initiatives [68]. Treatment rates range from

40% to 60% across Asia and Latin America, which reflects attempts to scale up the

management of chronic diseases and improve healthcare accessibility [69]. The glaring

disparity in treatment uptake highlights the persistent issues in Africa, such as the lack

of adequate healthcare infrastructure, the high cost of necessary pharmaceuticals, low

knowledge, and a paucity of qualified medical personnel, especially in underserved and

rural areas [60]. This low treatment rate has serious ramifications because untreated

CVDs greatly raises the risk of negative outcomes like heart attack, stroke, and early

death. Given the continued challenges with communicable diseases and the increasing

burden of CVDs in Africa, region-specific therapies are desperately needed. To close

this gap, it is crucial to address socioeconomic barriers, integrate CVD care into basic

healthcare, and provide access to cheap treatment.

In Africa, those with hypertension were seven times more likely than those without to

receive treatment for CVD, highlighting that hypertension not only serves as a crucial

starting point for CVD care but also aids in its diagnosis, given its well-established role

as a major risk factor. This pattern is consistent with findings from North America and

Europe, where clinical guidelines prioritise screening and treatment for hypertensive

people [70] [71]. However, missed chances for early detection and management of CVD

among other high-risk populations, such as those with diabetes, obesity, or tobacco

use, are reflected in the poorer treatment uptake among non-hypertensive adults in

Africa [69]. To close this gap and guarantee fair treatment access for all at-risk persons,

thorough risk assessment instruments and integrated care pathways must be

expanded. In order to meet global goals like the WHO's "25 by 25" program to reduce

early deaths from non-communicable illnesses and lessen the rising burden of CVD in

Africa, these measures must be scaled up.

CVD care cascade

A full understanding of CVD care requires looking at the entire cascade, from screening

and diagnosis through to treatment initiation and long-term adherence. The current

WHO STEPS dataset does not include information on CVD screening, which limits the

ability to assess the full pathway of care. Even so, our analysis shows major gaps: only

11% of participants reported receiving any form of CVD preventive therapy, and just

22% of those with CVD were on treatment. These figures point to large losses at

multiple points in the continuum. The gaps are especially evident among rural

populations and men, who were less likely to be receiving care [38]. To strengthen

future analyses, WHO STEPS surveys should collect data not only on diagnosis and

treatment but also on screening and adherence [38]. Alongside this, complementary

qualitative work or topic modelling could help explain country-specific barriers, giving

clearer insight into how to improve both equity and coverage in CVD care.

To address these gaps, targeted interventions should be prioritized, including

integrating CVD care into primary healthcare systems, task-shifting responsibilities to

community health workers, and ensuring the availability of low-cost essential

medications. When compared to other LMICs, such as countries in Asia and Latin

America, Africa's CVD preventive and treatment uptake remains substantially lower,

underscoring persistent structural and socioeconomic barriers [38]. Future WHO STEPS

iterations could further enhance policy relevance by including detailed care cascade

indicators and socioeconomic measures, enabling researchers and policymakers to

identify critical points for intervention and design context-specific strategies to improve
CVD outcomes.

All stages of the CVD cascade (screening, diagnosis, treatment, etc.) depend on people's recall of prior
care and on each prior step being completed, so country differences often reflect how many people ever
got checked rather than just true disease rates [72]. For example, in Morocco's 2017 survey many older
adults may have had routine blood-pressure or cholesterol tests in strong primary care settings and thus
report knowing their diagnosis, whereas in Ethiopia's 2015 survey a similar person in a rural area may
never have been screened at all. That means Morocco can show a higher percentage "diagnosed" or
"treated" simply because more cases were detected, while underdiagnosis in low-resource settings like
Malawi or Benin shrinks all subsequent steps. Using self-reported pathways also invites recall and
selection biases – for instance, educated or urban patients who seek care are more likely to remain in
the cascade – as experts have warned of "recall bias in self-reported information" and poor
comparability across surveys [73]. In practice, therefore, differences in cascade figures between
countries (e.g. Algeria vs Sudan or Kenya vs Uganda) should be seen as reflecting both epidemiological
variation and health-system capacity (screening reach, diagnostic availability), not as direct measures of
one country "performing" better than another.

Strengths and limitations

This is to our knowledge the first multi-country study analyzing prevalence and uptake
of CVD treatment in Africa using WHO STEPS. There are some analyses at country
level using other sources, but such datasets, unlike WHO STEPS, are not usually
nationally representative[5] [35]. Therefore, the major strengths of this study were the
large sample size and its representativeness, which potentially allows the analysis to
improve operational practice and inform policy change. Another strength is that the
study covered twelve countries and thus provides a good picture of CVD prevalence
and treatment in Africa in general.

However, several limitations should be noted. First, the analysis did not incorporate
country-level health system indicators, such as the availability of essential medicines

(e.g., aspirin or statins) or the existence of national NCD strategies, which may partly

explain inter-country differences. Second, the study relied on self-reported information

regarding heart attacks, CVD treatment, and related risk and protective factors. This

approach may introduce diagnostic and recall bias, particularly in settings with limited

healthcare access, where individuals may be unaware of or unable to accurately report

their condition or treatment history. These factors should be considered when

interpreting the results and may provide avenues for future research. We were,

however, unable to estimate the extent of underestimation or overestimation of the

reported CVD-related risk and protective factors with heart attack and CVD treatment

uptake. Although treatment-related analyses incorporated STEPS individual sampling

weights, primary sampling units and strata could not be specified due to errors arising

from the pooled, harmonized dataset, which may have led to underestimated standard

errors and overly narrow confidence intervals, and no formal sensitivity analyses using

alternative survey specifications were conducted. Furthermore, the data were available

for only one survey round per country, limiting assessment of within-country temporal

trends. Although survey years varied across countries, survey year was not significantly

associated with CVD risk; however, residual temporal heterogeneity may still affect

between-country comparisons.

The study only looked at the general CVD conditions and does not provide information

on the type of CVD that the individuals had. Having such information could help to

choose the most suitable risk reduction interventions. Another limitation is that the WHO

STEPS does not capture information on screening for CVDs although such information

would better help understand the care cascade from those eligible for screening to

uptake of treatment. Besides, wealth quintile is not captured in most of the WHO

STEPS surveys despite wealth index being paramount in determining equity and

inequity of health outcomes. For example, there is evidence from multiple countries

across the world that wealthier individuals had higher prevalence of CVD than poorer

ones [74] [75].

Although wealth quintile data were unavailable, urban/rural residence, gender, and
education were used as proxies to explore socioeconomic gradients in CVD prevention
and treatment uptake. Urban residents and women were more likely to receive care,
reflecting disparities in healthcare access and health-seeking behavior. While country-
specific differences (e.g., Uganda vs. Sudan) were observed, they were not fully
contextualized. Future work could explore the underlying reasons such as healthcare
access, survey methodology, or cultural factors using qualitative approaches or text-
based analyses like topic modeling to better understand country-level patterns.

Conclusion

In conclusion; using a sizable and representative dataset from twelve countries, our
work offers a crucial insight into the prevalence of CVDs and the uptake of its treatment
in Africa. The African context highlights gaps in diagnosis and treatment for CVD
compared to Europe, Asia, and North America. People with hypertension are seven
times more likely to undergo treatment, making hypertension a crucial point of entry for
CVD therapy. Important risk factors, such as smoking and excessive salt consumption,
highlight the need for multi-sectoral treatments like smoking cessation programs and
nutritional recommendations. Notwithstanding its advantages, such as its sizable
sample size and representativeness, the study's dependence on self-reported data and
its absence of details on wealth indices and particular forms of CVDs point to areas that
require more investigation. Achieving global health goals like the Sustainable
Development Goals and lowering the rising burden of CVDs in Africa depend on closing
these disparities and putting equitable, evidence-based interventions into place.
710

Acknowledgement: The authors would like to thank the World Health Organisation for
granting access to use the WHO STEPS data sets for Africa.

REFERENCES

[1]   U. Nations, "Transforming our world: The 2030 agenda for sustainable development," *N. Y. U. N.*
Dep. Econ. Soc. Aff., 2015.

[2]   G. A. Roth, G. A. Mensah, and V. Fuster, "The Global Burden of Cardiovascular Diseases and Risks:
A Compass for Global Action," *J. Am. Coll. Cardiol.*, vol. 76, no. 25, pp. 2980–2981, Dec. 2020, doi:
10.1016/j.jacc.2020.11.021.
[3]   S. Kaptoge *et al.*, "World Health Organization cardiovascular disease risk charts: revised models to
estimate risk in 21 global regions," *Lancet Glob. Health*, vol. 7, no. 10, pp. e1332–e1345, Oct. 2019,
doi: 10.1016/S2214-109X(19)30318-3.
[4]   M. Di Cesare *et al.*, "The Heart of the World.," *Glob. Heart*, vol. 19, no. 1, p. 11, 2024, doi:
10.5334/gh.1288.
[5]   N. W. Minja *et al.*, "Cardiovascular diseases in Africa in the twenty-first century: Gaps and priorities
going forward.," *Front. Cardiovasc. Med.*, vol. 9, p. 1008335, 2022, doi:
10.3389/fcvm.2022.1008335.
[6]   J. Sun, Y. Qiao, M. Zhao, C. G. Magnussen, and B. Xi, "Global, regional, and national burden of
cardiovascular diseases in youths and young adults aged 15–39 years in 204 countries/territories,
1990–2019: a systematic analysis of Global Burden of Disease Study 2019," *BMC Med.*, vol. 21, no.
1, p. 222, Jun. 2023, doi: 10.1186/s12916-023-02925-4.
[7]   Vaduganathan Muthiah, Mensah George A., Turco Justine Varieur, Fuster Valentin, and Roth
Gregory A., "The Global Burden of Cardiovascular Diseases and Risk," *J. Am. Coll. Cardiol.*, vol. 80,
no. 25, pp. 2361–2371, Dec. 2022, doi: 10.1016/j.jacc.2022.11.005.
[8]   L. J. Jack, "Advancing Health Equity, Eliminating Health Disparities, and Improving Population
Health," *Prev. Chronic. Dis.*, vol. 18, p. E79, 2021, doi: 10.5888/pcd18.210264.
[9]   N. Pronk, D. V. Kleinman, S. F. Goekler, E. Ochiai, C. Blakey, and K. H. Brewer, "Promoting Health
and Well-being in Healthy People 2030.," *J. Public Health Manag. Pract. JPHMP*, vol. 27, no. Suppl
6, pp. S242–S248, Dec. 2021, doi: 10.1097/PHH.0000000000001254.
[10]   L. A. Eberly *et al.*, "Cost of integrated chronic care for severe non-communicable diseases at
district hospitals in rural Rwanda," *BMJ Glob. Health*, vol. 4, no. 3, p. e001449, 2019.
[11]   Mbewu A, Mbanya JC., "Cardiovascular Disease. In: Jamison DT, Feachem RG, Makgoba MW, et al.,
editors. Disease and Mortality in Sub-Saharan Africa. 2nd edition. Washington (DC):" The
International Bank for Reconstruction and Development / The World Bank, 2006. Accessed: Jan.
31, 2024. [Online]. Available: https://www.ncbi.nlm.nih.gov/books/NBK2294/
[12]   A. K. Keates, A. O. Mocumbi, M. Ntsekhe, K. Sliwa, and S. Stewart, "Cardiovascular disease in
Africa: epidemiological profile and challenges," *Nat. Rev. Cardiol.*, vol. 14, no. 5, pp. 273–293, May
2017, doi: 10.1038/nrcardio.2017.19.
[13]   M. Palmer *et al.*, "The effectiveness of smoking cessation, physical activity/diet and alcohol
reduction interventions delivered by mobile phones for the prevention of non-communicable
diseases: A systematic review of randomised controlled trials.," *PloS One*, vol. 13, no. 1, p.
e0189801, 2018, doi: 10.1371/journal.pone.0189801.
[14]   M. Palmer *et al.*, "The effectiveness of smoking cessation, physical activity/diet and alcohol
reduction interventions delivered by mobile phones for the prevention of non-communicable
diseases: A systematic review of randomised controlled trials," *PLOS ONE*, vol. 13, no. 1, p.
e0189801, Jan. 2018, doi: 10.1371/journal.pone.0189801.
[15]   J. Stewart, G. Manmathan, and P. Wilkinson, "Primary prevention of cardiovascular disease: A
review of contemporary guidance and literature.," *JRSM Cardiovasc. Dis.*, vol. 6, p.
2048004016687211, Dec. 2017, doi: 10.1177/2048004016687211.
[16]   "World Health Organization (WHO). STEPS: A Framework for Surveillance. The WHO STEPwise
Approach to Surveillance of Noncommunicable Diseases (STEPS). Geneva, Switzerland: WHO;
2003."
[17]   WHO, "STEPwise approach to NCD risk factor surveillance (STEPS)." 2022. Accessed: Jul. 20, 2022.
[Online]. Available: https://www.who.int/teams/noncommunicable-
diseases/surveillance/systems-tools/steps

[18]   "Ministry of Health. . Malawi National STEPwise Survey for Non-Communicable Diseases Risk
767   Factors 2017 Report 2017 Lilongwe, Malawi Ministry of Health."

768   [19]   W. Mutale, R. Chilengi, and Z. M. of Health, *Zambia STEPS Survey for Non-communicable Diseases
769   Risk Factors: Zambia Report for 2017*. Republic of Zambia, Ministry of Health, 2018. [Online].
770   Available: https://books.google.mw/books?id=LqXO0AEACAAJ

771   [20]   World Health Organization, *WHO Package of Essential Noncommunicable (PEN) Disease
772   Interventions for Primary Health Care in Low-Resource Settings*. World Health Organization, 2021.
773   [Online]. Available: https://www.knowledge-action-portal.com/sites/default/files/2025-
774   03/WHO%20PEN%20package%20of%20essential%20NCD%20interventions%202021.pdf

775   [21]   K. Bush, D. R. Kivlahan, M. B. McDonell, S. D. Fihn, and K. A. Bradley, "The AUDIT alcohol
776   consumption questions (AUDIT-C): an effective brief screening test for problem drinking.
777   Ambulatory Care Quality Improvement Project (ACQUIP). Alcohol Use Disorders Identification
778   Test.," *Arch. Intern. Med.*, vol. 158, no. 16, pp. 1789–1795, Sep. 1998, doi:
779   10.1001/archinte.158.16.1789.

780   [22]   F. C. Bull *et al.*, "World Health Organization 2020 guidelines on physical activity and sedentary
781   behaviour.," *Br. J. Sports Med.*, vol. 54, no. 24, pp. 1451–1462, Dec. 2020, doi: 10.1136/bjsports-
782   2020-102955.

783   [23]   E. Osei *et al.*, "Prevalence and predictors of selected risk factors of non-communicable diseases in
784   Ghana: evidence from a sub-national survey," *J. Glob. Health Sci.*, vol. 3, no. 2, pp. 1–14, 2021, doi:
785   10.35500/jghs.2021.3.e13.

786   [24]   O. Harel *et al.*, "Multiple Imputation for Incomplete Data in Epidemiologic Studies.," *Am. J.
787   Epidemiol.*, vol. 187, no. 3, pp. 576–584, Mar. 2018, doi: 10.1093/aje/kwx349.

788   [25]   K. Kim, "Risk stratification of cardiovascular disease according to age groups in new prevention
789   guidelines: a review," *J. Lipid Atheroscler.*, vol. 12, no. 2, p. 96, 2023.

790   [26]   J. I. Davies *et al.*, "Estimating the burden of cardiovascular risk in community dwellers over 40
791   years old in South Africa, Kenya, Burkina Faso and Ghana," *BMJ Glob. Health*, 2021, doi:
792   10.1136/bmjgh-2020-003499.

793   [27]   F. Agyekum *et al.*, "Atherosclerotic cardiovascular disease risk among Ghanaians: A comparison of
794   the risk assessment tools.," *Am. J. Prev. Cardiol.*, vol. 18, p. 100670, Jun. 2024, doi:
795   10.1016/j.ajpc.2024.100670.

796   [28]   M. R. Mahanani *et al.*, "Change in WHO cardiovascular risk prediction over three years in PLWH on
797   efavirenz-based ART.," *South. Afr. J. HIV Med.*, vol. 26, no. 1, p. 1697, 2025, doi:
798   10.4102/sajhivmed.v26i1.1697.

799   [29]   N. M. Tapela *et al.*, "Integrating noncommunicable disease services into primary health care,
800   Botswana.," *Bull. World Health Organ.*, vol. 97, no. 2, pp. 142–153, Feb. 2019, doi:
801   10.2471/BLT.18.221424.

802   [30]   *World Health Organization. WHO STEPS: R package [software]. Geneva: World Health
803   Organization; 2023. Available from: https://github.com/WorldHealthOrganization/WHO_STEPS*.

804   [31]   South African National Department of Health, "Primary Healthcare Chapter 4: Cardiovascular
805   Conditions, with Supporting NEMLC Report Appendix Reviews 2020-4 Review, Version 1.0, 1
806   November 2024," South African National Department of Health, Pretoria, South Africa, 2024.
807   [Online]. Available: https://www.health.gov.za/wp-content/uploads/2024/12/Primary-Healthcare-
808   Chapter-4_Cardiovascular-Conditions-with-supporting-NEMLC-report-appendix-reviews-2020-4-
809   review_Version-1.0-1-November-20.pdf

810   [32]   WHO, "The WHO STEPwise approach to noncommunicable disease risk factor surveillance." WHO,
811   Geneva, Switzerland, Jan. 26, 2017. Accessed: Apr. 20, 2024. [Online]. Available:
812   https://www.who.int/docs/default-source/ncds/ncd-surveillance/steps/steps-
813   manual.pdf?sfvrsn=c281673d_5

[33] "Simmonds, M. and Higgins, J. P. T. (2016). A general framework for the use of logistic regression
models in meta-analysis. Statistical Methods in Medical Research, 25(6), 2858-2877.
https://doi.org/10.1177/0962280214534409."
[34] "Ripley, B. and Venables, B. (2009). Mass: support functions and datasets for venables and ripley's
mass. CRAN: Contributed Packages. https://doi.org/10.32614/cran.package.mass."
[35] V. N. Agbor, M. Essouma, N. A. Ntusi, U. F. Nyaga, J. J. Bigna, and J. J. Noubiap, "Heart failure in
sub-Saharan Africa: a contemporaneous systematic review and meta-analysis," *Int. J. Cardiol.*, vol.
257, pp. 207–215, 2018.
[36] G. A. Mensah, V. Fuster, C. J. L. Murray, and G. A. Roth, "Global Burden of Cardiovascular Diseases
and Risks, 1990-2022.," *J. Am. Coll. Cardiol.*, vol. 82, no. 25, pp. 2350–2473, Dec. 2023, doi:
10.1016/j.jacc.2023.11.007.
[37] C. W. Tsao *et al.*, "Heart Disease and Stroke Statistics—2022 Update: A Report From the American
Heart Association," *Circulation*, vol. 145, no. 8, pp. e153–e639, Feb. 2022, doi:
10.1161/CIR.0000000000001052.
[38] N. W. Minja *et al.*, "Cardiovascular diseases in Africa in the twenty-first century: Gaps and priorities
going forward.," *Front. Cardiovasc. Med.*, vol. 9, p. 1008335, 2022, doi:
10.3389/fcvm.2022.1008335.
[39] B. J. North and D. A. Sinclair, "The Intersection Between Aging and Cardiovascular Disease," *Circ.
Res.*, vol. 110, no. 8, pp. 1097–1108, Apr. 2012, doi: 10.1161/CIRCRESAHA.111.246876.
[40] Cho Leslie *et al.*, "Summary of Updated Recommendations for Primary Prevention of
Cardiovascular Disease in Women," *J. Am. Coll. Cardiol.*, vol. 75, no. 20, pp. 2602–2618, May 2020,
doi: 10.1016/j.jacc.2020.03.060.
[41] *Cardiovascular disease: risk assessment and reduction, including lipid modification. London:
National Institute for Health and Care Excellence (NICE); 2023 Dec 14. (NICE Clinical Guidelines, No.
238.) Available from: https://www.ncbi.nlm.nih.gov/books/NBK602380/.*
[42] A. K. McClain, P. P. Monteleone, and J. Zoldan, "Sex in cardiovascular disease: Why this biological
variable should be considered in in vitro models," *Sci. Adv.*, vol. 10, no. 19, p. eadn3510, doi:
10.1126/sciadv.adn3510.
[43] F. D. Fuchs and P. K. Whelton, "High Blood Pressure and Cardiovascular Disease.," *Hypertens.
Dallas Tex 1979*, vol. 75, no. 2, pp. 285–292, Feb. 2020, doi:
10.1161/HYPERTENSIONAHA.119.14240.
[44] J. D. Bundy *et al.*, "Systolic blood pressure reduction and risk of cardiovascular disease and
mortality: a systematic review and network meta-analysis," *JAMA Cardiol.*, vol. 2, no. 7, pp. 775–
781, 2017.
[45] J. D. Williamson *et al.*, "Effect of intensive vs standard blood pressure control on probable
dementia: a randomized clinical trial," *Jama*, vol. 321, no. 6, pp. 553–561, 2019.
[46] G. Gallucci, A. Tartarone, R. Lerose, A. V. Lalinga, and A. M. Capobianco, "Cardiovascular risk of
smoking and benefits of smoking cessation.," *J. Thorac. Dis.*, vol. 12, no. 7, pp. 3866–3876, Jul.
2020, doi: 10.21037/jtd.2020.02.47.
[47] A. Mambo, Y. Yang, E. Mahulu, and Z. Zihua, "Investigating the interplay of smoking, cardiovascular
risk factors, and overall cardiovascular disease risk: NHANES analysis 2011–2018," *BMC Cardiovasc.
Disord.*, vol. 24, no. 1, p. 193, Apr. 2024, doi: 10.1186/s12872-024-03838-7.
[48] R. Ferrari *et al.*, "Geographical variations in the prevalence and management of cardiovascular risk
factors in outpatients with CAD: Data from the contemporary CLARIFY registry," *Eur. J. Prev.
Cardiol.*, vol. 22, no. 8, pp. 1056–1065, Aug. 2015, doi: 10.1177/2047487314547652.
[49] A. Timmis *et al.*, "Cardiovascular disease care and outcomes in West and South European
countries.," *Lancet Reg. Health Eur.*, vol. 33, p. 100718, Oct. 2023, doi:
10.1016/j.lanepe.2023.100718.

[50] O. Oyebode, S. Oti, Y.-F. Chen, and R. J. Lilford, "Salt intakes in sub-Saharan Africa: a systematic
863   review and meta-regression," *Popul. Health Metr.*, vol. 14, no. 1, p. 1, Jan. 2016, doi:
864   10.1186/s12963-015-0068-7.
865   [51] J. A. Santos *et al.*, "A Systematic Review of Salt Reduction Initiatives Around the World: A Midterm
866   Evaluation of Progress Towards the 2025 Global Non-Communicable Diseases Salt Reduction
867   Target," *Adv. Nutr.*, vol. 12, no. 5, pp. 1768–1780, Sep. 2021, doi: 10.1093/advances/nmab008.
868   [52] A. K. Amegah, "Tackling the Growing Burden of Cardiovascular Diseases in Sub-Saharan Africa,"
869   *Circulation*, vol. 138, no. 22, pp. 2449–2451, Nov. 2018, doi:
870   10.1161/CIRCULATIONAHA.118.037367.
871   [53] *Challa HJ, Ameer MA, Uppaluri KR. DASH Diet To Stop Hypertension. [Updated 2023 Jan 23]. In:*
872   *StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from:*
873   *https://www.ncbi.nlm.nih.gov/books/NBK482514/?utm_source=chatgpt.com.*
874   [54] K. T. Mills *et al.*, "Global Disparities of Hypertension Prevalence and Control: A Systematic Analysis
875   of Population-Based Studies From 90 Countries.," *Circulation*, vol. 134, no. 6, pp. 441–450, Aug.
876   2016, doi: 10.1161/CIRCULATIONAHA.115.018912.
877   [55] "Worldwide trends in blood pressure from 1975 to 2015: a pooled analysis of 1479 population-
878   based measurement studies with 19·1 million participants.," *Lancet Lond. Engl.*, vol. 389, no.
879   10064, pp. 37–55, Jan. 2017, doi: 10.1016/S0140-6736(16)31919-5.
880   [56] S. Ma, L. Yang, M. Zhao, C. G. Magnussen, and B. Xi, "Trends in hypertension prevalence,
881   awareness, treatment and control rates among Chinese adults, 1991-2015.," *J. Hypertens.*, vol. 39,
882   no. 4, pp. 740–748, Apr. 2021, doi: 10.1097/HJH.0000000000002698.
883   [57] K. Kario, A. Okura, S. Hoshide, and M. Mogi, "The WHO Global report 2023 on hypertension
884   warning the emerging hypertension burden in globe and its treatment strategy," *Hypertens. Res.*,
885   vol. 47, no. 5, pp. 1099–1102, May 2024, doi: 10.1038/s41440-024-01622-w.
886   [58] G. A. Roth *et al.*, "Global, Regional, and National Burden of Cardiovascular Diseases for 10 Causes,
887   1990 to 2015.," *J. Am. Coll. Cardiol.*, vol. 70, no. 1, pp. 1–25, Jul. 2017, doi:
888   10.1016/j.jacc.2017.04.052.
889   [59] B. Zhou *et al.*, "Worldwide trends in hypertension prevalence and progress in treatment and
890   control from 1990 to 2019: a pooled analysis of 1201 population-representative studies with 104
891   million participants," *The Lancet*, vol. 398, no. 10304, pp. 957–980, Sep. 2021, doi: 10.1016/S0140-
892   6736(21)01330-1.
893   [60] R. Khatib *et al.*, "Availability and affordability of cardiovascular disease medicines and their effect
894   on use in high-income, middle-income, and low-income countries: an analysis of the PURE study
895   data.," *Lancet Lond. Engl.*, vol. 387, no. 10013, pp. 61–69, Jan. 2016, doi: 10.1016/S0140-
896   6736(15)00469-9.
897   [61] Á. Avezum *et al.*, "Challenges and Opportunities to Scale Up Cardiovascular Disease Secondary
898   Prevention in Latin America and the Caribbean.," *Glob. Heart*, vol. 13, no. 2, pp. 83–91, Jun. 2018,
899   doi: 10.1016/j.gheart.2017.05.002.
900   [62] M. F. Yuyun, K. Sliwa, A. P. Kengne, A. O. Mocumbi, and G. Bukhman, "Cardiovascular Diseases in
901   Sub-Saharan Africa Compared to High-Income Countries: An Epidemiological Perspective.," *Glob.*
902   *Heart*, vol. 15, no. 1, p. 15, Feb. 2020, doi: 10.5334/gh.403.
903   [63] H.-L. Kim, "Differences in Risk Factors for Coronary Atherosclerosis According to Sex.," *J. Lipid*
904   *Atheroscler.*, vol. 13, no. 2, pp. 97–110, May 2024, doi: 10.12997/jla.2024.13.2.97.
905   [64] A. Peerson and M. Saunders, "Men's health literacy: advancing evidence and priorities," *Crit. Public*
906   *Health*, vol. 19, no. 3–4, pp. 441–456, Sep. 2009, doi: 10.1080/09581590902906229.
907   [65] N. H. Phelps *et al.*, "Worldwide trends in underweight and obesity from 1990 to 2022: a pooled
908   analysis of 3663 population-representative studies with 222 million children, adolescents, and
909   adults," *The Lancet*, vol. 403, no. 10431, pp. 1027–1050, Mar. 2024, doi: 10.1016/S0140-
910   6736(23)02750-2.

[66] R. Gupta, "Convergence in urban-rural prevalence of hypertension in India.," *J. Hum. Hypertens.*,
vol. 30, no. 2, pp. 79–82, Feb. 2016, doi: 10.1038/jhh.2015.48.
[67] S. Dunachie and P. Chamnan, "The double burden of diabetes and global infection in low and
middle-income countries.," *Trans. R. Soc. Trop. Med. Hyg.*, vol. 113, no. 2, pp. 56–64, Feb. 2019,
doi: 10.1093/trstmh/try124.
[68] S. S. Virani *et al.*, "Heart Disease and Stroke Statistics-2021 Update: A Report From the American
Heart Association.," *Circulation*, vol. 143, no. 8, pp. e254–e743, Feb. 2021, doi:
10.1161/CIR.0000000000000950.
[69] *Rao H. Use of secondary prevention drugs for cardiovascular disease in the community in high-*
income, middle-income, and low-income countries (the PURE Study): a prospective epidemiological
survey. Indian Heart J. 2012 Jan;64(1):113. doi: 10.1016/S0019-4832(12)60034-0. PMCID:
PMC3861210.
[70] K. Kotseva *et al.*, "Lifestyle and risk factor management in people at high risk of cardiovascular
disease. A report from the European Society of Cardiology European Action on Secondary and
Primary Prevention by Intervention to Reduce Events (EUROASPIRE) IV cross-sectional survey in 14
European regions.," *Eur. J. Prev. Cardiol.*, vol. 23, no. 18, pp. 2007–2018, Dec. 2016, doi:
10.1177/2047487316667784.
[71] K. T. Mills *et al.*, "Global Disparities of Hypertension Prevalence and Control: A Systematic Analysis
of Population-Based Studies From 90 Countries.," *Circulation*, vol. 134, no. 6, pp. 441–450, Aug.
2016, doi: 10.1161/CIRCULATIONAHA.115.018912.
[72] A. O. Diallo *et al.*, "Multiple cardiovascular risk factor care in 55 low- and middle-income countries:
A cross-sectional analysis of nationally-representative, individual-level data from 280,783 adults.,"
PLOS Glob. Public Health, vol. 4, no. 3, p. e0003019, 2024, doi: 10.1371/journal.pgph.0003019.
[73] M. E. Marcus *et al.*, "Unmet need for hypercholesterolemia care in 35 low- and middle-income
countries: A cross-sectional study of nationally representative surveys.," *PLoS Med.*, vol. 18, no. 10,
p. e1003841, Oct. 2021, doi: 10.1371/journal.pmed.1003841.
[74] C. Brindley, T. Van Ourti, J. Capuno, A. Kraft, J. Kudymowa, and O. O'Donnell, "Risk factor
contributions to socioeconomic inequality in cardiovascular risk in the Philippines: a cross-sectional
study of nationally representative survey data," *BMC Public Health*, vol. 23, no. 1, p. 689, Apr.
2023, doi: 10.1186/s12889-023-15517-x.
[75] A. Rosengren *et al.*, "Socioeconomic status and risk of cardiovascular disease in 20 low-income,
middle-income, and high-income countries: the Prospective Urban Rural Epidemiologic (PURE)
study," *Lancet Glob. Health*, vol. 7, no. 6, pp. e748–e760, Jun. 2019, doi: 10.1016/S2214-
109X(19)30045-2.
945

947 Sample simulation code

(a) *Simulation of discrete variables [(just a few)] like sex, physical activity, smoking status, harmful alcohol use*

948
949

950   i.   *Physical activity*

```
951    #Physical activity level
952    set.seed(123)  # Set seed for reproducibility
953
954    # Proportions for each physical activity level
955    pa_proportions <- c("1) Low Level" = 0.16, "2) Moderate level" = 0.20, "3) High level" =
956    0.64)
957
958    # Get the indices of NA values in cvd_africa_2024$pa
959    na_indices_pa <- which(is.na(cvd_africa_2024$pa))
960
961    # Number of NA values
962    n_na_pa <- length(na_indices_pa)
963
964    # Sample replacement values based on the given proportions
965    replacement_pa_values <- sample(
966      names(pa_proportions),     # Possible values ("Low Level", "Moderate Level", "High
967    Level")
968      size = n_na_pa,           # Number of values to sample
969      replace = TRUE,           # Sampling with replacement
970      prob = pa_proportions     # The given probabilities
971    )
972
973    # Replace the NA values with the sampled values
974    cvd_africa_2024$pa[na_indices_pa] <- replacement_pa_values
975
976    # Check the updated column
977    table(cvd_africa_2024$pa)
978
979    ii.      Harmful alcohol use
980    cvd_africa_2024$alcharm <- factor(cvd_africa_2024$alcharm)
981    set.seed(123) ##  Set seed for reproducibility
```

```
982      #
983      #Proportions for each category
984      proportions_alcharm <- c("No" = 0.97, "Yes" = 0.03)
985      #
986      #Get the indices of NA values in cvd_africa_2024$alcharm
987      na_indices_alcharm <- which(is.na(cvd_africa_2024$alcharm))
988      #
989      #Number of NA values
990      n_na_alcharm <- length(na_indices_alcharm)
991      #
992      #Sample replacement values based on the given proportions
993      replacement_values_alcharm <- sample(
994        names(proportions_alcharm),
995        size = n_na_alcharm,
996        replace = TRUE,
997        prob = proportions_alcharm
998      )
999
1000     iii.      Smoking history
1001     set.seed(123) ## Set seed for reproducibility
1002     #
1003     #Proportions for each education level
1004     proportions <- c("Current" = 0.1, "Never" = 0.83, "Previous" = 0.07)
1005     #
1006     #Get the indices of NA values in cvd_africa_2024$edlev
1007     na_indices <- which(is.na(cvd_africa_2024$smokehis))
1008     #
1009     #Number of NA values
1010     n_na <- length(na_indices)
1011     #
1012     #Sample replacement values based on the given proportions
1013     replacement_values <- sample(
1014       names(proportions),        #Possible values ("None", "Primary", etc.)
1015       size = n_na,               #Number of values to sample
1016       replace = TRUE,            #Sampling with replacement
1017       prob = proportions         #The given probabilities
```

```
1018            )
1019            #
1020            #Replace the NA values with the sampled values
1021            cvd_africa_2024$smokehis[na_indices] <- replacement_values
1022            #
1023            #Check the updated column
1024            table(cvd_africa_2024$smokehis)
1025            cvd_africa_2024$smokehis <- factor(cvd_africa_2024$smokehis)
1026            var_label(cvd_africa_2024$smokehis) <- "Smoking history"
1027
1028
1029            iv.      Sex
1030            set.seed(123) ## Set seed for reproducibility
1031            #
1032            #Proportions for gender
1033            gender_proportions <- c("Female" = 0.49, "Male" = 0.51)
1034            #
1035            #Get the indices of NA values in cvd_africa_2024$gender
1036            na_gender_indices <- which(is.na(cvd_africa_2024$gender))
1037            #
1038            #Number of NA values
1039            n_na_gender <- length(na_gender_indices)
1040            #
1041            #Sample replacement values based on the given proportions
1042            gender_replacements <- sample(
1043             names(gender_proportions),   #Possible values ("Female", "Male")
1044             size = n_na_gender,          #Number of values to sample
1045             replace = TRUE,              #Sampling with replacement
1046             prob = gender_proportions    #The given probabilities
1047            )
1048

1049    (b) Simulation of continuous variables like age, BMI and cholesterol levels
1050            i.       BMI
1051                    set.seed(123)
1052                    cvd_africa_2024$bmi[is.na(cvd_africa_2024$bmi)] <- rnorm(
```

```
1053          sum(is.na(cvd_africa_2024$bmi)),  #Number of NA values
1054           mean = 24.47,              # Mean value
1055           sd = 11.88                 #  Standard deviation
1056          )
1057
1058    ii.   Cholesterol levels
1059          ##  Set seed for reproducibility
1060          set.seed(123)
1061          # Define the mean and#  Standard deviation
1062          mean_cholesterol <- 2.827
1063          sd_cholesterol <- 1.502
1064          # Find the indices of NA values in cvd_africa_2024$cholesterol
1065          na_indices_cholesterol <- which(is.na(cvd_africa_2024$cholesterol))
1066          #
1067          #Number of NA values
1068          n_na_cholesterol <- length(na_indices_cholesterol)
1069          #Generate random values from the normal distribution
1070          replacement_cholesterol_values <- rnorm(n_na_cholesterol, mean =
1071          mean_cholesterol, sd = sd_cholesterol)
1072          #Replace the NA values with the generated values
1073          cvd_africa_2024$cholesterol[na_indices_cholesterol] <-
1074          replacement_cholesterol_values
1075          # Replace cholesterol values with their absolute values
1076          cvd_africa_2024$cholesterol <- abs(cvd_africa_2024$cholesterol)
1077          # Check the updated cholesterol column
1078          summary(cvd_africa_2024$cholesterol)
1079
1080    iii.  Age
1081          set.seed(123)
1082          cvd_africa_2024$age[is.na(cvd_africa_2024$age)] <- rnorm(
1083           sum(is.na(cvd_africa_2024$age)),  #Number of NA values
1084           mean = 37.93,              # Mean value
1085           sd = 13.52                 #  Standard deviation
1086          )
1087
1088
```

1089

1_Create_dataset_for_analysis.R: Code for generating the dataset for analysis

2_who_steps_ncd_data_cleaning_code_2024.R: Code for data cleaning

3_CVD_Analysis_SSA_2024_FINAL.Rmd: R markdown file for data analysis

4_CVD sensitivity analysis 2025.R: Code for performing sensitivity analysis of the complete case analysis
vs imputed data

Functions_rmph.R: Miscellaneous functions used for data analysis