Inborn errors of immunity: manifestation, treatment, and outcome – an ESID registry 1994-2024 report on 30,628 patients

Gerhard Kindle[1, 2, 42], Mickaël Alligon[3, 42], Michael H. Albert[4], Matthew Buckland[5, 6], J. David Edgar[7], Benjamin Gathmann[1], Sujal Ghosh[8], Antonios Gkantaras[9], Alexandra Nieters[1, 2], Claudio Pignata[10], Peter Robinson[11, 12], Stephan Rusch[1, 2], Catharina Schuetz[13, 14], Svetlana Sharapova[15], Benjamin Shillitoe[16], Fabio Candotti[17], Andrew J. Cant[18, 19], Jean-Laurent Casanova[20, 21, 22, 23, 24], Amos Etzioni[25], Alain Fischer[26, 21], Isabelle Meyts[27], Luigi D. Notarangelo[28], Martine Pergent[29], C. I. Edvard Smith[30, 31, 32], ESID Registry Working Party, Lennart Hammarström[33, 43], Bodo Grimbacher[1, 34, 35, 36, 37, 43], Mikko Seppänen[38, 39, 43], Nizar Mahlaoui[3, 40, 43], Stephan Ehl[1, 43], and Markus G. Seidel[41, 43, 44]

1. Institute for Immunodeficiency, Center for Chronic Immunodeficiency, Medical Center, University of Freiburg, Faculty of Medicine, University of Freiburg, Freiburg, Germany
2. Centre for Biobanking FREEZE, Medical Center-University of Freiburg, Faculty of Medicine, University of Freiburg, Freiburg, Germany
3. French National Reference Center for Primary Immune Deficiencies (CEREDIH), Necker Enfants Malades University Hospital, Assistance Publique-Hôpitaux de Paris (AP-HP), Paris, France.
4. Department of Pediatrics, Dr. von Hauner Children's Hospital, Ludwig-Maximilians Universität München, Munich Germany
5. Barts Health National Health Service Trust, London, United Kingdom
6. Molecular and Cellular Immunology Section, Immunity and Inflammation Department, Great Ormond Street Institute of Child Health, London, United Kingdom
7. Department of Immunology, St James's Hospital, & School of Medicine, Trinity College Dublin, Ireland
8. Department of Pediatric Oncology, Hematology and Clinical Immunology, Medical Faculty, Center of Child and Adolescent Health, Heinrich-Heine-University and University Hospital, Duesseldorf, Germany
9. Pediatric Immunology and Rheumatology Referral Center, 1st Department of Pediatrics, Aristotle University of Thessaloniki, Hippokration General Hospital, Thessaloniki, Greece
10. Department of Translational Medical Sciences, Federico II University, Naples, Italy
11. Berlin Institute of Health, Charité - Universitätsmedizin Berlin, Berlin, Germany
12. The Jackson Laboratory for Genomic Medicine, Farmington, CT, USA
13. Department of Pediatrics and University Center for Rare Diseases, Medizinische Fakultät Carl Gustav Carus, Technische Universität, Dresden, Germany
14. German Center for Child and Adolescent Health (DZKJ), partner site Leipzig/Dresden, Dresden, Germany
15. Research Department, Belarusian Research Center for Pediatric Oncology, Hematology and Immunology, Minsk Region, Belarus
16. Paediatric Immunology and Infectious Diseases, Sheffield Children's NHS Foundation Trust, Sheffield, United Kingdom
17. Division of Immunology and Allergy, Lausanne University Hospital and University of Lausanne, Lausanne, Switzerland
18. Translational and Clinical Research Institute, Newcastle University
19. Department of Paediatric Immunology and Infectious Diseases, Great North Children's Hospital, Newcastle upon Tyne, England, UK
20. Laboratory of Human Genetics of Infectious Diseases, INSERM U1163, Necker Hospital for Sick Children, Paris, France

21. Imagine Institute, University Paris Cité, Paris, France

22. St Giles Laboratory of Human Genetics of Infectious Diseases, Rockefeller Branch, The Rockefeller University, New York, NY, USA

23. Howard Hughes Medical Institute, New York, NY, USA

24. Department of Pediatrics, Necker Hospital for Sick Children, AP-HP, Paris, France

25. Faculty of Medicine, Technion, Haifa, Israel

26. Unité d'immunologie pédiatrique, Hôpital Necker Enfants Malades, Paris

27. Department of Pediatrics, Department of Microbiology, Immunology and Transplantation, University Hospitals Leuven, KU Leuven, Leuven, Belgium, ERN-RITA Core Center

28. Laboratory of Clinical Immunology and Microbiology, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD, USA

29. International Patient Organisation for Primary Immunodeficiencies (IPOPI), Brussels, Belgium

30. Department of Laboratory Medicine, Clinical Immunology, Karolinska Institutet, Stockholm

31. Department of Infectious Diseases, Karolinska University Hospital, Stockholm, Sweden

32. Karolinska ATMP Center, Karolinska Institutet, Karolinska University Hospital, Stockholm, Sweden

33. Division of Immunology, Dept of Medical Biochemistry and Biophysics, Karolinska Institutet, Stockholm, Sweden

34. Clinic of Rheumatology and Clinical Immunology, Center for Chronic Immunodeficiency (CCI), Medical Center, Faculty of Medicine, Albert-Ludwigs-University of Freiburg, Germany

35. DZIF - German Center for Infection Research, Satellite Center Freiburg, Germany

36. CIBSS - Centre for Integrative Biological Signalling Studies, Albert-Ludwigs University, Freiburg, Germany

37. RESIST - Cluster of Excellence 2155 to Hanover Medical School, Satellite Center Freiburg, Germany

38. Rare Disease Center and Pediatric Research Center, Children and Adolescents, University of Helsinki, HUS Helsinki University Hospital, ERN-RITA Core Center, RITAFIN, Helsinki

39. Translational Immunology Research Program, University of Helsinki, Finland

40. Pediatric Immuno-Hematology and Rheumatology Unit, Necker Enfants Malades University Hospital, Assistance Publique-Hôpitaux de Paris (AP-HP), Paris, France

41. Styrian Children's Cancer Research Unit for Cancer and Inborn Errors of the Blood and Immunity in Children; Division of Pediatric Hematology and Oncology, Department of Pediatric and Adolescent Medicine, Medical University of Graz, Graz, Austria

42. *These authors contributed equally and should be considered aequo loco.*

43. *These authors jointly designed, coordinated, and supervised the study.*

44. *Corresponding author: Univ.-Prof. Dr. Markus G. Seidel,* Division of Pediatric Hematology and Oncology, Medical University of Graz, Auenbruggerplatz 38, 8036 Graz, Austria; Email: markus.seidel@medunigraz.at

Key words: Inborn error of immunity (IEI); primary immunodeficiency (PID); primary immune disorder (PID); online database; patient registry; overall survival.

Word count: abstract: 173; summary: 51; main text: 3463; tables: 1; figures 5; references: 30; supplementary material.

Conflict of interest statement at the end of the article.

ORCID identifiers: GK ORCID: 0000-0003-0894-5327; MHA ORCID: 0000-0001-9143-3263; JDE ORCID: 0000-0003-2969-4475; AG ORCID: 0000-0003-3609-1824; BS ORCID: 0000-0003-4407-5998; CIES ORCID: 0000-0003-1907-3392; MGS ORCID: 0000-0003-0981-8661.

Abstract

The *European Society for Immunodeficiencies* patient registry (ESID-R), established in 1994, is one of the world's largest databases on inborn errors of immunity (IEI). IEI are genetic disorders predisposing patients to infections, autoimmunity, inflammation, allergies and malignancies. Treatments include antimicrobial therapy, immunoglobulin replacement, immune modulation, stem cell transplantation and gene therapy. Data from 194 centers in 33 countries capture clinical manifestations and treatments from birth onward, with annually expected updates. This report reviews the ESID-R's structure, data content, and impact.

The registry includes 30,628 patient datasets (aged 0–97.9 years; median follow-up: 7.2 years; total 825,568.2 patient-years), with 13,550 cases in 15 sub-studies. It has produced 84 peer-reviewed publications (mean citation rate: 95). Findings highlight real-world observations of IEI diagnoses, genetic causes, treatments, and survival trends. The ESID-R fosters global collaboration, advancing IEI research and patient care. Early declining survival probabilities for many IEI subcategories suggest expanding newborn screening beyond severe combined immune deficiencies. This report highlights the key role of the multi-national ESID-R, led by an independent medical society, in evidence-based discovery.

Summary

Worldwide, most patient registries for inborn errors of immunity are national, with limited geographical and temporal scope. This 30-year ESID registry analysis of 30,628 patients' longitudinal datasets enables robust epidemiological studies on natural disease courses including diagnosis, treatment, and survival, supporting expanded newborn screening and future AI applications in IEI research.

Introduction

Inborn errors of immunity (IEI) are genetic disorders affecting immunity. They almost invariably increase the susceptibility to infections and may cause autoimmune, inflammatory, allergic and atopic diseases, and predispose individuals to malignancy(1-6). IEI can manifest at any age and are characterized by a spectrum of symptoms related to impaired or uncontrolled immune responses. These often have a serious impact on the health and quality of life of those affected. Many IEI are rare diseases, including ultra-rare and hyper-rare forms (*i.e.*, frequently <<1/2000 persons affected). Thus, an international strategy was needed to collect a meaningful number of datapoints for cross-sectional analyses and longitudinal clinical observations of natural courses of specific IEI.

The *European Society for Immunodeficiencies* registry (ESID-R) was founded in 1994 by ESID, a non-profit medical specialist society, creating a central database for primary immunodeficiencies. These are currently referred to as IEI or primary immune disorders (PID), differentiating them from secondary immune disorders (SID)(7). The ESID-R mainly serves to contain, store, and enable the analysis of IEI/PID data to improve the understanding of these diseases and their underlying immunobiology. The ESID-R was operated for about 10 years as a hard copy-based database, and data were submitted by fax to the first chairs of the ESID-R, Lennart Hammarström and Mohammad Abedi in Huddinge, Sweden. From 2000–2004, Bodo Grimbacher, then head of the ESID-R, Barbara Frisch and Viviane Knerr, all based in Freiburg, Germany, developed and implemented the first web-based version of the ESID-R with the help of Dominik Veit and Stephan Rusch (programmers) and financial support from the *Plasma Protein Therapeutics Association* (PPTA). Since 2004, the online ESID-R has been operated by the Medical Center of the University of Freiburg(8 9). In 2014, a major revision was carried out by Stephan Ehl in Freiburg and programmers Benjamin Gathmann and Stephan Rusch. The migration to the new system also included quality checks and a validation process to meet the clinical working criteria for IEI/PID diagnoses compiled and published by a group of experts(10). The two key drivers of this re-design were the goal to allow registry participation at three different levels according to center resources and the need to improve data quality so that a high level of confidence in the accuracy of the data could be inferred for clinical application and publication. In late 2024, ESID modernized the technical foundation and moved ESID-R to a commercial clinical trials operator (Castor-edc, NL) to improve data-, system-, and access security. The new version is expected to facilitate the generation of data modules by research groups, allowing decentralized sub-study (eCRF) programming, independent data exports for center or sub-study analyses, automated center dashboards and add-on features such as patient reporting.

Here, we present major findings from clinical observations of 30,628 patients, based on the current ESID-R dataset. These findings indicate epidemiologically relevant disease distributions and the calculated prevalence of diagnoses in registered patients, their clinical manifestations, diagnostic delays, treatment, disease course and survival probabilities across all IEI/PID categories. Furthermore, we describe the organizational and technical evolution of the ESID-R, its relationship with the international registry landscape, and its role as a research (sub-)study platform. These data are of the utmost relevance to anyone affected by immune disorders or involved in patient care, management and therapy, drug and policy development for patients with IEI/PID around the world.

Methods

Technical background and operating mode of the ESID-R

The study protocol, the patient informed consent template and the center data transfer agreement plus amendments thereof were approved by the IRB and the data protection officer at the Medical University of Freiburg, Germany (Albert-Ludwigs-University). A substantial amendment was approved by the Medical University of Graz, Austria (24-334 ex 11/12, IRB00002556) and implemented in 2024. Details of the operational structure and technical background are found in the *Supplementary Material*.

Data and statistical analyses

The study population included all patients with IEI/PID recorded in the ESID-R on March 19, 2024. Patients considered as discharged (*n*=538), patients with secondary immunodeficiency who were documented as part of one national subregistry only (*n*=417), or without definitive IEI diagnosis (n=306) were excluded from the study, leaving 30,628 patients. IEI were classified using the latest classification of the *International Union of Immunological Societies* (IUIS)(6), with abbreviations of the category names using, *e.g.*, "PIRD", typically referred to as primary immune regulatory diseases, as synonym for "Diseases of immune dysregulation" (category *IV*; *i.e.*, including familial hemophagocytic lymphohistiocytosis, FHL). The statistical analyses were conducted using R (version 4.3.2, Vienna, Austria). Continuous and categorical variables were reported using the medians and interquartile ranges, raw effectives, and percentages, respectively. Survival probabilities were estimated by applying the cumulative incidence function. Overall survival (OS) was defined as the time between birth and death from any cause. Curative therapies, such as allogeneic hematopoietic stem cell transplantation and gene therapy, were considered as competing events. As only patients with an eligible diagnosis are included, they are not considered to be at risk of dying before they are diagnosed(11). The prevalence was calculated based on the European population in 2019 (R package *rnaturalearth* version 1.0.1).

Role of the funding source

The present and former funding entities of ESID (*i.e.*, PPTA Europe [Baxter, Biotest, Grifols, Kedrion, Octapharma], Novartis, UCB, Celltech, GSK, Pharming, Takeda [and corporate predecessors] and Chiesi) had no role in the study design, study conduct, data collection, data management, data analysis, data interpretation, or writing of this report.

Results

On the end date chosen for inclusion, the ESID-R contained data for 30,628 IEI/PID patients from 194 participating centers in 33 European and other, mostly neighboring countries. There was a steady increase in the registration of patients over time (supplementary figure 1) apart from a temporary decline in patient numbers and, to a lesser extent, in center numbers shortly after 2014, due to structural platform changes in 2014. These required centers to verify their patients to improve the accuracy of existing data. This allowed us to derive patient distribution by country and the minimal prevalence of PID/IEI (supplementary table 1 and supplementary figure 2).

Patient characteristics

Figure 1 shows the ages at onset, clinical and genetic diagnoses and main clinical manifestations that led to diagnosis. A steady decline in new diagnoses was observed with age, although a small second peak and plateau were seen in adulthood, and a trend for earlier identification of genetic diagnoses was observed. As reported previously(12), the majority of 80.3% of patients have infections on their way to diagnosis, but only in 61.8% as sole recorded manifestation of IEI, followed by features of immune dysregulation in 11.1% and syndromic manifestations in 7.3% as sole clinical presentation at diagnosis (n=15,360; 2,746; 1,803, respectively; figure 1B). Malignancies were among the first manifestations of IEI in 479 patients, and the only initial manifestation in 117 patients (0.5%; figure 1B).

Table 1 shows the main patient demographic and diagnosis categories, comorbidities such as malignancy, COVID-19 and living status. The interactive supplementary figure 3 shows the patient numbers and their distribution according to their IUIS classification, disease name, and genetic diagnosis (https://esid.org/html-pages/Suppl%20Fig%203_ESID_30k_sunburst_PID.html). About half of all patients suffer from primary antibody deficiencies (PAD; n=15,123), followed in descending order, by combined immunodeficiencies with syndromic features ("syndromic"; n=4,239), phagocytic disorders ("phagocyte"; n=2,548), combined immunodeficiencies (CID; n=2,531), and primary diseases of immune dysregulation (PIRD; n=2,171; table 1 and supplementary figure 3). Table 1 presents the ages at onset, clinical diagnosis, genetic confirmation, and last follow-up, and the diagnostic delay. The proportion of patients reported to have suffered from malignancies was 8.9% (n=1,783). Malignancies were reported as occurring in all subgroups of patients with IEI/PID with moderately varying frequencies, corroborating the notion that IEI/PID are cancer predisposition syndromes. The proportion of patients reported to have been affected by COVID-19 was highest in PAD (35.9%; total cohort: 29.7%). More than half of CID patients received curative therapy (52.6%), followed by those with bone marrow failure syndromes (BMF; 39.8%), PIRD, and phagocyte disorders (30.3 and 25.8%, respectively). Supplementary Table 2 shows the causes of death.

Representation of genetic IEI diagnoses

Figure 2 shows the absolute numbers of patients with a documented genetic diagnosis (n=12,774, 44.5%,) versus those without (n=15,905, 55.5%, including patients not genetically tested) by IEI category. As expected, within the largest patient subgroup (PAD), the genetic diagnosis is lacking for

the majority of patients (figure 2A) and the clinical diagnosis of CVID is attributed, whereas patients with CID with syndromic features had the highest proportion of genetic diagnoses. The top five genes mutated per IEI category are shown in figure 2B, with *IL2RG* being the most frequently reported germline genetic cause of CID; *22q11.2 deletion syndrome,* of CID with syndromic features; *BTK*, of PAD; *TNFRSF6*, of PIRD; *CYBB*, of phagocyte disorders; *STAT1*, of innate immune disorders; *MEFV*, of autoinflammatory syndromes; C1 inhibitor, of complement deficiencies; *RTEL*, of BMF; and somatic *TNFRSF6*, of phenocopies. The proportion of genetic IEI diagnoses represented by patient entries in the ESID-R for the total number of genes known to underlie IEI based on the then available 2022 IUIS classification is shown in figure 2C; the top 50 genetic causes of IEI from the ESID-R are listed in figure 2D.

Treatment modalities according to IEI categories

Immunoglobulin replacement therapy (IGRT), hematopoietic stem cell transplantation (HSCT), gene therapy (GT), and splenectomy are recorded in the ESID-R and listed in supplementary table 3. As expected, the highest number and proportion of IGRT-receiving patients is seen in the subgroup of patients with PAD (figure 3 left- and right-hand panels, respectively). The highest absolute numbers of HSCT procedures were performed in patients with CID, followed by those with PIRD and phagocytic disorders. IGRT and HSCT were documented in patients with IEI/PID in any category. GT, an evolving curative treatment option devoid of some risks associated with HSCT, such as alloreactivity, was recorded in a descending order for CID, syndromic and phagocyte disorders (figure 3). Splenectomy was recorded relatively frequently in patients with phenocopies, but it was also documented for any IEI category. The use of immune-modulatory treatments such as anti-inflammatory, cell-depleting, or pathway-directed (targeted) therapies and the rare cases of thymus or solid organ transplantation recorded are not shown here, as the heterogeneity of the data exceeds the scope of this general report.

Survival probabilities of patients with IEI

We hypothesized that the survival probabilities of patients in the 10 IUIS categories of IEI differed due to their varying predispositions to life-threatening infections, malignancies, autoimmunity, or other manifestations and complications associated with their underlying conditions. Although the data granularity in the entire ESID-R with respect to patient follow-up intervals is not comparable with that in disease-specific prospective cohort studies, we could plot the reversed cumulative incidence function, due to the presence of competing risks, based on the relatively large patient numbers in each IEI category (figure 4). Of note, natural biases such as underreporting of patients who died from IEI before diagnosis or of patients with a mild phenotype exist, and numbers at risk increased over the first few years of the observation period (0–97.9 years of age) due to the later time points of inclusion or diagnosis. We detected a steep early decline in survival in many IE categories while curves plateaued, *i*, methodologically, in some where definitive treatments exist, or, *ii*, in genotypes with variable penetrance (*e.g.*, CID, PIRD). Additionally, while patients with PAD showed a continuous decline in survival probability from a young age across all age groups, the decline in the survival of patients with phagocyte and innate immune disorders showed an initial drop. This finding suggests that a proportion of patients are at very high risk during their first five

years of life (see also supplementary figure 4: curves for ages 0–5 years). Diagnoses of early
deceased patients with PIRD (n=70 under 5 years of age) were mostly due to disorders with a high
risk of hemophagocytic lymphohistiocytosis (HLH; 81.4%); premature deaths in "innate" IEI were
frequently due to IRAK4 or MyD88 deficiencies (supplementary table 4). Patients with IEI/PID with
syndromic features had a triphasic survival probability. After observing an initial decline in the first
four years of life, we detected a second pronounced decline in survival probability in the patients'
2^{nd} and 3^{rd} decades of life, most likely due to the increased risk of malignancies in many patients in
this subgroup (>25%, see supplementary table 2), and a third, relatively steep decline in patients in
their 6^{th}–7^{th} decades.

The ESID-R in the international registry landscape

Other continental or cross-regional IEI/PID registries of varying temporal and geographical depth
include the global *Jeffrey Modell Foundation* Centers Network (>94,000 patients), the *USIDNET*
registry in the USA (>5,000), the *LASID* registry in Latin America (>9,000), the Australian *AusPIPS*
registry (>1,500) as part of the Australasian network, *JSIAD* (>1,200) in Japan as part of the Asian-
Pacific (APSID) network, the Canadian registry (*CIEINR*) founded in 2024 and the registries of the
Primary Immune Deficiency Treatment Consortium (PIDTC) of North America; in addition, many
national registries exist inside or outside of ESID(13). Those reported in Europe are shown in
supplementary figure 5 and respective centers in supplementary table 5.

Scientific impact of the ESID-R and sub-studies

Up to the end date chosen for data inclusion in this manuscript, 84 peer-reviewed publications
resulted from projects deriving data directly from the ESID-R (Figure 5 and ESID website(14)). We
evaluated this scientific output by categorizing and counting the publications and their citations as
follows: disease-specific natural history studies (*n*=25; 3,658 citations), country-specific
epidemiological studies (*n*=19; 2,575 citations), six reviews (422 citations), two on technical aspects
(104 citations) and 15 of our own registry-conducted studies (*e.g.*, on first manifestations or on
working definitions for the clinical diagnosis of IEI/PID, 1,221 citations), plus 17 studies that could
have been assigned to multiple or overlapping categories (figure 5), resulting in a mean citation rate
of 95 per publication.

Discussion

The ESID-R is a large and growing real-world database enabling powerful analyses that have continuously generated epidemiological and disease-specific observational results since 1994. It contributes to knowledge and improvements in patient care in IEI in Europe and beyond. The present analysis illustrates the current distribution of: *i)* IEI diagnoses recorded in contributing centers and countries, *ii)* major treatment modalities between all IEI categories, *iii)*, ages at onset, at clinical and at genetic diagnosis and *iv)* survival probabilities. Our results underline the recommendation to implement newborn screening, potentially even for certain additional IEI other than severe combined immunodeficiencies (SCID). The 7,980 citations of 84 peer-reviewed ESID-R-based publications as of early 2024 demonstrate the success of the ESID-R as platform for research sub-studies and their substantial scientific impact. Hence, the ESID-R provides a clear example of how data collection and collaboration can benefit patients with rare diseases.

Survival probabilities, early diagnosis, and newborn screening

Newborn screening (NBS) for SCID by measuring T-cell receptor with or without kappa-deleting recombination excision circles in dried blood spots has been implemented in many countries around the world[15-17]. A substantial survival benefit for patients diagnosed and treated by early HSCT was demonstrated and recently confirmed in a longitudinal study of the PIDTC[18]. The steep early decline in survival we observed in patients with disorders of immune regulation and "innate" IEI may argue for an extension of NBS to other immediately life-threatening IEI (*e.g.*, FHL, XLP1, MyD88/IRAK4 deficiencies and IPEX syndromes). To extend the spectrum of early IEI–NBS, current techniques may be supplemented by RNA sequencing or germline genetic testing, such as targeted sequencing panels, whole exome or genome sequencing. Feasibility studies of such extended NBS are ongoing in regional or national programs[19]. However, the vast ethical and social implications of any omics-based addition like baby genome screening will require careful consideration of the risks and benefits[20].

Methodological and organizational limitations

Physicians and other documentarists regularly collect data for the ESID-R on a voluntary basis, with data quality and quantity (*i.e.*, data depth and accuracy) varying substantially between centers and across countries. While all centers can participate in the ESID-R and scientific sub-studies for free, minimum infrastructure is required. With the rare exception of per-patient fees in pharma-sponsored level-3 studies, ESID does not financially reimburse data entry. The participants mainly benefit from the international academic collaboration and representation, which increases their awareness for disease phenotypes and current clinical research questions. Also, each center may obtain their own (local and regional) epidemiological data within a legally and ethically approved and financially sustainable technical framework (see survey results in the supplementary material). Furthermore, ESID has no means of monitoring the data quality other than by inbuilt checks for logic and completeness at the time point of data entry. So far, automated transfers of data from electronic health records (eHR) into the ESID-R have not been attempted due to regulatory obstacles and the heterogeneity of eHRs in the 194 participating centers. All the above, along with the fact

328 that some monogenic IEIs have only been described recently while others have been recognized for
329 decades, introduces a systemic ascertainment bias into the dataset when comparing, *e.g.*, numbers
330 of diagnoses and time of survival. For example, X-linked agammaglobulinemia (caused by mutations
331 in *BTK*)(21) is probably not eight times more prevalent than, *e.g.*, the autosomal-dominant CTLA4-
332 (haplo)insufficiency(22 23) or NFKB1-(haplo)insufficiency(24).

Challenges of reporting genetic data

Collecting data from the heterogenous patient population of individuals with IEI/PID and storing
these in a registry presents many challenges. Scientists estimate that current routine next-
generation sequencing (NGS) technologies can reveal a molecular cause in approximately 30–35% of
IEI cases(19)*.* It is noteworthy that information classifying detected variants by applying ACMG
criteria or performing functional validation is not collected in the ESID-R. The incorporation of
genetic data beyond known pathogenic mutations, ideally linked to precisely defined phenotypes
according to the human phenotype ontology (HPO)(25), represents a biologically interesting, future
challenge. This includes the verification of variants of unknown significance, intronic changes,
somatic mutations and mosaicism, epigenetic factors, variable penetrance and other less-studied
factors as non-coding modifier alleles and monoallelic expression(26).

Link to other registries, synergies, overlapping efforts

From a scientific perspective, it has become increasingly important to compare outcomes for IEI/PID
patients with and without cellular therapy (HSCT or gene therapy). The registry of the *European*
Society for Blood and Marrow Transplantation (EBMT) currently registers definitive cellular therapies
in about 700 patients with IEI per year(27), many of whom are also registered in the ESID registry.
Hence, to optimize future research, datasets from the ESID and the EBMT registries for the same
patients should be combined. Better alignment of the registries and their data fields is urgently
needed to facilitate such studies. Furthermore, the IUIS classification has gradually incorporated
more diseases from overlapping specialties (*e.g.,* hematology-oncology, rheumatology,
gastroenterology), many of which are (also) covered by other medical societies and patient
registries. Thus, although data from many of these patients are stored in the ESID-R, their proportion
in the ESID-R does not reflect the real-world distribution. The registry of the *European Reference*
Network for Rare Immunological and Autoimmune Diseases (ERN-RITA), an EU sponsored network of
healthcare providers from reference centers for PID/IEI, rheumatology, autoinflammatory and
autoimmune diseases, aims to collect common data elements from patients across these disease
areas. This initiative may help to avoid redundancy. Finally, using independent (meta-)identifiers
such as the *European SPIDER-ID(28)* in all registries would help scientists disentangle the registry
landscape.

Future: AI in registry work for data analysis and interpretation

363 Advanced software tools and artificial intelligence (AI, *e.g.*, large language models, natural language
364 processing, and machine learning), will certainly transform clinical decision-making and other crucial
365 processes in medicine including the field of IEI.(29 30) Automated eHR scanning and data harvesting

processes, ideally for specific terms in accordance with HPO, medical reports, disease classification codes (*e.g.*, ICD-11 or ORPHA codes), but also for free text may be used to identify currently undiagnosed IEI/PID patients. These individuals might benefit from early screening by preventing complications. Accordingly, as a first step, data from an IEI/PID registry like the ESID-R that includes data from patients with an established diagnosis could be used to train AI models and to predict a monogenic IEI or at least the most likely affected pathway in patients who lack a genetic diagnosis. Consequently, patients may undergo screening for disease-specific risks and receive appropriate therapy early in their course. Complementing the ESID-R with AI-based tools has the potential to transform this data-collection platform to an electronic IEI/PID patient management assistant in the future.

Conclusions and Perspectives

The rarity and complexity of many IEI grants them an orphan status regarding pharmaceutical research and development. The feasibility of clinical research, including drug trials and post-authorization efficacy and safety studies, thus depends on large networks of academic institutions and medical specialist societies. As one of the largest registries, containing extensive longitudinal datasets and having a pan-European reach, the ESID-R is likely to remain one of the most relevant scientific registries for patients with IEI.

Acknowledgements

The authors would like to thank all contributors and supporters not already mentioned among the collaborator list of the ESID-R working party (*e.g.*, documentarists, study nurses, data analysts) for their continuous support. Scientific language editing was performed by Dr. Sara Crockett, Graz, Austria (www.saras-science.com). MGS was in part funded by the Styrian Children's Cancer Aid Foundation (Steirische Kinderkrebshilfe). MHA represents the PID stream of the European Reference Network for Rare Immunological and Autoimmune Diseases (ERN-RITA). CEREDIH thank colleagues from Université de Paris, Imagine Institute, Data Science Platform, INSERM UMR 1163, F-75015, Paris, France for their contribution in the interface between ESID-R and the CEREDIH registry. LDN is supported by the Division of Intramural Research, NIAID, NIH. IM is a senior clinical researcher at FWO Vlaanderen.

Conflict of interest statement

MGS received advisory board honoraria from Pharming. MHA received research support, consulting fees, and speaker honoraria from Octapharma. GK, FC declare no conflict of interest. LRD KU Leuven receives research funding and advisory board honoraria for IM from CSL-Behring and Boehringer-Ingelheim. BS has received travel grants from Pharming, and speaker honoraria from Takeda.

Contributors' statement

MGS designed the study; MGS and GK wrote the manuscript with contributions from MHA, BG, SG, CS, SE, NM, FC, JLC, LDN, CIES; CS, SS and MP collected data about the global registry landscape; AG performed literature search and analyzed ESID-R-related publication records; MA, MGS and GK analyzed the data and prepared the figures. These and all other authors contributed in varying proportions to the technical, organizational, or structural basis of ESID and the ESID-R and to the collection of patient data. All co-authors contributed to and edited drafts of the original manuscript and figures, and approved the final version.

List of collaborators (ESID registry working party)

Shahrzad; Bancé, Renate; Bangs, Catherine; Barfusz, Katerina; Baris, Safa; Barlan Isil, B; Bartsch, Michaela;
Baumann, Helge; Baumann, Ulrich; Baumeister, Veronika; Baxendale, Helen; Bazen, Suzanne; Beaurain,
Beatrice; Beauté, Julien; Bechar-Makhloufi, Mounia; Beck, Norbert; Becker, Brigitta; Becker, Christian;
Behrends, Uta; Beider, Renata; Beier, Rita; Belkacem, Amel; Belke, Luisa; Bellert, Sven; Belohradsky, Bernd H.;
Ben Slama, Lilia; Ben-Bouzid, Aouatef; Benoît, Vincent; Berdous-Sahed, Thamila; Bergils, Jan; Berglöf, Anna;
Bergman, Peter; Bernat-Sitarz, Katarzyna; Bernatoniene, Jolanta; Bernatowska, Ewa; Bernbeck, Benedikt;
Bethune, Claire; Beuckmann, Kai; Bhole, Malini; Biegner, Anika-Kerstin; Bielack, Stefan; Bienemann, Kirsten;
Bigl, Arndt; Bigorgne, Amélie; Bijl, Marc; Binder, Nadine; Bitzenhofer-Grüber, Michaela; Blanchard Rohner,
Geraldine; Blank, Dagmar; Blattmann, Claudia; Blau, Julia; Blaziene, Audra; Blazina, Stefan; Blume, Roswitha;
Boardman, Barbara; Bode, Sebastian; Boelens, Jaap-Jan; Boesecke, Christoph; Bogaert, Delfien; Bogner,
Johannes; Bohynikova, Nadezda; Booth, Claire; Bordon, Victoria; Borkhardt, Arndt;Börries, Melanie; Borte,
Michael; Borte, Stephan; Bossaller, Lukas; Bossard, Madeleine; Botros, Jeannet; Boucherit, Soraya; Boutros,
Jeannette; Boyarchuk, Oksana; Boyman, Onur; Boztug, Kaan; Braschler, Thomas; Bravo, Sophie; Bredius,
Robbert; Bright, Philip; Brito de Azevedo Amaral, Carolina; Brodszki, Nicholas; Brodt, Grit; Brolund, Allan;
Brosselin, Pauline; Brummel, Bastian; Brun-Schmid, Sonja; Brunner, Jürgen; Bruns, Roswitha; Buchta, Christina;
Buck, Dietke; Bücker, Aileen; Bührlen, Martina; Burdach, Stefan; Burns, Siobhan; Burton, Janet; Caminal, Luis;
Cancrini, Caterina; Canessa, Clementina; Cantoni, Nathan; Capilna, Brindusa; Caracseghi, Fabiola; Caragol,
Isabel; Carbone, Javier; Carrabba, Maria; Chamberlain, Latanya;Chandra, Anita; Chantrain, Christophe; Chapel,
Helen; Chassot, Julie; Chee, Ronnie; Chinello, Matteo; Chopra, Charu; Chovancova, Zita; Christmann, Martin;
Chrzanowska, Krystyna; Ciznar, Peter; Claes, Karlien; Classen, Carl Friedrich; Classen, Martin; Cochino, Alexis-
Virgil;Condliffe, Alison M.; Corbacioglu, Selim; Cordeiro, Ana Isabel; Cordier Wynar, Donatienne; Core, Claire;
Costes, Laurence; Coulter, Tanya; Courteille, Virginie; Cristina, Maria; Cucuruz, Maria; Dabrowska, Leonik Nel;
Dähling, Mandy; Daly, Mary Louise; Daniel, Claudia-Sabrina; Danieli, Maria Giovanna; Darroch, James; Davies,
Graham; de Baets, Frans; De Boeck, Christiane; De Gracia Roldan, Javier; de Nadai, Narimene; de Schutter, Iris;
De Vergnes, Nathalie; de Vries, Esther; de Witte, Josine ; Debert, Theo; Defila, Corina; Deimel, Judith;
Delaplace, Diane; Dellepiane, Rosa Maria; Dellert, Nelli; Delliera, Laura; Delor, Anita; Demel, Ulrike; Dempster,
John; den Os, M.M.; Dengg, Rosmarie; Desisa, Sora Asfaw; Desta, Alexandra; Detkova, Drahomira; Dewerchin,
Maite; Dieli Crimi, Romina; Dilloo, Dagmar; Dimitriou, Florentia; Dinser, Jasmin; Dipani, Nabila; Dirks, Johannes;
Dittrich, Anna-Maria; Djermane, Lylia; Dogru, Yagmur; Dogu Esin, Figen; Dombrowski, Angelika; Döring,
Michaela; Drabe, Camilla Heldbjerg; Drerup, Susann; Drexel, Barbara; Driessen, G.J.A.; Dückers, Gregor;
Dudoit, Yasmine; Duppenthaler, Andrea; Ebetsberger-Dachs, Georg; Ecser, Mate Barnabas; Eekman, Maartje;
Ehrat, Rosanna; Eisl, Eva; Ekwall, Olov; El Hawary, Rabab; El-Helou, Sabine M.; El-Marsafy, Aisha; Elbe, Sarina;
Elcombe, Suzanne; Eldash, Alia; Eldeen, Youssif Alkady Radwa Salah; Ellerbroek, P.M.; Elling, Roland; Elliott,
Jane; Engelhardt, Angelika; Ernst, Diana; Ersoy, Fügen; Esper, Stefanie; Esteves, Isabel; Exley, Andrew; Faber,
Martin; Fabio, Giovanna; Fahrni, Gaby; Faletti, Laura Eva;Faletti, Laura Eva; Farber, Claire-Michele; Farela
Neves, João ; Faria, Emilia; Farkas, Henriette; Farmaki, Evangelia; Faßhauer, Maria; Fasth, Anders; Fätkenheuer,
Gerd; Faustmann, Stefanie; Fecker, Gisela; Feighery, Conleth; Feiterna-Sperling, Cornelia; Fernandez-Cruz,
Perez Eduardo; Ferreira Concalo, Cordeiro; Ferster, Alice; Feuchtinger, Tobias; Feyen, Oliver; Finke, Daniela;
Fitter, Sigrid; Flaschberger, Stefan; Fleckenstein, Lucia; Föll, Dirk; Fontana, Adriano; Forino, Concetta; Förster-
Waldl, Elisabeth; Franchet, Nora; Freitag, Dagmar; Frey, Urs P.; Friedel, Elisabeth; Friedrich, Wilhelm; Frisch,
Barbara; Frischknecht, Lukas; Fritzemeyer, Stephanie; Gagro, Alenka; Gahr, Manfred; Gambineri, Eleonora;
Gamper, Agnes; Gams, Franziska; Ganzow, Astrid; Garcelon, Nicolas; Garcez, Tomaz; Garcia Prat, Marina;
Gardulf, Ann; Garibay, Janine; Garwer, Birgit; Gathmann, Jonathan; Gebauer, Corinna; Geberzahn, Linda;
Geikowski, Tilman; Geisen, Ulf; Gemander, Christiane; Gennery, Andrew R.; Gerisch, Marie; Gernert, Michael;
Gerrer, Katrin; Gerschmann, Stev; Giannini, Carolin; Gil Herrera, Juana; Girndt, Matthias; Girrbach, Ramona;
Girschick, Hermann; Gkougkourelas, Ioannis; Gładysz, Dominika; Gnodtke, Elisabeth; Goda, Vera; Goddard,
Sarah; Goebel, Daniela; Goffard, Jean-Christophe; Goldacker, Sigune; Gollowitsch, Eva Maria; Gomes,
Manuella; Gompels, Mark; González, Míriam; Gonzalez Granado, Luis Ignacio; Gordins, Pavels; Gößling,
Katharina; Göschl, Lisa; Gossens, Lucy; Gowin, Ewelina; Graafen, Lea; Graca, Leo; Gradauskiene (Sitkauskiene),

Brigita; Graf, Dagmar; Graf, Norbert; Grange, Elliot; Grashoff, H.Anne; Greil, Johann; Grigoriadou, Sofia;
Gronlund, Helen; Groß-Wieltsch, Ute; Guerra, Teresa; Guevara-Hoyer, Kissy; Gueye Mor, Seny; Guibert,
Noemie; Gülnur, Birgit; Güngör, Tayfun; Guseva, Marina; Guzman, David; Haag, Marcel; Haase, Gabriele;
Haenicke, Henriette; Haerynck, Filomeen; Hafsa, Ines; Hagin, David; Hallek, Michael; Hancioglu, Gonca;
Handgretinger, Rupert; Hanitsch, Leif G.; Hansen, Susanne; Hariyan, Tetyana; Harrer, Thomas; Hassunah, Pia;
Hatzistilianou, Maria; Hauck, Fabian; Hauser, Thomas; Haverkamp, Margje H.; Hayman, Grant; Heath, Paul;
Hedrich, Christian; Heeg, Maximilian; Heike, Michael; Heimbrodt, Martin; Heine, Sabine; Heininger, Ulrich;
Heinrich, Christian; Heinz, Valerie; Heitger, Andreas; Helbert, Matthew; Helbling, Arthur; Hellige, Antje;
Hempel, Julya; Henderson, Karen; Henes, Jörg; Henneke, Philipp; Hennig, Christian; Henrichs, Karin; Herbst,
Martin; Hermann, Walter; Hernandez, Manuel; Hernández, Anja; Heropolitanska, Edyta; Herriot, Richard;
Herrmann, Friedrich; Herwadkar, Archana; Hess, Christoph; Hess, Ursula; Hesse, Sebastian; Higgins, Sonja;
Hilfanova, Anna; Hilpert, Sophie; Hintze, Chantal; Hlaváčková, Eva; Hodl, Isabel; Hodzic, Adna; Hoernes,
Miriam; Hoffmann, Christina; Holbro, Andreas; Höllinger, Christiane; Holtsch, Lisa; Holzer, Ursula; Holzinger,
Dirk; Hönig, Manfred; Hönscheid, Andrea; Horn, Julia; Horneff, Gerd; Hoyoux, Claire; Hristova, Nataliya; Hübel,
Kai; Hübner, Angela; Huemer, Christian; Huissoon, Aarnoud; Hülsmann, Brigitte; Hundsdörfer, Patrick;
Huppertz, Hans-Iko; Huß, Kristina; Hussain, Sadia; Husson, Julien; Hüttner-Foehlisch, Tanja; Ijspeert, Hanna;
Ikinciogullari, Aydan; İlknur, Kökçü; Irga, Ninela; Jablonka, Alexandra; Jahnz-Rozyk, Karina; Jakob, Marcus;
Jakoby, Donate; Jandus, Peter; Jansson, Annette; Jaquet, Melanie; Jardefors, Helene; Jargulinska, Edyta; Jauk,
Barbara; Jesenak, Milos; Jilka, Katharina; Jolles, Stephen; Jones, Alison; Jones, Regina; Jonkman-Berk, Birgit;
Jönsson, Göran; Joyce, Hilary J.; Juliana, Pricillia; Kabesch, Michael; Kager, Leo; Kahlert, Christian; Kaiser-
Labusch, Petra; Kakkas, Ioannis; Kamitz, Dirk; Kanariou, Maria; Kanz, Lothar; Karakoc-Aydiner, Elif; Karanovic,
Boris; Kartal-Kaess, Mutlu; Käser, Elisabeth; Katzenstein, Terese L.; Kayserova, Hana; Kelleher, Peter; Kerre,
Tessa; Kilic, Sara Sebnem; Kirchner, Martina; Kiwit, Simone; Kiykim, Ayca; Klasen, Jessica; Klaudel-Dreszler,
Maja; Klein, Ariane; Klein, Christoph; Klein-Franke, Andreas; Kleine, Ilona; Kleinert, Stefan; Klemann, Christian;
Klima, Marion; Kobbe, Robin; Kocacik Uygun, Dilara Fatma; Koch, Melanie; Kochler, Yvonne; Kohistani, Naschla;
Kojic, Marina; Kolios, Antonio; Kölsch, Uwe; Koltan, Sylwia; Kondratenko, Irina; Königs, Christoph;
Konoplyannikova, Julia; Kopac, Peter; Kopp, Jana; Körholz, Dieter; Körholz, Julia; Korte, Pauline; Kostyuchenko,
Larysa; Kötter, Ina; Kracker, Sven; Králícková, Pavlina; Kramm, Christof; Kramme, Philipp;Krausz, Máté; Kreuz,
Wolfhart; Kriván, Gergely; Kropshofer, Gabriele; Krüger, Renate; Krystufkova, Olga; Ktistaki, Maria; Kühl, Jörn-
Sven; Kühn, Alexander; Kuijpers, Taco W.; Kuis, Wietse; Kullmann, Silke; Kulozik, Andreas; Kumararatne,
Dinakantha; Kümmler, Ria; Kündgen, Andrea; Kurenko – Deptuch, Magdalena; Kuss Paula, Cosima; Kütükcüler,
Necil; Lafoix-Mignot, Cécile; Lamers, Beate; Lanbeck, Peter; Landais, Paul; Landwehr-Kenzel, Sybille;
Langemeyer, Vanessa; Langer, Thorsten; Lankisch, Petra; Lanz, Nadia; Lara, Manrique de; Lara-Villacanas,
Eusebia; Laubenthal, Lisa; Laws, Hans-Jürgen; Le Mignot, Loic; Leahy, Ronan; Lee, Jae-Yun; Lehmann, Andrea;
Lehmberg, Kai; Lehner, Patricia; Leibfrit, Hans; Leistner, Leoni; Lesch, Petra; Leutner, Simon; Liatsis, Manolis;
Libai Véghová, Linda; Liebel, Johanna; Liese, Johannes G.; Linauskiene, Kotryna; Linde, Richard; Linßner,
Martina; Lippert, Conrad Ferdinand; Litzman, Jiri; Llobet, Pilar; Lo, Babacar; Lodin, Tariq; Lokaj, Jindrich;
Longhino, David; Longhurst, Hilary; Lopes da Silva, Susana; Lorenzen, Catharina; Lougaris, Vassilios; Löw, Doris;
Lubatschofski, Annelie; Lucas, Mary; Lutz-Wiegers, Verena; Maaß, Sabine; Maccari, Maria Elena; Macura-
Biegun, Anna; Maerz, Vanessa; Maggina, Paraskevi; Mahrenholz, Hannah; Maier, Sarah; Malfroot, Anne;
Malinauskiene, Laura; Mannhardt-Laakmann, Wilma; Manson, Ania; Mantkowski, Felicia; Manzey, Petra;
Marasco, Carolina; Marcus-Mandelblit, Nufar; Marg, Wolfgang; Markelj, Gasper; Marodi, Laszlo; Marques, José
Goncalo; Marques, Laura; Marschall, Karin; Martínez, Natalia; Martinez de la Ossa Saenz-Lopez, Rafael;
Martinez-Saguer, Inmaculada; Martire, Baldassarre; Marzollo, Antonio; Masekela, Refiloe; Masjosthusmann,
Katja; Masmas, Tania Nicole; Matamoros, Nuria; Mattern, Jutta; Mau-Asam, Pearl; McDermott, Elizabeth;
McIntosh, Nichole; Meglic, Karmen Mesko; Meijer, Ruben; Meinhardt, Andrea; Meshaal, Safa; Messaoud,
Yasmina; Meyer, Björn; Meyer-Olson, Dirk; Micol, Romain; Micoloc, Bozena; Mielke, Gudrun; Milito, Cinzia;
Misbah, Siraj; Mohr, Michael; Mohrmann, Karina; Moin, Mostafa; Molinos, Luis; Möller, Jana; Möller-Nehring,
Sarah; Morbach, Henner; Moschese, Viviana; Moser, Olga; Moshous, Despina; Motkowski, Radoslaw;

Motwani, Jayashree; Mouftah Galal, Nermeen; Müglich, Carmen; Mukhina, Anna; Muller, Eva; Müller,
Christiane; Müller, Gabriele; Müller, Hedi; Müller, Ingo; Müller, Thomas; Müller, Zoe; Müller-Ladner, Ulf;
Müller-Stöver, Sarah; Münstermann, Esther; Murtra Garrell, Núria; Muschaweck, Moritz; Mutert, Miriam;
Nademi, Zohreh; Naik, Paru; Näke, Andrea; Nalda, Andrea Martin; Nasrullayeva, Gulnara; Naumann-Bartsch,
Nora; Nemitz, Verena;Neth, Olaf; Neubauer, Andreas; Neubert, Jennifer; Neumann, Carla; Niehues, Tim;
Niemuth, Mara; Nieuwhof, Chris; Nolkemper, Daniela; Noorani, Sadia; Noorlander, Budde Adya; Notheis,
Gundula; Nowatsh, Sanam Amelie; Obenga, Gaelle; Ocak, Suheyla; Oker, Mehmet; Olbrich, Peter; Olipra, Anna;
Omran, Heymut; Oommen, Prasad; Orosova, Jaroslava; Oskarsdottir, Solveig; Özsahin, Hülya; Pac, Malgorzata;
Pachlopnik Schmid, Jana; Pandolfi, Franco; Papadopoulou-Alataki, Efimia; Papatriantafillou-Schmieder, Anna;
Parra-Martinez, Alba; Paschenko, Olga; Pašić, Srdjan; Pasnik, Jarek; Patel, Smita; Pavlík, Martin; Paz Artal,
Estela; Peeters, Anouk; Pereira da Silva, Sara Branco; Perez-Becker, Ruy; Perlhagen, Markus; Peter, Hans-
Hartmut; Pfreundschuh, Michael; Philippet, Pierre; Picard, Capucine; Pietrucha, Barbara; Pietzsch, Leonora;
Piquer Gibert, Monica; Pirolt, Kerstin; Plebani, Alessandro; Pleguezuelo, Daniel E.; Pollok, Katrin;
Pommerening, Helena; Popihn, Daniela; Poplonek, Aleksandra; Popp, Marina; Porta, Fulvio; Portegys, Jan;
Posfay-Barbe, Klara; Potjewijd, Judith; Poulheim, Sebastian; Prader, Seraina; Prämassing-Scherzer, Petra;
Prelog, Martina; Prevot, Johan; Price, Arthur; Price, Timothy; Proesmans, Marijke; Provot, Johan; Quinti,
Isabella; Raab, Anna; Rack, Anita; Raffac, Stefan; Ramos Oviedo, Eduardo; Randrianomenjanahary, Philippe;
Ranohavimparany, Anja; Raptaki, Maria; Rashidzadeh, Roonaka; Rathwallner, Margit; Reda, Shereen;
Redouane, Nahida; Regateiro, Frederico S.; Reichenbach, Janine; Reimers, Bianca; Reinhardt, Cornelia;
Reinhardt, Dirk; Reinprecht, Anne; Reiß, Tamara; Reisli, Ismail; Renner, Eleonore; Rezaei, Nima; Richter, Alex;
Richter, Darko; Rieber, Nikolaus Peter; Rieckehr, Nadja; Riedel, Marion; Riescher, Heidi; Rischewski, Johannes;
Ristl, Nicole; Ritterbusch, Henrike; Ritz, Tanja; Rivier, Francois; Rockstroh, Jürgen K.; Roesler, Joachim; Rofiah,
Himatur; Rogerson, Elizabeth; Rolfes, Elisabeth; Roller, Beate; Romanyshyn, Yaryna; Rondelli, Roberto;
Roosens, Fien; Rösen-Wolff, Angela; Rösler, Valentina; Roth, Johannes; Rothoeft, Tobias; Roubertie, Agathe;
Rübsam, Gesa; Rutgers, Abraham; Ryan, Paul; Sach, Gudrun; Sadeghi, Kambis; Sahrbacher, Ulrike; Saidi,
Angelika; Sanal Tezcan, Özden; Sanchez-Ramon, Silvia; Santos, Juan Luis; Sargur, Ravishankar; Savchak, Ihor;
Savic, Sinisa; Schaaf, Bernhard; Schaefer, Marzena; Schäfe, Christina; Scharbatke, Eva; Schatorje, Ellen;
Schauer, Uwe; Scheibenbogen, Carmen; Scheible, Raphael; Scheinecker, Clemens; Schiller, Romana; Schilling,
Beatrice; Schilling, Freimut; Schlieben, Steffi; Schmalbach, Thilo; Schmalzing, Marc Thomas; Schmid, Pirmin;
Schmidt, Nadine; Schmidt, Reinhold Ernst; Schmitz, Monika; Schneider, Dominik; Schneider, Dominik T.;
Schneppenheim, Reinhard; Scholtes, Cathy; Schölvinck, E.H.; Schönberger, Stefan; Schreiber, Stefan; Schrijvers,
Rik; Schroll, Andrea; Schruhl, Simone; Schrum, Johanna; Schubert, Ralf; Schuh, Sebastian; Schulz, Ansgar;
Schulz, Claudia; Schulze, Ilka; Schulze-Koops, Hendrik; Schulze-Sturm, Ulf; Schumacher, Eva-Maria; Schürmann,
Elvira; Schürmann, Gesine; Schuster, Volker; Schwaneck, Eva; Schwarz, Klaus; Schwarz, Tobias; Schwarze-
Zander, Carolynne; Schweigerer, Lothar; Schwinger, Wolfgang; Sediva, Anna; Seebach, Jörg; Seger, Reinhard;
Segerer, Florian; Selle, Barbara; Seneviratne, Suranjith; Shabanaj, Hatidje; Shcherbina, Anna; Siepelmeyer,
Anne; Siepermann, Kathrin; Simon, Anna; Simon, Arne; Simon-Klingenstein, Katja; Simonovic, Marija; Simsen-
Baratault, Merlin;Sindram, Elena; Skapenko, Alla; Skarke, Maiken; Skomska-Pawliszak, Malgorzata; Slatter,
Mary; Smet, Julie; Sobh, Ali; Sobik, Bettina; Sogkas, Georgios; Sohm, Michael; Solanich-Moreno, Xavier; Soler
Palacín, Pere; Sollinger, Franz; Somech, Raz; Sonnenschein, Anja; Soresina, Annarosa; Sornsakrin, Marijke;
Soura, Stavrieta; Spaccarotella, Sabrina; Spadaro, Guiseppe; Sparber-Sauer, Monika; Specker, Christof;
Speckmann, Carsten; Speidel, Lisa; Speletas, Matthaios; Stachel, Klaus-Daniel; Stadon, Catherine; Stanislas,
Aurélie; Stapornwongkul, Cynthia; Staus, Paulina; Steck, Regina; Steele, Cathal; Steffin, Herbert; Steiner, Urs;
Steinmann, Sandra; Stevens, Wim; Stiefel, Martina; Stieger, Sarah; Stiehler, Sophie; Stimm, Hermann; Stojanov,
Silvia; Stoll, Matthias; Stoppa-Lyonnet, Dominique; Strapatsas, Tobias; Strauß, Gabriele; Streiter, Monika; Strik-
Albers, Riet; Strotmann, Gaby; Suárez Casado, Héctor; Subiza, Jose Luis; Sundin, Mikael; Süß, Birgit; Sutter,
Fabienne; Szaflarska, Anna; Szemkus, Monika; Tamary, Hannah; Tantou, Sofia; Tarzi, Michael D.; Taschner,
Helga; Tedgard, Ulf; Teixeira, Carla; ten Berge, RJM; Tenbrock, Klaus; Tester, Sabine; Tezcan, Ilhan; Thalguter,
Sonja; Thalhammer, Julian; Thoma, Katharina; Thomas, Moira; Thomczyk, Fabian; Thon, Vojtech; Thrasher,

Adrian; Tierney, Patricia; Tietsch, Nadine; Tommasini, Alberto; Tönnes, Beate; Tony, Hans-Peter; Trachana,
Maria; Trapp, Carmen; Tricas, Lourdes; Trindade Neves, Maria Conceicao; Trischler, Jordis; Ubieto, Hugo;
Uelzen, Anett; Uhlmann, Annette; Ullrich, Jan; Ullrich, Kurt; Ünal, Ekrem; Urbanski, Gerhard; Urschel, Simon;
Uszynska, Aleksandra; Vacca, Angelo; Vaganov, N.N.; Vagedes, Daniel; Valicevic, Stefanie; Vallelian, Florence;
van Beem, Rachel T; van Damme, Charlotte; van de Ven, Annick; van den Berg, J. Merlijn; van der Flier, Michiel;
van Dissel, J.T.; van Hagen, P.M.; van Montfrans, J.M.; van Ogtrop, Geoffrey; van Rens, Jacqui; van Riel, Christel
A.M.P.; van Royen-Kerkhof, Annet; van Well, G.Th.J.; Vasiliki, Antari; Velbri, Sirje; Vencken, Jo; Ventura,
Alessandro; Vermeulen, François; Vermylen, Christiane; Viemann, Dorothee; Viereck, Anja; Villa, Anna; Vincke,
Jeroen; Vinnemeier-Laubenthal, Lisa; Vo Thi, Kim Duy; Vollbach, Kristina; Volokha, Alla; von Bernuth, Horst;
von Bismarck, Philipp; Voss, Rebecca; Voß, Sandra; Vural, Yüksel; Wachuga, Heike; Wagner, Norbert;
Wagström, Per; Wahle, Matthias; Wahn, Volker; Wapp, Nadine; Warnatz, Klaus; Warneke, Monika; Warris,
Adilia; Wasmuth, Jan-Christian; Wasserfallen, Jean-Blaise; Wawer, Angela; Weber, Manfred; Weemaes, Corrie
MR;Wehrle, Julius; Weidinger, Stephan; Weiß, Michael; Weißbarth-Riedel, Elisabeth; Werner, Antje; Wessel,
Sara; Westkemper, Marco; Wicher, Monika; Wickmann, Lutz; Wiegert, Sabine; Wiehe, Monique; Wiehler,
Katharina; Wiesböck, Lydia; Wiesik-Szewczyk, Ewa; Williams, Anthony; Winkler, Beate; Winkler, Christel;
Winkler, Martina; Winkler, Melanie; Wintergerst, Uwe; Wisgrill, Lukas; Witte, Torsten; Wittkowski, Helmut;
Wolf, Barbara; Wölke, Sandra; Wolschner, Christina; Wolska-Kusnierz, Beata; Wood, Philip; Workman, Sarita;
Worth, Austen; Wortmann, Michaela; Wuillemin, Walter Alfred; Wulffraat, Nico M; Wustrau, Katharina;
Wyndham-Thomas, Chloé; Yegin, Olcay; Yildiran, Alisan; Yilmaz, Denise; Young, Patrick; Yucel, Esra; Zečević,
Milica; Zepp, Fred; Zetzsche, Klaus; Zeuner, Rainald; Zielen, Stefan; Zimmermann, Martina; Želimir, Erić; Abou-
Chahla, Wadih; Aladjidi, Nathalie; Armari-Alla, Corinne; Barlogis, Vincent; Bayart, Sophie; Blanche, Stéphane;
Bodet, Damien; Bonnotte, Bernard; Borie, Raphaël; Boutard, Patrick; Boutboul, David; Briandet, Claire; Brion,
Jean-Paul; Brouard, Jacques; Castelle, Martin; Cathebras, Pascal; Carausu, Liana; Catherinot, Emilie; Cheikh,
Nathalie; Cheminant, Morgane; Comont, Thibault; Couderc, Louis-Jean; Cougoul, Pierre; Deville, Anne;
Devoldere, Catherine; Dore, Eric; Dulieu, Fabienne; Durieu, Isabelle; Entz-Werle, Natacha; Fieschi, Claire;
Galicier, Lionel; Gandemer, Virginie; Gardembas, Martine; Gourguechon, Clément; Grosbois, Bernard; Guffroy,
Aurélien; Guitton, Corinne; Guillerm, Gaëlle; Hamidou, Mohamed; Haro, Sophie; Hatchuel, Yves; Hermine,
Olivier; Hoarau, Cyrille; Humbert, Sébastien; Jaccard, Arnaud; Jais, Jean-Philippe; Jannier, Sarah; Jacquot,
Serge; Jaussaud, Roland; Jeandel, Pierre-Yves; Jeziorski, Eric; Kebaïli, Kamila; Korganow, Anne-Sophie;
Lambotte, Olivier; Lanternier, Fanny; Larroche, Claire; Launay, David; Le Guenno, Guillaume; Le Moigne,
Emmanuelle; Lecuit, Marc; Lefèvre, Guillaume; Lelièvre, Jean-Daniel; Levy, Romain; Li-Thiao-Te, Valérie;
Lortholary, Olivier; Luca, Luminita; Mallebranche, Coralie; Malphettes, Marion; Marie-Cardine, Aude; Martin-
Silva, Nicolas; Masseau, Agathe; Merlin, Etienne; Millot, Frédéric; Miot, Charline; Mirgot, Floriane; Moussouni,
Nabila; Mouthon, Luc; Néel, Antoine; Neven, Bénédicte; Nouar, Dalila; Nove-Josserand, Raphaële; Ouachée-
Chardin, Marie; Pagnier, Anne; Paillard, Catherine; Pasquet, Marlène; Pellier, Isabelle; Perlat, Antoinette;
Piguet, Christophe; Plantaz, Dominique; Rivière, Sophie; Raffray, Loïc; Roblot, Pascal; Rohrlich, Pierre-Simon;
Royer, Bruno; Salle, Valéry; Salvator, Hélène; Sarrot-Reynauld, Françoise; Servettaz, Amélie; Stephan, Jean-
Louis; Schleinitz, Nicolas; Suarez, Felipe; Swiader, Laure; Taque, Sophie; Thomas, Caroline; Tournilhac, Olivier;
Thumerelle, Caroline; Vannier, Jean-Pierre; Viallard, Jean-François.

References

1. Gray PE, David C. Inborn Errors of Immunity and Autoimmune Disease. *J Allergy Clin Immunol Pract* 2023;**11**(6):1602-22 doi: 10.1016/j.jaip.2023.04.018.

2. Hagele P, Staus P, Scheible R, Uhlmann A, Heeg M, Klemann C, Maccari ME, Ritterbusch H, Armstrong M, Cutcutache I, Elliott KS, von Bernuth H, Leahy TR, Leyh J, Holzinger D, Lehmberg K, Svec P, Masjosthusmann K, Hambleton S, Jakob M, Sparber-Sauer M, Kager L, Puzik A, Wolkewitz M, Lorenz MR, Schwarz K, Speckmann C, Rensing-Ehl A, Ehl S, group As. Diagnostic evaluation of paediatric autoimmune lymphoproliferative immunodeficiencies (ALPID): a prospective cohort study. *Lancet Haematol* 2024;**11**(2):e114-e26 doi: 10.1016/S2352-3026(23)00362-9.

3. Hauck F, Voss R, Urban C, Seidel MG. Intrinsic and extrinsic causes of malignancies in patients with primary immunodeficiency disorders. *The Journal of allergy and clinical immunology* 2018;**141**(1):59-68 e4 doi: 10.1016/j.jaci.2017.06.009.

4. Seidel MG, Hauck F. Multilayer concept of autoimmune mechanisms and manifestations in inborn errors of immunity: Relevance for precision therapy. *The Journal of allergy and clinical immunology* 2024;**153**(3):615-28 e4 doi: 10.1016/j.jaci.2023.12.022.

5. Vaseghi-Shanjani M, Smith KL, Sara RJ, Modi BP, Branch A, Sharma M, Lu HY, James EL, Hildebrand KJ, Biggs CM, Turvey SE. Inborn errors of immunity manifesting as atopic disorders. *The Journal of allergy and clinical immunology* 2021;**148**(5):1130-39 doi: 10.1016/j.jaci.2021.08.008.

6. Poli MC, Aksentijevich I, Bousfiha A, Cunningham-Rundles C, Hambleton S, Klein C, Morio T, Picard C, Puel A, Rezaei N, Seppänen MRJ, Somech R, Su HC, Sullivan KE, Torgerson TR, Meyts I, Tangye SG. Human inborn errors of immunity: 2024 Update on the classification from the International Union of Immunological Societies Expert Committee. Secondary Human inborn errors of immunity: 2024 Update on the classification from the International Union of Immunological Societies Expert Committee 2024. https://wp-iuis.s3.eu-west-1.amazonaws.com/app/uploads/2025/01/08170257/IEI-Final-Update-of-2024-Report-Jan-2025.pdf.

7. Seidel MG. Rethinking PIDs: Why the distinction between primary and secondary immune disorders is more frequently relevant than that between inborn and acquired errors of immunity. *The Journal of allergy and clinical immunology* 2024;**153**(6):1543-45 doi: 10.1016/j.jaci.2024.01.018.

8. Grimbacher B, Party ERW. The European Society for Immunodeficiencies (ESID) registry 2014. *Clinical and experimental immunology* 2014;**178 Suppl 1**(Suppl 1):18-20 doi: 10.1111/cei.12496.

9. Guzman D, Veit D, Knerr V, Kindle G, Gathmann B, Eades-Perner AM, Grimbacher B. The ESID Online Database network. *Bioinformatics* 2007;**23**(5):654-5 doi: 10.1093/bioinformatics/btl675.

10. Seidel MG, Kindle G, Gathmann B, Quinti I, Buckland M, van Montfrans J, Scheible R, Rusch S, Gasteiger LM, Grimbacher B, Mahlaoui N, Ehl S, Party ERW, collaborators. The European Society for Immunodeficiencies (ESID) Registry Working Definitions for the Clinical Diagnosis of Inborn Errors of Immunity. *J Allergy Clin Immunol Pract* 2019;**7**(6):1763-70 doi: 10.1016/j.jaip.2019.02.004.

11. Alligon M, Mahlaoui N, Bouaziz O. Pitfalls in time-to-event analysis of registry data: a
tutorial based on simulated and real cases. *Front Epidemiol* 2024;**4**:1386922 doi:
10.3389/fepid.2024.1386922.

12. Thalhammer J, Kindle G, Nieters A, Rusch S, Seppanen MRJ, Fischer A, Grimbacher B,
Edgar D, Buckland M, Mahlaoui N, Ehl S, European Society for Immunodeficiencies
Registry Working P. Initial presenting manifestations in 16,486 patients with inborn
errors of immunity include infections and noninfectious manifestations. *The Journal of*
allergy and clinical immunology 2021;**148**(5):1332-41 e5 doi:
10.1016/j.jaci.2021.04.015.

13. Abolhassani H, Azizi G, Sharifi L, Yazdani R, Mohsenzadegan M, Delavari S, Sohani M,
Shirmast P, Chavoshzadeh Z, Mahdaviani SA, Kalantari A, Tavakol M, Jabbari-Azad
F, Ahanchian H, Momen T, Sherkat R, Sadeghi-Shabestari M, Aleyasin S,
Esmaeilzadeh H, Al-Herz W, Bousfiha AA, Condino-Neto A, Seppanen M, Sullivan
KE, Hammarstrom L, Modell V, Modell F, Quinn J, Orange JS, Aghamohammadi A.
Global systematic review of primary immunodeficiency registries. *Expert Rev Clin*
Immunol 2020;**16**(7):717-32 doi: 10.1080/1744666X.2020.1801422.

14. European Society for Immunodeficiencies Registry WP. ESID-R Publications. Secondary
ESID-R Publications 2025. https://esid.org/working-parties/registry-working-
party/registry-publications/.

15. Brown L, Xu-Bayford J, Allwood Z, Slatter M, Cant A, Davies EG, Veys P, Gennery AR,
Gaspar HB. Neonatal diagnosis of severe combined immunodeficiency leads to
significantly improved survival outcome: the case for newborn screening. *Blood*
2011;**117**(11):3243-6 doi: 10.1182/blood-2010-08-300384.

16. Chan K, Puck JM. Development of population-based newborn screening for severe
combined immunodeficiency. *The Journal of allergy and clinical immunology*
2005;**115**(2):391-8 doi: 10.1016/j.jaci.2004.10.012.

17. Routes JM, Grossman WJ, Verbsky J, Laessig RH, Hoffman GL, Brokopp CD, Baker
MW. Statewide newborn screening for severe T-cell lymphopenia. *JAMA*
2009;**302**(22):2465-70 doi: 10.1001/jama.2009.1806.

18. Thakar MS, Logan BR, Puck JM, Dunn EA, Buckley RH, Cowan MJ, O'Reilly RJ, Kapoor
N, Satter LF, Pai SY, Heimall J, Chandra S, Ebens CL, Chellapandian D, Williams O,
Burroughs LM, Saldana BD, Rayes A, Madden LM, Chandrakasan S, Bednarski JJ,
2nd, DeSantes KB, Cuvelier GDE, Teira P, Gillio AP, Eissa H, Knutsen AP, Goldman
FD, Aquino VM, Shereck EB, Moore TB, Caywood EH, Lugt MTV, Rozmus J, Broglie
L, Yu LC, Shah AJ, Andolina JR, Liu X, Parrott RE, Dara J, Prockop S, Martinez CA,
Kapadia M, Jyonouchi SC, Sullivan KE, Bleesing JJ, Chaudhury S, Petrovic A, Keller
MD, Quigg TC, Parikh S, Shenoy S, Seroogy C, Rubin T, Decaluwe H, Routes JM,
Torgerson TR, Leiding JW, Pulsipher MA, Kohn DB, Griffith LM, Haddad E, Dvorak
CC, Notarangelo LD. Measuring the effect of newborn screening on survival after
haematopoietic cell transplantation for severe combined immunodeficiency: a 36-
year longitudinal study from the Primary Immune Deficiency Treatment Consortium.
Lancet 2023;**402**(10396):129-40 doi: 10.1016/S0140-6736(23)00731-6.

19. I. Jspeert H, Edwards ESJ, O'Hehir RE, Dalm V, van Zelm MC. Update on inborn errors
of immunity. *The Journal of allergy and clinical immunology* 2024 doi:
10.1016/j.jaci.2024.12.1075.

20. Seidel MG. Baby genome screening: paving the way to genetic discrimination? *Bmj*
2017;**358**:j3294 doi: 10.1136/bmj.j3294.

21. Vetrie D, Vorechovsky I, Sideras P, Holland J, Davies A, Flinter F, Hammarstrom L,
Kinnon C, Levinsky R, Bobrow M, et al. The gene involved in X-linked
agammaglobulinaemia is a member of the src family of protein-tyrosine kinases.
Nature 1993;**361**(6409):226-33 doi: 10.1038/361226a0.

22. Kuehn HS, Ouyang W, Lo B, Deenick EK, Niemela JE, Avery DT, Schickel JN, Tran DQ,
Stoddard J, Zhang Y, Frucht DM, Dumitriu B, Scheinberg P, Folio LR, Frein CA,
Price S, Koh C, Heller T, Seroogy CM, Huttenlocher A, Rao VK, Su HC, Kleiner D,
Notarangelo LD, Rampertaap Y, Olivier KN, McElwee J, Hughes J, Pittaluga S,
Oliveira JB, Meffre E, Fleisher TA, Holland SM, Lenardo MJ, Tangye SG, Uzel G.
Immune dysregulation in human subjects with heterozygous germline mutations in
CTLA4. *Science* 2014;**345**(6204):1623-27 doi: 10.1126/science.1255904.

23. Schubert D, Bode C, Kenefeck R, Hou TZ, Wing JB, Kennedy A, Bulashevska A,
Petersen BS, Schaffer AA, Gruning BA, Unger S, Frede N, Baumann U, Witte T,
Schmidt RE, Dueckers G, Niehues T, Seneviratne S, Kanariou M, Speckmann C, Ehl
S, Rensing-Ehl A, Warnatz K, Rakhmanov M, Thimme R, Hasselblatt P, Emmerich F,
Cathomen T, Backofen R, Fisch P, Seidl M, May A, Schmitt-Graeff A, Ikemizu S,
Salzer U, Franke A, Sakaguchi S, Walker LSK, Sansom DM, Grimbacher B.
Autosomal dominant immune dysregulation syndrome in humans with CTLA4
mutations. *Nat Med* 2014;**20**(12):1410-16 doi: 10.1038/nm.3746.

24. Fliegauf M, Bryant VL, Frede N, Slade C, Woon ST, Lehnert K, Winzer S, Bulashevska
A, Scerri T, Leung E, Jordan A, Keller B, de Vries E, Cao H, Yang F, Schaffer AA,
Warnatz K, Browett P, Douglass J, Ameratunga RV, van der Meer JW, Grimbacher
B. Haploinsufficiency of the NF-kappaB1 Subunit p50 in Common Variable
Immunodeficiency. *Am J Hum Genet* 2015;**97**(3):389-403 doi:
10.1016/j.ajhg.2015.07.008.

25. Haimel M, Pazmandi J, Heredia RJ, Dmytrus J, Bal SK, Zoghi S, van Daele P, Briggs
TA, Wouters C, Bader-Meunier B, Aeschlimann FA, Caorsi R, Eleftheriou D,
Hoppenreijs E, Salzer E, Bakhtiar S, Derfalvi B, Saettini F, Kusters MAA, Elfeky R,
Truck J, Riviere JG, van der Burg M, Gattorno M, Seidel MG, Burns S, Warnatz K,
Hauck F, Brogan P, Gilmour KC, Schuetz C, Simon A, Bock C, Hambleton S, de
Vries E, Robinson P, van Gijn M, Boztug K. Curation and Expansion of Human
Phenotype Ontology for Defined Groups of Inborn Errors of Immunity. *The Journal of*
allergy and clinical immunology 2021 doi: 10.1016/j.jaci.2021.04.033.

26. Akalu YT, Bogunovic D. Inborn errors of immunity: an expanding universe of disease and
genetic architecture. *Nat Rev Genet* 2024;**25**(3):184-95 doi: 10.1038/s41576-023-
00656-z.

27. Passweg JR, Baldomero H, Ciceri F, de la Camara R, Glass B, Greco R, Hazenberg
MD, Kalwak K, McLornan DP, Neven B, Peric Z, Risitano AM, Ruggeri A, Snowden
JA, Sureda A. Hematopoietic cell transplantation and cellular therapies in Europe
2022. CAR-T activity continues to grow; transplant activity has slowed: a report from
the EBMT. *Bone marrow transplantation* 2024;**59**(6):803-12 doi: 10.1038/s41409-
024-02248-9.

28. European Commission E-E. SPIDER pseudonymisation tool. Secondary SPIDER
pseudonymisation tool 2024. https://eu-rd-platform.jrc.ec.europa.eu/spider/.

29. Riviere JG, Soler Palacin P, Butte MJ. Proceedings from the inaugural Artificial
Intelligence in Primary Immune Deficiencies (AIPID) conference. *The Journal of*
allergy and clinical immunology 2024;**153**(3):637-42 doi: 10.1016/j.jaci.2024.01.002.

30. Kohler S, Gargano M, Matentzoglu N, Carmody LC, Lewis-Smith D, Vasilevsky NA,
Danis D, Balagura G, Baynam G, Brower AM, Callahan TJ, Chute CG, Est JL, Galer
PD, Ganesan S, Griese M, Haimel M, Pazmandi J, Hanauer M, Harris NL, Hartnett
MJ, Hastreiter M, Hauck F, He Y, Jeske T, Kearney H, Kindle G, Klein C, Knoflach K,
Krause R, Lagorce D, McMurry JA, Miller JA, Munoz-Torres MC, Peters RL, Rapp
CK, Rath AM, Rind SA, Rosenberg AZ, Segal MM, Seidel MG, Smedley D, Talmy T,
Thomas Y, Wiafe SA, Xian J, Yuksel Z, Helbig I, Mungall CJ, Haendel MA, Robinson
PN. The Human Phenotype Ontology in 2021. *Nucleic Acids Res*
2021;**49**(D1):D1207-D17 doi: 10.1093/nar/gkaa1043.

Legends

Table 1. *European Society for Immunodeficiencies* **(ESID) registry patient characteristics.**

Figure 1. Ages and main manifestations at onset of inborn errors of immunity (IEI) or primary immune disorders (PID)**.** ***A,*** age groups of age at disease onset, clinical diagnosis of IEI/PID, and of genetic diagnosis are shown. ***B,*** Venn diagram of main manifestations of IEI/PID with absolute patient numbers and proportions.

Figure 2. Representation of genetic diagnoses of known IEI/PID in the ESID registry. *A,* the number and proportion of patients with a genetic versus those without a genetic diagnosis is shown in descending order. ***B,*** the top 5 genetic defects or deletions registered in the ESIDR per all 10 IUIS categories of IEI are presented; the nomenclature in registry diagnosis and gene fields was not regularly updated/changed, showing, *e.g*., GP91-phox instead of *CYBB* and p47-phox instead of *NCF1*. ***C,*** at the time point of data analysis in March 2024, the number of monogenic IEI documented in the ESID-R as proportion of the number of all known genes in the respective 2022 IUIS category is shown following the order of IUIS categories from left to right. ***D,*** the top 50 genes mutated in ESID-R patients with monogenic IEI/PID are shown with respective patient numbers in descending order.

Figure 3. Major treatment modalities of patients with IEI/PID as recorded in the ESID-R. The left panel shows absolute numbers, the right panel shows the relative proportions of major treatments such as immunoglobulin replacement therapy (data on route, interval and dose were recorded but are not shown), allogeneic hematopoietic stem cell transplantation (AlloSCT), autologous gene therapy (GT) and on splenectomy. "No therapy" and IgRT refer to the latest follow-up time point.

Figure 4. Survival probabilities of main categories of IEI/PID and the living status at last news. *A,* Inverse cumulative incidence curves as described and referenced in the supplementary material. Start = age at diagnosis, stop = age at last news, event = living status (0= censored, 1 = deceased first, 2 = curative therapy first). There were 21,206 patients censored, 1,960 patients who deceased first, and 2,901 patients who had a curative therapy first (not showing deaths after curative therapy); ***B,*** Livings status in relation to the age at last news is shown.

Figure 5. Number, categories and citation counts of ESID-R-based publications. A total of 7,980 citations of 84 peer-reviewed ESID-R-based publications until early 2024 was recorded. See also the website for regular updates of ESID-R-related publications at https://esid.org/working-parties/registry-working-party/registry-publications/

Table 1. *European Society for Immunodeficiencies* (ESID) registry patient characteristics.

Covariate	Values	Whole cohort N=30628	CID N=2531	Syndromic N=4239	PAD N=15123	PIRD N=2171	Phagocyte N=2548	Innate N=823	AIS N=1042	Complement N=1482	BMF N=85	Phenocopies N=30	other N=554
Gender (male)	M	16767 (54.8%)	1470 (58.1%)	2473 (58.3%)	7782 (51.5%)	1329 (61.2%)	1693 (66.5%)	387 (47%)	575 (55.2%)	681 (46%)	57 (67.1%)	19 (63.3%)	301 (54.3%)
	F	13855 (45.2%)	1061 (41.9%)	1766 (41.7%)	7337 (48.5%)	841 (38.8%)	854 (33.5%)	436 (53%)	467 (44.8%)	801 (54%)	28 (32.9%)	11 (36.7%)	253 (45.7%)
Year of birth		1997 [1976;2008]	2007 [1997;2014]	2005 [1995;2012]	1984 [1963;2001]	2004 [1993;2011]	2002 [1991;2010]	2003 [1991;2011]	2008 [1999;2013]	1991 [1972;2006]	2004 [1987;2011]	2001.5 [1995;2008.5]	1998 [1974;2007]
Familial case	N	21854 (76.6%)	1725 (72.9%)	3002 (74.1%)	11949 (85.4%)	1266 (61.1%)	1468 (65.4%)	492 (62.2%)	771 (76.6%)	645 (46.5%)	57 (75%)	28 (93.3%)	451 (87.7%)
	Y	6681 (23.4%)	641 (27.1%)	1047 (25.9%)	2049 (14.6%)	805 (38.9%)	778 (34.6%)	299 (37.8%)	236 (23.4%)	742 (53.5%)	19 (25%)	2 (6.7%)	63 (12.3%)
Consanguinity	N	24731 (87.2%)	1426 (60.8%)	3247 (82.1%)	13381 (95.5%)	1602 (78.8%)	1699 (75.6%)	589 (77.3%)	920 (93.2%)	1303 (93.1%)	66 (84.6%)	28 (100%)	470 (92.7%)
	Y	3624 (12.8%)	921 (39.2%)	708 (17.9%)	631 (4.5%)	432 (21.2%)	547 (24.4%)	173 (22.7%)	67 (6.8%)	96 (6.9%)	12 (15.4%)	0 (0%)	37 (7.3%)
Age at onset of symptoms		3 [0.4;14.1]	0.3 [0;1.6]	0.5 [0;2.2]	8 [2;30]	2.2 [0.3;7.5]	0.7 [0.1;3]	1.2 [0.2;3.8]	2.8 [0.7;5]	8 [3;18]	3 [0.3;8.1]	2.8 [0.6;4.9]	3 [0.7;15.4]
Immune dysregulation at onset	N	21482 (77.2%)	1712 (77.5%)	3242 (84.1%)	11175 (80.3%)	616 (31.6%)	1944 (86.2%)	632 (84.9%)	536 (55%)	1182 (88.3%)	49 (67.1%)	4 (13.3%)	390 (80.7%)
	Y	6347 (22.8%)	498 (22.5%)	613 (15.9%)	2741 (19.7%)	1335 (68.4%)	311 (13.8%)	112 (15.1%)	438 (45%)	156 (11.7%)	24 (32.9%)	26 (86.7%)	93 (19.3%)
Infection at onset	N	8133 (28.9%)	436 (19.6%)	2195 (56.6%)	1920 (13.6%)	1161 (59.6%)	590 (26%)	108 (14.4%)	705 (72.5%)	827 (61.5%)	44 (60.3%)	19 (67.9%)	128 (26.3%)
	Y	19961 (71.1%)	1786 (80.4%)	1680 (43.4%)	12202 (86.4%)	787 (40.4%)	1681 (74%)	643 (85.6%)	267 (27.5%)	518 (38.5%)	29 (39.7%)	9 (32.1%)	359 (73.7%)
Malignancy at onset	N	27294 (98.3%)	2180 (98.6%)	3766 (97.8%)	13620 (98.1%)	1885 (97.1%)	2235 (99.4%)	741 (99.7%)	969 (99.8%)	1333 (99.8%)	69 (94.5%)	25 (89.3%)	471 (97.7%)
	Y	479 (1.7%)	30 (1.4%)	85 (2.2%)	269 (1.9%)	57 (2.9%)	13 (0.6%)	2 (0.3%)	2 (0.2%)	3 (0.2%)	4 (5.5%)	3 (10.7%)	11 (2.3%)
Syndromal manifestations at onset	N	24549 (88.3%)	2046 (92.6%)	1651 (42.7%)	13574 (97.7%)	1798 (92.5%)	2075 (92.3%)	683 (91.8%)	919 (94.6%)	1274 (95.4%)	40 (54.8%)	28 (100%)	461 (95.4%)
	Y	3253 (11.7%)	164 (7.4%)	2220 (57.3%)	320 (2.3%)	145 (7.5%)	174 (7.7%)	61 (8.2%)	52 (5.4%)	62 (4.6%)	33 (45.2%)	0 (0%)	22 (4.6%)
Other onset	N	24466 (87.9%)	1992 (90.1%)	3298 (85.5%)	12947 (93.1%)	1782 (91.6%)	2086 (92.7%)	685 (92.1%)	545 (55.6%)	647 (48.1%)	59 (80.8%)	28 (96.6%)	397 (81.9%)
	Y	3366 (12.1%)	220 (9.9%)	559 (14.5%)	963 (6.9%)	163 (8.4%)	164 (7.3%)	59 (7.9%)	436 (44.4%)	699 (51.9%)	14 (19.2%)	1 (3.4%)	88 (18.1%)

Covariate	Values	Whole cohort N=30628	CID N=2531	Syndromic N=4239	PAD N=15123	PIRD N=2171	Phagocyte N=2548	Innate N=823	AIS N=1042	Complement N=1482	BMF N=85	Phenocopies N=30	other N=554
No clinical symptoms	N	27605 (94.1%)	2188 (91%)	3770 (92.5%)	13875 (95.9%)	1926 (91.5%)	2238 (92.3%)	735 (91.9%)	966 (97.2%)	1326 (91.2%)	71 (91%)	28 (100%)	482 (97.6%)
	Y	1725 (5.9%)	217 (9%)	306 (7.5%)	595 (4.1%)	180 (8.5%)	187 (7.7%)	65 (8.1%)	28 (2.8%)	128 (8.8%)	7 (9%)	0 (0%)	12 (2.4%)
Age at clinical diagnosis (CD)		8.2 [2;30.1]	0.7 [0.3;4.6]	2.4 [0.4;6.9]	22 [6.2;44]	5.1 [1;13]	2 [0.5;7]	4.2 [1.1;14]	5 [2.7;11.4]	15 [5.2;28.8]	5 [1.3;13.7]	5.2 [3;9.8]	7.5 [2.3;34.1]
Delay between onset and CD		1.3 [0.1;5.1]	0.2 [0;1.3]	0.8 [0;3.9]	2.7 [0.7;7.5]	0.4 [0;3.4]	0.5 [0.1;2.2]	1 [0.1;5.6]	1.5 [0.5;4]	1.1 [0.1;7]	0.5 [0.1;2.1]	0.9 [0.3;3.9]	1.7 [0.3;4.7]
Genetics	Mutation found	12774 (44.5%)	1470 (63.1%)	3647 (88.8%)	2251 (16%)	1671 (80.8%)	1693 (71.8%)	529 (68.1%)	662 (65.3%)	772 (55.6%)	49 (71%)	30 (100%)	0 (0%)
	No mutation found	2852 (9.9%)	319 (13.7%)	79 (1.9%)	1850 (13.1%)	195 (9.4%)	106 (4.5%)	81 (10.4%)	71 (7%)	27 (1.9%)	9 (13%)	0 (0%)	115 (24.7%)
	Not tested	11956 (41.7%)	403 (17.3%)	325 (7.9%)	9398 (66.8%)	125 (6%)	478 (20.3%)	97 (12.5%)	250 (24.7%)	569 (41%)	8 (11.6%)	0 (0%)	303 (65%)
	Pending	1097 (3.8%)	136 (5.8%)	57 (1.4%)	576 (4.1%)	76 (3.7%)	80 (3.4%)	70 (9%)	31 (3.1%)	20 (1.4%)	3 (4.3%)	0 (0%)	48 (10.3%)
Age at genetic diagnosis (GD)		5.5 [1.1;15]	0.9 [0.3;6.7]	3.2 [0.4;9.7]	9.4 [2.9;25]	8.3 [1.8;17.3]	4.4 [1.1;12.7]	10 [3.2;20.4]	8.4 [4;22]	13.2 [5;26.9]	8.2 [3.6;17.4]	9.8 [5;14.1]	NA [NA;NA]
Delay between CD and GD		0.3 [0;2.9]	0.2 [0;1.2]	0.2 [0;1.7]	1.3 [0.1;8.8]	0.2 [0;2.2]	0.3 [0;2.2]	0.8 [0.1;5.9]	0.1 [0;1.1]	0.2 [0;1.6]	0.4 [0.1;2.6]	1 [0.2;4.9]	NA [NA;NA]
Reason of GD	Clinical	8787 (88%)	975 (87.7%)	2601 (90.8%)	1629 (89.2%)	1192 (82.2%)	1081 (90.6%)	380 (85.4%)	515 (88.6%)	350 (77.8%)	39 (92.9%)	25 (96.2%)	0 (0%)
	Family	975 (9.8%)	77 (6.9%)	153 (5.3%)	170 (9.3%)	245 (16.9%)	100 (8.4%)	63 (14.2%)	65 (11.2%)	99 (22%)	3 (7.1%)	0 (0%)	0 (0%)
	Neonatal	114 (1.1%)	41 (3.7%)	45 (1.6%)	20 (1.1%)	2 (0.1%)	3 (0.3%)	0 (0%)	1 (0.2%)	1 (0.2%)	0 (0%)	1 (3.8%)	0 (0%)
	Prenatal	112 (1.1%)	19 (1.7%)	64 (2.2%)	7 (0.4%)	11 (0.8%)	9 (0.8%)	2 (0.4%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Sequencing method of GD	Gene sequencing	7693 (83.3%)	848 (80.1%)	2122 (82%)	1421 (83.5%)	1154 (85.3%)	970 (87.7%)	323 (77.5%)	459 (85.5%)	347 (84.6%)	26 (70.3%)	23 (95.8%)	0 (0%)
	Non-genetic definitive test	441 (4.8%)	35 (3.3%)	284 (11%)	11 (0.6%)	12 (0.9%)	59 (5.3%)	3 (0.7%)	0 (0%)	36 (8.8%)	1 (2.7%)	0 (0%)	0 (0%)
	Whole exome/genome sequencing	1100 (11.9%)	176 (16.6%)	183 (7.1%)	270 (15.9%)	187 (13.8%)	77 (7%)	91 (21.8%)	78 (14.5%)	27 (6.6%)	10 (27%)	1 (4.2%)	0 (0%)
Malignancy	N	18176 (91.1%)	1578 (93.4%)	2491 (88.9%)	8759 (89.4%)	1595 (91.4%)	1609 (95.6%)	569 (94.7%)	604 (98.1%)	632 (97.7%)	66 (90.4%)	16 (72.7%)	257 (89.9%)
	Y	1783 (8.9%)	112 (6.6%)	312 (11.1%)	1034 (10.6%)	150 (8.6%)	74 (4.4%)	32 (5.3%)	12 (1.9%)	15 (2.3%)	7 (9.6%)	6 (27.3%)	29 (10.1%)

Covariate	Values	Whole cohort	CID	Syndromic	PAD	PIRD	Phagocyte	Innate	AIS	Complement	BMF	Phenocopies	other
		N=30628	N=2531	N=4239	N=15123	N=2171	N=2548	N=823	N=1042	N=1482	N=85	N=30	N=554
Covid19	N	5380 (70.3%)	456 (81%)	801 (76.1%)	2613 (64.1%)	434 (74.2%)	363 (81%)	130 (77.4%)	260 (74.9%)	177 (76.3%)	15 (78.9%)	6 (75%)	125 (79.6%)
	Y	2273 (29.7%)	107 (19%)	251 (23.9%)	1461 (35.9%)	151 (25.8%)	85 (19%)	38 (22.6%)	87 (25.1%)	55 (23.7%)	4 (21.1%)	2 (25%)	32 (20.4%)
Living status	Alive	20680 (67.5%)	1419 (56.1%)	2580 (60.9%)	10740 (71%)	1519 (70%)	1534 (60.2%)	546 (66.3%)	729 (70%)	1127 (76%)	52 (61.2%)	22 (73.3%)	412 (74.4%)
	Deceased	3216 (10.5%)	615 (24.3%)	676 (15.9%)	1117 (7.4%)	356 (16.4%)	269 (10.6%)	87 (10.6%)	23 (2.2%)	18 (1.2%)	22 (25.9%)	3 (10%)	30 (5.4%)
	Lost to follow-up	6732 (22%)	497 (19.6%)	983 (23.2%)	3266 (21.6%)	296 (13.6%)	745 (29.2%)	190 (23.1%)	290 (27.8%)	337 (22.7%)	11 (12.9%)	5 (16.7%)	112 (20.2%)
Age at last follow-up/death		20.2 [9.4;42.6]	8.7 [2;18.7]	12.5 [6.3;19.9]	35.4 [16.9;55.9]	14.8 [6.5;24.2]	13.7 [5.8;23.6]	15 [7;27.3]	11.2 [6;20.9]	27.3 [13.6;46.4]	14.6 [6.3;23.7]	18.2 [12;22.8]	19 [8.4;44.1]
Living status (stops at CT)	Alive at last follow-up	24222 (81.1%)	928 (37.1%)	3144 (76.5%)	13522 (92%)	1313 (61%)	1680 (67%)	695 (86.1%)	956 (95.1%)	1463 (99.1%)	41 (49.4%)	25 (83.3%)	455 (90.6%)
	Deceased	2226 (7.5%)	258 (10.3%)	496 (12.1%)	966 (6.6%)	186 (8.6%)	181 (7.2%)	74 (9.2%)	17 (1.7%)	11 (0.7%)	9 (10.8%)	3 (10%)	25 (5%)
	Curative therapy	3431 (11.5%)	1315 (52.6%)	470 (11.4%)	216 (1.5%)	653 (30.3%)	648 (25.8%)	38 (4.7%)	32 (3.2%)	2 (0.1%)	33 (39.8%)	2 (6.7%)	22 (4.4%)
Age at last follow-up (stops at CT)		19.4 [8;42.6]	1.3 [0.5;11]	11.4 [5.2;19.3]	35.5 [16.8;55.9]	12.1 [3.2;22.7]	11.6 [4;21.8]	14.6 [6.6;27]	11.1 [5.9;20.7]	27.4 [13.6;46.5]	10 [5;20.5]	18.2 [11;22.8]	19 [7.8;44.1]
Follow-up duration (years)		7.2 [2.6;14]	4.6 [0.8;12]	8 [2.9;14]	7.9 [3.4;14.5]	5.6 [1.7;11.6]	7.7 [2;17.2]	6.6 [1.7;13.1]	4 [1.6;7.9]	7.7 [2.1;16.9]	6.7 [3.4;10.6]	11.2 [5.2;17.1]	5.2 [1.5;11]

Quantitative covariates: Median [Q1;Q3]; qualitative covariates: Effective (Percentage); the order of subgroups from left to right follows the IEI categories I-X of the 2022 IUIS classification available on the end date chosen for inclusion; values *Y* for "yes/present" for real positive values, *N* for "no" is shown to reflect the size of the correct comparison group, excluding "unknown" replies; ages in years; CT, curative / definitive therapy; diagnosis via newborn screening is recorded as subcategory of "no clinical symptoms / lab abnormalities only", where not all choices of the submenu are shown.

Figure 1A.

Figure 1B.

Figure 2A.

Figure 2C.

Figure 2D.

Figure 2B.

Figure 3.

Figure 4A.

Number at risk											
CID	113	222	128	70	51	44	27	12	4	0	0
Synd.	286	1487	678	249	105	46	13	1	0	0	0
PAD	78	2796	2354	1977	2117	1868	1433	946	295	28	0
PIRD	29	384	296	139	73	37	20	8	2	0	0
Phag.	80	801	498	258	118	46	17	4	2	0	0
Innate	13	221	124	63	50	42	17	8	1	0	0
AIS	6	244	80	47	32	16	10	3	2	0	0
Compl.	8	253	293	264	217	147	76	37	16	0	0

Figure 4B.

Figure 5.

Supplementary Material

 Inborn errors of immunity: manifestation, treatment, and outcome

 – an ESID registry 1994-2024 report on 30,628 patients

Supplementary Text

Historical evolution of the ESID registry, objectives, structure, and content

The biggest hurdles faced during the first restructuration 2000-2004 (from a hardcopy to the first online version) were ethical considerations, data protection, and obtaining industry funding. The ESID registry (ESID-R) steering group, led by the elected chairperson of the ESID-R working party, oversees all aspects of the evolution of the registry, including the design, operations, financing, and set-up of dedicated studies.

Specific objectives of the ESID-R include:

1. *Collection of epidemiological data:* The registry contains basic information on patients with IEI, including their medical history, diagnosis, genetics, treatment, and disease progression. Data on the prevalence of these rare diseases and the associated health burden are important for political decision-makers.
2. *Analysis and research:* By analyzing the collected data on these rare diseases through multicenter collaboration, researchers can gain new insights into the pathophysiology and treatment of primary immunodeficiencies.
3. *Support of clinical studies and trials:* Registry data can serve as the basis for clinical studies or trials aimed at developing new therapies, improving the treatment of patients affected by those diseases, and for post-marketing surveillance studies of new drugs and medicinal products.
4. *Improving patient care:* Through the analyses and publication of registry data, the registry helps improve IEI patient care by providing physicians and healthcare providers/decision-makers with access to up-to-date information and practices.

Both versions of the ESID-R operating from 2004-2014 and 2014-2024 had the following key features[1]:

1. *Database Architecture:* The ESID-R utilizes a sophisticated database architecture to manage the vast amount of data collected from patients. This database is designed to be scalable, ensuring it can handle the increasing volume of patient records and research findings over time.
2. *Data Security:* Ensuring the security and privacy of patient data is paramount. The ESID-R implements stringent security measures to protect sensitive information, including encryption protocols and access controls

3. *Web-Based Platform:* The ESID-R is accessed through a web-based platform, allowing
authorized users to input and access data remotely. This platform is user-friendly and
intuitive, facilitating seamless data entry and retrieval processes. Only an up-to-date web-
browser and stable internet connection are required to access the system. No additional on-
site software needs to be installed/maintained at the documentation centers.
4. *Framework and Technologies:* The ESID-R is built using a combination of programming
languages, frameworks, and technologies suited to its specific requirements. Common
technologies include SQL databases for data storage, web development frameworks for the
user interface, and data analysis tools for reporting and research purposes.

The three-level content structure

The old version of the online ESID-R (2004-2014) was closed for documentation on June 25, 2014.
During the first main revision of the online registry in 2014, a three-level structure was implemented.
Level 1 addresses the task of the registry as a simple epidemiological tool. To achieve patient
registration by as many centers as possible, the dataset is kept short and simple such that caregivers
who know the patient well can fill in the data without consulting the patient chart in much detail. It
contains basic epidemiological information, information about main categories of symptoms at initial
presentation and information about some key therapies (immunoglobulins, HSCT, gene therapy). The
yearly update is restricted to survival and implementation of major therapies. The Level 1 registration
also serves to identify patients for additional in-depth studies including interventional trials. Level 2
was designed for long-term studies on certain diseases or disease groups without a predefined end.
Level 2 datasets allow registration of a limited number of clinical manifestations and lab values, that
are provided on a yearly basis. It was particularly designed to capture longitudinal data in disease
groups, for which the molecular basis has not been fully resolved (such as CVID or specific antibody
deficiencies). Again, the datasets are kept simple to limit the burden of registration. Level 3 was
designed for studies on particular genetically defined diseases based on a study protocol with a
defined endpoint. The level 3 dataset can contain very detailed information including grading of
symptoms, more sophisticated lab values, detailed treatment information and quality of life data.
Registration intervals are at least yearly. Level 3 studies usually offer financial support for the
documenting centers.

The level 2 and 3 projects or 'sub-studies' are designed by independent research groups. To apply
for implementation of a study in the ESID registry, a statistical evaluation plan, a time and budget
forecast, need to be defined in a standardized study proposal, which is assessed for feasibility and to
avoid redundancy or competing interests by the registry steering committee and the elected
chairperson in regular meetings. There should also be a financial contribution from the PI or research
institution to (at least partly) cover the costs associated with the set-up, coordination, data
maintenance, and analyses of these registry sub-studies (details on ESID-website, fee structure).
When signing the informed consent, patients may choose to have their anonymized data be included
either in sub-studies of academic research groups or of pharmaceutical industry-sponsored
consortia, or none of them.

Supplementary Methods

The ESID-R operates based on a technical infrastructure designed to securely receive, store, and enable the analysis of data related to IEI. These data cover, for example, various aspects of the disease, including genetic mutations, clinical symptoms, laboratory test results, imaging findings, and treatment outcomes. Patients with IEI are registered in the web-based database when their disease data are submitted by an authorized physician or healthcare professional or documenting personnel. Centers or sites where patients are cared for are typically university clinics/hospitals and have a data transfer and general data protection regulation (GDPR)-compliant handling agreement and a local IRB approval in place. The data are de-identified and kept strictly confidential to protect patient privacy.

The Kaplan-Meier curves were generated as described(2). The Kaplan-Meier is done using the start and stop notation to take care of the left truncation effect. Start = age at diagnosis, stop = age at last news, event = living status (0= censored, 1 = deceased first, 2 = curative therapy first). There were 21,206 patients censored, 1,960 patients who deceased first, and 2,901 patients who had a curative therapy first.

Supplementary Results and Discussion

Newborn screening and longitudinal patient documentation in the ESID-R. The ESID-R data structure does not (yet) allow an analysis of improved diagnostic delay and survival in patients diagnosed with SCID in the ESID-R in the NBS (newborn screening) era. Firstly, NBS for SCID has been performed sequentially over the last decade, but is still not generally available; and secondly, collecting the intended annual follow-up information may cause a system-inherent delay, including underreporting of SCID patients following definitive therapy (or death), before their data are stored in the ESID-R. The ESID-R working party continues to face significant challenges while aiming to integrate data collected over a patient's entire disease course into a single registry, as this ranges from NBS or clinical diagnosis, through different clinical events, such as a form of definitive therapy, and/or chronic treatment, and/or their optional participation in observational sub-studies and on to a cure or until death.

The ESID-R in the international landscape of registries for patients with rare immunological diseases. To depict the current situation of the global IEI registry landscape and participants' motivations, we searched PubMed, communicated with representatives of sister societies around the globe, and performed an online survey in 2024. Of 533 publications retrieved by searching for *primary immunodeficiency patient registry* and exclusion of disease-specific or other secondary sub-studies, the remaining publications showed that a vast majority of IEI/PID registries are national epidemiological databases with a limited temporal and geographical scope regarding longitudinal patient data(2-20). They, naturally, contain a substantially smaller number of patient datasets than the continental ESID registry, which also includes data from patients of some neighboring countries. As such, they are valuable and necessary platforms for the connection of healthcare providers and clinical experts, as well as for health politicians and economists. A relatively recently published overview of the global registry landscape and the burden of IEI/PID from 80 countries evaluated the

distribution of IEI/PID diagnoses and other factors such as the diagnostic delay or immunoglobulin
treatment in relation to the countries' resources and societal factors such as the rate of
consanguinity(21). These data show that the key drivers to implement and maintain an IEI/PID
patient registry range from health policy, warranting access to expert care and modern treatments,
over patient advocacy, to clinical, biological, and pharmaceutical research and development.
Importantly, IPOPI, the global patient organization for IEI, has identified the existence of registries as
one of the six key principles of care as part of its PID LIFE Index, a tool to measure the quality of
IEI/PID healthcare in a country(22). To analyze the policy of patient inclusion and obtain feedback
from documenting physicians or study nurses, we designed a survey distributed through ESID in
2024. A total of 43 physicians from 29 different European and non-European countries participated
in this survey (Supplementary Table 5). Seventy percent of respondents (30/43) said they would
exclusively register their patients into the ESID-R; 30% of the survey respondents (13/43) said they
would also enter patient data into one of the following disease-specific (other than general IEI/PID)
registries: *Eurofever, Severe Chronic Neutropenia International Registry (SCNIR), Juvenile*
Inflammatory Rheumatism (JIR) Cohort, and the registry of the *EBMT/Center for International Blood*
and Marrow Transplant Research (CIBMTR). Motivations for entering data into the ESID-R included
collaborative research (88%), networking (84%), taking part in academic or pharmaceutical studies
(84% and 72%), and improvements in patient care (72%) and for IEI education purposes. Reasons
given for not entering patient data into the ESID-R were: lack of personnel (84%), legal challenges
regarding data transfer (37%), no obvious local benefit (23%), and concerns about small countries
being underrepresented (14%). Survey participants stated that they would like a simplified version of
the patient consent form, easy interoperability between national registries and the ESID-R, and grant
incentives for entering patient data, especially if these were needed at a higher level of granularity.

Human Phenotype Ontology (HPO) in IEI/PID, and potential benefit of integrating HPO into the
ESID-R. The Human Phenotype Ontology (HPO) comprehensively organizes and defines the
phenotypic features of human disease, supporting genomic and phenotypic analyses through
semantic similarity and machine learning algorithms(23). Currently, the HPO terminology for IEI/PID
still needs to be further developed with help of the community of clinical immunologists and
researchers focusing on IEI/PID: *i.e.*, new HPO terms for relevant phenotypic features to be added,
and existing HPO terms to be revised, such that the definitions and hierarchical placement in the HPO
reflect current knowledge(24). The HPO can be used with the Global Alliance for Genomics and
Health (GA4GH) Phenopacket Schema to represent not only phenotypic features, but also
measurements, biopsies, and treatments over the entire time course of a disease(25). Collections of
phenopackets may support translational research in immunology by providing standardized input
data for software programs designed to identify genotype-phenotype correlations, and characterize
disease subtypes(26). Integrating the HPO terminology into the ESID-R, would inspire studies within
or across IUIS categories, especially in monogenic entities. To make full use of the HPO, better
systems for simple but comprehensive and accurate data entry are needed.

Supplementary Material (Tables)

Supplementary Table 1. Contributing countries, sorted by the calculated prevalence per 100,000 inhabitants.

Country	Number of registered patients	Registered patients/100,000 inhabitants	Calculated prevalence/100,000 inhabitants
Slovakia	794	14.56	14.32
France	9,113	13.59	10.99
Belgium	1,333	11.61	11.36
Switzerland	658	7.67	7.55
Slovenia	129	6.18	6.18
Germany	5,056	6.08	5.73
United Kingdom	3,949	5.91	5.51
Ireland	280	5.67	5.48
Portugal	569	5.54	5.00
Czechia	573	5.37	4.98
Netherlands	715	4.13	3.96
Greece	387	3.61	3.46
Hungary	315	3.22	2.68
Austria	286	3.22	3.03
Bulgaria	175	2.51	2.42
Belarus	228	2.41	1.92
Lithuania	63	2.26	2.12
Spain	1,008	2.14	2.01
Poland	813	2.14	1.86
Estonia	26	1.96	1.88
Italy	1,036	1.72	1.58
Iran	1,284	1.55	1.39

Country	Number of registered patients	Registered patients/100,000 inhabitants	Calculated prevalence/100,000 inhabitants
Denmark	77	1.32	1.31
Serbia	82	1.18	1.17
Turkey	874	1.05	0.98
Croatia	36	0.89	0.79
Egypt	512	0.51	0.41
Ukraine	140	0.32	0.29
Romania	50	0.26	0.20
Israel	5	0.06	0.06
Russia	59	0.04	0.03
Bosnia and Herzegowina	1	0.03	0.03
Sweden	2	0.02	0.02

Of note, the ESID-R has no means to influence the balance of data entry per country or center,
resulting in differences of the proportion of children, adolescent and adult patients, the "weight"
given to certain diseases at some centers and their potential overrepresentation. Although measures
were taken and found to be successful to unify the classification of non-genetic diseases such as
common variable immune deficiency or selective IgA deficiency by applying the ESID registry working
definitions for making a clinical diagnosis, the variance in the "epidemiological activity" of some
centers as compared to others remains.

Supplementary Table 2. Underlying and main causes of death.

Covariate	Whole cohort	CID	Syndromic	PAD	PIRD	Phagocyte	Innate	AIS	Complement	BMF	Phenocopies	other
	N=30628	N=2531	N=4239	N=15123	N=2171	N=2548	N=823	N=1042	N=1482	N=85	N=30	N=554
Underlying cause of death: Infection	989 (32.8%)	302 (51.3%)	192 (31.7%)	232 (22.2%)	99 (29.3%)	89 (34.6%)	47 (54%)	10 (43.5%)	4 (28.6%)	9 (40.9%)	0 (0%)	5 (17.9%)
Underlying cause of death: Malignancy	456 (15.1%)	34 (5.8%)	154 (25.5%)	199 (19%)	28 (8.3%)	22 (8.6%)	8 (9.2%)	0 (0%)	3 (21.4%)	1 (4.5%)	3 (100%)	4 (14.3%)
Underlying cause of death: Immune dysregulation	350 (11.6%)	47 (8%)	54 (8.9%)	90 (8.6%)	122 (36.1%)	16 (6.2%)	7 (8%)	7 (30.4%)	2 (14.3%)	1 (4.5%)	0 (0%)	4 (14.3%)
Underlying cause of death: transplantation	371 (12.3%)	154 (26.1%)	48 (7.9%)	30 (2.9%)	87 (25.7%)	33 (12.8%)	9 (10.3%)	4 (17.4%)	2 (14.3%)	3 (13.6%)	0 (0%)	1 (3.6%)
Underlying cause of death: Other	434 (14.8%)	46 (8.6%)	104 (17.3%)	197 (18.9%)	22 (6.9%)	30 (12.1%)	22 (25.6%)	3 (13%)	1 (7.1%)	5 (22.7%)	0 (0%)	4 (14.8%)
Main cause of death: Septic Shock	405 (13.4%)	82 (13.9%)	75 (12.4%)	127 (12.1%)	37 (10.9%)	38 (14.8%)	33 (37.9%)	8 (34.8%)	3 (21.4%)	2 (9.1%)	0 (0%)	0 (0%)
Main cause of death: Renal Failure	54 (1.8%)	10 (1.7%)	12 (2%)	20 (1.9%)	6 (1.8%)	3 (1.2%)	2 (2.3%)	0 (0%)	0 (0%)	1 (4.5%)	0 (0%)	0 (0%)
Main cause of death: Neurological Complications	183 (6.1%)	28 (4.8%)	24 (4%)	48 (4.6%)	43 (12.7%)	18 (7%)	15 (17.2%)	2 (8.7%)	1 (7.1%)	1 (4.5%)	0 (0%)	3 (10.7%)
Main cause of death: Relapse of Malignancy	200 (6.6%)	17 (2.9%)	55 (9.1%)	89 (8.5%)	16 (4.7%)	12 (4.7%)	5 (5.7%)	0 (0%)	1 (7.1%)	2 (9.1%)	2 (66.7%)	1 (3.6%)
Main cause of death: GvHD	82 (2.7%)	44 (7.5%)	10 (1.7%)	5 (0.5%)	12 (3.6%)	10 (3.9%)	1 (1.1%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Main cause of death: PTLD	25 (0.8%)	10 (1.7%)	7 (1.2%)	1 (0.1%)	5 (1.5%)	1 (0.4%)	1 (1.1%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Main cause of death: Heart Failure	125 (4.2%)	17 (2.9%)	23 (3.8%)	49 (4.7%)	13 (3.8%)	14 (5.4%)	3 (3.4%)	3 (13%)	1 (7.1%)	0 (0%)	2 (66.7%)	0 (0%)
Main cause of death: Multiple Organ Failure	368 (12.2%)	89 (15.1%)	64 (10.6%)	83 (7.9%)	74 (21.9%)	31 (12.1%)	19 (21.8%)	3 (13%)	0 (0%)	4 (18.2%)	0 (0%)	1 (3.6%)
Main cause of death: Surgical Complications	19 (0.6%)	1 (0.2%)	7 (1.2%)	5 (0.5%)	0 (0%)	3 (1.2%)	2 (2.3%)	1 (4.3%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Main cause of death: Veno Occlusive Disorder	22 (0.7%)	11 (1.9%)	2 (0.3%)	0 (0%)	6 (1.8%)	2 (0.8%)	0 (0%)	1 (4.3%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)

Covariate	Whole cohort	CID	Syndromic	PAD	PIRD	Phagocyte	Innate	AIS	Complement	BMF	Phenocopies	other
Main cause of death: Graft Rejection	45 (1.5%)	21 (3.6%)	5 (0.8%)	3 (0.3%)	7 (2.1%)	5 (1.9%)	2 (2.3%)	1 (4.3%)	0 (0%)	0 (0%)	0 (0%)	1 (3.6%)
Main cause of death: Respiratory Failure	640 (21.2%)	156 (26.5%)	174 (28.8%)	160 (15.3%)	70 (20.7%)	49 (19.1%)	10 (11.5%)	6 (26.1%)	3 (21.4%)	6 (27.3%)	3 (100%)	3 (10.7%)
Main cause of death: Haemorrhage	119 (4%)	15 (2.5%)	30 (5%)	36 (3.4%)	20 (5.9%)	7 (2.7%)	6 (6.9%)	3 (13%)	0 (0%)	1 (4.5%)	0 (0%)	1 (3.6%)
Main cause of death: Drug Toxicity	11 (0.4%)	2 (0.3%)	3 (0.5%)	4 (0.4%)	2 (0.6%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Main cause of death: Liver Failure	72 (2.4%)	10 (1.7%)	14 (2.3%)	34 (3.3%)	8 (2.4%)	3 (1.2%)	3 (3.4%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Main cause of death: Thrombosis	5 (0.2%)	1 (0.2%)	0 (0%)	4 (0.4%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Main cause of death: Other	238 (7.9%)	24 (4.1%)	35 (5.8%)	103 (9.8%)	41 (12.1%)	20 (7.8%)	8 (9.2%)	2 (8.7%)	3 (21.4%)	0 (0%)	0 (0%)	2 (7.1%)

Effective (Percentage); the order of subgroups from left to right follows the IEI categories I-X of the 2022 IUIS classification available on the end
date chosen for inclusion with abbreviations for combined immunodeficiencies (CID), CID with syndromic features (Syndromic), primary antibody
deficiencies (PAD), diseases with immune dysregulation or primary immune regulatory disorders (PIRD), diseases of phagocyte number or function
(Phagocyte), disorders of innate immunity (Innate), autoinflammatory syndromes (AIS), complement deficiencies (Complement), bone marrow
failure syndromes (BMF), and phenocopies.

Supplementary Table 3. Therapy.

Covariate	Values	Whole cohort	CID	Syndromic	PAD	PIRD	Phagocyte	Innate	AIS	Complement	BMF	Phenocopies	other
		N=30628	N=2531	N=4239	N=15123	N=2171	N=2548	N=823	N=1042	N=1482	N=85	N=30	N=554
Currently under IgRT	N	13701 (57.5%)	1039 (63.8%)	2280 (74.9%)	4316 (34.9%)	1290 (79.2%)	1864 (96.5%)	525 (82.9%)	770 (93.4%)	1269 (98.2%)	48 (85.7%)	20 (95.2%)	280 (73.9%)
	Y	10107 (42.5%)	589 (36.2%)	764 (25.1%)	8054 (65.1%)	339 (20.8%)	68 (3.5%)	108 (17.1%)	54 (6.6%)	23 (1.8%)	8 (14.3%)	1 (4.8%)	99 (26.1%)
IgRT: route	Intravenous	6410 (51%)	572 (65.3%)	600 (58.8%)	4643 (48.2%)	324 (63.9%)	87 (73.1%)	66 (43.7%)	41 (57.7%)	11 (33.3%)	8 (66.7%)	2 (66.7%)	56 (41.2%)
	Subcutaneous	6119 (48.7%)	297 (33.9%)	416 (40.8%)	4973 (51.6%)	181 (35.7%)	32 (26.9%)	85 (56.3%)	30 (42.3%)	21 (63.6%)	4 (33.3%)	1 (33.3%)	79 (58.1%)
	Intramuscular	17 (0.1%)	1 (0.1%)	2 (0.2%)	12 (0.1%)	1 (0.2%)	0 (0%)	0 (0%)	0 (0%)	1 (3%)	0 (0%)	0 (0%)	0 (0%)
	Push (s.c.)	24 (0.2%)	6 (0.7%)	2 (0.2%)	14 (0.1%)	1 (0.2%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	1 (0.7%)
Age at first IgRT		19.3 [5.2;41.8]	2 [0.5;10.3]	5 [1.3;11.5]	28.6 [8.7;46.7]	10.2 [4.2;19.1]	6.3 [2.1;11.8]	6.7 [2.3;13.3]	5.6 [1.9;15.9]	9.7 [5.6;18]	1.6 [0.7;7.3]	7 [6.1;14]	17.1 [5.9;45.2]
AlloSCT	N	27263 (89%)	1261 (49.8%)	3783 (89.2%)	14907 (98.6%)	1518 (69.9%)	1907 (74.8%)	785 (95.4%)	1010 (96.9%)	1480 (99.9%)	52 (61.2%)	28 (93.3%)	532 (96%)
	Y	3365 (11%)	1270 (50.2%)	456 (10.8%)	216 (1.4%)	653 (30.1%)	641 (25.2%)	38 (4.6%)	32 (3.1%)	2 (0.1%)	33 (38.8%)	2 (6.7%)	22 (4%)
Age at first AlloSCT		2 [0.6;8.1]	0.7 [0.4;1.7]	2.3 [1.1;6.5]	9.5 [3.5;17.7]	2.4 [0.5;8.9]	6.1 [2.2;13.1]	7.8 [3.3;10.7]	4 [1.5;7]	11.4 [10.5;12.3]	10 [5.1;16]	7.9 [6.6;9.2]	7.4 [4.9;14.4]
Gene therapy	N	29070 (99.7%)	2327 (97.9%)	4011 (99.6%)	14465 (100%)	2086 (100%)	2452 (99.6%)	793 (100%)	923 (100%)	1411 (100%)	81 (100%)	30 (100%)	491 (100%)
	Y	77 (0.3%)	50 (2.1%)	17 (0.4%)	0 (0%)	0 (0%)	10 (0.4%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)
Age at first gene therapy		2.3 [0.8;4.6]	0.9 [0.6;2.9]	3.6 [2.8;7.4]	NA [NA;NA]	NA [NA;NA]	15.1 [7.7;22.9]	NA [NA;NA]	NA [NA;NA]	NA [NA;NA]	NA [NA;NA]	NA [NA;NA]	NA [NA;NA]
Splenectomy	N	19732	1565	2681 (96.8%)	9846	1620	1485	603	753 (99.1%)	789 (99.9%)	55	21 (84%)	314 (98.7%)

Covariate	Values	Whole cohort	CID	Syndromic	PAD	PIRD	Phagocyte	Innate	AIS	Complement	BMF	Phenocopies	other
		(97.8%)	(98.1%)		(97.9%)	(94.8%)	(99.3%)	(99.7%)			(100%)		
	Y	453 (2.2%)	30 (1.9%)	90 (3.2%)	216 (2.1%)	89 (5.2%)	10 (0.7%)	2 (0.3%)	7 (0.9%)	1 (0.1%)	0 (0%)	4 (16%)	4 (1.3%)
Age at splenectomy		17.9 [7.3;33.3]	16 [9;24]	5.4 [2.3;8]	31.3 [20.1;42]	11.3 [6.7;20.2]	8.9 [4.5;21.3]	2.8 [2.8;2.8]	21.2 [17.9;36.1]	33.9 [33.9;33.9]	NA [NA;NA]	7.2 [4.6;8.6]	19.3 [15.7;25.3]

175 Quantitative covariates: Median [Q1;Q3]; qualitative covariates: Effective (Percentage)

178 **Supplementary Table 4. IEI/PID diagnoses in patients who died before the age of 5 years in two**
179 **subcategories of IEI/PID.**

IEI/PID diagnoses underlying death before the age of 5 years in the category *diseases with immune dysregulation* (synonymously used as primary immune regulatory disorders, *PIRD*) | *n*=	
Familial hemophagocytic lymphohistiocytosis syndromes (FHLH)*	44
X-linked lymphoproliferative syndrome (XLP)*	10
FOXP3 deficiency (IPEX)	4
Autoimmune lymphoproliferative syndrome (ALPS)	3
Early-onset inflammatory bowel disease	2
Griscelli syndrome, type 2*	2
Unclassified disorders of immune dysregulation	2
APECED / APS1 with CMC - Autoimmune polyendocrinopathy candidiasis ectodermal dystrophy (APECED)	1
Chediak Higashi syndrome*	1
IPEX-like disease	1
*proportion of patients with risk of hemophagocytic lymphohistiocytosis (HLH)	81·4%
IEI/PID diagnoses underlying death before the age of 5 years in category *disorders of innate or intrinsic immunity (*abbreviated *Innate) | n*=	
Defects of TLR/NFkappa-B signalling	7
Isolated congenital asplenia	7
Defects with susceptibility to mycobacterial infection (MSMD)	4
ZNFX1 deficiency	2
Asplenia syndrome (Ivemark syndrome)	1
Chronic mucocutaneous candidiasis (CMC)	1
IRF9 deficiency	1
Unclassified defects in innate immunity	1

182   **Supplementary Table 5. Institutions of participants of the 2024 ESID-R survey on the**
183   **international IEI/PID registry landscape.**

Western Europe	Eastern Europe	Other countries
Denmark (Copenhagen)	Armenia (Yerevan)	Australia (Melbourne)
Finland (Helsinki)	Belarus (Minsk)	Brazil (São Paulo)
Germany (Dresden, Freiburg, Leipzig, Munich)	Bulgaria (Sofia)	Canada (Halifax)
Ireland (Dublin)	Czech Republic (Prague)	Egypt (Cairo)
Italy (Milan)	Georgia (Tbilisi)	Iran (Teheran)
Portugal (Coimbra)	Hungary (Budapest)	Sudan (Khartoum)
Spain (Barcelona)	Latvia (Riga)	Turkey (Istanbul)
Sweden (Jönköping)	Romania (Bucharest)	USA (Philadelphia)
Switzerland (Geneva, Zurich)	Russia (Moscow)	
United Kingdom (Sheffield)	Serbia (Belgrade)	
	Ukraine (Kyiv)	

184   *The survey participants chose to reply (or not) to a survey which had been distributed via the ESID-R*
185   *listserv and was also posted on the website during the summer of 2024.*

187
188

Supplementary Material (Figures)

Supplementary Figure 1A. Cumulative number of patients registered in the ESID-R.

The total number shown here is 31,889 patients (male n=17,515; female n=14,363; unknown n=11), before exclusion of those discharged (n=538), those with secondary immunodeficiency (n=417) and those without a definitive diagnosis of IEI/PID (n=306). Data shown for the **online** Registry only. Data from the very first version (hardcopy) was not transferred to the first online registry, but newly entered.

According to "The Source https://esid.org/wp-content/uploads/2024/03/ESID_TheSource_2003.pdf" the total number in 2003 was 9707 pts.

Supplementary Figure 1B. Cumulative number of ESID-R participating centers.

202 **Supplementary Figure 2. IEI/PID minimal prevalence map according to ESID-R patient numbers and**
203 **the 2019 population (same data as in Supplementary Table 1).**

Prevalence map of PID
for 100,000 inhabitants (based on 2019 population)

Supplementary Figure 3. IEI/PID distribution, interactive nested pie chart (see https://esid.org/html-pages/Suppl%20Fig%203_ESID_30k_sunburst_PID.html).

210 **Supplementary Figure 4. Survival probabilities of IEI/PID until 5 years of age as documented in the**
211 **ESID-R.**

	Number at risk					
CID	113	256	223	231	236	247
Synd.	286	1114	1243	1408	1488	1572
PAD	78	598	1078	1447	1880	2268
PIRD	29	201	238	266	297	347
Phag.	80	646	781	844	885	882
Innate	13	124	176	211	224	231
AIS	6	77	127	193	250	280
Compl.	8	61	110	142	171	210

213 As defined in the main manuscript, the term PIRD refers to all disorders with immune dysregulation,
214 as in IUIS category IV, including hemophagocytic syndromes.

216 **Supplementary Figure 5.** Countries with national patient registries or national sub-registries within
217 the ESID-R according to the 2024 ESID survey (all participating IEI/PID centers worldwide are listed in
218 Suppl. Table 5).

Supplementary References

1. Scheible R, Rusch S, Guzman D, Mahlaoui N, Ehl S, Kindle G. The NEW ESID online database network. *Bioinformatics* 2019;**35**(24):5367-69 doi: 10.1093/bioinformatics/btz525.

2. Alligon M, Mahlaoui N, Bouaziz O. Pitfalls in time-to-event analysis of registry data: a tutorial based on simulated and real cases. *Front Epidemiol* 2024;**4**:1386922 doi: 10.3389/fepid.2024.1386922.

3. Abd Hamid IJ, Azman NA, Gennery AR, Mangantig E, Hashim IF, Zainudeen ZT. Systematic Review of Primary Immunodeficiency Diseases in Malaysia: 1979-2020. *Frontiers in immunology* 2020;**11**:1923 doi: 10.3389/fimmu.2020.01923.

4. Aghamohammadi A, Moein M, Farhoudi A, Pourpak Z, Rezaei N, Abolmaali K, Movahedi M, Gharagozlou M, Ghazi BM, Mahmoudi M, Mansouri D, Arshi S, Trash NJ, Akbari H, Sherkat R, Hosayni RF, Hashemzadeh A, Mohammadzadeh I, Amin R, Kashef S, Alborzi A, Karimi A, Khazaei H. Primary immunodeficiency in Iran: first report of the National Registry of PID in Children and Adults. *Journal of clinical immunology* 2002;**22**(6):375-80 doi: 10.1023/a:1020660416865.

5. Al-Herz W, Al-Ahmad M, Al-Khabaz A, Husain A, Sadek A, Othman Y. The Kuwait National Primary Immunodeficiency Registry 2004-2018. *Frontiers in immunology* 2019;**10**:1754 doi: 10.3389/fimmu.2019.01754.

6. El-Helou SM, Biegner AK, Bode S, Ehl SR, Heeg M, Maccari ME, Ritterbusch H, Speckmann C, Rusch S, Scheible R, Warnatz K, Atschekzei F, Beider R, Ernst D, Gerschmann S, Jablonka A, Mielke G, Schmidt RE, Schurmann G, Sogkas G, Baumann UH, Klemann C, Viemann D, von Bernuth H, Kruger R, Hanitsch LG, Scheibenbogen CM, Wittke K, Albert MH, Eichinger A, Hauck F, Klein C, Rack-Hoch A, Sollinger FM, Avila A, Borte M, Borte S, Fasshauer M, Hauenherm A, Kellner N, Muller AH, Ulzen A, Bader P, Bakhtiar S, Lee JY, Hess U, Schubert R, Wolke S, Zielen S, Ghosh S, Laws HJ, Neubert J, Oommen PT, Honig M, Schulz A, Steinmann S, Schwarz K, Duckers G, Lamers B, Langemeyer V, Niehues T, Shai S, Graf D, Muglich C, Schmalzing MT, Schwaneck EC, Tony HP, Dirks J, Haase G, Liese JG, Morbach H, Foell D, Hellige A, Wittkowski H, Masjosthusmann K, Mohr M, Geberzahn L, Hedrich CM, Muller C, Rosen-Wolff A, Roesler J, Zimmermann A, Behrends U, Rieber N, Schauer U, Handgretinger R, Holzer U, Henes J, Kanz L, Boesecke C, Rockstroh JK, Schwarze-Zander C, Wasmuth JC, Dilloo D, Hulsmann B, Schonberger S, Schreiber S, Zeuner R, Ankermann T, von Bismarck P, Huppertz HI, Kaiser-Labusch P, Greil J, Jakoby D, Kulozik AE, Metzler M, Naumann-Bartsch N, Sobik B, Graf N, Heine S, Kobbe R, Lehmberg K, Muller I, Herrmann F, Horneff G, Klein A, Peitz J, Schmidt N, Bielack S, Gross-Wieltsch U, Classen CF, Klasen J, Deutz P, Kamitz D, Lassay L, Tenbrock K, Wagner N, Bernbeck B, Brummel B, Lara-Villacanas E, Munstermann E, Schneider DT, Tietsch N, Westkemper M, Weiss M, Kramm C, Kuhnle I, Kullmann S, Girschick H, Specker C, Vinnemeier-Laubenthal E, Haenicke H, Schulz C, Schweigerer L, Muller TG, Stiefel M, Belohradsky BH, Soetedjo V, Kindle G, Grimbacher B. The German National Registry of Primary Immunodeficiencies (2012-2017). *Frontiers in immunology* 2019;**10**:1272 doi: 10.3389/fimmu.2019.01272.

7. group CTFPs. The French national registry of primary immunodeficiency diseases. *Clinical immunology* 2010;**135**(2):264-72 doi: 10.1016/j.clim.2010.02.021.

8. Lougaris V, Pession A, Baronio M, Soresina A, Rondelli R, Gazzurelli L, Benvenuto A, Martino S, Gattorno M, Biondi A, Zecca M, Marinoni M, Fabio G, Aiuti A, Marseglia G,

Putti MC, Agostini C, Lunardi C, Tommasini A, Bertolini P, Gambineri E, Consolini R, Matucci A, Azzari C, Danieli MG, Paganelli R, Duse M, Cancrini C, Moschese V, Chessa L, Spadaro G, Civino A, Vacca A, Cardinale F, Martire B, Carpino L, Trizzino A, Russo G, Cossu F, Badolato R, Pietrogrande MC, Quinti I, Rossi P, Ugazio A, Pignata C, Plebani A. The Italian Registry for Primary Immunodeficiencies (Italian Primary Immunodeficiency Network; IPINet): Twenty Years of Experience (1999-2019). *Journal of clinical immunology* 2020;**40**(7):1026-37 doi: 10.1007/s10875-020-00844-0.

9. Ludviksson BR, Sigurdardottir ST, Johannsson JH, Haraldsson A, Hardarson TO. Epidemiology of Primary Immunodeficiency in Iceland. *Journal of clinical immunology* 2015;**35**(1):75-9 doi: 10.1007/s10875-014-0107-3.

10. Massaad MJ, Zainal M, Al-Herz W. Frequency and Manifestations of Autoimmunity Among Children Registered in the Kuwait National Primary Immunodeficiency Registry. *Frontiers in immunology* 2020;**11**:1119 doi: 10.3389/fimmu.2020.01119.

11. Matamoros Flori N, Mila Llambi J, Espanol Boren T, Raga Borja S, Fontan Casariego G. Primary immunodeficiency syndrome in Spain: first report of the National Registry in Children and Adults. *Journal of clinical immunology* 1997;**17**(4):333-9 doi: 10.1023/a:1027382916924.

12. Mellouli F, Mustapha IB, Khaled MB, Besbes H, Ouederni M, Mekki N, Ali MB, Largueche B, Hachicha M, Sfar T, Gueddiche N, Barsaoui S, Sammoud A, Boussetta K, Becher SB, Meherzi A, Guandoura N, Boughammoura L, Harbi A, Amri F, Bayoudh F, Jaballah NB, Tebib N, Bouaziz A, Mahfoudh A, Aloulou H, Mansour LB, Chabchoub I, Boussoffara R, Chemli J, Bouguila J, Hassayoun S, Hammami S, Habboul Z, Hamzaoui A, Ammar J, Barbouche MR, Bejaoui M. Report of the Tunisian Registry of Primary Immunodeficiencies: 25-Years of Experience (1988-2012). *Journal of clinical immunology* 2015;**35**(8):745-53 doi: 10.1007/s10875-015-0206-9.

13. Michos A, Raptaki M, Tantou S, Tzanoudaki M, Spanou K, Liatsis M, Constantinidou N, Paschali E, Varela I, Moraloglou O, Bakoula C, Kanariou M. Primary immunodeficiency diseases: a 30-year patient registry from the referral center for primary immunodeficiencies in Greece. *Journal of clinical immunology* 2014;**34**(7):836-43 doi: 10.1007/s10875-014-0066-8.

14. Moore A, Blumenthal KG, Chambers C, Namazy J, Nowak-Wegrzyn A, Phillips EJ, Rider NL. Improving Clinical Practice Through Patient Registries in Allergy and Immunology. *J Allergy Clin Immunol Pract* 2024;**12**(10):2599-609 doi: 10.1016/j.jaip.2024.05.003.

15. Mukhina AA, Kuzmenko NB, Rodina YA, Kondratenko IV, Bologov AA, Latysheva TV, Prodeus AP, Pampura AN, Balashov DN, Ilyina NI, Latysheva EA, Deordieva EA, Shvets OA, Deripapa EV, Abramova IN, Pashenko OE, Vahlyarskaya SS, Zinovyeva NV, Zimin SB, Skorobogatova EV, Machneva EB, Fomina DS, Ipatova MG, Barycheva LY, Khachirova LS, Tuzankina IA, Bolkov MA, Shakhova NV, Kamaltynova EM, Sibgatullina FI, Guseva MN, Kuznetsova RN, Milichkina AM, Totolian AA, Kalinina NM, Goltsman EA, Sulima EI, Kutlyanceva AY, Moiseeva AA, Khoreva AL, Nesterenko Z, Tymofeeva EV, Ermakova A, Proligina DD, Kalmetieva LR, Davletbaieva GA, Mirsayapova IA, Richkova OA, Kuzmicheva KP, Grakhova MA, Yudina NB, Orlova EA, Selezneva OS, Piskunova SG, Samofalova TV, Bukina TV, Pechkurova AD, Migacheva N, Zhestkov A, Barmina EV, Parfenova NA, Isakova SN, Averina EV, Sazonova IV, Starikova SY, Shilova TV, Asekretova TV, Suprun RN, Kleshchenko EI, Lebedev VV, Demikhova EV, Demikhov VG, Kalinkina VA, Gorenkova AV, Duryagina SN, Pavlova TB, Shinkareva VM, Smoleva IV,

Aleksandrova TP, Bambaeva ZV, Philippova MA, Gracheva EM, Tcyvkina GI, Efremenkov AV, Mashkovskaya D, Yarovaya IV, Alekseenko VA, Fisyun IV, Molokova GV, Troitskya EV, Piatkina LI, Vlasova EV, Ukhanova O, Chernishova EG, Vasilieva M, Laba OM, Volodina E, Safonova EV, Voronin KA, Gurkina MV, Rumyantsev AG, Novichkova GA, Shcherbina AY. Primary Immunodeficiencies in Russia: Data From the National Registry. *Frontiers in immunology* 2020;**11**:1491 doi: 10.3389/fimmu.2020.01491.

16. Pickett G, Motazedi T, Kutac C, Cahill G, Cunnigham-Rundles C, Fuleihan RL, Sullivan KE, Rider NL. Infection Phenotypes Among Patients with Primary Antibody Deficiency Mined from a US Patient Registry. *Journal of clinical immunology* 2021;**41**(2):374-81 doi: 10.1007/s10875-020-00916-1.

17. Rhim JW, Kim KH, Kim DS, Kim BS, Kim JS, Kim CH, Kim HM, Park HJ, Pai KS, Son BK, Shin KS, Oh MY, Woo YJ, Yoo Y, Lee KS, Lee KY, Lee CG, Lee JS, Chung EH, Choi EH, Hahn YS, Park HY, Kim JG. Prevalence of primary immunodeficiency in Korea. *J Korean Med Sci* 2012;**27**(7):788-93 doi: 10.3346/jkms.2012.27.7.788.

18. Ruffner MA, Group UBW, Sullivan KE. Complications Associated with Underweight Primary Immunodeficiency Patients: Prevalence and Associations Within the USIDNET Registry. *Journal of clinical immunology* 2018;**38**(3):283-93 doi: 10.1007/s10875-018-0492-0.

19. Volokha A, Bondarenko A, Chernyshova L, Hilfanova A, Stepanovskiy Y, Boyarchuk O, Kostyuchenko L. Impact of the J Project on progress of primary immunodeficiency care in Ukraine. *Cent Eur J Immunol* 2021;**46**(2):250-57 doi: 10.5114/ceji.2021.108183.

20. Seminario G, Gonzalez-Serrano ME, Aranda CS, Grumach AS, Segundo GRS, Regairaz L, Cardona AA, Group LR, Becerra JCA, Poli C, King A, Fernandes FR, Leiva L, Franco JL, Espinosa-Rosales FJ, Sorensen R, Costa Carvalho BT, Bezrodnik L, Condino-Neto A. The Latin American Society for Immunodeficiencies Registry. *Journal of clinical immunology* 2024;**45**(1):28 doi: 10.1007/s10875-024-01822-6.

21. Abolhassani H, Azizi G, Sharifi L, Yazdani R, Mohsenzadegan M, Delavari S, Sohani M, Shirmast P, Chavoshzadeh Z, Mahdaviani SA, Kalantari A, Tavakol M, Jabbari-Azad F, Ahanchian H, Momen T, Sherkat R, Sadeghi-Shabestari M, Aleyasin S, Esmaeilzadeh H, Al-Herz W, Bousfiha AA, Condino-Neto A, Seppanen M, Sullivan KE, Hammarstrom L, Modell V, Modell F, Quinn J, Orange JS, Aghamohammadi A. Global systematic review of primary immunodeficiency registries. *Expert Rev Clin Immunol* 2020;**16**(7):717-32 doi: 10.1080/1744666X.2020.1801422.

22. Chapel H, Prevot J, Gaspar HB, Espanol T, Bonilla FA, Solis L, Drabwell J, Editorial Board for Working Party on Principles of Care at I. Primary immune deficiencies - principles of care. *Frontiers in immunology* 2014;**5**:627 doi: 10.3389/fimmu.2014.00627.

23. Kohler S, Gargano M, Matentzoglu N, Carmody LC, Lewis-Smith D, Vasilevsky NA, Danis D, Balagura G, Baynam G, Brower AM, Callahan TJ, Chute CG, Est JL, Galer PD, Ganesan S, Griese M, Haimel M, Pazmandi J, Hanauer M, Harris NL, Hartnett MJ, Hastreiter M, Hauck F, He Y, Jeske T, Kearney H, Kindle G, Klein C, Knoflach K, Krause R, Lagorce D, McMurry JA, Miller JA, Munoz-Torres MC, Peters RL, Rapp CK, Rath AM, Rind SA, Rosenberg AZ, Segal MM, Seidel MG, Smedley D, Talmy T, Thomas Y, Wiafe SA, Xian J, Yuksel Z, Helbig I, Mungall CJ, Haendel MA, Robinson PN. The Human Phenotype Ontology in 2021. *Nucleic Acids Res* 2021;**49**(D1):D1207-D17 doi: 10.1093/nar/gkaa1043.

24. Haimel M, Pazmandi J, Heredia RJ, Dmytrus J, Bal SK, Zoghi S, van Daele P, Briggs TA,
Wouters C, Bader-Meunier B, Aeschlimann FA, Caorsi R, Eleftheriou D, Hoppenreijs
E, Salzer E, Bakhtiar S, Derfalvi B, Saettini F, Kusters MAA, Elfeky R, Truck J,
Riviere JG, van der Burg M, Gattorno M, Seidel MG, Burns S, Warnatz K, Hauck F,
Brogan P, Gilmour KC, Schuetz C, Simon A, Bock C, Hambleton S, de Vries E,
Robinson P, van Gijn M, Boztug K. Curation and Expansion of Human Phenotype
Ontology for Defined Groups of Inborn Errors of Immunity. *The Journal of allergy and
clinical immunology* 2021 doi: 10.1016/j.jaci.2021.04.033.

25. Jacobsen JOB, Baudis M, Baynam GS, Beckmann JS, Beltran S, Buske OJ, Callahan
TJ, Chute CG, Courtot M, Danis D, Elemento O, Essenwanger A, Freimuth RR,
Gargano MA, Groza T, Hamosh A, Harris NL, Kaliyaperumal R, Lloyd KCK, Khalifa A,
Krawitz PM, Kohler S, Laraway BJ, Lehvaslaiho H, Matalonga L, McMurry JA, Metke-
Jimenez A, Mungall CJ, Munoz-Torres MC, Ogishima S, Papakonstantinou A, Piscia
D, Pontikos N, Queralt-Rosinach N, Roos M, Sass J, Schofield PN, Seelow D, Siapos
A, Smedley D, Smith LD, Steinhaus R, Sundaramurthi JC, Swietlik EM, Thun S,
Vasilevsky NA, Wagner AH, Warner JL, Weiland C, Consortium GPM, Haendel MA,
Robinson PN. The GA4GH Phenopacket schema defines a computable
representation of clinical data. *Nat Biotechnol* 2022;**40**(6):817-20 doi:
10.1038/s41587-022-01357-4.

26. Danis D, Bamshad MJ, Bridges Y, Cacheiro P, Carmody LC, Chong JX, Coleman B,
Dalgleish R, Freeman PJ, Graefe ASL, Groza T, Jacobsen JOB, Klocperk A, Kusters
M, Ladewig MS, Marcello AJ, Mattina T, Mungall CJ, Munoz-Torres MC, Reese JT,
Rehburg F, Reis BCS, Schuetz C, Smedley D, Strauss T, Sundaramurthi JC, Thun S,
Wissink K, Wagstaff JF, Zocche D, Haendel MA, Robinson PN. A corpus of GA4GH
Phenopackets: case-level phenotyping for genomic diagnostics and discovery.
medRxiv 2024 doi: 10.1101/2024.05.29.24308104.