Epigenetic entropy, social disparity, and health and lifespan in the Women's

Health Initiative

Khyobeni Mozhui[1,2,*], Athena Starlard-Davenport[2], Yangbo Sun[1], Aladdin H. Shadyab[3], Ramon

Casanova[4], Fridtjof Thomas[1], Robert B Wallace[5], Jay H Fowke[1], Karen C Johnson[1]

[1]Department of Preventive Medicine, College of Medicine, University of Tennessee Health

Science Center, Memphis, TN, USA.

[2]Department of Genetics, Genomics and Informatics, College of Medicine, University of

Tennessee Health Science Center, Memphis, TN, USA.

[3]Herbert Wertheim School of Public Health and Human Longevity Science and Division of

Geriatrics, Gerontology, and Palliative Care, Department of Medicine, University of California

San Diego, La Jolla, CA, USA.

[4]Department of Biostatistics and Data Science, Wake Forest University School of Medicine,

Winston-Salem, NC, USA.

[5]College of Public Health, University of Iowa, Iowa City, IA, USA.

[*]Corresponding author: kmozhui@uthsc.edu

ABSTRACT

The pace of aging varies between individuals and is marked by changes in DNA methylation (DNAm) including an increase in randomness or entropy. Here, we computed epigenetic scores of aging and entropy using DNAm datasets from the Women's Health Initiative (WHI). We investigated how different epigenetic aging metrics relate to demographic and health variables, and mortality risk. Income and education, two proxies of socioeconomics (SE), had consistent associations with epigenetic aging and entropy. Notably, stochastic increases in DNAm at sites targeted by the polycomb proteins were significantly related to both aging and SE. While higher income was associated with reduced age-related DNAm changes in White women, the protective effect of income was diminished in Black and Hispanic women, and on average, Black and Hispanic women had relatively more aged epigenomes. Faster pace of aging, as estimated by the DunedinPACE, predicted higher mortality risk, while the maintenance of methylation at enhancer regions was associated with improved survival. Our findings demonstrate close ties between social and economic factors and aspects of epigenetic aging, suggesting potential biological mechanisms through which societal disparities may contribute to differences in health outcomes and lifespan across demographic groups.

INTRODUCTION

Epigenetic clocks such as the Horvath, Hannum, PhenoAge , and DunedinPACE are machine-learning (ML) based predictive models that estimate the biological age, or the rate-of-aging, of an individual.[1-4] The underlying biomolecular data is DNA methylation (DNAm), the epigenetic process that entails a chemical modification to the DNA—specifically, whether a cytosine residue that is adjacent to a guanine (a CpG site) has a methyl- tag or is unmethylated. These epigenetic markers can be measured from readily accessible biofluids such as blood or saliva and are informative of whether a person is aging faster or slower relative to their chronological age. The first-generation epigenetic clocks (e.g., Horvath, Hannum) are accurate predictors of chronological age because the algorithms were mainly trained on the age variable.[1,2] The second-generation clocks (e.g., PhenoAge, GrimAge) were trained on a broader panel of age-related health parameters and perform remarkably well at assessing overall health, disease states, and life expectancy.[3,5,6] The more recent DunedinPACE model was developed by tracking longitudinal changes in health traits from age-matched individuals over decades, and it directly measures the pace-of-aging.[4]

These existing models estimate biological age using preselected sets of CpGs that were picked by the respective ML models and assigned specific weights, and the final value of aging is a weighted combination of methylation levels at these distinct CpGs sites. The lists of preselected CpGs are typically unique for each model, and it is unclear as to why the "mind of the algorithm" selected those specific sites and were assigned those weights.[7-9] While there are now several such ML-based biomarkers that are widely utilized in epidemiological research, the mechanistic pathways and the extent to which these represent stochastic versus programmatic

processes of aging remain largely undefined.[10,11] However, several of the DNAm clocks do share

some common themes that hint at the underlying biology. For instance, relative to the genome-

wide background, bivalent chromatin states and DNA sequences bound by the polycomb

repressive complex (particularly PRC2) are consistently overrepresented among both the first-

and second-generation clocks.[3,6,12] The bivalent and PRC2 bound sequences are evolutionarily

conserved and play critical roles during embryonic development and cell fate determination.[13]

A common feature of aging is for CpGs in these chromatin states to drift from a low methylation

to a high methylation state, and the gain in methylation with aging has been implicated in

cancers and other health risks.[14] This is closely related to the concept of epigenetic entropy, a

term borrowed from Shannon's Information theory, that refers to the level of randomness that

can be computed from DNAm data either at the global genome-wide scale, or at regional

scale.[14-16] These stereotypical shifts in the epigenetic landscape and the accumulation of

stochastic variability likely contribute to some of the signal captured by the DNAm aging

biomarkers.[9,14,16,17]

The Women's Health Initiative (WHI) is a long-term prospective study of postmenopausal

women that began recruitment in 1993 and has been collecting extensive health, medical, and

lifespan data since that time.[18-20] Previous epigenetic studies of DNAm clocks in the WHI have

shown that a higher rate of epigenetic aging is predictive of lifespan. Prior studies also report

associations with lung cancer,[21] insomnia and immune aging,[22] cognitive impairment and

dementia,[23] diabetes related traits, and cardiovascular health.[24-27] Notably, the second-

generation PhenoAge clock has strong associations with social disparities, and WHI participants

with lower education exhibited more advance epigenetic aging. Liu et al. also found that an

accelerated PhenoAge partly explain the disparity in life expectancy between racial and ethnic

groups.[27] There is growing evidence that these epigenetic biomarkers not only tell us about the

intrinsic aging of cells but are strongly influenced by the larger social and environmental

context. In the United States, recent studies in young as well as older adults have shown that

the DunedinPACE detects more rapid aging among Black participants and among participants

with low education and income levels.[28,29] This sensitivity to social factors should not be

entirely surprising since variables such as education, income inequity, and experiences of

societal biases have profound and long-lasting impact on health, stress, and mental and

emotional well-being, and are likely linked to the differential rates of biological aging. In fact,

these close ties between the epigenetic models of aging and social variables have been

interpreted in light of the "weathering hypothesis", which proposes that the chronic exposure

to socioeconomic disadvantage, and to race-based and other stressors lead to accelerated

aging and higher disease burden among African Americans and other marginalized groups.[27-30]

However, a caveat to keep in mind is that these ML-based epigenetic biomarkers were initially

trained in cohorts that were predominantly, or in the case of the DunedinPACE, almost

exclusively of European ancestry.[4] The training conditions likely introduce some algorithmic

biases and may not generalize in full to other populations. In the present work, we attempt to

unlink the epigenetic readouts from potential training-based effects by computing measures of

epigenetic entropy and variability in CpG methylation that do not rely on training algorithms.

For the present study, the only pre-selection or sub-setting of CpGs we performed was based

on biologically informed chromatin states. The questions we ask are: (1) *how do these readouts*

of epigenetic entropy and stochasticity, and gross chromatin states relate to the training-based

models of aging? And, *are these non-ML readouts as informative of health and socioeconomic*

variables as the ML-based models, and predictive of life expectancy? Notably, we present novel

evidence linking measures of epigenetic stochasticity with social and racial disparities.

Furthermore, we uncover intriguing connections between methylation maintenance at active

enhancer regions and lifespan.

RESULTS

Datasets and participant characteristics

The present work is a secondary analysis of DNAm data generated by three WHI ancillary

studies: (1) WHI-EMPC/AS315, [26,31] (2) BA23, [21,32] and (3) AS311.[33] All three datasets measured

blood DNA methylation using the Illumina Infinium Methylation 450K BeadChip (Illumina Inc.).

Brief summaries of baseline characteristics are provided in **Table 1**, and sample

inclusion/exclusion chart is in **Fig S1**. BA23 and AS311 used blood collected at the time of

eligibility screening visit (SV) and prior to randomization to a study arm in the clinical trials (CT)

or enrollment in the observational study (OS). In AS315, DNA methylation was measured either

at SV (64%), annual visit 3 (29%), or annual visit 6 (7%) in the CT only. We used the larger and

more diverse AS315 dataset to describe the large-scale topology of the methylome, and all

analyses were initially performed in AS315 (including a sensitivity analysis using only blood

collected at the SV). The main findings were then tested for replication in BA23 and AS311 after

excluding participants who were also part of AS315 (**Fig S1**).

Epigenetic stochasticity and how it relates to chromatin states and aging

The methylation beta-values show the expected bimodal distribution with most CpGs at beta-

values close to 0 (i.e., most cells are unmethylated at that CpG) or 1 (most cell are methylated

at that CpG) (**Fig 1a**). With aging, cells begin to drift from their initial methylation states,[34] and

this results in an increase in entropy and a shift towards a beta = 0.5 (a "hemi-methylated"

state). We computed the methylome-wide entropy using all CpGs that had complete data in all

participants. To illustrate the "landscape erosion" towards a hemi-methylated and presumably

more random state (i.e., beta ~ 0.5), **Fig 1a** displays 5 participants with Shannon entropy > 0.85,

and 5 with entropy <0.4. This global measure of epigenetic discordance showed a wide

variation in the AS315 cohort (**Fig 1b**) but had only a weak positive correlation with age (r=0.08,

p = 0.0002; **Fig 1c**).

As methylation levels are highly dependent on the genomic context, we annotated each CpG

for the predicted chromatin state based on the Roadmap Epigenomics Consortium's

ChromHMM 15-states model.[35-38] This overlays the DNAm data with histone-based predicted

epigenetic states and regulatory factors (**Table 2**).[35] For each participant, we computed the

average beta-values at each of the 15 states thereby reducing the ~450,000 features to 15

values. The mean methylation at the chromatin states showed varying levels of correlation with

chronological age (**Fig 1d**; **Table 2**). The highest positive correlate of age was the repressed

bivalent transcription start site (TssBiv; r = 0.16, **Fig S2a**), and the highest negative correlate was

for enhancer regions (Enh; r = -0.19, **Fig 1f**).

Whether a CpG positively or negatively correlates with chronological age is partly dependent on

the average methylation of its chromatin state, and this is depicted in **Fig 1e**. For instance, CpGs

Fig 1. (a) Density plots of methylome-wide beta-values for 10 AS315 participants (all identified as non-Hispanic White). 5 have entropy ≤0.4 (cyan), and 5 have entropy ≥0.85 (soft red). **(b)** Histogram of methylome-wide entropy shows wide variability in AS315. **(c)** Weak but positive correlation between chronological age and methylome entropy in AS315. **(d)** Pair-wise correlations between age and the different non-training based epigenetic readouts including mean methylation by chromatin states (due to the large number of comparisons, the graph only displays correlations with p < 0.001). **(e)** The x-axis is the mean beta-values for the 15 chromatin states; y-axis is the Pearson r between these states and chronological age. Mean methylation at enhance CpGs decrease **(f)**, while levels of stochastic epimutations at the three bivalent sites (TssBiv, BivFlnk, EnhBiv) increase with chronological age **(g)**. **(h)** Pair-wise correlation for the non-training based measures (overall entropy, stochasticity at bivalent sites, and average beta-values for the bivalent CpGs) and four train-based measures of aging (only displays correlations with p < 0.001). All these are residual values adjusted for chronological age.

at the bivalent sites—TssBiv, BivFlnk, and EnhBiv— are known to have low mean methylation

and gain methylation with aging, and this pattern is also seen in the WHI (plots to the top left

quadrant of **Fig 1e**).[39] These bivalents states also had the highest positive r with entropy (e.g.,

Fig S2c). For this reason, we explored the idea of "stochastic epimutations" (SEpiM) at these

sites. We implemented an outlier detection approach,[40,41] and counted for positive outliers at

TssBiv, BivFlnk, and EnhBiv. As these CpGs gain methylation with aging, a positive outlier with

higher beta-value is presumed to represent a more aged methylome. The outlier counts (what

we refer to as BivSEpiM) showed strong positive correlation with chronological age (**Fig 1g**). In

contrast to the bivalent sites, CpGs that flank actively transcribed genes (TxFlnk) had mean

methylation of beta-value ~0.5 and had the strongest negative correlation with entropy **(Fig**

S2d) and had modest inverse correlation with age (bottom right quadrant in **Fig 1e**). Weakly

repressed polycomb (ReprPCWk) and Quiescent (Quies) were also regions with higher mean

methylation (beta-values >0.70) and negatively correlated with both entropy and age.

Collectively, the results indicate that CpGs in low methylation sites (e.g., TssA, TssBiv, etc.) gain

methylation with aging, and contribute to higher entropy. In contrast, CpGs in sites with higher

mean methylation (TxFlnk, ReprPCWk and Quies) tend to lose methylation with aging, and

methylation maintenance at these sites may contribute to lower entropy.

Relating epigenetic stochasticity to training-based DNAm biomarkers

To evaluate how the non-training based DNAm readouts relate to the training-based models of

biological aging, we compared pace or rates of biological aging (i.e., age-deviation or age-

acceleration) estimated by the original Horvath and Hannum clocks, and the newer PhenoAge

and DunedinPACE.[1-4] After adjusting for chronological age, higher BivSEpiM was consistently

associated with a higher pace of aging as defined by the prior models (**Fig 1h**). The methylome-

wide entropy was positively correlated with higher age-acceleration as estimated by the

Horvath or Hannum clocks (**Fig 1h**). This indicates that the overall buildup of discordance in the

methylome partly, but not fully, contributes to these training-based biomarkers of aging.

Epigenetic stochasticity and social and health variables at baseline

We treated the age-regressed residuals of the DNAm readouts (i.e., age-deviation) as outcome

variables, and explored associations with baseline health and social variables. For rigor, we

performed the analysis with multiple regression models in the AS315 that included a sensitivity

analysis; for nominally significant associations (uncorrected $p \leq 0.05$), we tested for replication

in BA23 and AS311. Regarding the interpretation of age-deviation, as all the DNAm readouts

listed in **Table 3** increase in value with aging, a positive age-deviation indicates higher rate of

aging, and negative indicates slower rate of aging. Model 1 was a multivariable regression with

race/ethnicity, socioeconomic index (SE index, a combination of self-reported income and

education), smoking status, body weight index (BMI), and total weekly recreation energy

expenditure (TEXPWK; MET-hours/week) as predictors. To make the magnitudes of effects

comparable, **Fig 2a** displays the standardized regression estimates. For the three non-training-

based readouts (i.e., entropy, BivSEpiM, and BivCpGAvg), the most consistent association was

with SE index (**Table 3**). SE index also had significant negative associations with the

DunedinPACE and the PhenoAge clock (**Fig 2a**; unstandardized regression estimates in **Data**

S1.). A unit increase in SE index resulted in -0.004 lower entropy, -0.02 lower BivSEpiM, and -

0.005 slower DunedinPACE of aging (**Table 3**). The epigenetic clocks, which are in units of years

    provide a more intuitive interpretation. For the PhenoAge, one unit increase in SE index

Fig 2. (a) Forest plots of standardized regression coefficients (95% confidence intervals) for the baseline predictor variables used in Model 1 for the AS315 group. The outcome variables are the age-adjusted DNAm readouts. Smoking1 and Smoking2 are past and present smokers, respectively, and regressions are related to never smokers. Estimates for race/ethnicity are relative to non-Hispanic White. (b) Bar graphs of age-adjusted entropy, BivSEpiM, BivAvg, and DunedinPACE plotted separately by self-reported race/ethnicity, and overlaid by income groups in AS315, and (c) in BA23 (bars shades denote income levels). Positive values indicate higher biological aging relative to chronological age. Error bars are standard errors.

resulted in roughly 0.25 years lower biological age relative to chronological age. We note that

the bivalent readouts and DunedinPACE showed higher aging rates in all other race/ethnicity

groups relative to White participants. Consistent with previous reports, the Hannum clock

detected a significantly slower rate of aging among the Black participants.[42]

We computed these same readouts in BA23, and in the much smaller AS311. After excluding

participants who were also part of the AS315, the negative associations between SE index and

bivalent readouts and entropy remained consistent in both data sets (**Figs S3** and **Data S2, S3**).

For example, in BA23, a unit increase in SE index resulted in 0.005 lower entropy, and 0.001

lower beta BivCpGAvg, a 0.009 slower DunedinPACE per chronological age, and a 0.46 years

younger biological age relative to chronological age for the PhenoAge (**Table 3**). To illustrate the

gradational effect of income on these measures of aging, we display bar plots of the non-

training epigenetic readouts and the DunedinPACE (after adjusting for chronological age) by

self-reported income for AS315 and BA23 (**Fig 2b, c**). The distribution and variance are shown as

violin and box plots in **Fig S4**. The potential age slowing effect of high income appears

particularly pronounced in the White population, whereas, for the Black participants, most of

the DNAm readouts had positive age-adjusted residual values even among the highest income

group (**Fig 2b**, **c**).

In Model 2, we included additional baseline health variables: history of hysterectomy, diabetes,

CVD, hypertension, and cancer; and in Model 3, we included all the variables in Model 2, plus

alcohol intake and the healthy eating index (2010 HEI) score to account for differences in

dietary intake (**Data S1–S3; Fig S5**). These models were implemented to verify that the negative

associations with SE index is robust. As sensitivity analysis, we repeated Model 2 in AS315 after

excluding all participants with DNA from follow-up annual visits (**Data S1**). All these showed

that the non-training based DNAm readouts, and the DunedinPACE and PhenoAge have

negative associations with SE index. The entropy and bivalent readouts were not significantly

associated with baseline health variables aside from slightly higher entropy for women who had

undergone hysterectomy in AS315 and BA23 (but not replicated in AS311; **Fig S5**). The training-

based biomarkers were more strongly related to baseline health and lifestyle variables. The

DunedinPACE and PhenoAge showed the expected age-accelerating effect of smoking, and the

DunedinPACE was slowed by higher total energy expenditure from recreational activity (MET-

hours/week) and higher HEI score. Higher BMI was positively associated with all the training-

based biomarkers, but not with entropy, BivSEpiM, or BivCpGAvg. In AS315, the BivCpGAvg

showed an unexpected significant negative regression estimate for prior smokers relative to

never smokers, but no difference between never and current smokers was observed; however,

this was not replicated in BA23 or AS311.

Social disparity and methylation at the chromatin states

To examine if the association with SE index was specific to the polycomb and bivalent CpGs, we

performed the Models 3 regressions with the age-adjusted mean methylation values for the

remaining 12 chromatin states in the AS315 and BA23 datasets. At a p threshold of 0.004

(Bonferroni corrected alpha = 0.05 for 12 tests), only the polycomb repressed state, ReprPC,

had significant negative association with SE index in AS315 and this was replicated in BA23 (**Fig**

3a; **Table S1**). Notably, like the bivalent states, ReprPC is a PRC2 targeted domain, and these

sites also have typically low methylation when young and accrue methylation with aging (all

plot to top left quadrant of **Fig 1e**). This pattern for the bivalent and polycomb sites suggests

that higher SE index is associated with a lower and presumably more "youthful" methylation

state. In both AS315 and BA23, higher SE index also had negative associations with methylation

levels at the active chromatin states, TssA and TssAFlnk (transcription start sites at or flanking

actively transcribed genes), which are also regions with low average methylation. However, at

the Bonferroni corrected alpha = 0.05, the negative association of TssA and TssAFlnk with SE

index was significant only in BA23 (**Fig 3a; Table S1**).

Fig 3. Forest plots of regression coefficients (95% confidence intervals) for SE index as predictor variable. Dependent variables are age-adjusted mean methylation at chromatin states (each fitted separately) after adjustment for other baseline variables (Model 3). One standard deviation higher SE index was associated with lower methylation at TssA, TssAFlnk, and ReprPC, and higher methylation at TxFlnk in both datasets.

Another chromatin state, TxFlnk (downstream of transcribed genes), had a positive association

with SE index in both AS315 and BA23; however, the association was not significant after

multiple test correction. Nonetheless, it is notable that in contrast to the bivalent states, CpGs

in TxFlnk typically exist in hemi-methylated states (beta close to 0.5), loses methylation with

aging, and contribute negatively to entropy (r=-0.41, **Fig 1d**). The positive association between

TxFlnk and SE index suggests that higher SE index may be related to methylation maintenance

at these CpGs.

DNAm readouts and all-cause mortality

Next, we tested whether these DNAm readouts predict life expectancy. Due to the low sample

numbers, we performed this analysis only for the Black, White and Hispanic participants for

AS315 and BA23. The PhenoAge clock is already known to be predictive of lifespan[3,27], and for

the present work, we examined only the non-training measures, and the DunedinPACE as a

training-based reference. We first performed a race/ethnicity stratified Cox regression for

censored survival time, and the DNAm readouts as primary predictors (entropy, BivSEpiM,

BivAvg, DunedinPACE; each fitted separately), and with SE index and baseline health variables

and the Clinical Trial (CT) dietary modification and hormone replacement arms as covariates. In

both AS315 and BA23, only the DunedinPACE was a significant predictor of lifespan, and one SD

increase in the pace of aging increased the risk of death by about 20% with hazard ratio (HR) of

1.23 (95% confidence interval CI = 1.13–1.33) in AS315 and HR = 1.16 (1.07–1.25) in BA23 (**Fig**

4a, 4b; **Table S2**). For SE index, one SD increase was associated with a modest decreased risk of

mortality of HR of 0.95 (CI = 0.87–1) in AS315, and HR 0.89 (0.83—0.96) in BA23. We further

examined if SE index and DunedinPACE were associated with lifespan when race/ethnicity are

analyzed separately (**Fig S6**; **Table S2**). In both AS315 and BA23, higher DunedinPACE increased

risk of mortality in all groups, and the added lifespan privilege of higher income was seen

mainly among the White participants.

We applied the same Cox regression to examine whether the average methylation levels of the

12 remaining chromatin states are predictive of all-cause mortality. Generally, higher

methylation at chromatin states that were negatively correlated with chronological age were

associated with reduced mortality risk (**Table S3**). This was nominally significant in both AS315

and BA23 for enhancer (Enh) CpGs (**Fig 4c**; **Table S3**). Higher mean methylation of Enh states

predicted lower mortality risk with HR of 0.90 (0.84—0.98) in AS315, and HR of 0.93 (0.88—

0.99) in BA23. Notably, the mean methylation of Enh regions also had the strongest negative

correlation with chronological age (**Fig 1f**). While the association with mortality risk is not

significant after Bonferroni correction, the overall pattern suggests that maintaining higher

methylation at these regulatory CpGs that typically lose methylation with aging could be related

to longer life expectancy.

Fig 4. Effects of DNAm readouts on all-cause mortality risk in **(a)** AS315 and **(b)** BA23 based on race/ethnicity stratified multivariable Cox regression. The predictor variables (entropy, BivSEpiM, BivCpGAvg, and DunedinPACE) were fitted separately and the forest plots displays the hazard ratios associated with each. Only the DunedinPACE is predictive of survival time in both datasets. **(c)** HR associated with mean methylation at chromatin states. Higher methylation at Enh (enhancers) has a nominally significant association with reduced risk of death in both AS315 and BA23.

DISCUSSION

To summarize: **(1)** We found that nearly all the large-scale methylation patterns (i.e., entropy,

overall methylation averages of chromatin states, and outlier counts at the bivalent sites)

showed some degree of change with chronological age among postmenopausal women and the

typical pattern was for the low methylation CpGs to gain, and high methylation CpGs to lose

methylation with aging. **(2)** Age-dependent changes in the bivalent and polycomb targeted sites

contributed to the increase in epigenetic entropy. **(3)** The bivalent and polycomb CpGs were

consistently associated with SE index and varied between self-reported race/ethnicity. **(4)**

However, the untrained readouts were not significantly associated with baseline health

variables or predictive of mortality risk. **(5)** In contrast, biomarkers trained on health-related

parameters (i.e., the DunedinPACE and PhenoAge) were significantly associated with baseline

variables such as BMI, smoking, and energy expenditure; furthermore, a higher pace of aging

estimated by the DunedinPACE was associated with a higher risk for all-cause mortality. **(6)**

Among the chromatin states, we found evidence that maintaining methylation levels at

enhancer CpGs could be associated with a lower risk of death.

Of the DNAm readouts, the most "global" measurement we computed was the methylome-

wide entropy that was defined using nearly the full set of CpGs. This value estimates the overall

randomness and is related to the phenomenon of "epigenetic drift", the increase in

stochasticity and variance in DNAm with aging.[14] In cohorts with a wide age range, epigenetic

entropy has a strong positive correlation with chronological age.[1] Among the WHI participants,

the positive correlation between entropy and age was modest, albeit significant. This is possibly

because the WHI represents an older cohort, and the relationship between epigenetic aging

and chronological age is not perfectly linear and there is evidence that it begins to plateau

among older individuals.[9,43] Initially we had anticipated that health variables such as BMI and

history of cancer would show some association with epigenetic entropy. However, this was not

the case. Instead, the combination of self-reported education and income (what we refer to as

SE index) was the only variable at baseline that had a consistent inverse association with global

entropy in all three WHI datasets. Scoring high on the SE index was linked to lower age-adjusted

entropy. Since the increase in DNAm entropy will partly reflect the increase in cellular

heterogeneity with aging, we also attempted to account for blood cell heterogeneity by

including the DNAm-based estimated proportions of blood cell types, and in doing so, the link

between epigenetic entropy and SE index became stronger (see **Data S2–S3**). The lowest age-

adjusted entropy was seen among the White and Hispanic women at the highest income grade,

while the highest age-adjusted entropy was seen among the low-income minority groups. This

suggests that while higher income and the numerous psychosocial, environmental, and health

privileges of wealth could be protective against the buildup of epigenetic discordance with

aging, the impact of income is not the same across all racial groups. In fact, the effect is blunted

especially among Black women. This is interesting, yet not surprising as Black women who

achieve high education and socioeconomic status continue to experience chronic stressors over

their lifetime due systemic racism, discrimination, microaggressions and cultural

expectations.[44,45]

SE index also had a similar inverse relationship with methylation levels at bivalent CpGs, which

were the strongest positive correlates of methylome-wide entropy. Having a low SE index was

characterized by higher levels of stochastic epimutations and age-dependent gains at these

bivalent sites. Here, stochastic epimutation simply refers to the counts of outlier CpGs and this

is another way of quantifying the levels of epigenetic discordance.[40,41] Our finding is consistent

with previous work by Fiorito et al.[40] Theirs was a multi-cohort study that included both males

and females, and participants from a wide age-range. They also used education level as a proxy

of socioeconomic position, and found that methylome-wide stochastic epimutation was higher

among those with lower education.[40]

We must note that both education and income are readily recordable variables that are part of

an extremely complex social construct of ones status in society and referred to as

socioeconomic status.[46] The SE index therefore serves only as a global proxy that tags along

with a multitude of unmeasured variables such as differences in access to health care and

education, food and nutrition options, neighborhood and housing conditions, environmental

pollutants and toxins, chronic stress, etc. These exposures—the social and environmental

determinants of health—exert a strong influence on interrelated biological processes such as

the stress and glucocorticoid pathway, composition of circulating immune cells and

inflammatory state, metabolic health, etc. [47-49] Our results suggest that these large-scale non-

training based readouts from the epigenome are influenced by the larger socio-environmental

context, but lack the sensitivity to serve as predictors of specific health and lifestyle conditions

such as BMI, energy expenditure, smoking, and disease status.

In contrast to the global entropy and bivalent readouts, the widely used epigenetic clocks and

pace-of-aging biomarkers are derived from extensive training algorithms. For the initial training,

both the PhenoAge and DunedinPACE included blood biomarkers that are related to health and

correlated with aging (e.g., C-reactive protein, blood cell counts, albumin, cholesterol).[3,4,50] The

DunedinPACE also included waist-hip ratio, dental health, lung function, cardiorespiratory

fitness, etc. Notably, the DunedinPACE was trained on the New Zealand (NZ) based Dunedin

Longitudinal Cohort, which consists predominantly of European ancestry participants, with only

a very small minority of participants identifying as indigenous people of New Zealand.[4,50] It is

quite remarkable that a biomarker panel of aging developed on a NZ cohort can generalized to

the societal structure and inequities in the United States. The DunedinPACE also revealed a

potential "anti-aging" effect of higher income among the White women, but the effect of

income was not as pronounced among the minority women. Except for the small group of self-

identified Asians on the higher income scale, all other minority groups had positive pace of

aging according to the DunedinPACE. While the training population could partly contribute to

the differences between race/ethnic groups, there is evidence that DunedinPACE captures the

effects of adversity and negative experiences that is independent of genetic ancestry.[51] Having

a higher pace of aging likely has important health consequences, and in both AS315 and BA23,

higher pace of aging as measured by DunedinPACE predicted shorter lifespan independent of SE

index.

Segregating the CpGs by their predicted chromatin states and computing the average

methylation levels was a very broad stroke approach. However, we applied this data reduction

to determine the general patterns by which chromatin states contribute to the increase in

epigenetic entropy. The strongest positive correlates of entropy were the methylation levels at

the bivalent chromatin domains and regions bound by the PRC2 (TssBiv, BivFlnk, EnhBiv,

ReprPC). These sites mark repressed states with low average methylation, and have an age-

dependent drift towards higher methylation. The bivalent and polycomb target sites are highly

conserved regulatory regions, and are crucially involved in embryonic development and in

maintaining cellular identity and function.[13] Similar to the bivalent and polycomb domains,

CpGs at active promoters (TssA) also typically have low methylation levels, and in the WHI, the

mean methylation at TssA was positively correlated with entropy. Notably, the chromatin states

that had significant positive correlations with entropy (i.e., higher methylation related to higher

entropy) were all negatively associated with SE index. An interpretation we draw from this is

that higher SE index is associated with the age-dependent increase in discordance and drift at

these CpG sites. In contrast, the chromatin state with the strongest negative correlation with

entropy was TxFlnk, and this was the only chromatin domain that showed a significant and

replicated positive association with SE index (means higher SE index related to higher

methylation levels at TxFlnk). TxFlnk marks CpGs located at 3' and 5' ends of transcribed genes,

and the mean methylation of TxFlnk had a weak inverse correlation with age (i.e., methylation

loss with aging), and maintaining a higher methylation reduced the level of entropy. The

positive association with SE index suggests that high SE index could help maintain a higher

methylation level at these CpGs, and thereby contribute to lower entropy.

Intriguingly, when it came to predicting survival time, the general trend was for higher

methylation at active chromatin states (e.g., TssA, TssAFlnk, EnhG, and Enh) to be linked to

lower risk of all-cause mortality, and this reached statistical significance for Enh. Except for

TssA, which was uncorrelated with chronological age, all these chromatin states tended to have

a negative correlation with aging, and this was strongest for Enh. Enhancers are important

regulatory elements, and the ENCODE 15-states model divides them into three categories

based on histone marks: bivalent enhancers in repressed sites (EnhBiv) that have low

methylation, and enhancers in genic (EnhG) and non-genic (Enh) sites.[35] Both Enh and EnhG

mark actively transcribed regions, and other studies have also shown that these have higher

methylation and lose methylation with aging.[52-54] Taken together, our results suggest that being

able to maintain higher methylation levels at these enhancer CpGs may extend lifespan. To our

knowledge, this is the first time the global average methylation of Enh CpGs has been

associated with all-cause mortality in humans. There is however evidence from mouse studies

that interventions such as caloric restriction, which extends lifespan, suppress the age

associated methylation loss at enhancer CpGs.[52] Furthermore, in humans, aberrant methylation

at enhancer CpGs have been implicated in cancer progression and metastasis.[55] The work by

Cole et al.[55] also showed that higher methylation at the enhancer CpG of the metastasis gene

KIT increased survival time among cancer patients. This direction of association is therefore in

agreement with our observation that higher methylation at Enh is associated with a decreased

risk of all-cause mortality.

In the present work, we only examined time to all-cause mortality but did not perform a more

detailed analysis to test whether these DNAm readouts predict time to adjudicated diseases or

disease-related deaths. Given the links between enhancer methylation and cancer progression

and mortality, this will be particularly relevant in a follow-up analyses. We also acknowledge

that the results we present are all based on older postmenopausal women (age 50-79 at

baseline), and the significant associations between the bivalent and polycomb sites and SE

index may represent the effects of lifelong exposures to social inequities and weathering.

Further work is needed to examine whether the link between SE index and the bivalent CpGs is

also seen in younger populations. For the DunedinPACE, there is strong and growing evidence

that it serves as a sensitive biomarker of the age-accelerating effects of social inequities and

discrimination in a wide age range, and in both males and females.[28,29,56-58] Based on this corpus

of work, we also anticipate that the association between the social variables and the non-

training based DNAm readouts present in this work will also be generalizable, but that is yet to

be tested. The link between enhancers and lifespan also needs further replication. To facilitate

such follow-up studies, we have provided all the computational codes, and these can be easily

computed from existing datasets and can be adapted to the newer DNAm microarray datasets.

In conclusion, we have presented alternative ways to quantify aspects of epigenetic aging that

are based on epigenetic entropy and stochasticity, and have related the large-scale methylation

features to aging, as well as to extrinsic variables, and potentially to lifespan. The presented

work highlights the deep links between the larger social environment and the aging of the

epigenome and provides evidence that methylation at active chromatin states could be related

to life expectancy.

METHODS

WHI DNA methylation datasets and sample inclusion

The WHI is a multicenter long-term prospective study of postmenopausal women that began in

1993.[18-20] Women between the ages 50–79 were recruited in 1993–1998. Ancillary studies have

generated genomic data, and the present work uses the genome-wide DNA methylation

dataset by the three ancillary studies : (1) WHI-EMPC/AS315,[26,31] (2) BA23,[21,32] and (3) AS311.

These 3 datasets have been previously analyzed jointly as part of large meta-studies, and

detailed descriptions can be found in these publications.[3,59-61]

The EMPC study (Epigenetic Mechanisms of PM-mediated CVD Risk; AS315) randomly selected

2200 participants from within the Clinical Trial arm and measured DNA methylation at SV for

the majority or at annual visit 3 or 6, and with repeated measurements for a subset of the

samples [31]. We used only the first sampling for the present work, and after excluding eight

samples with low methylation calls, the primary analyses were done for 2192 AS315

participants. The statistical analysis for AS315 was repeated in the 1396 SV samples (**Fig S1**).

BA23 was designed as a case/control study of risk for future CHD. We received epigenetic data

for 2107 BA23 participants collected at SV, and after excluding samples that overlap with WHI-

AS315, we performed the analyses in 1989 participants. AS311 is a matched case-control study

of bladder cancer, and like BA23, used blood DNA collected at SV and prior to disease diagnosis.

We received 882 array data; after excluding an array with too many missing values (~262,970

CpGs had NA values) and samples that overlap with AS315, we performed analyses in 868

participants.

Computing entropy, chromatin annotations, and estimates of biological aging

All analyses were done using the processed data that were provided in NetCDF format. For

uniformity, we used the beta-mixture quantile (BMIQ) normalized data, which applied the same

data quality control and processing steps implemented by WHI-EMPC [31]. The R package "ncdf4"

was used to interface with the data matrices [62]. Before any downstream analyses, we first

counted how many CpGs had missing values per sample, and excluded one sample from AS311,

and eight samples from BA23 as those had unusually high missingness (>50%).

For methylome-wide discretized entropy calculation, we used the same method described in a

prior study.[16] The optimal number of bins was estimated to be 50 using the Freedman-Diaconis

rule [63]. The R codes used for each step are in supplemental file **DataS5**. We then used the R

package "entropy" [64,65] to compute the discretized entropy value for each participant and this

was scaled to a number between 0–1. Example density plots for the beta-values were

generated in R with ggplot2 [66] for ten participants of AS315; these were all non-Hispanic Whites

and five had low entropy, and five had high entropy.

To annotate for chromatin states, we downloaded the archived REMCChromHMM file from the

Zhou github page [38,67]. This contains the 15-states chromHMM consensus chromatin

annotations for each of the ~480,000 CpGs in the 450K platform [35]. These annotations were

matched by CpG probe IDs to the WHI BMIQ data matrices, and average methylation levels for

each chromatin state was computed for each participant. As an alternate measure of

randomness in the epigenetic data, we counted CpGs that were outliers in the three bivalent

states to compute epigenetic stochasticity: bivalent/poised transcription start site (TssBiv),

flanking bivalent TSS (BivFlnk), and bivalent enhancer (EnhBiv). While the entropy measures the

shift in the genome-wide methylome landscape for each participant and is not relative to the

study cohort, the outlier counts are estimates of how many CpGs are outliers for an individual

relative to the study cohort. We adapted the outlier detection method described in Gentilini

but counted only the positive outliers as these sites gain methylation with aging and a higher

beta-value is presumed to represent a more aged methylome, i.e., beta-values over 3-times the

interquartile range of the 3^{rd} quartile [40,41]. The codes used to compute the bivalent stochastic

epimutations (BivSEpiM) are in **Data S4**.

For the first-generation epigenetic clocks (i.e., Horvath and Hannum), we used the pre-

computed clocks that were shared by the WHI. Specifically, we used the linear transformed

Horvath and Hannum DNAm age predictions [1,2]. We computed the PhenoAge (aka, Levine2018)

and the DunedinPACE using the dnaMethyAge R package [3,4,68,69].

Statistical analyses

Pearson correlations were used for bivariate comparisons of chronological age with the DNAm

readouts. The pair-wise correlation plots were generated using the corrplot R package [70], and

display significance threshold was set at p=0.001. For epigenetic clocks, which correlate

strongly with chronological age, the deviation of the predicted biological age from chronological

age (referred to as age-acceleration or age-deviation) is computed as the residuals of predicted

age regressed on chronological age [2]. As most of the DNAm readouts, and the DunedinPACE

also show varying levels of correlation with chronological age, we used the residuals to make

the values independent of chronological age.

We used the linear regression lm() function in R to test associations between the age-residuals

of the DNAm readouts and baseline variables. These baseline variables included BMI computed

from weight and height at baseline (kg/m^2), medical history at baseline, and personal habits

including smoking status (0 = never smoked; 1 = past smoker; and 2 = present smoker), and

recreational energy expenditure computed as measured by the TEXPWK variable (MET-

hours/week). TEXPWK was computed from questionnaire-based reports of recreational physical

activity that included walking, mild, moderate and strenuous physical activity in kcal/week/kg.[71]

Among the demographic variables, WHI collected self-reported education and income at time

of screening and are presented on an ascending scale from 1–11 for education, and 1–8 for

income (higher numbers represent higher education or income). We derived a single variable,

SEindex as the average of the two after excluding unreported or missing data.

We applied a simple model, Model 1: lm(y_i ~ race/ethnicity + SEindex + Smoking + BMI +

TEXPWK), and y_i is each of the DNAm readouts fitted independently. We limited the analysis to

participants who identified as non-Hispanic White (White), non-Hispanic Black (Black), Hispanic,

and Asian or Pacific Islander (Asian/PI) in AS315, and White, Black and Hispanic for BA23 and

AS311. Model 2 was the same as Model 1 and included additional medical history terms

collected at baseline: hysterectomy, diabetes, cardiovascular disease (CVD), hypertension, and

cancer ever. An expanded version of Model 1 included blood cell proportions (lymphocytes,

monocytes, granulocytes) that were indirectly estimated from the DNAm data. The DNAm data

for BA23 and AS311 were from baseline SV, and majority of the AS315 blood DNA was from the

SV prior to assignment of a study arm (n=1396). Model 2 was repeated for AS315 after

excluding the 633 blood DNA that were sampled at visit year 3, and 163 at visit year 4. Model 3

included all the variables in Model 2, plus the alcohol intake variable, and 2010 HEI total

score.[72,73] The lm() R codes and details on the variables are provided in **Data S4**.

We performed Cox regressions to test predictive value of the DNAm readouts using the 2022

versions of All Discovered Deaths, and Adjudicated Outcomes. This was done using the survival

R package [74,75]. For all-cause mortality, we used the variable DEATHALL (all discovered deaths

including from National Death Index) in conjunction with ENDFOLLOWALLDY (days enrollment

to end of follow up including uncensored death). Covariates for Cox regression were baseline

variables age, smoking status, BMI, diabetes, cancer, SE index, hormone therapy arm, dietary

modification arm, alcohol intake, 2010 HEI total score, and with race/ethnicity in the strata()

term. The coxph() R codes are provided in **DataS4**.

Data availability: All data are available upon request through the Women's Health

Initiative Study (https://www.whi.org).

Acknowledgement: The WHI program is funded by the National Heart, Lung, and Blood

Institute, National Institutes of Health, U.S. Department of Health and Human Services through

contracts HHSN268201600018C, HHSN268201600001C, HHSN268201600002C,

HHSN268201600003C, and HHSN268201600004C. WHI EMPC (AS315) was supported by NIEHS

grant R01-ES020836 (Whitsel, Eric A and other PIs). WHI AS311 was supported by American

Cancer Society award 125299-RSG-13–100-01-CCE (Parveen Bhatti). WHI-BAA23 was supported

by NHLBI Broad Agency Announcement contract HHSN268201300006C.

Author contribution: KM: study design, analyses and interpretation, initial

manuscript preparation, review and editing. AD, YS, AHS, RC, FJT, RBW, JHF: part

of WHI writing group and contributed to manuscript reviewing, editing, and

interpretation. KCJ: WHI sponsor and contributed to manuscript reviewing,

editing, and interpretation. All authors approved final manuscript.

REFERENCES.

Hannum, G. *et al.* Genome-wide methylation profiles reveal quantitative views of human aging rates. *Mol Cell* **49**, 359-367, doi:10.1016/j.molcel.2012.10.016 (2013).

Horvath, S. DNA methylation age of human tissues and cell types. *Genome Biol* **14**, R115, doi:10.1186/gb-2013-14-10-r115 (2013).

Levine, M. E. *et al.* An epigenetic biomarker of aging for lifespan and healthspan. *Aging (Albany NY)* **10**, 573-591, doi:10.18632/aging.101414 (2018).

Belsky, D. W. *et al.* DunedinPACE, a DNA methylation biomarker of the pace of aging. *Elife* **11**, doi:10.7554/eLife.73420 (2022).

Lu, A. T. *et al.* DNA methylation GrimAge strongly predicts lifespan and healthspan. *Aging (Albany NY)* **11**, 303-327, doi:10.18632/aging.101684 (2019).

Horvath, S. & Raj, K. DNA methylation-based biomarkers and the epigenetic clock theory of ageing. *Nat Rev Genet* **19**, 371-384, doi:10.1038/s41576-018-0004-3 (2018).

Liu, Z. *et al.* Underlying features of epigenetic aging clocks in vivo and in vitro. *Aging Cell* **19**, e13229, doi:10.1111/acel.13229 (2020).

Dabrowski, J. K. *et al.* Probabilistic inference of epigenetic age acceleration from cellular dynamics. *bioRxiv*, 2023.2003.2001.530570, doi:10.1101/2023.03.01.530570 (2023).

Meyer, D. H. & Schumacher, B. Aging clocks based on accumulating stochastic variation. *Nat Aging* **4**, 871-885, doi:10.1038/s43587-024-00619-x (2024).

Ikram, M. A. The use and misuse of 'biological aging' in health research. *Nat Med* **30**, 3045, doi:10.1038/s41591-024-03297-9 (2024).

Gems, D., Virk, R. S. & de Magalhaes, J. P. Epigenetic clocks and programmatic aging. *Ageing Res Rev* **101**, 102546, doi:10.1016/j.arr.2024.102546 (2024).

Moqri, M. *et al.* PRC2-AgeIndex as a universal biomarker of aging and rejuvenation. *Nat Commun* **15**, 5956, doi:10.1038/s41467-024-50098-2 (2024).

Schuettengruber, B., Bourbon, H. M., Di Croce, L. & Cavalli, G. Genome Regulation by Polycomb and Trithorax: 70 Years and Counting. *Cell* **171**, 34-57, doi:10.1016/j.cell.2017.08.002 (2017).

Teschendorff, A. E. On epigenetic stochasticity, entropy and cancer risk. *Philos Trans R Soc Lond B Biol Sci* **379**, 20230054, doi:10.1098/rstb.2023.0054 (2024).

Jenkinson, G., Pujadas, E., Goutsias, J. & Feinberg, A. P. Potential energy landscapes identify the information-theoretic nature of the epigenome. *Nat Genet* **49**, 719-729, doi:10.1038/ng.3811 (2017).

Mozhui, K. *et al.* Genetic loci and metabolic states associated with murine epigenetic aging. *Elife* **11**, doi:10.7554/eLife.75244 (2022).

Sziraki, A., Tyshkovskiy, A. & Gladyshev, V. N. Global remodeling of the mouse DNA methylome during aging and in response to calorie restriction. *Aging Cell* **17**, e12738, doi:10.1111/acel.12738 (2018).

NIH. Design of the Women's Health Initiative clinical trial and observational study. The Women's Health Initiative Study Group. *Control Clin Trials* **19**, 61-109, doi:10.1016/s0197-2456(97)00078-0 (1998).

19   Anderson, G. L. *et al.* Implementation of the Women's Health Initiative study design.
Ann Epidemiol **13**, S5-17, doi:10.1016/s1047-2797(03)00043-7 (2003).
20   WHI. <www.whi.org>
21   Levine, M. E. *et al.* DNA methylation age of blood predicts future onset of lung cancer in
the women's health initiative. *Aging (Albany NY)* **7**, 690-700, doi:10.18632/aging.100809
(2015).
22   Carroll, J. E. *et al.* Epigenetic Aging and Immune Senescence in Women With Insomnia
Symptoms: Findings From the Women's Health Initiative Study. *Biol Psychiatry* **81**, 136-
144, doi:10.1016/j.biopsych.2016.07.008 (2017).
23   Shadyab, A. H. *et al.* Association of Epigenetic Age Acceleration With Incident Mild
Cognitive Impairment and Dementia Among Older Women. *J Gerontol A Biol Sci Med Sci*
77, 1239-1244, doi:10.1093/gerona/glab245 (2022).
24   Quach, A. *et al.* Epigenetic clock analysis of diet, exercise, education, and lifestyle
factors. *Aging (Albany NY)* **9**, 419-446, doi:10.18632/aging.101168 (2017).
25   Pottinger, T. D. *et al.* Association of cardiovascular health and epigenetic age
acceleration. *Clin Epigenetics* **13**, 42, doi:10.1186/s13148-021-01028-2 (2021).
26   Grant, C. D. *et al.* A longitudinal study of DNA methylation as a potential mediator of
age-related diabetes risk. *Geroscience* **39**, 475-489, doi:10.1007/s11357-017-0001-z
(2017).
27   Liu, Z. *et al.* The role of epigenetic aging in education and racial/ethnic mortality
disparities among older U.S. Women. *Psychoneuroendocrinology* **104**, 18-24,
doi:10.1016/j.psyneuen.2019.01.028 (2019).
28   Harris, K. M. *et al.* Sociodemographic and Lifestyle Factors and Epigenetic Aging in US
Young Adults: NIMHD Social Epigenomics Program. *JAMA Netw Open* **7**, e2427889,
doi:10.1001/jamanetworkopen.2024.27889 (2024).
29   Shen, B. *et al.* Association of Race and Poverty Status With DNA Methylation-Based Age.
JAMA Netw Open **6**, e236340, doi:10.1001/jamanetworkopen.2023.6340 (2023).
30   Geronimus, A. T. The weathering hypothesis and the health of African-American women
and infants: evidence and speculations. *Ethn Dis* **2**, 207-221 (1992).
31   Gondalia, R. *et al.* Methylome-wide association study provides evidence of particulate
matter air pollution-associated DNA methylation. *Environ Int* **132**, 104723,
doi:10.1016/j.envint.2019.03.071 (2019).
32   Levine, M. E. *et al.* Menopause accelerates biological aging. *Proc Natl Acad Sci U S A* **113**,
9327-9332, doi:10.1073/pnas.1604558113 (2016).
33   Jordahl, K. M. *et al.* Genome-Wide DNA Methylation in Prediagnostic Blood and Bladder
Cancer Risk in the Women's Health Initiative. *Cancer Epidemiol Biomarkers Prev* **27**, 689-
695, doi:10.1158/1055-9965.EPI-17-0951 (2018).
34   Teschendorff, A. E., West, J. & Beck, S. Age-associated epigenetic drift: implications, and
a case of epigenetic thrift? *Hum Mol Genet* **22**, R7-R15, doi:10.1093/hmg/ddt375 (2013).
35   Roadmap Epigenomics, C. *et al.* Integrative analysis of 111 reference human
epigenomes. *Nature* **518**, 317-330, doi:10.1038/nature14248 (2015).
36   Ernst, J. & Kellis, M. Chromatin-state discovery and genome annotation with
ChromHMM. *Nat Protoc* **12**, 2478-2492, doi:10.1038/nprot.2017.124 (2017).

37   Kaur, D. *et al.* Comprehensive Evaluation of The Infinium Human MethylationEPIC v2
BeadChip. *Epigenetics Commun* **3**, doi:10.1186/s43682-023-00021-5 (2023).
38   Zhou, W., Laird, P. W. & Shen, H. Comprehensive characterization, annotation and
innovative use of Infinium DNA methylation BeadChip probes. *Nucleic Acids Res* **45**, e22,
doi:10.1093/nar/gkw967 (2017).
39   Rakyan, V. K. *et al.* Human aging-associated DNA hypermethylation occurs preferentially
at bivalent chromatin domains. *Genome Res* **20**, 434-439, doi:10.1101/gr.103101.109
(2010).
40   Fiorito, G. *et al.* Socioeconomic position, lifestyle habits and biomarkers of epigenetic
aging: a multi-cohort analysis. *Aging (Albany NY)* **11**, 2045-2070,
doi:10.18632/aging.101900 (2019).
41   Gentilini, D. *et al.* Stochastic epigenetic mutations (DNA methylation) increase
exponentially in human aging and correlate with X chromosome inactivation skewing in
females. *Aging (Albany NY)* **7**, 568-578, doi:10.18632/aging.100792 (2015).
42   Horvath, S. *et al.* An epigenetic clock analysis of race/ethnicity, sex, and coronary heart
disease. *Genome Biol* **17**, 171, doi:10.1186/s13059-016-1030-0 (2016).
43   Snir, S., Farrell, C. & Pellegrini, M. Human epigenetic ageing is logarithmic with time
across the entire lifespan. *Epigenetics* **14**, 912-926,
doi:10.1080/15592294.2019.1623634 (2019).
44   Baker, T. A., Buchanan, N. T., Mingo, C. A., Roker, R. & Brown, C. S. Reconceptualizing
successful aging among black women and the relevance of the strong black woman
archetype. *Gerontologist* **55**, 51-57, doi:10.1093/geront/gnu105 (2015).
45   Henry, C. J. & Song, M. K. Use of the Strong Black Woman Construct in Research: An
Integrative Review. *ANS Adv Nurs Sci* **47**, E110-E120,
doi:10.1097/ANS.0000000000000501 (2024).
46   Antonoplis, S. Studying Socioeconomic Status: Conceptual Problems and an Alternative
Path Forward. *Perspect Psychol Sci* **18**, 275-292, doi:10.1177/17456916221093615
(2023).
47   Smarr, M. M. *et al.* Broadening the Environmental Lens to Include Social and Structural
Determinants of Women's Health Disparities. *Environ Health Perspect* **132**, 15002,
doi:10.1289/EHP12996 (2024).
48   Palma-Gudiel, H., Fananas, L., Horvath, S. & Zannas, A. S. Psychosocial stress and
epigenetic aging. *Int Rev Neurobiol* **150**, 107-128, doi:10.1016/bs.irn.2019.10.020
(2020).
49   Simons, R. L. *et al.* Discrimination, segregation, and chronic inflammation: Testing the
weathering explanation for the poor health of Black Americans. *Dev Psychol* **54**, 1993-
2006, doi:10.1037/dev0000511 (2018).
50   Belsky, D. W. *et al.* Quantification of the pace of biological aging in humans through a
blood test, the DunedinPoAm DNA methylation algorithm. *Elife* **9**,
doi:10.7554/eLife.54870 (2020).
51   Bourassa, K. J. *et al.* Demographic characteristics and epigenetic biological aging among
post-9/11 veterans: Associations of DunedinPACE with sex, race, and age. *Psychiatry Res*
336, 115908, doi:10.1016/j.psychres.2024.115908 (2024).

52   Cole, J. J. *et al.* Diverse interventions that extend mouse lifespan suppress shared age-
associated epigenetic changes at critical gene regulatory regions. *Genome Biol* **18**, 58,
doi:10.1186/s13059-017-1185-3 (2017).
53   Jain, N. *et al.* DNA methylation correlates of chronological age in diverse human tissue
types. *Epigenetics Chromatin* **17**, 25, doi:10.1186/s13072-024-00546-6 (2024).
54   Slieker, R. C., Relton, C. L., Gaunt, T. R., Slagboom, P. E. & Heijmans, B. T. Age-related
DNA methylation changes are tissue-specific with ELOVL2 promoter methylation as
exception. *Epigenetics Chromatin* **11**, 25, doi:10.1186/s13072-018-0191-3 (2018).
55   Bell, R. E. *et al.* Enhancer methylation dynamics contribute to cancer plasticity and
patient mortality. *Genome Res* **26**, 601-611, doi:10.1101/gr.197194.115 (2016).
56   Schmitz, L. L. *et al.* The Socioeconomic Gradient in Epigenetic Ageing Clocks: Evidence
from the Multi-Ethnic Study of Atherosclerosis and the Health and Retirement Study.
Epigenetics **17**, 589-611, doi:10.1080/15592294.2021.1939479 (2022).
57   Raffington, L. *et al.* Associations of socioeconomic disparities with buccal DNA-
methylation measures of biological aging. *Clin Epigenetics* **15**, 70, doi:10.1186/s13148-
023-01489-7 (2023).
58   Cuevas, A. G. *et al.* Multi-discrimination exposure and biological aging: Results from the
midlife in the United States study. *Brain Behav Immun Health* **39**, 100774,
doi:10.1016/j.bbih.2024.100774 (2024).
59   Do, W. L. *et al.* Associations between DNA methylation and BMI vary by metabolic
health status: a potential link to disparate cardiovascular outcomes. *Clin Epigenetics* **13**,
230, doi:10.1186/s13148-021-01194-3 (2021).
60   Agha, G. *et al.* Blood Leukocyte DNA Methylation Predicts Risk of Future Myocardial
Infarction and Coronary Heart Disease. *Circulation* **140**, 645-657,
doi:10.1161/CIRCULATIONAHA.118.039357 (2019).
61   Skinner, H. G. *et al.* Stressful life events, social support, and epigenetic aging in the
Women's Health Initiative. *J Am Geriatr Soc* **72**, 349-360, doi:10.1111/jgs.18726 (2024).
62   Pierce, D. *ncdf4*, <https://CRAN.R-project.org/package=ncdf4> (2023).
63   Freedman, D. & Diaconis, P. On the histogram as a density estimator:L2 theory.
Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete **57**, 453-476,
doi:10.1007/BF01025868 (1981).
64   Hausser, J. & Strimmer, K. Entropy Inference and the James-Stein Estimator, with
Application to Nonlinear Gene Association Networks. *Journal of Machine Learning*
Research **10**, 1469-1484 (2009).
65   Hausser, J. & Strimmer, K. *entropy*, <https://CRAN.R-project.org/package=entropy >
(2021).
66   Wickham, H. *ggplot2: Elegant Graphics for Data Analysis.* (Springer-Verlag 2016).
67   Zhou, W. *zhou-lab github*, <https://github.com/zhou-
lab/ARCHIVED_KYCG_knowledgebases_hg38> (
68   Wang, Y., Grant, O. A., Zhai, X., McDonald-Maier, K. D. & Schalkwyk, L. C. Insights into
ageing rates comparison across tissues from recalibrating cerebellum DNA methylation
clock. *Geroscience* **46**, 39-56, doi:10.1007/s11357-023-00871-w (2024).
69   yiluyucheng. *dnaMethyAge*, <https://github.com/yiluyucheng/dnaMethyAge> (

70   Wei, T. & Simko, V. *R package 'corrplot': Visualization of a Correlation Matrix. (Version*
0.92), <https://github.com/taiyun/corrplot> (2021).
71   Meyer, A. M., Evenson, K. R., Morimoto, L., Siscovick, D. & White, E. Test-retest
reliability of the Women's Health Initiative physical activity questionnaire. *Med Sci*
Sports Exerc **41**, 530-538, doi:10.1249/MSS.0b013e31818ace55 (2009).
72   Guenther, P. M. *et al.* Update of the Healthy Eating Index: HEI-2010. *J Acad Nutr Diet*
113, 569-580, doi:10.1016/j.jand.2012.12.016 (2013).
73   Prentice, R. L. *et al.* Mortality Associated with Healthy Eating Index Components and an
Empirical-Scores Healthy Eating Index in a Cohort of Postmenopausal Women. *J Nutr*
152, 2493-2504, doi:10.1093/jn/nxac068 (2022).
74   Therneau, T. M. & Grambsch, P. M. *Modeling Survival Data: Extending the Cox Model*.
(Springer, 2000).
75   Therneau, T. M. *A Package for Survival Analysis in R*, <https://CRAN.R-
project.org/package=survival> (2024).
715

Main Tables

Table 1. Datasets and baseline characteristics

	EMPC/AS315	BA23[1]	AS311[1]
N	2192	1989	868
Age ± SD (years)	64 ± 7	65 ± 7	65.6 ± 7
BMI ± SD	29.4 ± 5.9	29.9 ± 6.1	28.0 ± 6.1
TEXPWK[2] (MET-hrs/wk)	9.7 ± 12.3	9.9 ± 12.6	11.5 ± 12.5
WHI study arm			
Clinical Trial (CT)	2191	1546	412
Observational Study (OS)		443	456
Race/Ethnicity			
Asian/Pacific Islander	133		15
Hispanic	318	402	25
Native American	51		1
Non-Hispanic Black	558	635	68
Non-Hispanic White	1097	952	749
Other	35		10
Education			
No data	18	19	5
Less Than HS	82	96	7
High School or GED	517	539	188
Vocation Degree	263	274	99
Some College or Graduate	793	654	341
Post College Graduate	519	407	228
Income ($)			
No data	124	119	55
≤19,999	460	589	148
20K–34,999	541	499	192
35K–49,999	414	366	174
50K–99,999	531	335	235
≥100K	122	81	64
SE index (range)[3]	5.4 ± 1.6 (1–11)	5.1 ± 1.6 (1–10)	5.7 ± 1.5 (1–11)
Smoking			
No data	29	31	17
Never	1159	1048	389
Past	821	713	389
Current	183	197	73
CVD ever			
No data	223	140	52
No	1687	1572	633
Yes	282	277	183
Hypertension ever			

No data	18	24	1
No	1362	1092	558
Yes	812	873	309
Cancer ever			
No data	24	14	
No	2088	1867	868
Yes	80	108	0
Diabetes ever			
No data		3	
No	2018	1734	814
Yes	174	252	54
Hysterectomy ever			
No	1201	1037	525
Yes	991	952	343

719 [1]Excludes participants sampled in AS315. [2]Total energy expenditure from recreational activity (TEXPWK or MET-
720 hours/week). [3]Computed as the average of education level (WHI EDUC variable ranging from 1 = *Didn't go to*
721 *school*" to 11= "*Doctoral degree*") and income (WHI INCOME variable ranging from 1 = "*Less than $10,000*" to 8 =
722 "*150,000 or more*").

Table 2. Predicted chromatin states for the >450,000 CpG

ChromHMM	Counts[1]	Repressed/ Active	Description	Non-coding conserved elements	Mean beta in AS315	r (age)[2]	p (age)[2]
1_TssA	124139	Active	Active TSS	Strongly enriched	0.08	0.04	0.0476
2_TssAFlnk	23774	Active	Flanking Active TSS	Strongly enriched	0.29	-0.15	<.0001
3_TxFlnk	1001	Active	Transcribed state at gene 5' and 3'	Moderately enriched	0.52	-0.08	<.0001
4_Tx	32162	Active	Strong transcription	No enrichment	0.89	-0.08	0.0001
5_TxWk	50018	Active	Weak transcription	No enrichment	0.83	-0.11	<.0001
6_EnhG	1545	Active	Genic enhancers	Moderately enriched	0.75	-0.15	<.0001
7_Enh	20980	Active	Enhancers	Strongly enriched	0.59	-0.19	<.0001
8_ZNF/Rpts	987	Active	ZNF genes & repeats	No enrichment	0.74	0.11	<.0001
9_Het	3468	Repressed	Heterochromatin	No enrichment	0.72	0.08	0.0002
10_TssBiv	6833	Repressed	Bivalent/Poised TSS	Strongly enriched	0.10	0.16	<.0001
11_BivFlnk	8290	Repressed	Flanking Bivalent TSS/Enh	Strongly enriched	0.11	0.11	<.0001
12_EnhBiv	4869	Repressed	Bivalent Enhancer	Strongly enriched	0.16	0.09	<.0001
13_ReprPC	32422	Repressed	Repressed PolyComb	Strongly enriched	0.36	0.07	0.0007
14_ReprPCWk	33551	Repressed	Weak Repressed PolyComb	Weakly enriched	0.73	-0.11	<.0001
15_Quies	136857	Repressed	Quiescent/Low	No enrichment	0.80	-0.10	<.0001
NA	1339						

[1]Number of CpG probes on the Illumina 450K platform located in these predicted chromatin states. [2]Pearson correlation and p with chronological age in AS315

Table 3. Association between epigenetic readouts and socioeconomic index (SE index)

DNAm outcomes (age residuals)	AS315					BA23				
	Min	Max	SD	SE index coef[1]	p	Min	Max	SD	SE index coef[1]	p
DNAm Entropy	-0.655	0.344	0.10	-0.004	0.018	-0.548	0.452	0.13	-0.005	0.008
BivSEpiM (log10 outlier counts)	-1.437	2.312	0.50	-0.02	0.009	-1.453	2.373	0.51	-0.009	0.216
BivCpGAvg (beta)	-0.032	0.060	0.01	-0.001	4.20E-07	-0.059	0.098	0.02	-0.001	7.00E-07
DunedinPACE	-0.358	0.706	0.13	-0.005	0.002	-0.414	0.485	0.13	-0.009	4.91E-07
Horvath (years)	-27.71	93.40	6.22	-0.13	0.16	-29.16	54.18	6.64	-0.11	0.263
Hannum (years)	-21.07	48.44	5.35	-0.15	0.058	-23.86	31.59	5.83	-0.30	0.0007
PhenoAge (years)	-25.86	57.20	6.60	-0.25	0.012	-38.29	36.32	7.31	-0.46	3.30E-05

[1]Unstandardized regression estimates based on Model 1