

1 **Penetration of Inner Scleral Fibers into Peripapillary**

2 **Sclera Revealed by Polarization-Sensitive OCT**

3 Jianping Xiong¹, MD, Hongshuang Lu¹, MD, Nan Zhou¹, MD, Yijin Wu¹, MD,
4 Masahiro Yamanari², PhD, Hiroyuki Takahashi¹, MD, PhD, Michiaki Okamoto²,
5 BSc, Keigo Sugisawa¹, MD, Takeshi Yoshida^{1,3}, MD, PhD, Changyu Chen¹,
6 MD, Yining Wang¹, MD, Ziyue Wang¹, MD, Kyoko Ohno-Matsui¹, MD, PhD

7 ¹Department of Ophthalmology and Visual Science, Institute of Science Tokyo,
8 Tokyo, 113-8510, Japan

9 ²Tomey Corporation, Nagoya, Aichi-ken, 451-0051, Japan

10 ³Department of Advanced Ophthalmic Imaging, Institute of Science Tokyo,
11 Tokyo, Japan

12 **Corresponding Author:** Professor Kyoko Ohno-Matsui, Department of

13 Ophthalmology and Visual Science, Institute of Science Tokyo, 1-5-45

14 Yushima, Bunkyo-ku, Tokyo 1138510, Japan; Phone: 81-3-5803-5302, Fax:

15 81-3-3818-7188,

16 e-mail: k.ohno.oph@tmd.ac.jp

17

18 **Running title:** Inner Scleral Penetration into Circular Pattern

19 **Word count:** 3255 words

20 **Key words:** peripapillary sclera, collagen, optic nerve, polarization-sensitive

21 optical coherence tomography, myopia

22 **Funding/Support:** This work was supported by JST SPRING, Grant Number
23 JPMJSP2120, and received partial research funding from Tomey Corporation.
24 The funding organization and Tomey Corporation had no role in the design and
25 conduct of the study; collection, management, analysis, and interpretation of
26 the data; preparation, review, or approval of the manuscript; and decision to
27 submit the manuscript for publication.

28 **Conflict of Interest/Financial Disclosures:** Dr Xiong reported receiving a
29 research grant from JST SPRING. Dr Ohno-Matsui reported receiving a
30 research grant from Tomey Corporation and being a consultant for Santen and
31 Cooper Vision outside the submitted work. Dr Yamanari reported having
32 patents for JP 6463051 B2, US 9593936 B2, EP 2995245 B1, JP 6542178 B2,
33 and JP 7332131 B2 issued to Tomey Corporation. Dr Yoshida belongs to the
34 Department of Advanced Ophthalmic Imaging at the Institute of Science Tokyo,
35 which is funded by NIDEK Corporation. No other disclosures were reported.
36

37 **Abstract**

38 **Purpose:** To determine the structural characteristics of the peripapillary sclera
39 in myopic eyes using polarization-sensitive optical coherence tomography
40 (PS-OCT).

41 **Methods:** Patients with myopia who underwent PS-OCT imaging between
42 April and June 2023 were studied. The intensity, optic axis, and birefringence
43 of PS-OCT images were analyzed to determine the surface of the lamina
44 cribrosa (LC). The direction and arrangement of the scleral collagen fibers in
45 the peripapillary region and scleral ridge were determined.

46 **Results:** One hundred and ten eyes of 59 patients with myopia (mean axial
47 length, 30.00 ± 1.98 mm) were studied. The scleral fibers around the optic
48 nerve were arranged circularly in all eyes regardless of the presence or size of
49 the peripapillary atrophy. Inner scleral fibers that penetrated beneath this
50 circular arrangement were detected in 98 eyes (89.1%). There was a Y-shaped
51 split at the point of penetration in 77 eyes (70.0%). The integrity of the circular
52 arrangement and penetrating scleral fiber complexes remained intact even in
53 cases of severe peripapillary scleral deformation. A scleral ridge was identified
54 in 27 eyes (24.5%), and it appeared as a full-thickness protrusion composed
55 primarily of outer scleral fibers. The ridge was consistently located outside the
56 circular arrangement.

57 **Conclusions:** The penetration of the radial inner scleral fibers into the circular
58 arrangement and extension to the LC suggest that they act as stabilizing
59 anchors for the circular structure. The differences of the scleral ridges from
60 those in eyes with dome-shaped macula suggest that unique mechanisms
61 cause the scleral differences.
62

63 **Introduction**

64 The peripapillary sclera plays a critical role in the biomechanics of the eye.¹⁻³

65 Its structure (shape and thickness) and material properties (stiffness,

66 brittleness, and compliance) affect the mechanical sturdiness of the optic

67 nerve (ON) significantly. These properties thereby affect its susceptibility to

68 increased intraocular pressure (IOP) and other internal and external

69 pressures.⁴

70

71 Earlier studies examined the structure of the peripapillary sclera by wide-angle

72 X-ray scattering (WAXS), small-angle light scattering (SALS), and second

73 harmonic generation microscopic images of normal human and rat eyes.^{5,6}

74 These studies identified a key feature of the posterior scleral structure: a ring

75 of collagen fibers encircling the scleral canal. Gogola et al. used polarized light

76 microscopy (PLM) to show a radial arrangement of the inner scleral fibers and

77 circumferential arrangement of the outer scleral fibers. These findings were

78 consistent in humans, monkeys, pigs, cows, goats, and sheep⁷, and they have

79 greatly enhanced our understanding of the anatomy of the peripapillary sclera.

80

81 Polarization-sensitive optical coherence tomography (PS-OCT) is an

82 advanced OCT technology that uses polarized light as an additional image

83 contrast mechanism.^{8,9} It uses the birefringent properties of the sclera to image

84 and quantify the variations of birefringence across different regions of the
85 sclera. PS-OCT can also make visible the axis of orientation, or optic axis, of
86 the birefringence associated with the direction of the fiber bundles.¹⁰ Based on
87 these capabilities of PS-OCT, Baumann et al.¹¹ in rats and Willemse et al.¹² in
88 healthy humans eyes to confirm the peripapillary scleral 3D structure. Most
89 earlier studies suggested that the radial inner scleral fibers terminated at the
90 circular arrangement adjacent to the scleral canal.^{5-7,11-14}

91
92 The peripapillary circular scleral structure is known to provide protection and
93 support to the ON.^{5,7,13,14} Thus, Hua et al. created seven posterior pole models
94 demonstrating that radial fibers also contribute to the support of the posterior
95 ocular surface.¹ Most importantly, they demonstrated that the combination of
96 the radial and circumferential fibers provided enhanced ON protection at
97 elevated IOP. However, the structural relationship between the peripapillary
98 scleral fibers of different orientations has not been well explored using *in vivo*
99 imaging. This deficiency is due to the difficulties in obtaining detailed images of
100 the peripapillary scleral structures *in vivo*. However, the choroidal thinning in
101 myopic eyes facilitates the ability to record clear images of all the scleral layers
102 by PS-OCT.

103

104 Thus, the purpose of this study was to determine the structural construction of
105 the peripapillary sclera in myopic human eyes. To accomplish this, we
106 analyzed the structure of the sclera in the PS-OCT images of myopic eyes.

107

108 **Methods**

109 The procedures used in this study conformed to the tenets of the Declaration
110 of Helsinki and were approved by the Ethics Committee of the Institute of
111 Science Tokyo (Science Tokyo). The procedures complied with the standards
112 for case series studies which allowed us to present the results thoroughly and
113 transparently.

114

115 We reviewed the medical records of highly myopic patients who had
116 undergone PS-OCT examinations between April 1, 2023, and June 16, 2023 at
117 the Advanced Clinical Center for Myopia at Science Tokyo. Cases with
118 poor-quality PS-OCT images were excluded. All patients had undergone a
119 comprehensive ophthalmological examination including measurements of the
120 axial length (IOL Master; Carl Zeiss Meditec AG, Jena, Germany),
121 best-corrected visual acuity (BCVA), refractive error (spherical equivalent),
122 color fundus photography, and swept-source OCT (SS-OCT, Triton; Topcon,
123 Tokyo, Japan).

124

125 **PS-OCT Examinations and Image Processing**

126 To assess the characteristics of the birefringence of the posterior segment of
127 the eye, we developed a PS-OCT system using a swept-source laser centered
128 at 1050 nm (ROCTIA, Tomey Corporation, Japan). The hardware of this
129 system is commercially available in Japan. A comprehensive description of the
130 technical concepts and the system has been presented.^{15,16} Retinal images
131 were obtained over scan areas of 81 mm² (9 × 9 mm) or 144 mm² (12 × 12 mm)
132 using a raster scanning protocol with 1024 A-scans along the horizontal axis
133 and 256 B-scans along the vertical axis. In addition, a 6-line radial scan
134 protocol was also used. This system has an axial resolution of 7.3 μm with an
135 axial measurement range of 4.49 mm.

136
137 Already developed algorithms^{10,17-19} were adapted to process the measured
138 raster scan data that yielded the polarization-diverse OCT intensities^{20,21}, local
139 phase retardation or magnitude of birefringence, and the orientation of the
140 optic axis (OA) of the birefringence. These examinations resulted in three
141 corresponding images: referred to as the intensity, birefringence, and OA
142 images. To further enhance the image quality, the axial pixel separation was
143 increased which improved the performance in localization of the axially
144 accumulated Jones matrix.¹⁰ A necessary correction was used to resolve the
145 OA at each axial depth.¹⁰

146

147 The birefringence of collagen fibers occurs when light passes obliquely or
148 perpendicularly to the OA as in fundus imaging. This causes the collagen
149 fibers to exhibit two refractive indices that alters the polarization state along the
150 axial depth. In perfectly aligned fibers, the PS-OCT can mathematically
151 determine the orientation of the OA on a plane perpendicular to the direction of
152 the light.¹⁰ However, scleral collagen fibers have an interwoven structure that
153 exceeds the resolution of OCT,²² and this results in a net birefringence from
154 partially canceled birefringence of the individual fibers. PS-OCT provides
155 depth-resolved measurements of this net birefringence *in vivo*. The OA, which
156 corresponds to the preferential orientation of the net birefringence created by
157 interwoven collagen fibers, reflects the preferential orientation of the fibers
158 despite their complex interwoven structure.

159

160 For the data analysis, we processed the horizontal raster scan data and
161 obtained three types of images with 256 cross-sections for each eye. The
162 intensity images were used to locate the anterior surface of the lamina cribrosa
163 (LC), the OA images were used to determine the direction of the scleral
164 collagen fibers especially in the peripapillary area and at the scleral ridge. A
165 scleral ridge was defined as an anterior protrusion of the sclera that is located
166 temporal to the ON with a height greater than 150 μm .^{23,24} Radial scan images

167 centered on the ON were used to determine the presence and height of the
168 ridge. More specifically, the ridge measurements were made along the
169 horizontal meridian using built-in tools (ROCTIA, Tomey Corporation, Japan).
170 Birefringence images were used to assist in determining the alignment of the
171 scleral fibers. High birefringence corresponded to more aligned scleral fibers²⁵,
172 and low birefringence corresponded to less aligned scleral fibers such as
173 interwoven fibers.

174

175 Streamline rendering is a method for viewing the vectorial data and effectively
176 makes visible the orientation of fibrous tissues such as the OA. This technique
177 has been applied to imaging the OA of the heart muscles, articular cartilage,
178 and scleral collagen fibers using PS-OCT.^{10,18,26} We used the ParaView
179 software (version 5.12.0, Kitware Inc.) for volumetric streamline rendering of
180 the OA following the methods of earlier research.¹⁰ The streamline images
181 obtained directly and showed clearly the course of the scleral fibers.

182

183 **Statistical Analyses**

184 Statistical analyses were performed using SPSS version 25.0 (IBM Corp.,
185 2017. IBM SPSS Statistics for Windows, Version 25.0. Armonk, NY: IBM
186 Corp.). Normally distributed data are expressed as the means \pm standard

187 deviations (SD), and non-normally distributed data are presented as medians
188 with corresponding quartiles.

189

190 **Results**

191 One hundred and ten eyes of 59 patients with myopia that had undergone
192 PS-OCT imaging at the Advanced Clinical Center for Myopia at Science Tokyo
193 between April 1, to June 16, 2023, were studied. There were 35 women and 24
194 men with a mean (\pm SD) age of 55.1 ± 10.2 years and range of 26.8 to 76.3
195 years. The mean axial length was 30.00 ± 1.98 mm with a range of 24.63 to
196 34.13 mm. The mean refractive error (spherical equivalent) was -13.24 ± 4.62
197 diopters (D) with a range of -1.63 to -24.75 D excluding 30 eyes with an
198 implanted intraocular lens (IOL) and 1 eye with a contact lens. The median
199 best-corrected visual acuity (BCVA) was 0.0 logMAR units with a range -0.3 to
200 1.0 logMAR units.

201

202 The *en face* OA images were revised by a one-dimensional moving average
203 for the 15 axial pixels ($67 \mu\text{m}$) in the scleral slab, and the resulting images
204 showed that the fibers in the inner layer of the sclera were distributed radially
205 (Figure 1). The inner and outer layers of the sclera could be clearly
206 distinguished in the horizontal cross-sectional OA images. The collagen fibers
207 of the inner and outer layers of the sclera surrounding the optic nerve (ON) had

208 different orientations: the inner scleral fibers extended radially from the ON to
209 the periphery, while the outer scleral fibers were oriented vertically and
210 obliquely (Figure S1).

211

212 **Circular arrangements of scleral fibers around optic nerve (ON)**

213 In the *en face* OA images, the scleral fibers were noted to be arranged
214 circularly around the ON in all eyes. The size and shape of this circular
215 arrangement were determined solely by the ON, irrespective of the presence
216 or size of the peripapillary atrophy (PPA; Figure 1). Even in eyes with a large
217 PPA or deformed optic discs, the circular arrangement remained confined to
218 the area immediately surrounding the ON. The circular arrangement appeared
219 to be a projection of the outer sclera at the edge of the ON in the OA images
220 (Figures 2–5) as reported in an earlier study.¹² The fibers were composed
221 essentially of the outer circumferentially oriented sclera fibers and were only
222 seen and distinguishable in the OA images.

223

224 **Penetration of inner sclera fibers within and below the circular** 225 **arrangement**

226 We found in the OA image that the inner scleral fibers did not terminate outside
227 the circular arrangement. In 98 of 110 eyes (89.1%), the inner scleral fibers
228 penetrated beneath the circular arrangement (Figure 2). The penetrating fibers

229 ran toward the ON and probably toward the LC. In the remaining eyes, the
230 penetration of the inner scleral fibers could not be detected because of an
231 extremely tilted ON in 5 eyes, thick choroid in 4 eyes, and loss of the typical
232 circular arrangement in 3 eyes.

233

234 Additionally, the inner scleral layer was observed to split axially into a Y-shape
235 in 77 of 110 eyes (70.0%) at the initial point where the fibers penetrated the
236 circular arrangement. However, in these eyes, the complete course of
237 penetrating fibers was not consistently detected.

238

239 In representative cases (Figure 3), the superior branch passed over the body
240 of the circular arrangement and extended toward the anterior portion of the LC.
241 On the other hand, the inferior branch passed beneath the circular
242 arrangement and extended into or beyond the LC.

243

244 The inner scleral layer was also found to split into three or more branches. It
245 was not divided into only two fiber bundles but it could be further subdivided
246 into thinner bundles. Notably, 73 of 110 eyes (66.4%) had at least one small
247 superior penetrating branch with its complete course extending through the
248 main body of the circular arrangement. Because each eye had 256 PS-OCT

249 image sections, some values overlapped, allowing a simultaneous detection of
250 the above phenomena in a single eye.

251

252 In the birefringence images, all of the penetrating inner scleral fiber bundles
253 had low birefringence.

254

255 **Resilience of circular and penetrating scleral fiber complexes**

256 We also observed that even in cases of severe deformation of the peripapillary
257 sclera, the circular arrangements and the penetrating inner scleral fibers
258 remained unaffected (Figure 4). Various peripapillary abnormalities including
259 conus pits and scleral ridges were commonly observed in eyes with pathologic
260 myopia. Conus pits are thought to originate from scleral stretch-associated
261 schisis or openings for the short posterior ciliary arteries that form pit-like clefts
262 in the adjacent scleral crescent.^{27,28}

263

264 Scleral ridges are anterior protrusions of the sclera located temporal to the
265 optic disc.^{23,28} Among the 110 eyes, nine had conus pits with a circular
266 arrangement around the ON in all of these eyes. In eight of them, the
267 penetration of the inner scleral fibers remained in place, and one eye could not
268 be examined in detail because of the loss of the typical circular arrangement.

269

270 **PS-OCT images of eyes with scleral ridge**

271 Scleral ridges were observed in 27 of 110 (24.5%) eyes. In all 27 eyes, the
272 protrusions involved the entire scleral thickness rather than only the inner
273 sclera (Figure 5). In addition, the ridges were composed mainly of outer scleral
274 fibers rather than the inner scleral fibers. The ridges were always located
275 outside the circular arrangement of the scleral fibers.

276

277 **Discussion**

278 The OA in the posterior sclera have been studied in rats, guinea pigs, and
279 humans.^{10-12,25} These images consistently revealed a two-layer structure in the
280 peripapillary sclera: an inner layer of radial fibers and an outer layer of
281 circumferential fibers. The OA images from PS-OCT were needed to show the
282 scleral fiber alignment. This imaging approach showed a clear difference
283 between the inner and outer scleral layers (Figure S1) which then provided
284 valuable information on the physiological and pathological states of the sclera
285 *in vivo*. Building on the results of our laboratory findings on the characteristics
286 of dome-shaped maculas (DSMs),¹⁰ we examined the fiber structure
287 surrounding the ON in this study.

288

289 Our images showed a circular arrangement of the collagen fibers around the
290 ON in all eyes, and they appeared as projections of the outer sclera at the

291 edge of the ON (Figures 2-5). The circumferential arrangement of the fibers
292 around the ON has been documented earlier by *in vitro* using techniques such
293 as WAXS⁵, SALS⁶, PLM^{7,13}, and more recently by *in vivo* PS-OCT images.^{11,12}
294 The circumferential fibers adjacent to the ON form a proximal ring which we
295 refer to as a circular arrangement. Our observations closely agree with
296 previous findings, and the circular arrangement appeared more distinct and
297 easier to identify in our cross-sectional OA images compared to those obtained
298 by earlier algorithms.^{7,12,13}

299

300 The schematic illustration in Figure S2 is based on our observations and
301 shows the course of the peripapillary scleral fibers and the penetration pattern
302 of the inner radial fibers. Previous histological studies reported that the outer
303 sclera transitions into the dura mater while the inner sclera forms the LC and
304 contributes to the pia mater.^{29,30} On the other hand, the earlier studies reported
305 that the radial inner fibers emerged outside the circular arrangement slightly
306 farther from the optic canal.^{5-7,11-14} However, how the inner sclera is
307 connected to the LC has not been conclusively determined.

308

309 In 89.1% of the eyes studied, cross-sectional OA images showed that the inner
310 radial scleral fibers penetrated beneath the circular arrangement and ran
311 toward the ON and the LC (Figure 2). A Y-shaped split was seen in 70.0% of

312 eyes at the initial point of penetration. In addition, the radial fibers were seen to
313 divide into multiple bundles, and 66.4% of the eyes had at least one fully
314 visible superior bundle (Figures 3 and 4).

315

316 Our findings are the first detailing the unique patterns of inner scleral fibers
317 that penetrated the ring and extended toward the ON rather than terminating
318 outside. These findings provide new information on the contribution of the
319 inner sclera to the formation of the LC. All penetrating bundles had low
320 birefringence (Figures 2H, 2L, 3H, 3L, 4G), indicating less aligned and more
321 interwoven scleral fibers.

322

323 The structural and functional relationships between the circumferential and
324 radial fibers have not been determined. Ji et al. reported discrete LC insertions
325 into the sclera with significant variations in the depth and quadrant
326 distribution.³¹ Our findings showed that the inner scleral fibers penetrated the
327 circular arrangement and split into superior and inferior or multiple branches,
328 and closely aligned with these depth-dependent variations in the LC insertions.

329 In addition, the inconsistent observations of this penetration across
330 cross-sections reflect the discrete nature of the LC insertions. These spatial
331 correlations strongly support our suggestion that the radial fibers of the inner
332 sclera directly connect with the LC by this penetration mechanism. Most

333 previous studies focused on *en face* observations, with few employing
334 continuous cross-sectional approaches by PS-OCT. Our study has bridged this
335 methodological gap, leading to novel findings.

336

337 The circumferential fibers are widely accepted as protection for neural tissues
338 by resisting canal expansion and providing structural support for the ON.^{5,7,13,14}

339 Hua et al.¹ showed that the radial fibers reduced the displacement of the LC
340 and the peripapillary scleral tension under elevated IOP through finite element
341 models. The combination of radial and circumferential fibers protected the ON
342 better than either alone.

343

344 The *en face* OA images showed that the size and shape of the circular
345 arrangement depended exclusively on the ON, which was unaffected by the
346 presence or size of PPA (Figure 1). Cross-sectional OA images showed that
347 even significant deformations of the peripapillary sclera, such as a scleral pit,
348 did not alter the integrity of the complexes of the circular arrangements and the
349 penetrating inner scleral fibers (Figure 4). These findings suggested that the
350 structural stability of the circular arrangement, and emphasized the critical
351 support resulting from its interweaving with the penetrating fibers. The
352 penetrating fibers may act as anchors, stabilizing the circular arrangement
353 through a fabric-like interweaving which would provide mechanical support.

354

355 Scleral ridges were observed in 24.5% of eyes as full-thickness protrusions
356 and were not limited to the inner sclera. They consisted primarily of outer
357 scleral fibers (Figure 5). Previous research from our laboratory using PS-OCT
358 showed that the inner scleral fibers aggregated without a thickening of the
359 outer sclera in highly myopic eyes with DSM sites.¹⁰ Although both scleral
360 ridges and DSMs involved an inward protrusion of the sclera, their structural
361 components were markedly different. The distinct compositions of the scleral
362 abnormalities suggested different pathogenic mechanisms potentially requiring
363 individualized management for each type of scleral abnormality.

364

365 Our study has several limitations. First, it included only myopic eyes and
366 mostly highly myopic eyes as full-thickness scleral imaging was feasible only
367 with significant choroidal thinning. Because scleral fiber penetrations were
368 observed in most cases, we propose that the complexes likely represented a
369 physiological structure in all eyes. Future basic studies are needed to confirm
370 these findings. Second, the LC region was not suitable for study because of
371 the spatially unresolved birefringent signals from the retinal nerve fibers.
372 Consequently, a direct confirmation that the penetrating fibers form the LC was
373 not achieved, but SS-OCT indirectly validated this relationship by locating the
374 anterior surface of the LC. Third, the imaging principles of OA, relying on the

375 preferential orientation of net birefringence, could not fully represent the
376 complex microstructure of the sclera, such as the individual collagen
377 lamellae²² and the waviness of the collagen fibers¹⁴. Fourth, consistent with
378 Willemse et al.¹², we also failed to detect regions of interweaving fibers
379 throughout the scleral depth described by Gogola et al.⁷ Last, the pia mater
380 that is situated parallel to the incident light, could not be viewed through its OA,
381 blocking an exploration of its relationship^{29,30} with the inner radial scleral fibers.
382 SS-OCT has difficulties in identifying the posterior LC surface, although some
383 of the penetrating fibers appeared to extend posteriorly to the LC (Figure S2).
384 These fibers possibly linked the inner scleral fibers to the formation of the pia
385 mater. However, current technology cannot verify this. These technical
386 limitations underscore the need for future advancements in the algorithms to
387 optimize PS-OCT which would then enable a broader application in clinical
388 practice.

389

390 In conclusion, we examined the structural characteristics of the peripapillary
391 sclera by PS-OCT, and we found that the circular arrangement of the scleral
392 fibers around the ON is stable. The inner radial fibers penetrate and extend
393 toward the LC and act as anchors of the circular arrangement. Scleral ridges
394 had distinct compositional differences compared to DSMs, suggesting that the

395 mechanisms underlying peripapillary scleral abnormalities differ from those in
396 other regions. These findings offer insights for future clinical management.
397

398 **References**

- 399 1. Hua Y, Voorhees AP, Jan NJ, et al. Role of radially aligned scleral
400 collagen fibers in optic nerve head biomechanics. *Exp Eye Res.* Oct
401 2020;199:108188. doi:10.1016/j.exer.2020.108188
- 402 2. Sigal IA, Ethier CR. Biomechanics of the optic nerve head. *Exp Eye Res.*
403 Apr 2009;88(4):799-807. doi:10.1016/j.exer.2009.02.003
- 404 3. Safa BN, Wong CA, Ha J, Ethier CR. Glaucoma and biomechanics.
405 *Curr Opin Ophthalmol.* Mar 1 2022;33(2):80-90.
406 doi:10.1097/ICU.0000000000000829
- 407 4. Ren R, Wang N, Li B, et al. Lamina cribrosa and peripapillary sclera
408 histomorphometry in normal and advanced glaucomatous Chinese
409 eyes with various axial length. *Invest Ophthalmol Vis Sci.* May
410 2009;50(5):2175-2184. doi:10.1167/iovs.07-1429
- 411 5. Pijanka JK, Spang MT, Sorensen T, et al. Depth-dependent changes in
412 collagen organization in the human peripapillary sclera. *PLoS One.*
413 2015;10(2):e0118648. doi:10.1371/journal.pone.0118648
- 414 6. Girard MJ, Dahlmann-Noor A, Rayapureddi S, et al. Quantitative
415 mapping of scleral fiber orientation in normal rat eyes. *Invest*
416 *Ophthalmol Vis Sci.* Dec 28 2011;52(13):9684-9693.
417 doi:10.1167/iovs.11-7894

- 418 7. Gogola A, Jan NJ, Lathrop KL, Sigal IA. Radial and Circumferential
419 Collagen Fibers Are a Feature of the Peripapillary Sclera of Human,
420 Monkey, Pig, Cow, Goat, and Sheep. *Invest Ophthalmol Vis Sci*. Oct 1
421 2018;59(12):4763-4774. doi:10.1167/iovs.18-25025
- 422 8. Baumann B. Polarization Sensitive Optical Coherence Tomography: A
423 Review of Technology and Applications. *Appl Sci-Basel*. May
424 2017;7(5)doi:ARTN 474
425 10.3390/app7050474
- 426 9. de Boer JF, Hitzengerger CK, Yasuno Y. Polarization sensitive optical
427 coherence tomography - a review [Invited]. *Biomed Opt Express*. Mar 1
428 2017;8(3):1838-1873. doi:10.1364/BOE.8.001838
- 429 10. Ohno-Matsui K, Igarashi-Yokoi T, Azuma T, et al. Polarization-Sensitive
430 OCT Imaging of Scleral Abnormalities in Eyes With High Myopia and
431 Dome-Shaped Macula. *JAMA Ophthalmol*. Mar 7
432 2024;doi:10.1001/jamaophthalmol.2024.0002
- 433 11. Baumann B, Rauscher S, Glosmann M, et al. Peripapillary rat sclera
434 investigated in vivo with polarization-sensitive optical coherence
435 tomography. *Invest Ophthalmol Vis Sci*. Oct 28 2014;55(11):7686-7696.
436 doi:10.1167/iovs.14-15037
- 437 12. Willemse J, Grafe MGO, Verbraak FD, de Boer JF. In Vivo 3D
438 Determination of Peripapillary Scleral and Retinal Layer Architecture

- 439 Using Polarization-Sensitive Optical Coherence Tomography. *Transl Vis*
440 *Sci Technol.* Oct 2020;9(11):21. doi:10.1167/tvst.9.11.21
- 441 13. Jan NJ, Lathrop K, Sigal IA. Collagen Architecture of the Posterior Pole:
442 High-Resolution Wide Field of View Visualization and Analysis Using
443 Polarized Light Microscopy. *Invest Ophthalmol Vis Sci.* Feb 1
444 2017;58(2):735-744. doi:10.1167/iovs.16-20772
- 445 14. Boote C, Sigal IA, Grytz R, Hua Y, Nguyen TD, Girard MJA. Scleral
446 structure and biomechanics. *Prog Retin Eye Res.* Jan 2020;74:100773.
447 doi:10.1016/j.preteyeres.2019.100773
- 448 15. Yamanari M, Mase M, Obata R, et al. Melanin concentration and
449 depolarization metrics measurement by polarization-sensitive optical
450 coherence tomography. *Sci Rep.* Nov 11 2020;10(1):19513.
451 doi:10.1038/s41598-020-76397-4
- 452 16. Yamanari M, Uematsu S, Ishihara K, Ikuno Y. Parallel detection of
453 Jones-matrix elements in polarization-sensitive optical coherence
454 tomography. *Biomed Opt Express.* May 1 2019;10(5):2318-2336.
455 doi:10.1364/BOE.10.002318
- 456 17. Todorovic M, Jiao S, Wang LV, Stoica G. Determination of local
457 polarization properties of biological samples in the presence of
458 diattenuation by use of Mueller optical coherence tomography. *Opt Lett.*
459 Oct 15 2004;29(20):2402-2404. doi:10.1364/ol.29.002402

- 460 18. Fan C, Yao G. Imaging myocardial fiber orientation using polarization
461 sensitive optical coherence tomography. *Biomed Opt Express*. Mar 1
462 2013;4(3):460-465. doi:10.1364/BOE.4.000460
- 463 19. Li Q, Karnowski K, Noble PB, et al. Robust reconstruction of local optic
464 axis orientation with fiber-based polarization-sensitive optical
465 coherence tomography. *Biomed Opt Express*. Nov 1
466 2018;9(11):5437-5455. doi:10.1364/BOE.9.005437
- 467 20. Yamanari M, Lim Y, Makita S, Yasuno Y. Visualization of phase
468 retardation of deep posterior eye by polarization-sensitive swept-source
469 optical coherence tomography with 1-microm probe. *Opt Express*. Jul
470 20 2009;17(15):12385-12396. doi:10.1364/oe.17.012385
- 471 21. Ueno Y, Mori H, Kikuchi K, Yamanari M, Oshika T. Visualization of
472 Anterior Chamber Angle Structures With Scattering- and
473 Polarization-Sensitive Anterior Segment Optical Coherence
474 Tomography. *Transl Vis Sci Technol*. Dec 1 2021;10(14):29.
475 doi:10.1167/tvst.10.14.29
- 476 22. Li Q, Karnowski K, Untracht G, et al. Vectorial birefringence imaging by
477 optical coherence microscopy for assessing fibrillar microstructures in
478 the cornea and limbus. *Biomed Opt Express*. Feb 1
479 2020;11(2):1122-1138. doi:10.1364/BOE.382755

- 480 23. Xie S, Kamoi K, Igarashi-Yokoi T, et al. Structural Abnormalities in the
481 Papillary and Peripapillary Areas and Corresponding Visual Field
482 Defects in Eyes With Pathologic Myopia. *Invest Ophthalmol Vis Sci.* Apr
483 1 2022;63(4):13. doi:10.1167/iovs.63.4.13
- 484 24. Xiong J, Du R, Xie S, et al. Papillary and Peripapillary Hemorrhages in
485 Eyes With Pathologic Myopia. *Invest Ophthalmol Vis Sci.* Nov 1
486 2022;63(12):28. doi:10.1167/iovs.63.12.28
- 487 25. Liu X, Jiang L, Ke M, et al. Posterior scleral birefringence measured by
488 triple-input polarization-sensitive imaging as a biomarker of myopia
489 progression. *Nat Biomed Eng.* Jun 26
490 2023;doi:10.1038/s41551-023-01062-w
- 491 26. Yao G, Duan D. High-resolution 3D tractography of fibrous tissue based
492 on polarization-sensitive optical coherence tomography. *Exp Biol Med*
493 *(Maywood)*. Feb 2020;245(4):273-281.
494 doi:10.1177/1535370219894332
- 495 27. Ohno-Matsui K, Lai TY, Lai CC, Cheung CM. Updates of pathologic
496 myopia. *Prog Retin Eye Res.* May 2016;52:156-187.
497 doi:10.1016/j.preteyeres.2015.12.001
- 498 28. Ohno-Matsui K, Akiba M, Moriyama M, et al. Acquired optic nerve and
499 peripapillary pits in pathologic myopia. *Ophthalmology.* Aug
500 2012;119(8):1685-1692. doi:10.1016/j.ophtha.2012.01.047

- 501 29. Hogan MJ, Alvarado JA, Weddell JE. *Histology of the Human Eye: An*
502 *Atlas and Textbook*. Saunders; 1971.
- 503 30. Jonas JB, Spaide RF, Ostrin LA, Logan NS, Flitcroft I, Panda-Jonas S.
504 IMI-Nonpathological Human Ocular Tissue Changes With Axial Myopia.
505 *Invest Ophthalmol Vis Sci*. May 1 2023;64(6):5. doi:10.1167/iovs.64.6.5
- 506 31. Ji F, Islam MR, Wang B, Hua Y, Sigal IA. Lamina Cribrosa Insertions
507 Into the Sclera Are Sparser, Narrower, and More Slanted in the Anterior
508 Lamina. *Invest Ophthalmol Vis Sci*. Apr 1 2024;65(4):35.
509 doi:10.1167/iovs.65.4.35

510

511

512 **Figure legends**

513

514 **Figure 1.** Circular arranged scleral fibers around the optic nerve (ON), and
515 their relationship with the ON size rather than the peripapillary atrophic (PPA)
516 size.

517 A: Fundus photograph of the right eye of a 71-75-year-old woman with a
518 refractive error (spherical equivalent) of -7.6 diopters (D) and an axial length
519 (AL) of 25.7 mm with a tessellated fundus. Scarred macular neovascularization
520 can be seen.

521 B: *En face* optical coherence tomographic (OCT) intensity projection image
522 shows a small PPA temporal to the ON.

523 C: The optic axis (OA) projection image has a circular arrangement of the
524 scleral fibers around the ON (arrow). The circular arrangement exists just
525 around the ON within the PPA. Outside the circular arrangement, the inner
526 scleral fibers extend radially from the ON toward the fovea. The OA images (C,
527 F, and I) are colored based on the color wheel in the lower right corner, with
528 blue, yellow, pink, and green indicating the horizontal, vertical, and 2 oblique
529 scanning directions, respectively.

530 D: Fundus photograph of the left eye of a 51-55-year-old woman with a
531 refractive error of -17.8 D and an AL of 31.4 mm showing diffuse choroidal
532 atrophy with large PPA.

533 E: *En face* OCT intensity projection image showing a large PPA.

534 F: OA projection image shows a relatively small circular arrangement just

535 around the ON (arrow) within the PPA.

536 G: Fundus photograph of the right eye of a 51-55-year-old man with a

537 refractive error of -10.6 D and an AL of 28.9 mm showing diffuse choroidal

538 atrophy.

539 H: *En face* OCT intensity image shows horizontally oval ON surrounded by

540 annular PPA.

541 I: OA projection image shows horizontally arranged oval fibers just around the

542 ON (arrow) within the PPA.

543

544 **Figure 2.** The circular arrangement of the scleral fibers around the optic nerve

545 (ON) consists of the outer sclera with the inner sclera penetrating through this

546 circle to continue toward the lamina cribrosa (LC).

547 A: Fundus photograph of the left eye of a 61-65-year-old man with an

548 implanted intraocular lens and an AL of 30.5 mm showing diffuse choroidal

549 atrophy. The black line is the scanned line for the PS-OCT images in B to H.

550 B:, C:, and D: Horizontal scan PS-OCT images across the fovea.

551 B: Intensity image.

552 C: OA image shows that the inner and outer layers of the sclera around

553 the ON have different orientations of the scleral fibers. The inner sclera is

554 composed mainly of radial fibers which are horizontally oriented in this
555 cross-sectional image (blue; arrows). In contrast, the outer sclera is composed
556 of circumferentially oriented fibers and is obliquely oriented in this section
557 (mixed colors; arrowhead).

558 D: Birefringence image showing that the inner scleral fibers have high
559 birefringence (arrows). Peripapillary structures outlined by the squares are
560 magnified in images E to H.

561 E: to H: Magnified images of the squares in images B to D.

562 E: and F: Intensity images show the anterior surface of LC
563 (arrowheads). Peripapillary sclera appears to protrude anteriorly (arrow).

564 G: In the OA image, the circular arrangement is located at the edge of
565 the ON and is seen as the projection of the outer scleral fibers (white arrow). A
566 penetration of almost the entire thickness of the inner sclera is observed (red
567 arrow) beneath the circular arrangement and runs toward the LC.

568 H: In the birefringence image, the circular arrangement shows high
569 birefringence (white arrow), while penetrating inner scleral fibers have low
570 birefringence (red arrow).

571 I: to L: Peripapillary magnified PS-OCT images scanned at 235 μm inferior to
572 the scan line shown in A.

573 I: and J: Intensity images show the anterior surface of the LC
574 (arrowheads). Peripapillary sclera appear to protrude anteriorly (arrow).

575 K: In the OA image, the circular arrangement is located at the edge of
576 the ON and is seen as a projection of the outer sclera (white arrow). A
577 penetration of almost the entire thickness of the inner sclera is observed (red
578 arrow) beneath the circular arrangement and runs toward the LC.

579 L: In the birefringence image, the penetrating inner scleral fibers have
580 low birefringence (red arrow).

581 The OA images (C, G, and K) are colored based on the color wheel in image C,
582 with blue, yellow, pink, and green indicating the horizontal, vertical, and 2
583 oblique scanning directions, respectively. The birefringence images (D, H, and
584 L) are colored using the color bar in the upper right corner and are intentionally
585 saturated to show the low birefringence structure more clearly.

586

587 **Figure 3.** The circular arrangement of the scleral fibers around the optic nerve
588 (ON) consists of the outer sclera. The inner sclera can be seen to penetrate
589 through this circle to continue toward the lamina cribrosa (LC). The color wheel
590 and bar labeling are the same as in Figure 2.

591 A: Fundus photograph of the left eye of a 56-60-year-old woman with a
592 refractive error of -19.0 D and an AL of 31.6 mm showing diffuse choroidal
593 atrophy. Black line is the scanned line of PS-OCT images in B to H.

594 B:, C:, and D: PS-OCT images correspond to the scan line shown in A.

595 B: Intensity image.

596 C: OA image showing that the inner sclera is horizontally oriented in this
597 cross-sectional image (blue; arrows). The outer sclera is generally vertical in
598 this section (yellow; arrowhead).

599 D: Birefringence image shows that the inner scleral fibers are highly
600 birefringent (arrows). Peripapillary structures in the squares are magnified and
601 shown in images E to H.

602 E: to H: Magnified images of the squares in images B to D.

603 E: and F: Intensity images show the anterior surface of the LC
604 (arrowheads).

605 G: In the OA image, the inner scleral fibers do not terminate outside the
606 circular arrangement; they split in a Y-shape pattern and penetrate and extend
607 toward the LC. More specifically, the superior branch (white arrow) passes
608 through the body of the circular arrangement and runs toward the anterior part
609 of the LC. The inferior branch (red arrow) passes beneath the circular
610 arrangement and extends into or deeper into the LC.

611 H: Birefringence image showing that both the penetrating inner scleral
612 fiber bundles have low birefringence (arrows).

613 I: to L: Peripapillary magnified images of PS-OCT images scanned at 47 μm
614 inferior to the scan line shown in A.

615 I: and J: Intensity images show the anterior surface of the LC
616 (arrowheads).

617 K: In the OA image, the superior branch (white arrow) of the split inner
618 scleral fibers penetrates within the body of circular arrangement and runs
619 toward the anterior part of the LC. The inferior branch (red arrow) passes
620 beneath the circular arrangement and extends into or deeper into the LC.

621 L: Birefringence image. The circular arrangement shows high
622 birefringence (arrowhead) but both of the penetrating inner scleral fiber
623 bundles have low birefringence (arrows).

624

625 **Figure 4.** The inner scleral penetrating fibers remain even in eyes with a
626 scleral conus pit. The color wheel and bar labeling are the same as in Figure 2.

627 A: Left fundus of a 71-75-year-old woman with an implanted intraocular lens
628 and an AL of 34.9 mm. Multiple areas of patchy choroidal atrophy can be seen
629 surrounding the macula. A large area of peripapillary atrophy (PPA) can be
630 seen around the optic nerve (ON). The scleral ridge can be seen temporal to
631 the ON (arrowheads). The solid and dotted lines show the locations of two
632 PS-OCT scans.

633 B: *En face* OCT intensity image. Multiple conus pits can be clearly seen
634 (arrowheads).

635 C: A horizontal scan of swept-source OCT image across the macula shows a
636 large and deep conus pit (arrow). The sclera at the conus pit is posteriorly

637 bowed while the remaining retinal vessels remain protruded (arrowheads).

638 Macular sclera shows a dome-shaped macula (DSM) appearance.

639 D: to G: PS-OCT images corresponding to the dotted scan line shown in A.

640 D: The intensity image shows a deep conus pit (arrow). DSM is also
641 seen.

642 E: OA images. The inner scleral layer (arrowheads) is thick at the DSM
643 and continues until the conus pit (arrow).

644 F: This is a magnified image of the square area in E. A projection of the
645 outer scleral oblique fibers is seen just around the ON (white arrow).

646 Penetration of horizontally running fibers just below the circular arrangement
647 (red arrow) is seen toward the ON.

648 G: Birefringence image shows high birefringence at the circular
649 arrangement around the ON (white arrow) while penetrating inner scleral fibers
650 show low birefringence (red arrow).

651 H: to M: PS-OCT images corresponding to the solid scan line shown in A.

652 H:, I:, and J: Intensity images. A conus pit is seen (arrow in H). I and J
653 are the same magnified images of square area in H. The anterior surface of
654 the LC is shown (arrowheads in J).

655 K: OA images. A conus pit is seen (arrow in K). L and M are the same
656 magnified images of the square area in K.

657 L: Inner scleral layer is thin in this scan and ends before the conus pit.

658 However, just around the ON, the projection of outer sclera (consisting of

659 circular arranged fibers) is seen (arrow).

660 M: The inner scleral fibers that penetrate the circular arrangement

661 remain in place (two branches; blue and red arrows). The superior branch

662 (blue arrow) passes through the body of the circular arrangement and runs

663 toward the anterior part of the LC, and the inferior branch (red arrow) passes

664 beneath the circular arrangement and extends into or beneath the LC.

665

666 **Figure 5.** PS-OCT images of an eye with a scleral ridge. The color wheel and

667 bar labeling are the same as in Figure 2.

668 A: Left fundus of a 36-40-year-old woman with a refractive error of -17.4 D and

669 an AL of 30.8 mm showing diffuse choroidal atrophy. Peripapillary atrophy is

670 seen temporal to the optic nerve (ON). A scleral ridge can be seen temporal to

671 the ON (arrowheads). The black line is the scanned line of the PS-OCT images

672 in B to F.

673 B: and C: In the intensity images of PS-OCT, the scleral ridge is clearly seen

674 (white arrow). A magnified image of the square is shown in C. A slight

675 protrusion of the sclera is also seen just around the ON (blue arrow).

676 D: and E: In the OA images, the scleral ridge is seen as a protrusion of the

677 entire scleral thickness rather than just the inner sclera. The scleral ridge is

678 composed primarily of the outer rather than the inner scleral fibers. In the
679 magnified image E, the scleral ridge is located outside the circular
680 arrangement (white arrow). The inner sclera over the ridge is significantly
681 thinned (arrowheads). There is another protrusion just around the ON which is
682 the circular arrangement of the scleral fibers (blue arrow). The inner scleral
683 fibers that penetrate posterior to this circle can also be seen (red arrow).
684 F: Birefringence image of the same area. Scleral ridge and the circular
685 arrangement have generally high birefringence.
686









