Title Page

Manuscript Title: Genomic Insights into Opioid Addiction: Identification of genomic variants from Gene Expression data

Swati Ajmeriya[1], Biswadip Chatterjee[2], Subhradip Karmakar*[1]

1. Department of Biochemistry, All India Institute of Medical Sciences, Ansari Nagar, New Delhi-110029, India

2. National Drug Dependence Treatment Centre (NDDTC), All India Institute of Medical Sciences, Ansari Nagar, New Delhi-110029, India

Swati Ajmeriya, M.Sc
Department of Biochemistry, All India Institute of Medical Sciences, Ansari Nagar, New Delhi - 110029
Email: ajmeria.swati@gmail.com
ORCID ID: 0000-0002-2015-0335
Phone: +91 9990616801

Dr. Biswadip Chatterjee, MBBS, MD
Additional Professor, National Drug Dependence Treatment Centre,
All India Institute of Medical Sciences, New Delhi
Email:biswadip.c@gmail.com
ORCID ID: 0000-0002-3812-9147
Phone: +91 7042804218

Corresponding Author
Subhradip Karmakar, PhD
Additional Professor
Department of Biochemistry, All India Institute of Medical Sciences, Ansari Nagar, New Delhi - 110029
Email: subhradip.k@aiims.edu
ORCID ID: 0000-0002-4757-8729
Phone - +91 9999162564Phone: +91-11-26589556, Fax: +91-11-26588662

Abstract

Next-generation sequencing (NGS) has transformed high-throughput DNA and RNA analysis, facilitating the rapid identification of clinically relevant genetic variants. Opioid Use Disorder (OUD), a chronic condition characterized by relapse and remission cycles, poses significant challenges for genomic investigations. While whole-exome sequencing (WES) and whole-genome sequencing (WGS) serve as robust approaches for variant detection, their high cost restricts widespread use. RNA sequencing (RNA-Seq) presents a viable alternative; however, the complexity of the transcriptome complicates reliable variant identification. Despite these challenges, RNA-Seq has emerged as a valuable tool for detecting single nucleotide polymorphisms (SNPs) in conditions with limited WES data, such as OUD. In this study, RNA-Seq data from postmortem ventral midbrain specimens of chronic opioid users (PRJNA492904) were analyzed to identify variants associated with OUD. Given its established involvement in opioid addiction and impulsivity, we hypothesized that the *NRXN3* gene would harbor a significant number of variants. Variant analysis was conducted across eight genes: *BDNF, DRD2, DRD3, NRXN3, OPRD1, OPRM1,* and *NGFB*—with a primary focus on *NRXN3*. Our results revealed that *NRXN3* exhibited the highest variant burden among the analyzed genes, highlighting its potential role in OUD pathogenesis and reinforcing its association with opioid addiction.

Keywords: Opioid addiction, opioid use disorder, addiction, genomics, transcriptomics

Introduction

Substance use disorders are characterized by frequent relapses, with relapse rates ranging from 56.8% to 81.8%(Chen 2010; Peacock et al. 2018). The World Drug Report published by the CDC shows that the number of drug overdose deaths increased by nearly 30% from 2019 to 2020 and nearly 75% of the 91,799 drug overdose deaths in 2020 involved an opioid; from 2019 to 2020, Opioid-involved death rates increased by 38%(World Drug Report 2019). The past several decades have seen parallel increases in the prescription of opioid drugs and prescription-related opioid overdose deaths, followed by a resurgence in heroin abuse and deaths, and, most recently, there has been an abrupt rise in opioid deaths involving fentanyl or fentanyl analogs in the United States(Rudd et al. 2016). Thus, despite best and continuous efforts, the outcome of opioid use remains poor and it remains an important cause of premature death among the users. It has been established that genetic factors contribute to the vulnerability to developing drug addiction and to the effectiveness of its treatment (Bierut 2011; AL-Eitan et al. 2021). Genetic factors majorly contribute to the individual variations of the risk of developing opioid addiction with □~□60% of the population(Kendler et al. 2003; Goldman et al. 2005). Identification of these factors may increase our understanding of the disorders, help in the development of new treatments, and advance personalized medicine. Considerable effects on protein function and gene expression can be caused by variants occurring in coding regions and regulatory sequences, respectively(Zhao et al. 2019). Opioid exposure and OUD are linked to widespread gene expression changes in the brain and opioid-related variants have been reported to be associated with specific drug addictions(Levran et al. 2012; Seney et al. 2021). The significance of utilizing human postmortem brains for molecular characterization of neuropsychiatric disorders has been

thoroughly reviewed (Egervari et al. 2019a) This approach holds the potential to greatly enhance

our understanding of psychiatric disorders.

Drugs of abuse share the property of acutely increasing the signaling of dopamine (DA)-

synthesizing midbrain neurons to several forebrain targets(Koob and Volkow 2010) (Volkow and

Morales 2015). With continued drug use, neuroadaptations such as drug dependence, craving,

and relapse are thought to arise from persistent changes in gene expression(Koob and Volkow

2010; Volkow and Morales 2015). Gene expression changes have been identified in human

forebrain targets of dopamine (DA) signaling, such as the nucleus accumbens and prefrontal

cortex(Albertson et al. 2006; Ribeiro et al. 2017; Egervari et al. 2019b). However, there is

limited information regarding both gene expression changes and SNPs in midbrain human DA

neurons, despite their crucial role in the circuitry of addiction. Today, in the era of biomedical

research, the accessibility of Next-generation sequencing (NGS) offers a streamlined means to

comprehensively uncover all variants including SNPs and insertion and deletion variants

(indels).   Several methods have been implemented for initial SNPs discovery in a high-

throughput manner, such as whole-genome sequencing, exome capture, RNA sequencing,

methylated DNA sequencing, and restriction enzyme (RE) digestion(Davey et al. 2011).

Recently, transcriptome sequencing, or RNA-seq, has become one of the most representative

high-throughput sequence-based techniques because of its high accuracy and cost-

effectiveness(Zhao et al. 2019). However, the identification of variants from existing RNA-Seq

remains a challenge because of the transcriptome complexity. At present, Illumina with good

sequencing coverage and read quality, is applied to many plant species for variant detection by

RNA-seq(Zhao et al. 2019). Some existing studies prove the reliable and accurate identification

of variants from RNA-Seq data. One such study has evaluated the accuracy of SNPs identified

from RNA-Seq data that has utilized its application to produce massive SNPs at relatively low

cost(Rogier et al. 2018). Furthermore, RNA-Seq has been shown to detect genomic variations in

expressed regions like DNA-seq, as elucidated by(Piskol et al. 2013). Furthermore, a Trinity-

GATK pipeline has been developed in a research study to identify reliable variants from RNA-

Seq data (Zhao et al. 2019). These findings demonstrate the prospective benefits of utilizing

RNA-Seq or transcriptome sequencing as high-throughput sequence-based technology for

variant discovery. Previously, the applications and advantages of RNA-Seq to discover variants

have been reviewed and described. RNA-Seq has been described to be able to discover

thousands of variants as well as the expression levels of functional genes with sequence

variations, all at a reasonable cost(Zhao et al. 2019). For these reasons, RNA-seq is developing

into an extensive application in genetic polymorphism analysis(Zhao et al. 2019). There is a

significant paucity of genomic data related to OUD and other psychiatric disorders, resulting in a

limited understanding of the molecular and genetic mechanisms underlying these conditions.

This research gap persists despite the established association between gene expression changes

and brain physiology, particularly in opioid addiction. In the current study, we employed

PRJNA492904 gene expression data to detect variants derived from ventral midbrain specimens

from chronic opioid users due to the limited information on OUD-related variants. We have

elucidated the significance of variants in the genes previously found to have a crucial role in

OUD. The current investigation represents, to our knowledge, the first unbiased examination of

variants in the midbrain associated with human opioid abuse in the expression region of the

human genome. Our analyses revealed variants across seven candidate genes (BDNF, DRD2,

DRD3, NRXN3, OPRD1, OPRM1, and NGFB), among which NRXN3 exhibited the highest

frequency of genetic variation. Some pieces of evidence demonstrate the indispensable role of α-

NRXNs in the regulation of normal neurotransmitter release mechanisms(Missler et al. 2003).

NRXNs have been considered a potential candidate gene for drug addiction(Lachman et al.

2007a). In European and African American samples, 38 loci containing SNPs were strongly

associated with the amount of opioid abuse, including NRXN3 (Liu et al. 2005). Furthermore,

the involvement of NRXN3 in normal neurotransmitter release and synaptic integrity suggests

that variation in this gene may have widespread and substantial effects on key behavioral

phenotypes such as impulsivity(Stoltenberg et al. 2011a). Therefore, by utilizing the RNA-Seq

data this study examined the genetic polymorphisms in *NRXN3* relative to the other candidate

genes in transcriptome expression data. The pronounced variants observed in NRXN3 support

the notion that NRXN3 may be involved in mediating impulsive and addictive behavior among

individuals with opioid addiction as well as the reliability of RNA-Seq to detect genetic variants.

Material and Methods

Acquisition of RNA-Seq data for variant calling

We used publicly available RNA-seq data on humans with opioid abuse from the Sequence Read

Archive (SRA) on postmortem ventral midbrain specimens from chronic opioid users (N $\square$ = $\square$ 30)

and drug-free control subjects (N $\square$ = $\square$ 20), as described in the Methods with project ID

PRJNA492904(Saad et al. 2019a).

Case Inclusion and Exclusion Criteria for the dataset PRJNA492904

The inclusion criteria for the opioid abuse cohort (n=30) included: (1) a verifiable history of

opioid abuse, corroborated by a toxicology report indicating the presence of opioids and a

forensic analysis attributing death directly to opioid consumption. Inclusion was also extended to

a subset of cases with concurrent detection of opioids and cocaine, reflecting the prevalent

pattern of polydrug use with opioids. (2) Subjects with a positive toxicological screening for

non-opioid substances of abuse (e.g., alcohol, cannabinoids, anxiolytics, barbiturates) were

systematically excluded from participation. For the control group (n=20), the inclusion criteria

were: (a) absence of a documented drug abuse history and negative toxicological tests for

opiates, cocaine, and other substances of abuse or central nervous system (CNS) medications.

The causes of death in control cases were primarily cardiovascular incidents or gunshot injuries.

Exclusion criteria for either group encompassed: (a) a documented history of neurologic or

psychiatric disorders, death by suicide, evidence of neuropathological conditions, severe chronic

diseases, an estimated postmortem interval exceeding 20 hours, or biochemical indicators of

compromised tissue sample integrity or extended perimortem agonal phase, such as a brain pH

below 6.2 or an RNA integrity number (RIN) less than 6.0.

Pre-processing of data

We performed the quality control analysis of the RNA-Seq raw sequence data using fastp for

trimming of low-quality data, and trimming of adapters(Chen et al. 2018). FastQC

(https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and multiqc(Ewels et al. 2016)

were used to assess the quality of data.

Alignment of reads and RNA-Seq variant identification

The mapping of RNA-Seq reads to a UCSC reference human genome hg38 reference genome

was done using a STAR aligner with 2-pass mode because it increases sensitivity(Veeneman et

al. 2016). STAR is also recommended in GATK best practices workflow (RNAseq short variant

discovery (SNPs + Indels) – GATK (broadinstitute.org) for the identification of short variants

from RNA-Seq data and to get better alignments around novel splice junctions. From the STAR

alignments's Ist pass mode, *SJ.out.tab* was generated consisting of all the junctions obtained

from the 1-pass mapping step and was used as the input for the 2^{nd} pass mode. Sorted BAM files

based on genomic coordinates were obtained from the STAR output. The base quality score

recalibration was done to recalibrate base quality scores. GATK HaplotypeCaller was used to

find variants in the genome, including SNV and indels(McKenna et al. 2010a). Picard suite was

used to mark duplicates. The process of splitting reads containing N cigar string into multiple

supplementary alignments, incorporating hard clips to address mismatching overhangs, was

executed using the "SplitNCigarReads" function within GATK4(McKenna et al. 2010b). The

software tool, ANNOVAR was used to study the effect of variants on gene and protein function,

for the interpretation and prioritization of single nucleotide variants and Indels(Wang et al.

2010). SnpEff, an independent open-source platform used for annotating these variants using

databases, filters, and manipulating genomic annotated variants based on their genomic

locations, such as intronic, untranslated region (5′ UTR or 3′UTR), upstream, downstream, splice

site, or intergenic regions(Cingolani et al. 2012). SnpSift CaseControl tool was used for a case-

control study of opioid(n=30) and non-opioid(n=20) subjects and filter data to find out the most

significant variants(Cingolani et al. 2012). dbSNP was utilized for the retrieval of genetic

variants, along with the acquisition of allele population frequencies sourced from databases such

as 1000 Genomes, gnomAD, dbGaP, and TOPMED, among others(Sherry et al. 2001).

Variant Prioritization

For variant prioritization, variant scores were normalized by applying the quality filtration using

the "VariantFiltration" function of the Genome Analysis Toolkit (GATK), including Variant

confidence normalized by unfiltered depth of variant samples QualByDepth (QD) <2, Strand

bias estimated using Fisher's Exact Test FisherStrand (FS) <60, Strand bias estimated by the

symmetric odds ratio test SOR<9.0 and root mean square mapping quality over all the reads at

the site MQ>40. Subsequently, variants within specified genes (BDNF, DRD2, DRD3, NRXN3,

OPRD1, OPRM1, and NGFB) were extracted using the SnpSift filter function. Amino acid

substitution effects on protein function were predicted using various scores including SIFT,

PolyPhen,   LRT,   MutationTaster,   MutationAssessor_pred,   FATHMM_pred,

fathmm_MKL_coding_pred, PROVEAN_pred, MetaSVM_pred, and MetaLR_pred sourced

from the dbnsfp database(Liu et al. 2016). Potentially deleterious non-synonymous(nsSNPs)

were filtered out if they were predicted to have harmful effects by mutation annotators such as

Mutation Taster, FATHMM, and PROVEAN. [26]. Further, variants were prioritized if they

were exclusively present in Cases (both homozygous and heterozygous) and absent in controls.

Variants affecting both coding and non-coding regions, as well as both known and novel

variants, were prioritized for further analysis. Variants were classified as novel if they were

absent in dbSNP database. Variants were categorized as SNP and were later sorted into missense,

nonsense, and silent mutations if they were present. Utilizing the bcftools plot-vcfstats, a bar plot

elucidating   the   number   of   substitutions   for   each   gene   was   created

(https://samtools.github.io/bcftools/bcftools.html). Bcftools query function was utilized for

counting the total number of SNPs and Indels(https://samtools.github.io/bcftools/bcftools.html).

Results

RNA-Seq Variant frequency analysis

In the study aimed at identifying genetic variants associated with opioid addiction, a Case-

Control analysis was conducted, comparing opioid-addicted subjects (n=30) to non-opioid

subjects (n=20). Analysis using the CaseControl tool of SnpSift revealed a total of 60,13,139

annotated variants, including 3778 SNPs, 485 indels, and variants of unknown class and

significance. Subsequently, variants segregating exclusively in case samples, rather than control

samples, were prioritized to mitigate false positives. Further, investigation focused on variants

within the NRXN3 gene and seven other genes (BDNF, DRD2, DRD3, OPRD1, OPRM1, and

OPRK1 previously linked to opioid addiction(Mistry et al. 2014). We found the absence of any

variant in the NGFB gene. After variant annotation with ANNOVAR and SnpFff, the variants

were filtered based on predetermined thresholds, resulting in variant distributions across various

genomic regions. Among the eight analyzed genes, NRXN3 exhibited the highest number of

variants. Following variant filtration, the number of variants in NRXN3 decreased from 8,944 to

8,939, whereas the variant numbers for the other genes remained unchanged, as detailed in **Table**

I. The variant profiles for the genes, present exclusively in the cases cohort and absent in the

control group, were analyzed. The total number of variants in cases was determined both before

and after applying thresholds, to evaluate the significance of these thresholds on the variants

derived from the RNA-Seq data. **Fig. I** illustrates the variants identified in all seven genes, both

before and following the application of thresholds as described in the Materials and Methods

section. Further, to provide a comprehensive understanding, the frequency of variants present in

control samples was assessed and compared with variants observed in case samples. **Fig. II**

depicts the total variants identified in case and control samples. The bar graph indicates that the

maximum number of variants were found in the case samples, particularly in the NRXN3 gene.

This comparison aimed to elucidate the differences in variant profiles between opioid-addicted

and non-opioid subjects, shedding light on potential genetic contributors to opioid addiction

susceptibility. This also demonstrates the ability of the transcriptome data to identify the

variations accurately. We further present a detailed analysis of the variant profiles for each gene.

In this study we designate variants rsId with known varaints and the varaints with absence of rsId

with novel variants.

Table. I Elucidates variants in the genes before and after applying quality threshold filters

Total variants	60,13,139
Total SNPs	3778
Total Indels	485

Genes	Total variants before applying filter	Total variants after applying thresholds
NRXN3	8944	8939
OPRM1	423	423
DRD2	292	292
OPRK1	104	104
BDNF	192	192
OPRD1	97	97
DRD3	13	13

Known Variants present only in Cases (opioid subjects in at least 25 subjects) and controls

Genes	Total Variants in Cases	Total variants in controls	Known variants (Cases)	Known variants (Controls)
NRXN3	1406	572	438	183
OPRM1	47	43	13	13
BDNF	31	25	2	5
OPRD1	9	9	0	3
DRD3	1	1	0	1

OPRK1	**12**	**13**	**4**	**7**
DRD2	**42**	**24**	**13**	**8**

Fig. I The total number of variants before and after applying thresholds reveals that the gene NRXN3 exhibits the highest number of variants in both conditions. Conversely, the gene DRD3 shows the lowest number of variants.

Fig. II The bar plot displays the count of known variants in both cases and controls, as well as the total variants in both groups. The NRXN3 gene consistently exhibits the highest number of variants.

OPRM1 variants profile

Within the OPRM1 gene located on chromosome 6, our investigation unveiled a total of 47 variations with 13 known mutations, including two Insertion/Deletion variants (INDELs) and various Single Nucleotide Variations (SNVs). The spectrum of these variations encompassed diverse variant types, comprising 3 prime UTR variants, an intron variant, a downstream gene variant, an upstream gene variant, and a non-synonymous variant concurrently exhibiting a stop gained mutation, thereby classified as a HIGH IMPACT mutation. Notably, all mutations, except for one instance of the rs34427887 variant, manifested a heterozygous nature. In contrast, rs34427887 displayed both homozygous and heterozygous manifestations of the stop-gained

mutation. Utilizing Gene Set Enrichment Analysis (GSEA) with annotations sourced from the

MSigDb database, our findings consistently associated all identified variations with the Hallmark

Reactive Oxygen Species Pathway. Concerning the rs34427887 variant, identified with a stop

gained mutation, the Clinical Disease Database in ClinVar underscored its correlation with

Tramadol response. This clinical insight implies the variant's impact on drug response rather than

a specific disease, emphasizing its role in pharmacological responses. The rs34007356 variant

exhibited characteristics of a retained intron. Notably, there were 34 novel variants, including 24

3 prime UTR variants, one missense variant, 5 intron variants, and 3 downstream gene variant

and upstream gene variant instances. The Impact of all these variants was classified as

MODIFIER, except for one with a MODERATE impact due to a missense mutation resulting in

an amino acid change from glutamine to glycine. Intriguingly, all these novel variants exhibited

the same GSEA pathway of Hallmark Reactive Oxygen Species Pathway as that observed in

known variants. **Fig. III** shows the substitution of SNPs for this gene. The maximum number of

substitutions can be seen from A>G and C>T.

Table. II Variants Category based on their genomic region presence

Total variants	Variants in Cases	SNV	INDELS	Variants category				
423	47	10	3	3_prime_UTR_variant	intron_variant	downstream_gene_variant	upstream gene variant	non-synonymous
				32	46	31	4	3

Fig. III The figure shows the number of substitutions from the reference allele to the alternate allele for the gene OPRM1.

DRD2 variants profile

In the investigation of the DRD2 gene on chromosome 11, 13 variants were identified, consisting of one insertion-deletion (INDEL) and 12 single nucleotide variants (SNVs). All observed variants were manifested as heterozygous genotypes. The majority of variants were categorized with the consequence of "MODIFIER," except for one variant characterized with "MODERATE" impact, denoting a missense variant, and two variants with low impact. Notably, the variant, rs114403746, identified a non-synonymous change at exon 4, substituting guanine with thymine at position 414, leading to an amino acid alteration from methionine (M) to isoleucine (I). This variant was classified as of uncertain significance by Intervar automated analysis and predicted as deleterious by multiple tools including PolyPhen, LRT, MutationTaster, and ClinPred within ClinVar. Additionally, 29 novel variants were discovered,

characterized as upstream_gene_variant, downstream_gene_variant, and intron_variant, all

attributed to "MODIFIER" impact. Interestingly, one variant of MODIFIER impact had a

mutation in the exonic region and, was non-synonymous, replacing guanine(G) with thymine(T)

at nucleotide position 414. Among these novel variants, two were annotated as pseudogenes by

SnpEff, while two others were identified as belonging to the biotype of retained_intron. The

remaining novel variants were identified as protein-coding. Utilizing the bcftools query, a total

of 1012 SNPs and 392 INDELs were identified within the examined dataset. **Fig. IV** shows the

maximum number of SNPs of G>A and equal numbers of C>T and G>T. **Table. III** shows the

variant category in the different regions in the genome.

Table. III Variants category based on different regions in the genome for DRD2

Total variants	Variants in Cases	SNV	INDELS	Variants category				
423	47	10	3	3_prime_UTR_variant	intron_variant	downstream_gene_variant	upstream gene variant	non-synonymous
				32	46	31	4	3

Fig. IV The figure shows the number of substitutions from the reference allele to the alternate allele for the gene DRD2.

NRXN3 variants profile

Variants within the NRXN3 gene locus on chromosome 14 were extensively analyzed. Out of the total 8939 variants obtained post-threshold filtration, 1406 were observed in case subjects. Among these, 438 variants were annotated with known dbSNP rsIds, while the remaining 968 were classified as novel variants. The variants were categorized into four biotypes: protein_coding, pseudogene, processed_transcript, and retained_intron. Using the bcftools query, it was determined that among the known variants, 400 were single nucleotide variants (SNVs) and 37 were insertions/deletions (INDELs). These variants encompassed various types, including downstream_gene_variant, intron_variant, upstream_gene_variant, missense_variant, 5_prime_UTR_premature_start_codon_gain_variant, and 3_prime_UTR_variant. One notable finding was the presence of a 5_prime_UTR_premature_start_codon_gain_variant at position

72451988 with rsId rs1022111304, albeit with a low impact. Retained introns emerged as a

significant category of variants, with 170 instances identified in the known variants. No variants

were classified as having a HIGH impact mutation. However, one missense variant was

identified with rs2098424152 of MODERATE impact nature changing from serine to isoleucine

at position 187. Functional impact predictions revealed that the majority of variants were

intron_variant type (419), followed by downstream_gene_variant (28), and

3_prime_UTR_variant (17). The distribution of variants included both homozygous and

heterozygous mutations, with the latter being more prevalent. Overall, the analysis highlighted

the diverse nature of variants within the NRXN3 gene and their potential implications in

molecular functions and disease mechanisms. **Fig. V** shows the maximum number of

substitutions for SNPs of A>G. **Table. IV** shows the variation profile for NRXN3.

Table. IV Variants profile for NRXN3 gene

chromo some	Positi on	dbSNP ID	Muta tion	Functional Impact	Zygosit y	Functi onal Impac t Assess ment	Interv ar autom ated	Pathoge nicity predicti on
14	72451 988	rs10221 11304	T>C	5_prime_UTR_premature_start_ codon_gain_variant	Heteroz ygous	Low		
14	73088 078	rs20991 22312	C>T	Intron variant	Heteroz ygous	Modifi er	Pathog enic	
14	74067 484	rs71530 54	A<C	missense_variant(p.Ser187Ile)	Heteroz ygous	Moder ate	Likely pathog enic	

14	72923 353	rs20984 24152	G>T	missense_variant(p.Ser187Ile)	Heteroz ygous	Modifi er

Fig. V The figure shows the number of substitutions from the reference allele to the alternate allele for the gene NRXN3.

BDNF

Using a defined threshold as mentioned, 192 variants were identified within this gene on chromosome 11. Out of the 192 variants, 2 known variants that were SNVs were obtained on chromosome 11. The rest of the novel variants were 3_prime_UTR and upstream_gene_variants. And the impact of these variants was of MODIFIER type. Out of the 29 novel variants, 3 variants, had HIGH impact mutation, and all three variants had frameshift mutation having loss of function mutation. The frameshift had a change in amino acid from in Arginine at position

307, an amino acid change in serine at position 227, and tyrosine at position 172. The gene ID

for these frameshift variants was CHM13_G0008566, with 17 transcripts affected having a total

transcript affected percentage of 0.88. For novel variants as well as the known ones, the MsigDB

showed the Hallmark epithelial-mesenchymal transition and Hallmark UV response pathways.

All variants both in known and novel variants had heterozygous mutation. The nature of variants

detected in novel variants were 12 3_prime_UTR_variants, 3 frameshift variants and 3 intron

variants, 3 splice_donor_variants & Intron_variant, 5 downstream_gene_variant, and 1

splice_region_variant & non_coding_transcript_exon_variant. The 3 prime UTR variants, intron

variants, and downstream gene variants showed MODIFIER impact. While splice region variants

& intron variants showed a LOW impact. **Fig. VI** shows the maximum number of SNPs of C>T

and G>C.

Fig. VI The figure shows the number of substitutions from the reference allele to the alternate

allele for the gene BDNF.

OPRD1 variants profile

All novel variants were observed was this gene that were nine in number with all 3_prime_UTR_variant but one category of downstream gene variant situated on chromosome 1 All these novel variants exhibited a uniform "MODIFIER" impact characterized by heterozygous genotypes.

DRD3 variants profile

Only one variant was found in case subjects that were novel found on chromosome 3. The type of variant was found to be 3_prime_UTR_variant. The impact of this gene was MODIFIER. This mutation was heterozygous.

OPRK1 variants profile

For OPRK1, the variants were identified on chromosome 8, totaling 12 occurrences exclusively observed in case samples. Among these, 4 variants were recognized as known, while 8 were novel. Within the novel variants, all but one were classified as having MODIFIER impact, except for a single variant characterized by a HIGH impact due to a frameshift mutation affecting leucine at position 348. This frameshift mutation resulted in a loss of function and pertained to the GENE ID CHM13_G0053747, affecting four transcripts, constituting a total proportion of 0.75. Among the known variants, apart from the major MODIFIER, one LOW impact mutation, being synonymous was also identified. For this synonymous variant, the alteration involved a change from valine at position 282. Furthermore, both known and novel variants were annotated to the Hallmark Inflammatory Response pathway. All known variants exhibited heterozygous mutations.

Fig. VII The figure shows the number of substitutions from the reference allele to the alternate allele for the gene OPRK1.

Discussion

In the study done by (Saad et al. 2019b), high-throughput RNA-sequencing on post-mortem midbrain was done to examine the genome-wide changes in the mid-brain gene expression associated with human opioid abuse from chronic opioid users (N□=□30) and drug-free control subjects (N□=□20), as described in the Methods (Saad et al. 2019b). In the current study, we report exploratory variants identified from this gene expression data. Although the original study examined the differentially expressed genes in the mid-brain associated with opioid abuse(Saad et al. 2019b). We exploited this gene expression data (PRJNA492904) to study the variant type associated with opioid abuse. To our knowledge, this study represents the first attempt to identify variants using Case-Control association analysis from RNA-seq gene expression data of human

opioid abusers. We performed variant calling using GATK haplotyecaller and used Snpsift to

perform Case-Control association analysis from 30 Cases and 20 Control Cases. We focused on

the variants (both homozygous and heterozygous) present exclusively in Cases with chronic

opioid use. Variants present in Control samples were excluded from this primary analysis.

However, we analyzed the variants in the Control samples to facilitate a comparison between the

variants in the Control and Case samples. Our findings indicated that out of the eight (DRD2,

DRD3, OPRM1, OPRD1, OPRK1, BDNF, NRXN3, NGFB) genes that were previously

associated with human opioid abuse as reviewed by (Mistry et al. 2014), all genes exhibited

variants except for NGFB.

The analysis revealed a predominance of heterozygous variants across all genes, mainly located

in non-coding regions, suggesting a regulatory role in opioid use disorder (OUD). Additionally,

non-synonymous variants in exonic regions were observed for some novel variants, showing

deleterious mutations. Notably, a frameshift mutation was identified in the OPRK1 gene, and

three frameshift mutations with high impact, as annotated by SnpEff, were found in the BDNF

gene. A heterozygous mutation with rs1022111304 in the NRXN3 gene was exclusively present

in case samples and absent in controls, highlighting its potential significance. Similarly, the

missense mutation in NRXN3 with dbSNP ID rs2098424152 was oberved. These mutations

require further analysis in different populations and experimental validation. The maximum

number of variants were found in NRXN3. The number of substitutions for genes shows the

maximum number of susbstitutions for NRXN3 gene with A>G substitution highest in number.

The *NRXN3* gene is one of the largest genes in the human genome containing 1,826,818 base

pairs(Rowen et al. 2002a). Neurexin-3-alpha is a protein that in humans is encoded by

the *NRXN3* gene(Rowen et al. 2002b). Neurexins (NRXNs) are presynaptic transmembrane

proteins that play a role in the development and function of synapses(Stoltenberg et al. 2011b),

and this gene has also been found to be associated with impulsiveness in patients with substance

abuse(Stoltenberg et al. 2011c). We further did annotation of variants associated with genes

exhibiting variants to determine the functional impact of variants on gene function. Existing

suggestive evidence indicates that polymorphisms associated with NRXN3 induce impulsivity

that leads to risky behaviors, that may give rise to behavioral disorders such as

addiction(Stoltenberg et al. 2011c). This finding may prove to be significant in understanding the

genetic architecture of addiction because impulsivity is an important risk factor for addictions

and other behavioral disorders(Stoltenberg et al. 2011b). Evidence is accumulating that NRXN3

may be involved in a pathway that influences vulnerability to addiction in general(Stoltenberg et

al. 2011b). In that sense, our findings add to the growing evidence that implicates polymorphism

in NRXN3 in risk for substance use problems and/or dependence(Lachman et al. 2007b;

Hishimoto et al. 2007). Given the maximum number of variants found in the NRXN3 gene

compared to other genes, our findings suggest that NRXN3 polymorphisms may be modestly

associated with specific types of impulsivity and substance use problems. Impulsivity increases

the risk for several behavioral disorders such as addictions and therefore is an outstanding

candidate trait for further study. A more complete description of the genetic architecture of

impulsivity is likely to have a substantial impact on our understanding of the genetic architecture

of a variety of behavioral disorders. Furthermore, these patterns may vary among different

populations. Findings such as these, despite their limitations, can be critical for advancing our

understanding of the molecular mechanisms linking variants associated with genes to behavior.

NRXN3 is a promising candidate gene for addiction vulnerability and merits further study. The

majority of clinical genetic testing predominantly targets regions of the genome that encode

proteins. However, the significant role of variants in non-coding regions in penetrant diseases is

increasingly being recognized. Previously, several of the non-coding variants associated with the

substance use disorder were identified(Veerappa and Guda 2024). The application of whole

genome sequencing in clinical diagnostic settings is expanding across a broad spectrum of

genetic disorders. Despite this progress, there is currently no guidance on how existing

guidelines, primarily designed for variants in protein-coding regions, should be adapted for

variants identified in the gene expression region. Due to this reason, we applied variant filtering

thresholds to assess differences in variant counts before and after their application. As shown in

Table. I, the only notable difference in variant numbers is observed in the NRXN3 gene, with no

differences detected in the other genes. Table. I illustrate the differences in variants between

Cases and Controls. The only significant difference in variants is observed in the NRXN3 gene.

This difference is visualized in Fig. II, where both total and known variants are prominently

displayed for the NRXN3 gene alone. In the NRXN3 gene and other genes analyzed in this

study, most variants were located in non-coding regions, specifically of an intronic nature. This

finding suggests a potential involvement of intronic variants in opioid use addiction. Several

studies now indicate a possible association between intronic variants and various types of

addictions

Several studies have identified intronic variants implicated in addiction. For instance, (Ribeiro et

al. 2017) found intronic variants associated with heroin addiction. (Crist et al. 2013) reported a

significant association between an intronic variant and treatment outcomes in opioid dependence.

Additionally, (Haghighatfard et al. 2023) reported the associations of intronic single nucleotide

polymorphisms (SNPs) as a risk factor for internet addiction. These observations aim to increase

the number and range of non-coding region variants that can be clinically interpreted, which,

together with a compatible phenotype, can lead to new diagnoses and catalyze the discovery of

novel disease mechanisms. There is substantial evidence that genetic factors influence an

individual's susceptibility to addiction. However, identifying the specific genes that contribute to

this risk remains a challenging and ongoing endeavor. For a given substance use disorder, there

is an expectation that there are genes that affect addiction vulnerability to that particular

substance, and that there are also genes that influence traits that increase vulnerability to

addiction, in general. Our study offers suggestive evidence that the variants in the NRXN3 might

act as impulsivity factors as a key construct leading to risky behaviors and ultimately opioid

addiction. Further investigation with detailed phenotypic assessment that includes patterns of

substance use, substance use disorder diagnoses, multiple measures of impulsivity, and

additional NRXN3 polymorphisms should be conducted. We recognize that our study has certain

limitations due to the lack of standard guidelines for variant identification from RNA-Seq data

and needs to be interpreted with some caution. More detailed assessments of the potential

associations between important addiction-related phenotypes and genetic variation across

NRXN3 need to be conducted. From the current analysis, we suggest that transcriptomic data can

be used for variant identification which is cost-effective and offers the gene expression and

variants analysis at the same time.

Conclusion

Our comprehensive bioinformatics analysis from a case-control study identified some

characteristic known and unknown variations in opioid-addicted subjects. Although our study

shows characteristic variations in certain genes, an assay comprising multiple biomarkers that are

differentially expressed could be attempted in the future. If this is successful, the number of

biomarkers developed will depend on their validation in a large cohort of patients and the

translation of these findings to clinical practice. Our current study, we believe, is decisive in

understanding the significance of variations in opioid addiction as well as the utilization of

transcriptomic data for variant analysis. Based on the current SNP analysis results, further study

seems warranted to determine the potential association of NRXN3 variants as biomarkers and/or

therapeutic targets for substance abuse.

Legends to Figures and Tables

Figure legends

Fig. I The total number of variants before and after applying thresholds reveals that the gene

NRXN3 exhibits the highest number of variants in both conditions. Conversely, the gene DRD3

shows the lowest number of variants.

Fig. II The bar plot displays the count of known variants in both cases and controls, as well as

the total variants in both groups. The NRXN3 gene consistently exhibits the highest number of

variants.

Fig. III The figure shows the number of substitutions from the reference allele to the alternate

allele for the gene OPRM1.

Fig. IV The figure shows the number of substitutions from the reference allele to the alternate

allele for the gene DRD2.

Fig. V The figure shows the number of substitutions from the reference allele to the alternate

allele for the gene NRXN3.

Fig. VI The figure shows the number of substitutions from the reference allele to the alternate

allele for the gene BDNF.

Fig. VII The figure shows the number of substitutions from the reference allele to the alternate allele for the gene OPRK1.

Table legends

Table. I Variants for genes before and after applying quality threshold filters

Table.II Variants Category based on their genomic region presence

Table. III Variants category based on different regions in the genome for DRD2

Table. IV Variants profile for NRXN3 gene

Acknowledgments

The authors acknowledge the support extended by the All India Institute of Medical Sciences (AIIMS), New Delhi for providing computational facilities.

References

Albertson DN, Schmidt CJ, Kapatos G, Bannon MJ (2006) Distinctive profiles of gene expression in the human nucleus accumbens associated with cocaine and heroin abuse. Neuropsychopharmacology 31:2304–2312. https://doi.org/10.1038/SJ.NPP.1301089

AL-Eitan LN, Rababa'h DM, Alghamdi MA (2021) Genetic susceptibility of opioid receptor genes polymorphism to drug addiction: A candidate-gene association study. BMC Psychiatry 21:1–14. https://doi.org/10.1186/S12888-020-03006-Z/TABLES/5

Bierut LJ (2011) Genetic Vulnerability and Susceptibility to Substance Dependence. Neuron
69:618. https://doi.org/10.1016/J.NEURON.2011.02.015

Chen J-S (2010) A Brighter Side: Protective and Risk Factors in the Rehabilitation of Chronic
Drug Abusers in Hong Kong. http://dx.doi.org/101177/0094306110373238i 39:431–433.
https://doi.org/10.1177/0094306110373238I

Chen S, Zhou Y, Chen Y, Gu J (2018) fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics 34:i884–i890. https://doi.org/10.1093/BIOINFORMATICS/BTY560

Cingolani P, Platts A, Wang LL, et al (2012) A program for annotating and predicting the effects
of single nucleotide polymorphisms, SnpEff: SNPs in the genome of Drosophila
melanogaster strain w1118; iso-2; iso-3. Fly (Austin) 6:80.
https://doi.org/10.4161/FLY.19695

Crist RC, Clarke TK, Ang A, et al (2013) An Intronic Variant in OPRD1 Predicts Treatment
Outcome for Opioid Dependence in African-Americans. Neuropsychopharmacology 2013
38:10 38:2003–2010. https://doi.org/10.1038/npp.2013.99

Davey JW, Hohenlohe PA, Etter PD, et al (2011) Genome-wide genetic marker discovery and

genotyping using next-generation sequencing. Nat Rev Genet 12:499–510.

https://doi.org/10.1038/NRG3012

Egervari G, Kozlenkov A, Dracheva S, Hurd YL (2019a) Molecular windows into the human

brain for psychiatric disorders. Mol Psychiatry 24:653–673.

https://doi.org/10.1038/S41380-018-0125-2

Egervari G, Kozlenkov A, Dracheva S, Hurd YL (2019b) Molecular Windows into the Human

Brain for Psychiatric Disorders. Mol Psychiatry 24:653. https://doi.org/10.1038/S41380-

018-0125-2

Ewels P, Magnusson M, Lundin S, Käller M (2016) MultiQC: summarize analysis results for

multiple tools and samples in a single report. Bioinformatics 32:3047–3048.

https://doi.org/10.1093/BIOINFORMATICS/BTW354

Goldman D, Oroszi G, Ducci F (2005) The genetics of addictions: uncovering the genes. Nat

Rev Genet 6:521–532. https://doi.org/10.1038/NRG1635

Haghighatfard A, Ghaderi AH, Mostajabi P, et al (2023) The first genome-wide association study

of internet addiction; Revealed substantial shared risk factors with neurodevelopmental

psychiatric   disorders.   Res   Dev   Disabil   133:104393.

https://doi.org/10.1016/J.RIDD.2022.104393

Hishimoto A, Liu QR, Drgon T, et al (2007) Neurexin 3 polymorphisms are associated with

alcohol dependence and altered expression of specific isoforms. Hum Mol Genet 16:2880–

2891. https://doi.org/10.1093/HMG/DDM247

Kendler KS, Jacobson KC, Prescott CA, Neale MC (2003) Specificity of genetic and

environmental risk factors for use and abuse/dependence of cannabis, cocaine,

hallucinogens, sedatives, stimulants, and opiates in male twins. Am J Psychiatry 160:687–

695. https://doi.org/10.1176/APPI.AJP.160.4.687

Koob GF, Volkow ND (2010) Neurocircuitry of Addiction. Neuropsychopharmacology 35:217.

https://doi.org/10.1038/NPP.2009.110

Lachman HM, Fann CSJ, Bartzis M, et al (2007a) Genomewide suggestive linkage of opioid

dependence   to   chromosome   14q.   Hum   Mol   Genet   16:1327–1334.

https://doi.org/10.1093/HMG/DDM081

Lachman HM, Fann CSJ, Bartzis M, et al (2007b) Genomewide suggestive linkage of opioid

dependence   to   chromosome   14q.   Hum   Mol   Genet   16:1327–1334.

https://doi.org/10.1093/HMG/DDM081

Levran O, Yuferov V, Kreek MJ (2012) The genetics of the opioid system and specific drug

addictions. Hum Genet 131:823–842. https://doi.org/10.1007/S00439-012-1172-4

Liu QR, Drgon T, Walther D, et al (2005) Pooled association genome scanning: Validation and

use to identify addiction vulnerability loci in two samples. Proc Natl Acad Sci U S A

102:11864–11869.

https://doi.org/10.1073/PNAS.0500329102/SUPPL_FILE/INDEX.HTML

Liu X, Wu C, Li C, Boerwinkle E (2016) dbNSFP v3.0: A One-Stop Database of Functional

Predictions and Annotations for Human Nonsynonymous and Splice-Site SNVs. Hum

Mutat 37:235–41. https://doi.org/10.1002/humu.22932

McKenna A, Hanna M, Banks E, et al (2010a) The Genome Analysis Toolkit: A MapReduce

framework for analyzing next-generation DNA sequencing data. Genome Res 20:1297.

https://doi.org/10.1101/GR.107524.110

McKenna A, Hanna M, Banks E, et al (2010b) The Genome Analysis Toolkit: A MapReduce

framework for analyzing next-generation DNA sequencing data. Genome Res 20:1297.

https://doi.org/10.1101/GR.107524.110

Missler M, Zhang W, Rohlmann A, et al (2003) Alpha-neurexins couple Ca2+ channels to

synaptic vesicle exocytosis. Nature 423:939–948. https://doi.org/10.1038/NATURE01755

Mistry CJ, Bawor M, Desai D, et al (2014) Genetics of Opioid Dependence: A Review of the

Genetic Contribution to Opioid Dependence. Curr Psychiatry Rev 10:156.

https://doi.org/10.2174/1573400510666140320000928

Peacock A, Leung J, Larney S, et al (2018) Global statistics on alcohol, tobacco and illicit drug

use: 2017 status report. Addiction (Abingdon, England) 113:1905–1926.

https://doi.org/10.1111/ADD.14234

Piskol R, Ramaswami G, Li JB (2013) Reliable Identification of Genomic Variants from RNA-

Seq Data. Am J Hum Genet 93:641. https://doi.org/10.1016/J.AJHG.2013.08.008

Ribeiro EA, Scarpa JR, Garamszegi SP, et al (2017) Gene Network Dysregulation in Dorsolateral

Prefrontal Cortex Neurons of Humans with Cocaine Use Disorder. Scientific Reports 2017

7:1 7:1–10. https://doi.org/10.1038/s41598-017-05720-3

Rogier O, Chateigner A, Amanzougarene S, et al (2018) Accuracy of RNAseq based SNP

discovery and genotyping in Populus nigra. BMC Genomics 19:1–12.

https://doi.org/10.1186/S12864-018-5239-Z/FIGURES/6

Rowen L, Young J, Birditt B, et al (2002a) Analysis of the human neurexin genes: Alternative

splicing and the generation of protein diversity. Genomics 79:587–597.

https://doi.org/10.1006/geno.2002.6734

Rowen L, Young J, Birditt B, et al (2002b) Analysis of the human neurexin genes: Alternative

splicing and the generation of protein diversity. Genomics 79:587–597.

https://doi.org/10.1006/geno.2002.6734

Rudd RA, Seth P, David F, Scholl L (2016) Increases in Drug and Opioid-Involved Overdose

Deaths - United States, 2010-2015. MMWR Morb Mortal Wkly Rep 65:1445–1452.

https://doi.org/10.15585/MMWR.MM655051E1

Saad MH, Rumschlag M, Guerra MH, et al (2019a) Differentially expressed gene networks,

biomarkers, long noncoding RNAs, and shared responses with cocaine identified in the

midbrains of human opioid abusers. Scientific Reports 2019 9:1 9:1–9.

https://doi.org/10.1038/s41598-018-38209-8

Saad MH, Rumschlag M, Guerra MH, et al (2019b) Differentially expressed gene networks,

biomarkers, long noncoding RNAs, and shared responses with cocaine identified in the

midbrains of human opioid abusers. Sci Rep 9:. https://doi.org/10.1038/S41598-018-38209-

8

Seney ML, Kim SM, Glausier JR, et al (2021) Transcriptional Alterations in Dorsolateral

Prefrontal Cortex and Nucleus Accumbens Implicate Neuroinflammation and Synaptic

Remodeling   in   Opioid   Use   Disorder.   Biol   Psychiatry   90:550–562.

https://doi.org/10.1016/J.BIOPSYCH.2021.06.007

Sherry ST, Ward MH, Kholodov M, et al (2001) dbSNP: the NCBI database of genetic variation.

Nucleic Acids Res 29:308. https://doi.org/10.1093/NAR/29.1.308

Stoltenberg SF, Lehmann MK, Christ CC, et al (2011a) Associations among types of impulsivity,

substance use problems and Neurexin-3 polymorphisms. Drug Alcohol Depend 119:e31.

https://doi.org/10.1016/J.DRUGALCDEP.2011.05.025

Stoltenberg SF, Lehmann MK, Christ CC, et al (2011b) Associations among types of impulsivity,

substance use problems and Neurexin-3 polymorphisms. Drug Alcohol Depend 119:e31.

https://doi.org/10.1016/J.DRUGALCDEP.2011.05.025

Stoltenberg SF, Lehmann MK, Christ CC, et al (2011c) Associations among types of impulsivity,

substance use problems and Neurexin-3 polymorphisms. Drug Alcohol Depend 119:e31.

https://doi.org/10.1016/J.DRUGALCDEP.2011.05.025

Veeneman BA, Shukla S, Dhanasekaran SM, et al (2016) Two-pass alignment improves novel

splice junction quantification. Bioinformatics 32:43.

https://doi.org/10.1093/BIOINFORMATICS/BTV642

Veerappa A, Guda C (2024) Coordination among frequent genetic variants imparts substance use

susceptibility and pathogenesis. Front Neurosci 18:1332419.

https://doi.org/10.3389/FNINS.2024.1332419/BIBTEX

Volkow ND, Morales M (2015) The Brain on Drugs: From Reward to Addiction. Cell 162:712–

725. https://doi.org/10.1016/J.CELL.2015.07.046

Wang K, Li M, Hakonarson H (2010) ANNOVAR: functional annotation of genetic variants from

high-throughput sequencing data. Nucleic Acids Res 38:e164.

https://doi.org/10.1093/NAR/GKQ603

Zhao Y, Wang K, Wang WL, et al (2019) A high-throughput SNP discovery strategy for RNA-seq

data. BMC Genomics 20:1–10. https://doi.org/10.1186/S12864-019-5533-4/FIGURES/2

World Drug Report 2019 World Drug Report 2019. https://wdr.unodc.org/wdr2019/. Accessed 23

May 2023

Statements and Declarations

Funding

Research reported in this study was supported by the Indian Council of Medical Research (ICMR) under award number *BMI/11(36)/2022* to Swati Ajmeriya.

Competing interests

The authors have no relevant financial or non-financial interests to disclose.

Author Contributions

S.A. designed the analysis, applied methods, and wrote the main manuscript; B.C. reviewed and edited the manuscript; S.K. supervised, edited, and reviewed the manuscript. All authors read and approved the final manuscript.

Availability of data and materials

A publicly available dataset was analyzed in this study. This data can be found in the link NCBI Sequence Read Archive (SRA; www.ncbi.nlm.nih.gov/sra) under accession number PRJNA492904

Ethical approval

This declaration is not applicable

Acknowledgments

The authors acknowledge the support extended by the All India Institute of Medical Sciences

(AIIMS), New Delhi for providing computational facilities and Indian Council of Medical

Research(ICMR) for providing funding.

Author Information

Authors and Affiliations

Department of Biochemistry, All India Institute of Medical Sciences, Ansari Nagar, New

Delhi-110029

Swati Ajmeriya and Subhradip Karmakar

National Drug Dependence Treatment Centre, All India Institute of Medical Sciences, New

Delhi

Biswadip Chatterjee

719