

1 Hearing screening beyond the clinic: Childhood cancer 2 survivors' perceptions of a novel hearing screening 3 program

4 Authors

5 Philippa Jörger^{1,2}, Carina Nigg¹, Leonie Schreck^{1,2}, Christina Schindera^{1,3}, Katharina
6 Fessler¹, Nicolas Waespe^{1,4,5}, Zuzana Tomášiková⁶, Marc Ansari^{5,7}, Annika Frahsa^{1,*},
7 Claudia Kuehni^{1,4,*}✉

9 Affiliations

- 10 1. Institute of Social and Preventive Medicine, University of Bern, Bern, Switzerland
- 11 2. Graduate School for Health Sciences, University of Bern, Bern, Switzerland
- 12 3. Pediatric Oncology/Hematology, University Children's Hospital Basel, University of
13 Basel, Basel, Switzerland
- 14 4. Division of Pediatric Hematology/Oncology, Department of Pediatrics University
15 Children's Hospital Bern and University of Bern, Bern, Switzerland
- 16 5. CANSEARCH research platform in pediatric oncology and hematology Department of
17 Pediatrics, Gynecology and Obstetrics University of Geneva, Geneva, Switzerland
- 18 6. Childhood Cancer Switzerland, Basel, Switzerland
- 19 7. Division of Pediatric Oncology and Hematology, Department of Women, Child, and
20 Adolescent University Hospital of Geneva and University of Geneva, Geneva,
21 Switzerland

22
23 *These authors contributed equally

25 ✉ Corresponding author:

26 Correspondence to:

27 Claudia E. Kuehni, Prof. MD;

28 Institute of Social and Preventive Medicine, University of Bern,

29 Mittelstrasse 43, 3012 Bern, Switzerland;

30 email: claudia.kuehni@unibe.ch; telephone: +41 31 684 35 07

31
32

33 **Abstract**

34 **Purpose:** Childhood cancer survivors (CCS) are at risk for late effects including hearing loss
35 due to ototoxic treatments. It is crucial to detect late effects like hearing loss early, but many
36 adult CCS do not attend recommended follow-up care due to barriers such as accessibility
37 and emotional concerns associated with revisiting medical facilities. To address those
38 barriers, the HEAR-study piloted a new hearing screening program leveraging the extensive
39 network of hearing aid shops across Switzerland. This study explored perceptions of CCS
40 about this novel screening program.

41 **Methods:** We conducted semistructured interviews with 29 participants who completed a
42 hearing test at a hearing aid shop as part of the HEAR-study. We used thematic analysis of
43 interview transcripts, supported by MAXQDA for data analysis.

44 **Results:** Thematic analysis revealed two key themes: First, participants appreciated the
45 program's practicality but noted concerns about the effort of an additional appointment.
46 Second, participants valued the personal and approachable environment in hearing aid
47 shops, but some participants preferred hearing tests as part of an appointment with
48 physicians, valuing immediate advice and contextual knowledge, especially in case hearing
49 loss was found.

50 **Conclusions:** From the perspectives of CCS, this screening program shows promise as a
51 practicable, accessible way to evaluate hearing after childhood cancer. Further evaluation
52 from the perspectives of health care providers is needed to comprehensively assess its
53 feasibility.

54 **Implications for CCS:** For survivors prioritizing convenience or fearing medical institutions,
55 follow-up screenings at accessible, community-based locations like hearing aid shops could
56 provide a practical approach to monitoring hearing.

57

58 **Keywords: (4-6)**

59 Long-term follow-up, hearing screening, hearing problems, childhood cancer, ototoxicity

60 Introduction

61 Childhood cancer survivors are at risk for cancer- and treatment-related long-term
62 health conditions that can occur years after treatment completion [1]. Potential late effects
63 include hearing loss caused by ototoxic treatments such as cranial radiation and platinum
64 chemotherapy, whereby 10% of all childhood cancer survivors develop hearing loss [2]. To
65 detect such late effects early and take appropriate measures, risk-based follow-up care is
66 crucial [3–6].

67 Despite the importance of follow-up care, a significant proportion of adult CCS do not
68 attend care as recommended [7–12]. In Switzerland only half of participating childhood
69 cancer survivors (CCS) are reported to have attended follow-up care [11], while a study from
70 France found even lower follow-up attendance of 33% [7]. In North America and Europe,
71 follow-up care for pediatric and adolescent CCS is initially organized within the pediatric
72 oncology outpatient clinic where survivors received cancer treatment [13–15]. Survivors and
73 their parents discuss results of follow-up examinations with pediatric oncologists and
74 specialists in potentially involved organ systems. Once survivors reach adulthood, CCS are
75 transitioned to adult follow-up care, which is either carried out by a primary care physician,
76 an adult oncologist, or set up within a structured long-term follow-up care (LTFU) program.
77 LTFU programs consist of a multidisciplinary team lead by an internal medicine physician
78 and include specialized nurses, different medical disciplines, social workers, and
79 psychologists [13–15]. In Switzerland, structured LTFU programs for risk-adapted follow-up
80 care have been established in several centers, but they are resource intensive and currently
81 only available for a small proportion of CCS. Previous studies suggest various barriers to
82 follow-up attendance that include lack of knowledge about potential late effects, lack of
83 knowledge of follow-up care possibilities [12,16], long distances to follow-up care
84 appointments [10,12], costs associated with medical appointments [17,18], and emotional
85 barriers related to the cancer experience and fear of pathological findings [19,20].

86 To address some of these barriers, the HEAR-study piloted a new screening program
87 designed to provide a low-threshold option to screen for hearing loss after childhood cancer
88 [21]: In contrast to customary follow-up appointments, the program leverages the extensive
89 network of hearing aid shops distributed throughout Switzerland to offer an easily accessible,
90 alternative site at which survivors may undergo high quality hearing tests that require
91 minimal time, effort, and financial resources from individuals. By taking place outside a
92 hospital or doctor's office, this screening program may encourage follow-up attendance by
93 CCS who might otherwise be reluctant to revisit medical facilities.

94 The overall aim of the HEAR-study is to evaluate the feasibility, limitations, and
95 potential for improvement of this screening program. We therefore interviewed participating
96 CCS who completed a hearing test in a hearing aid shop to learn and interpret their
97 perspectives on such screening, which are crucial to further developing the program.

98

99 **Methods**

100 **Study design**

101 This study is part of the HEAR-study, a health service research project set up in 2021
102 in Switzerland. Details of the HEAR-study are published elsewhere [21]. The HEAR-study
103 piloted a new screening program for hearing loss after childhood cancer. This manuscript
104 presents qualitative data from interviews with HEAR-study participants that provide insights
105 into their experiences, perceived barriers, and reasons for favoring or not favoring this
106 screening program.

107 The Ethics Committee of the Canton of Bern approved the Childhood Cancer
108 Registry (ChCR) and the HEAR-study (166/2014; 2021-01624). We used the Consolidated
109 criteria for Reporting Qualitative research checklist (COREQ) as guidance for reporting [22].
110

111 **Study procedures and participant selection**

112 CCS eligible for the HEAR-study were identified through the Swiss ChCR. The ChCR
113 is a national registry of Swiss residents diagnosed with leukemias, lymphomas, central
114 nervous system (CNS) tumors, malignant solid tumors, or Langerhans cell histiocytosis
115 before 20 years of age and since 1976 [23,24]. We obtained cancer- and treatment-relevant
116 information from the ChCR.

117 All registered CCS who were ≥ 18 years at study start and who were at increased risk
118 for hearing loss due to their treatment—any chemotherapy and/or radiation to head, neck or
119 spine—were eligible for the HEAR-study. We distinguished between two risk groups, high
120 risk, treated with known ototoxic treatments including cranial radiation ≥ 30 Gray or
121 cisplatin/carboplatin, and standard risk, treated with any other chemotherapy or radiation to
122 head, neck or spine with any dose. We excluded CCS with a correspondence language
123 other than German or French due to limited resources, those for whom we had no current

124 postal address, and those who had recently been contacted for another study (January-July
125 2022) so as not to overburden them.

126 We invited eligible CCS (N=1604) to complete a hearing test at a hearing aid shop of
127 a specific branch that is widely distributed across Switzerland. Participants arranged an
128 appointment for themselves at a shop of their choice. Those who completed a hearing test
129 received the results immediately after the test and were encouraged to take them to their
130 physician for further consultation in case of marginal or pathological results. For ethical
131 reasons, acousticians in the hearing aid shops were not informed about participant health
132 history. Prior to participating in the hearing test, participants filled in a baseline questionnaire
133 on sociodemographic characteristics. After completing the hearing test, we sent participants
134 another survey in which they could express interest in being interviewed. Out of 296
135 participants who completed the survey after the hearing test, 169 expressed interest in an
136 interview. We used purposive sampling to select a diverse sample of participants regarding
137 current age, sex, education, corresponding language (German/French), and risk group for
138 hearing loss.

139

140 **Data collection**

141 Invited interview participants could choose face-to-face interviews to be conducted
142 online or in person at a location of their choice. We informed participants about the goals of
143 the study, explained data anonymity, and collected written consent before the interview. We
144 used a semistructured interview guide that focused on participants' experiences at the
145 hearing aid shop and advantages and disadvantages in conducting hearing tests as part of
146 follow-up care at hearing aid shops compared to medical institutions (e.g., hospitals). We
147 piloted the interview guide with study team members and iteratively refined it based on
148 feedback and insights from initial interviews with study participants; changes improved
149 question clarity, sequence, and wording.

150 We planned to interview between 20-30 participants. During the interviews, we
151 continuously analyzed the data. The thirtieth participant who agreed to participate
152 spontaneously canceled the meeting. We decided at this point that the data we had in hand
153 were sufficient to meaningfully explore the diverse perspectives of CCS. We interviewed
154 participants from October 2022 to April 2023, with the main author (PJ, female, PhD student)
155 conducting 24 interviews in German and a trained research assistant (female) conducting 5
156 interviews in French, with PJ providing support in the event of ambiguities or questions
157 arising from participants.

158

159 **Data analysis**

160 All interviews were recorded (length: 17-53 min) and, if conducted in French,
161 translated into German by a bilingual research assistant. During the interviews, the
162 interviewer made notes on interesting aspects discussed and the overall impression of the
163 interviewee for later consideration during data analysis. Another research assistant (KF,
164 female) did verbatim transcriptions into Microsoft Word before importing into MAXQDA 2022
165 (VERBI Software 2021. Consult. Sozialforschung GmbH, Berlin, Germany) for detailed
166 analysis. Another research assistant verified all transcripts against the audio recordings.
167 Braun and Clarke's (2021) approach to reflexive thematic analysis served as guiding
168 framework for data analysis [25], with the research team positioned in a critical realist
169 perspective.

170 Two researchers (PJ and KF), both trained in qualitative research methods,
171 inductively coded a subsample (20%) of interviews independently, guided by the research
172 question. Both discussed their coding from the subsample before agreeing upon a similar
173 coding scheme, which they then discussed with the rest of the research team. This included
174 an established expert in qualitative research (AF, female, professor in community health).
175 After refining the coding scheme with the study team, both researchers coded the remaining
176 transcripts, aligning them with existing themes and subthemes and identifying new ones.

Experiences of CCS with a low-threshold screening program
(Qualitative paper)

Philippa
V00 - 27-Feb-25

177 The ongoing coding was discussed regularly within the whole research team (eight women,
178 two men, with expertise in pediatrics, pediatric oncology, childhood cancer survivorship,
179 social and preventive medicine, community health) to check for trustworthiness of the
180 findings. We produced a brief description of each theme to clarify the content, check for
181 overlaps between themes, and establish final themes and subthemes to present findings
182 clearly and coherently within the narrative. The coders kept reflexive diaries, documenting
183 changes in codes and themes, along with discussions and decisions made throughout the
184 process, and personal thoughts and potential preconceptions. We analyzed the data on a
185 semantic level and identified themes that captured the diverse range of experiences and
186 perspectives of participants with the screening program.

187 Results

188 Sample characteristics

189 Twenty-nine people (15 women) with a median age of 36 years at study start and median
190 age of 12 years at diagnosis participated in the study. Eleven participants were in the high-
191 risk group for hearing loss (cf. further information Table 1). Fifteen interviews took place
192 online, eight were conducted in a meeting room at the University of Bern, while four were
193 conducted at participants' homes and one each in another university meeting room and a
194 workplace.

195 *Table 1: Participant demographic information and*
196 *disease characteristics. (N=29)*

Characteristics	n
Sex	
Male	14
Female	15
Age at study in years (median, range)	36 (18-56)
Age at diagnosis in years (median, range)	12 (0-17)
Diagnosis year	
<1986	3
1986-1995	8
1996-2005	12
>2005	6
Cancer diagnosis	
Leukemias	5
Lymphoma	8
CNS	2
Neuroblastoma	3
Malignant bone tumor	5
Other	6
Risk group	
High risk ^a	11
Standard risk	18
Correspondence language	
German	24
French	5
Education level	
Primary or secondary	16
Tertiary	13
Living area ^b	
Urban	18

Experiences of CCS with a low-threshold screening program
(Qualitative paper)

Philippa
V00 - 27-Feb-25

<i>Intermediate</i>	7
<i>Rural</i>	4
<i>Still visiting regular follow-up care</i>	
Yes	16
No	13
<i>Self-reported hearing loss after the hearing test</i>	
Yes	13
No	16

197 ^aOtotoxic treatments according to International Guideline
198 Harmonization Group guidelines [26]; cisplatin, carboplatin, or cranial
199 radiation ≥ 30 Gray.
200 ^bAccording to the Swiss Federal Office for Statistics; Typology of
201 municipalities and urban-rural typology [27].
202

203 **Themes:**

204 Two main themes represent key perspectives of CCS on the feasibility of the
205 program (Table 2). The first theme focused on practicality, which participants identified as a
206 crucial aspect of follow-up care appointments. The second theme concerned the different
207 circumstances of hearing tests in a hearing aid shop compared to a medical institution (e.g.,
208 hospital) and their advantages and disadvantages. In general, participants reported positive
209 experiences with the hearing test at the hearing aid shop. Only a few reported negative
210 experiences such as feeling tense during the test or experiencing slight discomfort during the
211 examination.

212

213

214

215

216

217 *Table 2: Overview of main themes and corresponding subthemes.*

Main themes	Subthemes
Theme 1	
Appreciating practicality but preferring no additional appointments	<ul style="list-style-type: none"> • Appreciating low effort, daily life compatibility, and simplicity • Preferring no additional appointment or combining it with other follow-up appointments • Preferring to be prompted for new appointment
Theme 2	
Perceiving hearing aid shops as being more accessible and personal but valuing physicians' expertise and familiarity with their medical history	<ul style="list-style-type: none"> • Appreciating personal and professional care competencies • Appreciating relaxed environment • Appreciating accessibility and availability • Appreciating cost free examination • Worrying about being judged in public space • Valuing physicians expertise and familiarity with medical history • Preferring medical infrastructure and wishing for soundproof chamber

218

219 **Appreciating practicality but preferring no additional appointments**

220 Participants prioritized efficiency and practicality in managing follow-up care
 221 appointments such as a hearing test. Participants' experiences at the hearing aid shop
 222 aligned with their wish for practicality. They described procedures at the hearing aid shop as
 223 efficient and uncomplicated: scheduling the appointment was easy, the examination was
 224 carried out swiftly, and it was overall, as a few participants said, "no big deal." They also
 225 liked that no prior consultation or long waiting times were needed, a complaint often raised
 226 when people attend doctors' appointments. Participants perceived the test as compatible
 227 with activities of daily life: they appreciated the options to have the hearing test before
 228 starting work, during their lunch break, or while taking care of their children. Some compared
 229 the effort of the hearing test to other appointments in daily life such as getting a haircut or
 230 buying shoes, and reported they completed their daily activities around the visit.

231

232 *“I found it very straightforward and very quick. I went into the room and after ten*
233 *minutes or so I was out again.”*

234 *18-29-year-old woman¹*

235

236 A few participants experienced the visit to the hearing aid shop as less practical and
237 compatible with everyday life. For them, the effort was higher when the hearing aid shop
238 they visited was not nearby. In these cases, participants visited the hearing aid shop on days
239 off work such as weekends.

240 Several survivors stated they preferred not having any additional appointments in
241 their schedule or preferred a structured LTFU program to avoid additional effort, i.e. having
242 to visit different locations on different days. Some also mentioned that in the event of an
243 abnormal result they would have to attend a doctor's appointment anyway, which would
244 mean even more effort for them.

245

246 *“If you're having a follow-up check-up with the oncologist anyway [...] and you can do it*
247 *all at the same time: a lab appointment, an X-ray appointment and a hearing test at the*
248 *same place on the same date, most people would certainly appreciate that. [...] Then*
249 *you don't have to drive to this place, that place and yet another place. You could do it*
250 *all in one go.”*

251 *30-39-year-old woman*

252

253 Some individuals expressed that the responsibility and organizational effort of
254 scheduling follow-up appointments was burdensome, highlighting the significance of
255 reminders and being invited to appointments. One person proposed a centralized system

¹ All quotes were translated from German or French to English and edited for language and clarity. Quotes in original language are in the appendix.

256 that would provide an overview of all their follow-up appointments, that would remind them of
257 appointments or even help with scheduling.

258

259 **Perceiving hearing aid shops as being more accessible and personal but**
260 **valuing physicians' expertise and familiarity with their health history**

261 Participants were content with personal and professional care competencies of
262 acousticians at the hearing aid shop. The majority particularly appreciated the thorough
263 explanations and information they received during the visit about hearing test procedures,
264 test results, and hearing-related topics raised during the test. Only a few participants were
265 left with open questions about specific topics (e.g., hearing hypersensitivity, tinnitus).

266

267 *"I was warmly welcomed and taken care of. They also told me a bit about what they*
268 *were doing and why, and explained how the hearing [physiology] works. They also*
269 *explained possible hearing impairments [...] and how the test is conducted. I really felt*
270 *very comfortable there and if I had had any questions, I would certainly have been*
271 *able to ask them, but they were actually answered by the person during the process."*

272 *30-39-year-old man*

273

274 Several participants perceived the atmosphere of the hearing aid shop as more
275 relaxed and personal than in a medical facility, which they valued. Some participants
276 appreciated that the acoustician was on their level and that there was more interaction
277 compared to a medical doctor. One person mentioned that this made them feel more
278 comfortable asking questions, indicating that there was a lower barrier to engage with
279 acousticians compared to medical doctors, who were perceived as being less approachable.
280 They considered this relaxed environment as advantageous compared to medical facilities,
281 considering some might have a negative association with medical facilities related to their
282 cancer history.

283

284 *“A hearing aid shop is low threshold. It's not: ‘Ah, I'm going back to the hospital*
285 *now’. Depending on the situation, you might not have the best associations with*
286 *hospitals. And if you're just like anyone else who goes there for counseling, it's*
287 *actually pleasant. It's not the feeling of being a patient, but simply a customer.”*
288 *30-39-year-old woman*

289

290 Survivors appreciated that the hearing aid shops were accessible. One respondent
291 highlighted how the hearing aid shop was more approachable and readily available
292 compared to medical appointments.

293

294 *“It's a place where I can go again at any time. Let's say I want to know something*
295 *more, or I still want to have this photo [otoscopy]. I can just write an email or call.*
296 *The shop is always available. I mean these specialists [doctors], just getting an*
297 *appointment at all... they are somehow so far away. Not so close. Not so easy to*
298 *reach. And that seems to me to be a clear advantage [of the hearing aid shop].”*
299 *40-49-year-old woman*

300

301 A few participants appreciated the test being free of cost. They emphasized that cost
302 of examinations was an important barrier to follow-up care appointments. While health
303 insurance is mandatory in Switzerland, there is a minimum deductible that patients must pay
304 themselves for examinations.

305

306 On the other hand, one participant noted that the hearing test at the hearing aid shop
307 took place in a public space, which could be a barrier for survivors exhibiting visible side or
308 late effects of their therapy. This relaxed environment as described by other participants
309 might lead to them feeling judged by other people who are not physicians.

309

Experiences of CCS with a low-threshold screening program
(Qualitative paper)

Philippa
V00 - 27-Feb-25

310 *"I see a disadvantage [of the hearing aid shop] for someone who [...] still has visible*
311 *sequelae of the illness [later he mentioned hair loss as an example]. That may really*
312 *hold someone back from going to a public place."*
313 *40-49-year-old man*

314

315 Although many participants felt well taken care of at the hearing aid shop, some still
316 perceived physicians as more knowledgeable and trustworthy. They argued that physicians
317 had extensive education and contextual knowledge that is important for conducting
318 examinations and interpreting results in relation to their cancer history.

319

320 *"[Examinations in hearing aid stores] have no medical components. They don't*
321 *have a doctor to interpret [the results]. Not to question the competence of a*
322 *hearing aid acoustician, but they don't have medical staff who know my diagnosis*
323 *and can reliably assess [the hearing]. And that's why I would say the hearing aid*
324 *shop is certainly quicker and better for people who just want a rough assessment*
325 *of how well they can hear. But it's not suitable for people who really need medical*
326 *follow-up treatment and follow-up checks like I do, because they don't meet all the*
327 *requirements. Even if it might be a bit more practical."*

328 *18-29-year-old man*

329

330 A few individuals specifically mentioned that they preferred "receiving bad results"
331 from a physician. Some emphasized the importance of always being examined by the same
332 people for several reasons. They found it burdensome to repeatedly educate new health
333 care professionals about their cancer history, worried about information loss if different
334 health care professionals were involved in the same examination, and saw value in having
335 an established relationship with health care staff, especially for sensitive issues. In relation to
336 this, some participants wished for a person with an overview of their medical history and

337 whom they could contact in case of questions. Nevertheless, many agreed that hearing aid
338 shops could serve as a useful opportunity for initial hearing screening, with referral to a
339 physician in case of unusual findings. For several, integrating the hearing test in everyday
340 life was still more important than the type of institution.

341

342 *“If I had to do such tests more often, the most decisive factor would probably be*
343 *where it is easiest to access... Locally. So it would be easy to integrate into*
344 *everyday life. I wouldn't care whether it was a clinic or a hearing aid shop. Where I*
345 *live, I would probably rather go to the [hospital name]. On the other hand, you wait*
346 *longer in the clinic. At [hearing aid shop name], they "timed" it nicely that it was just*
347 *my turn. That is a further aspect. So above all: what is easiest to integrate into*
348 *everyday life?”*

349 *30-39-year-old woman*

350

351 Some participants expressed concerns about the commercial nature of hearing aid
352 shops, believing that the acousticians performing the tests were motivated by the potential
353 for a sale. A few were worried that acousticians might try to sell them products that they
354 didn't need or want. Final comments concern the professionalism of the hearing test
355 infrastructure. Some perceived the infrastructure not to be as professional as in a medical
356 setting. Mainly, they missed a soundproof chamber for the test, and some were distracted by
357 background noise.

358 Table 3 summarizes some of the advantages and disadvantages of screening
359 hearing in hearing aid shops that were mentioned by participants.

360

361 *Table 3: Possible advantages and disadvantages of screening for hearing loss in hearing aid*
362 *shops mentioned by childhood cancer survivors.*

Advantages	Disadvantages
<ul style="list-style-type: none">• Accessible and easily available• Time efficient and convenient• Personal and approachable environment• No individual costs• Ideal for screening• Less medical environment	<ul style="list-style-type: none">• Burden of additional appointment• Perceiving a lack of contextual knowledge among acousticians• Preferring a physician in case of abnormal findings• Perceiving doctors as more knowledgeable• No soundproof chamber

363

364

365 Discussion

366 Participants reported overall positive experiences with the screening program based
367 in hearing aid shops as proposed within the HEAR-study. They appreciated the practicality
368 and accessibility of hearing aid shops, regarding the shops as a convenient venue for
369 screening. Visits were perceived as relaxed and personal. Participants' overall concerns
370 included the burden of organizing and attending multiple appointments at different
371 institutions, as well as the perception that physicians are better suited to interpret hearing
372 test results in the context of their medical history.

373 Practicality and local availability were important for participants when attending
374 follow-up appointments like hearing tests. These HEAR-study findings confirm those of
375 previous studies on barriers to follow-up care showing that local accessibility facilitates
376 participation [7,12]. Aligning with reports of adult CCS in previous qualitative studies on
377 follow-up care [16,28], we found that participants value structured LTFU programs: they are
378 reluctant to attend multiple appointments in different institutions due to organizational burden
379 and the effort associated with traveling to different locations. However, when structured
380 LTFU programs are not feasible for all CCS, whether due to limited admission capacity or
381 geographical barriers, the here proposed screening program could provide a practical and
382 straightforward option to screen for selected late effects, such as hearing loss. Participants
383 in the HEAR-study highlighted the practicality and simplicity of their visits at the hearing aid
384 shop. Reducing the need for travel and simplifying the appointment process could improve
385 adherence to recommended follow-up care. To implement any such program, a robust
386 organizational framework should be established and CCS should be adequately supported in
387 managing their follow-up appointments. The Survivorship Passport or Passport for Care,
388 electronic treatment summaries including individualized recommendations for follow-up,
389 could guide organization of different appointments within follow-up care [29,30]. Discussions
390 with other stakeholders including pediatric oncologists will shed further light on how to best
391 implement such a program.

392 Previous studies on preferences for and barriers to follow-up care report that adult
393 CCS value personal care and competent staff during examinations [10,20], which we also
394 found in our study. Most participants felt well cared for and professionally consulted during
395 their visits. Participants appreciated the approachability of staff who made it easy for CCS to
396 engage with acousticians during and after the hearing test. The relaxed atmosphere made
397 the experience more comfortable for some CCS, which is particularly important given
398 psychological burden often associated with follow-up visits. This burden, discussed in our
399 own and other studies on preferences and barriers to follow-up care among adult CCS
400 [12,19,20,28], can prevent CCS from returning to hospitals or healthcare providers; a
401 welcoming environment is crucial for encouraging attendance to follow-up care. CCS also
402 valued the hearing test being free of charge, which aligns with previous findings that
403 highlight costs as a considerable barrier to accessing healthcare [31]. Some participants
404 perceived acousticians as less knowledgeable and preferred physicians who could interpret
405 the results of the hearing test in the context of their cancer history. This aligns with findings
406 from earlier research that highlight survivors' apprehensions about new healthcare providers
407 lacking an understanding of their unique healthcare needs during the transition from
408 pediatric to adult follow-up care [20]. That said, a technician customarily will conduct a
409 hearing test in a hospital or general practitioner's office and regardless of whether the
410 hearing test is performed in a hearing aid store or at a medical facility abnormal or
411 inconclusive results in any case require follow-up consultation with a physician. In a setting
412 in which the likelihood of a pathological finding is relatively low (as in our study, with an
413 expected prevalence of 10%) [2], consultation with a physician need not take place in the
414 majority of cases. Implemented more broadly, such a program should be set up so that only
415 people with a pathological finding would then schedule a medical appointment with a
416 physician. This approach could reduce physician workload and thus help alleviate the
417 increasing shortage of doctors in Switzerland [32]. Many participants agreed that this

418 screening approach, when results are likely to be within a normal range, could be used for
419 initial screening.

420

421 **Strengths and limitations**

422 This study demonstrates the potential feasibility of a hearing screening program
423 conducted at hearing aid shops from the perspectives of CCS. Interviewing CCS allowed us
424 to gain insights into their experiences with this program, which are essential to further
425 develop and implement it. By using a semantic approach, we attempted to represent
426 participants' voices as accurately as possible. It is important to recognize that CCS are a
427 diverse group with varying needs and preferences regarding follow-up care. While some
428 may benefit from such a screening option, others may prefer follow-up care solely in clinical
429 settings. The program could be a complementary option available to CCS so that they can
430 choose the setting that best aligns with their needs. Although we used a thorough coding
431 process with multiple rounds of discussions to ensure reliability, there are always different
432 ways to interpret and code data due to the subjective nature of qualitative analysis. Our
433 study recruited participants from the overall HEAR study and includes only their perceptions.
434 We cannot report on the thoughts or perceptions of survivors who were not tested at a
435 hearing aid shop.

436

437 **Conclusion**

438 This study examined the potential of a new hearing screening program designed to
439 provide CCS with a low-threshold alternative to screenings in formal medical settings. Its
440 findings suggest that such a program based in hearing aid shops seems feasible for CCS,
441 and could facilitate greater participation in follow-up care for those who prioritize accessibility
442 and personal atmosphere. Informed by survivors' perspectives, the next step is to explore
443 program feasibility from the perspectives of other stakeholders: health care providers
444 including pediatric oncologists and ear, nose, and throat specialists, and the hearing aid

Experiences of CCS with a low-threshold screening program
(Qualitative paper)

Philippa
V00 - 27-Feb-25

445 company. This will allow us to work further on exploring how such a program could support

446 decentralized follow-up care in Switzerland.

447

448

449 **Acknowledgements**

450 We thank the study team of the Childhood Cancer Research Group, Institute of Social and
451 Preventive Medicine, University of Bern, the Swiss Childhood Cancer Survivor Study, and
452 the team of the Swiss Childhood Cancer Registry. We thank Leah Weber for her valuable
453 work. We thank all childhood cancer survivors participating in the study and for the
454 interesting discussions. We also thank Christopher Ritter for editorial assistance.

455 **Competing interests**

456 Authors PJ, CN, LS, KF, ZT, AF and CK have no relevant financial or nonfinancial interests
457 to disclose. NW reports a relationship with Swedish Orphan Biovitrum AB that includes
458 advisory board membership, consulting, and travel reimbursement and a relationship with
459 Novartis that includes advisory board membership. CS reports a relationship to Swedish
460 Orphan Biovitrum AB that includes travel reimbursement. MA reports a relationship with Jazz
461 Pharmaceutical and Novo Nordisk that includes travel reimbursement. None of these
462 relationships has any association with the current study.

463 **Funding information**

464 This work was financially supported by the Swiss Cancer League and Swiss Cancer
465 Research (grant number HSR-4951-11-2019, KLS/KFS-5711-01-2022, and KFS-5302-02-
466 2021). The CANSEARCH foundation, Kinderkrebs Schweiz Foundation, and Zoé4Life
467 Foundation supported NW.

468

469 **Author contributions**

470 CK, MA and NW contributed to the study conception and design. Data collection and
471 analysis were performed by PJ and KF and critically assessed by CK and AF. The first draft

Experiences of CCS with a low-threshold screening program
(Qualitative paper)

Philippa
V00 - 27-Feb-25

472 of the manuscript was written by PJ, CN, and LS, and all authors commented on previous
473 versions of the manuscript. All authors read and approved the final manuscript.

474

475

476 **References**

- 477 1 Robison LL, Hudson MM. Survivors of childhood and adolescent cancer: life-long
478 risks and responsibilities. *Nat Rev Cancer*. 2014;14:61–70. doi: 10.1038/nrc3634
- 479 2 Weiss A, Sommer G, Kasteler R, *et al*. Long-term auditory complications after
480 childhood cancer: A report from the Swiss Childhood Cancer Survivor Study. *Pediatr*
481 *Blood Cancer*. 2017;64:364–73. doi: 10.1002/pbc.26212
- 482 3 Winther JF, Kenborg L, Byrne J, *et al*. Childhood cancer survivor cohorts in Europe.
483 *Acta Oncol*. 2015;54:655–68. doi: 10.3109/0284186X.2015.1008648
- 484 4 Michel G, Mulder RL, van der Pal HJH, *et al*. Evidence-based recommendations for
485 the organization of long-term follow-up care for childhood and adolescent cancer
486 survivors: a report from the PanCareSurFup Guidelines Working Group. *J Cancer*
487 *Surviv*. 2019;13:759–72. doi: 10.1007/s11764-019-00795-5
- 488 5 International Guideline Harmonization Group. International Guideline Harmonization
489 Group. 2024. <https://www.ighg.org/> (accessed 1 February 2025)
- 490 6 Skinner R, Wallace WHB, Levitt G. Long-term follow-up of children treated for cancer:
491 why is it necessary, by whom, where and how? *Arch Dis Child*. 2007;92:257–60. doi:
492 10.1136/adc.2006.095513
- 493 7 Dumas A, Milcent K, Bougas N, *et al*. Predictive factors of long-term follow-up
494 attendance in very long-term childhood cancer survivors. *Cancer*. 2023;129:3476–89.
495 doi: 10.1002/cncr.34944
- 496 8 Rebholz CE, von der Weid NX, Michel G, *et al*. Follow-up care amongst long-term
497 childhood cancer survivors: a report from the Swiss Childhood Cancer Survivor Study.
498 *Eur J Cancer*. 2011;47:221–9. doi: 10.1016/j.ejca.2010.09.017
- 499 9 Michel G, Kuehni CE, Rebholz CE, *et al*. Can health beliefs help in explaining
500 attendance to follow-up care? The Swiss childhood cancer survivor study.
501 *Psychooncology*. 2011;20:1034–43. doi: 10.1002/pon.1823

Experiences of CCS with a low-threshold screening program
(Qualitative paper)

Philippa
V00 - 27-Feb-25

- 502 10 Michel G, Gianinazzi ME, Eiser C, *et al.* Preferences for long-term follow-up care in
503 childhood cancer survivors. *Eur J Cancer Care (Engl)*. 2016;25:1024–33. doi:
504 10.1111/ecc.12560
- 505 11 Baenziger J, Roser K, Mader L, *et al.* Can the theory of planned behavior help explain
506 attendance to follow-up care of childhood cancer survivors? *Psychooncology*.
507 2018;27:1501–8. doi: 10.1002/pon.4680
- 508 12 Ernst M, Brähler E, Faber J, *et al.* A Mixed-Methods Investigation of Medical Follow-
509 Up in Long-Term Childhood Cancer Survivors: What Are the Reasons for Non-
510 Attendance? *Front Psychol*. 2022;13:846671. doi: 10.3389/fpsyg.2022.846671
- 511 13 Guilcher GMT, Fitzgerald C, Pritchard S. A questionnaire based review of long-term
512 follow-up programs for survivors of childhood cancer in Canada. *Pediatr Blood
513 Cancer*. 2009;52:113–5. doi: 10.1002/pbc.21701
- 514 14 Eshelman-Kent D, Kinahan KE, Hobbie W, *et al.* Cancer survivorship practices,
515 services, and delivery: a report from the Children's Oncology Group (COG) nursing
516 discipline, adolescent/young adult, and late effects committees. *J Cancer Surviv*.
517 2011;5:345–57. doi: 10.1007/s11764-011-0192-8
- 518 15 Essig S, Skinner R, von der Weid NX, *et al.* Follow-Up Programs for Childhood
519 Cancer Survivors in Europe: A Questionnaire Survey. *PLoS One*. 2012;7:e53201. doi:
520 10.1371/journal.pone.0053201
- 521 16 Hendriks MJ, Harju E, Michel G. The unmet needs of childhood cancer survivors in
522 long-term follow-up care: A qualitative study. *Psychooncology*. 2021;30:485–92. doi:
523 10.1002/pon.5593
- 524 17 Cai J, Cheung YT, Hudson MM. Care Models and Barriers to Long-Term Follow-Up
525 Care Among Childhood Cancer Survivors and Health Care Providers in Asia: A
526 Literature Review. *JCO Glob Oncol*. 2024;10:e2300331. doi: 10.1200/GO.23.00331

Experiences of CCS with a low-threshold screening program
(Qualitative paper)

Philippa
V00 - 27-Feb-25

- 527 18 Howard AF, Kazanjian A, Pritchard S, *et al.* Healthcare system barriers to long-term
528 follow-up for adult survivors of childhood cancer in British Columbia, Canada: a
529 qualitative study. *J Cancer Surviv.* 2018;12:277–90. doi: 10.1007/s11764-017-0667-3
- 530 19 Casillas J, Kahn KL, Doose M, *et al.* Transitioning childhood cancer survivors to adult-
531 centered healthcare: insights from parents, adolescent, and young adult survivors.
532 *Psychooncology.* 2010;19:982–90. doi: 10.1002/pon.1650
- 533 20 Rosenberg-Yunger ZRS, Klassen AF, Amin L, *et al.* Barriers and Facilitators of
534 Transition from Pediatric to Adult Long-Term Follow-Up Care in Childhood Cancer
535 Survivors. *J Adolesc Young Adult Oncol.* 2013;2:104–11. doi:
536 10.1089/jayao.2013.0003
- 537 21 Jörger P, Nigg C, Mader L, *et al.* Protocol for a low-threshold screening program to
538 detect hearing loss in childhood cancer survivors in Switzerland: The HEAR-study
539 (Preprint). *JMIR Preprints.* 2024.
- 540 22 Tong A, Sainsbury P, Craig J. Consolidated criteria for reporting qualitative research
541 (COREQ): a 32-item checklist for interviews and focus groups. *Int J Qual Health Care.*
542 2007;19:349–57. doi: 10.1093/intqhc/mzm042
- 543 23 Childhood Cancer Registry. Childhood Cancer Registry (ChCR). 2025.
544 <https://www.childhoodcancerregistry.ch/> (accessed 1 February 2025)
- 545 24 Michel G, von der Weid NX, Zwahlen M, *et al.* The Swiss Childhood Cancer Registry:
546 rationale, organisation and results for the years 2001-2005. *Swiss Med Wkly.*
547 2007;137:502–9. doi: 10.4414/smw.2007.11875
- 548 25 Braun V, Clarke V. *Thematic Analysis: A Practical Guide.* SAGE 2021.
- 549 26 Clemens E, van den Heuvel-Eibrink MM, Mulder RL, *et al.* Recommendations for
550 ototoxicity surveillance for childhood, adolescent, and young adult cancer survivors: a
551 report from the International Late Effects of Childhood Cancer Guideline
552 Harmonization Group in collaboration with the PanCare Consortium. *Lancet Oncol.*
553 2019;20:e29–41. doi: 10.1016/S1470-2045(18)30858-1

Experiences of CCS with a low-threshold screening program
(Qualitative paper)

Philippa
V00 - 27-Feb-25

- 554 27 Federal Statistical Office. Typology of municipalities and urban-rural typology. Federal
555 Statistical Office. 2017. [https://www.bfs.admin.ch/bfs/en/home/statistics/territory-](https://www.bfs.admin.ch/bfs/en/home/statistics/territory-environment/nomenclatures/gemtyp.html)
556 [environment/nomenclatures/gemtyp.html](https://www.bfs.admin.ch/bfs/en/home/statistics/territory-environment/nomenclatures/gemtyp.html) (accessed 12 September 2024)
- 557 28 Lie HC, Mellblom A V, Brekke M, *et al.* Experiences with late effects-related care and
558 preferences for long-term follow-up care among adult survivors of childhood
559 lymphoma. *Support Care Cancer*. 2017;25:2445–54. doi: 10.1007/s00520-017-3651-6
- 560 29 Haupt R, Essiaf S, Dellacasa C, *et al.* The ‘Survivorship Passport’ for childhood
561 cancer survivors. *Eur J Cancer*. 2018;102:69–81. doi: 10.1016/j.ejca.2018.07.006
- 562 30 Horowitz ME, Fordis M, Krause S, *et al.* Passport for care: implementing the
563 survivorship care plan. *J Oncol Pract*. 2009;5:110–2. doi: 10.1200/JOP.0934405
- 564 31 Levesque J-F, Harris MF, Russell G. Patient-centred access to health care:
565 conceptualising access at the interface of health systems and populations. *Int J Equity*
566 *Health*. 2013;12:18. doi: 10.1186/1475-9276-12-18
- 567 32 FMH Swiss Medical Association. FMH-Aerztstatistik. 2023.
568 <https://www.fmh.ch/themen/aerztstatistik/fmh-aerztstatistik.cfm> (accessed 16
569 December 2024)

570

571