Development and clinical validation of blood-based multibiomarker models
for the evaluation of brain amyloid pathology

*Darren M. Weber,[1] *Matthew A. Stroh, [1] Steven W. Taylor,[1] Robert J. Lagier,[1] Judy Z. Louie,[1]
Nigel J. Clarke,[1] David E. Vaillancourt,[2,3,6,8] Sruti Rayaprolu,[4,5] Ranjan Duara,[3,7] Michael K.
Racke[1]

[1] Quest Diagnostics Nichols Institute, San Juan Capistrano, CA

[2] Department of Applied Physiology and Kinesiology, University of Florida, Gainesville, Florida,
USA

[3]1Florida ADRC, University of Florida, Gainesville FL

[4] Department of Neurology, University of Florida, Gainesville, FL

[5] Center for Translational Research in Neurodegenerative Disease, 1Florida ADRC, University of
Florida, Gainesville, FL

[6] Norman Fixel Institute for Neurological Diseases, University of Florida, Gainesville, Florida,
USA

[7] Wien Center for Alzheimer's Disease and Memory Disorders, Mount Sinai Medical Center,
Miami Beach, Florida, USA

[8] J. Crayton Pruitt Family Department of Biomedical Engineering, University of Florida,
Gainesville, Florida, USA

* These authors contributed equally to this work and share first authorship

Corresponding Author:

Darren Weber
darren.m.weber@questdiagnostics.com
1-949-728-4668
Quest Diagnostics Nichols Institute
33608 Ortega Hwy, B207
San Juan Capistrano, CA 92675
31

Word count: 4,487

ABSTRACT

Background and Objectives: Plasma biomarkers provide new tools to evaluate patients with

mild cognitive impairment (MCI) for Alzheimer's disease (AD) pathology. Such tools are needed

for anti-amyloid therapies that require efficient and accurate diagnostic evaluation to identify

potential treatment candidates. This study sought to develop and evaluate the clinical

performance of a multi-marker combination of plasma beta-amyloid 42/40 (Aβ42/40), ptau-217,

and *APOE* genotype to predict amyloid PET positivity in a diverse cohort of patients at a

memory clinic and evaluate >4,000 results from "real-world" specimens submitted for high-

throughput clinical testing.

Methods: Study participants were from the 1Florida AD Research Center (ADRC).

Demographics, clinical evaluations, and amyloid PET scan data were provided with plasma

specimens for model development for the intended-use cohort (MCI/AD: n=215). Aβ42/40 and

ApoE4 proteotype (reflecting high-risk *APOE* ☐4 alleles) were measured by mass spectrometry

and ptau-217 by immunoassay. A likelihood score model was determined for each biomarker

separately and in combination. Model performance was optimized using 2 cutpoints, 1 for high

and 1 for low likelihood of PET positivity, to attain ≥90% specificity and sensitivity. These

cutpoints were applied to categorize 4,326 real-world specimens and an expanded cohort

stratified by cognitive status (normal cognition [NC], MCI, AD).

Results: For the intended-use cohort (46.0% prevalence of PET-positivity), a combination of

Aβ42/40, ptau-217, and *APOE4* allele count provided the best model with a receiver operating

characteristic area under the curve (ROC-AUC) of 0.942 and with 2 cutpoints fixed at 91%

sensitivity and 91% specificity yielding a high cutpoint with 88% positive predictive value (PPV)

and 87% accuracy and a low cutpoint with 91% negative predictive value (NPV) and 85%

accuracy. Incorporating *APOE4* allele count also reduced the percentage of patients with

indeterminate risk from 15% to 10%. The cutpoints categorized the real-world clinical specimens

as having 42% high, 51% low, and 7% indeterminate likelihood for PET positivity and

differentiated between NC, MCI, and AD dementia cognitive status in the expanded cohort.

Discussion: Combining plasma biomarkers Aβ42/40, ptau-217, and *APOE4* allele count is a

scalable approach for evaluating patients with MCI for suspected AD pathology.

<u>**Key Takeaways:**</u>

1.) The approval of disease-modifying therapies for Alzheimer's disease ushers in the need

for accessible, affordable, and accurate blood-based testing for Alzheimer's pathology.

2.) Models implementing multiple analytes have demonstrated high performance in

identifying patients with brain amyloid pathology.

3.) We developed high-throughput, robust, multiple-analyte assays and models aimed at

predicting the likelihood of amyloid PET positivity.

4.) We report two models with excellent performance in alignment with current

recommendations for blood-based testing.

5.) Aβ42/40 + ptau-217 + *APOE4* allele count provided the best prediction for amyloid PET

positivity when sensitivity and specificity were both fixed at 91%.

1. INTRODUCTION

Alzheimer's disease (AD) is a progressive neurodegenerative disorder characterized by

beta-amyloid (Aβ) plaques and neurofibrillary tangles in the brain that are associated with

cognitive impairment.[1] Previously, PET imaging [2] or CSF evaluation were the only available

approaches to determine levels of Aβ and phosphorylated tau protein[3] (ptau, a marker of

neurofibrillary tangles) for diagnosis of AD pathology. However, these methods are perceived

as invasive and/or costly. Recently approved disease-modifying monoclonal antibody treatments

for AD target specific soluble aggregates of Aβ[4,5] and are indicated for individuals with mild

cognitive impairment (MCI) or mild AD. Eligibility for these treatments requires confirmation of

amyloid pathology and *APOE* □4 allele (*APOE4*) carrier status prior to administration. Scalable

diagnostic tests that are accurate, less invasive, and affordable can fulfill this need.[6]

Recent advances in blood-based biomarkers (BBM) provide clinicians with a more cost-

effective and accessible means to evaluate biomarkers such as Aβ and ptau.[7] High-throughput

mass spectrometry (MS) and immunoassay platforms combined with automated sample

preparation provide a scalable approach to measure large numbers of blood specimens with

fast turn-around times. Recently, we developed a liquid chromatography-tandem MS (LC-

MS/MS) plasma test to evaluate cerebral Aβ pathology using a ratio of peptides beta-amyloid 42

(Aβ42) to beta-amyloid 40 (Aβ40) (ie, Aβ42/40) to predict amyloid PET status.[8,9] In addition,

plasma tau protein phosphorylated at threonine 217 (ptau-217) is another excellent marker of

AD-related neuropathology.[10]

The Alzheimer's Association recently included Aβ42 and ptau-217 in their guidelines as

BBMs for core AD pathologies and processes.[11] Furthermore, combinations of plasma

biomarkers can provide greater accuracy in identifying and differentiating AD dementia from

other neurodegenerative disorders.[12-16] Additionally, some studies demonstrated that inclusion of

APOE4 allele carrier status or allele count, which confer elevated risk for AD, can improve

model performance when combined with the plasma Aβ42/40 ratio and/or ptau-217,[15,17] while

others showed no statistically significant improvement.[8,14]

For combinations of plasma biomarkers to have substantial clinical value, they should

have acceptable diagnostic performance on par with currently accepted CSF tests.[18] In addition,

BBMs should have high test-retest reliability and scalability for high throughput to be considered

useful for clinical implementation[19,20] as cost-effective diagnostic tests to identify potential

candidates for anti-amyloid therapies.[16,21]

In the current study, our objective was to develop a multibiomarker combination that (1)

achieved acceptable performance metrics while maintaining sensitivities and specificities ≥90%

using specimens from a well-characterized cohort of patients with MCI and AD; and (2) uses

robust high-throughput assays based on accessible cost-effective, low-resolution LC-MS/MS

and immunoassay methods.[9,16]

We note that these recommendations have been met previously using ptau-217 alone,

but in a racially (97% white) and ethnically homogeneous cohort with a high prevalence (64%)

of amyloid positivity[16] and with a model developed using a combination of ptau-217/np-tau217

and Aβ42/40 (94% white and 53% prevalence of amyloid positivity) on a high-resolution MS-

based platform.[14]

In our cohort, patients were age-, sex-, ethnicity-, and years-of-education matched as

PET-negative (n=116) and PET-positive (n=99), representing a lower prevalence (46%) of PET

positivity and greater ethnic diversity (54% Hispanic) than prior studies. In addition to using

plasma-based Aβ42/40 and ptau-217, we examined the effect of adding *APOE*4 genotype

information (ie, carrier or noncarrier vs allele count, ie, 0, 1, or 2 alleles) to the combination of

markers. Ultimately, we aim to assess the usefulness of these tests to effectively identify

candidates for anti-amyloid therapy.

2. METHODS

2.1 Study Design and clinical evaluation

This cross-sectional study included plasma specimens from 215 participants in the intended-use cohort enrolled with 1Florida Alzheimer's Disease Research Center (ADRC). All participants underwent amyloid PET imaging as well as cognitive testing and included 170 participants with MCI and 45 participants with AD (**Table 1**). Participants had a minimum of a 6th grade reading level and spoke English and/or Spanish as their primary language. Participant ethnicity was self-reported.

Participants underwent an extensive neurological and neuropsychological evaluation, including the clinical dementia rating sum of boxes scale (CDRsb)[22] and the mini-mental state examination (MMSE).[23] Diagnoses of MCI and AD dementia were determined using a previously described algorithmic procedure,[24] which combines the neuropsychological diagnosis with the CDRsb scores.

To evaluate the diagnostic performance of the biomarkers for identifying cognitive status, we evaluated and compared additional specimens from 56 participants in the same 1Florida ADRC cohort who were assessed to have normal cognition (NC; 49 PET-negative/7 PET-positive) (**eTable 1**).

In addition, we performed a retrospective analysis of 4,326 consecutive plasma specimens submitted to Quest Diagnostics for plasma Aβ42/40, ptau-217, and ApoE proteoform analysis testing. This was a limited dataset (as defined by https://www.hhs.gov/hipaa/for-professionals/special-topics/research/index.html) with only patient age and sex information available (**eTable 2**).

2.2 Amyloid PET imaging

Amyloid PET imaging was carried out using [18F]florbetaben or [18F]florbetapir tracers on a Siemens 16 Biograph PET/CT scanner, as described previously.[25] Global amyloid status (+/-) was determined based on visual read by an experienced rater who was blinded to diagnosis and cognitive status.[25] All participants underwent an amyloid PET scan within 12 months of plasma collection.

2.3 Plasma biomarker analysis

Blood specimens were collected by venipuncture into 10-mL tubes containing EDTA as anticoagulant and kept on ice (<1 h) until centrifugation at 1,200 relative centrifuge force for 12 min at room temperature. Plasma aliquots (0.5 mL) were transferred into polypropylene tubes and stored at –80 °C until analysis. All plasma specimens were deidentified, and results were blinded during the analysis.

2.3.1   Plasma Aβ42/40 ratio and ApoE proteoform assays

The LC-MS/MS-based assays for the plasma Aβ42/40 ratio and ApoE proteoforms have been described in detail elsewhere.[9] All sample preparation steps were fully automated using a Hamilton Star liquid handler (Hamilton, Reno, NV). LC-MS/MS was performed using a Transcend Vanquish TLX-4 TurboFlow UPLC, with a high-throughput 4-LC column configuration, coupled to a TSQ Altis Plus Triple Quadrupole MS (both Thermo Fisher Scientific, Waltham, MA); data were analyzed using TraceFinder Clinical Research v5.1 software (Thermo Fisher Scientific).

For the Aβ42/40 ratio, Aβ40 and Aβ42 were enriched by immunoprecipitation from plasma (0.5 mL), digested with Lys-C, and desalted and concentrated by solid phase extraction.

Characteristic peptides for Aβ40 (Aβ29-40) and Aβ42 (Aβ29-42) were chromatographically

separated, detected, and quantified by LC-MS/MS.[9]

For ApoE proteoform determination, plasma specimens (25 μL) were reduced, alkylated,

and proteolytically digested with trypsin. Unique tryptic peptides to the ApoE2 and ApoE4

proteoforms, as well as shared peptides between the ApoE2 and ApoE3 proteoforms, and the

ApoE3 and ApoE4 proteoforms, were chromatographically separated and detected by LC-

MS/MS. ApoE proteoforms were determined by the presence or absence of the unique and

shared peptides and used to establish ApoE proteotype, which has previously been shown to be

100% concordant with *APOE* genotype.[9]

Both assays were validated according to Clinical Laboratory Improvement Amendments

(CLIA) guidance. For the Aβ42/40 ratio, assay imprecision (random error) and bias (technical

error) was measured previously to establish the technical robustness[8] of the test per Rabe et

al.[26] Total error was calculated as described in section 2.4 Statistical analysis.

2.3.2   Plasma ptau-217 assay

On the day of the assay, frozen calibrators, quality control samples (QCs), and plasma

specimens were thawed and brought to room temperature. Plasma specimens were centrifuged

to remove any debris, and plasma ptau-217 levels were quantified using the Lumipulse G ptau-

217 - Plasma Immunoreaction Cartridges RUO (for research use only) assay kit on a Lumipulse

G System (Fujirebio Inc, Malvern, PA), which is a fully automated chemiluminescent enzyme

immunoassay (CLEIA) platform.

The assay was validated according to CLIA guidance and data were used to evaluate

technical robustness per Rabe et al.[26] Inter-assay imprecision (random error) was determined

by taking the average of 2 QCs analyzed in duplicate over 14 nonconsecutive days. Bias

(technical error) was experimentally determined by analyzing 100 previously assayed plasma

specimens using a different lot of reagents. Values for ptau-217 were compared as a percentile

difference calculated between the mean values of the original results and retest results, as has

been reported elsewhere.[8,26] Total error was calculated as described in Section 2.4.3.

2.4   Statistical analysis

2.4.1 Demographic, clinical, and biomarker comparisons

Participant demographics were summarized for the full cohort and further stratified by

amyloid PET status. Continuous variables were summarized as mean and standard deviation

(SD) or median and interquartile range. Categorical variables were summarized as percent and

count (N). Tests for difference between PET status were assessed by 2-sample t-test or

Wilcoxon Rank-Sum test for continuous variables and Fisher's Exact test for categorical

variables. Significance was determined with a threshold of $P \leq .05$.

2.4.2 Logistic regression

Logistic regression modeling was employed to assess the association between predictor

variables and amyloid PET status to estimate model linear predictors. The ptau-217 variable as

a predictor in the models was natural log transformed.[27] Akaike Information Criterion (AIC) was

used to select the final model. Amyloid PET positivity likelihood scores in probability scale were

calculated by transforming model linear predictors in natural exponential function. Tests for

differences in the distribution of amyloid PET positivity likelihood scores between amyloid PET

status were assessed by Wilcoxon Rank-sum test. Model performances of predictors in

predicting PET positivity were evaluated by ROC curve analysis. Differences between ROC

curves were assessed by the method of DeLong.[28] Comparisons of probability/likelihood scores

when stratified by amyloid PET status were conducted using a two-tailed paired t-test.

In addition, model performances were also evaluated by a 2-cutpoint approach with the

lower cutpoint establishing maximized sensitivity and the upper cutpoint establishing maximized

specificity. PPV and NPV were calculated at each classification threshold via Bayes' rule, with

estimated sensitivity, specificity, and prevalence as inputs. Recommended performance criteria

stipulate that those patients who fall in the indeterminate zone between the two cutpoints

represent ≤20% of the population.[18]

2.4.3 Technical robustness

To investigate the effects of total error (assay imprecision+potential bias) for the Aβ40, Aβ42,

and ptau-217 markers on the predictive performance of the best model, data simulations were

performed. Estimates for analytical imprecision and bias were previously established for each

marker (data not included). Based on those error estimates, Monte Carlo simulations were

performed to generate 10,000 replicate study data sets (N=215) with randomly assigned errors

drawn from a Gaussian distribution (mean=0, %CV=6%) and fixed bias offsets added to each

biomarker value. Aβ42/40 ratios were generated for each paired set of perturbed amyloid

markers. For each simulated data set, observations were scored by the best model (each

observation retained its original *APOE4* allele count for scoring) and assigned a new predicted

PET+ class (low, indeterminate, or high likelihood) based on the established classification

thresholds. Predicted PET+ classes for each simulated data set were tabulated against the

original (non-perturbed) data and reclassification rates were estimated. The average (95% CI)

reclassification rate due to overall assay error was estimated as the median (2.5[th] percentile,

97.5[th] percentile) of the 10,000 simulated reclassification rates.

All data analyses were performed using R version 4.3.1 (The R Foundation for Statistical

Computing).

2.5 Data availability

The raw data supporting the conclusions of this article will be made available by the authors

on request.

2.6 Ethics statement

The studies involving humans were approved by Mount Sinai Medical Center IRB. The

studies were conducted in accordance with the local legislation and institutional requirements.

The human samples used in this study were acquired as part of a previous study for which

ethical approval was obtained. Written informed consent for participation from the participants or

the participants' legal guardians/next of kin in accordance with the national legislation and

institutional requirements.

3. RESULTS

3.1 Demographic, clinical, and biomarker comparisons

The mean age of participants in the intended-use cohort was 79 years (SD, 8 years;

range 52-97) with almost 80% having MCI and the remaining having AD dementia (**Table 1**).

Women (56%) were slightly more represented than men and there was a slight majority (54%)

of participants with Hispanic ethnicity (56% of these participants were of Cuban origin, **Table 1**).

Most participants had a CDRsb score >0 (94%) and an MMSE score ≥25 (nearly 75%, **Table 1**).

Compared to PET-negative participants, PET-positive participants had a significantly

higher percentage with a CDRsb score >0 (P=.040) and significantly lower median MMSE

scores (P<.001, **Table 1**). No significant differences between patient age, sex, years-of-

education, and ethnicity were found between those who were PET-negative and PET-positive.

Compared to PET-negative participants, PET-positive participants had significantly lower

mean levels of plasma Aβ42, higher median levels of ptau-217, and a significantly lower mean

Aβ42/40 ratio (P<.001 for all, **Table 1**). No difference in plasma Aβ40 levels between PET-

positive and PET-negative groups was observed.

3.2 Prediction of amyloid PET status

As described below, the individual and multi-marker prediction of amyloid PET status

using either single or 2-cutpoint approaches were evaluated in terms of optimizing test

performance and to meet recommended 90% sensitivity/90% specificity criteria for confirmatory

testing.[18] For the 2-cutpoint approach, this included keeping the percentage of patients

classified as having indeterminate likelihood of PET positivity ≤20%.[18]

3.2.1 Diagnostic performance of individual and combined BBMs for predicting amyloid

PET status using a single cutpoint

ROC-AUCs for PET prediction using Aβ42/40 alone (0.851, 95%CI 0.796 to 0.905) and

ptau-217 alone (0.846, 95%CI 0.808 to 0.905) significantly improved (P<.001) when they were

combined (0.929, 95%CI 0.894 to 0.963) (**eTable 3, Figure 1)**. The AIC for the model that

combined Aβ42/40 and ptau-217 predictors was 195, lower than Aβ42/40 (AIC = 256) and ptau-

217 (AIC = 263) alone (**eTable 3)**. Further improvement on amyloid PET prediction was

achieved when *APOE4* results were added to the model, with *APOE4* carrier status

(positive/negative) yielding an AUC of 0.938 (95%CI 0.906 to 0.969), and *APOE4* allele count

yielding an AUC of 0.942 (95%CI 0.912 to 0.972). However, only the *APOE4* allele count model

AUC (AIC = 186, *P*=0.033), and not the carrier status model (AIC = 189, *P* =0.177), reached

statistical significance when compared to the model without *APOE4* results added. Despite high

AUCs for each combined BBM predictor, none of the models achieved ≥90%

sensitivity/specificity when using a single cutpoint (**eTable 3**, **Figure 1**).

3.2.2 Diagnostic performance of individual and combined BBMs for predicting amyloid

PET status using a 2-cutpoint method

At probability cutpoints establishing ≥90% sensitivity and specificity, no individual

biomarker met the criterion for ≤20% indeterminate classifications.[18] Aβ42/40 classified 34% and

ptau-217 classified 42% of the intended-use population as having an indeterminate result.

Combining Aβ42/40 and ptau-217 reduced the indeterminate results to 15%, and further

addition of *APOE4* allele count reduced them to 10% (**Figure 2, Table 2**). The reduction

represented 21 of 32 participants (66%) being reclassified in the correct direction relative to their

PET status (wrong classification to indeterminate [n=6] or indeterminate to correct classification

[n=15]). The distribution of results for the intended-use population for the Aβ42/40+ptau-

217+*APOE4* allele count model is summarized in **Figure 3A**.

The models with vs without *APOE4* allele count with the 2-cutpoints set for 91%

sensitivity and 91% specificity at the study prevalence of 46% of PET-positivity were compared.

PPVs were 88% with *APOE4* allele count and 87% without and NPVs were 91% for both

models (**Table 3**). At 50% prevalence, which is the prevalence expected in an MCI population,[29]

the estimated PPV and NPV would both reach 90% for the model with *APOE4* allele count

compared to 89% for both PPV and NPV for the model without. At lower prevalence of PET-

positivity, such as in a primary care setting where the prevalence would be closer to 20%,[30] the

model predictors with and without *APOE4* allele count show an estimated NPV of 97% (PPV

69% with, 66% without *APOE4* allele count). Conversely, at high prevalences of PET-positivity

(65%, 80%) the model predictors with and without *APOE4* allele count would have high PPVs

estimates (94%-97%) but lower NPVs (68%-82%, **Table 3**).

3.3 Stratification by cognitive status

Superimposing probability of PET positivity scores from the *APOE4* allele count model

onto cognitive status showed significantly higher probability scores in AD individuals compared

to MCI and NC (*P*<.001 for both), and significantly higher probability scores in MCI vs NC

(*P*=0.002, **eFigure 1A**).

ROC-AUCs for independent model predictors for cognitive status were 0.943 (95%CI

0.894 to 0.992) for NC vs AD and 0.864 (95%CI 0.817 to 0.911) for MCI vs AD (**eFigure 1B**).

However, the model predictor did not perform well by ROC analysis in distinguishing NC from

MCI (AUC = 0.650, 95%CI 0.568 to 0.740).

3.4 Real-world data performance by age

For the 4,326 plasma specimens submitted to Quest Diagnostics for plasma Aβ42/40,

ptau-217, and ApoE proteoform testing, the mean age of individuals was 73 years (SD 10 years,

age range 28-98), most (58%) were female, and 38% had 1 or 2 *APOE4* alleles (**eTable 2**). In

total, 42% (1,822) were classified as having a high likelihood, 51% (2,204) as having low

likelihood, and 7% (300) as having an indeterminate likelihood for PET positivity (**Figure 3B)**.

Based on their low likelihood for PET positivity, a PET scan or CSF testing would

potentially be unnecessary for the 2,204 patients (**Table 4**). Similarly, imaging or CSF testing

may not be required for the 1,822 high-likelihood patients based on the ≥90% sensitivity,

specificity, PPV, NPV (**Table 3**) and recent guidance for a projected a prevalence of 50% of

PET-positivity.[18]

When parsed by the 3-biomarker model predictor designation that uses *APOE4* allele

count, a significant difference in age between the low and high likelihood group (70.3 vs 75.5,

ANOVA *P*<.001) and the low and indeterminate likelihood group (70.3 vs 76.3, ANOVA *P*<.001)

was observed (**eTable 2**). No differences in age were seen between the indeterminate and high

likelihood group.

Further stratifying the results by age, we observed a significant positive relationship

between increasing median age and increasing probability of PET positivity (**eFigure 2**). For low

likelihood patients, the percentage who are likely PET negative decreases with age, with 90.2%

of patients younger than 50 years predicted to be amyloid PET negative compared to 37.2% for

those 90 years or older (**eTable 4**).

3.5 Assay technical robustness and effects of bias and imprecision on likelihood

reclassification

We evaluated the effects of total error, composed of analytical bias (technical error) and

imprecision (random error), on likelihood reclassification when using the model predictor with

APOE4 allele count. In the current study, measured bias was -1.1% (95%CI, -3.1% to 0.9%) for

ptau-217. Previously measured bias was 12% for Aβ42 and 11% for Aβ40.[8] In the current study,

the average measured imprecision for ptau-217, determined by taking the average of 2 QCs

analyzed in duplicate over 14 nonconsecutive days, was 6%. The average imprecision for

Aβ42/40, previously measured by taking the average imprecision across 5 study samples

analyzed in duplicate over 25 days, was also 6%.[8,9]

Reclassification rates are shown in **eTable 5** and visual representation of classification

comparisons with and without total error applied are shown in **eFigure 3**. Median reclassification

after total error was applied across 10,000 simulations was 13.5% (9.8% to 16.7%). Within

these data, most reclassifications were from indeterminate-to-negative: 3.3% (1.4% to 5.6%)

and fewest were from negative-to-positive: 0.9% (0% to 2.3%).

4. DISCUSSION

The need for affordable, scalable, and accessible blood-based testing for Alzheimer's

disease has never been more important than with the emergence of the first disease-modifying

therapies. We demonstrate that the combination of plasma Aβ42/40, ptau-217, and ApoE

proteoform test results, measured using high-throughput LC-MS/MS and immunoassay

methods, can predict the likelihood of amyloid pathology using an optimized model and a 2-

cutpoint approach (**Table 2**). Automated specimen preparation and analysis allows a 96-well

plate to be run in 2 to 8 hours, depending on the assay.

4.1 Model performance

The cutpoints were optimized for detecting amyloid PET status in patients with MCI and

AD, who represent the intended-use population. In addition, these optimized cutpoints are in

alignment with recently published performance recommendations to minimize the number of

indeterminate results to ≤20% while maintaining a sensitivity and specificity of ≥90%.[18] This

ensures that our test can be used to both rule-in and rule-out amyloid PET positivity consistent

with AD pathology as the likely culprit of cognitive impairment. The model minimized the

proportion of indeterminate results to ≤10%, thus limiting results that can be uninformative to

both patients and clinicians.

4.2 Model development

The optimized model was derived from logistic regression combining Aβ42/40, natural

log transformed ptau-217 concentrations, and *APOE4* allele count. Each component provided a

statistically significant contribution to the risk score, except for the homozygous *APOE4*

component. All 12 homozygous *APOE4* individuals in our cohort were PET positive, and the

absence of *APOE4* homozygotes in the PET-negative group provided no statistical control.

APOE4 homozygosity has a prevalence of approximately 2% in the general population.[31]

We found homozygosity to be slightly higher both in our intended-use cohort and in clinical

specimens (about 6%). However, *APOE4* homozygosity combined with PET-negativity would be

expected to be extremely rare. For example, Fortea et al found almost 99% of *APOE4*

homozygotes had some Alzheimer's disease neuropathologic change.[32] The same study

suggested that *APOE4* homozygosity is now being considered a genetically distinct form of

Alzheimer's disease.[32]

The optimized model was slightly improved over Aβ42/40 combined with ptau-217 in

terms of ROC-AUC and had lowest AIC score of all models (**Table 2**). Most importantly, the

model substantially reduced the number of participants with an indeterminate result from 15% to

10%, with about two-thirds of the reclassified participants having a result more predictive of their

actual PET status.

4.3 Comparisons with other models

The performance of combinations of biomarkers in predicting amyloid PET status has

been reported by several different groups,[14-17] including a study by Meyer et al[14] who reported

high predictive performance for a high-resolution MS-based assay incorporating ptau-217

normalized to non-phosphorylated tau (% ptau-217, ptau-217/np-tau-217) and Aβ42/40,

optimized by a single cutpoint. The combined cohort used to develop the model in this study

was less ethnically diverse than ours, although sub cohort analysis indicated that their model

had equivalent accuracy regardless of age, sex, race, ethnicity, or *APOE4* status.

In contrast to the study by Meyer et al, the design of our study incorporated PET-positive

and PET-negative participants, who were age-, sex-, ethnicity-, and years-of-education

matched. While they reported on model robustness for subgroups, Meyer et al did not report on

assay technical robustness, including the effects of technical and random error on patient

classification. Our robustness analysis, which measured total error for both the Aβ42/40 and

ptau-217 assays, demonstrated an overall reclassification rate of <14%, and just over 3% for

the most risk reclassifications between 2 categories.

Other studies showing high performance using fewer biomarkers established PET-

positivity using quantitative centiloid (CL) values with cutoffs of >25 CL or >24 CL indicating PET

positivity. In contrast, we established PET positivity by central visual read,[10,14,16] which is widely

recognized as being highly accurate and the standard to which CL quantification is compared.

Studies have demonstrated a 92% agreement between visual read and CL values. However, CL

values were considered supportive in only 70% of cases that were considered challenging (11%

of scans, expected rate in a memory clinic setting).[33] Results suggest that CL values can be

supportive when added to visual read assessment but have not necessarily been shown to be

superior to the highly accurate visual read approach, with the authors acknowledging that

"quantification should always be done in conjunction with visual assessment to avoid

misclassifications due to potential quantification errors." In addition, the authors note the recent

adoption of opinion from the Committee for Medicinal Products for Human Use (EMACDOC-

17005198181200791) stating that a "threshold of greater than 30 is reflective of established

amyloid pathology at the individual level with high certainty."[33]

4.3 Real-world data

Our real-world data set was composed of 4,326 consecutively run Aβ42/40, ptau-217,

and ApoE proteoform specimens, submitted by healthcare providers, presumably for suspected

dementia due to Alzheimer's disease. As expected, we observed an increase in the likelihood of

PET positivity with age, with predicted PET positivity rates ranging from 9% for individuals under

the age of 50 to 55% for those 90 or older (**eTable 4**). These results suggest that PPV/NPV

could be influenced by age-dependent trends in the prevalence of amyloid PET positivity,

particularly for individuals under the age of 70.

Using our fixed cutpoints to categorize individuals into low, indeterminate, and high

likelihood for amyloid PET positivity, we suggest that results may help guide physicians in

clinical decision making (**Table 4**). Projecting the proportion of PET positivity from the intended-

use population onto the 4,326 clinical specimens could potentially eliminate the need for PET or

CSF testing on 2,204 individuals who are likely to have non-AD etiology, or slightly over 50% of

tested individuals. Considering the cost of an amyloid PET scan in the United States (~$5,000

USD),[34] removing these individuals alone may potentially save $11,020,000 (less the cost of

testing) as well as eliminating travel and wait times to receive imaging services.

4.3 Strengths and limitations

Key strengths of this study include its use of scalable, cost-effective plasma biomarkers

in a cohort of ethnically diverse patients that are representative of the intended-use population.

However, the models were developed using a cohort of participants in a memory care setting,

which may not reflect the general clinical population. This concern is alleviated by similarity in

the distribution between our intended-use cohort and the "real world" clinical specimens across

the 3 risk classifications (**Figure 3**).

In addition, our study was not designed to evaluate the relevance or performance of

these markers in the context of other causes of dementia, including mixed dementia with

amyloid pathology or early onset Alzheimer's disease or to address the effects of ethnically

relevant genetic markers like *APOE4* on predictive performance of the tests. Also, the model will

falsely identify PET-negative *APOE4* homozygote individuals as PET-positives, although these

individuals are expected to be extremely rare. In our expanded cohort, we found one PET-

negative *APOE4* homozygote individual, but they had normal cognition, excluding them from the

intended-use population on which the model was built and predictors were trained.

Notably, *APOE4* status is also leading contributor to risk of amyloid-related imaging

abnormalities (ARIA) with anti-amyloid therapies, however *APOE4* was included solely as a

predictor of amyloid status in this study and results do not provide a quantitative measure of

ARIA risk.[35]

Another limitation is the potential for high false positive rates that may be observed in a

primary care setting (eg, PPV 69%) with low amyloid prevalence (eg, 20%). However, in such

settings, the clinical utility of the models is better suited to rule-out AD pathology, given the high

NPV's (eg, 97%) observed in these low-prevalence populations.

Finally, while the model works well for predicting PET status and differentiates

individuals with different cognitive status, models for predicting cognitive status only work well

for differentiating NC from AD.

Replication of these results in other ethnically and socioeconomically diverse groups will

support using these biomarkers for clinical assessment of potential candidates for anti-amyloid

therapies. The combination of plasma biomarkers described herein is likely to complement

clinical and radiologic diagnostic evaluations to further enhance the accuracy of AD diagnosis,

especially when incorporated into enhanced models. For example, combining these models with

assessments of cognitive function may offer a low-cost enhancement of model performance.

Combining these models with specific imaging methodology to assess diverse

neurodegenerative processes may be important in selecting patients who are most likely to

respond to specific therapeutic interventions.[36]

Study Funding

Funding for this work was provided by NIH Research Grants R01NS052318 and R01NS075012

(D.E. Vaillancourt).

Disclosures

D.M. Weber is an employee of Quest Diagnostics. He also holds patents for the detection of

beta-amyloid by mass spectrometry as well as for the detection of apolipoprotein E proteoforms

by mass spectrometry; M.A. Stroh is an employee of Quest Diagnostics; S.W. Taylor is an

employee of Quest Diagnostics; R.J. Lagier is an employee of Quest Diagnostics; J.Z. Louie is

an employee of Quest Diagnostics; N.J. Clarke is an employee of Quest Diagnostics. He also

holds patents for the detection of beta-amyloid by mass spectrometry as well as for the

detection of apolipoprotein E proteoforms by mass spectrometry; D.E. Vaillancourt reports no

disclosures relevant to the manuscript; S. Rayaprolu reports no disclosures relevant to the

manuscript; R. Duara reports no disclosures relevant to the manuscript; M.K. Racke is an

employee of Quest Diagnostics.

516   **References**

517
1.   Knopman DS, Amieva H, Petersen RC, et al. Alzheimer disease. *Nat Rev Dis Primers.*
2021;7:33.
2.   Chapleau M, Iaccarino L, Soleimani-Meigooni D, et al. The Role of Amyloid PET in
Imaging Neurodegenerative Disorders: A Review. *J Nucl Med.* 2022;63:13S-19S.
3.   Shaw LM, Arias J, Blennow K, et al. Appropriate use criteria for lumbar puncture and
cerebrospinal fluid testing in the diagnosis of Alzheimer's disease. *Alzheimers Dement.*
2018;14:1505-1521.
4.   van Dyck CH, Sabbagh M, Cohen S. Lecanemab in Early Alzheimer's Disease. Reply. *N*
Engl J Med. 2023;388:1631-1632.
5.   Sims JR, Zimmer JA, Evans CD, et al. Donanemab in Early Symptomatic Alzheimer
Disease: The TRAILBLAZER-ALZ 2 Randomized Clinical Trial. *JAMA.* 2023;330:512-
527.
6.   Mattke S, Jun H, Chen E, et al. Expected and diagnosed rates of mild cognitive
impairment and dementia in the U.S. Medicare population: observational analysis.
Alzheimers Res Ther. 2023;15:128.
7.   Canestaro WJ, Bateman RJ, Holtzman DM, et al. Use of a Blood Biomarker Test
Improves Economic Utility in the Evaluation of Older Patients Presenting with Cognitive
Impairment. *Popul Health Manag.* 2024;27:174-184.

8. Weber DM, Taylor SW, Lagier RJ, et al. Clinical utility of plasma Abeta42/40 ratio by LC-
MS/MS in Alzheimer's disease assessment. *Front Neurol.* 2024;15:1364658.
9. Weber DM, Kim JC, Goldman SM, et al. New plasma LC-MS/MS assays for the
quantitation of beta-amyloid peptides and identification of apolipoprotein E proteoforms
for Alzheimer's disease risk assessment. *J Investig Med.* 2024;72:465-474.
10. Ashton NJ, Brum WS, Di Molfetta G, et al. Diagnostic Accuracy of a Plasma
Phosphorylated Tau 217 Immunoassay for Alzheimer Disease Pathology. *JAMA Neurol.*
2024;81:255-263.
11. Jack CR, Jr., Andrews JS, Beach TG, et al. Revised criteria for diagnosis and staging of
Alzheimer's disease: Alzheimer's Association Workgroup. *Alzheimers Dement.*
2024;20:5143-5169.
12. Benedet AL, Brum WS, Hansson O, et al. The accuracy and robustness of plasma
biomarker models for amyloid PET positivity. *Alzheimers Res Ther.* 2022;14:26.
13. Palmqvist S, Stomrud E, Cullen N, et al. An accurate fully automated panel of plasma
biomarkers for Alzheimer's disease. *Alzheimers Dement.* 2023;19:1204-1215.
14. Meyer MR, Kirmess KM, Eastwood S, et al. Clinical validation of the PrecivityAD2 blood
test: A mass spectrometry-based test with algorithm combining %p-tau217 and
Abeta42/40 ratio to identify presence of brain amyloid. *Alzheimers Dement.*
2024;20:3179-3192.
15. Hu Y, Kirmess KM, Meyer MR, et al. Assessment of a Plasma Amyloid Probability Score
to Estimate Amyloid Positron Emission Tomography Findings Among Adults With
Cognitive Impairment. *JAMA Netw Open.* 2022;5:e228392.
16. Figdore DJ, Griswold M, Bornhorst JA, et al. Optimizing cutpoints for clinical
interpretation of brain amyloid status using plasma p-tau217 immunoassays. *Alzheimers*
Dement. 2024;20:6506-6516.
17. West T, Kirmess KM, Meyer MR, et al. A blood-based diagnostic test incorporating
plasma Abeta42/40 ratio, ApoE proteotype, and age accurately identifies brain amyloid
status: findings from a multi cohort validity analysis. *Mol Neurodegener.* 2021;16:30.
18. Schindler SE, Galasko D, Pereira AC, et al. Acceptable performance of blood biomarker
tests of amyloid pathology - recommendations from the Global CEO Initiative on
Alzheimer's Disease. *Nat Rev Neurol.* 2024;20:426-439.
19. Hansson O, Edelmayer RM, Boxer AL, et al. The Alzheimer's Association appropriate
use recommendations for blood biomarkers in Alzheimer's disease. *Alzheimers Dement.*
2022;18:2669-2686.
20. Teunissen CE, Verberk IMW, Thijssen EH, et al. Blood-based biomarkers for Alzheimer's
disease: towards clinical implementation. *Lancet Neurol.* 2022;21:66-77.
21. Tsoy E, La Joie R, VandeVrede L, et al. Scalable plasma and digital cognitive markers
for diagnosis and prognosis of Alzheimer's disease and related dementias. *Alzheimers*
Dement. 2024;20:2089-2101.
22. Morris JC. The Clinical Dementia Rating (CDR): current version and scoring rules.
Neurology. 1993;43:2412-2414.
23. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for
grading the cognitive state of patients for the clinician. *J Psychiatr Res.* 1975;12:189-
198.
24. Duara R, Loewenstein DA, Greig M, et al. Reliability and validity of an algorithm for the
diagnosis of normal cognition, mild cognitive impairment, and dementia: implications for
multicenter research studies. *Am J Geriatr Psychiatry.* 2010;18:363-370.
25. Duara R, Loewenstein DA, Lizarraga G, et al. Effect of age, ethnicity, sex, cognitive
status and APOE genotype on amyloid load and the threshold for amyloid positivity.
Neuroimage Clin. 2019;22:101800.

26. Rabe C, Bittner T, Jethwa A, et al. Clinical performance and robustness evaluation of
plasma amyloid-beta(42/40) prescreening. *Alzheimers Dement.* 2023;19:1393-1402.
27. Brum WS, Cullen NC, Janelidze S, et al. A two-step workflow based on plasma p-tau217
to screen for amyloid beta positivity with further confirmatory testing only in uncertain
cases. *Nat Aging.* 2023;3:1079-1090.
28. DeLong ER, DeLong DM, Clarke-Pearson DL. Comparing the areas under two or more
correlated receiver operating characteristic curves: a nonparametric approach.
Biometrics. 1988;44:837-845.
29. Jansen WJ, Janssen O, Tijms BM, et al. Prevalence Estimates of Amyloid Abnormality
Across the Alzheimer Disease Clinical Spectrum. *JAMA Neurol.* 2022;79:228-243.
30. Roberts RO, Aakre JA, Kremers WK, et al. Prevalence and Outcomes of Amyloid
Positivity Among Persons Without Dementia in a Longitudinal, Population-Based Setting.
JAMA Neurol. 2018;75:970-979.
31. Regy M, Dugravot A, Sabia S, et al. The role of dementia in the association between
APOE4 and all-cause mortality: pooled analyses of two population-based cohort studies.
Lancet Healthy Longev. 2024;5:e422-e430.
32. Fortea J, Pegueroles J, Alcolea D, et al. APOE4 homozygozity represents a distinct
genetic form of Alzheimer's disease. *Nat Med.* 2024;30:1284-1291.
33. Collij LE, Bischof GN, Altomare D, et al. Quantification Supports Amyloid PET Visual
Assessment of Challenging Cases: Results from the AMYPAD Diagnostic and Patient
Management Study. *J Nucl Med.* 2025;66:110-116.
34. Anderson J, Pierce C. PrecivityAD for Diagnosis of Alzheimer Disease. *Am Fam
Physician.* 2022;105:79-81.
35. Withington CG, Turner RS. Amyloid-Related Imaging Abnormalities With Anti-amyloid
Antibodies for the Treatment of Dementia Due to Alzheimer's Disease. *Front Neurol.*
2022;13:862369.
36. DeSimone JC, Wang WE, Loewenstein DA, et al. Diffusion MRI relates to plasma
Abeta42/40 in PET negative participants without dementia. *Alzheimers Dement.*
2024;20:2830-2842.

621 **Figure Captions**
622
623 **Figure 1:** Receiver operating characteristic-area under the curve (ROC-AUC) analysis
624 for individual and combined biomarkers for prediction of PET status. Abbreviations: Aβ,
625 beta-amyloid; *APOE4*, APOE ε4; AUC, area under the curve.
626
Figure 2: Boxplots of likelihood scores stratified by PET status for (A) ptau-217, (B)
Aβ42/40, (C) Aβ42/40 + ptau-217, (D) Aβ42/40 + ptau-217 +*APOE4* carrier status, and
(E) Aβ42/40 + ptau-217 + *APOE4* allele count. Orange dots represent individuals that
are MCI; blue dots represent individuals that are AD. Horizontal dashed lines indicate
high and low likelihood cutoffs. Results between these lines indicate neither a low nor
high likelihood (ie, an indeterminate result).
633

Figure 3: Distribution of model results for A) the intended use population, and B) real-world clinical specimens. Orange dots represent individuals that are amyloid PET negative; blue dots represent those that are amyloid PET positive in panel A. Grey dots represent individuals whose amyloid PET status is unknown. Horizontal dashed lines indicate high and low likelihood cutoffs.

Table 1: Baseline characteristics by amyloid PET status.

Characteristic	Intended Use Cohort [N = 215]	Aβ PET- [N=116]	Aβ PET+ [N = 99]	*P*-value
Demographics				
Age (years)				
Mean (SD) [N]	78.9 (8.4) [215]	78.6 (8.1) [116]	79.3 (8.8) [99]	0.552
Sex				
% female [N]	56.3% [121]	53.4% [62]	59.6% [59]	0.409
Ethnicity				
% Hispanic [N]	54.4% [117]	53.4% [62]	55.6% [55]	0.785
Hispanic Origin				
Central American	7	4	3	
Cuban	65	33	32	
Dominican	1	0	1	
Puerto Rican	9	3	6	
South American	34	22	12	
Missing	1	0	1	
Race: Hispanic				
Black	1	1	0	
Multiracial	1	0	1	
White	114	61	53	

Missing	1	0	1	
Race: Non-Hispanic				
Black	9	5	4	
Multiracial	0	0	0	
White	86	49	37	
Missing	3	0	3	
Education				
Mean (SD) [N]	15.1 (3.3) [214]	15.4 (3.3) [114]	14.8 (3.2) [99]	0.192
Cognitive Assessment				
Diagnosis				
% MCI [N]	79.1% [170]	100% [116]	54.5% [54]	
% AD [N]	20.9% [45]	0% [0]	45.5% [45]	
MMSE				
%MMSE ≥ 25 [N]	74.7% [165]	94.0 [109]	57.7 [56]	*< 0.001*
Median (IQR) [N]	27 (25 to 29) [213]	29 (27 to 30) [116]	25 (21 to 27) [97]	
CDRsb				
% CDRsb > 0 [N]	94.4% [203]	91.4% [106]	98.0% [97]	*0.040*
Plasma Biomarkers				
Plasma Aβ42/40				
Mean (SD) [N]	0.157 (0.028) [215]	0.172 (0.027) [116]	0.141 (0.018) [99]	*< 0.001*
Plasma Aβ42 (pg/mL)				
Mean (SD) [N]	60.4 (18.2)	65.6 (19.2) [116]	54.4 (14.8) [99]	*< 0.001*
Plasma Aβ40 (pg/mL)				
Mean (SD) [N]	383.2 (87.6)	380.5 (87.5) [116]	386.3 (88.2) [99]	0.631
Plasma ptau-217 (pg/mL)				
Median (IQR) [N]	0.26 (0.11 - 0.54) [215]	0.12 (0.06 - 0.25) [116]	0.51 (0.32 - 0.88) [99]	*< 0.001*
ApoE4				
% *APOE4*+ carriers [N]	38.1% [82]	21.6% [25]	57.6% [57]	
% *APOE4* allele count 1 [N]	32.6% [70]	21.6% [25]	45.5% [45]	*< 0.001*
% *APOE4* allele count 2 [N]	5.6% [12]	0% [0]	12.1% [12]	

Abbreviations: Aβ, beta-amyloid; AD, Alzheimer's disease; ApoE4, apolipoprotein E4; CDRsb, clinical dementia rating
sum of boxes; IQR, interquartile range; MMSE, mini-mental state examination; SD, standard deviation. Not all
demographic data were available for all participants; therefore N is the number of participants where data was
available. *P*-values were derived from two-sample t-test or Wilcoxon Rank-Sum test for continuous variables, and
Fisher's Exact test for categorical variables for the Aβ PET+ and Aβ PET- subgroups. *P*-values < 0.05 are highlighted
in   italics.

Table 2: Model performance for prediction of amyloid PET status at a defined sensitivity and specificity
664

Predictor: Two cutpoint

ptau-217	Cutpoint	Aβ PET+, N [%]	All, N [%]	Sensitivity (95% CI)	Specificity (95% CI)	Accuracy, % (95% CI)	AUC (95% CI)	AIC
Low likelihood	< 0.2490	8 [8%]	73 [34%]	92% (85% to 96%)		73% (66% to 78%)	0.846 (0.808 to 0.905)	
Indeterminant likelihood	0.2490 to 0.7554	49 [50%]	91 [42%]		-	-		262.7
High likelihood	> 0.7554	42 [42%]	51 [24%]		92% (86% to 96%)	69% (63% to 75%)		

Aβ42/40	Cutpoint	Aβ PET+, N [%]	All, N [%]	Sensitivity (95% CI)	Specificity (95% CI)	Accuracy, % (95% CI)	AUC (95% CI)	AIC
Low likelihood	< 0.3925	9 [9%]	95 [44%]	91% (84% to 95%)		82% (76% to 86%)	0.851 (0.796 to 0.905)	
Indeterminant likelihood	0.3925 to 0.7236	53 [74%]	72 [34%]		-	-		255.6
High likelihood	> 0.7236	37 [37%]	48 [22%]		91% (84% to 95%)	66% (60% to 72%)		

Model 1: pTau217 + Aβ42/40	Cutpoint	Aβ PET+, N [%]	All, N [%]	Sensitivity (95% CI)	Specificity (95% CI)	Accuracy, % (95% CI)	AUC (95% CI)	AIC
Low likelihood	< 0.3254	9 [9%]	97 [45%]	91% (84% to 95%)		83% (77% to 87%)	0.929 (0.894 to 0.963)	
Indeterminant likelihood	0.3254 to 0.6460	16 [16%]	33 [15%]		-	-		194.6
High likelihood	> 0.6460	74 [75%]	85 [40%]		91% (84% to 95%)	83% (78% to 88%)		

Model 2: ptau-217 + Aβ42/40 + APOE4 carrier status	Cutpoint	Aβ PET+, N [%]	All, N [%]	Sensitivity (95% CI)	Specificity (95% CI)	Accuracy, % (95% CI)	AUC (95% CI)	AIC
Low likelihood	<0.3709	9 [9%]	101 [47%]	91% (84% to 95%)	-	85% (79% to 89%)	0.938 (0.906 to 0.969)	
Indeterminant likelihood	0.3709 to 0.5626	9 [9%]	22 [10%]		-	-		189.2
High likelihood	>0.5626	81 [82%]	92 [43%]		91% (84% to 95%)	86% (81% to 90%)		

Model 3: pTau217 + Aβ42/40 + APOE4 allele count	Cutpoint	Aβ PET+, N [%]	All, N [%]	Sensitivity (95% CI)	Specificity (95% CI)	Accuracy, % (95% CI)	AUC (95% CI)	AIC
Low likelihood	< 0.3714	9 [9%]	101 [47%]	91% (84% to 95%)	-	85% (79% to 89%)	0.942 (0.912 to 0.972)	
Indeterminant likelihood	0.3714 to 0.5375	8 [8%]	21 [10%]		-	-		186.6
High likelihood	> 0.5375	82 [83%]	93 [43%]		91% (84% to 95%)	87% (82% to 91%)		

665 Abbreviations: Aβ, beta-amyloid; AIC, ApoE4, apolipoprotein E4; AIC, Akaike Information Criterion; AUC, area under the curve; CI, confidence interval; PET,
666 positron emission tomography
667

668   **Table 3:** PPV and NPV for each model at different prevalences.

Predictor	ptau-217 + Aβ42/40		ptau-217 + Aβ42/40 +*APOE4* allele count	
Prevalence	**PPV (95% CI)**	**NPV (95% CI)**	**PPV (95% CI)**	**NPV (95% CI)**
20%, as expected in a primary care setting	66% (54% to 79%)	97% (95% to 98%)	69% (56% to 80%)	97% (95% to 99%)
46%, observed study prevalence	87% (78% to 93%)	91% (83% to 95%)	88% (80% to 93%)	91% (84% to 95%)
50%, as expected in MCI population	89% (82% to 94%)	89% (82% to 94%)	90% (84% to 94%)	90% (83% to 95%)
65%, as expected in a general neurology setting	94% (90% to 96%)	82% (72% to 90%)	94% (91% to 97%)	82% (73% to 90%)
80%, as expected in a memory care setting	97% (95% to 98%)	68% (54% to 81%)	97% (95% to 98%)	69% (55% to 81%)

Table 4: Proposed linear predictor cutpoints for clinical decision making

Probability of PET positivity range	Interpretation	Intended use population *N (% of cohort)*	PET positivity *N* (%)	MCI, *N* (%)	AD, *N* (%)	*APOE4, N* (%)	Real world, *N* (% of cohort)	Real world, projections of PET positivity, *N (%)*[a]	Follow-up evaluation and management considerations
< 0.3714	Low likelihood	101 (47.0%)	9 (9.0%)	101 (100%)	0 (0%)	16 (15.8%)	2,204 (50.9%)	198 (9.0%)	Consider non-AD etiology.
0.3714 to 0.5375	Indeterminant likelihood	21 (9.8%)	8 (38.1%)	18 (85.7%)	3 (14.3%)	10 (47.6%)	300 (6.9%)	114 (38.0%)	Consider results within context of full differential; refer to a neurologist.
> 0.5375	High likelihood	93 (43.3%)	82 (88.2%)	51 (54.8%)	42 (45.2%)	56 (60.2%)	1,822 (42.1%)	1,607 (88.2%)	Consider results within context of full differential; refer to a neurologist; AD etiology likely.
All	All	215 (100%)	99 (46.0%)	170 (79.1%)	45 (20.9%)	82 (38.1%)	4,326 (100%)	1,919 (44.4%)	

685 Abbreviations: AD, Alzheimer's disease; ApoE4, apolipoprotein E4 proteoform; MCI, mild cognitive impairment; PET, positron emission tomography.
686 [a]Real world projections of PET positivity are based on percent PET positivity in the intended use population cohort.
687

- ptau-217: AUC = 0.846 [0.808 to 0.905]
- Aβ42/40: AUC = 0.851 [0.796 to 0.905]
- ptau-217 + Aβ42/40: AUC = 0.929 [0.894 to 0.963]
- ptau-217 + Aβ42/40+*APOE4* carrier status: AUC = 0.938 [0.906 to 0.969]
- ptau-217 + Aβ42/40+*APOE4* allele count: AUC = 0.942 [0.912 to 0.972]

A

High likelihood (47%)

Indeterminate likelihood (10%)

Low likelihood (43%)

Probability of PET positivity

Full model with *APOE4* allele count

B

High likelihood (42%)

Indeterminate likelihood (7%)

Low likelihood (51%)

Full model with *APOE4* allele count