Genome-wide association study meta-analysis brings monogenic hearing loss genes

into the polygenic realm

9 Clifford Royce[1,2], Johnson Jacquelyn[1,3], Mackey Caroline E[1,3], Mikita Elizabeth A[1,3], Ryan Allen

10 F[1,2], Million Veteran Program, Maihofer Adam X[1,3], Nievergelt Caroline M[1,3]

1. Veterans Affairs San Diego Healthcare System, Research Service, San Diego, CA, USA

2. University of California San Diego, Department of Otolaryngology – Head and Neck

Surgery, La Jolla, CA, USA

3. University of California San Diego, Department of Psychiatry, La Jolla, CA, US

Corresponding authors: Nievergelt and Clifford

Abstract

Background

Disabling age-related and noise-induced sensorineural hearing loss (SNHL) affects 5% of the

global population and is associated with isolation, depression, cognitive decline, and dementia.

SNHL is strongly heritable with a polygenic signature of common genetic variants conferring

small risks individually, thus requiring very large studies to achieve statistical power in genome-

wide association studies (GWAS).

Methods

We present the first report of a clinical SNHL GWAS in the Million Veteran Program (MVP)

cohort. GWAS findings are examined separately within MVP (210,240 cases and 265,275

controls), including multi-ancestry analysis, then combined and contrasted with the United

Kingdom Biobank (UKB) self-reported hearing loss study (87,056 cases and 163,333 controls).

We perform functional mapping and annotation, gene prioritization, gene-based and gene-set

analysis, and cochlear cell type enrichment, including human single-cell results, then compare

our results to known hereditary hearing loss genes. Summary results are leveraged to

characterize the genetic architecture of SNHL.

Results

Substantial genetic overlap is seen between MVP and UKB despite differences in phenotypes,

demographics, and environmental exposures. Individual GWAS and meta-analyses identify 108

loci, including 54 loci containing novel prioritized genes and/or protein-coding genes. Significant

gene ontology pathways include "sensory perception of mechanical stimulus," "ear

development," "actin-binding," and "cytoskeletal protein binding," indicative of mechanosensory

structure and function in the cochlea. GWAS and gene-based analyses implicate 17 known

familial hearing loss genes, expressing proteins in regions of hair cells, Type 1 neurons, stria

vascularis, and subcellular regions of stereociliae, rootlets, and notably, mechanoelectrical

transduction tip-links. Although risk for polygenic SNHL is predominantly captured by SNPs

outside of congenital HHL genes (HHL), SNP-based partitioned heritability estimates show a

3.26-fold enrichment of HHL relative to other genes.

Conclusion

In the largest GWAS to date, combining clinical diagnosis of hearing loss and self-report data

from two cohorts, we identify 108 loci with 54 novel genes. Despite the increased enrichment of

HHL genes, 97% of the risk for adult SNHL is captured by SNPs outside of HHL genes.

Although SNHL in the UKB and MVP were assessed using different phenotypes, genetic signals

between the two cohorts are predominantly shared, and locus discovery is boosted through

increased sample size in meta-analysis.

Keywords

Hearing loss, genome-wide association studies, congenital hearing loss

Introduction

The effects of permanent sensorineural hearing loss (SNHL) range from annoyance at a

partner's "mumbling" to a disabling disorder that currently affects 1.6 billion people globally, a

number that is predicted to increase by greater than 50% over the next three decades[1]. On a

personal level, SNHL is associated with isolation, depression, cognitive decline, frailty and

dementia[2-4].

The most common causes of SNHL are aging and noise exposure[5]. Twin-studies have

calculated a heritability for SNHL of 0.35-0.65[6,7], indicating that a large portion is accounted for

by genetic factors. Although early-onset hearing impairment has been linked to roughly 200

human hereditary hearing loss (HHL) genes,[8] in contrast, findings from genome-wide

association studies (GWAS) on adult onset SNHL demonstrate a highly polygenic genetic

architecture[9,10], characterized by common genetic variants individually conferring small risks.

Whereas the technology to cure hearing impairment caused by monogenic disorders utilizing

gene therapy is rapidly advancing with some syndromes currently undergoing clinical trials[11],

ultimate treatment of the more common polygenic forms of SNHL will require identification and

characterization of multiple genes, variants, and pathways using technology yet to be

determined. Given that statistical power to identify risk loci for polygenic disorders mandates

large genetic studies, much of the otologic research has focused on the use of one or two self-

reported items for disease identification[10,12]. While the audiogram remains the gold standard for

an objective measure of hearing loss, self-reported hearing loss questions have provided a

satisfactory measure to elicit relevant loci from large genome-wide association studies

(GWAS)[13-15]. The largest self-reported SNHL GWAS meta-analysis to date, including the UK

Biobank (UKB) and 8 smaller cohorts of clinically diagnosed and self-reported hearing

impairment on 723,266 individuals from the EARGEN consortium, identified 48 loci[16].

Here we report on a phenotype based on ICDs (international classification of disease) that

include the most common forms of SNHL, i.e., age-related and noise-induced, typically

ascertained by audiograms, from the Million Veteran Program (MVP), comparing the limitations

of measuring a complex and clinically heterogeneous trait based on self-report to a clinical

diagnosis of SNHL derived from more precise auditory measurements.

MVP consists of an aging population currently with over one million participants, who, during

their military career, have been exposed to multiple sources of loud noise[17]. To address the

question of 'minimal phenotyping', we contrast the GWAS based on clinical diagnoses from

MVP with a GWAS on self-reported hearing loss in the United Kingdom Biobank (UKB), which

has been included in most of the recent GWAS meta-analyses[10,12,16,18-22]. Our findings indicate

that despite the differences of phenotype, demographics, and noise exposure of the two

cohorts, they share a large proportion of their genetic architecture. Combination in meta-

analysis elicits a total of 108 loci in individual ancestry GWAS and meta-analysis with 54 novel

genes from prioritized analysis and 100 significant novel genes from gene-based examination.

These loci contain inferences of 17 congenital hearing loss genes as well as multiple other ear-

related genes, and we present a plausible explanation for a mechanism of injury.

Methods

SNHL Phenotype in MVP

MVP provided data from 963,753 participants (658,038 with genotype data) in version 23_1

(released Apr 29, 2024), recruited 2011 through Sep 30, 2023, consisting of ICD records from

the US Veterans Administration Corporate Data Warehouse[23]. Information was linked to

individual, de-identified electronic health records (EHR).

Hearing loss was assessed using ICD9/ICD10 codes; participants with the following codes were

excluded from analysis: otosclerosis (H80.0, H80.1, H80.2, H80.8, H80.9, 387.0, 398.1, 387.2,

387.8, and 387.9), Meniere's disease (H81.0 and 386.0), pulsatile and objective tinnitus (H93.A,

388.32), conductive hearing loss (H90.0, H90.1, H90.2, H90.A1, 389.0), mixed hearing loss

(H90.6, H90.7, H90.8, H90.A, H90.A3, 389.2), unilateral sensorineural hearing loss (H90.4,

H90.A2, 389.13, 389.15, 389.17), ototoxicity (H91.0), sudden hearing loss (H91.2, 388.2)

congenital malformations and anomalies (Q16, Q17.2, Q17.4, Q17.8, Q17.9, 744.0, 733.23,

744.29), intraoperative and postprocedural complications and disorders of ear and mastoid

process, including mastoiditis and related conditions (H95, 383.3, 383.8, 383.89), certain

degenerative and vascular disorders of the ear (H93.0, 388.0, 388.00, 388.02), disorders of the

acoustic and cranial nerves (D33.3, H93.3, 225.1, 388.5), acoustic neuritis in infectious and

parasitic diseases (H94.0), in addition to Deaf non-speaking and other specified forms of

hearing loss (H91.3, H91.8, H94.8, 389.7, 389.8).

From the remaining participants, SNHL cases (N = 236,597) had to have a diagnosis of

sensorineural hearing loss (H90.3, H90.5, H91.1, H91.10, H91.11, H91.12, H91.13, 388.01,

389.1, 389.10, 389.11, 389.12, 389.14, 389.16, and 389.18), while controls (N = 296,284)

consisted of those who had no evidence of the above ICDs nor ICDs for unspecified hearing

loss (H91.9, H91.90. H91.91, H91.92, H91.93), or for the fitting, adjustment, presence, or

management of a hearing aid or device (Z45.3, Z46.1, Z96.2, Z97.4, 389.9, and V53.2). All

participants provided written informed consent, and the study was approved by the University of

California San Diego and VA Central Institutional Review Boards.

Genotyping, quality control, and imputation

Analyses were based on release 4 of the MVP imputed genotype data containing 658,038

participants. Details of the genotyping, quality control, and imputation procedures used have

been reported in detail[23]. In brief, MVP samples were genotyped on a customized version of the

Affymetrix Axiom biobank array and standard genotype quality control procedures were

followed. Genotype data was phased using Eagle version 2.4[24] and imputed using minimac

version 4[25] with the Haplotype Reference Consortium[26] reference panel.

Assessment of ancestry

Ancestry was determined using SNPweights[27] with a global reference panel[28], including 93,172

SNPs that overlap between the reference data and MVP genotype data. Reference SNPs were

selected to have MAF > 1% in reference populations and are in low LD (based on independent

pairwise LD pruning, using a 1000 kb window, 50 SNP step size, and r-squared of 0.2, as

calculated in PLINK[29]. Participants were placed into the three following large groupings: 1.

European ancestry (EUA; individuals with ≥ 90% European ancestry); 2. African ancestry (AFA;

individuals with ≥ 5% African ancestry, < 90% European ancestry, < 5% East Asian, Native

American, Oceanian and Central-South Asian ancestry, or individuals with ≥ 50% African

ancestry, < 5% Native American, Oceanian and < 1% Asian ancestry); and 3. Indigenous

American Ancestry (IAA; individuals with ≥ 5% Native American ancestry, < 90% European, and

< 5% African, East Asian, Oceanian and Central-South Asian ancestry).

Calculation of relatedness and principal components (PCs)

Relatedness within samples was estimated using KING[30]. For each pair of related participants

(kinship coefficient > 0.0884), one was removed, giving preference to retain cases. If case

status was identical between the pair, one of the pair was removed at random (N = 21,923

excluded). PCs were calculated within unrelated subjects of the same ancestry using

FlashPCA2[31]. SNPs were excluded for MAF < 1%. Remaining SNPs were pruned for LD over 1

MB windows stepped over 50 variants at a time with an r^2 threshold of 0.05.

GWAS in MVP based on EHR data

GWAS for SNHL was performed for each of the 3 ancestry groups in MVP separately with

PLINK 2[29], using logistic regression including sex, age, age^2, and 10 PCs as covariates. Only

SNPs with MAF > 1% and imputation info score > 0.6 (calculated within analytic sample) were

analyzed. Genome-wide significance (GWS) was assessed at a p value < 5×10^{-8}.

GWAS in UK Biobank (UKB) based on self-reported hearing difficulty

GWAS summary results from Wells et al.[12] for self-reported hearing difficulty in the UKB were

downloaded from the UKB Returned Datasets Catalog. The cohort consisted of white British

ancestry (EUA only) recruited from 2007 - 2013 who volunteered to participate and completed a

questionnaire including field code 2247 and 2257 regarding hearing difficulties and difficulties

hearing in ambient noise. GWAS cases were those who answered "yes" to both questions ("Yes,

diagnosed by doctor or health professional" or "Yes, not diagnosed by health professional"; N =

87,056 cases), controls were those who answered "no" to both (N = 163,333 controls).

Association analysis was performed using a linear mixed-effects model implemented in BOLT-

LMM v.228 including age, sex, genotyping platform, and 10 PCs (see [12] for additional details).

Meta-analysis

For compatibility with MVP, UKB GWAS summary statistics were converted to the log odds ratio

scale[32] and inverse variance weighted fixed effects meta-analysis was performed in METAL[33].

Only SNPs present in two or more input datasets were included in meta-analyses. Three meta-

analyses were performed in total: 1. MVP multi-ancestry (EUA, AFA, IAA) meta-analysis; 2.

European ancestry meta- analysis in MVP and UKB; 3. meta-analysis of the full sample,

including MVP multi-ancestry + UKB EUA GWAS.

Functional mapping and annotation

Functional annotation of the GWAS results were performed with FUMA v1.6.1[34] using default

settings and the human genome assembly GRCh37 (hg19). The SNP2Gene module was used

to define independent genomic risk loci and variants in LD with lead SNPs ($r^2 > 0.6$, calculated

using ancestry appropriate 1KGPp3 reference genotypes (EUR, AMR, AFR). Genomic risk loci

were combined to have non-overlapping base pair positions and SNPs were annotated to the

nearest gene. Functional consequences of SNPs were obtained by mapping SNPs on their

chromosomal position and reference alleles to databases containing known functional

annotations, including ANNOVAR[35], Combined Annotation Dependent Depletion[36] and

RegulomeDB[37].

Gene prioritization

Gene prioritization was performed using FLAMES version 1.1.2[38]. To generate credible sets,

Bayesian fine-mapping of risk loci was performed using the coloc R package[39]. Convergence-

based gene prioritization scores were calculated using Polygenic Priority Scores v0.2[40], using

as inputs MAGMA gene-level z-scores and pathway naïve scores. FLAMES annotations were

generated for each credible set, using as inputs the functional annotation data, Polygenic

Priority Score output data, and FUMA generated MAGMA gene and gene-tissue analysis. Note,

while FUMA positionally maps SNPs to genes based on their proximity to gene boundaries

(within 10kb), FLAMES annotates risk loci to a potentially larger set of genes because, in

addition to positional mapping, it incorporates several other SNP-to-gene annotation databases.

FLAMES scoring was performed using the annotation data. Only genes above the previously

calibrated cumulative 75% precision threshold are reported.

Hearing genes from HereditaryHearingLoss.org

We extracted 196 protein-coding genes associated with non-syndromic and syndromic hearing

loss from the Hereditary Hearing Loss database (https://hereditaryhearingloss.org, accessed

8/9/2024)[8]. This excluded Mitochondrial Non-syndromic Hearing Loss genes and *MIR96,* which

encodes microRNA. Protein-coding genes were aligned with genes from GWS genomic risk loci

and gene-based analysis (MAGMA). Genes with multiple symbol aliases were consolidated

using the NIH gene database (https://www.ncbi.nlm.nih.gov/gene/ ; accessed 8/28/2024).

Identification of novel genes

In order to identify novel, previously not reported genes, the following criteria were used.

For GWS loci from the GWAS: 1. The locus did not have any genes that were present in the

HereditaryHearingLoss.org database (accessed 8/9/2024), 2. the locus did not have any genes

that were present in the NHGRI-EBI GWAS Catalog database

(https://www.ebi.ac.uk/gwas/home, accessed 7/2/2025; selected traits: sensory perception of

sound, hearing loss, sensorineural hearing loss, hearing threshold trait, sensorineural hearing

impairment, hearing physiology trait, noise-induced hearing loss, age-related hearing

impairment, hearing process quality, able to hear with hearing aids, deafness, but excluding

chemoradiation-induced hearing loss in nasopharyngeal carcinoma and speech-in-noise

perception traits, as well as SNP-by-SNP and SNP-by-environment interaction studies and

intergenic loci). 3. the locus does not overlap with a GWS locus from Wells 2019[12] hearing

difficulty (since this GWAS was used for the meta-analysis), 4. the locus did not overlap with

something that was disqualified due to #1, 2, or 3, and 5. the locus had at least one protein-

coding gene in the locus and/or a FLAMES prioritized gene.

For genes from the gene-based analysis: points 1-3 were applied.

Regional association plots

Regional visualizations of GWS loci were produced using LocusZoom 1.4[41]. Linkage

Disequilibrium (LD) was calculated using 1KGPp3 data, using ancestry-specific reference data

(EUR, AMR, AFR).

Gene-based and gene set analyses with MAGMA

The Multi-marker Analysis of GenoMic Annotation (MAGMA) v1.08[42] tool implemented in FUMA was used to perform gene-based, gene-pathway, and human-based tissue enrichment analyses. For gene-based analysis, SNPs were mapped to 18,809 protein coding genes. For each gene, its association with SNHL was determined as the weighted mean squared test statistic of SNPs mapped to the gene, where LD patterns were calculated using ancestry appropriate 1KGPp3 reference genotypes. The significance of genes was set at a Bonferroni-corrected threshold of $p < 2.658e-06$ (0.05/18,809). To identify specific biological pathways implicated, gene-based test statistics were used to perform a competitive set-based analysis of 17,012 pre-defined curated gene sets and GO terms obtained from MsigDB[43]. The significance of pathways was set at a Bonferroni-corrected threshold of $p < 2.94e-06$ (0.05/17,012).

Cochlear cell type enrichment analyses in human

To test if tissue-specific gene expression was associated with SNHL, enrichment analyses were performed via MAGMA gene-tissue analysis of the EUA gene-based GWAS results. Since human inner ear cell type data has not yet been integrated in the FUMA tissue collection, we paired the collection with human adult inner ear tissue [44]. Data was converted from h5ad to h5seurat using SeuratDisk (0.0.0.9021). The R package Seurat v5.3.0 function AggregateExpression() was used to log normalize and return pseudobulk gene expression across each cell type. MAGMA v1.10 was used to annotate the SNPs based on GRCh37 and perform gene-set analysis with GTEx v8 specific and general gene expression datasets downloaded from FUMA.[34] Gene-set analyses were conditioned on the mean of the normalized gene expressions (reads per kilo base per million, RPKM) of the GTEx v8 tissues.

Cochlear cell type enrichment analyses in mice

Cochlear cell type enrichment analyses were performed via MAGMA gene-tissue analysis of the

EUA gene-based GWAS results paired with mice expression data. Mouse gene expression data

was derived from cochlear cells and nuclei from Jean *et al.* (20-day old mice only)[45] and Hoa et

al[46]. Additional pre-processing steps were applied to data from Jean *et al.*; cells were retained if

the RNA count was between 1,000 and 30,000, gene/non-coding RNA between 500 and 5,000,

and mitochondrial genes less than 2% using the R package Seurat v4.9.9.9058. Pseudobulk

gene expression was calculated by log2 normalizing the mean raw expression across each cell

type. Data from Hoa et al. for the 5 tissues from the organ of Corti had been pre-processed into

reads per kilobase million by study authors. Mouse gene Ensembl IDs were converted to human

Entrez IDs using a homology map from the Mouse Genome Database[47]. Only genes with one-

to-one mappings were kept. MAGMA v1.10 was used to annotate the SNPs based on GRCh37

and perform gene-set analysis. Gene-set analyses were conditioned on the log2 mean

expression across the cochlear cell types for each gene.

SNP-based heritability and genetic correlation

SNP-based heritability (h^2_{SNP}) was evaluated in EUA data using LD score regression (LDSC)[48].

Input LD scores were computed from 1KGPp3 EUA samples. The LDSC intercept was used to

test for artifactual inflation of test statistics, and the attenuation factor was used to estimate the

proportion of inflation coming from polygenic signal. The h^2_{SNP} was converted to the liability

scale[49]. Cross-trait LDSC[48] was used to estimate the genetic correlation (r_g) between datasets.

Univariate and bi-variate Gaussian mixer model (MiXeR) analysis

We used univariate MiXeR v1.3[50] to estimate the genetic architecture of MVP EUA and UKB

SNHL phenotypes. MiXeR estimates SNP-based heritability and two subcomponents whose

product is proportional to heritability: the proportion of non-null SNPs (polygenicity) and variance

of effect sizes of non-null SNPs (discoverability). MiXeR was applied to GWAS summary

statistics under the default settings with the supplied EUA LD reference panel. The results for

the number of influential variants reflect the number of SNPs necessary to explain 90% of SNP

based heritability. Bivariate MiXeR[51] was used to estimate phenotype specific polygenicity and

the shared polygenicity between phenotypes. Goodness-of-fit of the MiXeR model relative to

simpler models of polygenic overlap was assessed using AIC values. Heritability, polygenicity

and discoverability estimates were contrasted between datasets using the *z*-test.

Results

Ancestry-specific GWAS and multi-ancestry meta-analysis for SNHL in MVP

GWAS in the MVP was performed based on ICD diagnosis of SNHL from the electronic health

record. After excluding participants for mixed, conductive, and uncommon forms of SNHL (e.g.

Meniere's disease and ototoxicity), 210,240 cases and 265,275 controls were available for

GWAS. The MVP cohort is predominantly (91%) male with an average age of 62.2 (13.9 SD)

years at enrollment (Supplementary Table 1). GWAS were stratified into three ancestry groups.

GWAS of European ancestry participants (MVP EUA; 176,393 cases and 172,117 controls)

identified 43 genome-wide significant (GWS) loci (Supplementary Table 2, Fig. 1a). The inflation

of test statistics (Genomic control λ = 1.33) was almost entirely accounted for by polygenic

effects (LDSC attenuation ratio: 14.7%; LDSC intercept = 1.06; SE = 0.009; Supplementary

Table 7). Protein-coding genes within GWS loci were identified using functional annotation with

FUMA and further prioritized using FLAMES, identifying 29 genes. GWAS of African ancestry

participants (MVP AFA; 22,757 cases and 74,842 controls) identified one GWS locus

(rs59174224, p = 4.12E-08) within the gene *TUB* (Fig. S1b. *Tub* is a known murine gene leading

to profound hearing loss. A modifier *Moth1* protects hearing, and *Mtap1* then mediates *Moth1* at

the synaptic terminal[52]. GWAS of MVP indigenous American ancestry participants (MVP IAA;

11,090 cases, 18,316 controls) did not identify any GWS loci (Fig. S1c). Meta-analysis of the

EUA, AFA, and IAA GWAS (MVP meta-analysis; N = 475,515 subjects) identified 52 loci; 14 of

which were not significant in individual ancestry analyses (Supplementary Table 2, Fig. S1d).

Gene prioritization using FLAMES identified 36 genes in the 52 loci. Annotations for SNPs in LD

with significant SNPs in the MVP multi-ancestry meta-analysis are summarized in

Supplementary Table 3. Of the 52 loci, 12 contained a GWS SNP with a CADD score predicted

to be deleterious to function (>12.37) that was within the exon region of a gene. A total of 57

independent GWS loci were identified across the four MVP GWAS performed. Regional

association plots comparing ancestry-specific analyses for these loci are shown in

Supplementary Data 1.

Gene-based, gene-set, and gene-tissue analyses in MVP

To aggregate genetic markers at the level of genes and allow integration with gene expression

data, we performed gene-based analyses using MAGMA[42]. Gene-based analyses identified a

total of 76 GWS genes across the EUA, AFA, IAA, and meta-analyses (Supplementary Table 4,

Fig. S2 and S3a), several of which had not already been implicated by the positional mapping of

SNPs to genes. To ascertain specific biological pathways, a competitive set-based analysis of

17,012 pre-defined curated gene sets and gene ontology (GO) terms obtained from MsigDB

was performed in the MVP meta-analysis (Supplementary Table 5). Significant curated gene

sets   identified   were   in   the   gene   ontology   biological   processes   area,   i.e.,

GOBP_EAR_DEVELOPMENT   (p   =   1.52E-07),   GOBP_SENSORY_PERCEPTION_OF_

MECHANICAL_STIMULUS   (p   =   3.19E-07),   and   NIKOLSKY_BREAST_CANCER_

14Q22_AMPLICON (p = 2.50E-06). We note that the Nikolsky breast cancer amplicon pathway

significance level was primarily driven by one locus with two mapped genes with SNPs in partial

LD, *NID2* (p-value 2.61E-14) and *C14orf116* (*FOXN3,* p = 7.25E-10) (Supplementary Table 6).

Genetic comparisons of MVP and UKB SNHL GWAS

We performed a comparison of the genetic architecture of SNHL based on the GWAS using ICD

codes in MVP with a similarly sized GWAS (cases: N = 87,056; controls: N = 163,333) based on

self-report in the previously published UKB GWAS[12], which had identified 41 GWS loci (Fig. 1b).

Only data of EUA individuals were included.

The h^2_{SNP} estimated from the MVP was 0.0761 (SE = 0.0039, on the liability scale at 30%

population prevalence) (Supplementary Table 7A). In contrast, h^2_{SNP} estimated from the UKB

was higher at 0.1162 (SE = 0.0059, liability scale at 30% population prevalence). Differences in

genetic architecture were further explored with univariate MiXeR, noting that h^2_{SNP} is

proportional to the number of causal variants and the effects sizes of these variants. Univariate

MiXeR analyses identified greater polygenicity within UKB (5,240 causal variants, SE = 250)

than in MVP (3,790 causal variants, SE = 198) (z = 4.5, p = 5.4 x 10^{-6}). However, variant effect

sizes were not significantly different between studies (z-test comparing causal effect size

variance estimates: z = 1.3, p = 0.18) (Fig. 1c and Supplementary Table 8a).

Shared and unshared genetic variation between studies was explored with bivariate MiXeR

analyses. MiXeR estimated that MVP and UKB shared 3,624 (SE = 204) causal variants,

comprising 95.6% of the total MVP causal variants, but only 69.2% of the total UKB causal

variants (Fig. 1c and Supplementary Table 8b). Correspondingly, MiXeR estimated that 1,617

(SE = 235) variants were unique to the UKB, but only 167 (SE = 126) variants were unique to

MVP.

Genetic correlation among shared variants was r_g = 0.73 (SE = 0.03), which was higher than the

standard genome-wide estimate of genetic correlation r_g = 0.59 (SE = 0.03) estimated by LDSC

(Supplementary Table 7B). A comparison of only GWS loci showed a correlation of r = 0.77

between effect sizes of leading variants, further establishing that the identified GWS loci have

similar effects across studies. In conclusion, while self-reported hearing loss in the UKB is

influenced by a broader set of variation compared to the clinically diagnosed SNHL in MVP,

there is a substantial core set of variation with comparable effect sizes across these cohorts. Of

note however, our analyses are not able to distinguish between differences in populations or the

assessment tools used.

Meta-analysis of MVP and UKB EUA GWAS

An inverse variance weighted fixed effects meta-analysis in 598,899 EUA individuals (cases: N

= 263,449; controls: N = 335,450) of the EUA MVP and UKB GWAS identified 99 GWS loci

(Supplementary Table 9). Of these, 43 were not previously identified in either the MVP or UKB

GWAS. These data will be used in downstream analyses that are based on linkage

disequilibrium (LD) patterns, as many of the non-EUA individuals in the MVP are admixed.

Multi-ancestry meta-analysis of MVP and UKB

A meta-analysis of the MVP multi-ancestry GWAS and the UKB GWAS identified a total of 108

GWS loci in 725,904 individuals; 54 of these loci have a novel protein-coding gene and/or

FLAMES prioritized gene (see definition of 'novel' in methods), while two loci (locus 15 and 49)

did not include a gene (Table 1, Supplementary Table 10 and Fig. 2b). Regional association

plots comparing ancestry-specific analyses for these loci are shown in Supplementary Data 1.

Functional mapping and annotation of risk loci using the FUMA pipeline are summarized in

Supplementary Table 11. Of the 108 loci, 26 (24%) contained an exonic GWS SNP with a

predicted deleterious CADD score (>12.37).

(Table 1 about here)

Gene-based analyses (MAGMA) identified 192 genome-wide significant genes, of which 100

are novel (Supplementary Table 12). Competitive set-based analysis of pre-defined curated

gene sets and gene ontology (GO) terms identified three significant gene sets,

GOBP_SENSORY_PERCEPTION_ OF_MECHANICAL_STIMULUS (p = 6.42E-09),

GOMF_ACTIN_BINDING (p = 2.55E-06) and GOMF_CYTOSKELETAL_PROTEIN_BINDING (p

= 1.63E-06) (Supplementary Tables 13 and 14).

To identify tissues and pathways underlying these gene-based associations, we performed a

MAGMA tissue expression analysis with 31 general human tissue types, including human inner

ear tissue from van der Valk *et al.*[44], and found significant enrichment in the inner ear ($p =$

2.66×10^{-8}), nerve ($p = 1.50 \times 10^{-5}$) and pituitary ($p = 8.66 \times 10^{-4}$) of EUA MVP, UKB, and in meta-

analysis. Analysis of 71 different tissue types identified significant enrichment in the cerebellum

($p = 2.33 \times 10^{-5}$), cerebellar hemisphere ($p = 6.22 \times 10^{-5}$), and tibial nerve ($p = 3.65 \times 10^{-4}$), as well

as 13 inner ear cell types (Supplementary Table 15 and Fig. S4).

Please note that these gene-tissue analyses were based on GTEx v8 data sets, which do not

include the human cochlea, the tissue most relevant to hearing loss. We enriched with human

data of one subject only[44] which had been labeled "utricle", however, this analysis identified

cochlear cell types as well as vestibular tissue. Because of the paucity of human cochlear data,

we performed additional enrichment analyses based on available mouse data from two recent,

large studies on the organ of Corti[46] and cochlear cells[45] (Supplementary Table 16 and Fig. S5).

The murine organ of Corti included outer and inner hair cells, pillar cells, Deiter's cells, and

melanocytes. The genetic signal associated with SNHL from the meta-analysis was significantly

enriched ($p < 0.01$) for expression in all cell types (except for Deiter's cells), conditional on the

average expression of cochlear tissues. The cochlear data included 34 cell types. No cell type

was significantly enriched when conditioned on the average expression of all cell types as the

baseline.

Examination of implicated genes in the context of congenital hearing loss

The Hereditary Hearing Loss database was used to compare findings from our analyses to

genes identified in the context of congenital hearing loss. Of the 196 protein-coding genes

(accessed August 9, 2024), 17 genes were implicated by our GWAS from either positional

mapping of GWS loci or gene-based analyses (Supplementary Table 17). Twelve out of the 17

are members of significant gene ontology (GO terms) from our gene-set analysis.

Finally, we performed a partitioned heritability analysis for the SNHL EUA MVP and UKB meta-

analysis (N = 598,899 European subjects from MVP and UKB), partitioning SNPs into 3 groups:

SNPs in protein coding genes (in HHLdb), SNPs in protein coding genes (not in HHLdb), and

SNPs outside of protein coding genes (Table 2). SNP-based heritability estimates including 196

known HHL genes showed a 3.26-fold enrichment relative to all other protein-coding genes.

However, the large majority of risk (99% of SNPs, explaining 97% of h^2_{SNP}) for polygenic SNHL

is captured by SNPs outside of congenital HHL genes.

(Table 2 about here)

Discussion

Million Veteran Program multi-ancestry analysis

This study introduces MVP hearing data based solely on a clinical diagnosis of sensorineural

hearing loss. MVP GWAS alone elicits a total of 57 distinct loci with 76 genes in gene-based

analysis and significant relevant otologic pathways implicated by gene-set analysis. We

demonstrate that this phenotype approach is broadly successful for variant identification. A

benefit of our method is the exclusion of many other diseases that lead to hearing loss, e.g.

acoustic neuroma, otosclerosis, sudden hearing loss, etc., which may have distinct genetic

architectures that can lead to inflation in the results.

In addition to a phenotypic difference between MVP and UKB, i.e., ICD versus self-report, the

two populations have contrasting demographics, environmental exposures, and slightly different

mean ages. Unlike the civilian population of the UKB, MVP is a group of ambulatory US

Veterans consisting of greater than 90% males who have been exposed to noise levels during

their military career[53] of up to 126 dB[17], as well as a 20% prevalence of head injury, another

cause for hearing loss[54]. Although we cannot distinguish the effects of noise and head trauma

from age in this study, the MVP cohort might have elicited the phenotypic expression of variants

related to these differences.

Nevertheless, despite these dissimilarities, there appears to be a commonality of hearing loss

genetic architecture between the MVP and UKB datasets where the studies estimated 3,624

(67.0%) shared causal variants. Importantly, this similarity indicates that the data are suitable for

aggregation, and meta-analysis of the two identified 108 loci[12,16], 54 with a novel prioritized

and/or protein-coding gene, and a total of 100 significant novel genes in gene-analysis.

Novel prioritized genes

Among our newly reported genes, at least five of the "top ten hits" within novel prioritized genes

are worth noting as cochlear-related (Supplementary Table 10). *DPT* has been identified in

familial Meniere's disease, a syndrome consisting of episodic hearing loss, vertigo and tinnitus,

and is an extracellular matrix protein that functions in cell matrix interactions and assembly[55].

SETBP1 mutations are a cause of Schinzel-Giedion syndrome, characterized by distinct facial

features, malformations, and SNHL[56]. *WNT7A* regulates axon outgrowth in the cochlea[57] during

inner ear development and is responsible for planarity of hair cell bundles[58]. *EVL* localizes at the

cuticular plate supporting inner ear hair cells[59]. Expression of the transcription factor *BACH2*,

under control of *miR-96* in the murine model, is altered in noise trauma, and plays a mediation

role in oxidative stress[60].

Monogenic hearing loss genes and a plausible explanation of significant GWAS findings

Partitioned SNP-based heritability based on our results of known monogenic hearing loss genes

showed a 3.26-fold enrichment relative to other protein-coding genes. This is in agreement with

other studies tying hereditary hearing loss (HHL) genes to the polygenic disorder of adult SNHL,

where polygenic risk scores from a self-reported hearing loss GWAS predicted higher odds of

children's hearing loss[61]. Separately, multiple studies of self-reported hearing difficulties have

also identified an increased burden of rare variants within HHL genes[16,20,62-65]. Thus, while

childhood hearing loss is known to be at least 50% monogenic in origin[66], it appears that adult-

onset SNHL has genetic signals in common with HHL.

To organize our results, it is helpful to examine the function of hereditary hearing loss genes and

additional genes identified as significant in this study that are directly related to inner ear

physiology (See Fig. 4), particularly within the context of the stereocilia of the inner and outer

hair cells. Parallel actin fibers in stereocilia are critical for stiffness in response to the fluid wave.

Within these structures, *ESPN* encodes an actin-binding protein required for assembly and

stabilization[67], *TMPRSS3* and *CLRN2* express stereociliary proteins critical for hair cell

survival[68], and *WNT7A* directs the uniform 3-dimensional polarity of outer hair cells.

In response to the fluid/sound wave, stereociliae of hair cells bend at their rootlets (Fig. 4b, 4c).

TRIOBP, CLIC5, and *ATP2B2*, all non-syndromal congenital hearing loss genes, are expressed

in this subcellular area[69-72]. *TRIOBP* wraps around actin providing stiffness and support, and

CLIC5 is a chloride channel protein that stabilizes the attachment of the stereocilia plasma

membrane to its base[73-75]. *ATP2B2* expresses a transmembrane ATPase at the lower half of the

stereocilia and cuticular plate, where it regulates Ca^{++} concentration within the cytosol. Although

not on the HHL list, *SPTBN1* is expressed as part of the support structure at the area of the

rootlet as well. *SPTBN1* is associated with an autosomal dominant neurodevelopmental

disorder which includes SNHL[76]. In support of the "sensory mechanical stimulus pathway" (*p*-

value 6.42E-09), our human-based transcription analysis of the organ of Corti indicates hair cell

enrichment (Supplementary Table 15).

Bending at the rootlet allows for increased tension on tip-links, which in turn controls the

opening of the mechanotransduction ion channel (MET). *CDH23*, prioritized by FLAME[38],

constitutes the upper half of the tip-link, and *MYO7A* tethers this link to an actin bundle within

the stereocilium. These genes are responsible for different forms of Usher's syndrome, the most

common genetic cause of congenital deafness. In this critical area of the sensory apparatus,

both *CLRN2* and *LOXHD1* are engaged in MET maintenance at the tip of the stereocilia[77].

OHCs are tethered at their apex and base. At their apex, the tectorial membrane (TM) (Fig.

4b)[77] is a gelatinous structure providing a static force to the shearing motion of the sound wave.

COL11A1 and *COL9A3* code for two α-chain collagens providing upper support within the

tectorial membrane. They individually cause different variants of Stickler's Syndrome[78]. Other

significant support genes include *GAS2*, identified in pillar and Deiters' cells as a microtubule

regulatory gene, maintaining arrangement of the actin skeleton[79].

In a separate micro-anatomical area, our human cochlear cell transcription analysis noted

enrichment within melanocytes, located in the stria vascularis, and functioning to maintain the

endocochlear potential. Although not on the HHL list, *ATP6V0A4,* identified within "sensory

perception of mechanical stimulus", is a known hearing loss gene[80] expressed in the basal stria

cells, where it acts as a proton pump, maintaining the pH of endolymph bathing the apical

region of the hair cell. *SOX10* (Waardenberg's Syndrome), also found within the stria vascularis,

expresses an aquaporin repressor[81].

Associated with the nerve signal through the spiral ganglion, other significant congenital hearing

loss genes were identified related to neurons, the ribbon synapse, and the myelin sheath (Fig.

4b). Studies of both age-related and noise-induced SNHL have found that after injury, up to 50%

of ribbon synapses can be lost prior to an objective audiogram threshold change[82,83] and this

finding has been correlated to loss of speech intelligibility[84,85]. These synapses connect the IHC

cells to Type I spiral ganglion nerve cells[86]. Located within this synaptic area are *LMX1A*, a

neuro-restrictive silencer[87] expressed in Type I spiral ganglion nerves. *QKI* is identified in a

murine model as essential for proper myelination of spiral ganglion neurons and auditory nerve

fibers[85]. In aggregate, these genes identified as significant in prioritization and gene-based

analysis provide putative points of injury to the hearing sensory mechanism to be explored in

future studies.

Limitations

This study has limitations. Although an ICD diagnosis is a step towards an objective phenotype,

it does not describe the state of individual audiogram frequencies, and heritability studies have

indicated that a good portion of the heritability of SNHL is frequency-specific[88]. Further, while the

genes identified provide biologically plausible explanations for SNHL, until now, expression data

in the human cochlea has not been available. While this study used expression data from one

human inner ear sample, further study for causative variants requires whole genome

sequencing and multiple human cochlear methylation/expression studies.

Conclusions

In GWAS of SNHL combining MVP and UKB, we have identified 108 loci containing 54 novel

protein-coding genes and/or prioritized genes, 100 novel genes in gene-based analysis, and

significant pathways related to the structure of the cochlea. Despite different phenotypes,

demographics, and exposures we find substantial overlapping genetic signals. Although over

95% of the risk for this polygenic disorder is captured by SNPs outside of congenital HHL

genes, SNP-based heritability estimates 3.26-fold enrichment of these genes relative to all other

protein-coding genes. These findings will provide new targets of investigation of druggable

genes and pathways for treatment of this pervasive and currently permanent condition. Our next

tasks are to analyze genetic signals from individual audiogram frequencies within larger studies

and search for phenotypes with increased heritability of this pervasive disorder.

Abbreviations

AFA – African Americans

AIC – Akaike information criterion

ANNOVAR – annotation variation software

BM – basement membrane

CADD – combined annotation dependent depletion

EUA – European ancestry

FUMA – functional mapping and annotation

GO – gene ontology

GWAS – genome-wide association study

GWS – genome-wide significant

h_2SNP – heritability of single nucleotide polymorphisms

IAA – indigenous American ancestry

ICD – international classification of diseases

IHC – inner hair cells

LD – linkage disequilibrium

LDSC – linkage disequilibrium score regression

MAF – minor allele frequency

MAGMA – multi-marker analysis of genomic annotation

MET – mechanoelectrical transduction channels

MsigDB – molecular signatures database

MVP – Million Veteran Program

OC – Organ of Corti

OHC – outer hair cells

PC – principal component

SNHL – sensorineural hearing loss

SNP – single nucleotide polymorphism

TM – tectorial membrane

UKB – United Kingdom Biobank

YLD – years lived with disability

Supplementary Information

Additional file 1: Supplementary Table 1. MVP demographics. **Supplementary Table 2.**

Genome-wide significant loci in MVP across ancestries. **Supplementary Table 3.** Annotations

for SNPs in LD with significant SNPs in the MVP multi-ancestry meta-analysis. **Supplementary**

Table 4. Genome-wide significant genes from gene-based analyses across MVP ancestries.

Supplementary Table 5. MAGMA gene-set analysis, including 17,012 pre-defined curated gene

sets and GO terms obtained from MsigDB in the MVP multi-ancestry meta-analysis.

Supplementary Table 6. Genes within the 3 significant gene-sets from the MAGMA gene-set

analysis (MVP multi-ancestry meta-analysis). **Supplementary Table 7.** SNP-based heritability

and genetic correlation across of MVP and UKB EUA data. **Supplementary Table 8.** MiXeR

analysis of SNHL GWAS in European ancestry data. **Supplementary Table 9.** Genome-wide

significant loci in the European ancestry meta-analysis (MVP EUA and UKB). **Supplementary**

Table 10. Extended Table 1: Genome-wide significant loci in meta-analysis of sensorineural

hearing loss (N=725,904 MVP multi-ancestry and UKB individuals). **Supplementary Table 11.**

Annotations for all SNPs in LD with significant SNPs in the MVP multi-ancestry and UKB meta-

analysis. **Supplementary Table 12.** Gene-based analyses in the MVP multi-ancestry and UKB

cohorts. **Supplementary Table 13.** MAGMA gene-set analysis, including 17009 pre-defined

curated gene sets and GO terms obtained from MsigDB in the MVP multi-ancestry and UKB

EUA meta-analysis. **Supplementary Table 14.** Genes within the 3 significant gene-sets from

the MAGMA gene-set analysis (MVP multi-ancestry and UKB meta-analysis). **Supplementary**

Table 15. Tissue enrichment analysis for 31 and 71 tissue types across the European ancestry

MVP and UKB cohorts. **Supplementary Table 16**. Cell type enrichment analyses in cochlear

cells based on mouse gene expression data in MVP EUA and UKB. Supplementary Table 17.

Hereditary Hearing Loss genes implicated across analyses (HereditaryHearingLoss.org

database)

Additional file 2: Supplementary Figure 1. Ancestry-stratified GWAS and meta-analysis for

SNHL in the MVP. **Supplementary Figure 2.** Ancestry-stratified gene-based analyses of SNHL

in the MVP. **Supplementary Figure 3.** Gene-based analyses of the MVP multi-ancestry meta-

analysis and the UKB SNHL GWAS. **Supplementary Figure 4.** Gene-tissue expression

analyses of SNHL. Supplementary Figure 5. Cochlear cell type enrichment analyses based on

mouse data.

Additional file 3: MVP Core Acknowledgements.

Acknowledgements

We thank all the participants in the UK Biobank and the Million Veteran Program. This research

is based on data from the Million Veterans Program, Office of Research and Development,

Veterans Health Administration, and was supported by MVP000 as well as awards

#I01RX004293 and #I01BX005920. This publication does not represent the views of the

Department of Veteran Affairs or the United States Government. See MVP Core

Acknowledgements in Additional file 3.

Authors' contributions

RC, AXM, and CMN designed the study. AXM, EAM, and CMN conducted GWAS data analysis.

RC, CEM, JJ, EAM, AXM, and CMN performed post-GWAS analysis. RC, JJ, CEM, and EAM

visualized the data. All of the authors participated in the interpretation of data and revision of the

manuscript for intellectual content. RC and CMN obtained primary funding. RC supervised the

phenotype and CMN supervised the statistical analysis of GWAS results. All authors read and

approved of the final manuscript.

Funding

This research is funded by VA Merit grants #I01RX004293 and #I01BX005920.

Availability of data and materials

Summary statistics for MVP analyses will be deposited upon publication on dbGaP under

accession   number   phs001672   (https://www.ncbi.nlm.nih.gov/projects/gap/cgi-

bin/study.cgi?study_id=phs001672.v11.p1). MVP summary data access can be obtained by

submitting a data access request through dbGaP; raw data are protected and are not available

due to privacy reasons.

The codes for the analysis are available on Github (https://github.com/nievergeltlab/)

Declarations

Ethics approval and consent to participate

The use of MVP individual-level data was conducted under project #MVP039 and #MVP060.

MVP is approved by the VA Central IRB, and the projects were also approved by local VA IRB in

San Diego. This research was performed using the UKB Resource under application 40951.

UKB has approval from the North West Multi-center Research Ethics Committee (MREC) as a

Research Tissue Bank (RTB) approval. This approval means that separate ethical approval is

waived and that researchers can operate under the RTB approval.

Consent for publication

This manuscript was approved for submission and publication by the MVP Publications

Committee.

Competing interests

No authors declare any competing interests.

Author details

[1]Research Department, Veterans Affairs San Diego Healthcare System, 3350 La Jolla Village

Drive, San Diego, CA, USA. [2]Department of Otolaryngology – Head and Neck Surgery, 9444

Medical Center Drive, La Jolla, CA, USA. [3]Department of Psychiatry, University of California

San Diego, 9500 Gilman Drive, La Jolla, CA, USA

References

1.   Nocini, R., Henry, B.M., Lippi, G. & Mattiuzzi, C. Estimating the worldwide burden of

health loss due to hearing loss. *Eur J Public Health* **33**, 146-148 (2023). PMID

36377968

2.   Dawes, P. *et al.* Hearing loss and cognition: the role of hearing AIDS, social isolation and

depression. *PLoS One* **10**, e0119616 (2015). PMID 25760329

3.   Yévenes-Briones, H. *et al.* Association Between Hearing Loss and Impaired Physical

Function, Frailty, and Disability in Older Adults: A Cross-sectional Study. *JAMA*

Otolaryngol Head Neck Surg **147**, 951-958 (2021). PMID 34554203

4.   Jafari, Z., Kolb, B.E. & Mohajerani, M.H. Age-related hearing loss and tinnitus, dementia

risk, and auditory amplification outcomes. *Ageing Res Rev* **56**, 100963 (2019). PMID

31557539

5.   Dobie, R.A. The burdens of age-related and occupational noise-induced hearing loss in

the United States. *Ear Hear* **29**, 565-77 (2008). PMID 18469718

6.   Duan, H. *et al.* Heritability of Age-Related Hearing Loss in Middle-Aged and Elderly

Chinese: A Population-Based Twin Study. *Ear Hear* **40**, 253-259 (2019). PMID

29794565

7.   Bogo, R. *et al.* The role of genetic factors for hearing deterioration across 20 years: a

twin study. *J Gerontol A Biol Sci Med Sci* **70**, 647-53 (2015). PMID 25665831

8.   Walls WD, A.H., Smith RJH. Hereditary Hearing Loss Homepage.

9.   Fransen, E. *et al.* Genome-wide association analysis demonstrates the highly polygenic

character of age-related hearing impairment. *Eur J Hum Genet* **23**, 110-5 (2015). PMID

24939585

10.   De Angelis, F. *et al.* Sex differences in the polygenic architecture of hearing problems in

679   adults. *Genome Med* **15**, 36 (2023). PMID 37165447

680   11.   Jiang, L., Wang, D., He, Y. & Shu, Y. Advances in gene therapy hold promise for treating

681   hereditary hearing loss. *Mol Ther* **31**, 934-950 (2023). PMID 36755494

682   12.   Wells, H.R.R. *et al.* GWAS Identifies 44 Independent Associated Genomic Loci for Self-

683   Reported Adult Hearing Difficulty in UK Biobank. *Am J Hum Genet* **105**, 788-802 (2019).

684   PMID 31564434

685   13.   McCullagh, M.C., Raymond, D., Kerr, M.J. & Lusk, S.L. Prevalence of hearing loss and

686   accuracy of self-report among factory workers. *Noise Health* **13**, 340-7 (2011). PMID

687   21959114

688   14.   Brennan-Jones, C.G. *et al.* Self-reported hearing loss and manual audiometry: A rural

689   versus urban comparison. *Aust J Rural Health* **24**, 130-5 (2016). PMID 26311193

690   15.   Cherny, S.S. *et al.* Self-reported hearing loss questions provide a good measure for

691   genetic studies: a polygenic risk score analysis from UK Biobank. *Eur J Hum Genet* **28**,

692   1056-1065 (2020). PMID 32203203

693   16.   Trpchevska, N. *et al.* Genome-wide association meta-analysis identifies 48 risk variants

694   and highlights the role of the stria vascularis in hearing loss. *Am J Hum Genet* **109**,

695   1077-1091 (2022). PMID 35580588

696   17.   Jokel, C., Yankaskas, K. & Robinette, M.B. Noise of military weapons, ground vehicles,

697   planes and ships. *J Acoust Soc Am* **146**, 3832 (2019). PMID 31795677

698   18.   Praveen, K. *et al.* Population-scale analysis of common and rare genetic variation

699   associated with hearing loss in adults. *Commun Biol* **5**, 540 (2022). PMID 35661827

700   19.   Naderi, E. *et al.* The genetic contribution of the X chromosome in age-related hearing

701   loss. *Front Genet* **14**, 1106328 (2023). PMID 36896235

20.   Ivarsdottir, E.V. *et al.* The genetic architecture of age-related hearing impairment

revealed by genome-wide association analysis. *Commun Biol* **4**, 706 (2021). PMID

34108613

21.   Liu, W., Johansson, A., Rask-Andersen, H. & Rask-Andersen, M. A combined genome-

wide association and molecular study of age-related hearing loss in H. sapiens. *BMC*

Med **19**, 302 (2021). PMID 34847940

22.   Kalra, G. *et al.* Biological insights from multi-omic analysis of 31 genomic risk loci for

adult hearing difficulty. *PLoS Genet* **16**, e1009025 (2020). PMID 32986727

23.   Gaziano, J.M. *et al.* Million Veteran Program: A mega-biobank to study genetic

influences on health and disease. *J Clin Epidemiol* **70**, 214-23 (2016). PMID 26441289

24.   Loh, P.R., Palamara, P.F. & Price, A.L. Fast and accurate long-range phasing in a UK

Biobank cohort. *Nat Genet* **48**, 811-6 (2016). PMID 27270109

25.   Das, S. *et al.* Next-generation genotype imputation service and methods. *Nat Genet* **48**,

1284-1287 (2016). PMID 27571263

26.   McCarthy, S. *et al.* A reference panel of 64,976 haplotypes for genotype imputation. *Nat*

Genet **48**, 1279-83 (2016). PMID 27548312

27.   Chen, C.Y. *et al.* Improved ancestry inference using weights from external reference

panels. *Bioinformatics* **29**, 1399-406 (2013). PMID 23539302

28.   Nievergelt, C.M. *et al.* International meta-analysis of PTSD genome-wide association

studies identifies sex- and ancestry-specific genetic risk loci. *Nat Commun* **10**, 4558

(2019). PMID 31594949

29.   Chang, C.C. *et al.* Second-generation PLINK: rising to the challenge of larger and richer

datasets. *Gigascience* **4**, 7 (2015). PMID 25722852

30.   Manichaikul, A. *et al.* Robust relationship inference in genome-wide association studies.

Bioinformatics **26**, 2867-73 (2010). PMID 20926424

31.　Abraham, G., Qiu, Y. & Inouye, M. FlashPCA2: principal component analysis of Biobank-

scale genotype datasets. *Bioinformatics* **33**, 2776-2778 (2017). PMID 28475694

32.　Cook, J.P., Mahajan, A. & Morris, A.P. Guidance for the utility of linear models in meta-

analysis of genetic association studies of binary phenotypes. *Eur J Hum Genet* **25**, 240-

245 (2017). PMID 27848946

33.　Willer, C.J., Li, Y. & Abecasis, G.R. METAL: fast and efficient meta-analysis of

genomewide association scans. *Bioinformatics* **26**, 2190-1 (2010). PMID 20616382

34.　Watanabe, K., Taskesen, E., van Bochoven, A. & Posthuma, D. Functional mapping and

annotation of genetic associations with FUMA. *Nat Commun* **8**, 1826 (2017). PMID

29184056

35.　Wang, K., Li, M. & Hakonarson, H. ANNOVAR: functional annotation of genetic variants

from high-throughput sequencing data. *Nucleic Acids Res* **38**, e164 (2010). PMID

20601685

36.　Rentzsch, P., Witten, D., Cooper, G.M., Shendure, J. & Kircher, M. CADD: predicting the

deleteriousness of variants throughout the human genome. *Nucleic Acids Res* **47**, D886-

D894 (2019). PMID 30371827

37.　Dong, S. *et al.* Annotating and prioritizing human non-coding variants with RegulomeDB

v.2. *Nat Genet* **55**, 724-726 (2023). PMID 37173523

38.　Schipper, M. *et al.* Prioritizing effector genes at trait-associated loci using multimodal

evidence. *Nat Genet* **57**, 323-333 (2025). PMID 39930082

39.　Hutchinson, A., Watson, H. & Wallace, C. Improving the coverage of credible sets in

Bayesian genetic fine-mapping. *PLoS Comput Biol* **16**, e1007829 (2020). PMID

32282791

40.　Weeks, E.M. *et al.* Leveraging polygenic enrichments of gene features to predict genes

underlying complex traits and diseases. *Nat Genet* **55**, 1267-1276 (2023). PMID

37443254

41.   Pruim, R.J. *et al.* LocusZoom: regional visualization of genome-wide association scan

754   results. *Bioinformatics* **26**, 2336-7 (2010). PMID 20634204

755   42.   de Leeuw, C.A., Mooij, J.M., Heskes, T. & Posthuma, D. MAGMA: generalized gene-set

756   analysis of GWAS data. *PLoS Comput Biol* **11**, e1004219 (2015). PMID 25885710

757   43.   Liberzon, A. *et al.* The Molecular Signatures Database (MSigDB) hallmark gene set

758   collection. *Cell Syst* **1**, 417-425 (2015). PMID 26771021

759   44.   van der Valk, W.H. *et al.* A single-cell level comparison of human inner ear organoids

760   with the human cochlea and vestibular organs. *Cell Rep* **42**, 112623 (2023). PMID

761   37289589

762   45.   Jean, P. *et al.* Single-cell transcriptomic profiling of the mouse cochlea: An atlas for

763   targeted therapies. *Proc Natl Acad Sci U S A* **120**, e2221744120 (2023). PMID

764   37339214

765   46.   Hoa, M. *et al.* Characterizing Adult Cochlear Supporting Cell Transcriptional Diversity

766   Using Single-Cell RNA-Seq: Validation in the Adult Mouse and Translational Implications

767   for the Adult Human Cochlea. *Front Mol Neurosci* **13**, 13 (2020). PMID 32116546

768   47.   Blake, J.A. *et al.* Mouse Genome Database (MGD): Knowledgebase for mouse-human

769   comparative biology. *Nucleic Acids Res* **49**, D981-D987 (2021). PMID 33231642

770   48.   Bulik-Sullivan, B.K. *et al.* LD Score regression distinguishes confounding from

771   polygenicity in genome-wide association studies. *Nat Genet* **47**, 291-5 (2015). PMID

772   25642630

773   49.   Lee, S.H., Wray, N.R., Goddard, M.E. & Visscher, P.M. Estimating missing heritability for

774   disease from genome-wide association studies. *Am J Hum Genet* **88**, 294-305 (2011).

775   PMID 21376301

776   50.   Holland, D. *et al.* Beyond SNP heritability: Polygenicity and discoverability of phenotypes

777   estimated with a univariate Gaussian mixture model. *PLoS Genet* **16**, e1008612 (2020).

778   PMID 32427991

779   51.   Frei, O. *et al.* Bivariate causal mixture model quantifies polygenic overlap between

780   complex traits beyond genetic correlation. *Nat Commun* **10**, 2417 (2019). PMID

781   31160569

782   52.   Johnson, K.R., Zheng, Q.Y. & Noben-Trauth, K. Strain background effects and genetic

783   modifiers of hearing in mice. *Brain Res* **1091**, 79-88 (2006). PMID 16579977

784   53.   Yankaskas, K. Prelude: noise-induced tinnitus and hearing loss in the military. *Hear Res*

785   **295**, 3-8 (2013). PMID 22575206

786   54.   Chen, J.X. *et al.* Systematic review of hearing loss after traumatic brain injury without

787   associated temporal bone fracture. *Am J Otolaryngol* **39**, 338-344 (2018). PMID

788   29506762

789   55.   Escalera-Balsera, A., Roman-Naranjo, P. & Lopez-Escamez, J.A. Systematic Review of

790   Sequencing Studies and Gene Expression Profiling in Familial Meniere Disease. *Genes*

791   *(Basel)* **11**(2020). PMID 33260921

792   56.   Hoischen, A. *et al.* De novo mutations of SETBP1 cause Schinzel-Giedion syndrome.

793   *Nat Genet* **42**, 483-5 (2010). PMID 20436468

794   57.   Munnamalai, V. & Fekete, D.M. Wnt signaling during cochlear development. *Semin Cell*

795   *Dev Biol* **24**, 480-9 (2013). PMID 23548730

796   58.   Dabdoub, A. *et al.* Wnt signaling mediates reorientation of outer hair cell stereociliary

797   bundles in the mammalian cochlea. *Development* **130**, 2375-84 (2003). PMID 12702652

798   59.   Yan, K. *et al.* EVL is not essential for cuticular plate and stereocilia development in

799   mouse auditory hair cells. *FEBS Lett* **599**, 330-339 (2025). PMID 39300480

800   60.   Xia, R. *et al.* Therapeutic restoration of miR-96 prevents hearing loss in mice through

801   modulation of noise-induced and genetic pathways. *iScience* **28**, 112355 (2025). PMID

802   40641557

803   61.   Wang, J. *et al.* Polygenic Risk Scores and Hearing Loss Phenotypes in Children. *JAMA*

804   *Otolaryngol Head Neck Surg* (2024). PMID 39509092

62.   Hui, D. *et al.* Gene burden analysis identifies genes associated with increased risk and
severity of adult-onset hearing loss in a diverse hospital-based cohort. *PLoS Genet* **19**,
e1010584 (2023). PMID 36656851

63.   Lewis, M.A. *et al.* Whole exome sequencing in adult-onset hearing loss reveals a high
load of predicted pathogenic variants in known deafness-associated genes and identifies
new candidate genes. *BMC Med Genomics* **11**, 77 (2018). PMID 30180840

64.   Lewis, M.A., Schulte, B.A., Dubno, J.R. & Steel, K.P. Investigating the characteristics of
genes and variants associated with self-reported hearing difficulty in older adults in the
UK Biobank. *BMC Biol* **20**, 150 (2022). PMID 35761239

65.   Cornejo-Sanchez, D.M. *et al.* Rare-variant association analysis reveals known and new
age-related hearing loss genes. *Eur J Hum Genet* **31**, 638-647 (2023). PMID 36788145

66.   Khela, H. & Kenna, M.A. Genetics of pediatric hearing loss: A functional perspective.
Laryngoscope Investig Otolaryngol **5**, 511-519 (2020). PMID 32596495

67.   Sekerkova, G., Zheng, L., Loomis, P.A., Mugnaini, E. & Bartles, J.R. Espins and the
actin cytoskeleton of hair cell stereocilia and sensory cell microvilli. *Cell Mol Life Sci* **63**,
2329-41 (2006). PMID 16909209

68.   Dunbar, L.A. *et al*. Clarin-2 is essential for hearing by maintaining stereocilia integrity
and function. *EMBO Mol Med* **11**, e10288 (2019). PMID 31448880

69.   Katsuno, T. *et al.* TRIOBP-5 sculpts stereocilia rootlets and stiffens supporting cells
enabling hearing. *JCI Insight* **4**(2019). PMID 31217345

70.   Fasquelle, L. *et al.* Tmprss3, a transmembrane serine protease deficient in human
DFNB8/10 deafness, is critical for cochlear hair cell survival at the onset of hearing. *J*
Biol Chem **286**, 17383-97 (2011). PMID 21454591

71.   Liu, W. *et al.* Expression of trans-membrane serine protease 3 (TMPRSS3) in the human
organ of Corti. *Cell Tissue Res* **372**, 445-456 (2018). PMID 29460002

72.   Pacentine, I., Chatterjee, P. & Barr-Gillespie, P.G. Stereocilia Rootlets: Actin-Based

831   Structures That Are Essential for Structural Stability of the Hair Bundle. *Int J Mol Sci*

832   **21**(2020). PMID 31947734

833   73.   Cooper, N.P., Vavakou, A. & van der Heijden, M. Vibration hotspots reveal longitudinal

834   funneling of sound-evoked motion in the mammalian cochlea. *Nat Commun* **9**, 3054

835   (2018). PMID 30076297

836   74.   Gagnon, L.H. *et al.* The chloride intracellular channel protein CLIC5 is expressed at high

837   levels in hair cell stereocilia and is essential for normal inner ear function. *J Neurosci* **26**,

838   10188-98 (2006). PMID 17021174

839   75.   Silverstein, R.S. & Tempel, B.L. Atp2b2, encoding plasma membrane Ca2+-ATPase type

840   2, (PMCA2) exhibits tissue-specific first exon usage in hair cells, neurons, and mammary

841   glands of mice. *Neuroscience* **141**, 245-57 (2006). PMID 16675132

842   76.   Cousin, M.A. *et al.* Pathogenic SPTBN1 variants cause an autosomal dominant

843   neurodevelopmental syndrome. *Nat Genet* **53**, 1006-1021 (2021). PMID 34211179

844   77.   Wang, P. *et al.* LOXHD1 is indispensable for coupling auditory mechanosensitive

845   channels to the site of force transmission. *Res Sq* (2024). PMID 38260480

846   78.   Shpargel, K.B., Makishima, T. & Griffith, A.J. Col11a1 and Col11a2 mRNA expression in

847   the developing mouse cochlea: implications for the correlation of hearing loss phenotype

848   with mutant type XI collagen genotype. *Acta Otolaryngol* **124**, 242-8 (2004). PMID

849   15141750

850   79.   Chen, T. *et al.* Cochlear supporting cells require GAS2 for cytoskeletal architecture and

851   hearing. *Dev Cell* **56**, 1526-1540 e7 (2021). PMID 33964205

852   80.   Stover, E.H. *et al.* Novel ATP6V1B1 and ATP6V0A4 mutations in autosomal recessive

853   distal renal tubular acidosis with new evidence for hearing loss. *J Med Genet* **39**, 796-

854   803 (2002). PMID 12414817

855   81.   Szeto, I.Y.Y. *et al.* SOX9 and SOX10 control fluid homeostasis in the inner ear for

856   hearing through independent and cooperative mechanisms. *Proc Natl Acad Sci U S A*

857   **119**, e2122121119 (2022). PMID 36343245

858   82.   Fernandez, K.A. *et al.* Noise-induced Cochlear Synaptopathy with and Without Sensory

859   Cell Loss. *Neuroscience* **427**, 43-57 (2020). PMID 31887361

860   83.   Kujawa, S.G. & Liberman, M.C. Synaptopathy in the noise-exposed and aging cochlea:

861   Primary neural degeneration in acquired sensorineural hearing loss. *Hear Res* **330**, 191-

862   9 (2015). PMID 25769437

863   84.   Bramhall, N.F. & McMillan, G.P. Perceptual Consequences of Cochlear Deafferentation

864   in Humans. *Trends Hear* **28**, 23312165241239541 (2024). PMID 38738337

865   85.   Panganiban, C.H. *et al.* Noise-Induced Dysregulation of Quaking RNA Binding Proteins

866   Contributes to Auditory Nerve Demyelination and Hearing Loss. *J Neurosci* **38**, 2551-

867   2568 (2018). PMID 29437856

868   86.   Jean, P., Ozcete, O.D., Tarchini, B. & Moser, T. Intrinsic planar polarity mechanisms

869   influence the position-dependent regulation of synapse properties in inner hair cells.

870   *Proc Natl Acad Sci U S A* **116**, 9084-9093 (2019). PMID 30975754

871   87.   Aoki, H., Hara, A. & Kunisada, T. White spotting phenotype induced by targeted REST

872   disruption during neural crest specification to a melanocyte cell lineage. *Genes Cells* **20**,

873   439-49 (2015). PMID 25818501

874   88.   Wingfield, A. *et al.* A twin-study of genetic contributions to hearing acuity in late middle

875   age. *J Gerontol A Biol Sci Med Sci* **62**, 1294-9 (2007). PMID 18000151

Figure Legends

Figure 1. Comparison of SNHL between the clinically diagnosed MVP and self-reported UKB

GWAS.

a-b) Manhattan plots of sensorineural hearing loss (SNHL) GWAS. The red line represents

genome-wide significance (GWS) at p < 5 × 10−8. a) MVP GWAS of clinically diagnosed SNHL

in European ancestry participants (176,393 cases and 172,117 controls) identified 43 GWS loci.

b) UKB GWAS of self-reported SNHL (87,056 cases and 163,333 controls) identified 41 GWS

loci. The X chromosome was not analyzed in the UKB GWAS. **c)** Quantification of the polygenic

overlap between SNHL in the MVP and UKB. Bivariate MiXeR modeling estimated 3,624 causal

variants are shared between cohorts (gray). There was only a small amount of polygenicity

unique to MVP (blue), such that shared polygenicity accounted for 95.6% of the causal variants

influencing MVP. The UKB had a larger unique polygenic component (green), such that shared

polygenicity accounted for 69.2% of the causal variants influencing UKB. The numbers (in

thousands, with standard errors in parenthesis) in the Venn diagram indicate the estimated

quantity of causal variants per component. The size of the circles reflects the degree of

polygenicity. Genetic correlation (rg) estimated between the two phenotypes is shown below the

Venn diagram.

Figure 2. Risk loci (N = 108) and gene discovery (N = 192) for SNHL in a multi-ancestry meta-

analysis.

Manhattan plots of SNP **a)** and gene-level **b)** associations from meta-analysis of the MVP multi-

ancestry and UKB sensorineural hearing loss (SNHL) GWAS (297,296 cases and 428,608

controls). The red line represents genome-wide significance at p < 5 × 10−8 (panel a) and gene-

wide significance p < 2.66 × 10−6 (panel b, Bonferroni correction for 18,809 genes tested),

respectively. A selection of GWS genes are labeled, and genes listed in

HereditaryHearingLoss.org are labeled in bold font.

Figure 3. Putative influences of significant hearing loss genes on cochlear function.

a) Cross-section of ear. Sound waves travel through the external auditory canal, vibrating the

tympanic membrane and ossicular chain, continuing as a fluid wave into the cochlea. The Organ

of Corti (OC) is suspended within the center of the cochlea (green). **b)** Cross-section of the OC.

supported by the basilar membrane below (BM). Movement is restricted by the tectorial

membrane above. The shearing motion of the fluid endolymph wave motion against OHC

stereocilia induces a dampening, focusing the wave towards the locally juxtaposed inner hair

cells (IHCs). IHCs then transduce this movement into an electric signal, releasing

neurotransmitter at the ribbon synapse. Post-synaptic neurons of the cochlear nerve transmit

signals to the auditory cortex. The stria vascularis maintains the electric potential of endolymph

that bathes the apices of hair cells. **c)** Expanded area of apex of hair cells from b., showing

stereociliary movement of both OHCs and IHCs. Fluid/sound wave movement pushes

stereocilia laterally, stretching tip-links attaching the taller stereocilium to its adjacent shorter

one. The stereocilium's parallel f-actin fibers are critical for stiffness and rigidity of the

stereocilia, while allowing for bending at the reinforced rootlet at the attachment of the

stereocilia to the apical surface of the hair cell. The rootlet is the focus area of vibration

secondary to sound as measured by high-resolution optics[73]. Tip-links consist of chained protein

threads connecting sequential rows of stereociliae, from the top of the shorter stereocilia to the

side of the next taller structure This movement opens a mechanoelectrical transduction (MET)

channel at its tip (black), allowing influx of Ca^{++} and K^+ and triggering depolarization at the

base of the cell. Tight junctions between hair cells and supporting cells separate endolymph

bathing the stereocilia from its cell bodies. Genes that are not color-coded are ubiquitous,

involved in development and/or maintenance.

a

Ossicular Chain

Tympanic Membrane

Organ of Corti

b

Structural support
COL11A1*
COL9A3
GAS2

Stria vascularis
(melanocytes)
SOX10*
ATP6V0A4*

TM

BM

Neurons and
ribbon synapse
LMX1A
REST*
TUB
QKI*
SVIP*

inner hair
cells
EYA4
CLRN2

pillar
cells

outer hair
cells
EYA4
CLRN2

Deiter cells

c

Stereociliary Orientation
WNT7A*

Sound Wave

TIP-LINK and MET
● *CDH23*
● *MYO7A**
● *LOXHD1*
   CLRN2

Actin Binding
ESPN
TMPRSS3
CLRN2
GAS2

Ca²⁺
● *ATP2B2*

Rootlet
TRIOBP
SPTBN1
● *CLIC5**

Cuticular Plate
*EVL

Tight junctions
ILDR1

Bold – known congenital hearing loss genes
* - genes novel to this study

Table 1. Genome-wide significant loci in meta-analysis of sensorineural hearing los

Locus	Chr	Start[a]	End[a]	Lead SNP	A1
1	1	6,355,065	6,505,159	rs112725535	A
2	1	16,367,202	16,424,085	rs2863455	T
3	1	25,580,095	25,863,694	rs2281179	T
4	1	45,768,792	46,644,026	rs201377643	C
5	1	61,858,621	61,868,766	rs2499507	T
7	1	102,902,818	103,757,081	rs12740738	A
8	1	165,086,665	165,126,202	rs7525101	T
9	1	168,745,350	168,902,339	rs7555365	T
10	1	170,058,829	170,196,129	rs501925	A
11	1	205,663,993	205,757,824	rs823116	A
13	2	43,450,843	43,602,415	rs67278917	CCA
14	2	54,687,254	55,062,865	rs2941580	A
15	2	112,046,160	112,069,485	rs2341098	T
16	2	208,000,822	208,118,374	rs2216374	T
17	3	10,411,827	10,521,964	rs511273	T
18	3	13,831,209	13,855,799	rs877306	T
19	3	24,267,011	24,267,842	rs6779690	T
20	3	71,353,372	71,628,286	rs75175086	T
23	3	121,351,315	121,729,262	rs3915060	T
24	3	127,710,474	128,095,278	rs4857868	A
25	3	156,243,368	156,303,912	rs9856073	T
26	3	181,935,178	182,210,347	rs7649191	A
28	4	3,285,389	3,433,782	rs3129327	A
30	4	5,860,906	5,919,670	rs6446406	T
31	4	10,011,996	10,412,191	rs10489074	A
32	4	17,517,558	17,530,692	rs13147559	C
33	4	57,709,860	57,903,482	rs13152711	T
34	5	2,549,211	2,564,723	rs7712395	T
35	5	32,323,623	32,449,816	rs2963985	A
36	5	68,047,930	68,253,056	rs4246760	T
37	5	72,823,715	73,199,539	rs6453022	A
38	5	82,299,375	82,547,602	rs2386237	A
40	5	92,684,177	93,571,190	rs167570	T
42	5	150,939,707	150,966,388	rs9688110	A
43	5	168,841,685	168,896,564	rs3074558	A
44	6	7,066,525	7,153,152	rs11243143	A
45	6	21,960,455	21,970,417	rs1928176	A
46	6	26,354,100	29,607,101	rs66868086	A
47	6	43,201,763	43,434,705	rs2242416	A
48	6	45,942,037	45,979,229	rs3777621	T
49	6	80,052,589	80,087,011	rs2038297	T
50	6	83,978,225	84,409,255	rs217308	T
51	6	90,645,615	90,673,538	rs9451298	T
52	6	133,273,015	133,869,809	rs9493627	A
53	6	147,946,333	147,986,716	rs9390488	T
54	6	158,497,717	158,619,409	rs3818457	T

55	6	163,789,736	164,005,951	rs1737332	T
56	7	19,548,842	19,654,201	rs10232942	A
57	7	50,734,630	50,882,081	rs1962390	C
58	7	86,198,818	86,340,497	rs274625	A
59	7	138,432,792	138,514,861	rs7780560	A
61	8	21,904,891	21,944,508	rs113973451	T
62	8	30,271,637	30,518,967	rs3824107	A
63	8	71,467,435	72,012,927	rs62506938	T
64	8	74,206,582	74,254,612	rs7819550	A
65	8	82,653,644	82,829,051	rs78724141	T
66	8	99,925,359	100,804,670	rs11784152	A
67	8	141,604,684	142,018,581	rs12156228	T
68	9	84,736,303	85,036,608	rs1409880	A
69	9	96,349,538	96,484,560	rs10821202	A
70	10	17,172,952	17,248,287	rs12355391	A
71	10	63,828,879	63,841,130	rs2393729	T
73	10	75,783,340	76,358,087	rs377150672	T
74	10	80,273,287	80,610,121	rs149979867	A
75	10	82,379,843	82,409,617	rs71483314	A
76	10	94,759,464	94,831,523	rs12771031	T
78	10	126,755,203	126,874,931	rs10901863	T
79	11	8,053,304	8,085,652	rs4483583	A
80	11	8,826,688	9,046,767	rs4910179	T
81	11	13,178,285	13,197,316	rs10831990	T
82	11	15,993,057	16,224,877	rs11023821	A
83	11	22,797,251	22,880,918	rs11543287	A
85	11	88,358,544	89,204,911	rs1126809	A
86	11	118,477,367	118,735,476	rs67307131	T
87	12	26,114,019	26,193,294	rs11048385	T
88	12	109,860,891	110,042,348	rs2111216	A
89	13	26,624,935	26,688,841	rs111516934	T
91	13	99,050,232	99,075,551	rs7329659	A
92	14	52,453,659	52,542,480	rs1566129	T
93	14	65,165,090	65,217,898	rs229650	T
94	14	73,961,821	74,018,555	rs35417585	A
95	14	75,503,759	75,668,384	rs35499335	G
96	14	85,288,673	85,409,941	rs12588043	A
97	14	100,571,762	100,611,949	rs1951486	T
99	15	57,132,863	57,586,614	rs59425097	CT
100	15	74,713,300	75,029,407	rs11858525	T
101	15	79,219,188	79,257,278	rs34593439	A
102	15	89,040,591	89,274,382	rs12594617	C
103	16	51,163,406	51,189,169	rs11643654	A
104	16	53,797,908	53,845,487	rs62033400	A
105	16	79,135,887	79,167,945	rs10492901	T
106	16	81,587,072	81,616,668	rs59164848	T
107	17	2,473,537	2,612,584	rs734957	A
108	17	6,193,984	8,130,452	rs34958987	T
109	17	17,628,678	18,140,308	rs117283876	A

111	17	43,573,649	44,351,285	rs2458203	T
113	17	65,822,573	66,098,154	rs56305452	A
114	17	79,585,385	79,689,749	rs58038388	T
115	18	6,674,194	6,755,963	rs2228712	A
116	18	41,994,736	42,298,323	rs2852779	T
117	18	52,573,169	52,669,732	rs11152090	A
118	18	53,210,302	53,466,338	rs1031829	A
120	20	1,328,766	1,382,035	rs6041774	T
122	20	3,217,989	3,415,566	rs6115827	A
123	20	61,440,005	61,455,532	rs61734651	T
124	21	43,809,092	43,831,005	rs45598239	T
125	22	38,077,995	38,503,479	rs5756795	T
126	22	50,952,735	51,157,353	rs36062310	A

[a]Locus start and end positions (in bp).

[b]Bolded p values indicate p values below genome-wide significance threshold ($p < 5 \times 10^{-8}$).

Abbreviations: GWS, Genome Wide Significant ($p < 5 \times 10^{-8}$); Locus, the index number identifier
SNP, single nucleotide polymorphism; Chr, Chromosome; Start, locus start position in base pairs
stop position; A1, coded allele (effects and allele frequencies are coded in terms of copies of this a
guanidine; T, thymidine; A1 Freq, frequency of allele 1.

A2	A1 Freq	Beta	SE	p Value[b]
G	0.1794	-0.0316	0.0049	9.86E-11
C	0.6774	-0.0257	0.0042	7.48E-10
C	0.5362	-0.0254	0.0036	2.33E-12
G	0.3229	0.0415	0.0039	4.20E-26
C	0.4596	0.0226	0.0038	3.71E-09
G	0.4487	-0.0337	0.0037	1.27E-19
C	0.4231	0.0275	0.0037	6.53E-14
C	0.6227	0.0346	0.0040	5.66E-18
G	0.6110	-0.0254	0.0038	1.78E-11
G	0.5209	-0.0206	0.0037	2.94E-08
C	0.7598	-0.0258	0.0043	2.56E-09
G	0.4731	-0.0307	0.0036	1.72E-17
G	0.4352	-0.0231	0.0042	2.74E-08
C	0.4022	0.0261	0.0038	7.82E-12
C	0.3401	0.0246	0.0041	2.15E-09
C	0.3997	-0.0255	0.0038	2.21E-11
C	0.8062	-0.0278	0.0050	2.13E-08
C	0.7695	-0.0284	0.0044	7.02E-11
C	0.7209	-0.0248	0.0042	2.59E-09
T	0.8352	-0.0318	0.0052	1.21E-09
G	0.7969	0.0300	0.0046	5.93E-11
G	0.2846	-0.0364	0.0043	1.18E-17
G	0.6176	0.0224	0.0040	2.86E-08
C	0.3576	0.0220	0.0038	5.53E-09
C	0.2456	0.0249	0.0042	4.34E-09
G	0.8729	-0.0368	0.0057	9.45E-11
C	0.2665	-0.0240	0.0042	1.13E-08
C	0.1196	0.0340	0.0056	9.64E-10
G	0.4183	-0.0237	0.0037	9.74E-11
C	0.6608	0.0284	0.0040	1.16E-12
C	0.5190	0.0435	0.0036	2.75E-33
G	0.6066	-0.0204	0.0037	4.38E-08
C	0.1273	-0.0354	0.0055	9.83E-11
G	0.3100	0.0243	0.0041	2.63E-09
AAG	0.6338	-0.0242	0.0039	6.70E-10
G	0.5951	0.0216	0.0038	1.39E-08
G	0.5314	-0.0214	0.0037	8.99E-09
T	0.9024	0.0381	0.0063	1.77E-09
G	0.4268	-0.0454	0.0037	5.21E-34
G	0.6368	0.0208	0.0038	4.91E-08
C	0.8133	0.0264	0.0047	2.31E-08
C	0.5841	-0.0241	0.0037	4.72E-11
C	0.7152	0.0270	0.0041	3.53E-11
G	0.3445	0.0444	0.0038	1.38E-31
C	0.5901	-0.0208	0.0037	2.62E-08
C	0.4838	-0.0240	0.0037	5.69E-11

C	0.7367	-0.0268	0.0042	**2.95E-10**
G	0.3886	-0.0251	0.0039	**9.44E-11**
G	0.2272	-0.0319	0.0044	**3.99E-13**
C	0.6072	-0.0210	0.0037	**2.13E-08**
G	0.4679	0.0255	0.0037	**3.28E-12**
G	0.2072	-0.0313	0.0047	**2.04E-11**
G	0.2264	-0.0244	0.0043	**1.81E-08**
G	0.9171	-0.0360	0.0065	**3.62E-08**
G	0.1878	0.0258	0.0046	**3.00E-08**
G	0.0662	0.0561	0.0073	**2.21E-14**
G	0.2487	-0.0241	0.0043	**1.84E-08**
G	0.6012	-0.0251	0.0040	**2.99E-10**
G	0.6194	0.0218	0.0039	**2.40E-08**
C	0.3817	-0.0209	0.0037	**1.54E-08**
C	0.5816	-0.0227	0.0038	**2.54E-09**
C	0.3972	-0.0231	0.0037	**6.00E-10**
TA	0.5980	-0.0257	0.0039	**5.92E-11**
G	0.1962	-0.0334	0.0047	**7.27E-13**
G	0.0211	0.0817	0.0145	**1.75E-08**
C	0.5120	-0.0228	0.0037	**1.23E-09**
C	0.2584	0.0501	0.0046	**5.45E-28**
C	0.8035	0.0360	0.0046	**2.54E-15**
G	0.3294	0.0230	0.0040	**6.30E-09**
C	0.7407	0.0298	0.0044	**9.96E-12**
G	0.4228	0.0202	0.0037	**3.59E-08**
G	0.3887	-0.0328	0.0038	**9.83E-18**
G	0.2838	0.0348	0.0042	**5.82E-17**
C	0.6240	-0.0300	0.0038	**5.25E-15**
C	0.5677	-0.0205	0.0037	**4.34E-08**
G	0.3891	-0.0211	0.0038	**2.40E-08**
C	0.1611	0.0324	0.0050	**1.19E-10**
G	0.6671	-0.0279	0.0040	**2.17E-12**
C	0.4221	0.0407	0.0036	**3.91E-29**
C	0.7543	-0.0300	0.0043	**1.78E-12**
G	0.7602	0.0249	0.0045	**3.50E-08**
GT	0.5835	0.0220	0.0039	**1.96E-08**
C	0.4712	-0.0219	0.0038	**5.18E-09**
C	0.4364	0.0248	0.0037	**2.99E-11**
C	0.7100	0.0252	0.0044	**1.32E-08**
G	0.8445	0.0380	0.0063	**1.62E-09**
G	0.0973	-0.0436	0.0064	**1.19E-11**
G	0.8041	0.0563	0.0047	**1.36E-32**
C	0.5703	-0.0208	0.0038	**3.89E-08**
G	0.6221	0.0249	0.0038	**4.42E-11**
C	0.6241	0.0246	0.0037	**4.91E-11**
TG	0.4407	0.0242	0.0037	**4.76E-11**
G	0.2317	0.0256	0.0043	**2.44E-09**
TG	0.4025	0.0249	0.0037	**1.53E-11**
G	0.0237	0.0806	0.0131	**7.00E-10**

C	0.6810	-0.0270	0.0043	**3.25E-10**
C	0.7191	0.0242	0.0042	**1.22E-08**
C	0.0251	0.0847	0.0155	**4.24E-08**
C	0.5154	0.0206	0.0037	**2.02E-08**
G	0.8384	-0.0396	0.0049	**4.52E-16**
C	0.7935	-0.0267	0.0046	**5.30E-09**
G	0.8327	0.0316	0.0050	**2.82E-10**
C	0.1053	0.0340	0.0060	**1.39E-08**
G	0.6673	0.0220	0.0039	**1.75E-08**
C	0.0650	0.0539	0.0082	**6.33E-11**
C	0.0524	0.0673	0.0084	**1.13E-15**
C	0.5702	-0.0360	0.0037	**1.18E-22**
G	0.0411	0.1147	0.0101	**8.65E-30**

 used to refer to independent loci identified by this GWAS;
(GR37 Human Genome Build/hg19 coordinates) , Stop, locus
allele); A2, non-coded allele; A, adenosine; C, cytosine; G,

Table 2. Partitioned heritability estimates for sensorineural hearing loss GWAS (N = 59

Category	SNPs	h^2_{SNP}
All SNPs	1.00	1.00
SNPs in protein coding genes (in HHLdb)	0.01	0.03
SNPs in protein coding genes (not in HHLdb)	0.38	0.54
SNPs outside of protein coding genes	0.61	0.43

[a] Calculated as the proportion of h^2_{SNP} accounted for by the category divided by the proportio

Abbreviations: HHLdb, Hereditary Hearing Loss Database; h^2_{SNP}, SNP-based heritability

8,899 European subjects from MVP and UKB)

SE	Enrichment[a]
0.00	1.00
0.01	4.58
0.02	1.40
0.02	0.71

n of the genome-wide set of SNPs that were annotated to the category