Estimating the potential malaria morbidity and mortality avertable by the President's Malaria Initiative in 2025: a geospatial modelling analysis

Tasmin L. Symons, PhD[1,2], Jailos Lubinda, PhD[2], Michael McPhail, DPhil[2], Adam Saddler, PhD[2], Mauricio van den Berg, MSc[2], Hunter Baggen, MPH[2], Yuval Berman, BPhil(Hons)[2], Sarah Hafsia, PhD[2], Rubi Jayaseelen, MSc[2], Punam Amratia, PhD[3,2,1], Annie Browne, DPhil[2], Ewan Cameron, PhD[1,2], Camilo Vargas-Ruiz, MSc[2], Susan F. Rumisha, PhD[3,2,1], Prof. Nick Golding, DPhil[4, 2, 5], Daniel J. Weiss, PhD[1,2], Prof. Peter W. Gething, PhD[1,2, *]

1. School of Population Health, Curtin University, Bentley, Western Australia.
2. Malaria Atlas Project, The Kids Research Institute Australia, Nedlands, Western Australia.
3. Malaria Atlas Project, Ifakara Health Institute, Dar es Salaam, Tanzania.
4. School of Physics, Mathematics and Computing, University of Western Australia, Western Australia.
5. Melbourne School of Population and Global Health, University of Melbourne, Victoria.

*Corresponding author: Prof. Peter W. Gething, peter.gething@thekids.org.au, +61 8 6319 1000

Abstract

Background

Since its inception in 2005, the President's Malaria Initiative (PMI) has played a major role in the reductions in malaria morbidity and mortality witnessed across Africa. With the status of PMI funding and operations currently uncertain, this study aimed to quantify the impact that a fully-functioning PMI would have on malaria cases and deaths in Africa during 2025.

Methods

We combined detailed spatio-temporal information on planned 2025 PMI and non-PMI malaria commodity procurement and distribution in Africa with spatio-temporal Bayesian models of intervention coverage and *Plasmodium falciparum* transmission and burden in Africa. By comparing coverage scenarios with and without planned PMI contributions we estimated the number of malaria cases and deaths PMI would avert in 2025.

Findings

We estimated that business-as-usual PMI contributions to vector control, seasonal chemoprevention, and routine malaria treatment in Africa would avert in 2025 13·6M (95% uncertainty interval 11·4M – 16·4M) malaria cases and 104,000 (69,000 – 161,000) deaths. This represents 11·3% (9·9 – 12·9%) and 37·5% (34·2 – 41·5%), respectively, of the total burden of malaria morbidity and mortality in PMI's focus geographies across 27 African countries. These estimates do not account for the additional impact of PMI-supported provision of diagnostics or severe case management commodities, nor preventive treatment for pregnant women which would add further to the averted burden.

Interpretation

PMI investment in supporting procurement and distribution of malaria control commodities would translate directly into millions of malaria cases averted and a hundred thousand lives saved across its focus geographies in Africa across 2025.

Funding

Research in context

Evidence before this study

We searched PubMed on 25 February 2025 with the search term "President's Malaria Initiative" AND (impact OR evaluation) AND (burden OR malaria cases OR malaria deaths). 102 results were returned, of which four directly addressed the question of PMI-attributable impact on malaria burden across all its focus geographies. One focussed solely on case-management and did not provide quantitative results[1]. A study focussing on mortality found a strong association between PMI investment and declines in all-cause infant mortality[2]. Another study used a generalised estimating equation approach to assess PMI's impact, concluding a strong association between PMI spending and reductions in malaria burden[3]. A 2017 study used a mechanistic malaria model to estimate PMI's historic contribution to reductions in malaria morbidity and mortality, with future projections to 2020[4].

Added value of this study

In this study we synthesised the most up-to-date information of all-funder volumes of key malaria control interventions (ITN, IRS, ACT, SMC) with PMI data on planned volumes and spatial targeting of funding in 2025 to derive near-real-time projections of malaria control intervention coverage in Africa under two scenarios: a business-as-usual scenario in which PMI commodities procured and distributed as previously planned versus a 'no-PMI' scenario in which PMI funding and technical assistance is absent. These scenario-based estimates of coverage were propagated through an empirical geospatial malaria model, coupled to mechanistic models of clinical incidence, in order to estimate the number of malaria infections, clinical cases, and deaths PMI activities could expect to avert in Africa in 2025.

Implications of all available evidence

PMI is a key contributor to malaria control efforts in Africa, and as such its impact on averting childhood mortality has been demonstrated repeatedly. Here, using empirically-derived estimates of malaria control impact, combined with detailed scenarios of intervention coverage with and without PMI support, we isolated the contemporary impact of PMI activities on malaria morbidity and mortality in Africa in 2025. Our findings suggest that PMI support is critical to maintaining suppression of malaria transmission in its focus geographies.

Introduction

Over the past 20 years, malaria control has greatly reduced malaria transmission, cases, and deaths in Africa[5,6]. A catalyst of this achievement has been the growth in international development assistance for health, including the launch in 2002 of the Global Fund followed in 2005 by the US President's Malaria Initiative (PMI). By 2023 PMI supported malaria control in 31 countries (27 in Africa), and disbursed USD$777·7M[7]. That investment enabled, in the 2023 US financial year, the procurement of 36·8M insecticide treated bednets (ITNs), 63·3M doses of the effective antimalarial drug artemisinin-based combination therapy (ACT), indoor residual spraying (IRS) of insecticides to protect 15·5M people, and 48M treatments of seasonal malaria chemoprevention (SMC) to protect children in locations with intense malaria seasonality[7]. Most of this financial support was directed to the highest burden regions of malaria-endemic Africa[6,7]. In addition to core vector control and treatment, PMI has also supported the large-scale deployment of preventive treatment for pregnant women, the scale-up of rapid diagnostic tests (RDTs) to support accurate point-of-care diagnosis, and other wide-ranging technical support to health personnel working in malaria-endemic countries. The immediate and longer-term future of PMI is currently uncertain, and the extent to which it's planned support for malaria control across Africa in 2025 will eventuate is not clear. Given the large proportional contribution of PMI to overall malaria control efforts in the highest-burden regions, the presence or absence of PMI financing and operations has the potential to significantly impact total malaria burden in Africa. Estimates of the overall potential impact of PMI in 2025, how this varies by country, and the relative contributions of different intervention tools, can therefore help inform ongoing decision making at international and national levels.

Various approaches have been developed to estimate - retrospectively or prospectively - the malaria burden averted by investments in malaria control efforts[8,9]. An initial challenge is to understand how those investments have translated into coverage of malaria control interventions. A common method is to assume that funds invested translate proportionally into coverage achieved, enabling coverage to be predicted using simple arithmetic based on commodity unit costs multiplied by funds available. While reasonable for some interventions, this assumption is not always valid and evidence demonstrates, for example, that incremental increases in coverage of ITNs becomes progressively more expensive to achieve above a certain threshold[10,11]. In this study, we deploy a range of coverage models that more explicitly

link investments to commodity volumes, to deployment, and ultimately to population coverage.

The second challenge is to estimate the likely impact interventions have on malaria at a given

level of coverage. In this study we build on the modelling architecture of the Malaria Atlas

Project (MAP)[12] used to produce annually updated estimates of malaria burden and trends for

the WHO World Malaria Report[6] and Global Burden of Disease study[13]. We use this framework

to generate modelled estimates of the potential of PMI to avert malaria cases and deaths in

2025. We define a business-as-usual scenario in which we reconstruct malaria control

commodity procurement and distribution as previously planned for 2025 and compare this to

a scenario in which activities supported by PMI do not take place.

Methods

Figure 1 provides a schematic overview of our analysis, which comprised the following steps.

First, we assembled data on past and planned malaria control commodity procurement and

distribution from both PMI and non-PMI funding sources and focusing on ITNs, ACTs, SMC,

and IRS. Second, we used coverage models to estimate subnational coverage of each of these

interventions in 2025 in scenarios with and without contributions from PMI. Third, we used an

established Bayesian geostatistical model to predict infection prevalence across Africa in 2025

under both scenarios and, fourth, converted these into corresponding estimates of malaria

clinical incidence and mortality. These steps are described in more detail below and with

supporting methodological descriptions in the appendix.

Determining malaria control coverage attributable to PMI in 2025

The Roll Back Malaria Initiative (RBM) hosts programmatic and financial gap analysis tables

that include data – by malaria-endemic country – detailing the number of ITNs financed for

planned distribution in 2024 and 2025, reflecting a business-as-usual (BAU) scenario[14]. These

numbers are inclusive of all sources of funding, including PMI. At the time of analysis, this

information was current as of 17 December 2024. Separately, we sourced historical ITN

distribution data from national malaria control programs (NMCPs) for the period 2000-2023,

primarily via World Health Organization reporting[6]. We then refined the financed 2025 annual

distribution volumes into inferred quarterly volumes and, where relevant, subnational target

regions based on information on the planned timings and locations of mass campaigns available

via the 'ITN Mass Campaign Tracker' resource maintained by the Alliance for Malaria

Prevention[15]. The fraction of ITNs likely to be distributed through mass-campaign distributions

versus routine distribution channels (for example via health facilities, antenatal clinics, schools,

or community health workers) was inferred based on this ratio for each country in historical

data. We then estimated for each country or subnational unit the fraction of all ITNs financed

for 2025 that are financed by PMI. Data on planned volumes of ITNs procured or distributed

by PMI versus other funders in 2025 were extracted from PMI Fiscal Year (FY) 2024 Malaria

Operational Plans (MOPs)[16], together with details on (i) the geographical extent of PMI support

where this was subnational, and (ii) where PMI provides financial support to facilitate

distribution of ITNs purchased by other donors (including in Cameroon and Nigeria). The

resulting inferred PMI-financed proportions were then subtracted from the BAU volumes to

define a 'no PMI' scenario (see appendix table S1). Next, both the BAU and no-PMI scenario

ITN volumes were entered into an established dynamic coverage model[10,11] which transformed

them into monthly estimates of ITN coverage (household access and use) at subnational level

for 2025. The model took into account ITNs already distributed to households in preceding

years and their likely rate of loss. In the no-PMI scenario, we did not explore possible

geographical reallocation of ITNs funded by other means.

Data on the volumes of ACTs financed for distribution in 2025 from all funding sources, and

the fraction supported by PMI, were extracted from the PMI FY2024 MOPs. These documents

also provided delineation, where relevant, of the subnational regions in which PMI provide this

support. The MOPs also detail past years distributions and end-of-year stock-on-hand.

Combining these data elements allowed us to generate inferred subnational ACT distribution

volumes for 2025 under the BAU (i.e. including planned PMI contributions) and no-PMI

(excluding those contributions) scenarios (see appendix figure s3). Stock-on-hand at the outset

of 2025 was considered available for distribution even in the no-PMI scenario, although only

within existing national or subnational units. We then converted these distribution scenarios

into estimated treatment coverages (i.e. the fraction of clinical malaria cases receiving effective

treatment) as follows. For the BAU scenario we used pre-existing coverage estimates

developed by MAP for the annual WHO World Malaria Report[12,6] that consider rates of care

seeking, access to antimalarial drugs, and the time-varying efficacy of different drug classes.

We used the estimates for 2024 as a proxy for 2025, assuming coverage would be similar under

BAU. Treatment coverage under the no-PMI scenario in 2025 was then computed by reducing

public-sector treatment rates in proportion to the fraction of all ACTs financed in 2025 by PMI

per subnational unit. We did not extend this analysis to include the possible effects of reduced

availability of rapid diagnostic tests (RDTs) under the no-PMI scenario. In reality, reduced

access to confirmatory diagnosis would further compromise levels of access to effective

antimalarial treatment by reducing appropriate targeting of limited ACT stock to confirmed

cases. In the no-PMI scenario, we did not explore possible geographical reallocation of ACTs

funded by other means.

Areas with planned SMC campaigns supported by PMI under BAU in 2025 were identified

using data from the FY2024 PMI MOPs, augmented with data reported by NMCPs to WHO[6].

Within those areas, coverage under BAU was assumed to be the same as that estimated for

2024 based on a compilation of post-campaign coverage surveys undertaken by NMCPs and

by Malaria Consortium[17,18]. The timing and number of cycles were assumed to be the same as

in 2024 and the target population was assumed to be strictly children under 59 months. The no-

PMI scenario was then simply defined as having zero SMC coverage in 2025 in the PMI-

supported SMC areas.

The most recent MAP estimates of IRS coverage (for the year 2023) were obtained, which are

based on subnational data on spray campaigns reported by NMCPs to WHO[6], reported by PMI

in national 'End of Spray Reports' (e.g. PMI VectorLink reports[19,20]), or from other country-

specific sources[21]. We used this 2023 coverage to represent our BAU scenario for 2025. Details

of the subset of planned IRS spraying supported by PMI in 2025 were extracted from PMI

FY2024 MOPs[16]. The no-PMI scenario was then defined as having zero IRS coverage in 2025

in those IRS target areas supported by PMI.

For all interventions, the no-PMI scenario was defined in the absence of any mitigation

strategies, such that coverage supported by PMI under BAU was assumed to be lost and not

replenished by other means.

Determining epidemiological impact of PMI-attributable coverage

We propagated the BAU and no-PMI intervention coverage scenarios for ITNs, ACTs, SMC,

and IRS through a series of established models to generate corresponding estimates of infection

prevalence, clinical case incidence, and malaria mortality at 5×5 km resolution and by month across 2025.

First, we fitted a Bayesian geostatistical model to historical data on malaria infection prevalence, intervention coverage, and environmental and socioeconomic covariates from 2000-2023. Prevalence data comprised 59,516 cross-sectional observations of malaria infection prevalence age-standardised to the 2-10 year old age group (i.e. *P. falciparum* parasite rate, $PfPR_{2-10}$), as compiled on a rolling basis by MAP[12] and collected predominantly through cross-sectional surveys such as those conducted by the Demographic and Health Surveys Program (DHS)[22]. Observations were standardised for diagnostic type[23] and age[24] before modelling. Geotemporal intervention coverage estimates from 2000-2023 were derived from the exiting MAP coverage models referenced above and described in detail previously[13]. Geotemporal data on environmental and socio-economic covariates capturing climatic, landcover, and infrastructural variables were also derived from existing MAP catalogues as described previously[13]. The geostatistical modelling framework used here is described extensively elsewhere[5,13]. In brief, the effect on $PfPR_{2-10}$ of: (i) the physical and insecticidal components of ITN coverage, as a function of baseline transmission intensity and trends in pyrethroid resistance; (ii) access to effective antimalarial treatment; and (iii) IRS coverage was determined empirically using a hierarchical Bayesian spatio-temporal model. The model included additional terms controlling for (iv) changing environmental and socioeconomic predictors; (v) underlying baseline endemicity; and (vi) unexplained variation accounted for using a spatio-temporal random field with separable Matérn-5/2-AR(1) covariance structure and independent country-level random effects. For this analysis, we updated the model to explicitly include the effect of insecticide resistance (see appendix section 2.2) and leverage new remote-sensing imagery to better resolve urban-rural gradients of risk. Fitted parameters are shared in the appendix, tables S3 and S4. After five-fold cross-validation, out-of-sample mean squared error was 2·1%, mean absolute error was 9·0%, and correlation between observed and predicted $PfPR_{2-10}$ was 0.80. The model was fit in R v.4.4.1 using INLA v.24.06.27[25,26]. The fitted model, which includes detailed terms characterising the effects of each intervention class, was then used in a final step to predict $PfPR_{2-10}$ at 5×5 km resolution for each month of 2025 across Africa. Two predictions were carried out to reflect intervention coverages under both the BAU and no-PMI scenarios described above. Malaria's transmission dynamics are uncertain and heterogeneous, but suggest the impact of changing intervention

coverage on PfPR$_{2\text{-}10}$ would not be immediate[27], so coverages were lagged by both one and two

months, before pooling to calculate summary statistics. For both 2025 scenarios, environmental

and socioeconomic covariates were fixed at 2023 levels, the most recent full year of available

data.

The BAU and no-PMI PfPR$_{2\text{-}10}$ predictions for 2025 were used to generate corresponding

geotemporal estimates of annual *P. falciparum* clinical incidence rate using an established

natural history model[28] that estimates age-structured clinical incidence as a function of PfPR,

seasonality, and exposure history. Annual estimates were disaggregated to monthly values

using a seasonality model based on compiled observations of clinical malaria seasonality across

Africa[29]. We additionally accounted for the impact on clinical incidence of SMC using our

modelled SMC coverage combined with observations of SMC effectiveness under routine

implementation pooled across sites in seven West African countries[30] (see appendix section

2.3.1). Malaria mortality in 2025 under the BAU and no-PMI scenarios was determined by first

combining the geotemporal estimates of clinical cases and (unlagged) treatment rates to derive

untreated cases. We then applied to those untreated case estimates an untreated case fatality

rate derived from the World Malaria Report[31], including upper and lower bound estimates. Note

that this approach captures two mechanisms by which antimalarial treatment averts mortality:

via direct impact on PfPR$_{2\text{-}10}$, which reduces clinical incidence rates, and then additionally by

reducing the fraction of clinical cases that progress to death.

Role of the funding source

The funder of the study had no role in study design, data collection, data analysis, data

interpretation, or writing of the report. The corresponding author had full access to all the data

in the study and had final responsibility for the decision to submit for publication.

Results

Intervention coverage

Our compiled data on planned procurement and distribution indicated that under business-as-

usual PMI would provide 44 million ITNs in 2025. An additional 15 million ITNs funded by

the Against Malaria Foundation for Nigeria and Cameroon are reliant on PMI logistical support

for distribution[32,33]. In total, these volumes represent 35% of the 169M ITNs planned for

distribution in 2025 across the 25 African PMI focus countries included in this study. We estimated that these ITNs would contribute 10.1 percentage points (pp) to population coverage (i.e. use – the percentage of people sleeping under an ITN) by the end of 2025 across the PMI focus countries. Impact varies considerably by country with effects exceeding 20 pp in Sierra Leone, Nigeria (PMI focus states), Burundi, Zimbabwe, and Cameroon (Figure 2A). Data on planned ACT procurement and distribution indicated that PMI would finance 71 million doses of the drug across its African focus regions in 2025 under business-as-usual, representing 52% of total ACT procurement in those regions. We estimated that these drugs would contribute 12·0 pp to the fraction of clinical malaria cases receiving effective treatment. Some countries are proportionally more reliant on PMI-funded ACTs than others, with PMI contributing over 20 pp to effective treatment rates in Zambia, Senegal, Burundi, and PMI focus provinces of Angola (Figure 2B). Compiled data indicates that under business-as-usual PMI would contribute 12·3 pp to under-five population coverage of SMC across all areas targeted for this intervention in Africa in 2025. In specific subnational geographies where PMI would concentrate SMC support, this figure is often greater than 75 pp (Figure 2C). PMI was scheduled to be a relatively minor contributor to IRS campaigns in Africa in 2025, with a business-as-usual contribution of 4·9 pp to annualised coverage across regions targeted continent-wide (Figure 2D).

Malaria cases and deaths

We estimate that PMI procurement and distribution of malaria control commodities, as planned under business-as-usual operations for 2025, would avert 13·6M (95% uncertainty interval 11·4M – 16·4M) clinical malaria cases and save 104,000 (69,000 – 161,000) lives across that year. (Table 1, Figure 3A,B). Burden averted by month is slightly greater in the latter half of the year, driven in part by the availability of pre-existing stock-on-hand for ACTs which reduce the impact of the no-PMI scenario in the first few months of 2025. In addition, the timing of planned PMI-funded ITN and SMC campaigns, which typically occur in quarters three and four for Sahelian countries due to seasonal variations in transmission intensity, means greater impact in these later months. Our decomposition by intervention type (Figure 3B,C) indicated that PMI-funded ACT provision formed the largest contribution to both cases (55%, 95% uncertainty interval 48 – 67%) and deaths (84% [82 – 85%]) averted, followed by ITNs (25% of cases [20 – 31%], 7% of deaths [6 – 8%]) and SMC (20% of cases [11 – 23%], 9% of deaths [8 –10%]). At national-level, we estimate business-as-usual PMI activities would save the

largest number of lives in PMI focus regions of Nigeria (20,000; 14,000 - 30,000), Burundi
(11,000; 7,100 - 18,000) and Democratic Republic of the Congo (11,000; 6,200 - 20,000),
corresponding to densely-populated high-burden countries with extensive PMI support (Table
1).

Discussion

At a time of ongoing policy reorientation, it is essential to make available to decision makers
reliable measures of the impact of investments in global health. Here we demonstrate that a
fully functioning PMI would likely avert nearly 14M malaria cases and around 100,000 malaria
deaths in Africa in 2025. These estimates do not rely on simple extrapolations of lives-saved-
per-dollar but are instead built up from a detailed reconstruction of planned procurement and
distribution activities for the major classes of malaria control commodity, the proportional
geographically-specific contribution of PMI to that commodity provision, the way those
commodities drive population coverage, and the way that coverage impacts malaria
transmission and burden. This granularity allows the necessary sophistication to capture the
differential effects of each intervention class, to take into account control interventions already
in place or available for distribution at the onset of the year, and to acknowledge heterogeneity
in underlying malaria risk across the continent and how this influences intervention impact.

The geospatial methodology used in this analysis does not explicitly account for immune
rebound and associated hysteresis in intervention impact. Rebound to above pre-intervention
burden levels of burden – due to the suppression of naturally acquired immunity – has been
observed in settings where sustained high-frequency chemoprophylaxis has been abruptly
removed[34], but the frequency and magnitude of this effect relative to resurgence to pre-
intervention levels is poorly understood[35,36,37]. A review of fifty studies suggested immune
rebound occurs infrequently (8/50 studies) under trial conditions[35]. Under real-world
conditions intervention coverage fluctuates (appendix Figures S5, S6), predominantly due to
cyclic vector control campaigns[11], leading to periods of reduced protection and increased
exposure. The extent to which the no-PMI scenario – a partial reduction of imperfect coverage
–could lead to a rebound effect is therefore unclear, but in this respect our morbidity and
mortality estimates may be conservative.

The results presented here must be interpreted in the context of several limitations. First, while
we focus on PMI-funded commodity procurement, we do not explicitly account for PMI
support to logistics and supply chain management, instead assuming that changes in coverage
stem only from reduced input commodity volumes. Second, we do not account for PMI support
for RDTs. In practice, RDT stock-outs would result in reversion to presumptive malaria
diagnosis which would translate into a higher fraction of ACTs being prescribed to non-malaria
cases. This would not only impair appropriate management of non-malaria causes but would
place further strain on ACT stocks. Reduced confirmatory diagnosis also means surveillance
data would decline in quality – impairing future monitoring of malaria trends. Third, when
estimating mortality effects, we take into account only changes to overall case incidence and
to rates of first-line treatment of presenting uncomplicated clinical cases, implicitly assuming
the case-fatality rate among untreated cases remains constant. This neglects any contribution
of PMI to supporting quality management of severe malaria cases - for example via
procurement of injectable artesunate. Fourth, we do not account for any present or planned
PMI support for the implementation of vaccination with the RTS,S/AS01 or R21/Matrix-M
vaccines. Fifth, we do not account for healthcare personnel directly supported by PMI. Most
importantly, this includes around 100,000 community health workers[7] whose efforts have
demonstrably improved access to timely malaria case management in rural and remote
communities in many PMI focus countries[38]. Sixth, we do not account for a range of additional
activities supported by PMI that include strengthening of disease and entomological
surveillance systems, and monitoring of insecticide and drug resistance. Seventh, we do not
capture the indirect effects of increased malaria burden on health systems, and its potential to
reduce capacity for care of other conditions.
374
It is important to note that nearly all the limitations listed above imply our estimates are
conservative with respect to the number of cases and deaths averted. Conversely, we do not
consider possible mitigation strategies in a scenario without PMI support – instead focusing on
PMI impact *ceteris paribus*. This likely neglects the resilience and increasing self-sufficiency
of African health agencies in many countries. In reality, a future with more constrained
international funding for malaria control would catalyse a range of mitigation and response
strategies by national government and health agencies. These could include emergency
domestic funding mechanisms, amplifying the existing trend towards greater domestic funding
for malaria control[6]. Limited control commodities could also be reprogrammed - either
geographically or across different intervention mixes – to maximise disease control within a

given resource envelope. However, the degree of readiness and the time required for these
structural responses to be enacted is likely to vary considerably across PMI focus countries.
Furthermore, given that PMI contributes nearly a quarter of the total $3.0 billion USD of
malaria funding spent in the WHO AFRO region[6,7], rapidly mitigating the absence of PMI at
the necessary scale would require significant reallocation of resources from other budgets. A
default scenario without substantial mediation remains, therefore, an appropriate benchmark to
consider for the forthcoming year. Compared to this scenario, our analysis indicates that PMI
investment in supporting procurement and distribution of malaria control commodities would
translate directly into millions of malaria cases averted and a hundred thousand lives saved in
its 27 countries of operation in Africa across 2025.
395

Contributors
PWG and TLS conceptualised the study, led data analysis and wrote the first draft of the
manuscript. JL, MM, AS, MV, HB, YB, SH, RJ, PA, AB, EC, CV, SFR, NG, DJW contributed
to data curation and compilation and contributed to data analysis. All authors had full access to
the data, contributed to review and approved the final version of the manuscript. The
corresponding author had full access to all the data in the study and had final responsibility for
the decision to submit for publication.
403

404 **Data sharing**
405 Data on commodity funding and distribution, malaria prevalence and intervention coverage
406 used in this analysis are available in the public domain via the cited sources. All PMI-focus-
407 geography level results are available within the main text and supplementary material. Other
408 summarisations are available upon request to the authors.
409

410 **Declaration of interests**
411 All authors declare no competing interests.
412

Acknowledgments
This work is supported by the National Health and Medical Research Council, Australia
(2025280).

Figures and Tables

Table 1. Estimated mean malaria cases and deaths averted in Africa by PMI in 2025, with 95%

uncertainty intervals. National level malaria case estimates are rounded to three significant

figures, deaths estimates to two. Africa-wide estimates are rounded to the nearly 100,000

(cases) or 1,000 (deaths). DRC = Democratic Republic of the Congo. Africa estimate

corresponds to the sum across all PMI focus geographies.

Country	Malaria cases averted	Malaria deaths averted
Angola	165,000 (117,000 - 221,000)	1,800 (1,000 - 3,200)
Burundi	1,040,000 (766,000 - 1,360,000)	11,000 (7,100 - 18,000)
Benin	251,000 (217,000 - 287,000)	1,200 (840 - 1,800)
Burkina Faso	888,000 (754,000 - 1,040,000)	7,200 (4,800 - 11,000)
Cote d'Ivoire	459,000 (344,000 - 599,000)	2,600 (1,800 - 3,900)
Cameroon (PMI focus)	1,510,000 (1,350,000 - 1,650,000)	4,400 (3,300 - 5,900)
DRC (PMI focus)	1,260,000 (983,000 - 1,610,000)	11,000 (6,200 - 20,000)
Ethiopia	190,000 (150,000 - 233,000)	400 (120 - 930)
Ghana	248,000 (219,000 - 273,000)	590 (500 - 700)
Guinea	200,000 (173,000 - 225,000)	450 (340 - 600)
Kenya	148,000 (120,000 - 177,000)	1,100 (950 - 1,400)
Liberia	120,000 (88,700 - 158,000)	1,200 (740 - 1,900)
Madagascar	36,000 (26,400 - 47,900)	220 (69 - 470)
Mali	515,000 (430,000 - 618,000)	2,800 (2,000 - 3,900)
Mozambique	419,000 (276,000 - 594,000)	4,900 (2,700 - 9,300)
Malawi	312,000 (221,000 - 428,000)	2,500 (1,800 - 3,600)
Niger	770,000 (563,000 - 1,060,000)	7,900 (5,200 - 12,000)
Nigeria (PMI focus)	2,850,000 (2,490,000 - 3,280,000)	20,000 (14,000 - 30,000)
Rwanda	98,100 (71,500 - 127,000)	690 (510 - 920)
Senegal	946,000 (676,000 - 1,250,000)	7,700 (2,700 - 17,000)
Sierra Leone	171,000 (140,000 - 201,000)	1,200 (700 - 1,900)
Togo	5,350 (4,140 - 6,460)	9 (6 - 13)
Tanzania	582,000 (417,000 - 767,000)	9,300 (7,400 - 12,000)
Uganda	107,000 (91,100 - 125,000)	140 (100 - 230)
Zambia	298,000 (218,000 - 402,000)	3,300 (2,500 - 4,600)
Zimbabwe	6,940 (4,820 - 9,310)	20 (6 - 47)
Africa (PMI focus)	**13.6M (11.4M – 16.4M)**	**104,000 (69,000 – 161,000)**

Figure 1. Schematic summary of methodological framework. Blue rectangles represent input datasets, some of which are modelled results from preceding analyses. Orange rectangles represent modelling steps. Green rectangles are the results of the analysis. ITN=insecticide-treated net. IRS = Indoor Residual Spraying, SMC = Seasonal Malaria Chemoprevention. *Pf*PR=*Plasmodium falciparum* parasite rate. S1 BAU = business-as-usual scenario. S2 noPMI = scenario with PMI-funded interventions removed.

A. Insecticide treated nets

0-5 | 10-15 | 20-25 | 30-35 | >40
5-10 | 15-20 | 25-30 | 35-40

Impact on ITN coverage (percentage point)

B. Antimalarial treatment

0-5 | 10-15 | 20-25 | 30-35 | >40
5-10 | 15-20 | 25-30 | 35-40

Impact on treatment rate (percentage point)

C. Seasonal malaria chemoprevention

SMC supported by PMI
SMC supported by other funders

D. Indoor residual spraying

IRS supported by PMI
IRS supported by other funders

Figure 2. Projected contributions of PMI in 2025 under business-as-usual (BAU) to each
intervention class considered in this study. (A) Percentage-point difference in access to
insecticide treated bednets between BAU and no-PMI scenarios, projected in December 2025.
Hashed areas correspond to regions of PMI focus countries where PMI does not procure ITNs
nor support their distribution. Inset shows BAU and no-PMI scenario projected ITNs in

households, January-December 2025. (B) Percentage-point difference in the proportion of

clinical malaria cases receiving effective treatment between BAU and no-PMI scenarios,

projected in December 2025. Hashed areas correspond to regions of PMI focus countries where

PMI does not procure ACTs. Inset shows BAU and no-PMI scenario projected effective

treatment rates of clinical malaria cases, January-December 2025. (C) Spatial extent of

Seasonal Malaria Chemoprevention projected in 2025, classified by funding support (PMI vs

other funders). Inset shows BAU and no-PMI scenario projected numbers of infants (under

five) protected by SMC, January-December 2025. (D) Spatial extent of IRS projected in 2025,

classified by funding support (PMI vs other funders). Inset shows BAU and no-PMI scenario

projected area (in km^2) protected by IRS, January-December 2025. SMC = Seasonal Malaria

Chemoprevention; BAU = business-as-usual; PMI = President's Malaria Initiative.

Figure 3. Estimates of malaria burden averted by PMI in Africa in 2025 with 95% uncertainty

intervals; (A) Number of *Plasmodium falciparum* clinical cases (in millions), by month, with

blue representing business-as-usual projection and red with PMI-funded interventions

removed; (B) the number of *P. falciparum* deaths, by month, with colour scheme as in (A).

Plotted monthly estimates are off-set in the no-PMI scenario for presentation and ease of

interpretation; (C) Number of *P. falciparum* clinical cases (in millions) averted by PMI-funded

interventions, by month of 2025, disaggregated by intervention class. (D) Number of *P.*

falciparum deaths averted by PMI-funded interventions, by month of 2025, disaggregated by

intervention class. Vector control consists of both coverage with Insecticide Treated Bednets

and Indoor Residual Spraying; SMC = Seasonal Malaria Chemoprevention.

462   References

1. Barat LM, Whitehurst N, Venkatesan M, *et al*. The US President's Malaria Initiative's support for improving the quality of malaria case management services: fifteen years of progress and learning. The American Journal of Tropical Medicine and Hygiene. 2023 Nov 27;110(3 Suppl):1.

2. Jakubowski A, Stearns SC, Kruk ME, Angeles G, Thirumurthy H. The US President's Malaria Initiative and under-5 child mortality in sub-Saharan Africa: A difference-in-differences analysis. PLoS medicine. 2017 Jun 13;14(6):e1002319.

3. Ye Y, Duah D. The President's Malaria Initiative contributed to reducing malaria burden in sub-Saharan Africa between 2004 and 2014: Evidence from generalized estimating equation analysis. PLoS one. 2019 May 24;14(5):e0217103.

4. Winskill P, Slater HC, Griffin JT, Ghani AC, Walker PG. The US President's Malaria Initiative, *Plasmodium falciparum* transmission and mortality: A modelling study. PLoS Medicine. 2017 Nov 21;14(11):e1002448.

5. Bhatt S, Weiss DJ, Cameron E, *et al*. The effect of malaria control on *Plasmodium falciparum* in Africa between 2000 and 2015. Nature. 2015 Oct 8;526(7572):207-11.

6. World Health Organization. World Malaria Report 2024: addressing inequity in the global malaria response. Geneva: World Health Organization; 2024. Available from: https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2024

7. President's Malaria Initiative. 18th Annual Report to Congress. Washington, DC: President's Malaria Initiative; 2024. Available from: https://stacks.cdc.gov/view/cdc/154755

8. White MT, Conteh L, Cibulskis R, Ghani AC. Costs and cost-effectiveness of malaria control interventions – a systematic review. Malar J. 2011;10:337.

9. Penny MA, Verity R, Bever CA, *et al*. Public health impact and cost-effectiveness of the RTS,S/AS01 malaria vaccine: a systematic comparison of predictions from four mathematical models. Lancet. 2016 Jan 23;387(10016):367-375. doi: 10.1016/S0140-6736(15)00725-4.

10. Bhatt S, Weiss DJ, Mappin B, *et al*. Coverage and system efficiencies of insecticide-treated nets in Africa from 2000 to 2017. Elife. 2015 Dec 29;4:e09672.

11. Bertozzi-Villa A, Bever CA, Koenker H, *et al*. Maps and metrics of insecticide-treated net access, use, and nets-per-capita in Africa from 2000-2020. Nature Communications. 2021 Jun 11;12(1):3589.

12. The Malaria Atlas Project. Datasets available from: https://data.malariaatlas.org. Last accessed 23/03/2025.

13. Weiss DJ, Dzianach PA, Saddler A, *et al*. Mapping the global prevalence, incidence, and mortality of *Plasmodium falciparum* and *Plasmodium vivax*, 2000–2022. Lancet, Volume 405, Issue 10483, 979—990.

14. Roll Back Malaria. Programmatic and Financial Gap Analysis(v17/12/2024). Available from: https://dashboards. endmalaria.org/en/gap-analysis. Last accessed 14/02/2025.

15. Alliance for Malaria Prevention. AMP ITN Mass Campaign Tracker. Available from: https://allianceformalariaprevention.com/itn-dashboards/mass-campaign-tracker. Last accessed 14/02/2025.

16. President's Malaria Initiative Malaria Operational Plans. Available at: http://pmi.gov/resources/

17. Nnaji C. Coverage and quality of seasonal malaria chemoprevention supported by Malaria Consortium with philanthropic funding or co-funding in 2023: results from Burkina Faso, Chad, Mozambique, Nigeria, South Sudan, Togo, and Uganda. Malaria Consortium; 2024. Available from: https://www.malariaconsortium.org/resources/publications/1774/coverage-and-quality-of-seasonal-malaria-chemoprevention-supported-by-malaria-consortium-in-2023. Last accessed 27/02/2025.

18. Loua KM, Milligan P. Seasonal malaria chemoprevention in Guinea in 2020: coverage survey results. Technical Report. London: London School of Hygiene & Tropical Medicine, Universite Gamal Abdel Nasser; 2021. Available from: https://doi.org/10.17037/PUBS.04663123.

19. The PMI VectorLink Project. August 2023. Ethiopia 2023 End of Spray Report. Spray Campaign: May 2, 2023 – July 7, 2023. Rockville, MD: The PMI VectorLink Project, Abt Associates Inc

20. The PMI VectorLink Ghana Project. May 2023. Ghana End of Spray Report. Spray Campaign: March 1– April 13, 2023. Rockville, MD, USA: The PMI VectorLink Project, Abt Associates.

21. AngloGold Ashanti (Ghana) Malaria Control (AGAMal). (2023). Annual report 2023. AGAMal. Retrieved from https://www.agamal.org

22. ICF. The DHS Program. Available from: https://dhsprogram.com.

23. Mappin B, Cameron E, Dalrymple U*, et al.* Standardizing *Plasmodium falciparum* infection prevalence measured via microscopy versus rapid diagnostic test. Malaria journal. 2015 Dec;14:1-0.

24. Smith DL, Guerra CA, Snow RW, Hay SI. Standardizing estimates of the *Plasmodium falciparum* parasite rate. Malaria journal. 2007 Dec;6:1-0.

25. Rue H, Martino S, Chopin N. Approximate Bayesian inference for latent Gaussian models using integrated nested Laplace approximations (with discussion). Journal of the Royal Statistical Society, Series B, 71(2):319-392, 2009.

26. Lindgren F, Rue H, Lindström J. An explicit link between Gaussian fields and Gaussian Markov random fields: the stochastic partial differential equation approach. Journal of the Royal Statistical Society Series B: Statistical Methodology. 2011 Sep;73(4):423-98.

27. Huber JH, Johnston GL, Greenhouse B, Smith DL, Perkins TA. Quantitative, model-based estimates of variability in the generation and serial intervals of Plasmodium falciparum malaria. Malaria journal. 2016 Dec;15:1-2.

28. Cameron E, Battle KE, Bhatt S, *et al*. Defining the relationship between infection prevalence and clinical incidence of *Plasmodium falciparum* malaria. Nature communications. 2015 Sep 8;6(1):8170.

29. Nguyen M, Howes RE, Lucas TCD, *et al*. Mapping malaria seasonality in Madagascar using health facility data. BMC Med. 2020 Feb 10;18(1):26. doi: 10.1186/s12916-019-1486-3. PMID: 32036785; PMCID: PMC7008536.

30. Baba E, Ndiaye JL, Uaene R, *et al*. Effectiveness of seasonal malaria chemoprevention at scale in West and Central Africa: an observational study. Lancet. 2020;396(10265):1829-1840.

31. World Health Organization. World malaria report 2023. Geneva: World Health Organization; 2023. Available from: https://www.who.int/publications/i/item/9789240078121.

32. U.S. President's Malaria Initiative. Cameroon Malaria Operational Plan FY2024. Washington, DC: U.S. Agency for International Development; 2024.

33. U.S. President's Malaria Initiative. Nigeria Malaria Operational Plan FY2024. Washington, DC: U.S. Agency for International Development; 2024.

34. Greenwood BM, David PH, Otoo-Forbes LN, Allen SJ, Alonso PL, Schellenberg JA, Byass P, Hurwitz M, Menon A, Snow RW. Mortality and morbidity from malaria after stopping malaria chemoprophylaxis. Transactions of the Royal Society of Tropical Medicine and Hygiene. 1995 Nov 1;89(6):629-33.

35. Technical consultation on the malaria rebound phenomenon: report on a virtual meeting, 22-23 March 2022. Geneva: World Health Organisation; 2022.

36. Cohen, J.M., Smith, D.L., Cotter, C. *et al.* Malaria resurgence: a systematic review and assessment of its causes. *Malar J* **11**, 122 (2012). https://doi.org/10.1186/1475-2875-11-122

37. Greenwood, B., Zongo, I., Dicko, A. *et al.* Resurgent and delayed malaria. *Malar J* **21**, 77 (2022). https://doi.org/10.1186/s12936-022-04098-6

38. Ashton RA, Hamainza B, Lungu C, *et al*. Effectiveness of community case management of malaria on severe malaria and inpatient malaria deaths in Zambia: a dose–response study using routine health information system data. Malar J 22, 96 (2023). https://doi.org/10.1186/s12936-023-04525-2