African-specific genetic loci determine iron status and risk of severe malaria and bacteremia in African children

John Muthii Muriuki[1], Alexander J Mentzer[2,3], Gavin Band[2], Amanda Y Chong[2], Alex W Macharia[1], Reagan M Mogire[1], Kelvin Mokaya Abuga[1], Ruth Mitchell[4], James J Gilchrist[5], Emily L Webb[6], Francis M Ndungu[1], Laura M Raffield[7], Lynette Ekunwe[8], Sodiomon B Sirima[9], Shabir A Madhi[10], Adrian V S Hill[2,11], Andrew Prentice[12], Philip Bejon[1,13], Gibran Hemani[4], George Davey Smith[4], Manjinder S Sandhu[14], Alison M Elliott[6,15], Thomas N Williams[1,16,17], Adebowale Adeyemo[18], and Sarah H Atkinson[1,16,19]

Authors' affiliations
[1]Kenya Medical Research Institute (KEMRI), Centre for Geographical Medicine Research – Coast, KEMRI-Wellcome Trust Research Programme, Kilifi, Kenya
[2]Centre for Human Genetics, Nuffield Department of Medicine, University of Oxford, Oxford, UK
[3]Big Data Institute, Li Ka Shing Centre for Health Information and Discovery, University of Oxford, Oxford, UK
[4]Medical Research Council (MRC) Integrative Epidemiology Unit, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK
[5]Oxford Vaccine Group, Department of Paediatrics, University of Oxford, Oxford, UK
[6]MRC Tropical Epidemiology Group, Department of Infectious Disease Epidemiology, London School of Hygiene and Tropical Medicine, London, UK
[7]Department of Genetics, University of North Carolina, Chapel Hill, North Carolina, USA
[8]Department of Medicine, University of MS Medical Center, Jackson, USA
[9]Groupe de Recherche Action en Sante (GRAS), 06 BP 10248, Ouagadougou 06, Burkina Faso
[10]South African Medical Research Council: Vaccines and Infectious Diseases Analytical Research Unit, Faculty of Health Sciences, University of the Witwatersrand, Johannesburg, South Africa
[11]Centre for Clinical Vaccinology and Tropical Medicine and the Jenner Institute Laboratories, University of Oxford, Oxford, UK
[12]MRC Unit The Gambia at London School of Hygiene and Tropical Medicine, Banjul, The Gambia
[13]Modernising Medical Microbiology Unit, Nuffield Department of Medicine, University of Oxford, Oxford, UK
[14]Wellcome Sanger Institute, Hinxton, Cambridge, UK
[15]MRC/Uganda Virus Research Institute and London School of Hygiene & Tropical Medicine Uganda Research Unit, Entebbe, Uganda
[16]Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of Oxford, Oxford, UK
[17]Department of Medicine, Imperial College, London, UK
[18]Centre for Research on Genomics and Global Health, National Human Genome Research Institute, National Institutes of Health, South Drive, MSC 5635, Bethesda, Maryland 20891-5635, USA
[19]Department of Paediatrics, University of Oxford, Oxford, UK

Correspondence to: John Muthii Muriuki; jmuriuki@kemri-wellcome.org and Sarah H Atkinson; satkinson@kemri-wellcome.org / sarah.atkinson@paediatrics.ox.ac.uk

Abstract

Iron is an essential trace element for both humans and pathogens, but its genetic determinants are understudied in Africa where iron deficiency and infectious diseases are highly prevalent. We conducted genome-wide association studies for six iron-related biomarkers in 3928 children from five sites across Africa with replication in 2868 African American adults. We identify a previously unreported loci for transferrin in *GTF3C5*, a gene regulating cellular iron-uptake, for soluble transferrin receptor in *FREM3*, the Dantu locus, and for hepcidin in *CHCHD7/SDR16C5*. The lead *GTF3C5* and *FREM3* variants, rs2905094, and rs141274959, were associated with protection against severe malaria and bacteremia in large case-control studies. The *CHCHD7/SDR16C5* lead variant, rs73596248, increased hepcidin concentrations and protected against bacteremia. We report limited transferability of polygenic risk scores derived from European studies to African populations. Our findings advance the understanding of the genetics of iron status in Africa and suggest an important link between iron and infection.

Introduction

Iron is an essential nutrient that is required for critical biological functions in almost all living organisms. Iron deficiency affects over half of young African children,[1] and is a leading cause of years lived with disability.[2] Iron status influences susceptibility to infection since pathogens require iron to proliferate and iron is essential for host immunity.[3,4] This is important in the context of Africa due to the high infectious disease burden. In 2023, there were an estimated 263 million cases of malaria, and 597,000 deaths globally with 95% of these occurring in the WHO African region.[5] Similarly, age standardised mortality rates due to invasive bacterial infections were highest in sub-Saharan Africa with an estimated 230 deaths per 100 000 population in 2019.[6]

Iron homeostasis is tightly regulated in humans since both iron deficiency and overload increase the risk of infection.[7] Transferrin, the main protein that carries iron in circulation, is upregulated in the acute phase response to sequester iron away from pathogens, contributing to 'nutritional immunity'[8] while cellular iron uptake through transferrin receptor 1 is critical for the development of host immunity.[9] Hepcidin regulates systemic iron homeostasis[10] and influences iron acquisition by malaria parasites and bacteria.[11,12] Iron plays a crucial role in the relationship between host and pathogen and the high infectious disease burden in Africa might have created different evolutionary pressures on iron-regulating genes.

The genetic architecture of iron status in African populations remains largely understudied, in contrast to the many large-scale genome-wide association studies (GWAS) of iron status conducted in European ancestry populations.[13,14] Studies of the genetics of iron status in Africa remain largely limited to candidate variant studies[15,16] and to our knowledge, no iron GWAS has been undertaken in continental African populations. A GWAS of iron status was conducted in 2347 middle-aged African-Americans,[17] however transferability to continental African populations might be complicated by admixture, limited representation of the greater genetic diversity found among continental Africans,[18] and profoundly different environmental exposures in terms of nutrition and infectious disease.[19,20] Even within Africa, there exist divergent patterns in LD which might yield region-specific variants.[21] Given the greater genetic variation in Africans,[18] an African iron GWAS is likely to reveal genetic variants that are specific to this population.

To enhance discovery of genetic determinants of iron status in an understudied ancestral population, we conducted a GWAS of six iron biomarkers (transferrin, hepcidin, soluble transferrin receptor (sTfR), ferritin, serum iron, and transferrin saturation (TSAT)) in 3,928 children from Kenya, Uganda, Burkina Faso, The Gambia and South Africa with replication in 2,868 African American adults. We identify three previously unreported African-specific genome-wide significant loci that are associated with iron homeostasis and risk of severe malaria and invasive bacterial infections.

Results

Characteristics of discovery sample

To assess the genetic contribution to iron status in continental Africans, we performed a discovery GWAS analysis using data from five sites in Africa (Fig. 1a).[22] A total of 3928 children (1059 Kenyan, 1360 Ugandan, 348 Burkinabe, 611 South African, and 550 Gambian), aged between birth and eight years were included in this study. The characteristics of the 3928 children are summarized in Supplementary Table 1. Prevalence of iron deficiency, and geometric mean concentrations of measured iron biomarkers (transferrin, ferritin, sTfR, hepcidin, serum iron, and TSAT) varied by study site (Fig. 1b). Inflammation, malaria parasitemia and undernutrition were highly prevalent among study participants (Supplementary Table 1). Principal component plots of the genotypes showed clustering of the study participants by geography and ethnolinguistic group (Fig. 1c).

A summary of quality control steps of genotypes for each population is shown in Supplementary Table 2. The number of variants imputed into individual datasets are shown in Supplementary Table 3. Only high-quality SNPs (imputation information score >0.3) with a minor allele frequency (MAF) >0.01 were included in genetic association analyses.

Discovery genetic association analyses

To identify variants associated with iron biomarkers, we performed individual GWAS within each of the African discovery sites (Supplementary Fig. 1) followed by meta-analysis. Fig. 2 shows the distribution of association statistics for the GWAS of the six iron biomarkers after fixed-effects meta-analyses of all five African sites. We applied a linear mixed model including a genetic relatedness matrix (GRM) as a random effect and age and sex as covariates and found minimal population stratification or inflation of the association statistics (lambda ranged from 0.98 to 1.05, Supplementary Fig. 2). Additional adjustment for inflammatory markers yielded similar results. Meta-analysis of GWAS results from all the African cohorts yielded a previously reported genome-wide significant locus for transferrin on chromosome 3 at the *TF* gene (Fig. 2a) and a previously unreported African-specific locus associated with hepcidin levels (Fig. 2b). We did not observe any loci at genome-wide significance (set at $P<5x10^{-8}$) after meta-analysis of all GWAS results for ferritin, sTfR, serum iron, and TSAT from all African study sites (Figs. 2c, 2d, 2e and 2f).

The *TF* locus is well characterized across ancestries. The lead SNP in our African transferrin GWAS was rs6762719 ($P=1.59x10^{-27}$) which is in strong LD ($r^2=0.9$) with the lead common SNP (rs3811647) associated with transferrin levels in European transferrin GWAS.[23] The rs3811647 SNP ($P= 3.45x10^{-16}$) was among the top SNPs in our African transferrin GWAS. After conditional analyses adjusting for rs6762719, we found another independent locus (lead SNP rs4854748) indicating at least two independent loci for transferrin (Table 1) as previously reported in African-Americans.[17] This SNP (rs4854748) was in strong LD ($r^2=0.9$) with the lead SNP, rs9872999, in the conditional analysis in African-Americans.[17] We also

found that rs4854748 was strongly associated with reduced total iron binding capacity (TIBC), a proxy for transferrin levels, in the largest European-ancestry TIBC GWAS (P=3.05x10^{-578}).[14] Genome-wide significant SNPs associated with transferrin levels in our discovery meta-analysis, replicated in the Jackson Heart Study (JHS) and the largest European-ancestry TIBC GWAS[14] (except rs113656013 which is rare in Europeans (minor allele frequency = 0.001 compared to 0.17 in Africans); Supplementary Table 4).

GTF3C5, an East African-specific locus for transferrin

Since genetic associations can be highly population specific across Africa, we performed meta-analysis by African region (i.e., East, West and South Africa; Supplementary Fig. 3). For transferrin levels, we observed a previously unreported East African (Kenya and Uganda) specific locus on chromosome 9 (Fig. 3a). Three SNPs reached genome-wide significance in East Africa only (Supplementary Table 5). The lead SNP, rs2905094, was in high LD with the other two SNPs (Fig. 3b) and explained 1.42% of variation in transferrin levels. The T allele frequency of rs2905094 in East Africa was 16% with allele frequencies of 14-23% across the African sites and in AfricanAmericans in the JHS, and of 33% in Europeans (Fig. 3c). The rs2905094 SNP was also associated with increased levels of ferritin, TSAT and serum iron, decreased levels of sTfR (Fig. 3d), and a 19% reduction in iron deficiency (OR=0.81, (95% CI 0.68, 0.97); P=0.02) in East Africa.

We then used external reference data to annotate and evaluate the functional or regulatory impact of genome-wide significant variants and nearby gene(s). The three genome-wide significant SNPs (Supplementary Table 5) mapped to the *GTF3C5* (general transcription factor IIIC, polypeptide 5) gene and were all intronic. Using the African Functional Genomics Resources (AFGR, https://github.com/smontgomlab/AFGR), rs2905094 was a significant eQTL for *GTF3C5* (P=0.04) in immortalized lymphoblastoid cell lines. *GTF3C5* is involved in RNA polymerase III-mediated transcription. It is an integral and tightly associated component of the DNA-binding transcription factor IIIC subcomplex that directly binds tRNA and virus-associated RNA promoters. *GTF3C5* is expressed ubiquitously, with a relatively higher expression in the cerebellum. Using a multiparametric profiling of the endocytic system, ablation of genes including *GTF3C5* in HeLa cells was shown to increase endocytosis and internalization of transferrin.[24]

Given the above observations, we next performed haplotype analyses to investigate why the *GTF3C5* locus association was East African-specific. We found two dominant haplotypes from chromosomes with the rs2905094 mutant (T) allele (Supplementary Fig. 4). Haplotype 1 accounted for approximately 70% and 60% of chromosomes in The Gambia and Burkina Faso, respectively and less than 20% in South Africa, Uganda and Kenya. Haplotype 2 accounted for 40%, 25% and 15% of chromosomes from Kenya, Uganda, and South Africa, respectively, but was absent in The Gambia and Burkina Faso. Given these population differences, we hypothesized that haplotype 2 might be responsible for the East-African specific transferrin signal. Consistent with this hypothesis, we found that rs2905094 had the largest effect size in Kenya, followed by Uganda, but was not associated with transferrin levels in South Africa, Burkina Faso, and The Gambia (Fig. 3c). This suggested that rs2905094 might be in LD with an unidentified causal SNP within the East-African specific haplotype 2.

Since transferrin levels influence risk of malaria and bacterial infections,[25,26] we investigated the association between rs2905094 and these infections using large case-control GWAS of severe malaria (7957 cases and 7746 controls),[27] and bacteremia (1536 cases and 2677

controls).[28,29] We compared fixed effects meta-analyses of severe malaria GWAS data from East Africa (i.e. Kenya, Malawi, and Tanzania) and West Africa (The Gambia, Mali, Burkina Faso, Ghana, Nigeria, and Cameroon; Supplementary Fig. 5) since the effect of rs2905094 on transferrin levels was restricted to East Africa. We found that rs2905094 was associated with 12% (OR=0.88, (95% CI 0.78, 0.98) protection against severe malaria in East Africa where it had an impact on transferrin levels but was not associated with protection in West Africa where it had no impact on transferrin levels (OR=0.95; 95% CI 0.87, 1.03 (Fig. 3f and Supplementary Fig. 5). The rs2905094 was associated with 15% (OR=0.85, (95% CI 0.75, 0.97; P=0.02) protection against overall bacteremia in Kenyan children (Fig. 3f).

CHCHD7/SDR16C5, an African-specific locus for hepcidin

Meta-analysis of hepcidin GWAS results from all five African sites yielded a previously unreported locus at genome-wide significance, found on chromosome 8 near the *CHCHD7* (Coiled-Coil-Helix-Coiled-Coil-Helix Domain Containing 7) and *SDR16C5* (Short Chain Dehydrogenase/Reductase Family 16C Member 5) genes (Fig. 2b). The regional plot suggests that this association is within an extended haplotype (Fig. 4a). This region shows evidence of positive selection among two 1000G African populations (Luhyas (Kenya) and Yorubas (Nigeria)); iHS = 1.5, XP-EHH =2.5, Fay & Wu's H =-7.6, and Tajima's D =-2.8) (https://pophumanscan.uab.cat/tables.php). The A allele of the lead SNP, rs73596248 occurred at a frequency of 6% in Africans with consistently increased hepcidin levels across all the African sites (Fig. 4b). rs73596248 explained 0.82% of variation in hepcidin levels. In contrast, the rs73596248 A allele was very rare in European and Latin American populations and was monomorphic in Asian populations (Supplementary Table 6). rs73596248 was associated with a 21% (OR=0.79, (95% CI 0.64, 0.97); P=0.03) protection against iron deficiency, but was not significantly associated with individual iron biomarkers (Supplementary Fig. 6a).

Based on available and limited evidence, the rs73596248 is downstream of *CHCHD7* and is an intergenic variant between the *CHCHD7* and *SDR16C5* genes. rs73596248 is a significant eQTL for *CHCHD7* (P=1.50x10^{-4}) in the AFGR. *CHCHD7* is a protein coding gene predicted to be in the mitochondrial intermembrane space and is overexpressed in fibroblasts. The *SDR16C5* gene encodes short chain dehydrogenase reductase which oxidizes retinol to retinoic acid and is overexpressed in the lung. Retinoic acid has been reported to suppress production of hepcidin.[30] Although the *CHCHD7/SDR16C5* locus is strongly associated with height in Open Targets (https://genetics.opentargets.org/), we did not find an association between rs73596248 and anthropometric indices including height (Supplementary Fig. 7).

Since hepcidin is implicated in the pathophysiology of malaria and bacterial infections,[11,12,31] we determined the association between rs73596248 and these infections in large case-control GWAS of severe malaria and bacteremia mentioned above. We found that rs73596248 was associated with protection against bacteremia due to *Klebsiella pneumoniae* and *Staphylococcus aureus*. Protection against overall bacteremia and severe malaria did not reach statistical significance (Fig. 4c).

FREM3, a Kenyan-specific locus for soluble transferrin receptor

We observed a genome-wide significant locus at the *FREM3* (FRAS1 Related Extracellular Matrix 3) gene that was associated with an increase in sTfR levels in Kenya (Fig. 5a). *FREM3* is within the complex Dantu region that is under strong selective pressure for protection against malaria along the Kenyan coast.[32,33] The regional plot suggests an extended haplotype underlying the association (Fig. 5b). The top SNPs were rare or monomorphic

outside Kenya (Supplementary Table 7). The G allele of the lead SNP, rs552439837, occurred at a frequency of 9% in Kenya (Table 1) and was also associated with an increase in TSAT and serum iron concentrations (Supplementary Fig. 6b). rs552439837 explained 3.77% of variation in sTfR levels. The causal malaria related Dantu SNP, rs186873296,[33] was in strong LD ($r^2>0.9$) with rs552439837 and was also associated with increased sTfR levels (effect in standard deviation=0.34, se=0.07, P= 1.3×10^{-6}, G allele frequency=11.3%). Owing to the Kenyan-specific nature of the Dantu region, we did not find any reported eQTL data for the top sTfR GWAS SNPs in GTex or AFGR databases.

The lead SNP, rs552439837 was missing in the large case-control GWAS of severe malaria and bacteremia. We therefore identified the closest available SNP, rs141274959, from among the genome-wide significant sTfR GWAS SNPs (Supplementary Table 7). rs141274959 is in strong LD with rs552439837 ($r^2=0.9$) and was highly protective against severe malaria and bacteremia (Fig. 5c).

Transferability of established European iron GWAS SNPs

To investigate the transferability of European iron GWAS findings to Africans, we identified the estimates of index SNPs that were associated with iron biomarkers in the largest GWAS of European-ancestry populations[13,14] in our meta-analyzed GWAS results of African children (Supplementary Table 8 and 9). We considered a GWAS result to be replicated if the estimate in the African iron GWAS was in the same direction as in the estimate in the European iron GWAS, and if the association in African iron GWAS was statistically significant (P< 0.05). Most of the variant allele frequencies differed substantially between European ancestry and continental African populations (Supplementary Table 8 and 9). The *TF* locus replicated while the other loci had limited replication. Notably, the *HFE* variants (rs1800562 and rs1799945) could not be replicated since the associated SNPs were very rare or monomorphic in Africans. The *TMPRSS6* variant, rs855791, replicated for TSAT and had similar effect size for the other iron biomarkers in Africans and Europeans (Supplementary Table 8 and 9). However, the effect allele (A) frequency of rs855791 is lower in African (7.7%) compared to European (43.1%) populations. In summary, of the 138 index SNPs available in our African iron GWAS, 69 (25 for ferritin, 20 transferrin, 9 serum iron, 6 hepcidin, 6 sTfR, and 4 TSAT) had the same direction of effect as in the estimate in the European iron GWAS but only the *TF* locus had a statistically significant P value (Supplementary Table 8 and 9). We then calculated polygenic risk scores (PRS) for each iron biomarker using the European iron biomarkers GWAS index SNPs and similarly found that only the transferrin PRS replicated in the current study (Fig. 6).

Discussion

African populations are substantially under-represented in human genomic studies.[34,35] In this study, we report the first GWAS of iron status in continental African populations. We replicated a previously reported signal at the *TF* gene that was associated with transferrin levels.[17,23] We further discovered three African-specific genetic loci that influence iron status in African children. We identified a previously unreported East African specific locus, *GTF3C5,* implicated in intracellular iron uptake, which was associated with reduced transferrin levels and protection against malaria and bacteremia. We also found a previously unreported locus within the *CHCHD7/SDR16C5* genes, associated with increased hepcidin levels and protection against bacteremia. The malaria protective *FREM3* SNP within the Dantu region,[33] was associated with increased sTfR levels in Kenyan children and protection against bacteremia. These region-specific findings underscore the existence of divergent

patterns in LD among African populations.[21] We further found limited transferability of findings from European iron GWAS to African populations. Africa's diverse lifestyle, cultural and environmental factors such as long-term burden of infections might have shaped the genomic architecture of its people.

Transferrin, synthesised in the liver, is the major iron-binding and transport protein that delivers iron to cells. It is highly conserved in humans and plays an important role in infection since it is upregulated in the acute phase response to sequester iron away from pathogens, contributing to 'nutritional immunity'.[8] Our transferrin GWAS analyses yielded three independent loci: two at the *TF* gene and one novel locus at the *GTF3C5* gene. The two independent loci at the *TF* gene have previously been reported and are well characterised across ancestries.[14,23,36,37] The *GTF3C5* locus was associated with reduced transferrin levels and is specific to East African populations. *GTF3C5* encodes transcription factor IIIC63 (TFIIIC63) which binds to DNA to recruit transcription factor IIIB and RNA polymerase III to mediate the transcription of small noncoding RNAs, such as tRNAs. *GTF3C5* has been implicated in transferrin endocytosis.[24] Through a phenotypic profiling of genes involved in transferrin endocytosis by genome-wide RNAi screening using a quantitative multiparametric image analysis, *GTF3C5* gene knockdown caused increased endocytosis and internalization of transferrin.[24] We observed SNPs at *GTF3C5* that were associated with a decrease in plasma transferrin levels suggesting that the SNPs might cause an increase in endocytosis and internalization of transferrin similar to the *GTF3C5* gene knockdown. We also found that the lead SNP, rs2905094 was associated with an increase in levels of ferritin, serum iron, and TSAT and a decrease in sTfR levels, in keeping with improved iron status. Indeed, we found that rs2905094 was associated with protection against iron deficiency suggesting that the mutation might play a role in iron absorption.

The *GTF3C5* locus appears to influence iron status in East Africa only despite the effect allele occurring at a similar frequency in other African ancestries and even at a higher frequency (33%) in Europeans. This finding might be explained by multiple factors. Owing to the distinct demographic histories of populations from different geographic regions in Africa, this locus might be characterized by distinct haplotype block structures. We found that East-African populations displayed a unique haplotype suggesting that an unknown causal SNP might be in LD with our index SNPs in East African populations but not in other populations. Gene environment interactions could also be at play since we found that the locus was associated with severe malaria in East Africa but was not associated with protection in West Africa. This population specific finding further underscores the high diversity of African genomes and existence of divergent patterns in LD among African populations.[18,21]

The *GTF3C5* locus was further associated with protection against severe malaria and bacteremia suggesting a link between iron metabolism and infection. We hypothesize three possible mechanisms. Firstly, an increase in intracellular iron within immune cells or improved iron status might play a role in improved immunity thereby protecting against infection.[4,38] Using a transgenic mouse model carrying a mutation in the transferrin receptor (thus limiting the ability of cells to internalise iron), Wideman et al reported impaired immune response and increased malaria parasitemia.[38] It is therefore possible that increased internalised iron improves immunity against pathogens. Secondly, by increasing endocytosis of transferrin-bound iron,[24] excess of iron within cells is potentially toxic to intracellular pathogens.[39] Indeed, we found that the mutation was associated with protection against severe *P. falciparum* an intracellular pathogen. Thirdly, since the malaria parasite can acquire iron

bound to transferrin,[26] a reduction in transferrin might deny the parasite of iron for growth and multiplication. Further research is required to test these hypotheses.

We found an African specific previously unreported locus between the *CHCHD7* and *SDR16C5* genes that was associated with increased hepcidin levels in populations across Africa. SNPs within this locus occurred at a frequency of ~6% in Africans but were very rare or monomorphic in non-African populations. Concordantly, we found evidence that the locus is under positive selective pressure probably to protect against pathogens. Hepcidin is the master regulator of systemic iron homeostasis and plays a crucial role in the body's response to infection by inducing 'nutritional immunity'.[10,31] The lead SNP, rs73596248, was associated with increased hepcidin levels and protection against *K. pneumoniae* and *S. aureus* bacteremia. This finding is also in agreement with our previous report that severely anemic children are unable to reduce hepcidin in response to infection and have increased risk of bacteremia.[40]

We found that the malaria-protective Dantu region at the *FREM3* locus was associated with increased sTfR levels and protection against bacteremia in Kenyan children. This locus is unique to the Kenyan population since SNPs were rare or monomorphic elsewhere. *FREM3* is in linkage disequilibrium with a complex structural mutation within the glycophorin gene region (comprising GYPA, GYPB, and GYPE) that encodes for the rare Dantu blood group antigen and is associated with protection against severe malaria in East Africa.[33] In our analyses, the locus was associated with protection against both severe malaria and bacteremia. The protection against bacteremia is not surprising since malaria infection strongly predisposes individuals to bacteremia in malaria-endemic areas.[41] Recent analyses from the same population showed that the Dantu polymorphism, rs186873296, is protective against bacteremia via its effect on malaria risk.[42] sTfR is a marker of erythropoiesis or younger RBCs and when increased it indicates iron deficiency. In an observational study, we previously showed that iron deficiency was protective against malaria infection.[43] This finding might also suggest that Dantu RBCs are younger than non-Dantu RBCs as previously shown.[44] However, further research is required to elucidate the complex interplay between iron and malaria infection.

We further found limited transferability of European iron GWAS findings to continental African populations. Using index variants identified in the largest iron GWAS in European populations,[13,14] we were only able to formally replicate other prior reported loci in the *TF* locus, in African children. Notably, the *HFE* polymorphisms were very rare or monomorphic in Africans and it is well known that the hemochromatosis-causing variants are not observed in African populations. The *TMPRSS6* SNP, rs855791, although not as common in African compared to European populations (allele frequency of 7.7% in Africans versus 43.1% in Europeans), replicated in terms of effect size but not significance (P>0.05) except for TSAT. Considering the low frequency of rs855791 in Africans, it is likely that we were underpowered to observe a statistically significant association. The *HFE* and *TMPRSS6* polymorphisms are often used in Mendelian randomization (MR) studies involving European populations,[45,46] but would have limited application in African MR and functional studies. All the other loci reported in European iron GWAS did not replicate (in terms of both direction of effect and P values) in our study populations. Similarly, only the transferrin PRS replicated in our African iron GWAS. However, our sample size was small compared to the large sample sizes reported in European GWAS. The European GWAS were also conducted in adults while the current study involved children, and regulation of iron homeostasis could differ across the life course. Moreover, most variants had very different effect allele frequencies

between African and European populations which might further explain the limited transferability and a need for African genomic studies.

The strengths of the present study include a focus on diverse understudied continental African ancestral groups and inclusion of a wide range of iron biomarkers. To our knowledge, this study is the first GWAS of iron biomarkers within continental Africa. We have identified previously unreported African-specific loci and further determined their impact of on infections using the largest case-control studies of severe malaria and bacteremia. A potential limitation is that variants with small effect sizes were not detectable with the present sample size. To overcome this limitation, there is a need for larger genomic studies of iron status within continental Africa.

In conclusion, we found African-specific genetic loci that are associated with iron status in African children. Malaria and bacteremia may have shaped the genetic architecture of iron status in Africans. However, mechanistic studies using the identified loci are required to infer the causal relationship between iron and pathogens. Finally, we found limited transferability of European iron GWAS to Africans. These findings further underscore the importance of conducting genomic studies in Africans to better address specific medical needs and global health inequalities.

References

1. Muriuki, J. M. *et al.* Estimating the burden of iron deficiency among African children. *BMC Med* **18**, 31 (2020).
2. GBD 2016 Disease and Injury Incidence and Prevalence Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 328 diseases and injuries for 195 countries, 1990-2016: A systematic analysis for the Global Burden of Disease Study 2016. *The Lancet* **390**, 1211–1259 (2017).
3. Ganz, T. & Nemeth, E. Iron homeostasis in host defence and inflammation. *Nat Rev Immunol* **15**, 500–10 (2015).
4. Frost, J. N. *et al.* Hepcidin-Mediated Hypoferremia Disrupts Immune Responses to Vaccination and Infection. *Med* **2**, 164–179 (2021).
5. World Health Organization. *World Malaria Report 2024*. (2024).
6. GBD 2019 Antimicrobial Resistance Collaborators. Global mortality associated with 33 bacterial pathogens in 2019: a systematic analysis for the Global Burden of Disease Study 2019. *The Lancet* **400**, 2221–2248 (2022).
7. Mottelson, M. *et al.* Iron, hemochromatosis genotypes, and risk of infections: a cohort study of 142188 general population individuals. *Blood* **144**, (2024).
8. Barber, M. F. & Elde, N. C. Escape from bacterial iron piracy through rapid evolution of transferrin. *Science (1979)* **346**, 1362–1366 (2014).
9. Jabara, H. H. *et al.* A missense mutation in TFRC, encoding transferrin receptor 1, causes combined immunodeficiency. *Nat Genet* **48**, 74–78 (2016).
10. Nemeth, E. *et al.* Hepcidin regulates cellular iron efflux by binding to ferroportin and inducing its internalization. *Science (1979)* **306**, 2090–2093 (2004).
11. Stefanova, D. *et al.* Hepcidin protects against lethal Escherichia coli sepsis in mice inoculated with isolates from septic patients. *Infect Immun* **86**, e00253-18 (2018).
12. Portugal, S. *et al.* Host-mediated regulation of superinfection in malaria. *Nat Med* **17**, 732–737 (2011).
13. Allara, E. *et al.* Novel loci and biomedical consequences of iron homoeostasis variation. *Commun Biol* **7**, 1631 (2024).

14.   Moksnes, M. R. *et al.* Genome-wide meta-analysis of iron status biomarkers and the effect of iron on all-cause mortality in HUNT. *Commun Biol* **5**, 591 (2022).

15.   Gichohi-Wainaina, W. N. *et al.* Associations between common variants in iron-related genes with haematological traits in populations of African ancestry. *PLoS One* **11**, e0157996 (2016).

16.   Jallow, M. W., Campino, S., Prentice, A. M. & Cerami, C. Association of common TMPRSS6 and TF gene variants with hepcidin and iron status in healthy rural Gambians. *Sci Rep* **11**, 8075 (2021).

17.   Li, J. *et al.* Genome-wide admixture and association study of serum iron, ferritin, transferrin saturation and total iron binding capacity in African Americans. *Hum Mol Genet* **24**, 572–581 (2015).

18.   Choudhury, A. *et al.* High-depth African genomes inform human migration and health. *Nature* **586**, 741–748 (2020).

19.   Parra, E. J. *et al.* Estimating African American admixture proportions by use of population- specific alleles. *Am J Hum Genet* **63**, 1839–1851 (1998).

20.   Tishkoff, S. A. *et al.* The Genetic Structure and History of Africans and African Americans. *Science (1979)* **324**, 1035 LP – 1044 (2009).

21.   Tarazona-Santos, E. & Tishkoff, S. A. Divergent patterns of linkage disequilibrium and haplotype structure across global populations at the interleukin-13 (IL13) locus. *Genes Immun* **6**, 53–65 (2005).

22.   Mentzer, A. J. *et al.* High-resolution African HLA resource uncovers HLA-DRB1 expression effects underlying vaccine response. *Nat Med* **30**, 1384–1394 (2024).

23.   Benyamin, B. *et al.* Variants in TF and HFE explain approximately 40% of genetic variation in serum-transferrin levels. *Am J Hum Genet* **84**, 60–65 (2009).

24.   Collinet, C. *et al.* Systems survey of endocytosis by multiparametric image analysis. *Nature* **464**, 243–249 (2010).

25.   Cross, J. H. *et al.* Oral iron acutely elevates bacterial growth in human serum. *Sci Rep* **5**, 16670 (2015).

26.   Clark, M. A., Goheen, M. M. & Cerami, C. Influence of host iron status on Plasmodium falciparum infection. *Front Pharmacol* **5**, 84 (2014).

27.   Malaria Genomic Epidemiology Network. Insights into malaria susceptibility using genome-wide data on 17,000 individuals from Africa, Asia and Oceania. *Nat Commun* **10**, 5732 (2019).

28.   Gilchrist, J. J. *et al.* Risk of nontyphoidal Salmonella bacteraemia in African children is modified by STAT4. *Nat Commun* **9**, 1014 (2018).

29.   Gilchrist, J. J. *et al.* BIRC6 modifies risk of invasive bacterial infection in Kenyan children. *Elife* **2022**, e77461 (2022).

30.   Han, L., Liu, Y., Lu, M., Wang, H. & Tang, F. Retinoic acid modulates iron metabolism imbalance in anemia of inflammation induced by LPS via reversely regulating hepcidin and ferroportin expression. *Biochem Biophys Res Commun* **507**, 280–285 (2018).

31.   Stefanova, D. *et al.* Endogenous hepcidin and its agonist mediate resistance to selected infections by clearing non–transferrin-bound iron. *Blood* **130**, 245–257 (2017).

32.   Leffler, E. M. *et al.* Resistance to malaria through structural variation of red blood cell invasion receptors. *Science (1979)* **356**, 1140–1152 (2017).

33.   Ndila, C. M. *et al.* Human candidate gene polymorphisms and risk of severe malaria in children in Kilifi, Kenya: a case-control association study. *Lancet Haematol* **5**, e333–e345 (2018).

34.   Popejoy, A. B. & Fullerton, S. M. Genomics is failing on diversity. *Nature* vol. 538 161–164 Preprint at https://doi.org/10.1038/538161a (2016).

35.   Fatumo, S. *et al.* A roadmap to increase diversity in genomic studies. *Nat Med* **28**, 243–250 (2022).

36.   Li, J. *et al.* Genome-wide admixture and association study of serum iron , ferritin , transferrin saturation and total iron binding capacity in African Americans. **24**, (2015).

37.   McLaren, C. E. *et al.* Associations between single nucleotide polymorphisms in iron-related genes and iron status in multiethnic populations. *PLoS One* **7**, (2012).

38.   Wideman, S. K. *et al.* Cellular iron governs the host response to malaria. *PLoS Pathog* **19**, (2023).

39.   Toh, S. Q., Glanfield, A., Gobert, G. N. & Jones, M. K. Heme and blood-feeding parasites: Friends or foes? *Parasit Vectors* **3**, 108 (2010).

40.   Abuga, K. M. *et al.* Hepcidin regulation in Kenyan children with severe malaria and non-typhoidal Salmonella bacteremia. *Haematologica* **107**, 1589–1598 (2022).

41.   Scott, J. A. G. *et al.* Relation between falciparum malaria and bacteraemia in Kenyan children: A population-based, case-control study and a longitudinal study. *The Lancet* **378**, 1316–1323 (2011).

42.   Kariuki, S. N. *et al.* Relation Between the Dantu Blood Group Variant and Bacteremia in Kenyan Children: A Population-Based Case-Control Study. *J Infect Dis* 10.1093/infdis/jiae339 (2024) doi:10.1093/infdis/jiae339.

43.   Muriuki, J. M. *et al.* Iron status and associated malaria risk among African children. *Clinical Infectious Diseases* **68**, 1807–1814 (2019).

44.   Kariuki, S. N. *et al.* Red blood cell tension protects against severe malaria in the Dantu blood group. *Nature* **585**, 579–583 (2020).

45.   Daghlas, I. & Gill, D. Genetically predicted iron status and life expectancy. *Clinical Nutrition* **40**, 2456–2459 (2021).

46.   Gill, D. *et al.* The Effect of Iron Status on Risk of Coronary Artery Disease: A Mendelian Randomization Study - Brief Report. *Arterioscler Thromb Vasc Biol* **37**, 1788–1792 (2017).

47.   Snow, R. W. *et al.* The prevalence of Plasmodium falciparum in sub-Saharan Africa since 1900. *Nature* **550**, 515–518 (2017).

48.   World Health Organization. *WHO Guideline on Use of Ferritin Concentrations to Assess Iron Status in Individuals and Populations.* https://apps.who.int/iris/bitstream/handle/10665/331505/9789240000124-eng.pdf?sequence=1&isAllowed=y (2020).

49.   Bejon, P. *et al.* Stable and Unstable Malaria Hotspots in Longitudinal Cohort Studies in Kenya. *PLoS Med* **7**, e1000304 (2010).

50.   Webb, E. L. *et al.* Effect of single-dose anthelmintic treatment during pregnancy on an infant's response to immunisation and on susceptibility to infectious diseases in infancy: A randomised, double-blind, placebo-controlled trial. *The Lancet* **377**, 52–62 (2011).

51.   Bliss, C. M. *et al.* Viral Vector Malaria Vaccines Induce High-Level T Cell and Antibody Responses in West African Children and Infants. *Molecular Therapy* **25**, 547–559 (2017).

52.   Nunes, M. C. *et al.* Duration of infant protection against influenza illness conferred by maternal immunization. *JAMA Pediatr* **170**, 840 (2016).

53.   Atkinson, S. H. *et al.* Seasonal childhood anaemia in West Africa is associated with the haptoglobin 2-2 genotype. *PLoS Med* **3**, e172 (2006).

54.   Taylor, H. The Jackson Heart Study: an overview. *Ethn Dis* **15**, S6-1–3 (2005).

55.   Wilson, J. *et al.* Study design for genetic analysis in the Jackson Heart Study. *Ethn Dis* **15**, S6-30–37 (2005).

56.   Wray, K. *et al.* Hepcidin detects iron deficiency in Sri Lankan adolescents with a high burden of hemoglobinopathy: A diagnostic test accuracy study. *Am J Hematol* **92**, 196–203 (2017).

57.   Yamanishi, H., Iyama, S., Yamaguchi, Y., Kanakura, Y. & Iwatani, Y. Total iron-binding capacity calculated from serum transferrin concentration or serum iron concentration and unsaturated iron-binding capacity. *Clin Chem* **49**, 175–178 (2003).

58.   Baichoo, S. *et al.* Developing reproducible bioinformatics analysis workflows for heterogeneous computing environments to support African genomics. *BMC Bioinformatics* **19**, 457 (2018).

59.   Brandenburg, J. T. *et al.* H3AGWAS: a portable workflow for genome wide association studies. *BMC Bioinformatics* **23**, 498 (2022).

60.   Haplotype-Reference-Consortium. A reference panel of 64,976 haplotypes for genotype imputation. *Nat Genet* **48**, 1279–1283 (2016).

61.   Yang, J., Lee, S. H., Goddard, M. E. & Visscher, P. M. GCTA: A tool for genome-wide complex trait analysis. *Am J Hum Genet* **88**, 76–82 (2011).

62.   Han, B. & Eskin, E. Random-effects model aimed at discovering associations in meta-analysis of genome-wide association studies. *Am J Hum Genet* **88**, 586–598 (2011).

63.   Choi, S. W., Mak, T. S. H. & O'Reilly, P. F. Tutorial: a guide to performing polygenic risk score analyses. *Nat Protoc* **15**, 2759–2772 (2020).

64.   Visscher, P. M. *et al.* 10 Years of GWAS Discovery: Biology, Function, and Translation. *The American Journal of Human Genetics* **101**, 5–22 (2017).

65.   World Health Organization. *Iron Deficiency Anaemia: Assessment, Prevention, and Control. A Guide for Programme Managers. Https://Www.Who.Int/Nutrition/Publications/Mic.* (2001).

66.   Camaschella, C. Iron-Deficiency Anemia. *N Engl J Med* **372**, 1832–1843 (2015).

Table 1. Top independent SNPs associated with iron biomarkers in African children

SNP	Chr	Hg19 BP	EA	OA	Gene	EA Freq	Beta	Se	P
Transferrin (all African sites)*									
rs6762719	3	133480817	G	A	*TF*	0.25	0.28	0.03	1.59×10^{-27}
rs4854748	3	133446881	T	C	*TF*	0.20	-0.24	0.03	1.12×10^{-16}
Transferrin (Uganda and Kenya)									
rs2905094	9	135902689	T	C	*GTF3C5*	0.16	-0.23	0.04	8.08×10^{-9}
Hepcidin (all African sites)									
rs73596248	8	57134187	A	G	*CHCHD7/ SDR16C5*	0.06	0.27	0.05	4.51×10^{-8}
sTfR (Kenya)									
rs552439837	4	144446525	G	A	*FREM3*	0.09	0.48	0.09	2.78×10^{-8}

Chr, chromosome; Hg19 BP, human genome build 19 base pair; EA, effect allele; OA, other allele; EA Freq, effect allele frequency; Se, standard error; P, two-sided P value from an additive model.
*R^2 between rs6762719 and rs4854748 was 0.03 in the African populations.

Study	N	ID (%)	Transferrin (g/L)	Ferritin (µg/L)	sTfR (mg/L)	Hepcidin (µg/L)	Iron (µmol/L)	TSAT (%)
Discovery studies (African sites)								
1 Kilifi, Kenya	1059	35.0	2.7 (1.3)	21.9 (3.1)	17.8 (1.5)	5.7 (3.6)	6.5 (2.0)	9.4 (2.2)
2 Entebbe, Uganda	1360	34.8	2.7 (1.3)	20.9 (2.9)	6.7 (2.0)	6.8 (3.3)	n/a	n/a
3 Soweto, South Africa	611	41.1	2.6 (1.3)	14.9 (2.6)	10.8 (1.5)	7.8 (3.5)	n/a	n/a
4 Banfora, Burkina Faso	348	35.5	2.7 (1.3)	22.2 (2.9)	17.7 (1.7)	5.3 (4.2)	6.0 (1.8)	8.9 (2.0)
5 West Kiang, The Gambia	550	21.5	2.6 (1.2)	24.4 (2.4)	3.5 (1.4)	5.5 (4.7)	8.6 (1.6)	12.8 (1.7)
Replication site								
6 Jackson Heart Study, USA	2868	10.4	2.0 (1.2)	112.2 (2.6)	n/a	n/a	74.1 (1.5)	25.7 (1.5)

Fig 1. Characteristics of study populations. a) Africa malaria map showing the location of the discovery sites included in the current study and predicted posterior predictions of age-standardized *P. falciparum* prevalence (PfPR$_{2-10}$) taken from Snow *et al* 2017.[47] **b)** Prevalence of iron deficiency (ID) and geometric means of iron markers by study site. Iron deficiency was defined as serum ferritin <12 µg/L with no inflammation, or <30 µg/L in the presence of inflammation (C-reactive protein>5mg/L or α1-antichymotrypsin >0.6g/dL) in children <5 years or <15µg/L with no inflammation or <70 µg/L in the presence of inflammation in children ≥5 years)[48] **c)** Principal components plot of genotypes of the discovery study participants. sTfR, soluble transferrin receptor; TSAT, transferrin saturation; PC (Principal Components); n/a, not available.

Fig 2. Manhattan plots of meta-analysis results of discovery samples. Meta-analysis included all five African sites except for serum iron and TSAT which were missing in Ugandan and South African samples. Genome-wide significant signals are annotated *TF*, for transferrin and *CHCHD7/SDR16C5* for hepcidin. Genetic variants were tested for association with iron biomarkers per study site using linear mixed models and evidence was combined across sites using inverse-variance weighted meta-analysis in a fixed-effects framework. The red line indicates threshold for genome-wide significance (set at $5\text{x}10^{-8}$).

Fig 3. ***GTF3C5*** **is a previously unreported locus associated with transferrin levels, severe malaria and bacteremia in East Africa. a)** Meta-analysis of transferrin GWAS results in East Africa (Kenya and Uganda). **b)** A regional plot of the *GTF3C5* locus in East Africa. The most strongly associated variant, rs2905094 (P=8.08x10[-9]), is represented by a purple diamond. Additional SNPs are colored by their correlation (r^2) with rs2905094. Location and direction of transcription of protein-coding genes within the region are shown. **c)** A forest plot by study site showing the minor allele frequency (MAF), beta in standard deviation, and additive model two-tailed GWAS p-values of the association between rs2905094 and transferrin levels. **d)** Association between rs2905094 and other iron biomarkers in Kenyan and Ugandan children. Linear regression models were adjusted for age, sex, and study site. Estimates were derived from an additive model. **e)** A forest plot showing odds ratios of rs2905094 for risk of severe malaria, overall bacteremia, and common bacterial isolates in East African children.

Fig 4. A previously unreported locus associated with hepcidin levels in African poulations a) A regional plot for the locus. The most strongly associated variant, rs73596248 (P=4.51x10^{-8}), is represented by a purple diamond. Additional SNPs are colored by their correlation (r^2) with rs73596248. Location and direction of transcription of protein-coding genes within the region are shown. **b)** A forest plot by study site showing the minor allele frequency (MAF), beta in standard deviation, and additive model two-tailed GWAS p-values of the association between rs73596248 and hepcidin levels. Hepcidin data was not available in the Jackson Heart Study so replication in that population was not tested. **c)** A forest plot showing odds ratio of rs73596248 and severe malaria, overall bacteremia, and common bacterial isolates.

Fig 5. Genome-wide association study of soluble transferrin receptor in Kenya. a) Manhattan plot of GWAS of soluble transferrin receptor (sTfR) in Kenya. **b)** A regional plot showing correlation between the lead SNP, rs552439837 (purple diamond) and SNPs within the region. **c)** A forest plot showing odds ratio of rs141274959 and severe malaria, overall bacteremia, and common bacterial isolates. rs141274959 had an r^2=0.9 with rs552439837 which was missing in the malaria and bacteremia GWAS.

Fig 6. Transferability of polygenic risk scores (PRS) derived from European iron GWAS lead SNPs to continental African populations. Association between European PRS and iron biomarkers in African children. Regression models were adjusted for age, sex, and study site. For each iron biomarker, we calculated a PRS weighted using effect sizes reported in European iron biomarker GWAS and reflecting an increase in levels of the biomarker (included SNPs are shown in Supplementary Table 8 and 9). In our African populations, we used transferrin as a proxy measure for total iron binding capacity (TIBC) that was reported in the European GWAS. TIBC indicates the amount of iron that can be bound to transferrin. TSAT=transferrin saturation; sTfR, soluble transferrin receptor.

Methods

Ethical Approvals
Individual study site ethical approvals were obtained. For Kilifi, Kenya (by the Scientific Ethics Review Unit of the Kenya Medical Research Institute (KEMRI/SERU/CGMR-C/046/3257/2983)); Entebbe, Uganda (locally by the Uganda Virus Research Institute (GC/127/12/07/32) and Uganda National Council for Science and Technology (MV625), and in the UK by the London School of Hygiene and Tropical Medicine (A340) and Oxford Tropical Research (OTR) (39-12, 42-14 and 37-15) Ethics Committees); Banfora, Burkina Faso (by Ministere de la Recherche Scientifique et de l'Innovation in Burkina Faso (2014-12-151) and the OTR Ethics Committees (41-12)); Soweto, South Africa (by the University of Witwatersrand Human Research (M130714) and the OTR Ethics Committees (1042-13 and 42-14)); and West Kiang, The Gambia (by the Gambian Government / Medical Research Council Ethics Committee (874/830)). For JHS, relevant data transfer agreements were signed.

Study populations
Our discovery studies included five community-based cohorts of children from sub-Saharan Africa in Kenya, Uganda, Burkina Faso, South Africa, and The Gambia. Our replication population included African Americans from the Jackson Heart Study (JHS) in Mississippi, USA. Below is a description of each of the study sites.

Discovery cohorts
Kilifi, Kenya: This included an ongoing rolling cohort evaluating malaria immunity in children in Kilifi, along the coast of Kenya.[49] Within this cohort, children were followed-up for malaria episodes up to 8 years of age with weekly follow-ups and annual cross-sectional surveys during which anthropometry measurements, auxiliary temperature, and blood samples were taken. Iron and inflammatory markers were measured from blood samples collected at a single cross-sectional survey based on the availability of plasma samples archived at -80^0C.

Entebbe, Uganda: The *Entebbe Mother and Baby Study (EMaBS)* is a prospective birth cohort that was originally designed as a randomized controlled trial to test whether anthelminthic treatment during pregnancy and early childhood was associated with differential response to vaccination or incidence of infections including malaria (http://emabs.lshtm.ac.uk/).[50] Pregnant women from Entebbe and in their second or third trimester of pregnancy were initially enrolled at Entebbe Hospital antenatal clinic. Livebirths from these mothers were enrolled into the EMaBS study. Blood samples were collected in vacutainer tubes containing ethylenediaminetetraacetic acid (EDTA) at birth and at subsequent birthdays up to five years of age. Iron and inflammatory biomarkers were measured from a single annual visit based on the availability of stored samples.

Banfora, Burkina Faso: The *VAC050 ME-TRAP Malaria Vaccine Trial* recruited infants between the ages of 6 and 18 months living in the Banfora region of Burkina Faso to participate in a Phase 1/2b clinical trial of the safety, immunogenicity and efficacy of a viral-vectored prime-boost liver-stage malaria vaccine.[51] Samples from a total of 350 infants were then selected based on suitability of samples for DNA extraction. Plasma samples were available from the infants at multiple time-points following receipt of the experimental vaccine. Samples from individuals taken at time points as close to the 12-month age as possible were prioritized for this study.

South Africa: The Soweto Vaccine Response Study consisted of infants born in Chris Hani Baragwanath Hospital living in the Soweto region of Johannesburg, South Africa who were recruited from vaccine trials[52] that are coordinated by the Respiratory and Meningeal Pathogens Unit (http://www.rmpru.com/). Mothers of the infants were approached if their infants had received all EPI vaccines up to six months of age. The infants were sampled prospectively at six months of age and at 12 months after receipt of measles vaccine at nine months. Single whole blood samples were collected in EDTA vacutainer tubes for measurement of iron and inflammatory markers and DNA extraction.

The Gambia: West Kiang study: All children aged two to six years (except those with chronic illness) were recruited from 10 rural villages in the West Kiang region of The Gambia during the start of a malaria season to initially investigate seasonal genetic effects of iron status, hemoglobin and haptoglobin concentrations between July and August 2001.[53] All children had a clinical examination, anthropometric measurements, and a three-day course of mebendazole for potential hookworm infection. A blood sample was collected for measurement of iron and inflammatory markers and DNA extraction.

Replication cohort
Jackson Heart Study: This is a population-based longitudinal study of African-American adults aged ≥21 years living in the Jackson, Mississippi metropolitan area in the USA.[54] This study was designed to evaluate risks of cardiovascular disease as described elsewhere.[54,55] Iron and inflammatory biomarkers were measured from blood samples collected at a single clinic visit. Whole blood was used to extract DNA using Puregene reagents (Gentra System, Minneapolis, USA).

Infection GWAS
MalariaGEN consortium: This was a GWAS study of severe malaria in 17,000 severe malaria cases and population controls from 11 countries.[27] For the East-Africa specific *GTF3C5* lead SNP, rs2905094, we retrieved population-specific results from all the nine African countries in the MalariaGEN database and applied fixed-effects meta-analyses by East (Kenya, Malawi, and Tanzania) and West (The Gambia, Mali, Burkina Faso, Ghana, Nigeria, and Cameroon) African sites. For the association between the hepcidin lead SNP, rs73596248 and severe malaria, we meta-analyzed all the nine African countries in the MalariaGEN database. We used the MalariaGen Kenyan GWAS to determine the association between the Kenyan specific sTfR GWAS top SNPs and severe malaria.

The Kenyan Bacteremia GWAS: This study comprised of 1536 hospitalized Kenyan children with positive blood cultures and 2677 matched community controls as previously described.[28,29] To determine associations between our iron-related conditionally independent SNPs and bacteremia, we identified the SNPs in the bacteremia GWAS and extracted estimates, standard errors, and p-values.

Assays for iron and inflammatory markers
The assayed biomarkers of iron (ferritin, soluble transferrin receptors (sTfR), hepcidin, serum iron, transferrin, and unsaturated iron binding capacity (UIBC) and inflammation (C-reactive protein (CRP) and α_1-antichymotrypsin (ACT) are shown in Supplementary Table 10 The Gambian hepcidin values were harmonized with the rest of the cohorts' hepcidin values (measured using high sensitive DRG kit) by converting to the old DRG hepcidin assay values and then to the new high sensitive DRG hepcidin assay values.[56] Transferrin saturation

(TSAT) was calculated as (serum iron (µmol/L)/transferrin (g/L) x 25.1) x 100 or as (serum iron/ UIBC + serum iron) x 100.[57] Since EDTA chelates iron, serum iron measurements (and TSAT) were missing in Uganda and South Africa where blood samples were collected in EDTA vacutainer tubes. Transferrin (g/L) in JHS was calculated as 0.7 x total iron binding capacity (TIBC) [µg/dL] x 0.01. Transferrin in The Gambia was similarly calculated but after calculating TIBC as UIBC + serum iron.

Genotyping and imputation
Genotyping for DNA extracted from Uganda, Burkina Faso and South Africa was performed using the Illumina HumanOmni 2.5M-8 ('octo') BeadChip array version 1.1 (Illumina Inc., San Diego, USA), at the Genotyping Core facilities at the Wellcome Trust Sanger Institute. Genotyping for Kenyan and Gambian samples was performed using H3Africa Custom Genotyping Array v1.0 at the Wellcome Centre for Human Genetics, University of Oxford. JHS samples were genotyped using the Affymetrix Genome-Wide Human SNP Array 6.0. Genotypes were called from intensities using Illuminus and GenCall clustering algorithms in GenomeStudio (Illumina Inc., San Diego, USA) incorporating data from pre-determined genotypes.

We performed quality control (QC) for samples and single nucleotide polymorphisms (SNPs) using SNPs mapped to Human Genome Build 37. Using the appropriate strand files (http://www.well.ox.ac.uk/~wrayner/strand/), we removed low quality variants (those that mapped to multiple regions within the human genome or did not map to any region) and duplicate SNPs. Only autosomal SNPs were included in downstream QC analyses. We used a standardized H3A GWAS pipeline (https://github.com/h3abionet/h3agwas) to perform sample and SNP QC separately for each cohort (using identical steps).[58,59] Samples having discordant sex information, call rates <98%, relatedness cutoffs of between 11 and 70% (accounts for cryptic relatedness), and heterozygosity <0.15 or >0.343 were removed. SNPs were retained if they passed thresholds of >99% call rate, minor allele frequency (MAF) >0.01, and Hardy–Weinberg equilibrium (HWE) adjusted p-value >0.008.

We imputed our quality-controlled genotypes using the African Genome Resources haplotype reference panel that contains 4,956 samples from 1000 Genomes Phase 3 and ~2000 Ugandan samples and ~100 Ethiopians/Egyptians/Namibians/South Africans (n= 93,421,145 biallelic SNPs). This reference panel is hosted by the Wellcome Sanger Institute (https://imputation.sanger.ac.uk/).[60] We filtered out the resulting imputation dataset for SNPs with information score ≥ 0.3 and MAF ≥ 0.01 for association analyses.

Statistical analyses
We used inverse-normal transformation to normalize the distribution of the iron biomarkers in R version 3.4.3. We performed association testing for each study site using genome-wide complex trait analysis (GCTA) software version 1.24.4[61] fitting a linear mixed model including a genetic relatedness matrix (GRM) as a random effect. We included age and sex as covariates. Inclusion of CRP as a covariate did not change the results. Inverse normal transformation measures were used throughout our analyses and all results are reported as such. We meta-analyzed our discovery study-specific summary statistics using METASOFT version 2.0.1.[62] Results were visualized using Manhattan and QQ plots generated using R packages *GenABEL* and *qqman*. Inflation factors (*lambda*) were calculated to check for population stratification. We used LocusZoom.js v0.12.0 (https://my.locuszoom.org/) to create GWAS regions and added LDs calculated using Plink version 1.9 from our discovery cohorts data.

Genome-wide significant SNPs ($P<5\times10^{-8}$) from discovery GWAS meta-analysis were selected for testing in the replication study of JHS. We considered a GWAS result to replicate if the effect in the replication was in the same direction as in the discovery sample, and if the association in the replication sample was statistically significant ($P< 0.05$).

To evaluate differences in the genetics of iron between European and African ancestry populations, we analyzed transferability of iron GWAS findings between continental African and European populations. We compared our discovery association results with previously reported variants from European iron GWAS in order to determine the importance of these same variants in an African population. We identified index SNPs from the largest genome-wide meta-analysis of European iron biomarkers GWAS[13,14] and compared their replication in our African iron GWAS. All lead/independent SNPs reported in the European iron GWAS were identified in African children with available iron GWAS data. We then compared effect sizes and P values between European and African populations. A SNP was considered to replicate if the effect in Africans was in the same direction as in the European iron GWAS, and if the association in Africans was statistically significant ($P< 0.05$). For each iron biomarker, we further calculated a PRS[63] weighted using effect sizes reported in European iron biomarkers GWAS and reflecting an increase in levels of the biomarker. A linear regression model adjusted for age, sex and study site was used to determine any association between the PRS and iron biomarkers measured in African children. In Africans, we used transferrin as a proxy measure for total iron binding capacity (TIBC) that was reported in the European iron GWAS. TIBC indicates the amount of iron that can be bound to transferrin.

Since iron status might influence risk of malaria and bacterial infections,[25,26] we investigated the association between our identified novel African-specific SNPs and these infections using the largest case-control GWAS of severe malaria[27] and bacteremia[28,29] described above. Since the *SDR16C5* lead SNP appeared across all five iron GWAS African sites, we meta-analysed severe malaria results from all nine MalariaGen African sites (The Gambia, Mali, Burkina Faso, Ghana, Nigeria, Cameroon, Malawi, Tanzania, and Kenya) using the '*meta*' package in R version 4.2.3 assuming fixed effects. Since the *GTF3C5* lead SNP, rs2905094, showed an East-African specific effect on transferrin levels, and was not significant in other African populations despite having a similar allele frequency, we meta-analyzed severe malaria GWAS results from East African countries (Kenya, Malawi and Tanzania). For the *FREM3* locus lead SNP, which is Kenyan specific and very rare in other tested populations from our cohorts or in reference panels such as 1000G, we identified its effect size in the Kenya severe malaria GWAS only.

Functional annotation

To explore mechanisms underpinning associations between iron status and severe infection and to identify biologically plausible instruments, we annotated genome-wide significant SNPs and investigated causal variants. We performed statistical fine-mapping by conducting conditional forward stepwise regression to determine whether there are independent signals. This entailed conditioning on the lead SNP from our GWAS by treating it as an adjusting covariate and testing the remaining SNPs. Once a new independent secondary signal was identified, we performed sequential testing by adjusting for the lead SNP until no conditional tests were statistically significant.

We performed comprehensive functional annotation using FUMA (https://fuma.ctglab.nl/) to annotate our previously unreported SNPs and determine genomic risk loci. We applied an R-

squared of 0.2 and used Africans ("AFR") in the 1000G phase 3 as a reference panel. We further used VEP (https://www.ensembl.org/Tools/VEP) to annotate and prioritize variants. To identify expression quantitative trait locus (eQTL) variants, we used GTex (https://gtexportal.org/home/) and the African Functional Genomics Resources (https://github.com/smontgomlab/AFGR) which is the largest African eQTL database.

Haplotype analyses
Haplotype patterns were derived from the imputed and phased genotypes using a custom R script that computed Manhattan distances followed by hierarchical clustering using the *hclust* function in R software (version 4.0.2). The resulting clusters were visualised with heat maps, and haplotype patterns were identified by visually assessing regions of homogeneity. We focused on a 72 kb region defined by 210 SNPs surrounding the rs2905094 SNP. Haplotype pattern analysis was performed separately for the ancestral and derived alleles of rs2905094, both across all countries combined and for each country individually.

GWAS sample size and power
Power for the GWAS studies was calculated using the formulae given in Visscher et al 2017,[64] implemented in: https://github.com/kaustubhad/gwas-power. Given an allele frequency ≥ 5% and effect size ≥ 0.33 standard deviation (SD), the overall discovery sample size of 3928 provided more than 80% power to observe true genotypic and phenotypic associations at a genome-wide significance level (type I error=$5x10^{-8}$). For allele frequency ≥10%, we had over 80% power to see effect sizes ≥ 0.24 SD.

Acknowledgments: We thank all study participants who contributed to this study and staff involved with consent, sample and data collection and preparation. This work was funded by Wellcome (Grant numbers [224317 to JMM], [226014 to SHA], [202800 to TNW], [103951 to AOE], and [106289 to AJM] and Grant numbers [064693, 079110, 095778 to AME]) and with core awards to the KEMRI-Wellcome Trust Research Programme (203077), The Wellcome Centre for Human Genetics (090532, 203141) and the Wellcome Sanger Institute (098051, 206194). The Jackson Heart Study (JHS) is supported and conducted in collaboration with Jackson State University (HHSN268201800013I), Tougaloo College (HHSN268201800014I), the Mississippi State Department of Health (HHSN268201800015I) and the University of Mississippi Medical Center (HHSN268201800010I, HHSN268201800011I and HHSN268201800012I) contracts from the National Heart, Lung, and Blood Institute (NHLBI) and the National Institute on Minority Health and Health Disparities (NIMHD). The authors also wish to thank the staffs and participants of the JHS. The views expressed in this manuscript are those of the authors and do not necessarily represent the views of the National Heart, Lung, and Blood Institute; the National Institutes of Health; or the U.S. Department of Health and Human Services. JJG was funded by a National Institute for Health and Care Research Clinical Lectureship. The views expressed are those of the author(s) and not necessarily those of the NIHR or the Department of Health and Social Care. This study was published with the permission of the Director of KEMRI.

Author contributions: JMM, AJM, AME, TNW, AA, and SHA conceptualized and designed the methods for the research project; JMM performed the analyses; JMM, AJM, GB, AYC, AWM, RMM, KMA, RM, JJG, ELW, FMN, LMR, LE, SBS, SAM, AVSH, AP, PB, GH, GDS, MSS, AME, TNW, AA, and SHA were involved in resources generation and curation of data. JMM, AJM, AME, TNW, AA, and SHA were responsible for funding acquisition. JMM and SHA wrote the manuscript and all co-authors reviewed the manuscript.

Competing interests: The authors declare no competing interests.

Data and materials availability: All data are available in the main text or the Supplementary Materials. All direct genotypes from individuals after QC along-side imputed data have been submitted to the European Genome-Phenome Archive under accession, with the datasets under EGAD00010002578 and EGAD00010002583 (Uganda); EGAD00010002582 and EGAD00010002580 (South Africa); EGAD00010002581 and EGAD00010002579 (Burkina Faso). Applications for access to primary individual level de-identified phenotype data for the Kilifi, Kenya; Entebbe, Uganda; Banfora, Burkina Faso; and West Kiang, The Gambia cohorts can be made through the Data Governance Committee cgmrc@kemri-wellcome.org.

Code availability: All analyses used preassigned code defined within software packages that are publicly available as described in Methods. Any other requests for clarifications may be sought from the corresponding authors.

Supplementary Figures

Supplementary Figure 1: Manhattan plots for each African study site. N/A, not available.

Supplementary Figure 2: QQ plot for the discovery GWAS of iron markers.

Supplementary Figure 3. Meta-analysis of transferrin GWAS results by African region.

Supplementary Figure 4: Haplotype structures, linkage disequilibrium, and genetic map of the *GTF3C5* gene region. This figure presents the haplotype structures within a 72 KB region surrounding the rs2905094 (9:135902689 (GRCh37)) allele. **Panel A** displays the structure for the ancestral allele, while **Panel B** illustrates the structure for the derived alleles across all countries combined. Additionally, haplotypes for the derived allele are shown

separately for each country, as labelled in the panels. In each panel, the rows represent individual phased haplotypes, and the columns denote SNPs, sorted and clustered based on Manhattan distance. The ancestral allele is depicted by black columns, and the derived allele is represented by white columns. The dominant haplotypes identified in the mutant chromosomes are labelled as "Hap 1" and "Hap 2." The LD map is included with the Y-axis indicating the recombination rate in cM/Mb, and the X-axis displaying the chromosomal position referenced to GRCh37. The genetic map highlights the positions of three genome-wide significant SNPs - rs2905089, rs2905094, and rs2905095 - using blue, red, and green lines, respectively.

Country	OR (95% CI)	%Weight	n
East Africa			
Malawi	0.81 (0.67, 0.98)	39.39	2471
Tanzania	1.04 (0.76, 1.42)	8.69	798
Kenya	0.91 (0.79, 1.06)	51.92	3068
EA Overall (I^2 = 0.0%, p = 0.39)	**0.88 (0.78, 0.98)**	100.00	6337
West Africa			
The Gambia	0.97 (0.88, 1.07)	66.74	4979
Mali	0.90 (0.60, 1.35)	4.28	422
Burkina Faso	0.91 (0.68, 1.21)	8.58	1294
Ghana	1.04 (0.77, 1.40)	6.07	706
Nigeria	0.51 (0.19, 1.32)	1.89	133
Cameroon	0.92 (0.73, 1.17)	12.44	1217
WA Overall (I^2 = 0.0%, p = 0.69)	**0.95 (0.87, 1.03)**	100.00	8751

rs2905094 reduces severe malaria   rs2905094 increases severe malaria

Supplementary Figure 5. Meta-analyses of the effect sizes of the *GTF3C5* lead SNP, rs2905094 and severe malaria. We applied fixed effects meta-analysis of severe malaria GWAS results from the MalariaGen consortium by East and West African countries.

Supplementary Figure 6. Association between **a**) hepcidin and **b**) soluble transferrin receptor GWAS lead SNPs and other iron biomarkers. Linear regression models were adjusted for age, sex, CRP, and study site. Effect sizes were derived from an additive model.

Supplementary Figure 7. Association between hepcidin GWAS lead SNP, rs73596248 and anthropometric indices. Estimates are derived from an additive linear regression model. HAZ06, height-for-age z-score; WAZ06, weight-for-age z-score; WHZ06, weight-for-height z-score. Stunting was defined as HAZ06 < -2, underweight as WAZ06 < -2 and wasting as WHZ06 < -2 using the WHO Growth Reference Standards.

Supplementary Tables

Supplementary Table 1. Characteristics of study participants

Characteristic	Kenya n=1059		Uganda n=1360		Burkina Faso n=348		South Africa n=611		The Gambia n=550		Jackson Heart Study n=2868	
Median age (IQR)[a]	21.8 m (15.2-40.0)		24.1 m (23.9-35.9)		23.3 m (19.7-26.2)		12.0 m (11.9-12.2)		46.0 m (34.4-58.5)		54 y (44-64)	
	n/total	%	n/total	%	n/total	%	n/total	%	n/total	%	n/total	%
Gender: Females	514/1059	48.5	675/1360	49.6	173/348	49.7	294/611	48.1	250/550	45.5	1759/2868	61.3
Inflammation[b]	275/1020	27.0	314/1325	23.7	112/330	33.9	95/611	15.6	78/550	14.2	830/2864	29.0
Malaria parasitemia[c]	215/954	22.5	19/1340	6.8	66/321	20.6	0/611	0.0	59/548	10.8	n/a	n/a
Iron deficiency[d]	350/999	35.0	436/1254	34.8	115/324	35.5	251/611	41.1	118/550	21.5	298/2868	10.4
Anemia[e]	420/639	65.7	646/1299	49.7	288/331	87.0	n/a	n/a	329/545	60.4	728/2818	25.8
Iron deficiency anemia[f]	140/609	23.0	253/1197	21.1	96/309	31.1	n/a	n/a	85/545	15.6	174/2142	8.1
Stunting[g]	89/183	48.6	207/1342	15.4	109/325	33.5	n/a	n/a	167/438	38.1	n/a	n/a
Underweight[h]	95/334	28.4	112/1355	8.3	60/327	18.4	n/a	n/a	111/439	25.3	n/a	n/a
Wasting[i]	24/180	13.3	66/1340	4.9	22/325	6.8	n/a	n/a	36/438	8.2	n/a	n/a
Biomarker	n	GMean (SD)	n	GMean (SD)	n	GMean (SD)	n	GMean (SD)	n	GMean (SD)	n	GMean (SD)
Ferritin, µg/L	999	21.9 (3.1)	1254	20.9 (2.9)	324	22.2 (2.9)	611	14.9 (2.6)	550	24.4 (2.4)	2868	112.2 (2.6)
sTfR, mg/L	1047	17.8 (1.5)	1330	6.7 (2.0)	342	17.7 (1.7)	610	10.8 (1.5)	510	3.5 (1.4)	n/a	n/a
Hepcidin, µg/L	976	5.7 (3.6)	1321	6.8 (3.3)	309	5.3 (4.2)	600	7.8 (3.5)	550	5.5 (4.7)	n/a	n/a
Serum iron, µmol/L	1009	6.5 (2.0)	n/a	n/a	337	6.0 (1.8)	n/a	n/a	550	8.6 (1.6)	2868	74.1 (1.5)
Transferrin, g/L	998	2.7 (1.3)	1320	2.7 (1.3)	327	2.7 (1.3)	611	2.6 (1.3)	550	2.6 (1.2)	2868	2.0 (1.2)
TSAT, %[j]	977	9.4 (2.2	n/a	n/a	325	8.9 (2.0)	n/a	n/a	550	12.8 (1.7)	2868	25.7 (1.5)
Hemoglobin, g/dL	639	10.2 (1.2)	1299	10.8 (1.1)	331	9.5 (1.1)	n/a	n/a	545	10.6 (1.1)	2818	12.9 (1.1)
Body mass index	243	15.1 (1.2)	1342	15.7 (1.1)	324	15.4 (1.2)	n/a	n/a	549	14.6 (1.1)	2864	31.2 (1.2)
Inflammatory marker[k]	1020	1.8 (5.2)	1325	1.4 (5.1)	330	2.8 (6.1)	611	0.9 (5.0)	550	0.4 (1.3)	2864	2.5 (3.5)

IQR, interquartile range; n/a, not available; sTfR, soluble transferrin receptors; TSAT, transferrin saturation; CRP, C-reactive protein; GMean, geometric means.

a age in months for children and age in years for African-American (Jackson Heart Study) adults.

b Inflammation was defined as C-reactive protein > 5mg/L or α1-antichymotrypsin > 0.6g/dL (in The Gambia). Inflammatory marker values indicate CRP (mg/L) for Kenya, Uganda, Burkina Faso and South Africa, and ACT (g/dL) for The Gambia.

c Malaria parasitemia was defined as *Plasmodium falciparum* parasitemia positive at any density. Participants in South Africa and Jackson Heart Study were not exposed to malaria.

d Iron deficiency defined using WHO definition as ferritin < 12 µg/L or <30 µg/L in presence of inflammation (C-reactive protein>5mg/L or α1-antichymotrypsin >0.6g/dL) in children < 5 years or <15µg/L in children >=5 years. In the JHS iron deficiency was defined as ferritin <30µg/L

e Anemia was defined as hemoglobin <11 g/dL in children aged <5 years, or hemoglobin <11.5 g/dL in children ≥5 years[65]. In the JHS and Nairobi, ID was defined as ferritin <30µg/L. In the Jackson Heart Study, anemia was defined as hemoglobin <12 g/dL in women or <13 g/dL in men[65,66].

f Iron deficiency anemia was defined as iron deficiency and anemia.

g Stunting was defined as height-for-age z-score < -2

h Underweight was defined as weight-for-age z-score < -2

i Wasting was defined as weight-for-height z-score < -2

j TSAT was calculated as ((Iron (µmol/L)/Transferrin (g/L) x 25.1) x 100) except in The Gambia where it was calculated using iron and unsaturated iron binding capacity. Iron and TSAT missing in Uganda and South Africa and transferrin missing in The Gambia.

k Inflammatory markers included CRP measured in Kenya, Uganda, Burkina Faso, South Africa, and Jackson Heart Study and ACT measured in The Gambia.

Supplementary Table 2. Summary of quality control (QC) steps for each population. The number of individuals and variants removed within each population are presented.

	Kenya	Uganda	Burkina Faso	South Africa	The Gambia	The Gambia	JHS
Pre-QC							
Total number sent for genotyping	1,312	1,429	354	777	574	46	3,021
Number of variants typed and mapping to Build 37	2,114,814	2,337,579	2,337,579	2,337,579	2,297,939	2,114,814	874,712
Individuals QC							
Individuals with missingness >0.03	3	14	-	2	46	-	135
Individuals with extreme heterozygosity (>0.343 or <0.15)	-	-	-	-	-	-	-
Individuals failing sex- check	58	3	-	2	13	-	4
Individuals IBD>0.9 and not twins / triplets	32	22	-	24	9	2	3
Autosomal variants QC							
Non-autosomal variants	36,413	51,101	51,101	51,101	36,847	36,413	34,771
Autosomal duplicate variants	28	9,248	9,248	9,248	182,365	28	2,830
Autosomal variants with missingness>0.01	93,959	262,676	263,935	226,503	113,215	89,922	122,182
Autosomal variants with MAF<0.01	211,678	392,392	399,486	416,608	222,631	231,339	342
Autosomal variants in Hardy-Weinberg disequilibrium (P<0.008)	29,735	29,266	20,089	25,174	21,989	6,544	17,258
Post-QC							
Total individuals remaining	1,219	1,390	354	749	506	44	2,879
Total autosomal variants remaining	1,743,001	1,592,896	1,593,720	1,608,945	1,720,892	1,750,568	697,329

Genotyping for the 574 and 46 Gambian samples was done separately and then merged after quality control giving total individual remaining=550 and total autosomal variants remaining 1,776,801. JHS, Jackson Heart Study.

Supplementary Table 3. Number of variants imputed into individual datasets.

Cohort	Total imputed	Total high quality	Total tested for association
Kenya	89,838,087	13,870,649	7,756,141
Uganda	89,838,087	13,982,934	13,765,898
Burkina Faso	89,838,087	13,128,631	13,042,768
South Africa	89,838,087	15,010,497	14,457,588
The Gambia	89,838,087	12,770,741	10,295,094
Jackson Heart Study	89,838,087	13,455,566	13,349,077

High quality SNPs are those only with an information score greater than 0.3 and those tested for association were only those of high quality with a minor allele frequency (MAF) greater than 0.01.

Supplementary Table 4. Genome-wide significant variants associated with transferrin levels at the *TF* locus in a meta-analysis of African sites, in African American and European populations

SNP	CHR	Hg19BP	EA	OA	Gene	African sites meta-analysis[a]				African Americans[b]				Europeans[c]				
						EAfreq	Beta	Se	P	EAfreq	Beta	Se	P	EA	EAfreq	Beta	Se	P
rs6762719	3	133480817	G	A	TF	0.25	0.28	0.03	1.59E-27	0.24	0.46	0.03	7.65E-56	A	0.69	-0.37	0.005	6.01e-1121
rs8177253	3	133480192	T	C	TF	0.25	0.28	0.03	2.54E-27	0.24	0.46	0.03	3.67E-56	T	0.30	0.37	0.005	4.85e-1116
rs8177252	3	133480174	A	C	TF	0.25	0.28	0.03	4.25E-26	0.24	0.47	0.03	7.13E-56	A	0.30	0.37	0.005	5.19e-1116
rs1880669	3	133483696	C	T	TF	0.35	0.24	0.02	4.88E-25	0.35	0.37	0.03	2.54E-44	T	0.43	-0.26	0.004	3.72e-776
rs8177260	3	133480458	A	T	TF	0.27	0.26	0.03	7.36E-25	0.26	0.42	0.03	2.98E-49	A	0.32	0.33	0.005	1.08e-1005
rs8177247	3	133479499	G	T	TF	0.23	0.28	0.03	2.66E-23	0.22	0.47	0.03	8.79E-54	T	0.70	-0.37	0.005	7.73e-1122
rs1358023	3	133484393	C	T	TF	0.27	0.25	0.03	1.72E-22	0.27	0.39	0.03	4.54E-43	T	0.70	-0.36	0.005	9.67e-1105
rs1405022	3	133481718	T	C	TF	0.3	0.21	0.02	2.91E-18	0.32	0.35	0.03	3.93E-37	T	0.56	0.26	0.004	1.13e-778
rs6439439	3	133481498	T	G	TF	0.22	0.23	0.03	4.25E-17	0.23	0.42	0.03	7.26E-44	T	0.30	0.37	0.005	3.48e-1121
rs8177261	3	133480588	C	A	TF	0.22	0.23	0.03	4.75E-17	0.23	0.42	0.03	9.63E-44	A	0.70	-0.37	0.005	4.55e-1121
rs8177262	3	133480652	A	G	TF	0.22	0.23	0.03	4.75E-17	0.23	0.42	0.03	1.02E-43	A	0.30	0.37	0.005	3.57e-1121
rs8177256	3	133480276	T	G	TF	0.22	0.23	0.03	5.65E-17	0.23	0.42	0.03	9.46E-44	T	0.30	0.37	0.005	1.02e-1116
rs8177259	3	133480388	C	T	TF	0.22	0.22	0.03	6.13E-17	0.23	0.42	0.03	1.08E-43	T	0.70	-0.37	0.005	1.90e-1117
rs4854748	3	133446881	T	C	TF	0.2	-0.24	0.03	1.12E-16	0.19	-0.27	0.03	2.10E-16	T	0.24	-0.25	0.005	3.05e-578
rs3811647	3	133484029	A	G	TF	0.22	0.22	0.03	3.45E-16	0.22	0.42	0.03	1.68E-43	A	0.31	0.36	0.005	1.96e-1105
rs8177185	3	133464618	A	G	TF	0.13	-0.28	0.03	6.81E-16	0.1	-0.27	0.04	1.02E-10	A	0.07	-0.29	0.008	9.09e-327
rs4854755	3	133452671	A	G	TF	0.16	-0.24	0.03	3.38E-14	0.15	-0.21	0.04	2.29E-09	A	0.27	-0.17	0.004	2.50e-370
rs147634092	3	133458182	G	A	TF	0.15	-0.25	0.03	4.70E-14	0.14	-0.23	0.04	3.23E-10	A	0.76	0.25	0.005	2.11e-585
rs6801244	3	133454847	A	G	TF	0.15	-0.25	0.03	5.11E-14	0.14	-0.23	0.04	3.19E-10	A	0.24	-0.25	0.005	2.09e-579
rs139374172	3	133458144	G	A	TF	0.15	-0.25	0.03	6.89E-14	0.14	-0.23	0.04	3.19E-10	A	0.76	0.25	0.005	3.46e-585
rs10935083	3	133453827	C	T	TF	0.13	-0.26	0.03	9.35E-14	0.1	-0.27	0.04	1.76E-10	T	0.93	0.29	0.008	7.18e-320
rs9824630	3	133455482	G	A	TF	0.11	-0.31	0.04	9.79E-14	0.1	-0.24	0.05	1.05E-07	n/a	n/a	n/a	n/a	n/a
rs4854749	3	133447397	A	G	TF	0.15	-0.24	0.03	1.01E-13	0.14	-0.23	0.04	1.55E-10	A	0.24	-0.25	0.005	1.23e-577
rs4854750	3	133447489	G	C	TF	0.15	-0.24	0.03	2.17E-13	0.14	-0.23	0.04	1.65E-10	C	0.76	0.25	0.005	6.85e-579
rs10935081	3	133451352	T	C	TF	0.15	-0.24	0.03	2.30E-13	0.14	-0.23	0.04	2.02E-10	T	0.24	-0.25	0.005	1.88e-578
rs10049093	3	133456056	T	A	TF	0.15	-0.24	0.03	2.33E-13	0.14	-0.23	0.04	3.14E-10	A	0.76	0.25	0.005	3.46e-583
rs16840812	3	133462807	C	T	TF	0.15	-0.24	0.03	2.74E-13	0.13	-0.22	0.04	3.32E-09	T	0.82	0.23	0.005	2.07e-509
rs79799101	3	133433773	T	C	TF	0.13	-0.26	0.04	2.99E-13	0.1	-0.27	0.04	2.45E-10	T	0.07	-0.29	0.008	1.23e-319
rs4241356	3	133452444	G	A	TF	0.15	-0.24	0.03	3.07E-13	0.14	-0.23	0.04	2.32E-10	A	0.76	0.25	0.005	6.11e-579
rs8177240	3	133477701	G	T	TF	0.17	0.21	0.03	1.02E-12	0.17	0.41	0.03	6.11E-34	T	0.69	-0.36	0.005	1.80e-1080
rs556138790	3	133484713	A	T	TF	0.06	0.34	0.05	1.14E-12	0.04	0.36	0.06	3.20E-08	n/a	n/a	n/a	n/a	n/a
rs4532136	3	133466885	A	G	TF	0.12	-0.24	0.03	1.26E-12	0.1	-0.27	0.04	1.96E-10	A	0.07	-0.29	0.008	7.57e-322
rs8177250	3	133479737	C	T	TF	0.06	0.33	0.05	1.87E-12	0.05	0.45	0.06	5.12E-16	n/a	n/a	n/a	n/a	n/a
rs8177276	3	133484336	C	T	TF	0.05	0.34	0.05	2.15E-12	0.03	0.51	0.07	4.16E-12	n/a	n/a	n/a	n/a	n/a
rs8177249	3	133479643	T	C	TF	0.06	0.33	0.05	3.91E-12	0.05	0.46	0.06	5.11E-16	n/a	n/a	n/a	n/a	n/a
rs8177184	3	133464496	G	A	TF	0.15	-0.25	0.04	3.95E-12	0.14	-0.21	0.04	9.04E-09	A	0.82	0.23	0.005	5.29e-512
rs386666076	3	133484712	A	G	TF	0.26	0.19	0.03	4.77E-12	0.27	0.37	0.03	1.23E-39	A	0.30	0.37	0.005	9.81e-1104
rs8177241	3	133477961	T	C	TF	0.06	0.32	0.05	5.02E-12	0.05	0.46	0.06	5.37E-16	n/a	n/a	n/a	n/a	n/a
rs8177181	3	133463798	A	T	TF	0.14	-0.25	0.04	5.45E-12	0.14	-0.21	0.04	9.62E-09	A	0.18	-0.23	0.005	1.65e-507
rs61099606	3	133484394	A	G	TF	0.06	0.31	0.05	1.69E-11	0.04	0.33	0.06	7.53E-08	n/a	n/a	n/a	n/a	n/a
rs2718798	3	133492088	C	A	TF	0.43	0.15	0.02	2.66E-11	0.43	0.22	0.03	3.02E-18	A	0.41	-0.24	0.004	5.15e-707
rs4582061	3	133438026	A	G	TF	0.12	-0.25	0.04	3.39E-11	0.1	-0.27	0.04	7.05E-10	A	0.07	-0.29	0.008	6.64e-323
rs3811658	3	133476852	T	C	TF	0.16	0.2	0.03	4.13E-11	0.17	0.41	0.03	2.31E-33	T	0.31	0.36	0.005	4.51e-1081
rs4428180	3	133466374	G	A	TF	0.14	-0.21	0.03	4.23E-11	0.13	-0.22	0.04	3.29E-09	A	0.84	0.25	0.005	5.74e-522

SNP	CHR	Hg19BP	EA	OA	Gene	EA Freq	Beta	Se	P	EA Freq	Beta	Se	P	EA	EA Freq	Beta	Se	P
rs8177258	3	133480387	A	C	*TF*	0.17	0.2	0.03	6.95E-11	0.17	0.41	0.03	2.50E-34	A	0.30	0.37	0.005	2.39e-1117
rs16840699	3	133437601	T	C	*TF*	0.15	-0.23	0.03	8.89E-11	0.13	-0.21	0.04	3.63E-08	T	0.07	-0.29	0.008	5.28e-317
rs8177243	3	133478578	C	T	*TF*	0.04	0.47	0.07	9.98E-11	0.02	0.74	0.1	4.26E-14	n/a	n/a	n/a	n/a	n/a
rs41295782	3	133478388	A	G	*TF*	0.04	0.47	0.07	1.01E-10	0.02	0.74	0.1	4.39E-14	n/a	n/a	n/a	n/a	n/a
rs1799852	3	133475722	T	C	*TF*	0.06	-0.36	0.06	1.16E-10	0.06	-0.4	0.05	8.98E-14	T	0.14	-0.31	0.006	3.38e-570
rs11717407	3	133438264	G	A	*TF*	0.12	-0.24	0.04	1.27E-10	0.1	-0.27	0.04	6.90E-10	A	0.93	0.29	0.008	2.00e-322
rs8177263	3	133480754	G	A	*TF*	0.03	0.47	0.07	1.41E-10	0.02	0.73	0.1	1.22E-13	n/a	n/a	n/a	n/a	n/a
rs8177266	3	133481255	T	C	*TF*	0.03	0.47	0.07	1.41E-10	0.02	0.73	0.1	1.22E-13	n/a	n/a	n/a	n/a	n/a
rs8177268	3	133481751	A	G	*TF*	0.03	0.47	0.07	1.41E-10	0.02	0.73	0.1	1.22E-13	n/a	n/a	n/a	n/a	n/a
rs8177271	3	133482230	A	G	*TF*	0.16	0.2	0.03	2.35E-10	0.17	0.41	0.03	4.64E-34	A	0.30	0.36	0.005	1.34e-1106
rs8177272	3	133482870	A	G	*TF*	0.16	0.2	0.03	3.77E-10	0.17	0.41	0.03	4.31E-34	A	0.30	0.36	0.005	4.94e-1105
rs8177280	3	133485664	G	T	*TF*	0.04	0.45	0.07	4.19E-10	0.02	0.72	0.1	3.11E-13	n/a	n/a	n/a	n/a	n/a
rs16840692	3	133433900	A	G	*TF*	0.12	-0.24	0.04	4.74E-10	0.1	-0.28	0.04	3.42E-10	A	0.07	-0.29	0.008	5.58e-323
rs16840751	3	133450433	A	G	*TF*	0.07	-0.31	0.05	5.16E-10	0.06	-0.28	0.05	2.43E-07	A	0.07	-0.29	0.008	1.21e-318
rs4854754	3	133448602	A	G	*TF*	0.08	-0.3	0.05	7.94E-10	0.06	-0.28	0.05	2.13E-07	A	0.07	-0.29	0.008	7.39e-321
rs74639761	3	133433523	G	T	*TF*	0.1	-0.24	0.04	1.08E-09	0.09	-0.28	0.05	2.06E-09	T	0.93	0.29	0.008	3.66e-314
rs4854757	3	133454304	A	G	*TF*	0.07	-0.3	0.05	1.18E-09	0.06	-0.28	0.05	2.22E-07	A	0.07	-0.29	0.008	3.10e-320
rs8177235	3	133476083	A	G	*TF*	0.07	-0.3	0.05	1.34E-09	0.06	-0.41	0.05	5.85E-15	A	0.08	-0.31	0.007	7.26e-428
rs8177303	3	133494678	T	C	*TF*	0.05	0.46	0.08	1.52E-09	0.01	0.7	0.11	5.75E-10	n/a	n/a	n/a	n/a	n/a
rs146105056	3	133492085	T	C	*TF*	0.05	0.46	0.08	1.56E-09	0.01	0.71	0.11	3.21E-10	n/a	n/a	n/a	n/a	n/a
rs2715634	3	133487807	C	T	*TF*	0.53	0.14	0.02	2.84E-09	0.53	0.25	0.03	9.59E-23	T	0.44	-0.26	0.004	1.31e-775
rs8177245	3	133479230	G	A	*TF*	0.16	0.19	0.03	3.23E-09	0.16	0.41	0.03	1.45E-33	A	0.70	-0.37	0.005	3.23e-1121
rs8177275	3	133484066	C	T	*TF*	0.04	0.39	0.07	4.48E-09	0.03	0.42	0.08	4.07E-07	n/a	n/a	n/a	n/a	n/a
rs6773891	3	133262438	C	T	*TF*	0.45	-0.13	0.02	4.58E-09	0.48	-0.13	0.03	2.74E-07	T	0.36	-0.10	0.004	1.2E-161
rs10935068	3	133262807	A	T	*TF*	0.48	-0.13	0.02	8.31E-09	0.49	-0.13	0.03	7.45E-07	A	0.64	0.10	0.004	2.03E-162
rs6762415	3	133478557	G	T	*TF*	0.42	0.14	0.02	1.15E-08	0.42	0.28	0.03	3.51E-28	T	0.50	-0.28	0.005	1.90e-867
rs8177248	3	133479626	T	C	*TF*	0.1	0.21	0.04	1.37E-08	0.13	0.38	0.04	1.36E-22	T	0.30	0.37	0.005	1.56e-1116
rs2002016	3	133265345	G	A	*TF*	0.49	-0.13	0.02	1.62E-08	0.53	-0.11	0.03	1.32E-05	A	0.37	-0.10	0.004	2.01E-158
rs9810872	3	133515796	C	T	*TF*	0.11	-0.2	0.04	2.85E-08	0.14	-0.18	0.04	1.54E-06	T	0.66	0.15	0.004	1.00e-320
rs7638018	3	133495461	G	A	*TF*	0.18	0.16	0.03	4.19E-08	0.19	0.33	0.03	5.99E-25	A	0.70	-0.36	0.005	2.71e-1103
rs113656013	3	133415112	T	C	*TF*	0.17	-0.17	0.03	4.66E-08	0.17	-0.14	0.03	2.23E-05	T	0.001	-0.10	0.202	0.609
rs34252038	3	133490621	T	G	*TF*	0.19	0.15	0.03	4.86E-08	0.2	0.31	0.03	6.31E-23	T	0.30	0.37	0.005	2.23e-1101

CHR, chromosome; Hg19BP, human build 19 base pair; EA, effect allele; OA, other allele; EA Freq, effect allele frequency; Se, standard error; P, additive model p value; n/a, not available.

[a]Beta reflects fixed effects meta-analysis

[b]Data from the Jackson Heart Study.

[b]Data from Moksnes et al, 2022, the largest European iron GWAS. TIBC (reported by Moksnes et al 2022) is compared with transferrin from current study.

Supplementary Table 5. Genome-wide significant SNPs associated with transferrin levels at *GTF3C5* locus in meta-analysis of Kenya and Uganda and how they compare with other populations

		rs2905094	rs2905095	rs2905089
	SNP	**rs2905094**	**rs2905095**	**rs2905089**
	CHR	9	9	9
	Hg19BP	135902689	135902793	135899844
	EA	T	T	C
	OA	C	C	A
Kenya+Uganda meta-analysis	**EAFreq**	0.16	0.18	0.25
	Beta	-0.23	-0.22	-0.18
	Se	0.04	0.04	0.03
	P	8.08E-09	1.75E-08	6.74E-08
Uganda	**EAFreq**	0.18	0.20	0.29
	Beta	-0.20	-0.19	-0.18
	Se	0.05	0.05	0.04
	P	9.82E-05	9.39E-05	3.60E-05
Kenya	**EAFreq**	0.14	0.16	0.21
	Beta	-0.29	-0.26	-0.20
	Se	0.07	0.06	0.06
	P	1.15E-05	3.17E-05	4.94E-04
South Africa	**EAFreq**	0.14	0.16	0.25
	Beta	-0.15	-0.13	-0.10
	Se	0.08	0.08	0.07
	P	7.35E-02	9.95E-02	1.59E-01
The Gambia	**EAFreq**	0.23	0.25	0.32
	Beta	0.12	0.06	-0.04
	Se	0.07	0.07	0.06
	P	1.06E-01	3.55E-01	5.81E-01
Burkina Faso	**EAFreq**	0.17	0.18	0.25
	Beta	-0.03	-0.06	0.09
	Se	0.10	0.09	0.09
	P	7.86E-01	5.08E-01	3.02E-01
Jackson Heart Study	**EAFreq**	0.18	0.19	0.27
	Beta	-0.02	-0.03	-0.02
	Se	0.04	0.04	0.03
	P	5.78E-01	4.62E-01	5.43E-01
Europe[a]	**EAFreq**	0.33	0.30	0.33
	Beta	0.004	0.01	0.001
	Se	0.004	0.007	0.004
	P	0.32	0.11	0.74

CHR, chromosome; Hg19BP, human build 19 base pair; EA, effect allele; OA, other allele; EA Freq, effect allele frequency; Se, standard error; P, additive model p value
[a]Effect of SNPs on transferrin iron binding capacity reported by Moksnes et al 2022 in Europeans.

Supplementary Table 6. Top SNPs associated with hepcidin levels in continental African populations and their frequencies in European, Asian and Latin American populations

									Africans	Europeans	Asians	Latin Americans
CHR	SNP	Hg19BP	EA	OA	EAFreq	Beta	Se	P		EAfreq	EAfreq	EAfreq
8	rs73596248	57134187	A	G	0.06	0.27	0.05	4.51E-08		0.00035	0.00	0.006
8	rs59493383	57199697	G	T	0.06	0.27	0.05	6.63E-08		0.0003	0.00	0.014
8	rs73596251	57135454	C	T	0.06	0.27	0.05	6.64E-08		0.0003	0.00	0.007
8	rs57580475	57199467	G	C	0.06	0.27	0.05	8.78E-08		0.0003	0.00	0.014
8	rs73596246	57133643	C	G	0.06	0.27	0.05	4.51E-08		0.0003	0.00	0.007

CHR, chromosome; Hg19BP, human build 19 base pair; EA, effect allele; OA, other allele; EAFreq, effect allele frequency; Se, standard error; P, additive model p value. Africans includes meta-analysed results from the five continental African sites included in this study. Frequencies for Europeans, Asians and Latin Americans are from dbSNP database (https://www.ncbi.nlm.nih.gov/snp) Allele Frequency Aggregator (ALFA).

Supplementary Table 7. Top SNPs associated with soluble transferrin receptor levels in Kenya and how they compare with other continental African populations

						Kenya				Uganda				South Africa				Burkina Faso			
SNP[a]	CHR	Hg19BP	EA	OA	Gene	EAFreq	Beta	Se	P	EAFreq	Beta	Se	P	EAFreq	Beta	Se	P	EAFreq	Beta	Se	P
rs552439837	4	144446525	G	A	*FREM3*	0.09	0.48	0.09	2.78E-08	Missing	NA	NA	NA	Missing	NA	NA	NA	Missing	NA	NA	NA
rs573724761	4	144449330	G	C	*FREM3*	0.09	0.48	0.09	2.78E-08	Missing	NA	NA	NA	Missing	NA	NA	NA	Missing	NA	NA	NA
rs141274959	4	144587704	A	G	*FREM3*	0.07	0.56	0.10	6.00E-08	Missing	NA	NA	NA	Missing	NA	NA	NA	Missing	NA	NA	NA
rs1040736918	4	144961711	T	C	*FREM3*	0.06	0.74	0.14	6.15E-08	Missing	NA	NA	NA	Missing	NA	NA	NA	Missing	NA	NA	NA
rs554867097	4	145087102	G	A	*FREM3*	0.06	0.57	0.11	6.22E-08	Missing	NA	NA	NA	0.02	-0.01	0.22	0.97	Missing	NA	NA	NA
rs113319340	4	144462931	G	T	*FREM3*	0.10	0.43	0.08	7.66E-08	Missing	NA	NA	NA	Missing	NA	NA	NA	Missing	NA	NA	NA
rs563199246	4	144504069	A	G	*FREM3*	0.09	0.46	0.09	7.96E-08	Missing	NA	NA	NA	Missing	NA	NA	NA	Missing	NA	NA	NA
rs111374053	4	144501489	T	C	*FREM3*	0.09	0.45	0.08	8.17E-08	Missing	NA	NA	NA	Missing	NA	NA	NA	Missing	NA	NA	NA
rs34330779	4	145124084	A	G	*FREM3*	0.09	0.42	0.08	9.01E-08	Missing	NA	NA	NA	0.02	-0.34	0.20	0.09	0.01	0.02	0.33	0.94
rs535317699	4	144532466	C	G	*FREM3*	0.08	0.52	0.10	9.28E-08	Missing	NA	NA	NA	Missing	NA	NA	NA	NA	NA	NA	NA

CHR, chromosome; Hg19BP, human build 19 base pair; EA, effect allele; OA, other allele; EAFreq, effect allele frequency; Se, standard error; P, additive model p value; NA, not applicable
[a]These SNPs were monomorphic in The Gambia, JHS, and in European populations.

Supplementary Table 8. Replication of European lead hepcidin and sTfR GWAS SNPs in current study of African children

Trait	SNP	CHR:Hg19BP	Gene	EA	OA	N	EAFreq	Beta	Se	P	N	EAFreq	Beta	Se	P
						European ancestry					**Continental Africa**				
Hepcidin	rs75965181	1:22584002	MIR4418	A	T	91675	0.023	-0.118	0.017	2.76E-12	n/a	n/a	n/a	n/a	n/a
Hepcidin	rs201255457	1:220291414	IARS2	A	ATC	75972	0.989	-0.156	0.026	3.15E-09	n/a	n/a	n/a	n/a	n/a
Hepcidin	rs6706968	2:121310269	AC073257.1	A	C	82690	0.424	0.03	0.005	1.43E-08	3756	0.115	0.056	0.037	0.131
Hepcidin	rs149682241	2:190390963	SLC40A1	C	G	91675	0.026	0.101	0.015	4.33E-11	n/a	0.011	n/a	n/a	n/a
Hepcidin	rs7568449	2:190521054	ASNSD1	T	C	89099	0.762	0.043	0.006	1.88E-13	3756	0.325	0.017	0.027	0.525
Hepcidin	rs885122	5:115331335	LVRN	A	G	91675	0.598	-0.043	0.005	2.28E-18	3756	0.65	-0.005	0.024	0.844
Hepcidin	rs79220007	6:26098474	HFE	T	C	91675	0.93	0.094	0.01	1.12E-22	n/a	n/a	n/a	n/a	n/a
Hepcidin	rs10090558	8:10577987	SOX7	A	G	78548	0.918	-0.06	0.01	4.64E-10	3756	0.761	-0.043	0.028	0.129
Hepcidin	rs9739943	12:51439858	LETMD1	T	C	91675	0.059	0.079	0.01	3.78E-14	3756	0.201	-0.005	0.029	0.854
Hepcidin	rs11609805	12:57735045	R3HDM2	A	G	91675	0.27	-0.032	0.006	5.24E-09	3756	0.066	0.041	0.054	0.45
Hepcidin	rs996347	14:34410892	EGLN3	T	C	91675	0.645	-0.033	0.005	6.06E-11	3756	0.849	0.012	0.033	0.718
Hepcidin	rs549876436	15:45274530	TERB2	T	C	83099	0.907	0.082	0.011	7.30E-15	n/a	n/a	n/a	n/a	n/a
Hepcidin	rs2600895	15:45319959	SORD	A	G	88174	0.913	0.082	0.012	2.73E-11	n/a	n/a	n/a	n/a	n/a
Hepcidin	rs113000093	15:45320493	SORD	C	G	85598	0.089	-0.078	0.013	7.01E-10	n/a	n/a	n/a	n/a	n/a
Hepcidin	rs199138	15:45387550	DUOX2	A	G	91675	0.076	-0.122	0.009	1.55E-40	3756	0.129	0	0.033	0.996
Hepcidin	rs71378510	16:353122	AXIN1	A	G	91675	0.907	0.05	0.009	1.62E-08	n/a	n/a	n/a	n/a	n/a
Hepcidin	rs34523089	17:56436109	RNF43	T	C	89817	0.162	0.063	0.007	8.15E-21	n/a	n/a	n/a	n/a	n/a
Hepcidin	rs104894696	19:35775902	HAMP	A	G	76690	0.003	-0.387	0.054	4.90E-13	n/a	n/a	n/a	n/a	n/a
Hepcidin	rs485073	19:49207255	FUT2	A	G	91675	0.453	0.046	0.005	4.64E-21	3756	0.504	0.026	0.023	0.261
Hepcidin	rs855791	22:37462936	TMPRSS6	A	G	91675	0.429	0.042	0.005	1.34E-17	3756	0.077	0.039	0.041	0.347
sTfR	rs6696846	1:205041952	CNTN2	T	C	45330	0.484	-0.043	0.007	6.22E-11	3839	0.214	-0.005	0.027	0.851
sTfR	rs35095338	2:191357694	TMEM194B	A	T	44341	0.556	-0.046	0.007	9.67E-12	3839	0.554	0.029	0.025	0.245
sTfR	rs715	2:211543055	CPS1	T	C	45330	0.688	-0.041	0.007	2.28E-08	3839	0.828	0.011	0.032	0.746
sTfR	rs11371594	3:133539500	SRPRB	G	GA	40837	0.845	0.058	0.01	2.54E-09	n/a	n/a	n/a	n/a	n/a
sTfR	rs112856048	3:195795618	TFRC	A	G	45330	0.243	-0.1	0.008	5.93E-38	3839	0.09	0.058	0.046	0.209
sTfR	rs9325434	3:195921311	ZDHHC19	A	G	45330	0.141	-0.062	0.01	2.27E-10	3839	0.016	0.115	0.151	0.444
sTfR	rs2290846	4:151199080	LRBA	A	G	45330	0.277	-0.041	0.007	3.11E-08	n/a	n/a	n/a	n/a	n/a
sTfR	rs12521868	5:131784393	C5orf56	T	G	45330	0.427	0.052	0.007	4.41E-15	3839	0.029	0	0.074	0.999

sTfR	rs116816795	5:141482333	*NDFIP1*	T	C	45330	0.168	0.127	0.009	3.13E-46	3839	0.028	0.005	0.105	0.965
sTfR	rs2906082	5:141602204	*NDFIP1*	T	C	44584	0.783	0.057	0.008	5.34E-12	3839	0.447	0.034	0.023	0.133
sTfR	rs72832593	6:25957426	*TRIM38*	T	C	45330	0.897	0.13	0.011	8.09E-33	n/a	n/a	n/a	n/a	n/a
sTfR	rs1800562	6:26093141	*HFE*	A	G	45330	0.075	-0.253	0.013	2.88E-88	n/a	n/a	n/a	n/a	n/a
sTfR	rs9389269	6:135427159	*HBS1L*	T	C	45330	0.729	0.059	0.007	3.10E-15	3839	0.961	-0.023	0.078	0.77
sTfR	rs143437464	6:159020121	*TMEM181*	A	G	43595	0.009	-0.234	0.037	3.81E-10	n/a	n/a	n/a	n/a	n/a
sTfR	rs200307986	6:159026327	*TMEM181*	A	G	42671	0.003	0.358	0.065	2.69E-08	n/a	n/a	n/a	n/a	n/a
sTfR	rs186044114	7:1080897	*C7orf50*	T	C	41188	0.852	-0.061	0.01	4.19E-10	n/a	n/a	n/a	n/a	n/a
sTfR	rs6592965	7:50427982	*IKZF1*	A	G	45330	0.458	-0.058	0.007	2.43E-18	3839	0.501	0.013	0.023	0.575
sTfR	rs7385804	7:100235970	*TFR2*	A	C	45330	0.627	-0.051	0.007	1.49E-13	3839	0.726	0	0.026	0.988
sTfR	rs7908745	10:45953767	*MARCH8*	A	G	45330	0.687	-0.054	0.007	5.61E-14	3839	0.677	0.027	0.024	0.262
sTfR	rs16926246	10:71093392	*HK1*	T	C	45330	0.133	0.056	0.01	1.05E-08	3839	0.164	-0.045	0.031	0.143
sTfR	rs187669805	11:117021097	*PCSK7*	C	G	43487	0.994	0.407	0.046	1.94E-18	n/a	n/a	n/a	n/a	n/a
sTfR	rs11216316	11:117081500	*PCSK7*	A	C	45330	0.892	-0.273	0.011	1.77E-137	3839	0.83	-0.041	0.032	0.196
sTfR	rs2238005	11:117088082	*PCSK7*	T	C	45330	0.061	0.118	0.014	2.71E-17	3839	0.031	-0.066	0.086	0.441
sTfR	rs10876169	12:51783420	*GALNT6*	A	T	44584	0.41	0.039	0.007	1.85E-08	3839	0.032	-0.005	0.081	0.955
sTfR	rs76944188	13:110401304	*IRS2*	T	C	45330	0.942	-0.09	0.015	5.45E-10	3839	0.794	0.006	0.028	0.818
sTfR	rs75922593	15:45395901	*DUOX2*	G	GT	40837	0.066	0.112	0.014	3.64E-15	n/a	n/a	n/a	n/a	n/a
sTfR	rs74035509	16:88567333	*ZFPM1*	T	C	44584	0.083	0.083	0.013	6.36E-11	3839	0.15	-0.02	0.033	0.551
sTfR	rs55925547	17:43556807	*PLEKHM1*	T	C	43244	0.801	-0.057	0.009	2.46E-11	3839	0.837	0.034	0.031	0.276
sTfR	rs1292072	17:57925649	*RNU6-450P*	A	G	45330	0.794	-0.048	0.008	7.13E-09	3839	0.945	-0.01	0.048	0.828
sTfR	rs1976703	17:76401328	*PGS1*	T	C	41826	0.493	0.039	0.007	1.57E-08	3839	0.3	-0.006	0.025	0.793
sTfR	rs13041	19:4502282	*PLIN4*	T	C	44584	0.518	0.048	0.007	8.53E-12	3839	0.332	-0.014	0.025	0.591
sTfR	rs855791	22:37462936	*TMPRSS6*	A	G	45330	0.434	0.134	0.007	2.70E-89	3839	0.077	0.001	0.041	0.973

European ancestry summary statistics were obtained from the most recent and largest GWAS of hepcidin and sTfR by Allara et al 2024.

CHR:BP(hg19), chromosome and human genome build 19 base pair; EA, effect allele; OA, other allele; EAFreq, effect allele frequency; Se, standard error; P, additive model p value; n/a, not available

Supplementary Table 9. Replication of European lead serum iron, transferrin saturation, TIBC and ferritin GWAS SNPs in current study of African children

Trait	SNP	CHR:Hg19BP	Gene	EA	OA	European ancestry				Continental Africa			
						EAFreq	Beta	Se	P	EAFreq	Beta	Se	P
Serum iron	rs35945185	1:66137239	LEPR;PDE4B	G	A	0.3618	0.0308	0.0042	1.54E-13	n/a	n/a	n/a	n/a
Serum iron	rs2228145	1:154426970	IL6R	A	C	0.4010	0.0254	0.0033	1.63E-14	0.0785	-0.0328	0.0580	0.5713
Serum iron	rs61804206	1:161658821	RPL31P11;FCRLA	A	G	0.1416	0.0302	0.0048	4.74E-10	n/a	n/a	n/a	n/a
Serum iron	rs7580634	2:113866854	IL1F10;IL1RN	G	T	0.4638	-0.0202	0.0032	4.74E-10	0.2085	-0.0495	0.0409	0.2265
Serum iron	rs13007705	2:239069196	ERFE	C	T	0.3852	0.0297	0.0033	4.14E-19	n/a	n/a	n/a	n/a
Serum iron	rs12633819	3:66448307	LRIG1	A	G	0.3792	0.0279	0.0034	1.22E-16	0.2913	0.0075	0.0357	0.8345
Serum iron	rs8177245	3:133479230	TF	A	G	0.2878	0.0553	0.0035	1.36E-55	0.1596	0.0253	0.0486	0.6032
Serum iron	rs571559677	3:140011552	CLSTN2	T	C	0.0023	-0.3512	0.0634	3.10E-08	n/a	n/a	n/a	n/a
Serum iron	rs1800562	6:26093141	HFE	G	A	0.0747	0.298	0.0061	7.66e-520	n/a	n/a	n/a	n/a
Serum iron	rs9399136	6:135402339	HBS1L;MYB	T	C	0.2708	0.0553	0.0037	8.93E-52	0.0208	n/a	n/a	n/a
Serum iron	rs12718598	7:50428445	IKZF1	T	C	0.4493	0.0229	0.0033	2.07E-12	0.5190	0.0331	0.0331	0.3168
Serum iron	rs2075672	7:100240296	ACTL6B	A	G	0.6288	0.0587	0.0033	6.04E-69	0.6881	0.0475	0.0349	0.1734
Serum iron	rs10740134	10:65315433	REEP3	T	C	0.5129	0.0211	0.0032	7.78E-11	0.2932	0.0376	0.0431	0.3827
Serum iron	rs59213222	10:81073027	ZMIZ1	CCCAC CCACCA	C	0.2231	-0.0264	0.0046	9.33E-09	n/a	n/a	n/a	n/a
Serum iron	rs2900478	12:21368797	SLCO1B1	T	A	0.1783	-0.0233	0.0042	3.86E-08	0.0205	n/a	n/a	n/a
Serum iron	rs4759827	12:131492288	ADGRD1	C	T	0.2793	-0.0209	0.0037	1.54E-08	0.2157	-0.0728	0.0454	0.1089
Serum iron	rs1958078	14:70354858	SMOC1	A	C	0.8647	0.0357	0.0048	9.96E-14	0.4931	0.0607	0.0394	0.1238
Serum iron	rs28929474	14:94844947	SERPINA1	C	T	0.0131	0.1046	0.0136	1.55E-14	n/a	n/a	n/a	n/a
Serum iron	rs374974760	15:45394467	DUOX2	G	GGT	0.0823	-0.0494	0.0081	1.32E-09	n/a	n/a	n/a	n/a
Serum iron	rs77542162	17:67081278	ABCA6	A	G	0.0228	0.0843	0.0114	1.28E-13	n/a	n/a	n/a	n/a
Serum iron	rs12975762	19:35950523	FFAR2;KRTDAP	A	G	0.3161	-0.0265	0.0035	4.26E-14	0.4589	0.0166	0.0339	0.6238
Serum iron	rs1800961	20:43042364	HNF4A	C	T	0.0472	0.0447	0.008	2.35E-08	n/a	n/a	n/a	n/a
Serum iron	rs855791	22:37462936	TMPRSS6	A	G	0.5805	0.1723	0.0035	8.00e-516	0.0768	0.0821	0.0629	0.1920
Serum iron	rs3747207	22:44324855	PNPLA3	G	A	0.2335	0.0271	0.0038	1.67E-12	0.1063	0.0884	0.0538	0.1004
TSAT	rs35945185	1:66137239	LEPR;PDE4B	G	A	0.3618	0.0268	0.0047	1.34E-08	n/a	n/a	n/a	n/a
TSAT	rs4133213	1:154395212	IL6R	C	A	0.4295	0.022	0.0037	2.26E-09	0.1943	0.0205	0.0484	0.6714
TSAT	rs6025	1:169519049	F5	C	T	0.0260	0.0727	0.0116	4.35E-10	n/a	n/a	n/a	n/a
TSAT	rs13007705	2:239069196	ERFE	C	T	0.3852	0.0334	0.0037	1.59E-19	n/a	n/a	n/a	n/a
TSAT	rs12633819	3:66448307	LRIG1	A	G	0.3792	0.0288	0.0037	1.42E-14	0.2913	0.0572	0.0361	0.1132
TSAT	rs8177253	3:133480192	TF	C	T	0.2890	-0.0915	0.0039	1.03E-122	0.2491	-0.0389	0.0385	0.3121
TSAT	rs41300395	3:195809135	TFRC	C	A	0.2457	0.0358	0.0042	1.36E-17	0.0902	-0.0379	0.0619	0.5398
TSAT	rs218264	4:55408875	LINC02283;LINC02260	A	T	0.2501	0.0229	0.0042	4.17E-08	0.6821	-0.0011	0.0370	0.9772
TSAT	rs79220007	6:26098474	HFE	T	C	0.0791	0.477	0.0078	3.90e-810	n/a	n/a	n/a	n/a
TSAT	rs9399136	6:135402339	HBS1L;MYB	T	C	0.2708	0.0653	0.004	6.29E-59	0.0208	n/a	n/a	n/a
TSAT	rs12718598	7:50428445	IKZF1	T	C	0.4493	0.0234	0.0036	6.54E-11	0.5190	0.0351	0.0334	0.2936
TSAT	rs2075672	7:100240296	ACTL6B	A	G	0.6288	0.0622	0.0037	1.67E-63	0.6881	0.0205	0.0353	0.5609
TSAT	rs112875651	8:126506694	TRIB1;LINC00861	G	A	0.3935	-0.0234	0.0037	2.98E-10	0.1512	0.0118	0.0475	0.8045
TSAT	rs10740134	10:65315433	REEP3	T	C	0.5129	0.0221	0.0036	8.91E-10	0.2932	0.0119	0.0432	0.7826
TSAT	rs2900478	12:21368797	SLCO1B1	T	A	0.1783	-0.0303	0.0047	1.38E-10	0.0205	n/a	n/a	n/a
TSAT	rs1958078	14:70354858	SMOC1	A	C	0.8647	0.0361	0.0054	1.72E-11	0.4931	0.0407	0.0395	0.3022
TSAT	rs7165401	15:45229327	SORD2P;TERB2	T	C	0.0941	-0.0541	0.0073	1.61E-13	0.9004	-0.0447	0.0547	0.4144
TSAT	rs77542162	17:67081278	ABCA6	A	G	0.0228	0.0698	0.0127	3.73E-08	n/a	n/a	n/a	n/a
TSAT	rs36152188	19:35949423	FFAR2;KRTDAP	A	T	0.3130	-0.0297	0.0039	3.50E-14	0.1985	-0.0401	0.0401	0.3170
TSAT	rs855791	22:37462936	TMPRSS6	A	G	0.5805	0.1748	0.0036	5.56e-510	0.0768	0.1552	0.0636	**0.0147**
TSAT	rs3747207	22:44324855	PNPLA3	G	A	0.2335	0.0244	0.0043	1.11E-08	0.1063	0.0582	0.0541	0.2824

TIBC	rs514595	1:46036960	*AKR1A1;NASP*	T	C	0.8451	0.0308	0.005	6.51E-10	0.7083	0.0229	0.0243	0.3454
TIBC	rs469721	1:91530001	*ZNF644;HFM1*	C	T	0.2129	-0.0359	0.0045	1.02E-15	0.1605	0.0559	0.0352	0.1121
TIBC	rs137891701	1:161468310	*SDHC;FCGR2A*	CTAACCACATTTTT	C	0.1291	0.0376	0.0064	4.86E-09	n/a	n/a	n/a	n/a
TIBC	rs6025	1:169519049	*F5*	C	T	0.0260	-0.1052	0.0119	1.21E-18	n/a	n/a	n/a	n/a
TIBC	rs61830291	1:221001142	*RNU6ATAC35P;LINC01352*	A	C	0.0992	0.0364	0.0062	4.52E-09	n/a	n/a	n/a	n/a
TIBC	rs1435167	1:231502244	*EGLN1*	T	A	0.6273	-0.0244	0.0038	1.57E-10	n/a	n/a	n/a	n/a
TIBC	rs11336847	2:27749001	*GCKR;C2orf16*	AT	A	0.6644	-0.0345	0.0046	3.86E-14	n/a	n/a	n/a	n/a
TIBC	rs11900910	2:46557600	*EPAS1*	C	T	0.4508	0.0247	0.0037	2.08E-11	0.4412	-0.0066	0.0232	0.7756
TIBC	rs1861408	2:64924037	*SERTAD2;LINC01800*	C	T	0.7034	0.0249	0.004	7.84E-10	0.5667	0.0016	0.0232	0.9467
TIBC	rs13008704	2:190387487	*WDR75;SLC40A1*	C	T	0.5075	0.0411	0.0037	5.49E-29	0.0770	0.0610	0.0477	0.2007
TIBC	rs715	2:211543055	*CPS1*	T	C	0.2852	0.0249	0.004	4.50E-10	0.1715	-0.0026	0.0324	0.9360
TIBC	rs781168893	3:124932695	*SLC12A8;ZNF148*	C	T	0.0026	-0.7468	0.0641	2.07E-31	n/a	n/a	n/a	n/a
TIBC	rs8177247	3:133479499	*TF*	T	G	0.3049	0.3665	0.0051	7.73e-1122	0.2279	0.2759	0.0277	**2.66E-23**
TIBC	rs760897340	3:145423900	*DIPK2A;LNCSRLR*	C	G	0.0011	-0.9733	0.1039	7.54E-21	n/a	n/a	n/a	n/a
TIBC	rs969672878	3:147672687	*LOC440982;LINC02032*	G	A	0.0011	-0.9008	0.103	2.29E-18	n/a	n/a	n/a	n/a
TIBC	rs11915082	3:195809139	*TFRC*	G	A	0.3886	0.0379	0.0037	1.30E-24	0.1449	0.0096	0.0358	0.7891
TIBC	rs11947277	4:3473751	*DOK7*	C	T	0.2661	0.0346	0.0041	5.16E-17	0.2925	0.0079	0.0254	0.7546
TIBC	rs185240714	5:52095934	*PELO*	C	A	0.0403	-0.0685	0.0094	2.38E-13	0.0184	0.0069	0.1139	0.9514
TIBC	rs1800562	6:26093141	*HFE*	G	A	0.0747	-0.4831	0.0079	6.94e-807	n/a	n/a	n/a	n/a
TIBC	rs56293029	6:135419039	*HBS1L;MYB*	C	A	0.2743	-0.0373	0.0043	9.21E-18	0.2291	-0.0328	0.0305	0.2819
TIBC	rs9987289	8:9183358	*LOC157273*	A	G	0.9050	0.0369	0.0063	3.94E-09	0.8067	0.0118	0.0290	0.6857
TIBC	rs1495743	8:18273300	*NAT2;PSD3*	G	C	0.7983	0.0494	0.0045	2.29E-28	n/a	n/a	n/a	n/a
TIBC	rs28601761	8:126500031	*TRIB1;LINC00861*	C	G	0.4215	0.0258	0.0037	3.09E-12	0.3226	0.0239	0.0239	0.3159
TIBC	rs41274050	10:52573772	*A1CF*	C	T	0.0081	0.1296	0.0221	4.83E-09	n/a	n/a	n/a	n/a
TIBC	rs174547	11:61570783	*FADS1*	T	C	0.3733	0.0492	0.0038	4.10E-39	0.0186	0.1171	0.1048	0.2642
TIBC	rs143878994	11:110468861	*ARHGAP20*	G	A	0.0460	0.0488	0.0089	3.63E-08	n/a	n/a	n/a	n/a
TIBC	rs76895963	12:4384844	*CCND2-AS1*	T	G	0.0165	-0.0808	0.0147	4.15E-08	n/a	n/a	n/a	n/a
TIBC	rs3782735	12:6885076	*LAG3*	G	A	0.5981	0.0232	0.0039	2.37E-09	0.8278	-0.0409	0.0300	0.1728
TIBC	rs1150975	12:32052422	*LINC02422;RESF1*	C	A	0.3474	0.0212	0.0039	4.08E-08	0.0895	-0.0082	0.0381	0.8289
TIBC	rs77371258	12:121029603	*POP5;CABP1*	C	A	0.0987	0.0437	0.006	3.53E-13	0.1971	0.0240	0.0284	0.3968
TIBC	rs1680695	14:34408083	*EGLN3*	T	G	0.3547	-0.0247	0.0038	1.16E-10	0.0717	-0.0356	0.0541	0.5105
TIBC	rs12879801	14:70770582	*ADAM21P1;SYNJ2BP-COX16*	A	G	0.4522	0.0311	0.0036	1.48E-17	0.2038	-0.0093	0.0286	0.7451
TIBC	rs11159099	14:75235397	*YLPM1*	A	G	0.4817	-0.0295	0.0036	5.22E-16	0.0345	-0.0876	0.0757	0.2477
TIBC	rs17580	14:94847262	*SERPINA1*	T	A	0.0334	0.0631	0.0097	9.79E-11	n/a	n/a	n/a	n/a
TIBC	rs4775744	15:45385496	*DUOX2*	A	T	0.0903	0.0596	0.0064	1.97E-20	n/a	n/a	n/a	n/a
TIBC	rs8062982	16:53375645	*CHD9;LOC643802*	G	C	0.4012	0.0256	0.0037	6.65E-12	0.6313	0.0303	0.0246	0.2185
TIBC	rs34682685	16:72096227	*HPR*	G	A	0.1003	0.0425	0.006	1.20E-12	n/a	n/a	n/a	n/a
TIBC	rs17767742	16:79749005	*MAF;MAFTRR*	C	G	0.3087	0.0272	0.0039	3.11E-12	n/a	n/a	n/a	n/a
TIBC	rs2287322	17:1641035	*WDR81*	A	G	0.2389	0.0309	0.0043	8.92E-13	0.2454	0.0548	0.0267	**0.0398**
TIBC	rs1292072	17:57925649	*VMP1;TUBD1*	A	G	0.2239	0.0289	0.0044	5.03E-11	0.0551	0.0133	0.0478	0.7800
TIBC	rs34284056	18:60203855	*ZCCHC2*	C	A	0.2635	-0.0232	0.0041	1.62E-08	0.0236	0.2139	0.0974	**0.0280**
TIBC	rs111234557	19:19436229	*MAU2*	C	G	0.1212	-0.033	0.0056	4.03E-09	0.0187	-0.0501	0.1115	0.6531
TIBC	rs1672992	19:35556729	*HPN-AS1*	C	T	0.8484	0.0356	0.0053	1.38E-11	0.3473	0.0087	0.0241	0.7171
TIBC	rs12976652	19:46396610	*MYPOP*	T	C	0.1046	-0.05	0.0058	6.52E-18	0.0338	0.0612	0.0698	0.3807
TIBC	rs111981233	19:50016479	*FCGRT*	T	G	0.0651	0.0736	0.0071	1.82E-25	0.0237	0.1439	0.0865	0.0964
TIBC	rs2236252	20:17597531	*RRBP1*	C	T	0.1762	0.0394	0.0048	1.99E-16	0.0314	-0.0331	0.0747	0.6576
TIBC	rs1800961	20:43042364	*HNF4A*	C	T	0.0472	0.0534	0.0091	4.00E-09	n/a	n/a	n/a	n/a
TIBC	rs855791	22:37462936	*TMPRSS6*	A	G	0.5805	-0.0335	0.0036	4.80E-20	0.0768	-0.0161	0.0408	0.6926

Ferritin	rs477190	1:20898514	FAM43B;CDA	C	A	0.4042	-0.0192	0.0034	1.26E-08	0.0632	-0.0481	0.0456	0.2918
Ferritin	rs79052526	1:22572707	WNT4;MIR4418	G	A	0.0267	-0.1229	0.0112	3.83E-28	n/a	n/a	n/a	n/a
Ferritin	rs10801913	1:116214279	VANGL1	A	G	0.6979	-0.0222	0.0036	9.06E-10	0.5331	0.0084	0.0234	0.7188
Ferritin	rs6025	1:169519049	F5	C	T	0.0260	0.1558	0.011	3.44E-45	n/a	n/a	n/a	n/a
Ferritin	rs551459670	1:220288690	IARS2	G	A	0.0102	0.1407	0.019	1.28E-13	n/a	n/a	n/a	n/a
Ferritin	rs1260326	2:27730940	GCKR	T	C	0.6544	-0.0234	0.0035	1.57E-11	0.0888	0.0053	0.0393	0.8925
Ferritin	rs6760824	2:29178319	WDR43;TOGARAM2	A	C	0.7529	-0.0311	0.0038	2.17E-16	0.6571	-0.0407	0.0239	0.0894
Ferritin	rs604126	2:111943200	BCL2L11;MIR4435-2HG	A	G	0.6856	0.02	0.0036	2.73E-08	0.9474	-0.0696	0.0591	0.2391
Ferritin	rs17050272	2:121306440	LINC01101;GLI2	G	A	0.4166	0.0209	0.0034	1.07E-09	0.0489	0.0359	0.0523	0.4931
Ferritin	rs17707216	2:188201256	ZSWIM2;CALCRL	G	A	0.0758	0.0381	0.0068	2.22E-08	n/a	n/a	n/a	n/a
Ferritin	rs7596205	2:190428193	SLC40A1	G	A	0.1021	0.0838	0.0055	1.32E-51	n/a	0.1262	0.2245	0.5740
Ferritin	rs1250244	2:216297796	FN1	G	C	0.6572	0.0254	0.0036	2.29E-12	0.8092	0.0247	0.0348	0.4778
Ferritin	rs553656123	3:47033788	NBEAL2	G	A	0.0080	0.1135	0.0199	1.09E-08	n/a	n/a	n/a	n/a
Ferritin	rs142350264	3:49594992	BSN	T	TGGA	0.0983	-0.0403	0.0067	1.73E-09	n/a	n/a	n/a	n/a
Ferritin	rs536826368	3:52453040	PHF7	C	T	0.0047	0.2905	0.0324	3.39E-19	n/a	n/a	n/a	n/a
Ferritin	rs533732725	3:54526076	CACNA2D3	C	G	0.0032	0.2602	0.0386	1.56E-11	n/a	n/a	n/a	n/a
Ferritin	rs10804630	3:133997989	RYK;LINC02004	T	C	0.1026	-0.0301	0.0055	4.15E-08	0.2060	-0.0329	0.0281	0.2403
Ferritin	rs6822746	4:56257107	SRD5A3-AS1;TMEM165	G	A	0.3165	-0.0206	0.0035	6.32E-09	0.1854	-0.0044	0.0297	0.8813
Ferritin	rs4865796	5:53272664	ARL15	G	A	0.6813	0.0207	0.0036	6.38E-09	0.7668	-0.0127	0.0274	0.6424
Ferritin	rs79694859	5:59642035	PDE4D	G	A	0.0456	0.0514	0.0087	3.23E-09	n/a	n/a	n/a	n/a
Ferritin	rs35107257	5:68805616	OCLN	G	A	0.0581	0.0437	0.0076	1.08E-08	n/a	n/a	n/a	n/a
Ferritin	rs970079	5:98119864	RGMB	A	G	0.4630	-0.0193	0.0033	7.00E-09	0.9380	-0.0095	0.0513	0.8527
Ferritin	rs144861591	6:26072992	HIST1H1C;LOC108783645	C	T	0.0669	0.1292	0.0066	6.76E-85	n/a	n/a	n/a	n/a
Ferritin	rs35486885	6:32763506	HLA-DQB2;HLA-DOB	A	G	0.0377	-0.0561	0.0081	5.25E-12	n/a	n/a	n/a	n/a
Ferritin	rs13215052	6:41681835	TFEB	G	A	0.1113	-0.0317	0.0054	5.74E-09	0.0286	-0.1510	0.1003	0.1322
Ferritin	rs36184164	6:43781092	VEGFA;LINC02537	T	G	0.1386	0.0356	0.0051	2.07E-12	n/a	n/a	n/a	n/a
Ferritin	rs2529440	7:30511794	NOD1	C	T	0.4496	-0.0347	0.0034	1.02E-24	0.9556	0.0613	0.0537	0.2534
Ferritin	rs4841429	8:10568529	C8orf74;SOX7	A	G	0.0788	0.0562	0.0062	1.59E-19	0.5762	0.0124	0.0241	0.6081
Ferritin	rs7009973	8:23374454	ENTPD4;SLC25A37	A	G	0.3378	0.0223	0.0036	3.74E-10	0.3167	-0.0063	0.0248	0.7993
Ferritin	rs189899297	8:87173917	ATP6V0D2;SLC7A13	T	A	0.2634	0.0214	0.0039	4.67E-08	n/a	n/a	n/a	n/a
Ferritin	rs681099	8:109218892	EIF3E	T	C	0.4547	-0.0189	0.0034	1.68E-08	0.5907	-0.0054	0.0253	0.8314
Ferritin	rs2954027	8:126485294	TRIB1;LINC00861	T	A	0.4896	-0.0235	0.0033	1.71E-12	0.5458	0.0382	0.0232	0.0996
Ferritin	rs7865362	9:33117965	B4GALT1	C	T	0.3468	0.0251	0.0035	7.47E-13	0.6516	-0.0035	0.0238	0.8840
Ferritin	9:136139907	9:136139907	ABO	G	GAA ACTGCC	#N/A	0.0617	0.0048	2.48E-37	n/a	n/a	n/a	n/a
Ferritin	rs72775768	9:139324574	INPP5E	C	T	0.2586	-0.022	0.0037	4.18E-09	0.2941	-0.0246	0.0251	0.3274
Ferritin	rs9423600	10:5395670	LINC02561;UCN3	A	G	0.2675	0.0207	0.0038	4.82E-08	0.2779	-0.0202	0.0255	0.4283
Ferritin	rs17476364	10:71094504	HK1	T	C	0.0947	0.0421	0.0055	2.32E-14	n/a	n/a	n/a	n/a
Ferritin	rs704017	10:80819132	ZMIZ1-AS1	A	G	0.6224	-0.019	0.0034	2.61E-08	0.5267	0.0044	0.0222	0.8423
Ferritin	rs17112021	10:101343726	NKX2-3;SLC25A28	T	G	0.2754	0.0245	0.0038	1.97E-10	0.1629	0.0160	0.0299	0.5916
Ferritin	rs12419620	11:2232553	MIR4686;ASCL2	T	G	0.1623	-0.0313	0.0045	5.65E-12	0.3094	0.0133	0.0252	0.5984
Ferritin	rs7102016	11:10127309	SBF2	C	T	0.5559	0.0184	0.0033	3.36E-08	0.4582	0.0071	0.0231	0.7597
Ferritin	rs12807014	11:47760078	FNBP4	T	C	0.2462	-0.0283	0.0038	7.34E-14	0.0133	n/a	n/a	n/a
Ferritin	rs4938939	11:60160838	MS4A7	G	A	0.2926	0.0221	0.0036	1.39E-09	0.3062	0.0320	0.0245	0.1926
Ferritin	rs10750215	11:122505344	MIR100HG;UBASH3B	G	T	0.4173	0.0211	0.0037	1.46E-08	0.2937	0.0322	0.0250	0.1973
Ferritin	rs161044	12:51382232	SLC11A2	T	C	0.9572	-0.0743	0.0078	1.14E-21	0.8033	-0.0083	0.0287	0.7720
Ferritin	rs9512463	13:27528519	GPR12;USP12	T	C	0.7745	-0.0237	0.004	4.77E-09	0.2524	-0.0139	0.0259	0.5907
Ferritin	rs28715334	14:23635213	SLC7A8	G	T	0.2029	0.0244	0.0042	6.44E-09	0.4323	0.0003	0.0226	0.9879
Ferritin	rs996347	14:34410892	EGLN3	T	C	0.3411	0.0492	0.0035	9.04E-45	0.1506	0.0331	0.0324	0.3061
Ferritin	rs764195359	15:41974309	MGA	T	G	0.0020	-0.3016	0.0506	2.57E-09	n/a	n/a	n/a	n/a

				EA	OA	EAFreq		Se	P				TSAT
Ferritin	rs10685744	15:45378096	*SORD;DUOX2*	G	GTAAA	0.0813	-0.1472	0.0065	8.22E-113	n/a	n/a	n/a	n/a
Ferritin	rs12917564	15:65923746	*SLC24A1*	G	C	0.2064	-0.0242	0.0042	7.26E-09	0.1607	-0.0004	0.0315	0.9904
Ferritin	rs11634990	15:78771186	*IREB2*	T	C	0.1701	-0.0268	0.0045	2.00E-09	0.1215	-0.0338	0.0336	0.3152
Ferritin	rs8047501	16:392318	*AXIN1*	A	G	0.5974	-0.0257	0.0034	5.18E-14	0.8142	-0.0281	0.0300	0.3481
Ferritin	rs3747602	16:4802386	*ZNF500*	G	T	0.6336	-0.019	0.0035	3.76E-08	0.4233	-0.0157	0.0247	0.5251
Ferritin	rs186021206	17:7069412	*ASGR2;ASGR1*	G	A	0.0047	0.2105	0.0243	4.30E-18	n/a	n/a	n/a	n/a
Ferritin	rs55789050	17:9793417	*GLP2R*	A	T	0.3541	-0.0264	0.0035	6.77E-14	0.0202	0.1895	0.1087	0.0812
Ferritin	rs62074125	17:44852612	*WNT3*	A	C	0.2884	0.0208	0.0038	3.75E-08	n/a	n/a	n/a	n/a
Ferritin	rs34523089	17:56436109	*RNF43*	C	T	0.1636	0.0662	0.0045	4.52E-48	n/a	n/a	n/a	n/a
Ferritin	rs2952290	17:66463383	*PRKAR1A*	G	A	0.7864	-0.0228	0.0041	1.91E-08	0.7375	-0.0786	0.0259	**0.0024**
Ferritin	rs4789111	17:72940481	*OTOP3*	T	C	0.8527	-0.0331	0.0047	1.28E-12	0.8089	0.0470	0.0321	0.1427
Ferritin	rs708686	19:5840619	*FUT6*	C	T	0.2346	-0.0312	0.0039	6.79E-16	0.8518	0.0036	0.0319	0.9103
Ferritin	rs141253118	19:15314444	*NOTCH3;EPHX3*	G	A	0.0045	-0.1245	0.0228	4.60E-08	n/a	n/a	n/a	n/a
Ferritin	rs4808802	19:18577873	*ELL*	G	C	0.2291	0.0272	0.004	1.35E-11	0.5434	-0.0208	0.0231	0.3680
Ferritin	rs192331981	19:49431144	*NUCB1;DHDH*	G	T	0.0169	0.1968	0.0138	2.11E-46	n/a	n/a	n/a	n/a
Ferritin	rs2595586	20:3607469	*ATRN*	A	G	0.4135	-0.019	0.0034	2.20E-08	n/a	n/a	n/a	n/a
Ferritin	rs6088374	20:30079334	*LINC00028;HM13*	T	C	0.1690	-0.0374	0.0045	1.94E-16	0.7966	0.0482	0.0279	0.0839
Ferritin	rs56206139	20:39108705	*LINC01370;MAFB*	A	C	0.0782	0.0465	0.0064	3.33E-13	0.1132	-0.0030	0.0351	0.9329
Ferritin	rs540828131	22:35782508	*HMOX1*	A	G	0.0069	-0.1595	0.0263	1.33E-09	n/a	n/a	n/a	n/a
Ferritin	rs2076085	22:37470041	*TMPRSS6*	C	A	0.5716	0.0413	0.0034	1.04E-34	0.7547	-0.0365	0.0264	0.1669

CHR:BP(hg19), chromosome and human genome build 19 base pair; EA, effect allele; OA, other allele; EAFreq, effect allele frequency; Se, standard error; P, additive model p value; TSAT, transferrin saturation; n/a, not available

Transferrin iron binding capacity, TIBC (reported by Moksnes et al 2022) is compared with transferrin from current study. Transferrin (g/L) = 0.7 x TIBC [μg/dL] x 0.01.

Supplementary Table 10. Laboratory assays for iron and inflammatory biomarkers by site

Study site	Transferrin	Ferritin	sTfR	Hepcidin	Iron	UIBC	TIBC	CRP	ACT
Kilfi, Kenya	Chemiluminescent Microparticle Immunoassay (Abbott Architect, USA)	Microparticle Enzyme Immunoassay (Abbott Architect, USA)	Human sTfR ELISA (BioVendor, Czech Republic)	DRG Hepcidin 25 [bioactive] high sensitive ELISA (DRG International, USA)	MULTIGENT iron calorimetric assay, Abbott Architect, USA	Not measured	Not measured	MULTIGENT CRP Vario assay, (Abbott Architect, USA)	Not measured
Banfora,Burkina Faso	Chemiluminescent Microparticle Immunoassay (Abbott Architect, USA)	Microparticle Enzyme Immunoassay (Abbott Architect, USA)	Human sTfR ELISA (BioVendor, Czech Republic)	DRG Hepcidin 25 [bioactive] high sensitive ELISA (DRG International, USA)	MULTIGENT iron calorimetric assay, Abbott Architect, USA	Not measured	Not measured	MULTIGENT CRP Vario assay, (Abbott Architect, USA)	Not measured
Entebbe, Uganda	Chemiluminescent Microparticle Immunoassay (Abbott Architect, USA)	Microparticle Enzyme Immunoassay (Abbott Architect, USA)	Human sTfR ELISA (BioVendor, Czech Republic)	DRG Hepcidin 25 [bioactive] high sensitive ELISA (DRG International, USA)	Not measured	Not measured	Not measured	MULTIGENT CRP Vario assay, (Abbott Architect, USA)	Not measured
Soweto, South Africa	Chemiluminescent Microparticle Immunoassay (Abbott Architect, USA)	Microparticle Enzyme Immunoassay (Abbott Architect, USA)	Human sTfR ELISA (BioVendor, Czech Republic)	DRG Hepcidin 25 [bioactive] high sensitive ELISA (DRG International, USA)	Not measured	Not measured	Not measured	MULTIGENT CRP Vario assay, (Abbott Architect, USA)	Not measured
West Kiang, The Gambia	Not measured	Microparticle Enzyme Immunoassay (Abbott Architect, USA)	Quantikine sTfR ELISA kit, (R&D Systems, USA)	Hepcidin-25 [human] Enzyme Immunoassay Kit (Bachem, Switzerland)	Ferrozine-based photometry and colorimetry analyser (Hitachi 911, Hitachi, Tokyo, Japan)	Ferrozine-based photometry and colorimetry analyser (Hitachi 911, Hitachi, Tokyo, Japan)	Not measured	Not measured	Immunoturbidimetry, Cobas Mira Plus Bio-analyser, Roche
Jackson Heart Study	Somascan	Roche immunoturbimetric assay	Not measured	Somascan	Ferrozine colorimetric assay (Roche)	Not measured	Ferrozine colorimetric assay (Roche)	ELISA	Somascan

sTfR, soluble transferrin receptor; UIBC, unsaturated iron binding capacity; TIBC, total iron binding capacity; CRP, C-reative protein; ACT, α1-antichymotrypsin