Gender inequities regarding the HPV vaccination coverage in high-income countries: a systematic review and meta-analysis

Sabeena S[1]*, Nagaraja Ravishankar[2]

Allure residency, Near the British School, Kathmandu, Bagmati Province, Nepal[1], Vallabhbhai Patel Chest Institute, University of Delhi, New Delhi[2]

Short title: Gender inequities in the HPV vaccination coverage

Affiliations

1 Dr Sasidharanpillai Sabeena[1], MBBS, DGO, PhD (Virology)

Clinical Virologist

Allure Residency,
Near the British School,
Kathmandu,

Bagmati Province,

Nepal.

Postal Code: 44600

Phone: +977-9808187821

Email: sabeenauthradam**@gmail.com**

ORCID id: https://orcid.org/ 0000-0002-2099-9312

2 Nagaraja Ravishankar [2]

MSc, PhD

Assistant Professor

Department of Biostatistics

Vallabhbhai Patel Chest Institute

University of Delhi

New Delhi, Delhi 110007,

India

Email:ravi8133@gmail.com

ORCID id: https://orcid.org/0000-0002-9533-039X

Corresponding author:

Dr Sasidharanpillai Sabeena[1], MBBS, DGO, PhD

Clinical Virologist

Allure Residency,

Near the British School,

Kathmandu,

Bagmati Province,

Nepal.

Postal Code: +44600

Email:sabeenauthradam@**gmail.com**

ORCID id: **https://orcid.org/ 0000-0002-2099-9312**

Gender inequities regarding the HPV vaccination coverage in high-income countries: a systematic review and meta-analysis

Abstract

Introduction

Globally, selected high-income countries have implemented the gender neutral HPV vaccination. Universal HPV vaccine approach promotes vaccine acceptance, minimizes vaccine misinformation, and reduces vaccine-related stigma.

Methods

An electronic search of the PubMed/MEDLINE, Scopus, Google Scholar, and official websites was carried out for articles reporting the HPV vaccination coverage in high income countries. Meta-analysis was performed in STATA version 13.0 (College Station, TX, USA). The forest plots were made using the midas package in STATA.

Results

The pooled prevalence of the adolescent HPV vaccine uptake among adolescent girls was 67.75% (95% CI: 62.81%,72.69%) and among teenage boys was 58.32% (95% CI 48.88%, 67.76%). The pooled prevalence among men having sex with men (MSM) was 27.94% (95% CI 14.35, 41.53) and the pooled prevalence in transgender women was 32.59% (95% CI 8.35%. 56.83%)

Conclusion

Significant gender inequities exist in the HPV vaccination coverage and less number of gender minorities were vaccinated in high income countries.

Keywords: adolescents; Human Papillomavirus Vaccine; HPV vaccine; gender diverse; gender minorities; transgender persons.

Introduction

Human Papillomavirus (HPV) infection is one of the most common sexually transmitted infections [1]. World Health Organization (WHO) has recommended the HPV vaccination coverage of 90% among adolescent girls before the age of 15 years, screening of 70% of women by 35 years, and management of cervical pre-invasive lesions and cervical cancers in 90% of women for global elimination of cervical cancer [2]. However, a persistent rise in HPV-associated oropharyngeal squamous cell cancers (OPSCC) among men has been reported in high-income countries over the past few decades [3]. Gender neutral or universal HPV vaccination contributes to higher herd immunity. Gender minorities also should be vaccinated against this oncogenic virus to reduce the global burden of HPV-associated cancers.

World Health Organization has listed the HPV vaccine as an essential medicine [4]. Currently, forty-seven countries such as Australia, Canada, France, Ireland, Hong Kong, Portugal, South Korea, Switzerland, the United Kingdom, and the United States of America have introduced gender neutral HPV vaccination in their national immunization programs. The updated data regarding the human papillomavirus (HPV) vaccination coverage among adolescents, men having sex with men (MSM), and transgender women (TGW) are essential to monitor the HPV vaccination programs and analyse the impact of universal HPV vaccination on HPV-related disease burden.

HPV is a gender-neutral sexually transmitted infection affecting all genders [5]. Even though many high-income countries have implemented the HPV vaccination program for women, gender-neutral HPV vaccine coverage is negligible [6]. HPV vaccination programs extending to adolescent boys and catch-up vaccination of men up to 26 years is more cost-effective than vaccinating only adolescent girls and women [7,8]. The countries implementing gender-neutral HPV vaccination are high-income or upper middle income countries except Bhutan

[9]. The HPV vaccination coverage is suboptimal and varies widely depending on the gender identity or sexual orientation in high income countries.

The gender neutral HPV vaccination approach ensures gender equity as gender diverse populations such as TGW and MSM are at risk of persistent HPV infection. There is sparse information regarding the HPV vaccination uptake among gender minorities. They often encounter distinct barriers to healthcare and preventive vaccination services. A recent US study reported higher prevalence of HPV infection among transgender women (TGW) than among men having sex with men (MSM) [10]. The reduced HPV vaccine uptake among sexual minorities was attributed to the low awareness among healthcare professionals, lack of recommendations by clinicians, and public mistrust of healthcare systems [11]. This systematic review aimed to analyse the studies from high income countries reporting the HPV vaccine uptake among adolescents, adult transgender women, and MSM up to the age of 26 years.

Description of condition

HPV vaccine coverage: As per the WHO definition, HPV vaccine coverage is defined as the percentage of the target population (9-14 year old girls) who have received the recommended doses of HPV vaccine in a given year [12]. The uptake of single dose HPV vaccine was considered complete coverage based on the revised HPV vaccination recommendation in 2023 [13]. The US Advisory Committee on Immunization Practices (ACIP) has recommended gender neutral HPV vaccination across all genders up to the age of 26 [14]. The vaccine effectiveness decreases with increasing age.

HPV vaccine uptake: The HPV vaccine uptake indicates at least one dose of the HPV vaccine in adolescents, transgender persons, and MSM up to 26 years.

Transgender person: Transgender person is an "umbrella term" with no accepted definition and describes an individual whose sexual identity differs from the sex assigned at birth [15].

2. Methods

The review was started after ruling out any registered or ongoing systematic reviews concerning gender inequities of the HPV vaccination coverage in PROSPERO database.

The study protocol was registered in the PROSPERO database (CRD42025641689) and may be accessed through the link https: // www. crd. york. ac. uk / prospero / display_record. php? Record ID= 641689. An electronic search of the PubMed/MEDLINE, Scopus, Google Scholar, and official websites was carried out. Articles published in English between January 2010 and January 2025 regarding the gender disparity in HPV vaccination coverage were screened. Relevant articles were identified electronically using specific search terms and related articles in peer-reviewed journals were manually searched. The references of retrieved articles were reviewed and the last date of the search was noted (**15th February 2024**).

Observational studies reporting the HPV vaccination coverage in adolescents and gender minorities were screened. The study population included adolescents, TGW, and MSM aged 18-26 years. Studies reporting HPV vaccination coverage only among girls or boys from a specific region were not included. Cost-effectiveness studies, modelling studies, studies carried out among intellectually disabled adolescents, and migrant adolescents were excluded. Studies carried out among adults above 26 years were also not included.

Data extraction

PubMed/MEDLINE, Scopus, and Google Scholar were searched for the articles published between January 2010 and 2015. The relevant articles in English were screened using search terms such as "adolescents" AND "Human Papillomavirus vaccines" OR "HPV vaccines" AND transgender women" AND "MSM" OR "men having sex with men" AND "high income countries".

A three-stage selection process was carried out for the final inclusion. One reviewer assessed the article titles and appropriate studies were moved to the second stage after eliminating irrelevant topics. In the second stage, the abstracts of the studies were independently assessed. After reviewing the abstracts, the full texts of the studies were retrieved which were scrutinised by two reviewers independently. The corresponding authors were contacted electronically if further interpretation was required. A manual library search for articles in peer-reviewed journals was carried out and references of retrieved articles were also reviewed to increase the search sensitivity. The study selection process was demonstrated in the PRISMA chart (**Figure 1**). A validated proforma including the first author, year of publication, region, study design, study population, sample size, gender, study period, HPV vaccination coverage, and quality of the studies was prepared.

Risk of bias (quality) assessment

To assess the risk of bias in individual studies (quality assessment) identified after the screening, the National Institutes of Health (NIH) checklist for observational, cohort, and cross-sectional studies was used [16]. The studies with a minimum score of eight or above, seven, and five or below "Yes responses" were considered good, fair, and poor quality respectively.

For cross-sectional studies, question numbers 1, 2, 3, 4, 5, and 11 were relevant. The responses to the remaining eight questions (6-10, 12, 13, 14) were marked as not applicable (NA). The studies with six "Yes" responses were considered good and those with four /five were qualified as fair. The studies with less than four "Yes responses" were qualified as of poor quality. The quality of the studies was assessed independently by two reviewers.

Meta-analysis

Meta-analysis was performed in STATA version 13.0 (College Station, TX, USA). The forest plots were made using the midas package in STATA. The pooled HPV vaccine uptake with a

95% confidence interval was reported among adolescent girls, adolescent boys, transgender women, and MSM up to the age of 26 years. As substantial methodological and statistical heterogeneity was anticipated, the random effects model was applied. The forest plots depict the pooled prevalence of HPV vaccination uptake with 95% CI and Chi-square statistic (Q statistic). The heterogeneity between qualified studies was reported using I^2 index. The values ranging between 25%-49%, 50-74%, and 75-100% imply low, moderate, and high heterogeneity respectively [17].

Assessment of Publication Bias

Egger's test was used to report the publication bias. Weighted linear regression with standardised effect estimate was the dependent variable and precision was the independent variable. In the present study, the $\log^e$ prevalence rate of HPV vaccine uptake among adolescents, transgender women, MSM, and precision were considered the effect estimate. The precision was reported as 1/standard error of $\log^e$ prevalence rate. Weights were allotted using the inverse variance approach (1/variance of the effect estimate). The statistically significant bias coefficient implies publication bias ($P < 0.05$).

3. Results

3.1 Identification and selection of sources of evidence

Using specific search terms, 153 articles from high income countries were screened from electronic databases and fifty-three articles were systematically reviewed (**Figure 1**). Twenty-three articles regarding the HPV vaccination coverage were qualified for the meta-analysis (quantitative synthesis) including eleven studies on gender neutral adolescent HPV vaccination [18–27] and six studies each among transgender women [10,28–32] and men having sex with men [29,30,33–36].

3.2 Characteristics of sources of evidence

Out of the eleven qualified studies on adolescent HPV vaccination uptake, five studies from the USA, two from Australia, and one each from Canada, Austria, France, and Germany were analysed **(Table 1)**. The study population included 2,780,307 adolescents including 1,975,425 girls and 804, 882 boys. All these studies were cross sectional studies except three cohort studies from Germany, Canada, and France [24,25,27]. The HPV vaccination uptake varied between 46%-97.4% among adolescent girls. Meanwhile, the uptake of the HPV vaccine varied between 27.9%-95.3% among adolescent boys. All the six qualified studies reporting the HPV vaccination coverage among 207 transgender women were from the USA **(Table 2)**. The study sample of TGW reporting to health clinics ranged between six to sixty-five. The HPV vaccination coverage among the study population varied between 4.3%-88%. The HPV vaccination data among 11,909 MSM were retrieved from six US studies **(Table 3)**. Three studies enrolled young men from twenty metropolitan statistical areas as part of National HIV Behavioural Surveillance (NHBS).[33,34,36] Meanwhile, two qualified studies were from the multicentre Vaccine Impact in Men (VIM) study [29,30]. Except for one study by Thomas et al, all the qualified studies included adult MSM [35]. Thomas et al included data from young boys aged 13 to 18 years and adults up to the age of 26 years attending a sexually transmitted disease clinic and teen health centre. The number of MSM attending the health clinics ranged between four hundred to three thousand one hundred and fifty-five. The HPV vaccination coverage varied between 4.9% to 47.6%. Quality of all the studies included for the meta-analysis was good.

3.3 Meta-analysis (Quantitative synthesis)

The pooled prevalence of the HPV vaccine uptake among 1,975,415 adolescent girls from eleven qualified studies was 67.75% (95% CI: 62.81%,72.69%). As shown in the **Figure 2** the I^2 value was 99.92%. The pooled prevalence of the HPV vaccine uptake among 804,882 adolescent boys was 58.32% (95% CI 48.88%, 67.76%) with an I^2 value of 99.98% (**Figure**

3). The pooled prevalence of the HPV vaccine coverage among 207 transgender women was 32.59% (95% CI 8.35%. 56.83%) with an I^2 value of 96.16% **(Figure 4).** The pooled prevalence of the HPV vaccine uptake among 11,909 men having sex with men (MSM) was 27.94% (95% CI 14.35, 41.53) and I^2 value was 99.72% **(Figure 5)**.

Publication bias

The *p*-value for the bias coefficient is above 0.05, which is statistically non-significant. There is no evidence of publication bias.

Discussion

There is strong evidence of gender disparity in the HPV vaccination coverage in high income countries. The present review systematically analysed the HPV vaccination coverage among adolescents, TGW, and MSM. Persistently high HPV vaccine coverage was observed among adolescent girls in all the qualified studies. Meanwhile, the HPV vaccination coverage was suboptimal among MSM and TGW. TGW and MSM were enrolled from sexually transmitted disease (STD) centres, gender diversity centres, or HIV surveillance units. Hence, the study findings may not be applicable to all the gender diverse persons in the USA. Few TGW sought medical care with minimal intention to take the HPV vaccine. In contrast to TGW, MSM in the USA were more aware of sexually transmitted diseases and were not hesitant to consult healthcare professionals if needed [33,34]. Generally, the rural health care providers were less likely to recommend adolescent HPV vaccination for cancer prevention [37]. Transgender women are not often included in the epidemiological studies as they encounter many barriers and challenges in accessing appropriate healthcare. Transgender individuals in the USA encounter higher morbidity and mortality than cisgender individuals [38]. As per the Swedish population based cohort study, sex reassigned transgender individuals demonstrate a higher mortality rate than the general population [39]. A higher incidence of sexually

transmitted diseases such as gonorrhoea, chlamydial infections, and Human Immunodeficiency Virus (HIV) infection were reported among transgender men and women [40]. Singh et al reported higher HPV prevalence among TGW than MSM in the USA [10]. A higher incidence of HPV-related cervical and anogenital cancers was reported among transgender and gender diverse persons [41].

Most of the published studies were carried out among adolescents before the COVID-19 pandemic with adequate response rates. All the included studies were qualified as good. Except for one study conducted at a medical centre, all the retrieved US studies on adolescent HPV vaccination reported provider verified data from the National Teen Survey (NIS-Teen). The studies from high income countries in the Europe, Australia, and the USA were included. Across Europe, higher vaccine hesitancy was reported in France and Germany due to the lack of trust in healthcare system and low vaccine confidence [42]. This meta-analysis has included one study each from France and Germany regarding adolescent HPV vaccine uptake.

Limitations

A limited number of studies were conducted during the pandemic with a low response rate due to COVID-19 related public health preventive measures. Both the COVID-19 pandemic and Ukraine war affected the response rate in a study from Germany [27]. Adolescent HPV vaccines administered by the general practitioners were not notified in many high income countries resulting in underestimation. Data among rural adolescents were lacking in some studies. Most of the US publications regarding adolescent vaccinations were based on the NIS Teen survey which collected information through telephone interviews. The households without telephones were not included in the survey underestimating the actual HPV vaccination coverage. As the major proportion of reviewed studies were cross-sectional

studies, confounding factors such as provider recommendation could not be adjusted and causal inference could not be reported.

All the qualified studies regarding the HPV vaccination coverage among TGW and MSM were from the USA. The sample size was small in studies enrolling transgender women, resulting in wider confidence interval while pooling the data. Hence the findings may not be generalisable to the entire US. Few studies restricted the inclusion to individuals having English language expertise. Recruitment of TGW from rural areas through social media platforms would have caused selection bias. Self-reporting of HPV vaccination status by adult men having sex with men and TGW would have generated recall bias and social desirability bias. The studies about other gender diverse groups such as transgender men, and lesbians were not qualified.

Conclusion

Significant gender inequities exist in the HPV vaccination initiation. The barriers to HPV vaccination uptake should be managed by targeted public health and clinical practice interventions. The major barriers were low awareness among healthcare providers and parents, lack of recommendations from trusted clinicians, and public mistrust towards all vaccines.

Recommendations

1. Policymakers should promote gender inclusive HPV vaccination services to all sexually naïve adolescents and young adults, irrespective of their gender identity.

2. Partnerships between health clinics and gender diverse units or organisations should be established to assist gender minorities by disseminating knowledge.

3. Personalised care aiming for tailored interventions to meet the patient preferences in an appropriate social context should be adopted.

4. Patient-physician shared decision making should be promoted in the HPV vaccination programs.

Conflict of interests

The authors declare that there is no conflict of interest.

Funding statement: No funding received.

Ethics statement

The author states that this manuscript does not involve any misconduct such as plagiarism, forgery, tampering, improper signature, multiple submission, repeated publication, split publication, etc. The systematic review and meta☐analysis used published data from peer-reviewed articles in indexed journals. As such, ethical approval to conduct the analysis was not sought.

Data availability statement

The data that support the findings of this study are available from the corresponding author upon reasonable request. (Dr Sasidharanpillai Sabeena, sabeenauthradam@gmail.com).

ORCID ID

Sasidharanpillai Sabeena https://orcid.org/0000-0002-2099-9312

Nagaraja Ravishankar https://orcid.org/0000-0002-9533-039X

All authors attest they meet the ICMJE criteria for authorship

Author contributions

Conceptualization, Data curation: S Sabeena

Formal analysis: S Sabeena, N Ravishankar

Methodology: S Sabeena

Validation: S Sabeena & N Ravishankar

Original draft; S Sabeena

Writing/Editing – S Sabeena & N Ravishankar

References

[1]   Wolf J, Kist LF, Pereira SB, Quessada MA, Petek H, Pille A, et al. Human papillomavirus infection: Epidemiology, biology, host interactions, cancer development, prevention, and therapeutics. Rev Med Virol 2024;34:e2537. https://doi.org/10.1002/rmv.2537.

[2]   World Health Organization. Cervical Cancer Elimination Initiative 2023. https://www.who.int/initiatives/cervical-cancer-elimination-initiative (accessed January 21, 2025).

[3]   Lechner M, Liu J, Masterson L, Fenton TR. HPV-associated oropharyngeal cancer: epidemiology, molecular biology and clinical management. Nat Rev Clin Oncol 2022;19:306–27. https://doi.org/10.1038/s41571-022-00603-7.

[4]   World Health Organization. The selection and use of essential medicines 2023: web annex A: World Health Organization model list of essential medicines: 23rd list (2023) 2023.

[5]   Dykens JA, Smith JS, Demment M, Marshall E, Schuh T, Peters K, et al. Evaluating the implementation of cervical cancer screening programs in low-resource settings globally: a systematized review. Cancer Causes Control 2020;31:417–29. https://doi.org/10.1007/s10552-020-01290-4.

[6]   Kaur KN, Niazi F, Nandi D, Taneja N. Gender-Neutral HPV Vaccine in India; Requisite for a Healthy Community: A Review. Cancer Control 2024;31:10732748241285184. https://doi.org/10.1177/10732748241285184.

[7]   Ng SS, Hutubessy R, Chaiyakunapruk N. Systematic review of cost-effectiveness studies of human papillomavirus (HPV) vaccination: 9-Valent vaccine, gender-neutral and multiple age cohort vaccination. Vaccine 2018;36:2529–44. https://doi.org/10.1016/j.vaccine.2018.03.024.

[8]   Simons JJM, Westra TA, Postma MJ. Cost-effectiveness of a male catch-up human papillomavirus vaccination program in the Netherlands. Prev Med Rep 2022;28:101872. https://doi.org/10.1016/j.pmedr.2022.101872.

[9]   Dorji T, Tshomo U, Gyamtsho S, Tamang ST, Wangmo S, Pongpirul K. Gender-neutral HPV elimination, cervical cancer screening, and treatment: Experience from Bhutan. Int J Gynaecol Obstet 2022;156:425–9. https://doi.org/10.1002/ijgo.13728.

[10]   Singh V, Gratzer B, Gorbach PM, Crosby RA, Panicker G, Steinau M, et al. Transgender Women Have Higher Human Papillomavirus Prevalence Than Men Who Have Sex With Men—Two U.S. Cities, 2012–2014. Sex Transm Dis 2019;46:657–62. https://doi.org/10.1097/OLQ.0000000000001051.

[11]   Amboree TL, Darkoh C. Barriers to Human Papillomavirus Vaccine Uptake Among Racial/Ethnic Minorities: a Systematic Review. J Racial Ethn Health Disparities 2021;8:1192–207. https://doi.org/10.1007/s40615-020-00877-6.

[12]   World Health Organization. Indicator Metadata Registry Details 2024. https://www.who.int/data/gho/indicator-metadata-registry/imr-details/5560 (accessed January 23, 2025).

[13]   World Health Organization. WHO adds an HPV vaccine for single-dose use 2024. https://www.who.int/news/item/04-10-2024-who-adds-an-hpv-vaccine-for-single-dose-use (accessed February 22, 2025).

[14]   CDC. ACIP Recommendations: Human Papillomavirus (HPV) Vaccine. ACIP Vaccine Recommendations and Guidelines 2024. https://www.cdc.gov/acip-recs/hcp/vaccine-specific/hpv.html (accessed January 28, 2025).

[15]   American Psychological Association. Defining transgender terms. Https://WwwApaOrg 2018. https://www.apa.org/monitor/2018/09/ce-corner-glossary (accessed January 24, 2025).

[16]   NHLBI/NIH. Study Quality Assessment Tools | NHLBI, NIH 2013. https://www.nhlbi.nih.gov/health-topics/study-quality-assessment-tools (accessed June 23, 2024).

[17]   Higgins JPT, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses. BMJ 2003;327:557–60. https://doi.org/10.1136/bmj.327.7414.557.

[18]   Borena W, Luckner-Hornischer A, Katzgraber F, Holm-von Laer D. Factors affecting HPV vaccine acceptance in west Austria: Do we need to revise the current immunization scheme? Papillomavirus Res 2016;2:173–7. https://doi.org/10.1016/j.pvr.2016.10.001.

[19]   Brotherton JM, Winch KL, Bicknell L, Chappell G, Saville M. HPV vaccine coverage is increasing in Australia. Med J Aust 2017;206:262. https://doi.org/10.5694/mja16.00958.

[20]   Swiecki-Sikora AL, Henry KA, Kepka D. HPV Vaccination Coverage Among US Teens Across the Rural-Urban Continuum. J Rural Health 2019;35:506–17. https://doi.org/10.1111/jrh.12353.

[21]   Elam-Evans LD, Yankey D, Singleton JA, Sterrett N, Markowitz LE, Williams CL, et al. National, Regional, State, and Selected Local Area Vaccination Coverage Among Adolescents Aged 13-17 Years - United States, 2019. MMWR Morb Mortal Wkly Rep 2020;69:1109–16. https://doi.org/10.15585/mmwr.mm6933a1.

[22]   C P, D Y, Ld E-E, Le M, Cl W, B F, et al. National, Regional, State, and Selected Local Area Vaccination Coverage Among Adolescents Aged 13-17 Years - United States, 2020. MMWR Morbidity and Mortality Weekly Report 2021;70. https://doi.org/10.15585/mmwr.mm7035a1.

[23]   Adjei Boakye E, Fedorovich Y, White M, Vohra S, Volle M, Osazuwa-Peters N, et al. Rural-Urban Disparities in HPV Vaccination Coverage Among Adolescents in the Central Part of the State of Illinois, USA. J Community Health 2023;48:24–9. https://doi.org/10.1007/s10900-022-01136-x.

[24]   de Pouvourville G, Guyot E, Farge G, Belhassen M, Bérard M, Jacoud F, et al. Human Papillomavirus (HPV) vaccination coverage among French adolescents: A claims data study. Vaccine 2024;42:126039. https://doi.org/10.1016/j.vaccine.2024.06.007.

[25]   Du C, Voaklander D, Meherali S, Paudel YR, MacDonald SE. Human papillomavirus vaccine coverage among immigrant adolescents in Alberta: a population-based cohort study. International Health 2024:ihae038. https://doi.org/10.1093/inthealth/ihae038.

[26]   McIndoe L, Wilson E, Ferson MJ, Sheppeard V. Beyond Averages: Unpacking Disparities in School-Based Vaccination Coverage in Eastern Sydney: An Ecological Analysis. Vaccines (Basel) 2024;12:888. https://doi.org/10.3390/vaccines12080888.

[27]   Singer R, Hübotter I, Hölzner F, Genedl C, Jasker L, Michalski N, et al. School vaccination programmes to increase HPV vaccination coverage – Experiences from Bremen, Germany. Vaccine 2025;45:126636. https://doi.org/10.1016/j.vaccine.2024.126636.

[28]   Bednarczyk RA, Whitehead JL, Stephenson R. Moving beyond sex: Assessing the impact of gender identity on human papillomavirus vaccine recommendations and uptake among a national sample of rural-residing LGBT young adults. Papillomavirus Res 2017;3:121–5. https://doi.org/10.1016/j.pvr.2017.04.002.

[29]   Amiling R, Winer RL, Newcomb ME, Gorbach PM, Lin J, Crosby RA, et al. Human papillomavirus vaccination coverage among young, gay, bisexual, and other men who have sex with men and transgender women — 3 U.S. cities, 2016–2018. Human Vaccines & Immunotherapeutics 2021.

[30]   Forward T, Meites E, Lin J, Hughes JP, Unger ER, Markowitz LE, et al. Sensitivity of Self-Reported Human Papillomavirus Vaccination History Among 18- to 26-Year-Old Men Who Have Sex With Men: Seattle, WA, 2016 to 2018. Sex Transm Dis 2022;49:81–5. https://doi.org/10.1097/OLQ.0000000000001503.

[31]   Domínguez-Riscart J, Ariza-Jimenez A-B, Baez-Castillo C, Mateo-Gavira I. Factors associated with knowledge and vaccination intention for human papillomavirus on trans girls by their main caregiver: A cross-sectional study. Front Immunol 2023;14:1097449. https://doi.org/10.3389/fimmu.2023.1097449.

[32]   Schwartz BI, Bear B, Kazak AE. Vaccination against Human Papilloma Virus in Transgender and Gender Diverse Patients. Journal of Pediatric and Adolescent Gynecology 2023;36:191. https://doi.org/10.1016/j.jpag.2023.01.132.

[33]   Meites E, Markowitz LE, Paz-Bailey G, Oster AM, NHBS Study Group. HPV vaccine coverage among men who have sex with men - National HIV Behavioral

Surveillance System, United States, 2011. Vaccine 2014;32:6356–9. https://doi.org/10.1016/j.vaccine.2014.09.033.

[34]   Oliver SE, Hoots BE, Paz-Bailey G, Markowitz LE, Meites E, NHBS Study Group. Increasing Human Papillomavirus Vaccine Coverage Among Men Who Have Sex With Men-National HIV Behavioral Surveillance, United States, 2014. J Acquir Immune Defic Syndr 2017;75 Suppl 3:S370–4. https://doi.org/10.1097/QAI.0000000000001413.

[35]   Thomas R, Higgins L, Ding L, Widdice LE, Chandler E, Kahn JA. Factors Associated With HPV Vaccine Initiation, Vaccine Completion, and Accuracy of Self-Reported Vaccination Status Among 13- to 26-Year-Old Men. Am J Mens Health 2018;12:819–27. https://doi.org/10.1177/1557988316645155.

[36]   McClung N, Burnett J, Wejnert C, Markowitz LE, Meites E, NHBS Study Group. Human papillomavirus vaccination coverage among men who have sex with men-National HIV Behavioral Surveillance, United States, 2017. Vaccine 2020;38:7417–21. https://doi.org/10.1016/j.vaccine.2020.08.040.

[37]   Walker TY, Elam-Evans LD, Williams CL, Fredua B, Yankey D, Markowitz LE, et al. Trends in human papillomavirus (HPV) vaccination initiation among adolescents aged 13–17 by metropolitan statistical area (MSA) status, National Immunization Survey – Teen, 2013 – 2017. Hum Vaccin Immunother 2019;16:554–61. https://doi.org/10.1080/21645515.2019.1671765.

[38]   Hughes LD, King WM, Gamarel KE, Geronimus AT, Panagiotou OA, Hughto JMW. Differences in All-Cause Mortality Among Transgender and Non-Transgender People Enrolled in Private Insurance. Demography 2022;59:1023–43. https://doi.org/10.1215/00703370-9942002.

[39]   Dhejne C, Lichtenstein P, Boman M, Johansson ALV, Långström N, Landén M. Long-Term Follow-Up of Transsexual Persons Undergoing Sex Reassignment Surgery: Cohort Study in Sweden. PLoS One 2011;6:e16885. https://doi.org/10.1371/journal.pone.0016885.

[40]   Van Gerwen OT, Jani A, Long DM, Austin EL, Musgrove K, Muzny CA. Prevalence of Sexually Transmitted Infections and Human Immunodeficiency Virus in Transgender Persons: A Systematic Review. Transgend Health 2020;5:90–103. https://doi.org/10.1089/trgh.2019.0053.

[41]   LoSchiavo C, Greene RE, Halkitis PN. Human Papillomavirus Prevalence, Genotype Diversity, and Risk Factors Among Transgender Women and Nonbinary Participants in the P18 Cohort Study. AIDS Patient Care STDS 2020;34:502–5. https://doi.org/10.1089/apc.2020.0213.

[42]   Verger P, Botelho-Nevers E, Garrison A, Gagnon D, Gagneur A, Gagneux-Brunon A, et al. Vaccine hesitancy in health-care providers in Western countries: a narrative review. Expert Review of Vaccines 2022;21:909–27. https://doi.org/10.1080/14760584.2022.2056026.

[43]   Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ 2021;372:n71. https://doi.org/10.1136/bmj.n71.

Figure Captions

Figure 1: The PRISMA 2020 flow diagram for systematic reviews [43]. The flow diagram illustrates the number of articles screened, abstracts retrieved and full texts included/ excluded for the systematic review and meta-analysis.

PRISMA: Preferred Reporting Items for Systematic Reviews and Meta-analyses.

Figure 2: Forest plot of the summary effect size (pooled prevalence of HPV vaccine uptake among adolescent girls) using random effects model. Squares imply the effect size of individual studies and extended lines represent 95% confidence interval. The size of the squares indicate the weight of the studies based on sample size using a random effects analysis. The diamond data indicates the pooled prevalence. Test of heterogeneity = 99.92%, *P*-value =0.00. CI confidence interval.

Figure 3: Forest plot of the summary effect size (pooled prevalence of HPV vaccine uptake among adolescent boys) using random effects model. Squares imply the effect size of individual studies and extended lines represent 95% confidence interval. Size of the squares indicate the weight of the studies based on sample size using a random effects analysis. The diamond data indicates the pooled prevalence. Test of heterogeneity =99.98%, *P*-value = 0.00. CI confidence interval.

Figure 4: Forest plot of the summary effect size (pooled prevalence of HPV vaccine uptake among transgender women) using random effects model. Squares imply the effect size of individual studies and extended lines represent 95% confidence interval. Size of the squares indicate the weight of the studies based on sample size using a random effects analysis. The diamond data indicates the pooled prevalence. Test of heterogeneity =96.16 %, *P*-value =0.00. CI confidence interval.

Figure 5: Forest plot of the summary effect size (pooled prevalence of HPV vaccine uptake among men having sex with men) using random effects model. Squares imply the effect size of individual studies and extended lines represent 95% confidence interval. Size of the squares indicate the weight of the studies based on sample size using a random effects analysis. The diamond data indicates the pooled prevalence. Test of heterogeneity =99.72 %, *P*-value =0.00. CI confidence interval

Figure 1: The PRISMA 2020 flow diagram for systematic reviews [43]. The flow diagram illustrates the number of articles screened, abstracts retrieved and full texts included/ excluded for the systematic review and meta-analysis.

Figure 2: Forest plot of the summary effect size (pooled prevalence of HPV vaccine uptake among adolescent girls) using random effects model. Squares imply the effect size of individual studies and extended lines represent 95% confidence interval. The size of the squares indicate the weight of the studies based on sample size using a random effects analysis. The diamond data indicates the pooled prevalence. Test of heterogeneity = 99.92%, *P*-value =0.00. CI confidence interval.

Figure 3: Forest plot of the summary effect size (pooled prevalence of HPV vaccine uptake among adolescent boys) using random effects model. Squares imply the effect size of individual studies and extended lines represent 95% confidence interval. Size of the squares indicate the weight of the studies based on sample size using a random effects analysis. The diamond data indicates the pooled prevalence. Test of heterogeneity =99.98%, *P*-value = 0.00. CI confidence interval.

Figure 4: Forest plot of the summary effect size (pooled prevalence of HPV vaccine uptake among transgender women) using random effects model. Squares imply the effect size of individual studies and extended lines represent 95% confidence interval. Size of the squares indicate the weight of the studies based on sample size using a random effects analysis. The diamond data indicates the pooled prevalence. Test of heterogeneity =96.16 %, *P*-value =0.00. CI confidence interval.

Figure 5: Forest plot of the summary effect size (pooled prevalence of HPV vaccine uptake among men having sex with men) using random effects model. Squares imply the effect size of individual studies and extended lines represent 95% confidence interval. Size of the squares indicate the weight of the studies based on sample size using a random effects analysis. The diamond data indicates the pooled prevalence. Test of heterogeneity =99.72 %, *P*-value =0.00. CI confidence interval.

Table 1: Characteristics of qualified studies on gender neutral HPV vaccination uptake in adolescents

No	Author (year)	Country	Study population	Age group	Study design	Study period	Sample (N)	Vaccination uptake Girls N (%)	Vaccination uptake Boys N(%)	Limitations	Quality
1	Singer et al. (2025)	Bremen, Germany	School students	13-14	Cohort study	2022-2023	2694 Girls-1418 Boys-1276	647 (46)	361(28)	COVID-19 and Ukraine war contributed to lower response rate. Small cohort size. Data from limited period was considered.	Good
2	McIndoe et al. (2024)	Australia	School students	11-12	Cross sectional	2019	9379 4572 girls 4807 Boys	4023 (88)	4134 (86)	Findings may not be generalizable, as the study was conducted in south eastern Sydney. Catch-up and general practitioner administered HPV vaccination coverage were not included.	Good
3	Crystal Du et al. (2024)	Alberta, Canada	non-immigrant adolescents	12-18	Cohort	2014-2018	208, 2 56 101,725 girls 106,531 boys	64,606 (63.51)	42,357 (39.76)	Coverage data was limited to Western Canadian Province and may not be generalizable.	Good
4	de Pouvourville et al.(2024)	France	adolescents	11-19	Cohort Claims study	2017-2022	2,366,067 Girls-1,773,900 boys 592,167	1,193,834 (67.3)	369,512 (62.4)	Number of vaccines dispensed was taken as actual vaccination data resulting in overestimation. Vaccinated during free medical visits was not recorded in the national database leading to underestimation.	Good
5	Boakye et al. (2022)	Illinois USA	adolescents	11-17	Cross sectional	2015-2020	9351 girls 4649 boys 4702	2241(48.2)	2076 (44.1)	Data from one medical centre was analysed. Hence may not be representative of the national situation.	Good
6	Pingali et al. (2021)	USA	adolescents	13-17	Cross sectional study NIS-Teen survey	2020	20,163 girls 9,576 boys 10,587	7383 (77.1)	7739 (73.1)	Low response rate may increase the bias. As per the survey error assessment, NIS Teen survey might underestimate true coverage. Households without telephone could	Good

No	Author	Country	Population	Age	Study design	Year	Sample size			Comments	Quality
										not be included.	
7	Elam Evans et al. (2020)	USA	adolescents	13-17	cross sectional study NIS-Teen survey	2019	18,788 girls 8916 boys 9872	6526 (73.2)	6890 (69.8)	Lower response rate due to COVID-19 pandemic. Increase in bias rates. Phone-less households could not be included.	Good
8	Swiecki-Sikora et al. (2019)	USA	adolescents	13-17	cross sectional study NIS-Teen survey	2012-2013	37,114 Girls-17,596 Boys-19,518	9783 (55.6)	5,445 (27.9)	Low response rate and minimal data were provider verified. Inclusion of provider verified records would have caused underestimation.	Good
9	Walker et al. (2018)	USA	adolescents	13-17	cross sectional study NIS-Teen Survey	2017	103,703 girls 50,711 boys 52,992	34,788 (68.6)	33,172 (62.6)	Biases in estimates due to low response rates.	Good
10	Brotherton et al (2017)	Australia	adolescents	12 years	Cross sectional study pf NHPVR	2015	4,343 Girls- 2108 Boys- 2235	2054 (97.4)	2129 (95.3)	HPV vaccination by general practitioners might not be notified resulting in underestimation.	Good
11	Borena et al. (2014)	Austria	adolescents	9-10	cross sectional	2015	449 girls 254 boys 195	150 (59)	101 (51.8)	Data from limited number of adolescents from western region adolescents from rural areas not included.	Good

Table 2: Characteristics of studies among Transgender women (TGW)

S No	Author (year)	Country	Study population	Age group	Study design	Study period	Sample (N)	Vaccination uptake among TGW N (%)	Limitations	Quality
1	Schwartz et al (2023)	USA	multidisciplinary paediatric gender program	11-18	Cross sectional	2015-2020	**25**	**22 (88)**	Small sample size	Good
2	Domínguez-Riscart J (2023)	USA	multidisciplinary Gender Diversity Units	9-16	Cross sectional	Jan-July 2022	**65**	**14 (21.5)**	Possibility of selection bias as participating mothers might be more health conscious.	Good
3	Forward et al (2023)	USA	Multicentre VIM study	18-26	Cross sectional	2016-2018	**6**	**3 (50)**	Small sample size. Eligible individuals requiring language translation are excluded. Recall bias owing to self-reporting of HPV vaccine uptake.	Good
4	Singh et al. (2019)	USA	YMHPV study at two community health clinics	18-26	Cross sectional	2012-2014	**49**	**5/49 (10.2)**	Small sample size. Not generalizable to all TGW in the US	Good
5	Amiling et al. (2021)	USA	VIM study (three urban centre clinics)	18-26	Cross sectional	2016-2018	**39**	**11 (28.2)**	Self-reporting of HPV vaccination status might be subjected to recall bias, study not representative of three cities.	Good
6	Bednarczyk et al. (2014)	USA	LGBTQ 185 TGW-15	18-26	Cross sectional	2014	**23**	**1 (4.3)**	Recruitment through social networking sites and targeted commercials to rural residents, Limited to tech savvy individuals leading to selection bias. Self-reported HPV vaccination status was not confirmed.	Good

LGBTQ Lesbian Gay Bisexual Transgender Queer **TGW** transgender women   **YMHPV** study Young men's HPV study **VIM** study Vaccine Impact in Men study

No	Author (year)	Country	Study population	Age group	Study design	Study period	MSM (N)	Vaccination Uptake N (%)	Limitations	Quality

1	Forward et al. (2023)	USA	Multicentre Vaccine Impact (VIM) study	18-26	Cross sectional	2016-2018	738	289 (39.2)	Hospital based studies are not generalizable. Recall bias due to self-reporting.	Good
2	Amiling et al. (2021)	USA	VIM study (three urban centre clinics)	18-26	Cross sectional	2016-2018	1342	640 (47.6)	As this was not a population based study, not representative of all MSM in three US cities. Self-reporting of data resulted in under or over estimation of vaccination coverage.	Good
3	McClung et al. (2020)	USA	Young MSM from twenty MSA as part of NHBS	18-26	Cross sectional	2017	2482	815 (32.8)	NHBS is not a national survey and the study population from MSA is not representative of all MSM in the USA. Recall bias due to self-reporting of data.	Good
4	Thomas et al. (2018)	USA	Young men attending STD clinic and a teen health centre.	13-26	Cross sectional	2013-2015	400	104 (26)	Hospital based study population included low income men and may not be representative of entire study area. Recall bias due to self-reporting.	Good
5	Oliver et al (2017)	USA	Young MSM from twenty MSA as part of NHBS	18-26	Cross sectional	2014	2892	497 (17.2)	Recall bias or social desirability bias due to self-reporting of data. Due to anonymous survey, self-reporting could not be confirmed with medical records. Study population at MSA was not representative of all MSM in the USA.	Good
6	Meites et al. (2014)	USA	Young MSM from twenty MSA as part of NHBS	18-26	Cross sectional	2006-2011	3155	157 (4.9)	Recall bias due to self-reporting of data. Venue based sampling at bars, clubs did not represent all MSM.	Good