Long title: Co-development of an acceptance and commitment

therapy-based intervention to increase intrinsic motivation of

adolescents to change weight: the AIM2Change study

Short title: AIM2Change Co-development

Witherden JS[1†], Searle AJ[1†], Iturbe I[2], Thornton GA[3], Wright I[4], Semple C[4], Clare K[3],

Hamilton-Shield JP[1], Hinton EC[1*]

[1]National Institute for Health and Care Research Bristol Biomedical Research Centre

Diet and Physical Activity Theme University Hospitals Bristol NHS Foundation Trust

and University of Bristol, Bristol, UK, BS8 1TU, UK.

[2]Department of Clinical and Health Psychology and Research Methodology, Faculty of

Psychology, University of the Basque Country (UPV/EHU), Donostia-San Sebastian,

Basque Country, Spain.

[3]Public contributor

[4]Bristol Royal Hospital for Children, University Hospitals Bristol & Weston NHS

Foundation Trust, Bristol, UK, BS8 2AE, UK

*Corresponding author

E-mail: elanor.hinton@bristol.ac.uk

[†]Denotes joint first authorship.

Abstract (max 300 words)

Childhood obesity levels continue to rise, with significant impacts to individuals and the NHS. The 'Complications of Excess Weight' (CEW) clinics provide support to young people with complications of their weight. Our objective was to co-develop, with young people, a new intervention (AIM2Change), to enable young people to develop their intrinsic motivation to manage weight, using Acceptance and Commitment Therapy (ACT), with a person-centred approach. Young people from the Care of Childhood Obesity (CoCO) clinic in Bristol, UK, were recruited to co-develop this intervention. The study was registered on ISRCTN (ISRCTN16607863). The seven-session, ACT-based intervention was delivered one-to-one, securely online. Qualitative interviews were conducted after each intervention session was delivered. Qualitative data were coded and reviewed regularly to make iterative changes to individual sessions and the overall programme. Fourteen co-developers were recruited, of whom nine completed the co-development process (female=4; median age (IQR)=15(1.5); 4 with a parent; Indices of Multiple Deprivation (IMD) range=1-10). Iterative changes made during co-development included introducing earlier a focus on eating behaviour and body image, with more practical activities to increase engagement. Thematic analysis of co-developer feedback identified four themes: theoretical understanding; delivery and receipt of therapy; view of strategies and engagement; real world benefits of co-development process. Framework analysis was conducted to map data pertaining to these themes into matrices according to each participant and session. Insights from the co-development process have shaped AIM2Change to optimise the intervention's value, relevance and acceptability. Findings suggest that AIM2Change meets an unmet need in delivery of current clinical services.

Introduction

The prevalence of childhood obesity remains alarmingly high(1), with over 340 million children and adolescents classified as overweight or obese worldwide(2). Moreover, among high-income Western countries, English-speaking and Mediterranean regions have the highest levels of obesity in children and adolescents aged 5-19 years(3). Childhood overweight or obesity is associated with several physical (e.g. metabolic risk factors) and psychological (e.g. anxiety) comorbidities(4), which are maintained into, or emerge in, adulthood(5). The early onset of obesity that persists into adulthood is associated with greater morbidity and mortality than obesity that arises in adulthood alone(6). To this is added the financial cost to the NHS(7).

The National Institute for Health and Care Excellence (NICE) guidance recommends multicomponent lifestyle interventions as the preferred option for tackling obesity in childhood and adolescence(8). These interventions promote changes in calorie and nutritional intake and physical activity, which have demonstrable benefits for reducing abdominal obesity(9). If treatments so far have been unsuccessful, and patients are over 12 years age and/or have comorbidities, the option of using pharmacological treatments is considered. Surgical interventions remain a rarely utilised option(8). Within pharmacological treatments, the US Food and Drug Administration (FDA) and the European Medicines Agency (EMA) have approved the administration of subcutaneous Glucagon-Like-Peptide 1 (GLP-1) agonists as an adjunct to lifestyle interventions in adolescents, under certain conditions(10).

In line with the recommendations, the Care of Childhood Obesity (CoCO) clinic at Bristol Children's hospital offers a comprehensive treatment approach delivered by a multidisciplinary team (MDT) to help children and young people with enduring obesity manage their weight. However, such an approach does not always yield the expected

benefits. Thus, a qualitative service review was conducted to explore the service-related

perceptions of families and patients who did not meet their weight-loss goals(11). They

reported feeling lost, unmotivated and hoped that the clinic could take responsibility for

their weight management(11).

According to Deci and Ryan's Self-determination theory(12), two types of

motivation are differentiated: intrinsic motivation (i.e. determined by internal factors,

such as personal reward) and extrinsic motivation (i.e. determined by external factors,

such as pleasing others). Intrinsic motivation is key to behavioural change maintenance

long term, as it is not contingent on external reinforcement that is likely to change (e.g.

being discharged from the clinical service). In this regard, Gillison et al.(13) found that

intrinsic motivation to exercise indirectly predicted exercise behaviour and quality of

life in adolescents who perceived themselves as overweight. In contrast, extrinsic

motivation showed a negative association with these outcomes and was connected to

social physique anxiety (13). Moreover, Jensen et al.(14) found that adolescents who

were successful in weight loss and maintenance reported intrinsic rather than extrinsic

motivations.

Acceptance and commitment therapy (ACT) is a therapeutic approach developed

within the third wave of cognitive-behavioural therapies based on functional

contextualism(15) and relational frame theory(16). ACT focuses on fostering

psychological flexibility and the ability to be in the present moment by deciding to

maintain or change behavioural patterns based on one's personal values(17, 18).

Personal values are fundamental to this therapeutic approach, and ACT guides the

individual to clarify what intrinsically motivates their behaviour and make

commitments to enact these behaviours(17). A variation of ACT that offers an

evidence-based framework for fostering psychological strength that has been

specifically targeted for young people, is the DNA-V (Discoverer, Noticer, Advisor-

Values) approach(19). The DNA-V model guides young people to make decisions using

the DNA-V specific skills, helping them to understand how to use those skills within

self and social contexts (Fig. 1)(19).

Fig 1. The DNA-V model (19).

Advisor relates to how a person moves in his/her 'context' (past, present, future); Discoverer relates to

what a person does (repeat patterns or try new behaviours); Noticer relates to how a person uses his/her

body, senses and awareness in various 'contexts' (internal, external); Values are the contexts in which a

person is empowered to build strength in day-to-day life to do what is important for the future (19).

Applying this to the context of weight management, ACT aims to promote

healthy behaviour patterns connected to one's personal values, through training to

develop mindfulness and the willingness to accept external or internal experiences that

may trigger unhealthy behaviours(20). Engaging in healthy behaviours sometimes

means making challenging decisions in the moment that may generate physical or

mental discomfort.

According to a network meta-analysis, ACT has the strongest evidence for adult

weight management among third-generation treatments(21). It has also been found to

improve variables related to the psychological well-being of people living with

overweight or obesity(22). Furthermore, a recent scoping review evaluated the

feasibility and acceptability of ACT in adolescent weight management(23). Treatment

acceptability, attendance, retention and appraisal were found to be high, whilst weight

and quality of life-related outcomes were promising, highlighting the need for further

studies in this area(23).

This study is the next step of an intervention development process (initial steps

reported in(24)), based on integrating the person-based approach to intervention

development(25), behaviour change theory (COM-B; Capability, Opportunity,

Motivation–Behaviour(26), and a Patient and Public Involvement (PPI) group, to co-

develop an ACT-based approach to increase intrinsic motivation in weight management

in young people(24). Specifically, this paper reports the outcomes of the co-

development work with young people, presenting the changes to the intervention that

were identified through 'Think aloud' interviews with participants, consensus meetings,

and PPI contributions.

Materials and methods

Study design and setting

This study was approved by the UK NHS North-West Research Ethics

Committee (ref: 22/NW/0337) and was registered in February 2023 in International

Standard Randomised Controlled Trial Number (ISRCTN) database

(https://www.isrctn.com/ISRCTN16607863). The study was conducted in the context of

tier-three, paediatric weight management services in Bristol (UK). Recruitment began

on 7th December 2022 and ended on 7th June 2023.

Participants

Young people attending the CoCO clinic at Bristol Royal Hospital for Children

were approached to collaborate in the intervention development if they (i) were between

11 and 18 years of age and (ii) had a body mass index (BMI) of > 95th percentile. For

recruitment, members of the MDT at the CoCO clinic at Bristol Royal Hospital for

Children offered the study information pack to young people who needed additional

support for weight management beyond the current multidisciplinary treatment in their

regular appointments. Next, the ACT therapist with the role of delivering the co-

development therapy sessions (JC), contacted the young people and their families to

answer any questions they had about the study and their participation. If the young

person was interested, written informed consent or assent was obtained online from

young people and/or their parents (dependent on age), and appointments were scheduled.

151   Procedures

152   PPI

Collaborators with lived experience of obesity provided input throughout the

research process: (i) pre-ethics committee approval (two sessions); (ii) during the data-

gathering and analysis stages (two sessions); (iii) design of next study (one session); (iv)

discussion of findings and to get feedback on our PPI approach. All PPI sessions were

facilitated by our Public Contributor research team member (GT), who has advised and

participated throughout the research process, and co-authored this paper.

The PPI Advisory Group meetings were evaluated using the CUBE

methodology(27, 28) to understand perceptions of attendees' experience in four areas:

how they could contribute (one-way to many ways), voice (strong to weak), agenda-

setting (public members to organisation) and potential to change the intervention

(organisation resistant to change intervention to willing to change). After the meetings,

those who attended answered each question using a slider under each question from a

score of 0 at one end of each dimension and a score of 1 at the other end. Contributors

could also add a comment regarding each dimension. All data were recorded

anonymously in a spreadsheet associated with the CUBE link. The mapping of

responses on the CUBE could be viewed by all contributors (after logging into the

online system).

The AIM2Change Intervention

The template intervention was based on the DNA-V model of ACT (19) and aimed to increase intrinsic motivation in young people for making and maintaining lifestyle changes by connecting them with their personal values(24). It comprised seven 45-minute sessions and was delivered on an individual basis (by JC) using the Microsoft Teams online platform, to minimise visits to the clinic, saving time and travel costs. Where personal circumstances required, arrangements were made to provide computer tablets and data packages, and participants were offered a suitable, private, space in which to hold the sessions.

Co-development with 'Think aloud' interviews

The initial therapy framework was iteratively co-developed with the participants, through 15-minute 'Think aloud' interviews. The interviews (n=63) were conducted by a qualitative researcher (AS) immediately following each therapy session to elicit feedback on that session. The 'Think aloud' interviews were appraised on an individual basis rather than on aggregated data from multiple participants/sessions, to enable real-time changes to be made. Pertinent feedback, shared during the therapy sessions and the 'Think aloud' interviews, was compiled into a 'Table of Changes', along with the insights from the PPI Advisory Groups and consensus meetings (see below). Once a point was raised, evidence in support of this suggestion, and opposing thoughts, were collated in the table. The research team (EH, AS, JC) met regularly to discuss the content of the Table of Changes and make iterative changes to the intervention accordingly. The MoScoW (Must do, Should do, Could do, Would like to do) coding scheme(25) was used to classify suggestions. Appropriate changes were also made to the interview topic guide to gather further opinions on the suggestions raised.

Consensus meetings

Once all participants had completed the intervention, two consensus meetings

were held to discuss the proposed changes and make additional refinements, with the

aim of coming to a consensus about the final intervention manual. The first meeting

involved participants in the co-development process (young people and their parents if

involved), and the second meeting involved members of the clinical MDT, collaborators

involved in the research and the management group. Agreed amendments were recorded

in the 'Table of Changes' (see Supporting information).

Data analysis

Coding of 'Think Aloud' Interviews and consensus meeting

Inductive coding was used to explore participants' overall experience of

receiving and engagement with the therapy. All interviews were transcribed by AS and

II and were managed in NVivo software. Independent 'open' coding was undertaken, by

AS and II, on a subsample of the interview transcripts (7 of 63), representing four

participants; one interview from each of the seven therapy sessions,. The initial coding

and any discrepancies in coding between the coders were discussed until consensus was

reached. A final coding frame was developed jointly, and then applied to the entire

dataset by AS. The codes were discussed further by AS, II and EH to inform the

development of four themes. The consensus meeting with young people and/or parents

was open coded using existing codes developed in the analysis of the 'Think-aloud'

interviews. (NB: the second consensus meeting was not subjected to coding and

analysis, as the focus of that meeting was on the practical delivery of the intervention

within the NHS. Insights were captured through inclusion in the Table of Changes).

Framework analysis

A framework approach enabled summarising the data pertaining to the four

themes in matrices that display the responses from (i) each participant and (ii) each

therapy/intervention session attended(29). These coded data from the consensus

meeting were also integrated by AS and II into the framework matrices, whereever they

were considered in relation to the existing themes (e.g. provided additional insight or

perspective to responses derived from the Think-aloud interviews).

Results

Patient and Public Involvement in the study

229   Seventeen young people with relevant personal experience contributed to the

230 study over the course of the overall project, and nine adults, with either personal or

231 parental experience of living with obesity. They came from the South-West of England,

232 Bristol, Sheffield and Birmingham.

Outcomes from the Patient and Public Involvement process

The advisory group members collectively provided feedback and suggestions on

the design and desirability of the research; practicalities of recruitment (including help

with the wording of recruitment materials and participant information both for young

people and their parents); and offered helpful guidance on how to approach recruitment

sensitively. Throughout, the contribution of the advisory group actively influenced the

conduct and delivery of the intervention.

240   The results of the evaluation of Advisory Group members' experience of their

241 involvement are given in Table 1, which presents the scores allotted to each CUBE

242 dimension(27, 28). In summary, the advisory group evaluated the meetings as

243 interesting, allowing different ways to participate and felt the team were highly likely to

244 make changes resulting from their input.

247 **Table 1**: CUBE evaluation of PPI sessions by advisory group members

Meeting	Purpose of PPI meeting	No. completed evaluation	CUBE dimensions				Comments
			Ways to be involved	Strong/Weak voice	Agenda	Changes	
1	Introduction to the project	9	0.76	0.72	0.50	0.80	"The session went quite quickly, but there was time to ask questions."
2	Intro to project and response to REC	4 (*2)	0.81	1.00	0.66	0.91	"They have already demonstrated change."
3	Findings so far and future grant	1 (*2)	0.83	0.84	0.64	0.82	"When people had comments to make our opinions were actively taken into account and discussed."
4	Project findings and end of grant	5 (*1)	0.92	0.93	0.71	0.87	"All participants given equal opportunity to be involved and included. never felt rushed."
	Median of meeting scores		**0.83**	**0.87**	**0.63**	**0.85**	

248 *Scores of 0.5 across the board excluded as appears no changes when person logged on

Fig 2. Screenshot of the CUBE.

Questions for each dimension appear on the left of the screen with the slider to move the data point in

blue. The CUBE appears on the right, with each dimension on a different axis: Contribute on the x☐axis,

Voice on the y☐axis, agenda on the z☐axis and change as colour of the data point (yellow is neutral; red

is resists change, green is willing to change). The cube can be rotated using the mouse so that contributors

can explore the data points in relation to the different dimensions (27, 28).

Co-development

Participants' characteristics

Fourteen co-developers were recruited to the study, although one did not start

due to access difficulties. Of the 13 remaining individuals (six female; mean age, 14.9

years; mean IMD score 4.9), neurodiversity was experienced by two participants with

Autism spectrum disorder (ASD) or ASD and Attention deficit hyperactivity disorder

(ADHD) and seven were receiving a GLP-1 agonist as adjuvant treatment. Eight

participants recruited had previous experience of psychological therapy. When asked

the question "How would you describe your ethnicity?", participants used the following

terms to describe themselves: Afro-Caribbean (n=1), Black African (n=1), Black British

(n=1), Eastern European White (n=1), Pakistani (n=2), Roma Gypsy (n=1), White

Romanian (n=1), White (n=2), and White British (n=4).

Fig 3. Participant flow diagram.

275 **Table 2:** Sample characteristics.

Variable	Enrolled ppts (n = 14) f/M (SD)	Completed ppts (n = 9) f/M (SD)
Female	6	4
Age	14.9 (1.58)	14.6 (1.54)
IMD	4.9 (3.16)	5.2 (3.31)
Neurodiversity	2	1
Prescribed GLP-1 Agonist	7	6
Psychological therapy experience	8	5

276 IMD = Indices of multiple deprivation; M = Mean; Ppts = participants; SD = Standard Deviation.

Co-development participation

Co-developers who started the intervention attended a mean of 5.1 sessions, and

rescheduled sessions twice on average. The completed intervention had a total mean

duration of 47 days. Parents involved in sessions attended an average of 4.8 sessions.

Two consensus meetings were held online in June and July 2023. All the young

people who participated, and their parents, were invited to attend. Six young people and

four parents attended, alongside members of the research team (JC, GT, EH). The

second meeting was attended by 10 clinical MDT members, collaborators and

management group and research team members (including a young PPI member of the

management group) to share views, experience and reach consensus on proposed

changes.

Outcomes from the co-development process

The complete 'Table of Changes' is included in Supporting information. The

key findings from the table of changes will be discussed below and how they interact

with the four themes from the 'Think Aloud' interviews (Table 3). The four themes are

1) participants' theoretical understanding, 2) factors around the delivery and receipt of

the therapy, 3) factors to do with specific strategies and 4) details of real-world benefits

of the therapy.

Table 3: Thematic structure and coding description

Themes	Codes informing theme	Description
Theoretical understanding	Identification and relevance of DNA-V; understanding and relevance of metaphors; assimilation of concepts	Participants' ability to understand the theory underpinning DNA-V approach and research team identifying examples and metaphors to facilitate delivery of the therapy
Delivery and receipt of therapy	Alignment with therapy aims; openness and receptivity to therapy; focus of sessions; structure and format of sessions; parental involvement; written materials and workbook; continuity and coherence of sessions; uniqueness of the contribution of therapy	Concurrence with aims of the study i.e. co-development. Issues with the structure and format of individual sessions, openness to therapist and suggested approaches / strategies. Practicalities – online or in-person
View of strategies and engagement	Putting strategies into practice; usefulness of strategies and metaphor; proposed changes to future sessions; satisfaction with session; continuity and coherence of sessions	Extent to which strategies and metaphors are received by participants and they feel able to, or have already been, put into practice. Evaluating understanding of strategies in context of perceived usefulness and how they relate to their own behaviour
Real world benefits of co-development process	Reported benefits of attending therapy sessions; view of therapy coming to an end	Self-reported positive outcomes associated with psychological strength and intrinsic motivation, both behavioural (e.g. eating behaviour, setting and appraising goals) relational / social (i.e. communication) outcomes

Key Learning 1. No "one-size-fits-all" approach to understanding DNA-V; instead, different activities worked for different people.

Qualitative theme: Theoretical Understanding

Whilst some young people liked an activity, others did not. It was agreed that the manual would provide several options, which the therapist could consider offering to the young person based on their current situation or receptivity:

Interviewer: *What was offered for your stress, what strategies were on offer?*

Ppt: *Oh, we did try breathing, counted breathing. It was very helpful.* (Ppt 8, Session 5, Male)

In previous ones [sessions], *we talked about the Advisor. But it wasn't about one big thing really. But today we talked about lots of different things. Yeah, we did where you shut your eyes and the leaves on the stream* [therapeutic exercise] *and everything. Yeah, that was quite good*. (Ppt 4, Session 5, Female)

Outcome: The manual will contain 'core content' that carries the fundamental DNA-V content followed by several different, flexible, exercises that explore the concepts in slightly different ways. Together the therapist and patient will experiment to see which activities help them to understand and embed the content in the context of their life and experience. Therefore, all patients will be guided using the same therapeutic principles led by DNA-V but will not experience an identical programme.

Key learning 2: Enhancing the support to connect DNA-V with their lives and context

Qualitative theme: Theoretical understanding

Whilst the videos and generic explanations were well received with regard to the

presentation of the concepts, the young people valued the support of the therapist

guiding them to understand, and apply the DNA-V concepts to their own lives, values

and experiences:

After that [video], *we started talking about those three parts of the brain and*

that went on for the rest of the time, talking about examples of it. OK. It was

really well explained. At first, I was thinking, well, how do these relate to how

people think? Because I didn't think how well it could be simplified into three

different aspects. But eventually realised it does fit perfectly, whether you're, like,

a little voice in your head, or what. You're just noticing things. It was familiar to

me, I never had quite a name for it, but it was recognisable. (Ppt 7, Session 1,

Male)

We went back on to what we did last week, which was starting off with the DNA.

And then we added the "V" in the middle, so it was like a geometric shape type

thing. And it was quite helpful because we looked at one and then we were like,

OK, Ohh, how can you apply it? (Ppt 1, Session 2, Female)

The space for self-reflection also supported the young person to link the

programme with their own experiences:

There's, like, a good amount of self-reflection, I suppose, and sort of placing

emotions on certain memories. It makes you feel stuff like that. (Ppt 2, Session,

Female)

Outcome: The intervention was adapted to enhance the 'core content' which consists of

the fundamental DNA-V materials, with specific activities that help to contextualise the

concepts for the individual, so they can understand how they could utilise them in their

life.

Key Learning 3. Earlier focus on eating, weight and self-image during the therapy

Qualitative theme: Delivery and Receipt of therapy

Throughout early PPI work, the whole-person approach to therapy was commended, as

opposed to an approach that solely looked at issues around weight. However, during the

intervention co-development itself, some young people requested that discussions of

weight, eating and self-image were introduced earlier in the therapy to have time to

expand on the issues raised:

However, other young people noted that the topic was difficult to discuss, and that

balance was important:

It's a hard thing for me to talk about. I just don't feel comfortable with it, but it's

sort of like putting in a bit of it to every single aspect that we talk about, that sort

of the right amount rather than being constantly talking about it (Ppt 7, Session

5, Male)

Outcome: The intervention was adapted to introduce discussions about eating, weight

and self-image earlier, but with the scope to focus more intently on it, throughout, being

optional. For example, many activities request that the young person offers an example

to work through, and the individual could choose to focus on an eating behaviour, body

image concern or a more general example for the exercises.

Key learning 4: The therapeutic alliance was of high importance to young people

Qualitative theme: Delivery and Receipt of Therapy

The therapeutic relationship felt important to how successful the therapy was likely to be, and for many the building of rapport was successful:

> *I think the environment when I was talking to JC* [therapist] *was quite comfortable and welcoming. I think at the start of the session I was quite stressed out and now I'm feeling quite calm through the techniques that JC taught me and that we went through today. Well, I guess I never really talked to someone about how I felt.* (Ppt 1, Session 1, Female)

> *First time ever having therapy, so I was a bit nervous about what to expect. I was quite shocked.... It was very well explained. I think it was structured and planned out quite well.* (Ppt 7, Session 1, Male)

Not all young people began the sessions feeling able to engage:

> *I don't like doing it* [talking] *with people like strangers. I just don't feel comfortable. But even if I do know the person, it's not like it's guaranteed that I'll tell them everything, like. Weird in a way to say everything you feel whenever you feel and whatnot.* (Ppt 12, Session 1, Male)

However, this resistance to engagement began to dissipate over the course of the sessions:

> *I mean, it's still, it's not, like, the most enjoyable thing ever. So still quite boring, but it's not as bad as I thought.* (Ppt 12, Session 3, Male).

Outcome: The programme structure of exercises that support rapport building within

the early sessions was retained, and further rapport building exercises were made

available for later sessions, should they be required to support the ongoing maintenance

of a strong patient-therapist relationship.

Key learning 5: Online therapy worked for the young people

Qualitative theme: Delivery and Receipt of therapy

The choice, in the design, of online therapy was made initially for pragmatic reasons, to

reduce the time commitment and cost to the co-developers of traveling to the hospital

weekly. Feedback suggested that this did not affect the rapport:

[I've] *had therapy before with someone online, and it never really flowed...*

Whereas this flows, it absolutely flows. It feels like you're in-person. (Ppt 4,

Session 3, Female)

Ppt 7: *I like doing it on Zoom as well, I felt quite safe and secure.*

Mother: *He was in his room. He was on his territory* (Ppt 7, Consensus meeting,

Male)

However, several young people were not able to have a private area at home to receive

the therapy, and experienced frequent interruptions:

Interviewer: *What about your situation at home, is it private enough for you?*

Ppt: *Yeah. Go[a]way* [shouting to Brother]. (Ppt 9, Session 7, Male)

This sibling interaction did not seem to affect this young person's experience of the

therapy, but is an important consideration for the therapist when session planning.

Outcome: Online delivery modality retained, with space at the hospital offered for young people unable to have a private space at home to ensure inclusivity (this offer was available but not taken up during the current research).

Key Learning 6: Young people preferred one-to-one therapy

Qualitative theme: Delivery and Receipt of therapy

The co-developers felt that the approach they received, to have one-to-one sessions was optimal:

> *I don't really like the idea of group therapy because everyone has different opinions on things and they have different things that they want to talk about, and, depending on how long the group therapy lasts for, I feel like not everyone gets to have a say.* (Ppt 1, Session 7, Female).

The earlier learnings regarding tailoring the explanations to the individual's circumstances, and adapting activities, also lend themselves to one-to-one therapy.

Outcome: One-to-one therapy to be recommended as the first-choice approach, both directly answering the co-developers' requests, and enabling the flexible manual approach highlighted in Key Learning 1.

Key learning 7: Optional inclusion of parents (or preferred companion) in sessions

Qualitative Theme: Delivery and receipt of therapy

The intervention design intentionally allowed young people to choose the level of

involvement their parents had in the therapy, and young people reflected on this

favourably:

I wouldn't be as open as I probably am. I think if I was to have my parents [it]

would probably be a big argument. (Ppt 1, Session 7, Female)

I like it better just me (Ppt 10, Session 7, Male)

I personally like my mum being with me, but because I'm quite open with those

sort things. But I guess it depends on what sort of relationship you have with

your parents. If you're quite open with them or not, in a way. So, for some

people it can be good, but for some people they may not like it. (Ppt 4,

Consensus meeting, Female)

Outcome: To retain the young people's ability to choose parental/companion

involvement.

Key Learning 8: Views on individual strategies

Qualitative theme: Views on strategies and engagement

Throughout the co-development process, the 'Think aloud' interviews facilitated

feedback on a strategy-by-strategy basis, providing an understanding of how activities

were perceived by the young people. These data were used in real-time to evolve the

therapy manual and were included in the table of changes (see Supplementary Material).

An example of this feedback:

Our bad moments are like thunder or something, but it's trying to remind

ourselves that despite that thunder, that bad weather, that there'll always be that

good bit of sky there. It was one of the best ones ... I didn't really understand it

at first, but after I thought about it a bit more it made more sense to me and I

think, as I think about that more often, and I put that in practical situations, it'll

become more useful. (Ppt 7, Session 5, Male)

Outcome: Data collected on individual strategies in this way, were used to adapt the

manual. This process was a fundamental part of the co-development process, and

enabled the young people to shape the programme of therapy.

Key learning 9: Benefits of the psychological therapy

Qualitative theme: Real world benefits of co-development process

The young people's reflections suggested that they experienced benefits from taking part

in the psychological therapy. For instance, by incorporating strategies into their lives:

I think urge surfing or surfing the urge [strategy]. I happen to like ride along

with it because there's no point trying to stop a wave because it would just be

wasting energy and have a more negative effect. (Ppt 7, Session 6, Male)

Becoming more aware of their internal experiences:

As the sessions have gone on, it's been easier to think about these sort of things.

These analogies have been more clear. And with, like, the time going on, it's

much more easier to think about feelings or easy examples to actually give. (Ppt

7, Session 3, Male)

And feeling that they were able to connect and share more with other people in their

social world:

Well, I'm opening up to people more. I'm talking more about me and how much

will work personally in my life. I thought, I really want to tell my friends I lost

weight. Yeah. (Ppt 8, Session 7, Male)

Outcome: The co-developers report anecdotal positive effects from experiencing the

therapy that should be explored quantitatively in future research trials.

Key Learning 10: Wanting more sessions

Qualitative Theme: Delivery and receipt of therapy

The young people concluded that they would like to have had more therapy sessions

than the seven they received:

I got the impression that it took a few sessions to kind of get going and then

when it was really getting good it was finished. (Ppt 2, Female)

So yeah, I think a bit longer might have, you know, that it might have been quite

good in the next four sessions. (Ppt 2, Female)

I think that it did take a couple of sessions to get used to what we were, like,

talking about and, you know, feeling comfortable to talk to [the therapist]

*because I feel like a few more sessions would have been good. (*Ppt 4, Female).

Expanding on this suggestion, we discussed the spacing of sessions and asked:

would offering a longer duration of therapy yield any additional benefits? However, the

participants favoured the weekly spacing of the current design:

I think if you spread stuff out too much then you kind of lose the kind of train of

thought. I guess there's a more of a struggle to connect stuff from different

sessions and if I thought something, like, on Monday and I had to wait all the

way until, like, Saturday to tell [the therapist]*, that's already a long time. But if it*

was, like, two weeks from then I just think it's, like, too long and there's not

really a lot of opportunity to address immediate problems that you're

facing. (Ppt2, Female)

I think the timing was about right. (Ppt 7, Male)

During the clinical team members consensus meeting, discussion concluded that

a focus on increasing the amount, and the way that participants practice between

sessions to integrate theory learnt during the therapy with life, could support a more

intrinsic approach.

YP do need to be prepared to 'act' (e.g. accepting difficult feelings in the service

of their values). *It's not homework, its facilitating change for them* (tweak

linguistically or the activities themselves to embed in their life not just about

coming back to therapy or clinic appointments). (Psychologist)

However, feedback from the 'Think-Aloud' interviews suggested that additional

work may not be favourably received:

Interviewer: *Would you like a workbook to have alongside these sessions?*

Ppt: *No, I actually don't feel like I would use it.* (Ppt 12, Male, Session 7)

Outcome: The length of the therapy programme and the spacing between sessions

would remain the same. However, there will be an increase to the expectation setting at

the beginning of the programme to clarify how time with the therapist and independent

time in the week to be implementing/practising the skills, come together to develop the

outcome.

Key Learning 11: Session content sometimes felt in contrast to

CEW/COCO approaches

Qualitative theme: Views on strategies and engagement

One young person felt that the sessions were offering a different approach from the main clinic, but that sometimes this felt confusing or that the approaches conflicted with one another.

> *I mean, the fact that you're at the CoCo clinic and kind of being told to kind of count calories and look at the diet and, you know, really focus on the food and take the drugs and so on, with the very clear focus about, you know, losing weight and focusing on that. And then in the counselling* [AIM2Change]*, it's much more about feeling good about yourself, being positive and it's those two things are almost in conflict with each other.* (Ppt 2, Female)

It was suggested that a key outcome of the therapy could be seen to be that attending the therapy helps the young person to navigate the CEW clinics, and the difficult feelings and thoughts that arise during them.

> *How YP may feel when they are weighed* (using ACT as a way to understand the world, clinic or family factors are part of that). (Psychologist)

Outcome: Each therapy session should include a check-in with how the young person is feeling about their weight-loss and the experiences they have had within the main CEW service that week, to ensure that they are being directly supported to understand any seemingly conflicting approaches and any difficult feelings that may have arisen.

Discussion

This paper reports the co-development of AIM2Change, in which young people attending a tier 3 weight management clinic received, and guided the development of, an ACT-based therapy.

Overall, this co-development process, and the young people's view therein, has provided preliminary evidence that an ACT/DNA-V approach is suitable for use in a paediatric weight management setting. In common with the wider paediatric weight management population, several of our co-developers experienced neurodiversity. Our findings are consistent with previous studies that indicate that ACT shows promise for individuals with ASD or ADHD(30, 31). The co-developers demonstrated a good comprehension of the model throughout, and even co-developers who were initially hesitant reported experiencing value in the model and therapeutic relationship during the period of the study.

The multifaceted co-development process yielded valuable insights, with some clear changes that were unanimously agreed and simple to implement, such as the suggestion to enhance the opportunities for self-reflection or to tailor the more generic videos and activities to the individual's context. The importance of this request was understood; tailoring the structure and content of the sessions to individuals' characteristics and preferences is known to be important to outcomes(32), and not meeting participants' individual needs and expectations is a common reason for dropping out of weight management treatments in children(33).

The more complex decisions were discussed during the consensus meetings. These involved discussions around the duration of the therapy, and how to balance the AIM2Change session content with the content already delivered by the clinical MDT. Regarding the duration of the therapy, the co-developers consistently recommended

more therapy sessions. This mirrors the requests for more clinician time raised by

participants during a qualitative review of the main service of the clinic(11). As it was

these requests that initially highlighted the extrinsic nature of the participants'

motivations, it was pertinent to readdress this request, first by establishing expectations

early on in the AIM2Change sessions for the level of engagement required by the young

people to gain maximum benefit from AIM2Change, then secondly, by altering the

content of the final sessions of the co-developed intervention to enhance the

individual's self-determination and confidence to continue the DNA-V approach

independently. Thus, the AIM2Change intervention will provide a strong exit strategy

to continue with the activities and skills participants learned during the intervention, as

opposed to facilitating further clinician (therapist) time. Furthermore, extended clinician

time may make the intervention financially unfeasible for adoption by the NHS.

The issue around the lack of consistency between the approach of AIM2Change

and the focus of behaviour change offered by the CoCO clinic, elicited a nuanced

discussion during the consensus meetings. The main CEW services are limited by time

and limited psychological input. Lillis and Kendra(20) highlight potential barriers

arising from the combination of ACT and standard behavioural therapy, referring to

philosophical and structural contradictions such as non-aligned treatment goals, and

ways of appraising thoughts and encouraging behavioural change. Nonetheless,

literature suggests that the simultaneous application of both approaches may be

feasible(20, 34). AIM2Change has the potential to give young people the time and space

to reflect on the often difficult conversations that they experience at the clinic, and to

better understand how they experience their weight loss journey. It was also advised

that the individual's goals from the CoCo clinic be incorporated into the therapeutic

process, in order to align outcomes, and support the young person to reach the goals set

by clinicians in a sustainable, intrinsically motivated way.

Strengths and limitations

Strengths of this research include the young co-developers' positive attitude

towards the therapist and the strong 'therapeutic alliance' that was built, with an online

format. The alliance is considered essential as it provides an equal and cooperative

footing in the process of change(32), which is aligned with both the ACT approach and

with the intention to foster self-determination rather than a reliance on the therapeutic

'expert'.

The flexible approach to both the content and timings of sessions was a strength

of this work that enabled participants to continue to participate throughout changeable

life circumstances. Participants from a wide range of backgrounds, neurodiversity and

readiness to engage in therapy were retained in this study. This is particularly

noteworthy given that it has previously been indicated that young people from

underserved communities (well represented in our co-developer sample) are more likely

to disengage with paediatric weight management services(33). However, it is regrettable

that delays within the University distribution system meant that loan computer tablets

were unavailable on time and consequently two participants withdrew, who were due to

participate using tablets provided through the study's funding for accessibility. Future

studies, if able to offer the loan of tablets, should ensure streamlined access to the

required technology.

The person-based approach, including co-development therapy sessions

followed by consensus meetings with the young people, their parents, as well as

clinicians and ACT experts, was deemed to be successfully applied by all contributors

including the PPI advisory group. Indeed, this may be the first time the young people

have engaged in such "meta-work" (i.e. experiencing and commenting on the

therapeutic intervention at the same time). The young people were asked to distinguish

between the therapy itself and their reflections and opinions on that therapy; this study

has shown that this process is both possible and successful.

Whilst this study was centred on co-design of a new intervention, participants

did appear to experience personal benefits from participating. In particular, some co-

developers suggested changes in the way they appraised situations, which increased

their self-worth and acceptance, in line with the findings of previous studies(35, 36).

Young people also reported improvements in engagement at school, family and social

interactions, consistent with Tronieri et al.(37), where benefits in family relations and

social life-related quality of life were observed. The young people generalised their

positive experiences, expecting others like them to feel the same way, which offers

promise for the outcomes of a future randomised controlled trial. Members of the PPI

advisory group found benefits in joining the project and agreed that contributing to the

project had been a positive experience. They stated that they had been alerted to a new

way of thinking about managing their weight, and had felt listened to and valued.

Conclusion

In conclusion, the co-developers approved of the use of the ACT DNA-V model,

tailored, jointly with them, for paediatric weight management in AIM2Change, within

this population. Plans for future work include a pilot study to test the feasibility of the

co-developed intervention across two CEW clinics, followed by an RCT in many CEW

clinics in the UK to evaluate the effectiveness of the intervention. The invaluable

insights provided by our co-developers and PPI members throughout this process have

shaped and guided the intervention to ensure it best meets the specific needs of this

population.

Acknowledgments

We thank the young people and their parents who participated in the

AIM2Change co-development project. We thank members of the steering group for

their input into this study (Dr N. Davis, Mr R. Courtney-Tucker, Miss L. Stancheris).

We also thank the members of the Patient and Public Advisory Group for their

invaluable contributions throughout the project. We would also like to acknowledge the

support of Professor A. Gibson and the CUBE team for facilitating the PPI evaluation

through the use of the CUBE, and Dr D. Giri and Dr K. Hawton during the recruitment

stage of the study, and Dr E. Maiz as our valued collaborator.

Competing interests

653   Dr Hinton is employed by both the University of Bristol as a Senior Research

654 Fellow (through which she completed this work) and also on a part-time basis with

655 Oxford Medical Products as a Head of Clinical Research & Operations. Please note the

656 work reported in this paper was conducted fully independently from the industry role.

657 All co-authors have no conflict of interest to declare.

Funding

659   This project is funded by the National Institute for Health and Care Research

660 (NIHR) under its Research for Patient Benefit (RfPB) Intervention (Grant Reference

661 Number NIHR203605). The views expressed are those of the authors and not

662 necessarily those of the NIHR or the Department of Health and Social Care.

References

1. (NCD-RisC) NRFC. Worldwide trends in body-mass index, underweight, overweight, and
obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies
in 128.9 million children, adolescents, and adults. . Lancet. 2017;390:2627–42.
2. World Health Organization. Obesity and overweight. 2024 [Available from:
https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight.
3. Di Cesare M, Sorić M, Bovet P, Miranda J, Bhutta Z, Stevens GA, et al. The
epidemiological burden of obesity in childhood: a worldwide epidemic requiring urgent action. .
BMC Medicine 2019;17:212.
4. Pulgaron ER. Childhood obesity: a review of increased risk for physical and
psychological comorbidities. Clin Ther. 2013;35(1):A18-32.
5. Darnton-Hill I, Nishida C, James WP. A life course approach to diet, nutrition and the
prevention of chronic diseases. Public Health Nutr. 2004;7(1A):101-21.
6. Park MH, Sovio U, Viner RM, Hardy RJ, Kinra S. Overweight in childhood, adolescence
and adulthood and cardiovascular risk in later life: pooled analysis of three british birth cohorts.
PLoS One. 2013;8(7):e70684.
7. Public Health England. Health matters: obesity and the food environment. 2017
[Available from: https://www.gov.uk/government/publications/health-matters-obesity-and-
the-food-environment/health-matters-obesity-and-the-food-environment--2.
8. National Institute for Health and Care Excellence. Obesity: identification, assessment
and management. 2014 [Available from: https://www.nice.org.uk/guidance/cg189.
9. Gungor NK. Overweight and obesity in children and adolescents. J Clin Res Pediatr
Endocrinol. 2014;6(3):129-43.
10. Apperley LJ, Blackburn J, Erlandson-Parry K, Gait L, Laing P, Senniappan S. Childhood
obesity: A review of current and future management options. Clin Endocrinol (Oxf).
2022;96(3):288-301.
11. Cox JS, Searle AJ, Hinton EC, Giri D, Shield JPH. Perceptions of non-successful families
attending a weight-management clinic. Arch Dis Child. 2021;106(4):377-82.
12. RM R, EL D. Self-determination theory: Basic psychological needs in motivation,
development, and wellness: Guilford Publications; 2017.
13. Gillison FB, Standage M, Skevington SM. Relationships among adolescents' weight
perceptions, exercise goals, exercise motivation, quality of life and leisure-time exercise
behaviour: a self-determination theory approach. Health Educ Res. 2006;21(6):836-47.
14. Jensen CD, Duraccio KM, Hunsaker SL, Rancourt D, Kuhl ES, Jelalian E, Wing RR. A
qualitative study of successful adolescent and young adult weight losers: implications for
weight control intervention. Child Obes. 2014;10(6):482-90.
15. Hayes SC, KD S, KG W. Acceptance and commitment therapy: An experiential approach
to behavior change. New York: Guilford Press; 1999.
16. Hayes SC, D B-H, B R. Relational frame theory: A post-Skinnerian account of human
language and cognition. Boston, MA: Springer; 2001.
17. Hayes SC, Luoma JB, Bond FW, Masuda A, Lillis J. Acceptance and commitment
therapy: model, processes and outcomes. Behav Res Ther. 2006;44(1):1-25.
18. Hayes SC, KD S, KG W. Acceptance and commitment therapy: The process and practice
of mindful change. New York, NY: Guilford Press; 2011.
19. Hayes LL, Ciarrochi JV. The thriving adolescent: Using acceptance and commitment
therapy and positive psychology to help teens manage emotions, achieve goals, and build
connection. : New Harbinger Publications; 2015.
20. Lillis J, Kendra KE. Acceptance and commitment therapy for weight control: Model,
evidence, and future directions. J Contextual Behav Sci. 2014;3(1):1-7.
21. Lawlor ER, Islam N, Bates S, Griffin SJ, Hill AJ, Hughes CA, et al. Third-wave cognitive
behaviour therapies for weight management: A systematic review and network meta-analysis.
Obes Rev. 2020;21(7):e13013.

22.   Iturbe I, Echeburua E, Maiz E. The effectiveness of acceptance and commitment
therapy upon weight management and psychological well-being of adults with overweight or
obesity: A systematic review. Clin Psychol Psychother. 2022;29(3):837-56.
23.   Cox JS, Iturbe I, Searle A, Maiz E, Hinton EC. The acceptability, feasibility and
preliminary efficacy of acceptance and commitment therapy for adolescents in the
management of overweight or obesity: A scoping review. Journal of Contextual Behavioral
Science. 2024;31(100713).
24.   Cox JS, Searle A, Thornton G, Hamilton-Shield JP, Hinton EC. Integrating COM-B and the
person-based approach to develop an ACT based therapy programme to raise self-
determination in adolescents with obesity. BMC Health Serv Res. 2023;23(1):1158.
25.   Yardley L, Morrison L, Bradbury K, Muller I. The person-based approach to intervention
development: application to digital health-related behavior change interventions. J Med
Internet Res. 2015;17(1):e30.
26.   Michie S, Atkins L, West R. The behaviour change wheel book - A guide to designing
interventions: Silverback Publishers; 2014.
27.   Gibson A, Welsman J, Britten N. Evaluating patient and public involvement in health
research: from theoretical model to practical workshop. Health Expectations. 2017;20:826–35.
28.   Hinton EC, Fenwick C, Hall M, Bell M, Hamilton-Shield JP, Gibson A. Evaluating the
benefit of early patient and public involvement for product development and testing with
small companies. Health Expect. 2023;26(3):1159-69.
29.   Gale NK, Heath G, Cameron E, Rashid S, Redwood S. Using the framework method for
the analysis of qualitative data in multi-disciplinary health research. BMC Med Res Methodol.
2013;13:117.
30.   Pahnke J, Hirvikoski T, Bjureberg J, Bölte S, Jokinen J, Bohman B, Lundgren T.
Acceptance and commitment therapy for autistic adults: An open pilot study in a psychiatric
outpatient context. Journal of Contextual Behavioral Science. 2019;13:34-41.
31.   Vanzin L, Mauri V, Valli A, Pozzi M, Presti G, Oppo A, et al. Clinical Effects of an ACT-
Group Training in Children and Adolescents with Attention-Deficit/Hyperactivity Disorder.
Journal of Child and Family Studies. 2019;29(4):1070-80.
32.   Walser RD, O'Connell M. Acceptance and commitment therapy and the therapeutic
relationship: Rupture and repair. J Clin Psychol. 2021;77(2):429-40.
33.   Skelton JA, Beech BM. Attrition in paediatric weight management: a review of the
literature and new directions. Obes Rev. 2011;12(5):e273-81.
34.   Forman EM, Butryn ML. A new look at the science of weight control: how acceptance
and commitment strategies can address the challenge of self-regulation. Appetite.
2015;84:171-80.
35.   Cardel MI, Lee AM, Chi X, Newsome F, Miller DR, Bernier A, et al.
Feasibility/acceptability of an acceptance-based therapy intervention for diverse adolescent
girls with overweight/obesity. Obes Sci Pract. 2021;7(3):291-301.
36.   Newsome FA, Cardel MI, Chi X, Lee AM, Miller D, Menon S, et al. Wellness Achieved
Through Changing Habits: A Randomized Controlled Trial of an Acceptance-Based Intervention
for Adolescent Girls With Overweight or Obesity. Child Obes. 2023;19(8):525-34.
37.   Tronieri JS, Wadden TA, Leonard SM, Berkowitz RI. A pilot study of acceptance-based
behavioural weight loss for adolescents with obesity. Behav Cogn Psychother. 2019;47(6):686-
96.

Supporting information

Table of changes

using the Zoom Hand Tool? and full afterwards etc.

One way to be
involved

Many ways to
be involved

Comment

Do you feel you were given a weak or strong voice in this session? A strong voice takes part in discussions and has and impact on decisions made. A weak voice takes part in the conversation but has little chance of making change.

Weak voice

Strong voice

Comment

Who set the agenda and topics of the meeting? Were your thoughts and issues, or the organisers thoughts and issues more important during the session?

Organisations
concerns

Your Concerns

Comment

Do you believe the organisers will make changes based on the comments made today? You can move the marker between 'organisers resistant (unwilling) to change' to 'organisers have been willing to change'

Organisation
not willing to
change

Organisation
willing to
change

Comment

Colour Profile | Default |

Show Other Users ☑

Strong Voice

Weak Voice

Organisation

One Way

Public

Many Ways