The effect of medical explanations from large language models on diagnostic decisions in radiology

Philipp Spitzer[1, ‡], Daniel Hendriks[1, ‡], Jan Rudolph[2], Sarah Schlaeger[2], Jens Ricke[2], Niklas Kühl[3,4†], Boj Friedrich Hoppe[2,5†], and Stefan Feuerriegel[*,5,6,†]

[1]Karlsruhe Institute of Technology, Karlsruhe, Germany

[2]Department of Radiology, LMU University Hospital, LMU Munich, Munich, Germany

[3]University of Bayreuth, Bayreuth, Germany

[4]Fraunhofer FIT, Bayreuth, Germany

[5]Munich Center for Machine Learning, Munich, Germany

[6]LMU Munich, Munich, Germany

[‡]Joint first author

[†]Joint last author

[*]Corresponding author: feuerriegel@lmu.de

Abstract

Large language models (LLMs) are increasingly used by physicians for diagnostic support. A key advantage of LLMs is the ability to generate explanations that can help physicians understand the reasoning behind a diagnosis. However, the best-suited format for LLM-generated explanations remains unclear. In this large-scale study, we examined the effect of different formats for LLM explanations on clinical decision-making. For this, we conducted a randomized experiment with radiologists reviewing patient cases with radiological images ($N = 2020$ assessments). Participants received either no LLM support (control group) or were supported by one of three LLM-generated explanations: (1) a *standard output* providing the diagnosis without explanation; (2) a *differential diagnosis* comparing multiple possible diagnoses; or (3) a *chain-of-thought* explanation offering a detailed reasoning process for the diagnosis. We find that the format of explanations significantly influences diagnostic accuracy. The chain-of-thought explanations yielded the best performance, improving the diagnostic accuracy by 12.2% compared to the control condition without LLM support ($P = 0.001$). The chain-of-thought explanations are also superior to the standard output without explanation (+7.2%; $P = 0.040$) and the differential diagnosis format (+9.7%; $P = 0.004$). Evidently, explaining the reasoning for a diagnosis helps physicians to identify and correct potential errors in LLM predictions and thus improve overall decisions. Altogether, the results highlight the importance of how explanations in medical LLMs are generated to maximize their utility in clinical practice. By designing explanations to support the reasoning processes of physicians, LLMs can improve diagnostic performance and, ultimately, patient outcomes.

Introduction

Large language models (LLMs), a form of artificial intelligence (AI) capable of producing human-like text, are becoming increasingly common in clinical workflows [1, 2], including tasks such as assisting with diagnosis [3–7]. A particular promise of LLMs is to improve patient outcomes by reducing diagnostic errors [8], a major source of preventable harm [9–11], but diagnostic errors still remain a major concern, with nearly 800,000 U.S. patients dying or suffering permanent disability each year as a result [9].

Existing research has extensively evaluated the diagnostic accuracy of LLMs in clinical practice, often finding that LLMs perform as well as human physicians and, in some cases, achieve superhuman performance [5, 6, 12–14]. Other studies have examined the diagnostic accuracy of physicians when supported by LLMs compared to when they work without LLM assistance [4, 7, 12, 15]. However, these studies primarily assess the correctness of LLM-generated diagnoses but neglect how explanations provided by the LLMs influence the decision-making of physicians. Notwithstanding, a key strength of LLMs is that they can explain in natural language why a certain diagnosis seems likely. Such explanations can help physicians understand and verify the rationale behind a suggested diagnosis. Yet, the impact of different explanation formats on medical decision-making is unclear. Thus, we analyze how different formats for LLM-generated explanations influence the diagnostic accuracy of physicians.

We hypothesize that the format of LLM-generated explanations affects the diagnostic accuracy of physicians. On the one hand, explanations may improve diagnostic accuracy by helping physicians understand the reasoning behind a diagnosis. For instance, explanations with a differential diagnosis align with standard clinical practice where differential diagnoses help in reducing biases and detect rare diseases [16, 17]. Similarly, LLMs can generate explanations through chain-of-thought reasoning [18], where the LLM is asked to provide a step-by-step explanation so that a physician can verify its plausibility against their domain knowledge. On the other hand, explana-

tions can also mislead physicians [19–21]. A key risk is automation bias, where humans overly trust machine-generated outputs such as LLM-generated advice [22, 23]. This risk is particularly pronounced for LLM-generated explanations, as LLMs tend to produce explanations that sound convincing but contain errors, a phenomenon known as "hallucinations". Consequently, explanations, especially if detailed but incorrect, may also lead to diagnostic errors. Overall, LLM-generated explanations may support or hinder medical decision-making. However, the best-suited design of LLM-generated explanations is unclear.

Here, we analyzed the effect of different formats of LLM-generated explanations on the diagnostic accuracy of physicians (Figure 1). To do so, we conducted a randomized experiment in which radiologists were tasked to diagnose patients based on clinical vignettes containing both text and radiological images ($N = 2020$ assessments). Using a between-subject design, we varied the *explanation format*. We used the following three treatments: (1) a *standard output* that offered a diagnosis without explanation; (2) a *differential diagnosis*, which outputted the five most likely differential diagnoses in descending order; and (3) a *chain-of-thought explanation*, which detailed the reasoning process behind the diagnosis. Additionally, a control group performed the task without LLM support. We examined how the explanation formats influenced the diagnostic accuracy of physicians. Furthermore, we assessed how often doctors followed LLM advice or overrode (in)correct diagnostic suggestions. The results show that chain-of-thought explanations led to the best overall diagnostic accuracy. With these findings, we aim to provide recommendations for *how* to design explanations in medical LLMs to maximize clinical utility.

Figure 1: **Research design. a**, We selected 20 real-world patient cases from the *New England Journal of Medicine Image Challenge* [24]. Each patient case includes a textual description and at least one radiological image (i.e., computed tomography [CT] or magnetic resonance imaging [MRI]). **b**, We recruited 101 radiologists from the U.S. who were asked to provide diagnoses for each case and who were randomly assigned to either a control group (no LLM support) or one of three treatment groups (with LLM support). **c**, We designed prompts to produce LLM-based explanations with different formats (i.e., *standard output*, *differential diagnosis*, and *chain-of-thought*). The explanations were generated via a state-of-the-art LLM able to handle multi-modal inputs (GPT-4). However, because the LLM can make errors, both the suggested diagnoses and explanations may sometimes be incorrect. **d**, Our primary outcome is diagnostic accuracy, measured by the proportion of correct diagnoses per radiologist. As secondary outcomes, we assess how frequently radiologists adhere to correct LLM advice or overrode incorrect LLM advice.

Results

Experiment

To analyze how different formats of LLM-generated explanations affect diagnostic accuracy, we conducted a randomized experiment with 101 radiologists from the U.S. The physicians had a mean of 13.6 years (standard deviation [SD] = 8.0) of medical experience. Each radiologist was presented with 20 real-world patient cases (mean length: 55.4 words; SD = 18.8), each consisting of a brief clinical description and at least one radiological image (i.e., computed tomography [CT] or magnetic resonance imaging [MRI]), and was then asked to provide a diagnosis for the patient ($N = 2020$ assessments overall). Participants were randomly assigned to either a control group with no LLM support or one of three treatment groups, each with a different form of LLM advice.

In the three treatment groups, physicians received advice through different types of explanations generated by GPT-4, a state-of-the-art multi-modal LLM [25, 26]. (1) *Standard output*. Here, a single best-guess diagnosis was provided, but the explanation is either absent, very brief, or paraphrasing the case description. (2) *Differential diagnosis*. Here, multiple plausible diagnoses were listed, each with a short justification, ordered by decreasing likelihood. This explanation format mirrors typical clinical reasoning, where multiple conditions are considered [27]. (3) *Chain-of-thought*. Inspired by best practice in LLM research [18], an explanation was produced that provides a step-by-step rationale of how the LLM arrives at the final recommendation, which can give physicians a deeper understanding of the reasoning process and help them identify potential errors. All outputs were kept as generated, including any diagnostic errors or hallucinations, to mimic real-world LLM use in medical practice. Example outputs for each treatment group are shown in Figure 1 (further example outputs are shown in Supplementary Figures S1 to S3). The length of the LLM-generated outputs exhibited substantial variation across the different output formats (standard output: mean = 62.7, SD = 12.5; differential diagnosis: mean = 208.6, SD = 20.2;

chain-of-thought: mean = 188.6, SD = 26.2; see Supplementary Figure S5).

The control group (with no LLM advice) achieved a diagnostic accuracy of 61.7%. The baseline diagnostic accuracy of the LLM advice remained relatively consistent across all 20 patient cases but with slight variability (standard output: 75%; differential diagnosis: 65% for top-1 answer and 80% for top-5; chain-of-thought: 80%; Supplementary Figure S4). This corresponds to an average diagnostic accuracy of 78.3% across all conditions (based on the top-5 accuracy for the differential diagnosis; 73.3% based on the top-1 accuracy of the differential diagnosis). To assess the benefits of different prompting strategies for generating correct LLM outputs, we estimated a logistic regression with conditions as independent variables. The LLM with chain-of-thought prompting performed better than the standard output (coef = -0.288, 95% CI = $[-1.779, 1.204]$, $P = 0.705$), and the LLM with chain-of-thought output performed similar to the LLM with differential diagnosis output (coef = 0.000, 95% CI = $[-1.549, 1.549]$, $P = 1.000$) (see Supplementary Table S5).

Effect on diagnostic accuracy

Physicians augmented with LLM advice outperformed those in the control group without LLM advice (Fig. 2 for statistical comparisons based on Welch's t-tests). To calculate effect sizes, we used an ordinary least squares (OLS) regression (Supplementary Table S6). Here, the diagnostic accuracy of physicians with standard output was comparable to the diagnostic accuracy of the control group (difference = 5.0 percentage points, 95% CI = $[-1.8; 11.8]$, $P = 0.150$). Similarly, the diagnostic accuracy of physicians supported by a differential diagnosis was comparable to the control group (difference = 2.5 percentage points, 95% CI = $[-4.0, 9.0]$, $P = 0.446$). In contrast, physicians in the chain-of-thought group performed best with a 12.2 percentage point improvement over the control group (95% CI = $[5.3, 19.2]$, $P = 0.001$). Separate comparisons for the chain-of-thought versus standard condition and chain-of-thought versus differential diagnosis condition showed improvements in the outcome measure by 7.2 percentage points (95% CI = $[0.3, 14.2]$,

$P = 0.040$) and 9.7 percentage points (95% CI = $[3.2, 16.3]$, $P = 0.004$), respectively, which confirms that the chain-of-thought condition performed best (Supplementary Table S7).

Physicians with chain-of-though explanations outperformed both the control group and treatment group supported by the standard output even after controlling for various physician-specific control variables such as years of medical experience, radiology-specific expertise, hours per week spent on visual inspections, IT skills, and experience with medical AI (see Supplementary Table S8 for the regression results). The findings also hold when accounting for the time physicians spent to complete the study, the length of the LLM outputs, and the length of the answers entered by participants (see Supplementary Table S9). A breakdown with the diagnostic accuracy for each patient case is in Supplementary Fig. S7.

Figure 2: **Diagnostic accuracy across different LLM explanations.** The figure shows the distribution of diagnostic accuracy for all participants by condition as violin plots (which illustrate the probability density of the data). The boxplots within each violin indicate the 25% and 75% quartiles, with the median represented by the thick center line. For the differential diagnosis condition, the top-5 diagnostic accuracy is reported (the top-1 diagnostic accuracy is lower and corresponds to 65%). Statistical significance was assessed using Welch's t-test to compare diagnostic accuracy between the treatment groups (in the main text, we report results from a regression analysis to obtain effect sizes).

To control for the baseline diagnostic accuracy across different prompting strategies, we performed two further analyses. (1) We restricted our analysis to the subset of patient cases where the LLM advice was correct (or incorrect) across all conditions. Here, the control group had the lowest diagnostic accuracy (77.60%), compared to the standard output group (89.58%), the differential diagnosis group (95.83%), and the chain-of-thought explanation (94.02%) (Supplementary Figure S8). To compare the chain-of-thought explanations with the other conditions, we again conducted an OLS regression. The diagnostic accuracy of the chain-of-thought explanation group was higher than the control group (difference = 16.4 percentage points, 95% CI = [9.1, 23.8], $P < 0.001$). The diagnostic accuracy of the chain-of-thought explanation group was similar to

the standard output group (difference = 4.4 percentage points, 95% CI = $[-2.9, 11.8]$, $P = 0.233$) and to the differential diagnosis group (difference = -1.8 percentage points, 95% CI = $[-8.8, 5.2]$, $P = 0.607$). The regression results are reported in Supplementary Table S23. (2) We also controlled for the correctness of the LLM diagnoses and the correctness of the LLM explanations in a logistic regression model. The standard group and the differential diagnosis group were outperformed by the chain-of-thought group (coef = 0.504, 95% CI = $[0.094, 0.914]$, $P = 0.016$; and coef = 1.014, 95% CI = $[0.646, 1.383]$, $P < 0.001$; respectively) (see Supplementary Table S24).

Mechanism of adherence and overriding

To explore adherence and overriding[1] behavior, we examined the extent to which participants followed correct or incorrect LLM advice (Fig. 3). For differential diagnosis, we assessed adherence based on the top-1 recommendation (results for top-5 recommendations are consistent; see Supplementary Figure S10). When the LLM output was incorrect, adherence was highest in the differential diagnosis group (74.3%), significantly exceeding that of the standard output (33.3%, $P < 0.001$ based on a two-sided Welch's t-test) and chain-of-thought groups (53.3%, $P < 0.001$). The adherence in the chain-of-thought group was also significantly higher than in the standard group ($P = 0.034$). Thus, physicians in the differential diagnosis group appeared more inclined to follow the LLM advice even when it was wrong. When the LLM output was correct, the mechanism for adherence vs. overriding was similar. Physicians again showed the highest adherence in the differential diagnosis group (84.7%) in comparison to both the chain-of-though group (64.7%, $P < 0.001$) and the standard output group (56.4%, $P < 0.001$). The comparison between the chain-of-thought group and the standard group was also significant ($P < 0.001$). In sum, physicians supported with differential diagnosis explanations had consistently high adherence regardless of whether the LLM was correct or incorrect, thus often failing to override when the LLM advice

---

[1]We follow the terminology of literature on human-computer interaction and use the term "override" to describe the scenario in which a human is not adhering to an AI-generated recommendation, regardless of whether this behavior is correct [28].

was incorrect. Conversely, the chain-of-thought explanation (in comparison to the standard output) led to a more selective adherence, where physicians were more likely to follow correct LLM advice.

Figure 3: **Adherence vs. overriding LLM advice.** Breakdown of adherence for **a**, correct LLM advice and **b**, incorrect LLM advice. Here, *adherence* was calculated as the proportion of patient cases in which participants' diagnoses matched the LLM-generated diagnoses. Correspondingly, *overriding* was defined as 1− adherence. Reported is the adherence to the top-1 answer in the differential diagnosis explanation. Results for the top-5 answers are in Supplementary Figure S10. We observe a strong adherence to LLM-generated differential diagnosis in both cases of correct and wrong advice (84.7% and 74.3%), in contrast to other explanation formats.

Heterogeneity across physicians and patient cases

To examine heterogeneity across physicians, we analyzed diagnostic accuracy across physician subgroups with different IT skills and medical tenure (Fig. 4a,b). Overall, the chain-of-thought

group tends to achieve a higher diagnostic accuracy than both the control and the differential diagnosis groups. A similar trend was observed when accounting for the complexity of patient cases (Fig. 4c).

Figure 4: **Heterogeneity across physicians and patient cases. a**, Diagnostic accuracy was compared between different levels of IT skills, as defined by a post-hoc survey. Participants were classified as having advanced IT skills ($n = 56$) when reporting "very good", while all others were classified as having basic IT skills ($n = 43$). **b**, Diagnostic accuracy was compared across different numbers of years with medical expertise in radiology ($n = 46$ have a tenure of < 12 years, while $n = 53$ have tenure of ≥ 12 years). **c**, Diagnostic accuracy for basic and complex patient cases was assessed by dividing cases into two equally sized subsets based on the median diagnostic accuracy observed in the control group. The boxplots within each violin indicate the 25% and 75% quartiles, with the median represented by the thick center line. Statistical significance was assessed using Welch's t-test. Estimated effect sizes using regression analysis are in Supplementary Tables S11 to S15.

Decision time

Physicians randomized to LLM support spent more time per case (see Supplementary Figure S6). The control group had the lowest decision time with 79.4 seconds per patient case. To calculate the effect on decision time for the different explanation formats compared to the control group, we again estimated an OLS regression model (Supplementary Table S19). The decision time per patient case was 92.5 seconds for the standard output, similar to the control group (difference = 13.1 seconds, 95% CI = $[-73.2, 99.4]$, $P = 0.763$). The differential diagnosis group and the chain-of-thought group had an average decision time per patient case of 147.6 and 137.2 seconds (difference = 68.3 seconds 95% CI = $[-18.0, 154.7]$, $P = 0.119$; difference = 57.8 seconds, 95% CI = $[-28.5, 144.2]$, $P = 0.186$; respectively). Additionally, we estimated an OLS regression model to compare the standard output group against (i) the differential diagnosis group and (ii) the chain-of-thought explanation group (Supplementary Table S20). The differential diagnosis group and the chain-of-thought group had a similar average decision time per patient case compared to the standard output group (difference = 61.8 seconds, 95% CI = $[-12.5, 136.1]$, $P = 0.102$; and difference = 51.3 seconds, 95% CI = $[-23.0, 125.6]$, $P = 0.173$; respectively).

Discussion

The rapid adoption of LLMs in clinical settings [29] raises the question of how physicians and LLMs can most effectively collaborate to improve patient outcomes. While LLMs can help physicians verify their advice by providing explanations, the best-suited design of such explanations remains unclear. To use LLMs effectively in medical practice, our experiment demonstrates that the format of explanation is crucial. Although our results echo prior research showing that LLM-augmented physicians ("human-in-the-loop") generally outperform those without LLM support [4, 7, 12, 15], we uncover a previously overlooked source of heterogeneity: when physicians receive carefully prompted LLM-generated explanations, their diagnostic accuracy significantly

exceeds that of a control group with no LLM support. Surprisingly, chain-of-thought explanations provide the largest benefit to diagnostic performance. Overall, the effects were consistent across physicians with varying levels of medical experience and IT proficiency.

The key mechanism behind the performance gain from chain-of-thought explanations appears to be the physicians' improved ability to assess the correctness of LLM output based on the provided reasoning. This aligns with cognitive psychology research, which suggests that structured reasoning can enhance judgment and decision-making by making thought processes more transparent [30]. The step-by-step reasoning in the chain-of-thought output helps users follow correct LLM advice (correct AI-reliance) and override flawed suggestions (correct self-reliance), consistent with research suggesting that chain-of-thought prompts help improve the reasoning of LLMs [18]. In contrast, explanations with a differential diagnosis seem to be misleading, sometimes inducing high adherence to LLM advice, even when the LLM is incorrect (often referred to as automation bias). Notwithstanding, the differential diagnosis format may be appropriate when the LLM accuracy is high and, thus, could be more relevant in future LLMs with overall higher baseline diagnostic accuracy. Here, it would be interesting to explore how careful training of physicians may help reduce existing wrong adherence and overriding behavior. Another potential factor is that diagnostic accuracy and decision time are interrelated [7]. Although decision times were longer for physicians supported by a differential diagnosis, the effects of increased performance from chain-of-thought explanations remained statistically significant after controlling for decision time. Finally, because the perceived helpfulness was similar across treatment groups (Supplementary Figure S11), the benefits from LLM advice seem unrelated to subjective factors or preferences. Instead, the benefits appear to originate from how the explanation format enables physicians to more effectively evaluate LLM advice and incorporate it into their own decision-making processes.

In general, prompt design is known to influence the correctness of LLM outputs [18]. In our study, diagnostic accuracy varied with the prompting strategy: the chain-of-thought strategy improved the accuracy of the LLM output by 5 percentage points relative to a naïve strategy with

a default prompt and thus no prompt tuning. This finding aligns with best practice recommendations [18, 31] and underscores the importance of carefully fine-tuning prompts for clinical applications. Another implication of this finding is the need to boost AI literacy among physicians to help them design effective prompts, especially in light of the fact that the standard strategy for LLM deployment is one without formal training in prompt strategies [32]. Further, while the varying levels of diagnostic accuracy across different prompting strategies may have initially complicated direct comparisons, this was done intentionally and is a key strength of our study because we mimic a real-world scenario in medical LLM deployment. Nonetheless, our robustness checks confirmed that the effect of explanation formats remained robust even after adjusting for the diagnostic accuracy of the prompting strategies (see Supplementary Table S24).

Previous medical research has primarily focused on comparing the diagnostic accuracy between physicians augmented with LLM support versus those without it [4, 7, 12, 15], yet without isolating the effect of prompt design. As a result, the impact of explanation formats on diagnostic decision-making has remained unclear. Even though the importance of *how* information is presented in medical AI applications is increasingly acknowledged [33–35], empirical research focusing on decision-making is limited. Some work has explored visual annotations generated by AI tools to assist in medical image interpretation [36–38], but these studies primarily address search-and-locate tasks (e.g., lesion detection) and use outputs different from ours (e.g., heatmaps). In contrast, our study examines how different LLM explanation formats can improve diagnostic decisions.

This study has multiple limitations that present interesting directions for future research. First, we focused on only one medical specialty; however, diagnostic decisions are central to radiology and common in routine medical practice [39, 40]. Moreover, the set of patient cases covers a broad range of typical radiological assessments. Future research could explore to what extent our findings extend to other medical specialties. Second, our analysis focused on a single time point; longitudinal studies are needed to analyze the long-term effect of LLM assistance on medical

practice. Third, while we evaluated diagnostic accuracy, we did not assess the potential harm that can arise from incorrect decisions or other patient outcomes. Fourth, although some physicians in the control group reported having accessed the Internet analogous to routine practice, none indicated using LLMs, suggesting that the risk of conflated results is low. Fifth, our findings rely on GPT-4, a state-of-the-art multi-modal LLM whose diagnostic accuracy is similar to that of clinical experts [6,41,42]. However, the LLM may have seen some of the cases during training but still produced incorrect diagnoses, indicating that the tasks were sufficiently challenging. Future LLMs may generate even more informative explanations, but we also acknowledge the inherent limitations of LLMs, such as their tendency to produce hallucinations or permeating biases [43,44]. Hence, the real-world implementation of LLM in medical settings requires a careful approach considering these risks.

Still, misdiagnosis is a common issue in medical practice [9], including in radiology [39,40]. While LLMs hold promise for reducing diagnostic errors [8], designing systems that effectively support collaboration with physicians remains challenging. Our findings show that the format of LLM explanations has a large impact on decision performance, thereby highlighting the need for a human-centered approach to designing real-world LLM implementations in order to maximize clinical utility and ultimately improve patient outcomes. Future LLMs may provide even more informative explanations and thereby further improve diagnostic accuracy.

Methods

This work analyzes how the diagnostic accuracy of physicians is affected by different explanation formats of LLMs. We preregistered our hypotheses (see https://aspredicted.org/4tgb-sr3z.pdf) and tested them in a randomized experiment using a between-subject design. We complied with all local ethical regulations. The research design was approved by the Ethics Commission of LMU Munich (EK-MIS-2024-320). All participants provided informed consent.

Procedure

To evaluate the effect of different explanation formats on the diagnostic accuracy of physicians, we designed the following experiment. First, participants received information about the study objectives. After giving informed consent, each participant was introduced to the diagnostic task and randomly assigned to one of four groups: a control group ($n = 24$) or one of three treatment groups—namely the standard output group ($n = 24$), the differential diagnosis group ($n = 30$), or the chain-of-thought group ($n = 23$). Only participants in the treatment groups received LLM-based advice. Once the participants completed the diagnostic tasks, they were asked to fill out a post-task survey. The survey collected socio-demographic information (e.g., experience in medicine in years, experience specifically in radiology in years) and asked general questions about the task (e.g., helpfulness of the LLM support). A full list of the survey items is in Supplementary Table S2.

After giving informed consent, participants had to pass a pre-registered attention check to ensure that all participants actively engaged with the task. Afterward, each participant was shown 20 cases and then asked to provide a diagnosis in the form of an open-ended text. The responses were manually coded by the author team to correct for minor typos. The cases were shown in random order to minimize bias from potential learning effects or fatigue. Each case consisted of a descriptive text plus at least one (and sometimes two) radiological images (i.e., computed to-

mography [CT] or magnetic resonance imaging [MRI]). Participants completed the tasks without a maximum time limit (a minimum time limit of 10 seconds was enforced for each task to ensure high-quality answers).

Patient cases

All patient cases included in this study were sourced from the *New England Journal of Medicine Image Challenge* [24], a high-quality collection of peer-reviewed cases, including both common and rare diagnoses. Each case consists of a short text describing the patient's characteristics and symptoms, along with at least one medical image, such as photographs of dermatological findings, endoscopic images, histopathological slides, or radiological images.

In the first step, a set of only radiological cases and images were randomly sampled and the difficulty of diagnosing each case was rated by a panel of three radiologists (J.Ru., S.S., and B.F.H.) on a 5-point Likert scale. Of those, we selected 20 cases to reflect a broad spectrum of possible diagnoses and levels of difficulty, but omitted cases that were rated as either too easy or too difficult (e.g., very rare diagnoses only based on case reports). This ensured a broad and diverse set of diagnostic challenges commonly encountered in radiology. Simpler patient cases included, e.g., disseminated brain metastasis or hepatic echinococcosis, while more complex cases included, e.g., disseminated mycobacterium avium complex (MAC) infection or infantile hepatic hemangiomas. The median word count of the texts was 54.5 words, with small variability (minimum: 24 words; maximum: 85 words). Of the 20 cases, 15 include one radiological image, while the remaining 5 include two. Detailed case descriptions are provided in Supplementary Table S3 and exemplary cases are shown in Supplementary Figures S1 to S3.

LLM advice

Participants in the treatment groups received advice generated by an LLM (GPT-4, version 2024-02-15-preview, temperature = 0.7, accessed via Microsoft Azure) based on the patient's text and

imaging data. We followed best practice in prompt design [18, 31] to produce different output formats depending on the treatment groups (see Supplementary Table S1 for the exact prompts and Supplementary Figures S1 to S3 for example outputs):

1. *Standard output* presents a single diagnosis along with either no explanation or only a concise rationale that typically mirrors the input prompt. This approach reflects LLM use cases in which no custom prompting strategy is employed. Furthermore, it aligns with the format commonly adopted in benchmarking studies from LLM research [5, 12], where the focus is primarily to assess the diagnostic accuracy of the LLM.

2. *Differential diagnosis* lists multiple possible diagnoses (here: the top five diagnoses) ordered by likelihood from most likely to less likely. This output format is motivated by that differential diagnosis is a common approach in routine medical practice [16, 17] and further acknowledges that LLMs should perform better when the top-k predictions are considered rather than the top-1 prediction. Intuitively, such a differential diagnosis should help physicians consider alternative diagnoses, especially for rare diseases. Unless stated otherwise, we report diagnostic accuracy for the top-5 answers.

3. *Chain-of-thought explanation* provides a detailed, step-by-step description of the reasoning process behind how the diagnosis is made. Intuitively, this is motivated by that, on the one hand, research in machine learning suggesting that chain-of-thought prompting can improve performance [18] and, on the other hand, such format should help physicians in comparing the diagnostic process against their domain knowledge and thus identify—and eventually— override errors in the LLM advise.

We selected GPT-4 (version: 2024-02-15-preview) due to its state-of-the-art performance for diagnostic reasoning capabilities [6, 12, 13]. Further, GPT-4 is designed to handle multi-modal inputs, which allows us to process both textual descriptions and radiological images. A compar-

ison regarding the diagnostic accuracy for another LLM is in Supplementary Figure S16, which suggests that, while other LLMs perform similarly overall, these are slightly suboptimal.

Outcome

To evaluate the effect of different explanation formats, the primary outcome for our study was the *diagnostic accuracy* at the physician level. Specifically, we mapped the answers onto a dichotomous value ($= 1$ if the diagnosis is correct and 0 otherwise) and then averaged the values across all 20 patient cases for each participant. When determining whether answers were correct, we only included fully correct answers. In an additional analysis, we looked at answers, which were partially correct, but missing important details for full diagnosis (e.g., in case 2, "metabolic toxicity" is partially correct, but the underlying cause of "manganese poisoning" is missing; or, in case 3, where "gastric outlet" is partially correct, but the "superior mesenteric artery (sma) syndrome" as the cause of the obstruction is missing) and performed a robustness check where such partially correct responses were counted as correct and arrived at similar findings (see Supplementary Figure S9).

We also collected additional measures through a post-task survey to assess the perceived interactions with the LLM support (e.g., helpfulness). The survey items are listed in Table S2, which we used to analyze the perceptions of different explanations (see Supplementary Figures S11 to S15).

Study participants

The study was conducted online (November 21, 2025 – December 1, 2025) using the online platform Qualtrics (Qualtrics International Inc.). We recruited physicians specializing in radiology from the U.S. via MSI-ACI. Participants were compensated by MSI-ACI with a fair payment above the U.S. minimum wage. No additional performance-based financial incentives were provided to preserve the ecological validity of the study and minimize potential biases due to extrinsic motivations, thereby mimicking the incentive structure of decision-making in medical practice [37].

Participants were required to meet the following inclusion criteria: They had to be licensed

physicians specialized in radiology, consent to the study, and have access to a device with internet

access to participate in the study. Following pre-registered exclusion criteria, we then excluded

participants who did not complete the study and failed the attention check.

A total of 156 participants consented to join the study, of whom 101 completed the tasks and

met the inclusion criteria, which thus forms the final sample for analysis. The reasons for exclusion

were incomplete participation ($n = 42$), failure to pass the attention checks ($n = 11$), or duplicate

entries ($n = 2$). Detailed socio-demographic information and professional characteristics are re-

ported in Supplement Table S4.

Statistical analysis

To compare the diagnostic accuracy across different conditions, we used analyses of variance

(ANOVA) and one-sided Welch's t-tests. This approach was selected due to its robustness against

unequal variances and different sample sizes between groups. A significance level of $\alpha = 0.05$

was used for all statistical comparisons.

To calculate effect sizes comparing the explanations, we estimated an OLS regression model

at the physician level. The outcome was the diagnostic accuracy, modeled as a function of expla-

nation type, with the control group as the reference and separate variables to capture the effects

for groups supported by standard output, differential diagnosis, and chain-of-thought explanations.

To estimate the effect of different prompting strategies on both the diagnostic accuracy of the

LLM-only approach (without human involvement) and the decision time, we estimate a logistic

regression at the assessment level. We report the coefficients on the log-odds scale for reasons of

comparability.

All analyses were conducted in Python (3.11.11) using the packages `numpy` (2.2.1), `pandas`

(2.2.3), `scipy` (1.15.1), `pymer4` (0.8.2), and `statsmodels` (0.14.4). The data visualizations

were created with `seaborn` (0.13.2) and `matplotlib` (3.10.0).

Robustness checks

We conducted an extensive series of robustness checks to ensure the validity and reliability of our findings. (1) We repeated the regression models while incorporating various participant-specific controls to benefit from increased power (Supplementary Table S8). (2) We further controlled for the length of participants' answers (in the number of characters), the length of the LLM-generated advice (in the number of words), and the average duration of participants to complete the 20 patient cases (in minutes). The analysis results are shown in Supplementary Table S9. (3) To mitigate the influence of extreme outliers in our regression, we winsorized the diagnostic accuracy of study participants by limiting the extreme values in the data to the 5th and 95th percentiles (Supplementary Table S16). (4) To account for diagnostic ambiguity (e.g., when a diagnosis appears accurate but omits a specific sub-diagnosis necessary for determining the appropriate treatment strategy), our initial coding scheme classified partially correct responses as incorrect. We now repeated the analysis and counted such responses as correct (Supplementary Figure S9). (5) We estimated a mixed-effects model to control for heterogeneity across physicians and tasks (see Supplementary Table S17). (6) We estimated a quasi-binomial regression, which may limit the interpretability of the estimated effect sizes but which can better accommodate for the fact that the diagnostic accuracy is in the range between 0% and 100% (see Supplementary Table S18). (7) We controlled for the characteristics of radiological images provided in the patient cases, including the number of images in the case, the source of image (i.e., CT or MRI), and the body region displayed in the image (i.e., abdomen, head, spine, or thorax) (see Supplementary Table S25). Across all robustness checks, the findings remained consistent with the main results.

In principle, different prompting strategies can influence the accuracy of LLM outputs. While such variability reflects real-world LLM use, potential differences in accuracy across different prompting strategies may decline with future LLMs. Thus, to remove bias from different prompting strategies, we performed two additional analyses. (1) We restricted our analysis to the subset of

patient cases where LLM outputs had comparable accuracy (Supplementary Table S23). While this approach allows for direct comparability, it can reduce both statistical power and representativeness of the case selection. (2) We controlled for the diagnostic accuracy of the LLM advice within each treatment arm, which allows us to yield "ceteris paribus" estimates by factoring out variability in LLM performance (Supplementary Table S24). Both approaches yielded qualitatively similar conclusions.

References

[1] Thirunavukarasu, A. J. *et al.* Large language models in medicine. *Nature Medicine* **29**, 1930–1940 (2023).

[2] Williams, C. Y. K., Miao, B. Y., Kornblith, A. E. & Butte, A. J. Evaluating the use of large language models to provide clinical recommendations in the Emergency Department. *Nature Communications* **15**, 8236 (2024).

[3] Eriksen, A. V., Möller, S. & Ryg, J. Use of GPT-4 to diagnose complex clinical cases. *NEJM AI* **1**, AIp2300031 (2024).

[4] Goh, E. *et al.* Large language model influence on diagnostic reasoning: a randomized clinical trial. *JAMA Network Open* **7**, e2440969 (2024).

[5] Hager, P. *et al.* Evaluation and mitigation of the limitations of large language models in clinical decision-making. *Nature Medicine* **30**, 2613–2622 (2024).

[6] Kanjee, Z., Crowe, B. & Rodman, A. Accuracy of a generative artificial intelligence model in a complex diagnostic challenge. *JAMA* **330**, 78–80 (2023).

[7] Goh, E. *et al.* GPT-4 assistance for improvement of physician performance on patient care tasks: A randomized controlled trial. *Nature Medicine* (2025).

[8] Topol, E. J. Toward the eradication of medical diagnostic errors. *Science* **383**, eadn9602 (2024).

[9] Newman-Toker, D. E. *et al.* Burden of serious harms from diagnostic error in the USA. *BMJ Quality & Safety* **33**, 109–120 (2024).

[10] Bates, D. W. *et al.* The safety of inpatient health care. *New England Journal of Medicine* **388**, 142–153 (2023).

[11] Agency for Healthcare Research and Quality. Diagnostic errors in the emergency department: A systematic review. URL https://effectivehealthcare.ahrq.gov/products/ diagnostic-errors-emergency-updated/research.

[12] Brodeur, P. G. *et al.* Superhuman performance of a large language model on the reasoning tasks of a physician (2024). URL http://arxiv.org/abs/2412.10849.

[13] Wu, S. *et al.* Benchmarking open-source large language models, GPT-4 and Claude 2 on multiple-choice questions in nephrology. *NEJM AI* **1**, AIdbp2300092 (2024).

[14] Rodman, A., Buckley, T. A., Manrai, A. K. & Morgan, D. J. Artificial intelligence vs clinician performance in estimating probabilities of diagnoses before and after testing. *JAMA Network Open* **6**, e2347075 (2023).

[15] Huang, A. S., Hirabayashi, K., Barna, L., Parikh, D. & Pasquale, L. R. Assessment of a large language model's responses to questions and cases about glaucoma and retina management. *JAMA Ophthalmology* **142**, 371–375 (2024).

[16] Sox, H. C. (ed.) *Medical decision making* (American College of Physicians, Philadelphia, 2007), ACP Press, 2007 edn.

[17] Richardson, W. S. *et al.* Users' guides to the medical literature: How to use an article about disease probability for differential diagnosis. *JAMA* **281**, 1214–1219 (1999).

[18] Wei, J. *et al.* Chain-of-Thought prompting elicits reasoning in large language models. In *Advances in Neural Information Processing Systems* (2022).

[19] Zack, T. *et al.* Assessing the potential of GPT-4 to perpetuate racial and gender biases in health care: A model evaluation study. *The Lancet Digital Health* **6**, e12–e22 (2024).

[20] Gaube, S. *et al.* Do as AI say: Susceptibility in deployment of clinical decision-aids. *npj Digital Medicine* **4**, 31 (2021).

[21] Yu, F. *et al.* Heterogeneity and predictors of the effects of AI assistance on radiologists. *Nature Medicine* **30**, 837–849 (2024).

[22] Lyell, D. & Coiera, E. Automation bias and verification complexity: A systematic review. *Journal of the American Medical Informatics Association* **24**, 423–431 (2017).

[23] Vered, M., Livni, T., Howe, P. D. L., Miller, T. & Sonenberg, L. The effects of explanations on automation bias. *Artificial Intelligence* **322**, 103952 (2023).

[24] The New England Journal of Medicine: Image Challenge (2025). URL https://www.nejm.org/image-challenge.

[25] Lu, M. Y. *et al.* A multimodal generative AI copilot for human pathology. *Nature* **634**, 466–473 (2024).

[26] Chen, D. *et al.* Performance of multimodal artificial intelligence chatbots evaluated on clinical oncology cases. *JAMA Network Open* **7**, e2437711 (2024).

[27] Liu, Y. *et al.* A deep learning system for differential diagnosis of skin diseases. *Nature Medicine* **26**, 900–908 (2020).

[28] Chen, V., Liao, Q. V., Wortman Vaughan, J. & Bansal, G. Understanding the role of human intuition on reliance in human-AI decision-making with explanations. *Proceedings of the ACM on Human-Computer Interaction* **7**, 1–32 (2023).

[29] Blease, C. R., Locher, C., Gaab, J., Hägglund, M. & Mandl, K. D. Generative artificial intelligence in primary care: An online survey of UK general practitioners. *BMJ Health & Care Informatics* e101102 (2024).

[30] Mellers, B. A., Schwartz, A. & Cooke, A. D. Judgment and decision making. *Annual Review of Psychology* **49**, 447–477 (1998).

[31] Giray, L. Prompt engineering with ChatGPT: A guide for academic writers. *Annals of Biomedical Engineering* **51**, 2629–2633 (2023).

[32] Nori, H. *et al.* Can generalist foundation models outcompete special-purpose tuning? Case study in medicine. *arXiv:2311.16452* (2023).

[33] Ghassemi, M. Presentation matters for AI-generated clinical advice. *Nature Human Behaviour* **7**, 1833–1835 (2023).

[34] Reis, M., Reis, F. & Kunde, W. Influence of believed AI involvement on the perception of digital medical advice. *Nature Medicine* **30**, 3098–3100 (2024).

[35] Nagendran, M., Festor, P., Komorowski, M., Gordon, A. C. & Faisal, A. A. Quantifying the impact of AI recommendations with explanations on prescription decision making. *npj Digital Medicine* **6**, 206 (2023).

[36] Gaube, S. *et al.* Non-task expert physicians benefit from correct explainable AI advice when reviewing X-rays. *Scientific Reports* **13**, 1383 (2023).

[37] Jabbour, S. *et al.* Measuring the impact of AI in the diagnosis of hospitalized patients: A randomized clinical vignette survey study. *JAMA* **330**, 2275–2284 (2023).

[38] Tschandl, P. *et al.* Human-computer collaboration for skin cancer recognition. *Nature Medicine* **26**, 1229–1234 (2020).

[39] Busby, L. P., Courtier, J. L. & Glastonbury, C. M. Bias in radiology: The how and why of misses and misinterpretations. *RadioGraphics* **38**, 236–247 (2018).

[40] Lee, C. S., Nagy, P. G., Weaver, S. J. & Newman-Toker, D. E. Cognitive and system factors contributing to diagnostic errors in radiology. *American Journal of Roentgenology* **201**, 611–617 (2013).

[41] Shea, Y.-F., Lee, C. M. Y., Ip, W. C. T., Luk, D. W. A. & Wong, S. S. W. Use of GPT-4 to analyze medical records of patients with extensive investigations and delayed diagnosis. *JAMA Network Open* **6**, e2325000 (2023).

[42] Cabral, S. *et al.* Clinical reasoning of a generative artificial intelligence model compared with physicians. *JAMA Internal Medicine* **184**, 581–583 (2024).

[43] Seyyed-Kalantari, L., Zhang, H., McDermott, M. B., Chen, I. Y. & Ghassemi, M. Underdiagnosis bias of artificial intelligence algorithms applied to chest radiographs in under-served patient populations. *Nature Medicine* **27**, 2176–2182 (2021).

[44] Wang, J. & Redelmeier, D. A. Cognitive biases and artificial intelligence. *NEJM AI* **1**, AIcs2400639 (2024).

Acknowledgments

The authors thank all participants.

Author contributions

P.S. and D.H. conducted the experiment and performed the data analysis. J.Ru., S.S., and B.F.H. designed the task and assessed the responses of the participants. P.S., D.H., and S.F. wrote the first draft of the manuscript. All authors contributed to conceptualization, results interpretation, and manuscript editing, and approved the final manuscript.

Data availability

De-identified data to reproduce the results is publicly available at https://github.com/ PhilippSpitzer/LLM-Explanations-for-Radiology.

Code availability

Code to reproduce the results is available via our GitHub at https://github.com/PhilippSpitzer/ LLM-Explanations-for-Radiology. The GitHub also includes the survey of the online experiment as a Qualtrics `.qsf` file.

Competing interests

The authors declare no competing interests.

Funding

S.F. acknowledges funding via the Swiss National Science Foundation (SNSF), Grant 186932. The funding body had no role in the design, analysis, and interpretation of the study.

Supplements

Supplementary Figures

Supplementary Tables

 # Supplementary Figures

Patient Information:

A 52-year-old woman with end-stage kidney disease that was being managed with peritoneal dialysis presented with a 1-month history of bloody dialysate. She had had 3 episodes of bacterial peritonitis in the past 12 years. Physical examination and laboratory studies were unremarkable. Computed tomography of the abdomen was performed.

AI advice:

The patient is likely suffering from encapsulating peritoneal sclerosis (EPS). This diagnosis is based on several key findings:

1. Clinical Presentation: A 52-year-old woman with end-stage kidney disease managed with peritoneal dialysis presented with a 1-month history of bloody dialysate, a common symptom of EPS.
2. Medical History: The patient has had 3 episodes of bacterial peritonitis in the past 12 years, which is a known risk factor for EPS.
3. Physical Examination and Laboratory Studies: Both were unremarkable, indicating a chronic rather than an acute condition.
4. Initial Imaging: Computed tomography of the abdomen was performed.
5. Current Imaging: The CT scan shows characteristic findings of EPS, such as a thickened peritoneum and encapsulated bowel loops.
6. Epidemiology: Long-term peritoneal dialysis and recurrent bacterial peritonitis are significant risk factors for developing EPS.

In conclusion, the combination of clinical symptoms, medical history, and characteristic findings on CT strongly supports the diagnosis of encapsulating peritoneal sclerosis.

Your diagnosis:

Figure S1: **Example output: Case 5 "Encapsulating peritoneal sclerosis (EPS)".** Screenshot of the user interface during the experiment with LLM advice in the form of chain-of-thought explanations. Shown is the patient case 5, involving encapsulating peritoneal sclerosis (EPS).

Patient Information:

A 28-year-old woman with vertigo, confusion, and falls 2 weeks after a surgical abortion at 11 weeks of gestation presents to the emergency department. Examination revealed spontaneous upbeat nystagmus, gaze-evoked nystagmus, and gait ataxia.

AI advice:

The patient's symptoms of vertigo, confusion, falls, spontaneous upbeat nystagmus, gaze-evoked nystagmus, and gait ataxia after a surgical abortion, along with the MRI findings of hyperintensities in the bilateral thalami, suggest a diagnosis of Wernicke encephalopathy. This condition can occur due to thiamine deficiency, which may be precipitated by hyperemesis gravidarum or malnutrition associated with pregnancy.

Your diagnosis:

Figure S2: **Example output: Case 14 "Wernicke's encephalopathy".** Screenshot of the user interface during the experiment with LLM advice in the form of a standard output. Shown is patient case 14, involving Wernicke's encephalopathy.

Patient Information:

A 30-year-old man presented with a 15-month history of intermittent discomfort in the right upper quadrant of the abdomen. He lived in a rural area of Morocco and had occasional contact with dogs. The physical examination revealed hepatomegaly with a palpable hepatic mass. Laboratory tests showed a normal white-cell count and a normal absolute eosinophil count. Ultrasonography and computed tomography of the abdomen revealed a large cyst in the right lobe of the liver.

AI advice:

List of possible diseases

1. Hydatid disease (Echinococcosis)
2. Hepatic cystadenoma
3. Pyogenic liver abscess
4. Hepatic hemangioma
5. Hepatocellular carcinoma (HCC)

Detailed assessment of each possible diagnosis

1. Hydatid disease (Echinococcosis):

   Best Fit: The patient's history of living in a rural area with contact with dogs, along with the imaging showing a large cyst in the liver, strongly suggests hydatid disease. The presence of a cyst with internal septations is characteristic of Echinococcosis caused by Echinococcus granulosus.

2. Hepatic cystadenoma:

   Hepatic cystadenomas can present with large cysts in the liver and may cause discomfort due to their size. However, they are usually not associated with a history of exposure to dogs or rural living, making this less likely.

3. Pyogenic liver abscess:

   A pyogenic liver abscess can present with a hepatic mass and discomfort, but it is typically associated with systemic signs of infection, such as fever and elevated white-cell count, which are not present in this case.

4. Hepatic hemangioma:

   Hepatic hemangiomas are common liver lesions that can be large and cause discomfort. However, they usually appear as well-circumscribed, hypervascular lesions on imaging, which does not match the cystic appearance seen in the provided image.

5. Hepatocellular carcinoma (HCC):

   Least Likely: HCC can present with a hepatic mass and hepatomegaly, but it is more commonly associated with underlying liver disease, such as cirrhosis or chronic hepatitis. The imaging findings of a large cyst with septations are not typical of HCC.

Your diagnosis:

Figure S3: **Example output: Case 18 "Cystic echinoccoccis".** Screenshot of the user interface during the experiment with LLM advice in the form of a differential diagnosis. Shown is patient case 18, involving cystic echinoccoccis.

Figure S4: **Baseline diagnostic accuracy of LLM advice.** The diagnostic accuracy reports how often the LLM advice was stating the correct diagnosis (regardless of whether the explanation was correct or not). For the differential diagnosis, top-1 refers to whether the first answer was correct, while top-k refers to whether any of the five answers contained the correct diagnosis.

Figure S5: **Length of LLM-generated explanations across different treatment groups (in words).** Bar plots show the mean word count across all patient cases (mean as text annotations). Whiskers denote the standard deviation.

Figure S6: **Decision time.** Distribution of average time per case over all participants in minutes. The figure shows the distribution as violin plots (which illustrate the probability density of the data). The boxplots within each violin indicate the 25% and 75% quartiles, with the median represented by the thick center line and the mean by a diamond.

Figure S7: **Diagnostic accuracy across patient cases.** For certain cases, such as *collagenous enterocolitis* and *superior mesenteric artery syndrome*, all conditions led to high diagnostic accuracy rates above 80%. However, other patient cases proved more challenging, with notably lower performance across all conditions. For instance, for the cases of *ischemic colitis* and *cerebral cavernous malformations*, diagnostic accuracy rates dropped below 40% for several conditions. The chain-of-though group showed particularly strong performance in complex cases like *drug-induced thrombocytopenia* and *disseminated mycobacterium avium-intracellulare infection*, where other conditions often led to incorrect diagnoses. Physicians supported by the standard output performed well only in cases such as *Wernicke's encephalopathy* and *cystic echinococcosis*, while the differential diagnostic explanation showed strengths in diagnosing *pulmonary-artery dissection* and *infantile hepatic hemangioma*. This variation in performance across different medical conditions highlights the complexity of medical diagnosis and suggests that different prompting strategies might be more effective for different types of cases. Whiskers refer to standard deviations.

Figure S8: **Diagnostic accuracy is shown for a subset of patient cases where LLM advice was uniformly correct (or incorrect) across all conditions.** Here, we restricted our analysis to the subset of patient cases where the LLM advice was correct (or incorrect) across all conditions. The figure shows the distribution of diagnostic accuracy for all participants as violin plots (which illustrate the probability density of the data). Whiskers show standard deviations. For the differential diagnosis condition, the top-5 diagnostic accuracy is reported. Statistical significance was assessed using Welch's t-test to compare diagnostic accuracy between the treatment groups (in the main text, we report results from a regression analysis to obtain effect sizes).

Figure S9: **Robustness check for the diagnostic accuracy when counting partial responses as correct.** To account for diagnostic ambiguity (for instance, when a diagnosis appears accurate but omits a sub-diagnosis necessary for determining an appropriate treatment strategy), we perform a robustness check. The coding scheme in the main paper classified such partial responses as incorrect, while we now present an analysis in which we classified such responses as correct. The figure shows the distribution of diagnostic accuracy for all participants as violin plots (which illustrate the probability density of the data). The boxplots within each violin indicate the 25% and 75% quartiles, with the median represented by the thick center line. For the differential diagnosis condition, the top-5 diagnostic accuracy is reported (the top-1 diagnostic accuracy is lower and numbers to 65%). Statistical significance was assessed using Welch's t-test to compare diagnostic accuracy between the treatment groups.

Figure S10: **Adherence vs. overriding LLM advice based on the top-k answers.** Breakdown of adherence for **a**, correct LLM advice and **b**, incorrect LLM advice. The *adherence* was calculated as the proportion of patient cases in which participants' diagnoses matched the LLM-generated diagnoses. Correspondingly, *overriding* was defined as $1-$ adherence. Reported is the adherence for the top-5 answers in the differential diagnosis explanation.

Figure S11: **Effect on helpfulness.** The survey items were combined by averaging the Likert-scale ratings per participant and normalizing the result to a percentage scale. Helpfulness was not collected for the control group, leading to $n = 77$. Helpfulness was assessed by participants after completing the diagnostic tasks. An ANOVA reveals no statistically significant differences between the conditions (F-value = 1.324, P-value = 0.271), indicating that the type of explanation had no effect on helpfulness. Pairwise comparisons using t-tests also showed no significant differences between any of the conditions (all P-values > 0.05). Whiskers denote standard deviations.

Figure S12: **Effect on usefulness.** The survey items were combined by averaging the Likert-scale ratings per participant and normalizing the result to a percentage scale. The sample size of the post-study questionnaire varies, as the questionnaire was not fully displayed in two cases where participants selected specific specialties ("neuroradiologist" and "interventional radiology"). Usefulness was not collected for the control group, leading to $n = 76$. An ANOVA analysis reveals no statistically significant differences between the conditions (F-value = 0.690, P-value = 0.505), indicating that the explanation type had no effect on usefulness. Pairwise comparisons using t-tests also showed no significant differences between any of the conditions (all P-values > 0.05). Whiskers denote standard deviations.

Figure S13: **Effect on ease of use.** The survey items were combined by averaging the Likert-scale ratings per participant and normalizing the result to a percentage scale. The sample size of the post-study questionnaire varies, as the questionnaire was not fully displayed in two cases where participants selected specific specialties ("neuroradiologist" and "interventional radiology"). Ease of use was not collected for the control group, leading to $n = 76$. An ANOVA analysis reveals no statistically significant differences between the conditions (F-value = 1.313, P-value = 0.275), indicating that the explanation type had no effect on ease of use. Pairwise comparisons using t-tests also showed no significant differences between any of the conditions (all P-values > 0.05). Whiskers denote standard deviations.

Figure S14: **Effect on trust.** The survey items were combined by averaging the Likert-scale ratings per participant and normalizing the result to a percentage scale. The sample size for the trust construct is smaller due to a technical error in the study's setup, resulting in missing data. An ANOVA reveals no statistically significant differences between the conditions (F-value = 0.333, P-value = 0.718), indicating that the explanation type had no effect on trust. Pairwise comparisons using t-tests also showed no significant differences between any of the conditions (all P-values > 0.05). Whiskers denote standard deviations.

Figure S15: **Effect on task load.** The survey items were combined by averaging the Likert-scale ratings per participant and normalizing the result to a percentage scale. The sample size of the post-study questionnaire varies, as the questionnaire was not fully displayed in two cases where participants selected specific specialties ("neuroradiologist" and "interventional radiology"), leading to $n = 99$. An ANOVA reveals no statistically significant differences between the conditions (F-value = 0.505, P-value = 0.679), indicating that the type of explanation had no effect on task load. Pairwise comparisons using t-tests also showed no significant differences between any of the conditions (all P-values > 0.05). Whiskers denote standard deviations.

Figure S16: **Comparison of GPT-4 and Claude in diagnosing patients.** To provide a justification for our choice of GPT-4, we now compare the diagnostic accuracy of GPT-4 against another state-of-the-art LLM, namely, Claude. The reason for the choice is that Claude is designed to handle multi-modal input (i.e., patient descriptions with radiology images). This is unlike many other open-source LLMs (e.g., Llama-3, DeepSeek-R1), which, to this date, are limited to text-only input. We generated advice for the 20 patient cases from our study with images and text using Claude (version claude-3-opus-20240229, temperature = 1.0, maximum token generations = 4096). We then assessed the accuracy of both the diagnoses and the explanations provided by the model. Overall, both GPT-4 and Claude are comparable in terms of diagnostic accuracy. The largest difference is observed in the accuracy of observations. Considering that the effect of incorrect explanations on the participants in our study was small, it is reasonable to assume that we would obtain similar results from our study using a different LLM. To further support this claim, we calculated the inter-rater reliability of the models in diagnosing patient cases. The analysis of the inter-rater agreement between GPT-4 and Claude using Cohen's kappa revealed a fair agreement on both diagnoses ($\kappa = 0.243$) and explanations ($\kappa = 0.348$).

 # Supplementary Tables

Table S1: **Prompts for generating explanations.** Shown are the three different prompting approaches for generating medical diagnosis explanations: standard output, differential diagnosis, and chain-of-thought reasoning

Condition	Prompt
Standard output	Based on the provided text and image, I want you to make a diagnosis and explain your decision in 2-3 sentences.
Differential diagnosis	Differential diagnosis is a systematic method used by medical professionals to identify the underlying cause of a patient's symptoms or conditions and to eventually come up with a diagnosis based on a patient description and an image. It involves considering various possible diagnoses that could explain the symptoms and then systematically narrowing down the list based on clinical findings, medical history, laboratory tests, imaging studies, and other relevant information. The process typically involves the following steps: 1. Create a simple list with five possible diseases. 2. Loop through the list and explain why a specific disease does not fit, starting with the best fit and ending up with the disease that has the most unlikely fit. Reason for each disease within two short sentences taking into account the patient's symptoms. I want you to apply the process of differential diagnosis by taking into account not only the textual information but also the image.
Chain-of-thought	I want you to diagnose a patient by providing a diagnosis and explain how you come to that diagnosis step by step. For that, there is a patient description and an image with relevant information about the patient.

Table S2: **Items in the post-task survey.** Survey items measure participants' socio-demographics, expertise, and perceptions toward the LLM advice in the medical diagnosis tasks.

Variable	Items	Scale
Explanation helpfulness	• How helpful were the explanations? • How can we make the AI advice more helpful for diagnosing patients?	7-point Likert-scale and free-form text answer respectively
Expertise in Generative AI	• How do you rate your expertise in using Generative AI like ChatGPT?	7-point Likert-scale
Specialization	• What is your specialization in medicine? • Which area do you specialize in?	Multiple-choice-question with option for free-form text answer
Medical experience	• How many years of experience do you have as a medical professional?	Number
Experience in radiology	• How do you rate your experience in radiology?	7-point Likert-scale
IT skills	• How strong would you consider your IT skills?	7-point Likert-scale
Experience with medical AI systems	• How much experience do you have in working with medical AI systems?	7-point Likert-scale
Weekly visual inspection time	• How many hours per week do you spend on visual inspections?	7-point Likert-scale
Perceived self-efficacy	• Please estimate the percentage (0–100%) of diagnoses that you made correctly. • How successful were you in accomplishing what you were asked to do?	Free-form text answer with check for numeric inputs and 7-point Likert-scale respectively
Perceived AI efficacy	• Please estimate the percentage (0–100%) of diagnoses that the AI system made correctly. • How likely is the AI system to make a bad estimate?	Free-form text answer with check for numeric inputs and 7-point Likert-scale respectively
AI expectations	• How well did the AI system perform in comparison to your expectations?	7-point Likert-scale

Usefulness	• Using the AI system would enable others to classify architectural styles of buildings more quickly. • Using the AI system would improve the performance when classifying architectural styles of buildings. • Using the AI system would increase productivity for classifying the architectural styles of buildings. • Using the AI system would enhance my effectiveness in classifying the architectural styles of buildings. • Using the AI system would make it easier to classify the architectural styles of buildings. • I would find the AI system useful for classifying the architectural styles of buildings.	7-point Likert-scale
Ease of use	• Learning to operate the AI would be easy for me. • I would find it easy to get the AI to do what I want it to do. • My interaction with the AI would be clear and understandable. • I would find the AI to be flexible to interact with. • It would be easy for me to become skillful at using AI. • I would find the AI easy to use.	7-point Likert-scale
Trust	• The AI is deceptive. • I am suspicious of the AI's intent, action, or outputs. • The AI's actions will have a harmful outcome. • I am confident in the AI. • The AI is reliable. • I can trust the AI.	7-point Likert-scale

Task load	- How mentally demanding was the task? - How physically demanding was the task? - How hurried or rushed was the pace of the task? - How hard did you have to work to accomplish your level of performance? - How insecure, discouraged, irritated, stressed, and annoyed were you?	7-point Likert-scale
Open-ended questions	- Have you used any assistive technology/tools (internet research, books, colleagues, etc.) in filling out this survey? - In case you clicked "yes", which assistive technology/tools did you use?	Free-form text answer

Table S3: **Patient cases included in the study.** Description of the 20 diverse patient cases used in the study, which include, for example, pediatric and geriatric patients with various presenting symptoms and diagnostic images.

Case 1	A previously healthy 5-year-old boy was brought to the surgery clinic with a 2-day history of intermittent abdominal pain. On palpation of the abdomen there was pain in the periumbilical region, but no rebound or guarding. An ultrasound was normal, and a computed tomography of the abdomen was performed (Panels A,B).
Case 2	A 55-year-old man presented with 10 years of progressive handwriting impairment and rapid, slurred speech. In his thirties, he had worked as a welder without access to personal protective equipment. Neurologic examination was notable for reduced facial expression, blepharospasm, and cluttered, dysarthric speech. Postural reflexes were mildly impaired. MRI imaging of the head showed a nonenhancing, T1-weighted, hyperintense signal in the basal ganglia on both sides. Ceruloplasmin and iron levels were normal.
Case 3	26-year-old man from Somalia presented with a 5-month history of dry cough, night sweats, and unintentional weight loss of 18 kg. During this period, epigastric pain and postprandial vomiting had also developed. His BMI was 11. On examination, he was cachectic with abdominal distention and diffuse tenderness to palpation. On the basis of chest imaging and sputum studies, a diagnosis of pulmonary tuberculosis was made, and intravenous antituberculous treatment was initiated. However, he continued to have postprandial vomiting. Contrast-enhanced CT of the abdomen was obtained.
Case 4	A 35-year-old man with IgA nephropathy presented with confusion, blurry vision, and seizures. Two weeks before presentation, he had started receiving cyclosporine. Physical examination was notable for a blood pressure of 160/80 mm Hg, drowsiness, and decreased visual acuity. A fundoscopic examinations was normal. T2-weighted magnetic resonance imaging (MRI) with fluid-attenuated inversion recovery sequencing of the head was performed.
Case 5	A 52-year-old woman with end-stage kidney disease that was being managed with peritoneal dialysis presented with a 1-month history of bloody dialysate. She had had 3 episodes of bacterial peritonitis in the past 12 years. Physical examination and laboratory studies were unremarkable. Computed tomography of the abdomen was performed.
Case 6	A 32-year-old man presented with a 6-week history of tingling in his arms and legs and a 2-week history of inability to walk. A positive Romberg test, sensory ataxia, impaired proprioception and vibratory sensation, and preserved nociception were noted. Magnetic resonance imaging of the whole spine showed hyperintensity in the posterior spinal cord from C1 to T12 and hyperintense lesions in the dorsal column on T2-weighted images. A vitamin B12 level was 107 pg per ml (reference value, >231) without macrocytic anemia.
Case 7	A 35-year-old woman with idiopathic pulmonary arterial hypertension and a pulmonary aneurysm presented with chest pain. Computed tomography (CT) of the chest is shown.
Case 8	A 38-year-old man presented to the otolaryngology clinic with chronic difficulty breathing through his right nostril. Physical examination showed nasal septal deviation, calcified septal spurs, and a 2-cm perforation in the posterior septum. On rhinoscopy, a hard, nontender, white mass was observed in the floor of the right nostril. CT of the paranasal sinuses showed a well-defined, radiodense mass.
Case 9	A 16-day-old girl was brought to the emergency department with lethargy. Physical exam showed tachypnea and marked hepatomegaly, as well as small hemangiomas on the skin. TSH was elevated. MRI showed numerous hepatic lesions and cardiomegaly.
Case 10	A 71-year-old man was hospitalized with altered mental status progressing over the preceding 3 weeks. The patient had a recent diagnosis of adenocarcinoma of the colon with known metastatic lesions in the lung and bones. A gadolinium-enhanced magnetic resonance image of the brain was performed and is shown.

Case 11	A 29-year-old man with perinatally acquired human immunodeficiency virus (HIV) infection and intermittent adherence to antiretroviral therapy presented to the hospital with abdominal pain and drenching night sweats. On presentation, his CD4 count was 18 cells per cubic millimeter (reference range, 500 to 1500), and the HIV viral load was undetectable. Physical exam showed severe abdominal distention, splenomegaly, and diffuse abdominal tenderness to palpation. Computed tomography of the abdomen confirmed massive splenomegaly with multifocal infarction of the splenic parenchyma.
Case 12	A 42-year-old man presented to the clinic with a 3-month history of worsening cough, shortness of breath, and fever. Physical examination showed inflamed nasal mucosa and nasal crusting. Wheezes and rales were heard on auscultation. A computed tomographic scan of the face showed extensive destruction of the structural bones of the midface, resulting in a large nasal cavity.
Case 13	A 63-year-old man presented to the emergency department with a 3-day history of abdominal pain that had started in the periumbilical area and subsequently shifted to the left lower quadrant. Initial laboratory tests showed a white-cell count of 12,000 per cubic millimeter (reference range, 4000 to 10,000) and a lactate level of 1.8 mmol per liter (normal value, <1.9). Contrast-enhanced computed tomography of the abdomen revealed edema of the sigmoid colon with thumbprinting.
Case 14	A 28-year-old woman with vertigo, confusion, and falls 2 weeks after a surgical abortion at 11 weeks of gestation presents to the emergency department. Examination revealed spontaneous upbeat nystagmus, gaze-evoked nystagmus, and gait ataxia.
Case 15	A 59-year-old previously healthy man presented with progressively worsening headaches and bluish nodular skin lesions. Fast-field echo MRI image of the brain showed this finding.
Case 16	A 44-year-old woman presented to the emergency department with acute chest pain after several months of progressive dyspnea. Her oxygen saturation was 92%, and she had diminished breath sounds on the right side. Chest CT revealed a large right-sided pneumothorax and diffuse, intraparenchymal pulmonary cysts.
Case 17	A 54-year-old man presented with a 3-week history of cognitive deterioration. Neurologic examination revealed disorientation, horizontal gaze-evoked nystagmus, hyperreflexia, startle myoclonus, and ataxia. Brain MRI with diffusion-weighted imaging revealed hyperintensity of the cortical gyri and caudate heads.
Case 18	A 30-year-old man presented with a 15-month history of intermittent discomfort in the right upper quadrant of the abdomen. He lived in a rural area of Morocco and had occasional contact with dogs. The physical examination revealed hepatomegaly with a palpable hepatic mass. Laboratory tests showed a normal white-cell count and a normal absolute eosinophil count. Ultrasonography and computed tomography of the abdomen revealed a large cyst in the right lobe of the liver.
Case 19	A 59-year-old woman presented to the emergency department with a 4-day history of inflammation and pain in the right eye. She had been blind in the eye for several years before presentation. Magnetic resonance imaging revealed a right orbital mass. Abdominal and thoracic imaging showed numerous hepatic masses, abdominal and thoracic lymphadenopathy, and vertebral sclerotic osseous disease. The right eye was enucleated for palliative relief and to obtain tissue for diagnosis.
Case 20	An 18-year-old man presented to the emergency department with generalized tonic–clonic seizures. On physical examination, the patient was confused. He had swelling over the right eye and tenderness in the right testis. Magnetic resonance imaging of the head showed numerous well-defined cystic lesions throughout the cerebral cortex.

| | | Condition | | | | |
Variable	Statistic	Control	Standard	Differential diagnosis	Chain-of-thought	Overall
Medical experience (in years)	Mean	15.087	13.104	10.800	16.500	13.621
	SD	6.557	6.617	7.364	10.327	7.981
Experience with medical AI systems	Mean	5.000	5.500	5.133	4.818	5.121
	SD	1.859	1.956	1.814	2.322	1.965
Experience in radiology	Mean	5.739	6.000	5.967	6.500	6.040
	SD	1.010	0.933	0.964	0.598	0.925
IT skills	Top	Very Good	Good	Good	Good	Good
	Freq	10	8	9	9	33
Weekly visual inspection time	Top	45-50h	>55h	35-40h	45-50h	45-50h
	Freq	8	7	7	5	25

Table S4: **Demographics of study sample by condition.** The table presents demographic statistics for study participants across the different experimental conditions. Variables include medical experience (in years), experience with medical AI systems (reported on a 7-point Likert scale), experience in radiology (reported on a 7-point Likert scale), and weekly time spent on visual inspection tasks (reported as discrete time ranges on a 7-item scale). In the table, we report the mean and standard deviation (SD) for continuous variables. For categorical variables, we report the most frequently occurring category (top) and the corresponding frequency (freq). The sample size of the post-study questionnaire varied, as the questionnaire was not fully displayed in two cases where participants selected specific specialties ("neuroradiologist" and "interventional radiology"), leading to $n = 99$ participants with demographic data.

	Coef.	s.e.	_P_-value	95% CI
Intercept	1.386	0.559	0.013	[0.291; 2.482]
Condition: Standard	−0.288	0.761	0.705	[−1.779; 1.204]
Condition: Differential diagnosis	0.000	0.791	1.000	[−1.549; 1.549]
AIC				68.526
Obs. (N)				60

Table S5: **Diagnostic accuracy of the LLM (without human involvement) across different prompting strategies.** We estimated a logistic regression at the diagnostic level (i.e., 20 diagnoses × 3 conditions). The chain of thought condition is used as the reference category. The odds ratios computed based on the coefficients are 0.750 for the standard condition and 1.000 for the differential diagnosis condition. Hence, considering the intercept, the chain-of-thought approach has a better diagnostic accuracy than the standard prompt and a similar diagnostic accuracy as the differential diagnosis. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	P-value	95% CI
Intercept	0.617	0.024	< 0.001	[0.568; 0.665]
Condition: Standard	0.050	0.034	0.150	[−0.018; 0.118]
Condition: Differential diagnosis	0.025	0.033	0.446	[−0.040; 0.090]
Condition: Chain-of-thought	0.122	0.035	0.001	[0.053; 0.192]
AIC				−138.835
Obs. (N)				101

Table S6: **Effect of explanation types on diagnostic accuracy.** OLS regression explaining the diagnostic accuracy at the physician level by the different conditions. The intercept represents the diagnostic accuracy of humans without LLM support (control condition). Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	P-value	95% CI
Intercept	0.739	0.025	< 0.001	[0.690; 0.789]
Condition: Control	−0.122	0.035	0.001	[−0.192; −0.053]
Condition: Standard	−0.072	0.035	0.040	[−0.142; −0.003]
Condition: Differential diagnosis	−0.097	0.033	0.004	[−0.163; −0.032]
AIC				−138.835
Obs. (N)				101

Table S7: **Benefit of chain-of-thought explanations on diagnostic accuracy.** OLS regression of diagnostic accuracy at the physician level on conditions with chain-of-thought as baseline (=intercept). Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	P-value	95% CI
Intercept	0.604	0.123	< 0.001	[0.359; 0.850]
Condition: Standard	0.070	0.040	0.081	[−0.009; 0.149]
Condition: Differential diagnosis	0.039	0.039	0.325	[−0.039; 0.117]
Condition: Chain-of-thought	0.151	0.041	< 0.001	[0.070; 0.232]
Medical experience	0.001	0.002	0.707	[−0.003; 0.004]
Experience in radiology	−0.001	0.001	0.301	[−0.003; 0.001]
Experience with medical AI systems	−0.000	0.000	0.519	[−0.001; 0.001]
Weekly visual inspection time: 30−35h	0.077	0.108	0.478	[−0.138; 0.292]
Weekly visual inspection time: 35−40h	0.043	0.101	0.667	[−0.157; 0.244]
Weekly visual inspection time: 40−45h	0.067	0.100	0.508	[−0.133; 0.266]
Weekly visual inspection time: 45−50h	0.076	0.097	0.441	[−0.118; 0.269]
Weekly visual inspection time: 50−55h	0.039	0.100	0.698	[−0.159; 0.237]
Weekly visual inspection time: >55H	0.047	0.102	0.648	[−0.156; 0.249]
IT skills: Good	0.004	0.046	0.935	[−0.088; 0.095]
IT skills: Very Good	0.068	0.048	0.156	[−0.027; 0.163]
IT skills: Advanced	0.043	0.051	0.403	[−0.058; 0.143]
IT skills: Expert	0.023	0.072	0.748	[−0.120; 0.167]
AIC				−116.208
Obs. (N)				99

Table S8: **Extended OLS regression of diagnostic accuracy with physician-level controls.** Reference category is the control group. In the post-survey questionnaire, the sample size of participants across conditions totals $n = 99$, as the questionnaire was not displayed in two cases where participants selected specific specialties ('neuroradiologist' and 'interventional radiology'). Hence, the two cases were dropped for the analysis. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	P-value	95% CI
Intercept	0.544	0.041	< 0.001	[0.463; 0.625]
Condition: Standard	0.021	0.028	0.467	[−0.035; 0.076]
Condition: Differential diagnosis	−0.048	0.017	0.005	[−0.081; −0.015]
Condition: Chain-of-thought	0.046	0.018	0.012	[0.010; 0.081]
Total time	0.000	0.000	0.125	[−0.000; 0.001]
Advice length	0.000	0.000	0.030	[0.000; 0.001]
Answer length	0.003	0.002	0.050	[−0.000; 0.006]
AIC				−141.197
Obs. (N)				101

Table S9: **Extended OLS regression of diagnostic accuracy with advice-level controls.** Here, we control for total time (measured in minutes), advice length (measured in the number of words), and answer length (of the reported diagnosis by the physicians in the free-text form, measured in the number of characters). The reference category is the control group. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	P-value	95% CI
Intercept	0.575	0.037	< 0.001	[0.500; 0.650]
Condition: Standard	0.025	0.051	0.628	[−0.079; 0.129]
Condition: Differential diagnosis	0.090	0.052	0.094	[−0.016; 0.196]
Condition: Chain-of-thought	0.167	0.050	0.002	[0.065; 0.268]
AIC				−58.545
Obs. (N)				43

Table S10: **Subgroup analysis for participants with basic IT skills.** OLS regression explaining the diagnostic accuracy at the physician level by the conditions for the subgroup of participants that have basic IT skills (i.e., lower or equal to "Good"). Reference category is the control group. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	P-value	95% CI
Intercept	0.646	0.033	< 0.001	[0.581; 0.711]
Condition: Standard	0.077	0.046	0.101	[−0.015; 0.169]
Condition: Differential diagnosis	−0.016	0.042	0.701	[−0.100; 0.068]
Condition: Chain-of-thought	0.094	0.049	0.063	[−0.005; 0.193]
AIC				−77.177
Obs. (N)				56

Table S11: **Subgroup analysis for participants with advanced IT skills.** OLS regression explaining the diagnostic accuracy at the physician level by the conditions for the subgroup of participants that have advanced IT skills (i.e., better than "Good"). Reference category is the control group. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	*P*-value	95% CI
Intercept	0.619	0.030	< 0.001	[0.558; 0.680]
Condition: Standard	0.072	0.040	0.077	[−0.008; 0.153]
Condition: Differential diagnosis	0.037	0.036	0.318	[−0.037; 0.110]
Condition: Chain-of-thought	0.109	0.042	0.012	[0.025; 0.193]
AIC				−91.631
Obs. (N)				46

Table S12: **Subgroup analysis for participants with short tenure.** OLS regression explaining the diagnostic accuracy at the physician level by the conditions for the subgroup of radiologists with short tenure (less than 12.0 years). Reference category is the control group. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	P-value	95% CI
Intercept	0.613	0.038	< 0.001	[0.537; 0.689]
Condition: Standard	0.033	0.055	0.556	[−0.078; 0.144]
Condition: Differential diagnosis	0.008	0.057	0.895	[−0.106; 0.121]
Condition: Chain-of-thought	0.137	0.055	0.017	[0.025; 0.248]
AIC				−49.587
Obs. (N)				53

Table S13: **Subgroup analysis for participants with long tenure.** OLS regression explaining the diagnostic accuracy at the physician level by the conditions for the subgroup of radiologists with long tenure (more than or equal to 12.0 years). Reference category is the control group. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	P-value	95% CI
Intercept	0.867	0.022	< 0.001	[0.823; 0.911]
Condition: Standard	−0.033	0.031	0.291	[−0.096; 0.029]
Condition: Differential diagnosis	−0.057	0.030	0.060	[−0.116; 0.002]
Condition: Chain-of-thought	0.051	0.032	0.113	[−0.012; 0.114]
AIC				−157.675
Obs. (N)				101

Table S14: **Subgroup analysis for basic patient cases.** OLS regression explaining the diagnostic accuracy at the physician level by the conditions for the subset of basic cases. We split patient cases into basic and complex subsets based on the mean diagnostic accuracy of the control group for each case, where the lower half corresponds to the basic cases. Reference category is the control group. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	*P*-value	95% CI
Intercept	0.500	0.031	< 0.001	[0.438; 0.562]
Condition: Standard	0.154	0.045	0.001	[0.066; 0.243]
Condition: Differential diagnosis	0.080	0.042	0.061	[−0.004; 0.164]
Condition: Chain-of-thought	0.200	0.045	< 0.001	[0.111; 0.289]
AIC				−87.045
Obs. (N)				101

Table S15: **Subgroup analysis for complex patient cases.** OLS regression explaining the diagnostic accuracy at the physician level by the conditions for the subset of complex cases. We split patient cases into basic and complex subsets based on the mean diagnostic accuracy of the control group for each case, where the upper half corresponds to the complex cases. Reference category is the control group. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	P-value	95% CI
Intercept	0.625	0.021	< 0.001	[0.584; 0.666]
Condition: Standard	0.052	0.029	0.079	[−0.006; 0.110]
Condition: Differential diagnosis	0.015	0.028	0.591	[−0.040; 0.070]
Condition: Chain-of-thought	0.114	0.030	< 0.001	[0.055; 0.173]
AIC				−171.332
Obs. (N)				101

Table S16: **Robustness checks controlling for outliers.** OLS regression explaining the winsorized diagnostic accuracy at the physician level by the different conditions. To improve the statistical reliability and reduce the potential bias from outliers or extreme values, we applied winsorization to the diagnostic accuracy at the participant level. We do this by replacing values outside the 5th and 95th percentiles with values at these percentiles, thus limiting the influence of extreme values. Reference category is the control group. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	*P*-value	95% CI
Intercept	1.036	0.823	0.208	[−0.577; 2.648]
Condition: Standard	0.548	0.248	0.027	[0.062; 1.034]
Condition: Differential diagnosis	0.095	0.242	0.695	[−0.379; 0.569]
Condition: Chain-of-thought	1.141	0.259	< 0.001	[0.633; 1.650]
Medical experience	0.001	0.011	0.922	[−0.021; 0.023]
Experience in radiology	−0.008	0.007	0.256	[−0.021; 0.006]
Experience with medical AI systems	−0.003	0.003	0.316	[−0.009; 0.003]
Weekly visual inspection time: 30−35h	0.738	0.663	0.265	[−0.561; 2.038]
Weekly visual inspection time: 35−40h	0.361	0.610	0.554	[−0.834; 1.556]
Weekly visual inspection time: 40−45h	0.551	0.607	0.364	[−0.639; 1.741]
Weekly visual inspection time: 45−50h	0.535	0.590	0.364	[−0.621; 1.691]
Weekly visual inspection time: 50−55h	0.299	0.604	0.621	[−0.884; 1.482]
Weekly visual inspection time: >55h	0.314	0.617	0.611	[−0.896; 1.524]
IT skills: Good	0.070	0.285	0.805	[−0.488; 0.628]
IT skills: Very Good	0.555	0.299	0.063	[−0.030; 1.140]
IT skills: Advanced	0.516	0.316	0.103	[−0.104; 1.136]
IT skills: Expert	0.330	0.452	0.465	[−0.556; 1.215]
AIC				1780.944
Obs. (N)				1980
Variance explained by participant IDs				0.235
Variance explained by case IDs				2.275

Table S17: **Mixed-effects model.** We estimated a mixed-effects model to control for heterogeneity across physicians and cases. Reference category is the control group. As we control for variables assessed in the questionnaire, which have not been displayed in two cases, the sample size is reduced to $N = 1980 = 2020 − 20 \times 2$ assessments. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	P-value	95% CI
Intercept	0.475	0.105	< 0.001	[0.269; 0.681]
Condition: Standard	0.218	0.151	0.149	[−0.078; 0.514]
Condition: Differential diagnosis	0.107	0.142	0.450	[−0.171; 0.385]
Condition: Chain-of-thought	0.566	0.159	< 0.001	[0.255; 0.877]
AIC				95.364
Obs. (N)				101

Table S18: **Quasi-binomial regression.** We estimated quasi-binomial regression, which may limit the interpretability of the estimated effect sizes but which can better accommodate for the fact that the diagnostic accuracy is in the range between 0% and 100%. Here, we fitted a generalized linear model with a logistic link function, using a binomial error distribution to model the diagnostic accuracy of responses based on the conditions from our experiment. We further adjusted for overdispersion by scaling the standard errors using a dispersion factor estimated from the Pearson χ^2-statistic. Reference category is the control group. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	*P*-value	95% CI
Intercept	79.369	30.649	0.012	[18.327; 140.411]
Condition: Standard	13.117	43.344	0.763	[−73.209; 99.443]
Condition: Differential diagnosis	68.332	43.344	0.119	[−17.994; 154.659]
Condition: Chain-of-thought	57.825	43.344	0.186	[−28.501; 144.152]
AIC				1018.199
Obs. (N)				80

Table S19: **Effect on decision time.** OLS regression explaining the average time to answer each case across the different conditions, leading to $N = 80$ observations (= 20 cases × 4 conditions). Reference category is the control group. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	*P*-value	95 % CI
Intercept	92.485	34.900	0.010	[22.599; 162.372]
Condition: Differential diagnosis	55.216	49.356	0.268	[−43.619; 154.050]
Condition: Chain-of-thought	44.709	49.356	0.369	[−54.126; 143.543]
AIC				779.238
Obs. (N)				60

Table S20: **Effect of decision time (compared to the standard group).** OLS regression explaining the average time to answer each case across the different conditions. The standard output group is set as the reference group, leading to $N = 60$ observations (= 20 cases $\times 3$ treatment groups). Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	P-value	95% CI
Intercept	0.578	0.022	< 0.001	[0.536; 0.621]
Condition: Standard	-0.095	0.030	0.002	[-0.154; -0.036]
Condition: Differential diagnosis	0.217	0.029	< 0.001	[0.160; 0.273]
AIC				2003.272
Obs. (N)				1540

Table S21: **Effect on adherence.** OLS regression explaining the adherence at the assessment level for the different conditions. The chain-of-thought group is set as the reference group; the control group is excluded as adherence is not applicable therein. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	P-value	95% CI
Intercept	0.769	0.098	< 0.001	[0.577; 0.961]
Condition: Standard	0.296	0.143	0.039	[0.015; 0.577]
Condition: Differential diagnosis	0.054	0.132	0.680	[−0.205; 0.314]
Condition: Chain-of-thought	0.672	0.154	< 0.001	[0.371; 0.974]
AIC				2340.802
Obs. (N)				2020

Table S22: **Logistic regression at the assessment level.** Logistic regression at the assessment level (=1 if the correct diagnosis was given by a physician for a specific case, 0 otherwise) explaining the diagnostic accuracy by the different conditions. Reference category is the control group. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	P-value	95% CI
Intercept	0.940	0.026	< 0.001	[0.888; 0.993]
Condition: Control	−0.164	0.037	< 0.001	[−0.238; −0.091]
Condition: Standard	−0.044	0.037	0.233	[−0.118; 0.029]
Condition: Differential diagnosis	0.018	0.035	0.607	[−0.052; 0.088]
AIC				−126.733
Obs. (N)				101

Table S23: **Subgroup analysis for patient cases with comparable LLM performance across conditions.** OLS regression explaining the diagnostic accuracy at the physician level across the different conditions but only for a subset of cases. The subset consists of cases for which the LLM advice was consistently correct or incorrect across all conditions for both the diagnosis and explanation. The chain-of-thought group is set as the reference group. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	P-value	95% CI
Intercept	−0.619	0.171	< 0.001	[−0.954; −0.285]
Condition: Standard	−0.504	0.209	0.016	[−0.914; −0.094]
Condition: Differential diagnosis	−1.014	0.188	< 0.001	[−1.383; −0.646]
Correct explanation ($True = 1$): 1	0.462	0.162	0.004	[0.144; 0.779]
Correct diagnosis ($True = 1$): 1	2.900	0.168	< 0.001	[2.571; 3.229]
AIC				1263.086
Obs. (N)				1540

Table S24: **Logistic regression controlling for correct explanations and correct diagnosis in the LLM advice.** Logistic regression at the assessment level (=1 if the correct diagnosis was given by a physician for a specific case, 0 otherwise) explaining the diagnostic accuracy by the different conditions. Here, we separately control for (a) whether the suggested diagnosis of the LLM is correct or (b) whether the explanation generated by the LLM is correct. The diagnosis in the differential diagnosis group is considered correct if the correct diagnosis is one of the five presented options. The chain-of-thought group served as the reference group. We omit the control group because the control did not receive LLM advice, making it impossible to control for the correctness of that advice. Abbreviations: s.e., standard error; CI, confidence interval, AIC, Akaike information criterion.

	Coef.	s.e.	*P*-value	95 % CI
Condition: Standard	0.470	0.137	0.001	[0.201; 0.739]
Condition: Differential diagnosis	0.216	0.125	0.085	[−0.030; 0.461]
Condition: Chain-of-thought	0.862	0.148	< 0.001	[0.572; 1.152]
Image: Count	0.045	0.075	0.546	[−0.102; 0.192]
Source: MRI	0.729	0.141	< 0.001	[0.451; 1.006]
Body region: Head	0.192	0.136	0.159	[−0.075; 0.460]
Body region: Spine	1.806	0.488	< 0.001	[0.849; 2.762]
Body region: Thorax	0.719	0.176	< 0.001	[0.375; 1.064]
AIC				2267.722
Obs. (*N*)				2020

Table S25: **Regression controlling for the characteristics of radiological images.** Logistic regression at the assessment level (=1 if the correct diagnosis was given by a physician for a specific case, 0 otherwise) explaining the diagnostic accuracy by the different conditions and image characteristics. Here, we include (a) the number of images shown for a patient case, (b) the source of the image(s) (i.e., MRI or CT [CT serves as a baseline in the regression]), and (c) the body region shown in the image(s) (abdomen serves as a baseline for the regression here). By exponentiating the coefficients, the odds ratios of 1.60 for the standard condition, 1.24 for the differential condition, and 2.37 for the chain-of-thought condition are obtained. Abbreviations: s.e., standard error; CI, confidence interval.