The diagnostic and prognostic utility of mitral annular plane systolic excursion (MAPSE) – a systematic review

Debbie Falconer*[1,2], Fredrika Fröjdh*[3], Daniel Brieger[4], Gaby Captur[1,2], Rebecca Kozor[4,5], Martin Ugander[3,4,5]

*Equal contribution

[1]University College London, Gower Street, London, United Kingdom

[2]Royal Free Hospital, Pond Street, London, United Kingdom

[3]Department of Clinical Physiology, Karolinska University Hospital, and Karolinska Institutet, Stockholm, Sweden

[4]Kolling Institute, Royal North Shore Hospital, Sydney, Australia

[5]University of Sydney, Sydney, Australia

Corresponding author:

Martin Ugander, MD, PhD

Professor of Cardiac Imaging, University of Sydney

Kolling Building, Level 12, Royal North Shore Hospital, St Leonards, Sydney, NSW 2065, Australia

Email: martin.ugander@gmail.com

Funding: This work is supported in part by the British Cardiology Society/ Heart Research UK clinical fellowship.

Abstract

Movement of the mitral annulus towards the left ventricular (LV) apex during systole, termed atrioventricular plane displacement (AVPD) or mitral annular plane systolic excursion (MAPSE), was first observed by Leonardo da Vinci in the 15[th] century. Longitudinal LV function accounts for 60% of the total stroke volume, so MAPSE reflects the majority contribution to systolic function. However, no major imaging guidelines advise routine measurement. We present a systematic review of studies appraising the diagnostic and prognostic performance of MAPSE from PubMed, Medline and Embase in accordance with the PRISMA statement. MAPSE values show good agreement between transthoracic echocardiography and cardiac magnetic resonance imaging (CMR), and can also be measured by transesophageal echocardiography and gated cardiac computed tomography. Manual measurement is fast, simple, and less reliant on good echocardiographic image quality than left ventricular ejection fraction (LVEF) or global longitudinal strain (GLS). Also, measurement can be easily automated, reducing reporting time. MAPSE correlates with both LVEF (r=0.60 [95% confidence interval 0.49–0.70]) and global longitudinal strain (r=0.45 [0.34–0.57]) and can act as an alternative marker of LV systolic dysfunction in patients with poor echocardiographic windows. MAPSE falls while LVEF remains preserved across a range of pathologies, allowing physicians to detect systolic impairment earlier than when using LVEF. MAPSE is also a powerful prognostic tool, outperforming LVEF and GLS in predicting adverse events in several studies. These results highlight a clinically useful and important role for MAPSE in routine cardiac imaging and care.

Highlights

- MAPSE reflects the systolic longitudinal movement of the atrioventricular valve plane, which accounts for 60% of the left ventricular stroke volume.

- MAPSE is easily and quickly attainable, shows good agreement between imaging modalities, and can be easily automated.

- MAPSE can be used to detect systolic dysfunction before left ventricular ejection fraction falls in many disease states, allowing clinicians to identify patients most at risk from progression of heart disease.

- MAPSE has an association with prognosis that is stronger than that of left ventricular ejection fraction and global longitudinal strain in several conditions.

Keywords

Atrioventricular plane displacement (AVPD), cardiovascular magnetic resonance (CMR), transesophageal echocardiography, mitral annular plane systolic excursion (MAPSE), left ventricular systolic dysfunction

Abbreviations and acronyms

ACS	acute coronary syndrome
AVPD	atrioventricular plane displacement
AS	aortic stenosis
AF	atrial fibrillation
AI	artificial intelligence
ARCD	anthracycline related cardiac dysfunction
BSE	British Society of Echocardiography
CABG	coronary artery bypass graft
CI	confidence interval
CMR	cardiovascular magnetic resonance
COVID-19	coronavirus disease 2019
CT	computed tomography
DD	diastolic dysfunction
DMD	Duchenne Muscular Dystrophy
EACVI	European Association of Cardiovascular Imaging
ED	emergency department
GLS	global longitudinal strain
HCM	hypertrophic cardiomyopathy
HFpEF	heart failure with preserved ejection fraction
HFrEF	heart failure with reduced ejection fraction
HTN	hypertension
HR	hazard ratio
IHD	ischemic heart disease
LAS	left atrial strain
LFLG	low-flow low-gradient
LGE	late gadolinium enhancement
LOA	limits of agreement
LV	left ventricle
LVEF	left ventricular ejection fraction
LVH	left ventricular hypertrophy
MACE	major adverse cardiovascular event
MAPSE	mitral annular plane systolic excursion
NYHA	New York Heart Association
OR	odds ratio
PAH	pulmonary artery hypertension
PE	pulmonary embolism
POCUS	point-of-care ultrasound
TAVI	Transaortic catheter valve implantation
TDI	tissue Doppler imaging
TMAD	tissue mitral annular displacement
TTE	transthoracic echocardiography
TEE	transesophageal echocardiography
RAVI	right atrial volume index
RV	right ventricle
SCMR	Society for Cardiovascular Magnetic Resonance
STE	speckle tracking echocardiography.

Introduction

Mitral annular plane systolic excursion (MAPSE), also termed atrioventricular plane displacement (AVPD), refers to the movement of the mitral annulus towards the apex during systole arising from ventricular contraction[1]. In the majority of patients, the cardiac apex is fixed to the chest wall meaning long axis contraction of the heart can be measured through the change in position of the mitral annulus. MAPSE, therefore, is a marker of longitudinal left ventricular (LV) function, which contributes 60% of the total stroke volume[2]. This proportion is constant across healthy hearts, elite athletes, and patients with dilated[3] and ischaemic cardiomyopathy, even when the absolute longitudinal function is decreased[4].

Despite reflecting an important component of cardiac function, the majority of contemporary imaging protocols do not advise routine measurement of MASPE, including the most recent guidelines from the British Society for Echocardiography[5] (BSE), American Society of Echocardiography (ASE)[6], European Association of Cardiovascular Imaging (EACVI)[7], and the Society for Cardiovascular Magnetic Resonance (SCMR)[8]. The BSE does advise a single MAPSE measurement as part of a comprehensive dataset[5], but this is not routinely performed in most institutions, and EACVI suggests MAPSE can be measured if other markers of longitudinal function are not available[7].

Left ventricular ejection fraction (LVEF), the mainstay for assessing systolic function, is based on the change in cavity size between end-diastole and end-systole resulting from both longitudinal and radial contraction of the LV. Some studies have suggest that since subendocardial longitudinal fibres are more sensitive to ischemia and may be more likely to be impaired in early stages of heart disease, this could potentially explain why longitudinal

function reduces before other indices [9-11]. LVEF can also be maintained until late stages of a disease by a compensatory increase in radial function [12,13], meaning early systolic impairment is missed when assessed by LVEF alone.

Global longitudinal strain (GLS) is a myocardial deformation analysis reflecting longitudinal contraction, which detects changes in systolic function before LVEF is impaired[14]. Strain can be expressed as a function of the entire LV (also termed long axis strain) or on a regional basis, the latter of which has been widely adopted in cardio-oncology. Both LVEF and GLS require excellent endocardial border definition, which is not always possible using echocardiography. GLS can also be time-consuming for practitioners, and requires additional training. Poor acoustic windows at echocardiography can be overcome with the use of CMR, but there remain patients in whom this is precluded due to ferromagnetic implants or severe claustrophobia.

Conversely, MAPSE can be measured almost universally, even with limited echocardiographic views, and using generic linear calliper measurement tools. Furthermore, measurements can be performed automatically using dedicated artificial intelligence (AI)-enabled software, without adding time to the reporting process[15]. As well as detecting early systolic impairment in longitudinal function, MAPSE has prognostic value across several cardiovascular diseases, often outperforming LVEF and GLS[15-17]. Therefore, excluding MAPSE from routine imaging may result in the loss of an easily attainable but highly informative biomarker.

This study aimed to systematically review evidence outlining the diagnostic and prognostic utility of MAPSE in adult cardiology patients, where possible comparing its performance to LVEF and GLS. The review addresses the following clinical questions:

1. When LVEF and GLS are not available, can MAPSE be used to diagnose systolic dysfunction (i.e. how closely does MAPSE correlate to these measurements)?

2. For what pathologies can MAPSE be used to detect early systolic dysfunction, in the presence of a normal LVEF?

3. In cardiovascular disease, how strongly is MAPSE associated with prognosis, and how does this compare with LVEF and GLS?

Of note, MAPSE has also been referred to as 'movement of the mitral ring' or 'atrioventricular plane displacement'. The term MAPSE came into common use in the early 2000s, and for simplicity, shall be used throughout this review.

Measurement of MASPE. Using TTE, MAPSE is measured using M-mode in the apical view (**Figure 1**). MAPSE can be measured from an annular edge or midpoint in the 2-, 3- or 4-chamber views, though commonest is the lateral annulus in the 4-chamber view. The M-mode cursor should be aligned parallel to the LV wall. MAPSE is the distance from the lowest point at end-diastole until the aortic valve closure. Movement after closure of the aortic valve, termed post-systolic shortening, is abnormal and should be excluded from the measurement[9]. MAPSE can also be measured by speckle tracking echocardiography, which automatically tracks and measures annular movement without angle dependence. This has proven to be fast and reproducible[18,19], but is only available with specific vendor software. Consequently, the majority of studies outlined have used the manual M-mode method.

Using CMR, and less commonly cardiac computed tomography (CT), MAPSE is measured on long axis views as the distance travelled by the mitral annulus between end-diastole and end-systole (**Figure 2**). Automated AI measurements have proven possible using both TTE and CMR with good agreement between manual and automated measurements (**supplementary table S2**)[15,20-22,23-25].

A brief history of MAPSE. Da Vinci provided the first documented recording of mitral annular movement in the 15[th] Century. He observed the movement of a probe-like instrument called a *spillo* after insertion through the chest wall and heart of a pig, noting the ventricles shortened during systole and lengthened in diastole by 'the thickness of a finger' [26,27]. The first brief reference to a non-invasive measurement came in 1965[28] followed by the first clinical study two years later[1]. Using A-mode, Zaky et al, described how the 'mitral ring echo' moved towards the transducer during systole and away in diastole. The mean distance of the movement of the mitral ring in 25 healthy subjects was 1.6 cm, near identical to normal values from recent TTE and CMR studies[18,29]. Later, using gated planar 201-Tl myocardial perfusion scintigraphy, Lundback found a similar normal value of approximately 19 mm [30].

M-mode allowed easier measurement in clinical practice[30], and attempts were made to define normal values in healthy subjects across age groups[31]. Through the 1980s and 90s, MAPSE was investigated in the context of various pathologies including heart failure with reduced ejection fraction[32] (HFrEF) and ischemic heart disease[33] (IHD). With advancements of 2-dimensional (2D) and 3D TTE, MAPSE appears to have been overlooked in favour of

LVEF, and more recently, GLS. However, recent years have seen a resurgence of interest in MAPSE as a marker of early systolic impairment and for prognostication.

Normal MAPSE values. Incorporation of MAPSE into imaging guidelines and routine practice requires validated normal values. The largest studies are summarised in **Table 1,** and a full summary of studies are presented in **supplementary Table S1**. All studies observed an inverse relationship with age[18,34-36].

Another consideration is from which views, annular positions and over how many cardiac cycles MAPSE should be measured. If manually measured, the addition of more views increases reporting time. However, automated measurements can be acquired in seconds[15], allowing multiple cycles to be averaged quickly, which may be of value particularly in patients with atrial fibrillation, where MAPSE is likely to vary from beat to beat.

Methods

This systematic review is presented in accordance with the PRISMA statement[37]. PubMed, Medline and Embase were searched (search terms presented in the **supplementary material**). Retrospective and prospective observational studies assessing the diagnostic and prognostic utility of MAPSE in adult humans were included. Studies of healthy cohorts and those with pathology were included. Studies were grouped into the following categories:

1. Diagnostic studies:
    a. Measuring association between MAPSE with other markers of systolic function.

b. Comparing MAPSE in populations with a known pathology vs healthy controls where LVEF is normal.

2. Prognostic studies measuring the association between MAPSE and outcomes in cardiovascular disease and comparison with LVEF and GLS where available.

Non-English publications and abstracts were excluded. Review articles were not included, but references were manually searched for additional publications. Titles were screened for suitability by two reviewers (DF and FF), and in cases of uncertainty a third reviewer (MU) was consulted. After initial screening, abstracts were independently screened by two reviewers (DF and FF). If abstracts were insufficient to determine whether the publication should be included or not, the full text was read to inform the final decision. Data from each study including author and year of publication, cohort size and underlying pathology, correlations, MAPSE values and outcome measures such as hazard rations was extracted by one reviewer (DF) and checked by another reviewer (FF). Meta-analysis of correlation coefficients (r) was calculated as a mean correlation [95% confidence interval] weighted by sample size to analyze overall correlation between MAPSE and LVEF or GLS using the Hunter-Schmidt method[38] in R (PsychMeta package, R version 4.2.3, 2023). Given the variations in MAPSE values used to measure outcomes, a meta-analysis of such values was neither advisable nor possible, and the synthesis of those review findings are described narratively.

Results

The search was performed December 2024, and 86 studies were included in the final review as shown in **Figure 3**. Summaries of included studies are presented in **Tables 2, 3 and 4**.

MAPSE as a method of diagnosing systolic impairment

1. Correlation between MAPSE and other markers of systolic function.

Limited echocardiographic windows can render accurate assessment of systolic function challenging. The relationship between MASPE and other markers of systolic function has been investigated to ascertain if MAPSE can be used when LVEF or GLS are not available.

Correlation between MAPSE and LVEF.

A meta-correlation of 18 studies (*n*=2126) reported an *r* of 0.60±0.21 [0.49–0.70] between MAPSE and LVEF (**Table 2**; **Figure 4**).

A strong correlation was reported in mixed outpatient cohorts (*r*=0.77–0.87, all *p*<0.05)[39-43], patients with IHD[33] (*r*=0.87, *p*<0.001), HFrEF[32] (*r*=0.96, *p*=<0.001), Duchenne muscular dystrophy[44] (*r*=0.71, *p*<0.001) and in ventilated patients with raised body mass index (BMI)[45] (*r*=0.87, *p*<0.001) and normal BMI[46] (*r*=0.61, *p*<0.001). Moderate correlation was seen in patients admitted to intensive care with septic shock[47,48] (*r*=0.59 and 0.52, *p*<0001), three outpatients cohorts (r=0.46–0.64[49-51], p<0.001) and in a general pre-operative clinic using point-of-care ultrasound (POCUS)[52] (*r*=0.56, *p*<0.001). The largest study (a mixed cohort all with preserved LVEF) showed, as expected, a weaker correlation (r=0.34, *p*<0.001)[17]. As LVEF incorporates radial, circumferential and longitudinal contraction, a perfect correlation with MAPSE would not be expected. These results indicate, however, that MAPSE can be used to guide systolic assessment if LVEF is not available. The optimal threshold for detecting impaired LVEF ranged from 8–12mm. An outpatient cohort (*n*=153) found mean MAPSE <12mm was associated with an LVEF<50% with 90% sensitivity and 88% specificity[41]. A retrospective analysis of transesophageal echocardiography (TEE) images in patients

undergoing coronary artery bypass grafting found the optimal cut-off was <9mm. This cutoff was associated with a reduced LVEF (<52% in men, <54% in women) with 86% sensitivity and 81% specificity[53]. During dobutamine stress TTE, MAPSE weakly correlated with LVEF at baseline (r=0.28, p<0.05), and an increase in MASPE of ≥2 mm was associated with contractile reserve of greater than 10% with 68% sensitivity and 59% specificity [54].

Correlation between MAPSE and tissue Doppler imaging (TDI).

There was a correlation between MAPSE and tissue Doppler imaging (TDI) in 92 patients (r=0.9, p<0.001)[55] and 56 healthy subjects (r=0.56, p<0.0001)[56]. Due to the limited number of studies, no meta-correlation performed was performed for this association.

Correlation between MAPSE and GLS

Meta-correlation of 11 studies (n=2847) reported a moderate correlation between MAPSE and GLS r of 0.45±0.17 [0.34–0.57] (**Table 2**; **Figure 5**).

A strong correlation was found in patients with hypertrophic cardiomyopathy (HCM)[57] (r=-0.86, p<0.001), a haemodialysis cohort[58] (r=-0.61, p<0.001), ventilated patients with septic shock[48,59] (r=-0.64 and -0.81, p<0.001) and a post-heart-transplant group[60] (r=-0.71, p<0.001). Moderate correlation was seen in patients with heart failure with preserved ejection fraction (HFpEF) both at rest (r=-0.43, p<0.001) and during exercise[61] (r=-0.59, p<0.001), patients with moderate-severe mitral regurgitation[62] (r=-0.51, p=0.0002) and moderate-severe AS [63] (r=-0.558, ,<0.001), a mixed outpatient cohort [64] (r=-0.54, p<0.001), and following treatment with anthracycline-based chemotherapy[65] (r=-0.53, p<0.001). The largest cohort (n=1683 heathy subjects) demonstrated a weaker correlation[18] (r=-0.35, p<0.001).

The heterogeneity between reported correlations likely reflects the differences in underlying pathology as well as the software used to measure GLS. Furthermore, the range of MAPSE and GLS values in patients with pathology tends to be greater than in healthy subjects, which can lead to higher correlation coefficients, hence the weakest correlation was seen in the healthy cohort[18].

2. MAPSE as a marker of reduced longitudinal systolic function when LVEF is normal.

Longitudinal function is impaired before radial or circumferential indices change. MAPSE can identify isolated longitudinal systolic dysfunction, allowing clinicians to identify patients at risk of progressive heart disease. 29 studies compared MAPSE in patients and controls, where LVEF was normal (**Table 3**).

Hypertension (HTN) or type 2 diabetes mellitus (T2DM). Six TTE studies of subjects with HTN and/or T2DM demonstrated MAPSE was reduced compared with age and sex-matched controls[23,66-70]. In one study, MAPSE was similar in cohorts with diabetes *or* HTN (13.5±2.3mm and 13.2±2.2mm, respectively) but was lower in those with both conditions (11.5±3mm, $p<0.05$), suggesting a cumulative effect. A study of 45 patients undergoing sleeve gastrectomy showed a significant increase in MAPSE 6 months post-surgery (median 16mm vs 18m, p<0.01) which may be related to the associated reduction in HTN and DM. However, several other cardiovascular risk factors were also reduced[71].

Aortic stenosis (AS). Mean MAPSE was lower in a cohort of 49 patients with AS (all grades) compared with 133 controls (9.2±2.3mm vs 10.7±2.8mm, $p<0.001$)[72]. A prospective study (*n*=78) found lateral MAPSE was lower in patients with severe AS compared with controls

(p<0.002) whilst septal MAPSE was lower in mild, moderate and severe AS (p<0.001)[73]. In 58 patients undergoing surgical replacement for severe AS, MAPSE was higher in patients with no myocardial fibrosis (10.7±1.2mm) compared with mild (8.8±1.4mm, p<0.05) or severe (4.9±1mm, p<0.001) changes on biopsy at the time of surgery. Patients with no or mild myocardial fibrosis were more likely to experience an improvement in symptoms to New York Heart Association (NYHA) grade I or II, whilst patients with severe fibrosis remained in NYHA class III or IV nine months post-surgery. LVEF did not predict improvement in symptoms[74]. Likewise, a study of 101 patients undergoing transcatheter aortic valve implantation reported an increase in MASPE, but not LVEF, post-procedure (9.1±3.2 to 10.2±3.3mm, p=0.006), though MAPSE was not related to an improvement in symptoms in this group[75]. A study of 205 patients with severe AS showed a reduction in MAPSE of >2mm per serial scan had an increased likelihood of requiring intervention (HR 1.95 [1.04–3.66] p=0.04), when LVEF remained preserved [76]. Depressed LVEF is an indication for aortic valve replacement in the absence of symptoms as per European Society of Cardiology guidelines[77], but this occurs at a late stage of the disease. As MAPSE demonstrates improvement after intervention, and may be associated with an improvement in functional class, it could act an alternative biomarker to guide timing of intervention.

Diagnosing true low-flow, low-gradient (LFLG) severe AS can be challenging. MAPSE was lower in LFLG severe AS compared with moderate (12.8±1.7mm vs 5mm±0.9mm, p<0.05)[78] and pseudosevere AS (9.8±2.6 vs 11.6±2.2mm, p<0.05) [79], and could assist in differentiating these groups where there is uncertainty.

GLS measured by gated-CT was shown to improve post transaortic valve implantation (TAVI) (-18.3 vs -18.7%, p<0.001) and was the strongest imaging biomarker associated with mortality (HR 1.08 per % decrease (1.03–1.14), p=0.005)[80]. Though MAPSE was not

specifically measured in this study, markers of longitudinal function could be easily

measured during a pre-TAVI CT to aid pre-intervention workup.

HFpEF. Five studies found MAPSE is lower in patients with diastolic dysfunction (DD) than

healthy controls[81-83 84 61]. Rather than acting as a marker of pure DD, reduced MAPSE likely

reflects concurrent longitudinal systolic dysfunction[81] as they are closely linked[9] and may

result from a higher prevalence of comorbidities associated with DD, such as HTN and DM.

In patients with HFpEF undergoing cardiopulmonary exercise testing, MAPSE was lower at

rest (10.9±2.1mm vs 12.1±2.2mm, *p*=0.008), and augmented less during exercise in patients

with DD compared with controls (12±2.2mm vs 16.2±2.7mm, *p*<0.001)[61].

MAPSE was also lower in two cohorts with LV hypertrophy (LVH) [85,86] a biomarker closely

associated with HFpEF. In disease states where the LV hypertrophies, less longitudinal

contraction is required to generate the same stroke volume, so a reduction in MAPSE

reflects increased LV mass.

HCM. 45 patients with HCM had a reduced MAPSE compared with controls (10.0±3.2 vs

13.3±3.8, *p*<0.001), and in those with outflow obstruction, septal MAPSE was reduced

compared with lateral MAPSE (9.2±3.7mm vs 11±3.6mm, *p*=0.04) reflecting greater

impairment in the septum, the area of greatest disease[87].

Marfan's syndrome. MAPSE was lower in all 5 measurement positions in a TTE study

compared with controls (*p*<0.001 for all measurements). Of note, LVEF was also lower in the

Marfan's group, but was well within normal range (66 vs 71%, *p*<0.001)[88].

Duchenne Muscular Dystrophy (DMD). In adolescents, MAPSE was reduced in the DMD cohort with and without late gadolinium enhancement (LGE) on CMR (11.6±1.9mm and 10.8±1.6mm respectively) compared with controls (13.7±2.7mm, *p*<0.01)[89].

ARCD. Baseline MAPSE was lower in those who did not recover systolic function compared with those with good recovery (11.7±1.5 vs 15.7±3.1mm, *p*=0.03) [90] from ARCD, suggesting MAPSE can identify those most at risk from ongoing impaired function when ARCD occurs. Two studies demonstrated no difference in pre-chemotherapy MAPSE in patients who developed ARCD compared with those who did not[91,92].

Hypothyroidism. A study of 81 patients with hypothyroidism found lateral MAPSE was lower than controls with similar rates of typical cardiovascular risk factors and normal LVEF (1.66±0.2 vs 1.49±0.3mm, p=0.002[93]). Impaired cardiac relaxation is well documented in hypothyroidism[94], but impaired systolic function may be masked as LVEF remains normal.

Mixed connective tissue disease (MCTD). MAPSE was significantly lower in 77 patients with MCTD and normal LVEF and GLS compared with controls[95] (15.3±2.3 vs 13.7±2.1, p<0.001).

MAPSE as a prognostic biomarker

The prognostic capability of MAPSE has been investigated in 28 studies across a range of pathologies (**Table 4**). Despite being a simpler measurement to obtain, it has been shown to outperform LVEF and GLS in several circumstances.

General outpatient cohorts. In the largest cohort including individuals with pathology and normal scans undergoing outpatient CMR (*n*=1572), MAPSE was a better predictor of death or hospitalisation for heart failure (HR 1.7 [1.5–2.0] per mm decrease, *p*<0.001) compared with GLS (HR 1.5 [1.3–1.7], *p*<0.001) or LVEF (HR 1.4 [1.2–1.6], *p*<0.001) on multivariable analysis[15]. In that study, a convolutional neural network was used for automated measurements of MAPSE from 6 positions across 2-chamber, 3-chamber and 4-chamber long axis views, taking on average 1.1 seconds per case. In a similarly heterogeneous cohort of 400 patients followed up for 14 months, lateral MAPSE was the strongest imaging-based predictor of MACE after multivariable analysis (HR 1.34 per 2mm decrease, *p*=0.02) whereas neither LVEF nor the presence of LGE remained associated[16]. In a TTE-based study of 512 patients, lateral MAPSE was the only imaging-based predictor of mortality (HR 0.01 [0.00–0.52], *p*=0.02)[17]. Mortality risk for the group with a MAPSE <12mm was higher as early as 30 days after imaging (*p*<0.001).

HFrEF. A multicentre study of 1040 patients undergoing CMR for left ventricular systolic dysfunction showed lateral MAPSE was the strongest imaging-based biomarker in predicting mortality (HR 2.1 [1.5–2.8] per mm decrease, *p*<0.001). Presence of LGE was a weaker predictor (HR 2.0 [1.3–3.0], *p*=0.02) whilst LVEF was not associated. Patients with median MAPSE >9mm had a similar survival regardless if their LVEF was above or below 35% (*p*=0.96)[96]. A TTE study of 228 patients with HFrEF showed MAPSE, but not LVEF using the Teichholz method, predicted mortality in multivariable analysis[97] (HR 0.80 [0.66–0.97], *p*<0.02). Another study of 181 patients admitted to hospital with heart failure demonstrated no deaths in the cohort with MAPSE>10mm, though no comparison was made to LVEF[98]. In patients with end-stage heart failure requiring extracorporeal membrane oxygenation,

MAPSE >8.9mm was associated with weaning success and survival after weaning (HR 6.7 [2.7–16.6], $p<0.001$), where LVEF was not associated[99].

IHD. MAPSE measured by CMR in 255 patients 2 days post-ST elevation myocardial infarction (STEMI) was a better predictor of major adverse cardiac events (MACE) (death, MI, stroke, heart failure) than LVEF in a cohort followed up for 3 years (area under curve 0.74 vs 0.61, $p<0.01$)[100]. MAPSE <9mm remained a predictor in models of clinical and imaging risk factors (HR 6.0 [2.5–14.7], $p<0.01$) where LVEF was not associated. Another study of 165 patients post-acute coronary syndrome (ACS) also found MAPSE was a stronger predictor for MACE than LVEF (HR 0.83 [0.73–0.94], $p=0.004$ vs HR 0.96 [0.94–0.99], $p=0.01$)[101]. The optimal cut-off for predicting survival was 9.7mm (sensitivity 79%; specificity 72%). In a study of 271 patients admitted to hospital with suspected ACS, the group in the lowest MAPSE quartile (<8mm) had a risk ratio of 3.0 ($p<0.001$) for death or heart failure hospitalisation during follow-up compared with the remainder of the cohort[102]. In patients admitted with suspected ACS (n=233) lateral MAPSE on CMR predicted MACE during follow up (HR 1.4 [1.1–1.8])[103] whereas LVEF did not. Conversely, another CMR study of 445 patients 2-4 days post-revascularisation for ST-elevation myocardial infarction found MAPSE was a predictor in univariable analysis (HR 0.80 [0.72–0.89]), but not when adjusted for other imaging biomarkers[104]. Possible reasons for this difference include use of CMR-based adjustment for infarct size or presence of microvascular obstruction, which were not adjusted for in other studies. In 333 patients with stable IHD, MAPSE was the only clinical or imaging marker that correlated independently with mortality (OR 1.6 [1.4–2.0] per mm decrease, $p<0.001$)[105]. MAPSE was lower in those who died compared with survivors (8.3±2.1mm vs 11.5±2.1mm, $p<0.001$) or those who died of non-cardiac causes (11.0±0.8mm, $p<0.001$).

Takotsubo cardiomyopathy. A study of 53 patients showed mortality was higher in patients with MAPSE <10mm on their presenting TTE after 12 months follow-up, but there was no mortality difference within the first year. MAPSE <10mm was a predictor of mortality (HR 5.1 [1.3–19.2], *p*<0.01), whilst LVEF <35% was not (*p*=0.23)[106].

HTN. In 1735 patients with hypertension followed up for 5.1 years after a CMR, lateral MAPSE was associated with increased risk of death (HR 1.4 [1.3–1.5] per mm decrease, *p*<0.001) in the whole cohort, as well as in in the subgroup analysis of patients with a preserved LVEF (HR 1.3 [1.2–1.5] per mm decrease, *p*<0.001)[107]. LVEF was a predictor in univariable, but not multivariable analysis. GLS was a predictor of MACE in multivariable analysis, but weaker than MAPSE (HR 1.18 [1.14–1.22], *p*<0.001). A TTE study of 156 patients with HTN found mean MAPSE was a predictor of MACE in multivariable analysis (HR 0.81 [0.67–0.98] per standard deviation of reduction, *p*<0.05)[108].

HCM. Mean MAPSE was lower in those with MACE, defined as death, transplant or ventricular arrhythmia (6.9±1.9mm vs 10.2±3.1mm, *p*=0.01) with no difference in LVEF (*p*=0.86)[87].

Amyloidosis In 191 patients with wildtype transthyretin amyloidosis, MAPSE was the only imaging-based predictor of mortality in multivariable analysis (HR 0.19 [0.04–0.83], *p*=0.003)[109]. A CMR-based study of 68 AL amyloid patients showed lateral MAPSE was a stronger predictor of death or heart transplant (HR 0.23 [0.10–0.56] per mm decrease, *p*=0.001) than LVEF or GLS[110]. However, two TTE studies of patients with AL amyloid found

MAPSE predicted death in uni-, but not multivariable analysis. [111] [112]. MAPSE was a stronger predictor than LVEF in both cohorts, but performed worse than GLS.

Heart transplant. Heart transplant patients require vigilant follow-up but often have challenging echocardiographic windows. MAPSE measured by STE at the midpoint of the mitral annulus in 155 patients was a stronger predictor of death and hospitalisation than GLS (HR 0.68 [0.56–0.83] vs 0.76 [0.66–0.87], p<0.001) or LVEF (not associated)[60]. Moreover, measurement of MAPSE was possible in a greater proportion of transplanted patients (86%) compared to LVEF or GLS (72%)[60].

Pulmonary arterial hypertension (PAH). In 71 patients with PAH, CMR-MAPSE was higher in the group that survived without transplant compared with non-survivors (12.3±3mm vs 10±3mm, p=0.02). MAPSE predicted death or transplantation (HR 1.2 [1.0–1.3] per mm decrease p=0.01). LVEF and right ventricular EF were not associated[113].

Septic shock. Studies assessing MAPSE as a prognostic marker in the setting of septic shock have generated mixed results. Three studies found MAPSE to be the sole or strongest imaging biomarker for predicting mortality in multivariable analysis[47,114,115]. A study of 152 patients showed MASPE was a predictor of myocardial injury (troponin T level >45nmol/ml), but not of mortality[116]. The largest intensive care-based study of 451 patients found MAPSE univariably associated with inpatient mortality (r=0.79, p<0.01), but did not remain associated in multivariable analysis[117]. LVEF was not a predictor of mortality in any study. The varying results may reflect the heterogeneity of patients admitted to the intensive care for sepsis, who vary markedly in baseline comorbidities, frailty and inotropic drug use. Most

intensive care-based studies follow-up patients for 30 days or less, and the prognostic ability of MAPSE seen may be over years, so longer follow up may be required which is often not pragmatic in intensive care-based studies.

Atrial fibrillation (AF). A study of 160 patients with AF showed MASPE was a predictor of mortality (OR 1.4 [1.1–1.7], p<0.01) where LVEF was not.[118].

Acute pulmonary embolism (PE). In 362 patients with CT-proven acute PE, MAPSE <11mm predicted mortality or the need for advanced therapies among patients with a preserved LVEF (OR 3.5 [1.4–9.0], *p*=0.01). MAPSE <11mm combined with tricuspid annular plane systolic excursion (TAPSE) <16mm was strongly associated with adverse outcomes (OR 10.8 [3.1–37.8], *p*<0.01). LVEF <50% was also an adverse prognostic marker (OR 4.1 [1.4–12.0], *p*=0.011)[119].

Coronavirus disease-2019 (COVID-19). In a retrospective TTE study of 68 patients admitted to hospital with severe COVID-19, MAPSE was the only left-heart imaging biomarker that differed in patients who died compared with survivors (11±3 vs 12±3mm, *p*=0.006)[120]. In multivariable analysis, MAPSE and right atrial volume index (RAVI) were predictors of mortality (MAPSE HR 1.6 [1.2–2.2], *p*=0.006; RAVI HR 1.1 [1.0–1.3], *p*=0.01). Difference in RAVI is likely to represent the higher ventilatory requirements of the more unwell patients.

Discussion

This review found a moderate-to-strong correlation between MAPSE and LVEF, and a moderate correlation between MAPSE and GLS. In the presence of a poor echocardiographic

acoustic window it is reasonable and appropriate to use MAPSE to assess LV function. In particular, a MAPSE of <8mm is associated with LVEF<50% in most studies.

In cohorts with a normal LVEF, MAPSE was lower in patients with HTN, T2DM, aortic AS, HCM, Marfan's syndrome, Duchenne Muscular Dystrophy, hypothyroidism and mixed connective tissue disease compared with matched healthy controls. Measuring MAPSE routinely in these conditions would allow clinicians to detect early cardiac impairment that may otherwise be missed. Although no evidence-based treatments currently exist for isolated longitudinal dysfunction, identifying patients with impaired MAPSE may highlight those most at risk of progressive disease. MAPSE also improved following surgery for AS, and may represent a biomarker that could guide timing of intervention when LVEF remains normal.

In the largest study of general outpatients, MAPSE was the strongest predictor of death or hospitalisation, outperforming LVEF and GLS[15]. MAPSE predicted mortality or MACE in cohorts with HFrEF, IHD, Takotsubo cardiomyopathy, HTN, amyloidosis, post-heart transplant, pulmonary hypertension, AF and COVID-19, where LVEF was either a weaker predictor or not associated. MAPSE was also a stronger predictor than GLS in cohorts with hypertension and amyloidosis. Routine incorporation of MAPSE into clinical studies with any of the above conditions would seem appropriate to identify the most at-risk patients.

Other clinical benefits of incorporating MAPSE into routine imaging

A TTE study in 284 children showed image quality did not affect reproducibility of MAPSE (p>0.24)[121], which otherwise hugely impacts interpretation of LVEF and GLS[122]. Less training is required to obtain or interpret MAPSE, making it a convenient biomarker in settings

where full accreditation in cardiac imaging is difficult to attain, such as emergency medicine or critical care[47,123].

High intra- and interobserver reproducibility has been reported for both TTE and CMR MAPSE measurements (TTE R^2=0.88 and 0.74; CMR R^2=0.76 and 0.72, p<0.01 for all)[124].There was no difference between TTE and CMR values in a study of 111 patients (p=0.1), meaning the same normal values can be applied to both modalities, and unlike LVEF, either can be used for follow-up[124]. MAPSE also correlates closely with left atrial strain (LAS) (r=0.65, p<0.001), a marker of left atrial function and diastolic dysfunction[125]. Routine measurement of MAPSE would ensure LAS results are not biased by longitudinal function. Automated measurement or tracking of MAPSE is relatively straight-forward compared with LVEF or GLS. Automated results show good agreement compared with manual measurements[20] and less variability than human measurements in scan-rescan datasets[15] (supplementary material **Table S2)**.

POCUS increasingly guides rapid decision-making in time-critical specialties such as emergency medicine and anaesthetics. MAPSE has been proposed for real-time monitoring of patients during high-risk periods of their care, and can be automatically tracked using algorithms such as autoMAPSE. In a retrospective offline study, autoMAPSE has been validated against manually-measured MAPSE in 40 patients following cardiac surgery, demonstrating 95% limits of agreement of –3.7 to 4.5mm[126]. Another study performed in real-time at the bedside of 30 stable and 50 critical care patients showed autoMAPSE was able to obtain measurements in 88% of stable and 89% of critical care patients. Compared with manual measurements, there was a difference of 0.8 [–2.9–4.7] mm in the stable patients and difference of 0.6mm [–2.3–3.5] mm in critical care patients[127]. Real-time monitoring of systolic function could detect early signs of deterioration before end-organ

perfusion is affected, and MAPSE could provide an objective quantification method, rather than relying on subjective visual estimation.

Limitations of using MAPSE

The main limitation with MAPSE is the lack of regional assessment of LV function[128] for which visual assessment and GLS will remain essential. Measuring MAPSE using M-mode on TTE is angle dependent, requiring on-axis apical views. Off-axis views may underestimate MAPSE, reducing specificity. However, MAPSE can readily be measured in an angle-independent fashion using a simple linear caliper applied to a 2D image. MAPSE interpretation is limited in patients with a large pericardial effusions, as some longitudinal movement of the annulus may arise from movement of the heart inside the pericardial sac. Although uncommon, caution must also be taken in patients with paradoxical septal motion secondary to constrictive pericarditis, as septal annular movement arises from the right ventricle, as the LV is fixed by the thickened pericardium, rendering MAPSE less useful in this context. MAPSE is also reduced in patients following mitral valve replacement[129], when it may moreso reflect post-surgical changes rather than inherent LV myocardial function. Finally, as mentioned, widespread application of MAPSE is currently limited by a lack of standardised practice. However, the generation of normal values from large datasets may lead to the inclusion of MAPSE in guidelines, and with it set the standards for acquisition, measurement, and interpretation.

Limitations of this review

Within the prognostic studies, different limits of MAPSE were used to determine the presence or absence of the outcome, and as such no meta-analysis of these results could be performed.

Conclusions

MAPSE is a marker of longitudinal LV function and offers a wealth of clinical information. Measurement of MAPSE can be performed almost universally, allowing assessment of LV function even with compromised image quality. Early systolic impairment that would otherwise be missed when using LVEF alone can be detected using MAPSE, highlighting patients most at risk of progressive systolic dysfunction. MAPSE also aids in prognostication across multiple cardiac conditions including hypertension, aortic stenosis, and ischaemic heart disease. Despite the advancement of more sophisticated techniques within cardiac imaging, new studies continue to highlight the clinical information that MAPSE has to offer clinicians and patients. Those contemplating using MAPSE in their clinical practice may consider both the robust data supporting its use presented herein, and da Vinci's timeless sage words that "simplicity is the ultimate sophistication" [130].

References

1.	Zaky A, Grabhorn L, Feigenbaum H. Movement of the mitral ring: a study in ultrasoundcardiography. *Cardiovasc Res* 1967; **1**(2): 121-31.

2.	Carlsson M, Ugander M, Heiberg E, Arheden H. The quantitative relationship between longitudinal and radial function in left, right, and total heart pumping in humans. *Am J Physiol Heart Circ Physiol* 2007; **293**(1): H636-44.

3.	Carlsson M, Ugander M, Mosen H, Buhre T, Arheden H. Atrioventricular plane displacement is the major contributor to left ventricular pumping in healthy adults, athletes, and patients with dilated cardiomyopathy. *Am J Physiol Heart Circ Physiol* 2007; **292**(3): H1452-9.

4.	Asgeirsson D, Hedstrom E, Jogi J, et al. Longitudinal shortening remains the principal component of left ventricular pumping in patients with chronic myocardial infarction even when the absolute atrioventricular plane displacement is decreased. *BMC Cardiovasc Disord* 2017; **17**(1): 208.

5.	Robinson S, Rana B, Oxborough D, et al. A practical guideline for performing a comprehensive transthoracic echocardiogram in adults: the British Society of Echocardiography minimum dataset. *Echo Res Pract* 2020; **7**(4): G59-G93.

6.	Mitchell C, Rahko PS, Blauwet LA, et al. Guidelines for Performing a Comprehensive Transthoracic Echocardiographic Examination in Adults: Recommendations from the American Society of Echocardiography. *J Am Soc Echocardiogr* 2019; **32**(1): 1-64.

7.	Galderisi M, Cosyns B, Edvardsen T, et al. Standardization of adult transthoracic echocardiography reporting in agreement with recent chamber quantification, diastolic function, and heart valve disease recommendations: an expert consensus document of the European Association of Cardiovascular Imaging. *Eur Heart J Cardiovasc Imaging* 2017; **18**(12): 1301-10.

8.	Kramer CM, Barkhausen J, Bucciarelli-Ducci C, Flamm SD, Kim RJ, Nagel E. Standardized cardiovascular magnetic resonance imaging (CMR) protocols: 2020 update. *J Cardiovasc Magn Reson* 2020; **22**(1): 17.

9.	Hu K, Liu D, Herrmann S, et al. Clinical implication of mitral annular plane systolic excursion for patients with cardiovascular disease. *Eur Heart J Cardiovasc Imaging* 2013; **14**(3): 205-12.

10.	Stanton T, Marwick TH. Assessment of subendocardial structure and function. *JACC Cardiovasc Imaging* 2010; **3**(8): 867-75.

11.	Henein MY, Gibson DG. Long axis function in disease. *Heart* 1999; **81**(3): 229-31.

12.	Vancheri F, Longo G, Henein MY. Left ventricular ejection fraction: clinical, pathophysiological, and technical limitations. *Front Cardiovasc Med* 2024; **11**: 1340708.

13.	Mizuguchi Y, Oishi Y, Miyoshi H, Iuchi A, Nagase N, Oki T. The functional role of longitudinal, circumferential, and radial myocardial deformation for regulating the early impairment of left ventricular contraction and relaxation in patients with cardiovascular risk factors: a study with two-dimensional strain imaging. *J Am Soc Echocardiogr* 2008; **21**(10): 1138-44.

14.	Slawinski G, Hawryszko M, Lizewska-Springer A, Nabialek-Trojanowska I, Lewicka E. Global Longitudinal Strain in Cardio-Oncology: A Review. *Cancers (Basel)* 2023; **15**(3).

15.	Xue H, Artico J, Davies RH, et al. Automated In-Line Artificial Intelligence Measured Global Longitudinal Shortening and Mitral Annular Plane Systolic Excursion: Reproducibility and Prognostic Significance. *J Am Heart Assoc* 2022; **11**(4): e023849.

16.   Rangarajan V, Chacko SJ, Romano S, et al. Left ventricular long axis function assessed during cine-cardiovascular magnetic resonance is an independent predictor of adverse cardiac events. *J Cardiovasc Magn Reson* 2016; **18**(1): 35.

17.   Ozer PK, Govdeli EA, Demirtakan ZG, et al. The relation of echo-derived lateral MAPSE to left heart functions and biochemical markers in patients with preserved ejection fraction: Short-term prognostic implications. *J Clin Ultrasound* 2022; **50**(5): 593-600.

18.   Wang YH, Sun L, Li SW, et al. Normal reference values for mitral annular plane systolic excursion by motion-mode and speckle tracking echocardiography: a prospective, multicentre, population-based study. *Eur Heart J Cardiovasc Imaging* 2023; **24**(10): 1384-93.

19.   Buss SJ, Mereles D, Emami M, et al. Rapid assessment of longitudinal systolic left ventricular function using speckle tracking of the mitral annulus. *Clin Res Cardiol* 2012; **101**(4): 273-80.

20.   Yoon SS, Fischer C, Amsel D, et al. Fully automated AI-based cardiac motion parameter extraction - application to mitral and tricuspid valves on long-axis cine MR images. *Eur J Radiol* 2023; **166**: 110978.

21.   Seemann F, Pahlm U, Steding-Ehrenborg K, et al. Time-resolved tracking of the atrioventricular plane displacement in Cardiovascular Magnetic Resonance (CMR) images. *BMC Med Imaging* 2017; **17**(1): 19.

22.   Tasken AA, Berg EAR, Grenne B, et al. Automated estimation of mitral annular plane systolic excursion by artificial intelligence from 3D ultrasound recordings. *Artif Intell Med* 2023; **144**: 102646.

23.   Li Y, Wu C, Li Y. Feasibility study of automated cardiac motion quantification to assess left ventricular function in type 2 diabetes. *Sci Rep* 2023; **13**(1): 1101.

24.   Gonzales RA, Seemann F, Lamy J, et al. MVnet: automated time-resolved tracking of the mitral valve plane in CMR long-axis cine images with residual neural networks: a multi-center, multi-vendor study. *J Cardiovasc Magn Reson* 2021; **23**(1): 137.

25.   Magelssen MI, Palmer CL, Hjorth-Hansen A, et al. Feasibility and Reliability of Automatic Quantitative Analyses of Mitral Annular Plane Systolic Excursion by Handheld Ultrasound Devices: A Pilot Study. *J Ultrasound Med* 2021; **40**(2): 341-50.

26.   Wandt B. Long-axis contraction of the ventricles: a modern approach, but described already by Leonardo da Vinci. *J Am Soc Echocardiogr* 2000; **13**(7): 699-706.

27.   Keele KD. Leonardo da Vinci's Corpus of Anatomical Studies. *Hist Med* 1980; **8**(4): 4-5.

28.   Edler I. Examination of the Cardiac Patient. New York: A. A. McGraw Hill; 1965.

29.   Grue JF, Storve S, Dalen H, et al. Automatic Measurements of Mitral Annular Plane Systolic Excursion and Velocities to Detect Left Ventricular Dysfunction. *Ultrasound Med Biol* 2018; **44**(1): 168-76.

30.   Lundback S. Cardiac pumping and function of the ventricular septum. *Acta Physiol Scand Suppl* 1986; **550**: 1-101.

31.   Hoglund C, Alam M, Thorstrand C. Atrioventricular valve plane displacement in healthy persons. An echocardiographic study. *Acta Med Scand* 1988; **224**(6): 557-62.

32.   Alam M, Hoglund C, Thorstrand C. Longitudinal systolic shortening of the left ventricle: an echocardiographic study in subjects with and without preserved global function. *Clin Physiol* 1992; **12**(4): 443-52.

33.   Alam M. The atrioventricular plane displacement as a means of evaluating left ventricular systolic function in acute myocardial infarction. *Clin Cardiol* 1991; **14**(7): 588-94.

34.   Grue JF, Storve S, Stoylen A, et al. Normal ranges for automatic measurements of tissue Doppler indices of mitral annular motion by echocardiography. Data from the HUNT3 Study. *Echocardiography* 2019; **36**(9): 1646-55.

35.   Ochs MM, Fritz T, Andre F, et al. A comprehensive analysis of cardiac valve plane displacement in healthy adults: age-stratified normal values by cardiac magnetic resonance. *Int J Cardiovasc Imaging* 2017; **33**(5): 721-9.

36.   Leng S, Zhao X, Koh AS, et al. Age-related changes in four-dimensional CMR-derived atrioventricular junction velocities and displacements: Implications for the identification of altered annular dynamics for ventricular function assessment. *Int J Cardiol Heart Vasc* 2019; **22**: 6-12.

37.   Moher D, Liberati A, Tetzlaff J, Altman DG, Group P. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009; **151**(4): 264-9, W64.

38.   Hunter JE SF. Methods of Meta-Analysis: Correcting Error and Bias in Research Findings. 2nd ed. Newbury Park, California.: Sage; 1990.

39.   Simonson JS, Schiller NB. Descent of the base of the left ventricle: an echocardiographic index of left ventricular function. *J Am Soc Echocardiogr* 1989; **2**(1): 25-35.

40.   Qin JX, Shiota T, Tsujino H, et al. Mitral annular motion as a surrogate for left ventricular ejection fraction: real-time three-dimensional echocardiography and magnetic resonance imaging studies. *Eur J Echocardiogr* 2004; **5**(6): 407-15.

41.   Elnoamany MF, Abdelhameed AK. Mitral annular motion as a surrogate for left ventricular function: correlation with brain natriuretic peptide levels. *Eur J Echocardiogr* 2006; **7**(3): 187-98.

42.   Tsang W, Ahmad H, Patel AR, et al. Rapid estimation of left ventricular function using echocardiographic speckle-tracking of mitral annular displacement. *J Am Soc Echocardiogr* 2010; **23**(5): 511-5.

43.   Bazaz R, Edelman K, Gulyasy B, Lopez-Candales A. Evidence of robust coupling of atrioventricular mechanical function of the right side of the heart: insights from M-mode analysis of annular motion. *Echocardiography* 2008; **25**(6): 557-61.

44.   Turan O, Kocabas A. Disturbed Atrial Conduction in Patients with Duchenne Muscular Dystrophy. *Anatol J Cardiol* 2024; **28**(10): 493-8.

45.   Zidan DH, Helmy TA. Usefulness of mitral annular plane systolic excursion in assessment of left ventricular systolic function in mechanically ventilated obese patients. *J Crit Care* 2016; **34**: 74-6.

46.   Shah A, Nanjayya V, Ihle J. Mitral Annular Plane Systolic Excursion as a predictor of Left Ventricular Ejection Fraction in mechanically ventilated patients. *Australas J Ultrasound Med* 2019; **22**(2): 138-42.

47.   Bergenzaun L, Ohlin H, Gudmundsson P, Willenheimer R, Chew MS. Mitral annular plane systolic excursion (MAPSE) in shock: a valuable echocardiographic parameter in intensive care patients. *Cardiovasc Ultrasound* 2013; **11**: 16.

48.   Brault C, Zerbib Y, Mercado P, et al. Mitral annular plane systolic excursion for assessing left ventricular systolic dysfunction in patients with septic shock. *BJA Open* 2023; **7**: 100220.

49.   Matos J, Kronzon I, Panagopoulos G, Perk G. Mitral annular plane systolic excursion as a surrogate for left ventricular ejection fraction. *J Am Soc Echocardiogr* 2012; **25**(9): 969-74.

50.   Lopez-Candales A, Hernandez-Suarez DF, Lopez Menendez F. Mitral Annular Dynamics and Left Ventricular Diastole. *Cardiol Res* 2017; **8**(5): 228-31.

51.   Hernandez Burgos PM, Lopez Menedez F, Candales MD, Lopez-Candales A. Is mitral annular ascent useful in studying left ventricular function through left atrio-ventricular interactions? *Indian Heart J* 2018; **70**(3): 368-72.

52.   Stenberg Y, Wallinder L, Lindberg A, Wallden J, Hultin M, Myrberg T. Preoperative Point-of-Care Assessment of Left Ventricular Systolic Dysfunction With Transthoracic Echocardiography. *Anesth Analg* 2021; **132**(3): 717-25.

53.   Borde DP, Joshi S, Asegaonkar B, et al. Mitral Annular Plane Systolic Excursion: A Simple, Reliable Echocardiographic Parameter to Detect Left Ventricular Systolic Dysfunction in Patients Undergoing Off-Pump Coronary Artery Bypass Grafting with Transesophageal Echocardiography. *J Cardiothorac Vasc Anesth* 2019; **33**(5): 1334-9.

54.   Magdy G, Hamdy E, Elzawawy T, Ragab M. Value of mitral annular plane systolic excursion in the assessment of contractile reserve in patients with ischemic cardiomyopathy before cardiac revascularization. *Indian Heart J* 2018; **70**(3): 373-8.

55.   Ballo P, Bocelli A, Motto A, Mondillo S. Concordance between M-mode, pulsed Tissue Doppler, and colour Tissue Doppler in the assessment of mitral annulus systolic excursion in normal subjects. *Eur J Echocardiogr* 2008; **9**(6): 748-53.

56.   Mondillo S, Galderisi M, Ballo P, Marino PN, Study Group of Echocardiography of the Italian Society of C. Left ventricular systolic longitudinal function: comparison among simple M-mode, pulsed, and M-mode color tissue Doppler of mitral annulus in healthy individuals. *J Am Soc Echocardiogr* 2006; **19**(9): 1085-91.

57.   Liu L, Tuo S, Zhang J, et al. Reduction of left ventricular longitudinal global and segmental systolic functions in patients with hypertrophic cardiomyopathy: Study of two-dimensional tissue motion annular displacement. *Exp Ther Med* 2014; **7**(6): 1457-64.

58.   Chiu DY, Abidin N, Hughes J, Sinha S, Kalra PA, Green D. Speckle tracking determination of mitral tissue annular displacement: comparison with strain and ejection fraction, and association with outcomes in haemodialysis patients. *Int J Cardiovasc Imaging* 2016; **32**(10): 1511-8.

59.   Johansson Blixt P, Chew MS, Ahman R, et al. Left ventricular longitudinal wall fractional shortening accurately predicts longitudinal strain in critically ill patients with septic shock. *Ann Intensive Care* 2021; **11**(1): 52.

60.   Ji X, Zhang Y, Xie Y, et al. Feasibility and prognostic value of tissue motion annular displacement in patients with heart transplantation. *Echocardiography* 2024; **41**(4): e15809.

61.   Wenzelburger FW, Tan YT, Choudhary FJ, Lee ES, Leyva F, Sanderson JE. Mitral annular plane systolic excursion on exercise: a simple diagnostic tool for heart failure with preserved ejection fraction. *Eur J Heart Fail* 2011; **13**(9): 953-60.

62.   Teraguchi I, Hozumi T, Takemoto K, et al. Assessment of decreased left ventricular longitudinal deformation in asymptomatic patients with organic mitral regurgitation and preserved ejection fraction using tissue-tracking mitral annular displacement by speckle-tracking echocardiography. *Echocardiography* 2019; **36**(4): 678-86.

63.   Luszczak J, Olszowska M, Drapisz S, et al. Assessment of left ventricle function in aortic stenosis: mitral annular plane systolic excursion is not inferior to speckle tracking echocardiography derived global longitudinal peak strain. *Cardiovasc Ultrasound* 2013; **11**: 45.

64.   Lopez-Candales A, Hernandez-Suarez DF, Menendez FL. Are Measures of Left Ventricular Longitudinal Shortening Affected by Left Atrial Enlargement? *Cardiol Res* 2018; **9**(1): 1-6.

65.   Zhang W, Azibani F, Libhaber E, et al. The role of conventional echocardiographic parameters on detecting subclinical anthracycline therapy related cardiac dysfunction-The SATRACD study. *Front Cardiovasc Med* 2022; **9**: 966230.

66.   Xiao HB, Kaleem S, McCarthy C, Rosen SD. Abnormal regional left ventricular mechanics in treated hypertensive patients with 'normal left ventricular function'. *Int J Cardiol* 2006; **112**(3): 316-21.

67.   Koulouris SN, Kostopoulos KG, Triantafyllou KA, et al. Impaired systolic dysfunction of left ventricular longitudinal fibers: a sign of early hypertensive cardiomyopathy. *Clin Cardiol* 2005; **28**(6): 282-6.

68.   Ballo P, Cameli M, Mondillo S, et al. Impact of diabetes and hypertension on left ventricular longitudinal systolic function. *Diabetes Res Clin Pract* 2010; **90**(2): 209-15.

69.   Salas-Pacheco JL, Lomeli-Sanchez O, Baltazar-Gonzalez O, Soto ME. Longitudinal systolic dysfunction in hypertensive cardiomyopathy with normal ejection fraction. *Echocardiography* 2022; **39**(1): 46-53.

70.   Loncarevic B, Trifunovic D, Soldatovic I, Vujisic-Tesic B. Silent diabetic cardiomyopathy in everyday practice: a clinical and echocardiographic study. *BMC Cardiovasc Disord* 2016; **16**(1): 242.

71.   Tromba L, Tartaglia F, Carbotta S, et al. The Role of Sleeve Gastrectomy in Reducing Cardiovascular Risk. *Obes Surg* 2017; **27**(5): 1145-51.

72.   Rydberg E, Gudmundsson P, Kennedy L, Erhardt L, Willenheimer R. Left atrioventricular plane displacement but not left ventricular ejection fraction is influenced by the degree of aortic stenosis. *Heart* 2004; **90**(10): 1151-5.

73.   Takeda S, Rimington H, Smeeton N, Chambers J. Long axis excursion in aortic stenosis. *Heart* 2001; **86**(1): 52-6.

74.   Weidemann F, Herrmann S, Stork S, et al. Impact of myocardial fibrosis in patients with symptomatic severe aortic stenosis. *Circulation* 2009; **120**(7): 577-84.

75.   Kempny A, Diller GP, Kaleschke G, et al. Longitudinal left ventricular 2D strain is superior to ejection fraction in predicting myocardial recovery and symptomatic improvement after aortic valve implantation. *Int J Cardiol* 2013; **167**(5): 2239-43.

76.   Matos JD, Kiss JE, Locke AH, Goldberg RI, Mukamal K, Manning WJ. Relation of the Mitral Annular Plane Systolic Excursion to Risk for Intervention in Initially Asymptomatic Patients With Aortic Stenosis and Preserved Systolic Function. *Am J Cardiol* 2017; **120**(11): 2031-4.

77.   Vahanian A, Beyersdorf F, Praz F, et al. 2021 ESC/EACTS Guidelines for the management of valvular heart disease. *Eur Heart J* 2022; **43**(7): 561-632.

78.   Herrmann S, Stork S, Niemann M, et al. Low-gradient aortic valve stenosis myocardial fibrosis and its influence on function and outcome. *J Am Coll Cardiol* 2011; **58**(4): 402-12.

79.   Liu D, Hu K, Liebner E, et al. Value of low-dose dobutamine stress echocardiography on defining true severe low gradient aortic stenosis in patients with preserved left ventricular ejection fraction. *Int J Cardiovasc Imaging* 2018; **34**(12): 1877-87.

80.   Fukui M, Hashimoto G, Lopes BBC, et al. Association of baseline and change in global longitudinal strain by computed tomography with post-transcatheter aortic valve replacement outcomes. *Eur Heart J Cardiovasc Imaging* 2022; **23**(4): 476-84.

81.   Dursunoglu D, Polat B, Evrengul H, Tanriverdi H, Kaftan A, Kilic M. Assessment of systolic function by atrioventricular plane displacement in patients with diastolic dysfunction. *Acta Cardiol* 2004; **59**(4): 409-15.

82.   Gromadzinski L, Pruszczyk P. Echocardiographic changes in patients with stage 3-5 chronic kidney disease and left ventricular diastolic dysfunction. *Cardiorenal Med* 2014; **4**(3-4): 234-43.

83.   Tasolar H, Mete T, Cetin M, et al. Mitral annular plane systolic excursion in the assessment of left ventricular diastolic dysfunction in obese adults. *Anatol J Cardiol* 2015; **15**(7): 558-64.

84.   Bytyci I, Bajraktari G. Left atrial changes in early stages of heart failure with preserved ejection fraction. *Echocardiography* 2016; **33**(10): 1479-87.

85.   Lopez-Candales A. Automated functional imaging for assessment of left ventricular mechanics in the presence of left ventricular hypertrophy. *Echocardiography* 2014; **31**(5): 605-14.

86.   Wierzbowska-Drabik K, Chrzanowski L, Kapusta A, et al. Severe obesity impairs systolic and diastolic heart function - the significance of pulsed tissue Doppler, strain, and strain rate parameters. *Echocardiography* 2013; **30**(8): 904-11.

87.   Doesch C, Sperb A, Sudarski S, et al. Mitral annular plane systolic excursion is an easy tool for fibrosis detection by late gadolinium enhancement cardiovascular magnetic resonance imaging in patients with hypertrophic cardiomyopathy. *Arch Cardiovasc Dis* 2015; **108**(6-7): 356-66.

88.   Kiotsekoglou A, Bajpai A, Bijnens BH, et al. Early impairment of left ventricular long-axis systolic function demonstrated by reduced atrioventricular plane displacement in patients with Marfan syndrome. *Eur J Echocardiogr* 2008; **9**(5): 605-13.

89.   Panovsky R, Pesl M, Machal J, et al. Quantitative assessment of left ventricular longitudinal function and myocardial deformation in Duchenne muscular dystrophy patients. *Orphanet J Rare Dis* 2021; **16**(1): 57.

90.   Harries I, Biglino G, Ford K, et al. Prospective multiparametric CMR characterization and MicroRNA profiling of anthracycline cardiotoxicity: A pilot translational study. *Int J Cardiol Heart Vasc* 2022; **43**: 101134.

91.   Ichikawa N, Nishizaki Y, Miyazaki S, et al. Efficacy of mitral annular velocity as an alternative marker of left ventricular global longitudinal strain to detect the risk of cancer therapy-related cardiac disorders. *Echocardiography* 2024; **41**(7): e15877.

92.   Botelho LFB, de Melo MDT, de Almeida ALC, Salemi VMC. Accuracy of mitral annular plane systolic excursion in diagnosing anthracycline-induced subclinical cardiotoxicity in patients with breast cancer - a retrospective cohort study. *Cardiooncology* 2024; **10**(1): 76.

93.   Tafarshiku R, Henein MY, Berisha-Muharremi V, et al. Left Ventricular Diastolic and Systolic Functions in Patients with Hypothyroidism. *Medicina (Kaunas)* 2020; **56**(10).

94.   Udovcic M, Pena RH, Patham B, Tabatabai L, Kansara A. Hypothyroidism and the Heart. *Methodist Debakey Cardiovasc J* 2017; **13**(2): 55-9.

95.   Berger SG, Witczak BN, Reiseter S, et al. Cardiac dysfunction in mixed connective tissue disease: a nationwide observational study. *Rheumatol Int* 2023; **43**(6): 1055-65.

96.   Romano S, Judd RM, Kim RJ, et al. Left Ventricular Long-Axis Function Assessed with Cardiac Cine MR Imaging Is an Independent Predictor of All-Cause Mortality in Patients with Reduced Ejection Fraction: A Multicenter Study. *Radiology* 2018; **286**(2): 452-60.

97.   Svealv BG, Olofsson EL, Andersson B. Ventricular long-axis function is of major importance for long-term survival in patients with heart failure. *Heart* 2008; **94**(3): 284-9.

98.   Willenheimer R, Cline C, Erhardt L, Israelsson B. Left ventricular atrioventricular plane displacement: an echocardiographic technique for rapid assessment of prognosis in heart failure. *Heart* 1997; **78**(3): 230-6.

99.   Tavazzi G, Colombo CNJ, Klersy C, et al. Echocardiographic parameters for weaning from Extracorporeal Membrane Oxygenation - the role of longitudinal function and cardiac time intervals. *Eur Heart J Cardiovasc Imaging* 2024.

100.   Mayr A, Pamminger M, Reindl M, et al. Mitral annular plane systolic excursion by cardiac MR is an easy tool for optimized prognosis assessment in ST-elevation myocardial infarction. *Eur Radiol* 2020; **30**(1): 620-9.

101.   Wang L, Yuan W, Huang X, Zhao X, Zhao X. Cardiac magnetic resonance-derived mitral annular plane systolic excursion: a robust indicator for risk stratification after myocardial infarction. *Int J Cardiovasc Imaging* 2024; **40**(4): 897-906.

102.   Brand B, Rydberg E, Ericsson G, Gudmundsson P, Willenheimer R. Prognostication and risk stratification by assessment of left atrioventricular plane displacement in patients with myocardial infarction. *Int J Cardiol* 2002; **83**(1): 35-41.

103.   Yan C, Chang Y, FangWu, et al. Evaluation of the prognostic value of lateral MAPSE in patients with suspected coronary artery disease. *Int J Cardiol Heart Vasc* 2025; **56**: 101567.

104.   Holzknecht M, Reindl M, Tiller C, et al. Global longitudinal strain improves risk assessment after ST-segment elevation myocardial infarction: a comparative prognostic evaluation of left ventricular functional parameters. *Clin Res Cardiol* 2021; **110**(10): 1599-611.

105.   Rydberg E, Erhardt L, Brand B, Willenheimer R. Left atrioventricular plane displacement determined by echocardiography: a clinically useful, independent predictor of mortality in patients with stable coronary artery disease. *J Intern Med* 2003; **254**(5): 479-85.

106.   El-Battrawy I, Ansari U, Lang S, et al. Risk stratification in Takotsubo syndrome: a role of mitral annular plane systolic excursion. *QJM* 2018; **111**(4): 231-6.

107.   Romano S, Judd RM, Kim RJ, et al. Prognostic Implications of Mitral Annular Plane Systolic Excursion in Patients with Hypertension and a Clinical Indication for Cardiac Magnetic Resonance Imaging: A Multicenter Study. *JACC Cardiovasc Imaging* 2019; **12**(9): 1769-79.

108.   Ballo P, Barone D, Bocelli A, Motto A, Mondillo S. Left ventricular longitudinal systolic dysfunction is an independent marker of cardiovascular risk in patients with hypertension. *Am J Hypertens* 2008; **21**(9): 1047-54.

109.   Siepen FAD, Bauer R, Voss A, et al. Predictors of survival stratification in patients with wild-type cardiac amyloidosis. *Clin Res Cardiol* 2018; **107**(2): 158-69.

110.   Ochs MM, Fritz T, Arenja N, et al. Regional differences in prognostic value of cardiac valve plane displacement in systemic light-chain amyloidosis. *J Cardiovasc Magn Reson* 2017; **19**(1): 87.

111.   Riffel JH, Mereles D, Emami M, et al. Prognostic significance of semiautomatic quantification of left ventricular long axis shortening in systemic light-chain amyloidosis. *Amyloid* 2015; **22**(1): 45-53.

112.   Liu D, Hu K, Stork S, et al. Predictive value of assessing diastolic strain rate on survival in cardiac amyloidosis patients with preserved ejection fraction. *PLoS One* 2014; **9**(12): e115910.

113.   Lindholm A, Kjellstrom B, Seemann F, et al. Atrioventricular plane displacement and regional function to predict outcome in pulmonary arterial hypertension. *Int J Cardiovasc Imaging* 2022; **38**(10): 2235-48.

114. Havaldar AA. Evaluation of sepsis induced cardiac dysfunction as a predictor of mortality. *Cardiovasc Ultrasound* 2018; **16**(1): 31.

115. Zou T, Yin W, Li Y, et al. Hemodynamics in Shock Patients Assessed by Critical Care Ultrasound and Its Relationship to Outcome: A Prospective Study. *Biomed Res Int* 2020; **2020**: 5175393.

116. Blixt PJ, Nguyen M, Cholley B, et al. Association between left ventricular systolic function parameters and myocardial injury, organ failure and mortality in patients with septic shock. *Ann Intensive Care* 2024; **14**(1): 12.

117. Yin W, Li Y, Zeng X, et al. The utilization of critical care ultrasound to assess hemodynamics and lung pathology on ICU admission and the potential for predicting outcome. *PLoS One* 2017; **12**(8): e0182881.

118. Rydberg E, Arlbrandt M, Gudmundsson P, Erhardt L, Willenheimer R. Left atrioventricular plane displacement predicts cardiac mortality in patients with chronic atrial fibrillation. *Int J Cardiol* 2003; **91**(1): 1-7.

119. Matos JD, Balachandran I, Heidinger BH, et al. Mitral annular plane systolic excursion and tricuspid annular plane systolic excursion for risk stratification of acute pulmonary embolism. *Echocardiography* 2020; **37**(7): 1008-13.

120. Jarori U, Maatman TK, Maatman B, Mastouri R, Sawada SG, Khemka A. Mitral Annular Plane Systolic Excursion: An Early Marker of Mortality in Severe COVID-19. *J Am Soc Echocardiogr* 2020; **33**(11): 1411-3.

121. Hensel KO, Roskopf M, Wilke L, Heusch A. Intraobserver and interobserver reproducibility of M-mode and B-mode acquired mitral annular plane systolic excursion (MAPSE) and its dependency on echocardiographic image quality in children. *PLoS One* 2018; **13**(5): e0196614.

122. Marwick TH. Ejection Fraction Pros and Cons: JACC State-of-the-Art Review. *J Am Coll Cardiol* 2018; **72**(19): 2360-79.

123. Schick AL, Kaine JC, Al-Sadhan NA, et al. Focused cardiac ultrasound with mitral annular plane systolic excursion (MAPSE) detection of left ventricular dysfunction. *Am J Emerg Med* 2023; **68**: 52-8.

124. Sepulveda-Martinez A, Steding-Ehrenborg K, Rodriguez-Lopez M, et al. Atrioventricular plane displacement versus mitral and tricuspid annular plane systolic excursion: A comparison between cardiac magnetic resonance and M-mode echocardiography. *Clin Physiol Funct Imaging* 2021; **41**(3): 262-70.

125. Nishikage T, Yamamoto H, Fukumoto N, et al. Significant dependency of left atrial strain on left ventricular longitudinal motion. *J Echocardiogr* 2023; **21**(4): 149-56.

126. Yu J, Tasken AA, Flade HM, et al. Automatic assessment of left ventricular function for hemodynamic monitoring using artificial intelligence and transesophageal echocardiography. *J Clin Monit Comput* 2024; **38**(2): 281-91.

127. Tasken AA, Yu J, Berg EAR, et al. Automatic Detection and Tracking of Anatomical Landmarks in Transesophageal Echocardiography for Quantification of Left Ventricular Function. *Ultrasound Med Biol* 2024; **50**(6): 797-804.

128. Stoylen A, Skjaerpe T. Systolic long axis function of the left ventricle. Global and regional information. *Scand Cardiovasc J* 2003; **37**(5): 253-8.

129. Jones CJ, Raposo L, Gibson DG. Functional importance of the long axis dynamics of the human left ventricle. *Br Heart J* 1990; **63**(4): 215-20.

130. Goodreads. Leonardo da Vinci > Quotes > Quotable Quote. 2025. https://www.goodreads.com/quotes/9010638-simplicity-is-the-ultimate-sophistication-when-once-you-have-tasted (accessed 14/02/2025 2025).

Central graphical abstract: Clinical utility of MAPSE

Figure 1. Examples of MAPSE measurement using M-mode echocardiograph in a person with (A) normal systolic function and (b) impaired systolic function. Reproduced from Hu et al[9], 2012.

Abbreviation: MAPSE, mitral annular plane systolic excursion

Figure 2. Example of MAPSE measurement using M-mode CMR. MAPSE measurement using CMR on 4-chamber cine view from lateral and septal annulus. Adapted from Mayr et al[100], 2020.

Abbreviation: MAPSE, mitral annular plane systolic excursion

Figure 3. Preferred Reporting Items for Systematic Reviews and Meta-Analyses flow chart outlining study selection.

Figure 4: Graph demonstrating individual study and meta-correlation between MAPSE and LVEF.

The size of the blue individual circles denotes the population size. The red diamond denotes the meta-correlation and 95% confidence interval.
Abbreviations: LVEF, ejection fraction, **MAPSE**, mitral annular plane systolic excursion

Figure 5: Summary of individual studies and meta-correlation between MAPSE and the absolute value of GLS.

The size of the blue individual circles denotes the population size. The red diamond denotes the meta-correlation and 95% confidence interval.
Abbreviations: GLS, global longitudinal strain; MAPSE, mitral annular plane systolic excursion

Table 1. MAPSE normal values proposed from observational studies on healthy cohorts. (Adapted from Wang 2023[18] and Grue 2019[34]).

		Grue, 2019, n=1157	
		Lower 95% limit of normal (mm)	Upper 95% limit of normal (mm)
Females	<40 years	11.6	18.8
	40-59 years	10.4	18.2
	>60 years	8.3	16.3
Males	<40 years	11.4	18.8
	40-59 years	9.7	18.2
	>60 years	9.0	17.4
		Wang, 2023, n=1952	
Females	Septal M-mode	9	19
	Lateral M-mode	10	21
Males	Septal M-mode	9	19
	Lateral M-mode	10	20

Table 2. Summary of studies comparing MAPSE with other measures of systolic function.								
First author's last name	Year	Cohort	N with pathology	N healthy control	Modality	Measurement (correlation with biplane LVEF unless stated)	Key findings*	
							r	Optimal cut off
Correlation between MAPSE and LVEF								
Simonson[39]	1989	Mixed cardiac pathologies	74	26	TTE	Midpoint A4C	0.84	<8mm -> LVEF<50%: sens: 98%; spec: 82%
Alam[33]	1991	Post-ACS	37		TTE	Mean of lateral, septal, anterior, inferior A2C and A4C (vs radionucleotide angiography LVEF)	0.87	≥10mm -> LVEF≥50% sens 95%; spec 82%
Alam[32]	1992	Post-ACS or HFrEF	34	45	TTE	Mean lateral, septal, anterior, inferior A2C and A4C	0.96	
Qin[40]	2004	Mixed cardiac pathologies	47		3D TTE/ CMR	Mean of 18 points on mitral annulus (vs CMR LVEF)	0.84	<12mm -> LVEF <50% Sens: 95% spec:85%
Elnoamany[41]	2006	Multiple cardiac pathologies	133	20	TTE	Mean of lateral, septal, anterior, inferior A2C and A4C	0.81	<12mm -> LVEF<50% sens: 90%; spec 88%
Bazaz[43]	2008	Mixed cohort	75		TTE	Lateral A4C	0.87	
Tsang[42]	2010	Multiple cardiac pathologies	60		TTE/CMR	Midpoint on 4AC (vs CMR LVEF)	0.77	
Matos[49]	2012	Mixed cohort	300		TTE	Mean of lateral and septal A4C	Males 0.64 Females 0.47	<6mm -> LVEF <30% sens: 92%; spec 85%
Zidan[45]	2016	Obese ventilated patients	60		TTE	Mean of lateral, septal, anterior, inferior A2C and A4C	0.87	
Lopez-Candelas[50]	2017	Mixed cohort	100		TTE	Lateral 4AC	0.6	
Magdy[54]	2018	Ischaemic cardiomyopathy LVEF<35%	50		Dobutamine stress TTE	Mean of lateral and septal A4C	Baseline 0.28 Low dose 0.35	Δ ≥2 mm -> contractile reserve >10% sens 68%; spec 59%
Borde[53]	2018	Patients undergoing CABG	116		TOE	Mean of lateral and septal A4C		<9mm -> LVEF<52% sens 90%; spec: 92%

Hernandez Burgos[51]	2018	Mixed cohort	150		TTE	Lateral A4C	0.46	
Shah[46]	2019	Ventilated patients	39		TTE	Lateral point 4AC	0.61	>12.5mm -> LVEF >50% sens: 82.1%; spec: 91.7%
Stenberg[52]	2021	POCUS pre-elective operations	100		TTE	Mean of lateral and septal A4C	0.56	<12mm -> LVEF<50% sens: 95%; spec: 80%
Ozer[17]	2022	Consecutive outpatients	512		TTE	Lateral point 4AC	0.34	
Schick[123]	2023	POCUS in emergency dept	61		TTE	Lateral A4C		≤8mm -> LVEF<50% sens: 42%; spec: 89%
Bergenzaun[47]	2013	ICU admissions septic shock	50		TTE	Mean lateral, septal, anterior, inferior A2C and A4C	0.59	
Brault[48]	2023	Ventilated patients with septic shock	71		TTE	Lateral A4C	0.52	
Turan[44]	2024	Duchenne Muscular Dystrophy	55	54	TTE	Mean lateral and septal A4C	0.71	
Correlation between MAPSE and TDI								
Mondillo[56]	2006	Healthy subjects	56		TTE	Mean of lateral, septal, anterior, inferior A2C and A4C	0.54	
Ballo[55]	2008	Healthy subjects	92		TTE	Lateral point A4C	0.90	
Correlation between MAPSE and GLS								
Luszczak[63]	2013	Moderate-severe AS	82		TTE	Mean of midpoint of A4C, A3C and A2C	−0.56	
Wenzelburger[61]	2014	HFpEF	62	36	TTE	Mean of lateral and septal A4C	Rest: −0.43 Exercise: −0.59	
Liu[57]	2014	HCM	65	48	TTE (speckle tracking)	Mean of lateral, septal, anterior, inferior, anteroseptal and inferolateral A4C A2C and A3C.	−0.86	
Chiu[58]	2016	Hemodialysis	198		TTE (speckle tracking)	Mean of midpoint of A4C, A3C and A2C	−0.61	
Lopez-Candelas[64]	2018	Mixed cohort	75		TTE	Lateral A4C	−0.54	

Teraguchi[62]	2019	Moderate-severe MR	50		TTE (speckle tracking)	Lateral A4C	−0.51	14.6% change in lateral MAPSE -> low GLS Sens: 94%; spec: 94%
Blixt[59]	2021	ICU admissions with septic shock	73		TTE		−0.81	
Wang[18]	2023	Healthy subjects	1683		TTE	Mid-point A4C (speckle tracking)	−0.35	
Zhang[65]	2023	Chemotherapy-related cardiotoxicity	200		TTE	Mean of lateral and septal A4C	Baseline: −0.38 Post-chemo: −0.53	Reduction in MAPSE ≥2mm -> 15% drop in GLS Sens: 74.5%; spec 54.9%
Brault[48]	2023	Ventilated patients with septic shock	71		TTE	Septal A4C	−0.64	
Ji[60]	2024	Post-heart transplant	155	49	TTE	Midpoint A4C	−0.71	9.1mm associated with mortality/ hospitalisation Sens: 94%; spec 58%

All results are statistically significant with a *p* value <0.05 unless stated otherwise. Where multiple results are provided (e.g. from multiple measurement points) the strongest correlation is given.

Abbreviations: A2C, apical 2 chamber; A3C, apical 3 chamber; A4C, apical 4 chamber; ACS, acute coronary syndrome; AUC, area under curve; CABG, coronary artery bypass graft; CMR, cardiac magnetic resonance; LVEF, ejection fraction; HCM, hypertrophic cardiomyopathy; HFpEF, heart failure with preserved ejection fraction; HFrEF, heart failure with reduced ejection fraction; MR, mitral regurgitation; POCUS, point of care ultrasound; sens, sensitivity; spec, specificity; TDI, tissue Doppler; TOE, transesophageal echocardiography; TTE, transthoracic echocardiogram; VTI, velocity time integral.

Table 3. Summary of studies using MAPSE to diagnose left ventricular systolic impairment when LVEF is normal									
							Key findings		
First author's last name	Year	Cohort	N with pathology	N control	Modality	Measurements taken	Study subgroup (comparison if not healthy controls)	MAPSE pathology (mm)	MAPSE control (mm)
Xiao[66]	2006	HTN	26	10	TTE	Septal 4AC		13.1±1.5	16.7±2.5
						Lateral 4AC		12.3±2.5	15.3±0.3
Koulouris[67]	2006	HTN	81	50	TTE	Mean lateral, septal, anterior, inferior A2C and A4C		10.3±3	13.1±1
Ballo[68]	2010	DM/HTN	163 with HTN/DM/both	70	TTE	Mean lateral, septal, anterior, inferior A2C and A4C	DM	13.5±2.3	15.1±2.4
							HTN	13.2±2.2	
							DM+HTN	11.5±3	
Loncarevic[70]	2016	DM (no HTN)	210	80	TTE	Mean lateral, septal, anterior, inferior A2C and A4C		16.4±1.6	15.4±2.1
Salas-Pacheco [69]	2021	HTN	226	101	TTE	Mean lateral and septal A4C		13.8±2.2	15.5±2.0
Tromba[71]	2017	Sleeve gastrectomy	45		TTE	Mean lateral and septal A4C	Pre- and post-sleeve gastrectomy	Median 16	Median 18
Li[23]	2023	T2DM	100	50	TTE		DM>15 years vs controls	13.98±1.1	15.6±0.45
Takeda[73]	2001	AS (all grades)	78	10	TTE	Lateral 4AC	Lower in severe AS vs healthy controls		
						Septal 4AC	Lower in mild, moderate and severe AS vs control		
Rydberg[72]	2004	AS (all grades)	49	133	TTE	Mean lateral, septal, anterior, inferior A2C and A4C		9.2±2.3	10.7±2.8
Weidemann[74]	2009	Symptomatic severe AS ± fibrosis on CMR	21 (no fibrosis)	21 (severe fibrosis)	TTE	Septal 4AC	Severe fibrosis vs no fibrosis	4.9±1.0	10.7±1.2

Herrmann[78]	2011	Low-flow low-gradient AS	86		TTE	Lateral and septal A4C	Low-flow low-gradient AS preserved LVEF (vs moderate AS)	5.8±2.2	12.8±1.7
							Low-flow low-gradient AS reduced LVEF (vs moderate AS)	5.0±0.9	
Kempny[75]	2013	Severe AS undergoing TAVI	101		TTE	Lateral A4C	Pre and post-TAVI	9.1±3.2	10.2±3.3
Matos[76]	2017	Severe asymptomatic AS	205		TTE	Mean lateral, septal, anterior, inferior A2C and A4C	Change in MAPSE requiring intervention scan (vs those not requiring intervention)	2mm reduction	<2mm reduction
Liu[79]	2018	Low-flow low-gradient AS	43	20	TTE	Lateral A4C	Low-flow low-gradient AS reduced LVEF (vs pseudosevere AS)	9.8±2.6	11.6±2.2
Dursunoglu[81]	2004	DD	62	40	TTE	Mean lateral, septal, anterior, inferior A2C and A4C		12.4±1.4	15.3±1.4
Gromadzinski[82]	2014	CKD±DD	36	34	TTE	Lateral A4C	CKD with DD vs CKD without DD median(range)	13(11–17)	14(11–16)
Wenzelburger[61]	2014	HFpEF (during exercise)	62	36	TTE	Lateral and septal A4C	At rest	10.9±2.1	12.1±2.2
							During exercise	12±2.2	16.2±2.7
Taşolar[83]	2015	High BMI ±DD	40 high BMI+ HFpEF	40 high BMI	TTE	Mean lateral, septal, anterior, inferior A2C and A4C		11.8±1.6	15.8±1.6
Bytyci[84]	2016	HFpEF	79	29	TTE	Septal		11.8±2.0	14.0±3.0
						Lateral		12.4±3	15.0±3.0
Wierbowska-Drabik[86]	2013	Obesity and LVH	27	27	TTE	Lateral A4C		20±4	22±2

Study	Year	Condition			Modality	View			
Lopez-Candale[85]s	2014	Patients with LVH	45	45	TTE	Lateral A4C		12±2	14±2
Doesch[87]	2015	HCM	88	50	TTE/CMR	Mean lateral and septal A4C		10.0±3.2	13.3±3.8
Kiotsekoglou[88]	2008	Marfan's	66	61	TTE	Anterior A2C		13.8±0.3	18.0±0.2
Panovsky[89]	2021	Duchenne Muscular Dystrophy	43	18	CMR	Lateral and septal A4C	With LGE	10.8±1.6	13.7±2.7
							Without LGE	11.6±1.9	
Harries[90]	2022	Anthracycline-related cardiotoxicity	24		CMR	Lateral A4C	Recovery of function post-anthracycline	11.7±1.5 (poor recovery)	15.7±3.1 (good recovery)
Botelho[92]	2024	Anthracycline-related cardiotoxicity	5 (with cardiotoxicity)	56 (without cardiotoxicity)				13.5 [8.8-16.3] *ns	13.3 [12.0-13.8] *ns
Ichikawa[91]	2024	Anthracycline-related cardiotoxicity	56 (with cardiotoxicity)	200 (without cardiotoxicity)	TTE	Mean lateral and septal A4C		14.0±1.5*ns	13.8±2.6*ns
Tafarshiku[93]	2020	Hypothyroidism	81	22	TTE	Lateral A4C		14.9±0.3	16.6±0.2
Berger[95]	2023	Mixed connective tissue disease	77	59	TTE	Mean lateral and septal A4C		13.7±2.1	15.3±2.3

* All results presented as mean±standard deviation unless stated; all results are statistically significant with a *p* value <0.05 unless stated otherwise

Written results provided where numerical results not available in manuscripts.

Abbreviations: A2C, apical 2 chamber; A3C, apical 3 chamber; A4C, apical 4 chamber; AS, aortic stenosis; BMI, body mass index; CAD, coronary artery disease; CKD, chronic kidney disease; DD, diastolic dysfunction; DM, diabetes mellitus; LVEF, left ventricular ejection fraction; HFpEF, heart failure with preserved ejection fraction; HFrEF, heart failure with reduced ejection fraction; HTN, hypertension; LGE, late gadolinium enhancement; LVH, left ventricular hypertrophy; ns, not significant; TAVI, transcatheter aortic valve implantation; TTE, transthoracic echocardiogram

First author's last name	Year	Cohort	n	Modality	Measurement	Outcome	Key findings*		
							MAPSE HR [95% CI] unless stated	Uni/ Multivariable	Comparison with LVEF or GLS
Rangarajan[16]	2016	General outpatients	400	CMR	Lateral A4C	Death, MI, hospitalisation, revasc	1.344 /2mm decrease	Multi	LVEF not associated
Ozer[17]	2022	General outpatients	512	TTE	Lateral A4C	Death, hospitalisation	Hospitalization: OR 0.12 [0.03–0.42] Death 0.01 [0.00–0.52]	Multi	LVEF not associated
Xue[15]	2023	General outpatients	1572	CMR	Mean lateral, septal, anterior, inferior, anteroseptal, inferoseptal A4C, A2C and A3C	Death, heart failure hospitalisation	Death: 1.6 [1.4–1.9]/mm decrease Hospitalisation: 2.1 [1.7–2.7]/mm decrease	Multi	Death: GLS: 1.3 [1.1–1.5] LVEF HR 1.2 [1.0–1.4] Hospitalisation: GLS: 1.8 [1.4–2.3] LVEF 1.7 [1.3–2.1]
Willenheimer[98]	1997	HFrEF	181	TTE	Mean lateral, septal, anterior, inferior A2C and A4C	Death	No HR		>10mm -> mortality 0% (no comparison with LVEF or GLS)
Svealv[97]	2008	HFrEF	228	TTE	Mean lateral and septal A4C	Death	0.8 [0.66–0.97]	Multi	
Romano[96]	2018	HFrEF	1040	CMR	Lateral A4C	Death	2.1 [1.5–2.8] / mm decrease	Multi	LVEF not associated
Tavazzi[99]	2024	HFrEF	76	TTE	Lateral A4C	Successful wean from ECMO	6.7 [2.7–16.6] (>8.9 vs <8.9mm)	Uni (multi not performed)	LVEF not associated
Brand[102]	2002	Post-ACS	271	TTE	Mean lateral, septal, anterior, inferior A2C and A4C	Death	Relative risk= 3 (lowest quartile vs upper three quartiles)	Multi	
Rydberg[105]	2003	Stable CAD	333	TTE	Mean lateral, septal, anterior, inferior A2C and A4C	Death	OR 1.6 [1.4–2.0]/mm decrease – also associated in multivariable analysis (value not reported)	Uni	LVEF not associated
Mayr[100]	2019	Day 2 post-STEMI	255	CMR	Lateral and septal A4C	Death; MI; stroke, heart failure	6.0 [2.5–14.7] MAPSE <9 vs >9mm.	Multi	LVEF not associated

Table 4. Studies comparing MAPSE with LVEF or GLS as a prognostic marker

Holzknecht[104]	2021	Post-STEMI (day 2-4 post PCI)	445	CMR	Septal point on A4C	Death; MI; heart failure	0.80 [0.72–0.89]	Uni	LVEF 0.95 [0.93–0.98], GLS associated in multivariable analysis: 1.13 [1.01–1.27]
Wang[101]	2024	Post-ACS	165	CMR	Mean lateral and septal A4C	Death; MI; stroke; heart failure; ventricular arrhythmia	0.83 [0.73–0.94]	Multi	LVEF 0.96 [0.94–0.99]
Yan[103]	2024	Suspected CAD	233	CMR	Lateral A4C	Death, MI, hospitalisation, revascularisation	1.4 [1.1–1.8]	Multi	LVEF not associated
El-Battrawy[106]	2018	Takotsubo	53	TTE	Lateral A4C	Death	5.1 [1.3–19.2] MAPSE <10 vs >10mm	Multi	LVEF not associated
Romano[107]	2019	HTN	1735	CMR	Lateral A4C	Death	1.4 [1.3–1.5] / mm decrease	Multi	GLS 1.180 [1.137–1.224) LVEF not associated
Ballo[108]	2008	HTN	156	TTE	Mean lateral, septal, anterior, inferior A2C and A4C	Death; MI; stroke; heart failure; revascularisation	0.81 [0.67–0.98] / 1 standard deviation reduction	Multi	
Dem Siepen[109]	2018	Wildtype-TTR amyloidosis	191	TTE		Death	0.19 [0.04–0.83] MAPSE <8.8 vs >8.8mm	Multi	LVEF and GLS not associated
Liu[112]	2014	AL amyloidosis	41	TTE	Lateral and septal A4C	Death	1.24 [1.02–1.50]	Uni (not associated in multivariable analysis)	GLS associated in multivariable: HR 7.3 [2.08–25.65]
Riffel[111]	2014	AL amyloidosis	120	TTE	Lateral A4C	Death	0.05 [0.018–0.11]	Uni (not associated in multivariable analysis)	Long axis strain associated in multivariable: HR 0.76 [0.63–0.91]
Ochs[110]	2017	AL amyloidosis	68	CMR	Lateral A4C	Death, heart transplant	0.23 [0.10–0.56] /mm increase	Uni (multi not given)	LVEF: 0.95 [0.93–0.98] GLS: 1.12 [1.05–1.19]
Ji[60]	2024	Post-heart transplant	155	TTE	Lateral and septal A4C by speckle tracking	Death, hospitalisation	0.68 [0.56–0.83]	Multi	LVEF not associated GLS: 0.758[0.657–0.874]

Lindholm[113]	2022	Pulmonary hypertension	71	CMR/TTE	Mean lateral, septal, anterior, inferior, anteroseptal, inferoseptal A4C, A2C and A3C	Death, transplantation	1.16 [1.04–1.29] /mm decrease	Multi	LVEF not associated
Bergenzaun[47]	2013	ICU admissions septic shock	50	TTE	Mean lateral, septal, anterior, inferior A2C and A4C	28-day mortality	OR 0.7 [0.5–0.9]	Multi	LVEF not associated
Havaldar[114]	2018	ICU admissions septic shock	48	TTE	Lateral A4C	Death in ICU	OR 0.02 [0.00–0.69]	Multi	LVEF not associated
Blixt[116]	2024	ICU admissions septic shock	152	TTE	Mean lateral and septal A4C	Myocardial injury	OR 0.84 [0.70–0.99]	Multi	LVEF not associated
Rydberg[105]	2003	Permanent A fib	160	TTE	Mean lateral, septal, anterior, inferior A2C and A4C	Death	OR 1.4 [1.1–1.7]	Multi	LVEF not associated
Matos[119]	2020	Acute PE	362	TTE	Mean lateral, septal, anterior, inferior A2C and A4C	Death, need for thrombolysis or pressors	OR 3.5 [1.4–9.0] MAPSE <11vs >11mm with preserved LVEF	Multi	LVEF <50%, OR 4.1 [1.4–12.0]
Jarori[120]	2020	Severe COVID-19	68	TTE	Lateral and septal A4C	Death	1.6 [1.2–2.2] MAPSE <10 vs >10mm	Multi	LVEF not associated

* All results presented are statistically significant with a p value <0.05 unless stated otherwise

Abbreviations: : A2C, apical 2 chamber; A3C, apical 3 chamber; A4C, apical 4 chamber; ACS, acute coronary syndrome; A fib: atrial fibrillation; CAD, coronary artery disease CMR, cardiac magnetic resonance; COVID-19, coronavirus disease-2019; LVEF, left ventricular ejection fraction; GLS, global longitudinal strain; HR, hazard ratio; HCM, hypertrophic cardiomyopathy; HFrEF, heart failure with reduced ejection fraction; HR, hazard ratio; ICU, intensive care unit; MACE, major adverse cardiac events; MAPSE, mitral annulus systolic plane excursion; OR, odds ratio; PE, pulmonary embolism; STEMI, ST elevation myocardial infarction; TTE, transthoracic echocardiogram; TTR, transthyretin