Monitoring Report: GLP-1 RA Prescribing Trends - December 2024 Data

Samuel Gratzl, PhD*

Brianna M Goodwin Cartwright, MS*

Patricia J Rodriguez, PhD, MPH

Karen Gilbert, PhD

Duy Do, PhD

Nina Masters, MPH, PhD

Nicholas L Stucky, MD, PhD

*Denotes co-first author

Affiliations: All authors are affiliated with Truveta, Inc.

Corresponding author:
Dr Nicholas Stucky
Truveta, Inc
1745 114th Ave SE, Bellevue, WA 98004
Email: nicholass@truveta.com

Abstract

Background: Limited recent data exist on prescribing patterns and patient characteristics for glucagon-like peptide 1 receptor agonists (GLP-1 RAs), an important drug class used as anti-diabetic medication (ADM) for patients with type 2 diabetes mellitus (T2D) and/or anti-obesity medication (AOM) in patients with overweight or obesity.

For brevity, we use the term GLP-1 RA to refer to both GLP-1 RA and dual GLP-1 RA/GIP medications.

Objective: To describe recent trends in prescribing and dispensing of GLP-1-based medications in the US.

Methods: Using a subset of real-world electronic health record (EHR) data from Truveta, a growing collective of health systems that provide more than 18% of all daily clinical care in the US, we identified people who were prescribed a GLP-1-based medication between January 01, 2018 and December 31, 2024. We describe prescribing volumes and patient characteristics over time, by medication, and by FDA-labeled use. Among the subset of patients for whom post-prescription dispensing data is available, we describe the proportion and characteristics of patients who were and were not dispensed a GLP-1 RA following their prescription.

Results: 1,828,203 patients were prescribed a GLP-1 RA between January 2018 and December 2024, with 8,412,584 total prescriptions during this period. Across the full study period, among first-time prescriptions for which use could be established, ADMs accounted for 77.2% and AOMs accounted for 22.8%. The most common first-time medication was semaglutide (n = 1,016,220). During December 2024, overall AOM prescribing was highest for tirzepatide (+21.95%). Among those with available data, 67.3% of first-time ADM prescriptions overall and 69.0% in September 2024 had a fill within 60 days. By comparison, 43.6% of first-time AOM prescriptions overall and 48.7% in September 2024 had a fill within 60 days.

Trends in Prescribing

	GLP-1 RA	Semaglutide	Tirzepatide	Liraglutide	Dulaglutide	Other
Overall						
September 2024	5.14%	2.85%	1.85%	0.07%	0.44%	0.03%
December 2024	5.24%	2.71%	2.10%	0.07%	0.42%	0.03%
Change (percentage)	+1.89%	-4.83%	+13.48%	-6.21%	-3.88%	-7.86%
Difference (percentage points)	+0.10%	-0.14%	+0.25%	-0.00%	-0.02%	-0.00%
ADM						
September 2024	2.83%	1.39%	0.99%	0.01%	0.44%	0.03%
December 2024	2.81%	1.34%	1.04%	0.01%	0.42%	0.03%
Change (percentage)	-0.88%	-3.92%	+4.49%	-14.40%	-3.88%	-7.86%
Difference (percentage points)	-0.02%	-0.05%	+0.04%	-0.00%	-0.02%	-0.00%
AOM						
September 2024	1.36%	0.75%	0.61%	0.02%	n/a	n/a
December 2024	1.46%	0.71%	0.75%	0.02%	n/a	n/a
Change (percentage)	+6.98%	-4.87%	+21.95%	-9.67%		
Difference (percentage points)	+0.10%	-0.04%	+0.13%	-0.00%		
Unknown						

	GLP-1 RA	Semaglutide	Tirzepatide	Liraglutide	Dulaglutide	Other
September 2024	1.19%	0.81%	0.34%	0.05%	n/a	n/a
December 2024	1.23%	0.75%	0.44%	0.04%	n/a	n/a
Change (percentage)	+2.89%	-7.17%	+27.58%	-3.69%		
Difference (percentage points)	+0.03%	-0.06%	+0.09%	-0.00%		

- Overall prescribing rates (GLP-1 RA prescriptions per total prescriptions) in December 2024 were similar to September 2024 (+1.89%).
- ADM prescribing rates in December 2024 were similar to those in September 2024 (-0.88%).
- Among ADMs, semaglutide continues to be the most prescribed medication (1.34% of all prescriptions), however prescription rates decreased (-3.92%)
- AOM prescribing increased slightly in December 2024 compared to September 2024 (+6.98%).
- Amongst AOMs, tirzepatide was the most prescribed medication (a rate increase of 21.95%).

Trends in First-Time Prescribing

	GLP-1 RA	Semaglutide	Tirzepatide	Liraglutide	Dulaglutide	Other
Overall						
September 2024	1.23%	0.74%	0.40%	0.01%	0.07%	0.01%
December 2024	1.10%	0.62%	0.41%	0.01%	0.05%	0.01%
Change (percentage)	-10.25%	-16.13%	+2.06%	-11.98%	-18.17%	-14.02%
Difference (percentage points)	-0.13%	-0.12%	+0.01%	-0.00%	-0.01%	-0.00%
ADM						
September 2024	0.51%	0.28%	0.16%	0.00%	0.07%	0.01%
December 2024	0.44%	0.23%	0.15%	0.00%	0.05%	0.01%
Change (percentage)	-13.81%	-16.44%	-7.21%	-24.41%	-18.17%	-14.02%
Difference (percentage points)	-0.07%	-0.05%	-0.01%	-0.00%	-0.01%	-0.00%
AOM						
September 2024	0.38%	0.21%	0.17%	0.00%	n/a	n/a
December 2024	0.35%	0.17%	0.18%	0.00%	n/a	n/a
Change (percentage)	-8.10%	-19.92%	+6.63%	-4.87%		
Difference (percentage points)	-0.03%	-0.04%	+0.01%	-0.00%		
Unknown						
September 2024	0.34%	0.26%	0.08%	0.01%	n/a	n/a
December 2024	0.32%	0.22%	0.09%	0.01%	n/a	n/a
Change (percentage)	-6.56%	-12.65%	+12.86%	-7.50%		
Difference (percentage points)	-0.02%	-0.03%	+0.01%	-0.00%		

- First-time prescribing rates (first-time GLP-1 RA prescriptions per total prescriptions) in December 2024 decreased slightly compared to rates in September 2024 (-10.25%).
- First-time prescribing of ADMs decreased in December 2024 relative to September 2024 (-13.81%).
- First-time prescribing of AOMs decreased slightly in December 2024 relative to September 2024 (-8.10%).

Trends in Dispensing

Changes in GLP-1 RA Initiation: Proportion of First-Time GLP-1 RA Prescriptions Dispensed within 60 days between May 2024 and September 2024

	GLP-1 RA	Semaglutide	Tirzepatide	Liraglutide	Dulaglutide	Other
Overall						
June 2024	55.69%	53.80%	56.87%	54.21%	72.79%	57.52%
September 2024	54.94%	51.94%	57.19%	59.26%	74.36%	60.58%
Change (percentage)	-1.34%	-3.46%	+0.56%	+9.32%	+2.17%	+5.32%
Difference (percentage points)	-0.74%	-1.86%	+0.32%	+5.05%	+1.58%	+3.06%
ADM						
June 2024	68.25%	67.48%	68.75%	61.06%	72.79%	57.52%
September 2024	68.99%	67.43%	69.65%	79.35%	74.36%	60.58%
Change (percentage)	+1.08%	-0.06%	+1.31%	+29.96%	+2.17%	+5.32%
Difference (percentage points)	+0.74%	-0.04%	+0.90%	+18.29%	+1.58%	+3.06%
AOM						
June 2024	48.13%	46.54%	51.50%	40.31%	n/a	n/a
September 2024	48.69%	44.66%	53.76%	43.56%	n/a	n/a
Change (percentage)	+1.16%	-4.04%	+4.38%	+8.07%		
Difference (percentage points)	+0.56%	-1.88%	+2.26%	+3.25%		
Unknown						
June 2024	43.47%	44.09%	40.27%	52.30%	n/a	n/a
September 2024	41.02%	41.46%	38.41%	56.76%	n/a	n/a
Change (percentage)	-5.63%	-5.97%	-4.61%	+8.52%		
Difference (percentage points)	-2.45%	-2.63%	-1.86%	+4.46%		

- During the full time period (2018 – 2024), 69.4% of GLP-1 RA prescriptions had a dispense within 60 days of their prescription (e.g., were initiated within 60 days).
- For patients first prescribed an ADM in September 2024, 68.99% filled a GLP-1 RA within 60 days. Initiation rates for ADM GLP-1 RA in September 2024 were similar to June 2024 (+1.08%).
- For patients first prescribed an AOM in September 2024, 48.69% filled a GLP-1 RA within 60 days. Initiation rates for AOM GLP-1 RA in September 2024 were similar to June 2024 (+1.16%).

About this report

This report provides recent trends in the prescribing and dispensing of glucagon-like peptide-1 receptor agonist (GLP-1 RAs)-based medications, including dual GLP-1 RA/gastric inhibitory polypeptide (GIP) over time, by medication, and by FDA-labeled use. We used near real-time, aggregated electronic health record (EHR) data from a collective of US healthcare systems to describe prescribing and dispensing of GLP-1 RA medications from January 2018 to December 2024.

Introduction

Obesity and type 2 diabetes mellitus (T2D) are interconnected epidemics with a high prevalence and individual and combined negative impacts on health (CDC, 2020, Hales, 2020). Glucagon-like peptide-1 receptor agonist (GLP-1 RAs) -based medications, including dual GLP-1 RA/gastric inhibitory polypeptide (GIP), have been shown to improve glycemic control in patients with T2D and reduce weight in patients with and without T2D (Davies, 2015, Davies, 2021, Garvey, 2023, Jastreboff, 2022, Wilding, 2023, Yeh, 2023). GLP-1 RA have also demonstrated cardiovascular benefit in patients with and without T2D (Lincoff, 2023, Ussher, 2023).

Interest in these medications has recently accelerated, largely for their weight-loss effects; however, limited empiric data exist on prescribing, dispensing, and patient characteristics. Access to and use of GLP-1 RA may be impacted by high cost, limited insurance coverage for patients without T2D and medication shortages.

Methods

Data

We evaluated prescribing and dispensing of GLP-1 RAs between January 2018 and December 2024 using a subset of Truveta Data. Truveta provides access to continuously updated, linked, and de-identified electronic health record (EHR) from a collective of US health care systems that provide 18% of all daily clinical care in the US, including structured information on demographics, encounters, diagnoses, vital signs (e.g., weight, BMI, blood pressure), medication requests (prescriptions), medication administration, laboratory and diagnostic tests and results (e.g., HbA1c tests and values), and procedures. Updated EHR data are provided daily to Truveta by member health care systems. In addition to EHR data for care delivered within Truveta member health care systems, medication dispensing and social drivers of health (SDOH)

information are made available through linked third-party data. Medication dispense (via e-prescribing data) includes fills for prescriptions written both within and outside Truveta member health care systems, providing greater observability into patients' medication history. Medication dispense histories are updated at the time of the encounter and include fill dates, NDC or RxNorm codes, quantity dispensed, and days of medication supplied.

Population

We identified GLP-1 RA prescriptions (medication requests) and dispenses among adults using RxNorm and NDC codes for dulaglutide, exenatide, liraglutide, lixisenatide, semaglutide, and tirzepatide. Exenatide and lixisenatide were grouped as 'other' given small relative sizes. Brands were identified when possible using RxNorm and NDC codes. Brands labeled for treatment of T2D were classified as ADMs and brands labeled for treatment of overweight and obesity were classified as AOMs. When brand could not be established, the labeled use was classified as unknown.

Prescribing Analysis

Characteristics of patients ever prescribed a GLP-1 RA were summarized, including demographics, social drivers of health (SDOH), and presence of T2D and/or overweight or obesity. Characteristics are provided by medication and by year of first prescription.

Overall and first-time prescribing rates of GLP-1 RA were summarized over time by medication and labeled use. The prescribing rate was calculated monthly as the number of patients with a GLP-1 RA prescription divided by the number of patients with any prescription in the month. Absolute differences and percentage changes in the rate of GLP-1 RA prescribing over the most recent 3-month period were summarized overall and by medication and labeled use.

Dispensing Analysis

We describe the proportion of GLP-1 RA prescriptions (overall and first-time) with a dispense occurring within 60 days. Dispensing analyses were restricted to prescriptions for which complete 60-day dispensing outcomes could be observed. Dispensing outcomes were considered observable if (1) a dispense was recorded in the 60 days following the prescription, and/or (2) an encounter took place after the 60-day period ended (at which time any dispenses would be observed). To allow sufficient follow-up, we report dispensing with a 90-day delay. Given shortages and expected substitutions at the pharmacy, we considered the dispense of any GLP-1 RA , which could differ from the prescribed GLP-1 RA.

We plot the proportion of prescriptions (overall and first-time) dispensed within 60 days over time, by medication and use. Differences in dispensing were expected between ADMs and AOMs given ADMs are typically covered by insurance, whereas insurance coverage for AOMs is limited. As such, we provide dispensing rates stratified by FDA labeled use. Further, we stratify dispensing by age 65+ vs. 18-64 as a proxy for Medicare coverage. Medicare never provides coverage for AOMs, whereas commercial insurers may.

Results

Overall population

The demographics of patients by first-prescribed medication are as follows:

Table 1: Patient Characteristics by First-Prescribed GLP-1 RA Medication.

	Semaglutide	Tirzepatide	Liraglutide	Dulaglutide	Other	Overall
N	1,016,220 (100.00%)	274,748 (100.00%)	156,470 (100.00%)	305,865 (100.00%)	41,416 (100.00%)	1,794,719 (100.00%)
Age Group						
18-44	266,209 (26.20%)	87,293 (31.77%)	42,141 (26.93%)	43,767 (14.31%)	4,759 (11.49%)	444,169 (24.75%)
45-64	497,115 (48.92%)	139,659 (50.83%)	77,805 (49.73%)	154,447 (50.50%)	19,580 (47.28%)	888,606 (49.51%)
65+	252,896 (24.89%)	47,796 (17.40%)	36,524 (23.34%)	107,651 (35.20%)	17,077 (41.23%)	461,944 (25.74%)
Sex						
Female	662,810 (65.22%)	188,572 (68.63%)	105,628 (67.51%)	165,028 (53.95%)	22,304 (53.85%)	1,144,342 (63.76%)
Male	353,410 (34.78%)	86,176 (31.37%)	50,842 (32.49%)	140,837 (46.05%)	19,112 (46.15%)	650,377 (36.24%)
Race						
White	708,632 (69.73%)	199,080 (72.46%)	110,531 (70.64%)	199,185 (65.12%)	29,467 (71.15%)	1,246,895 (69.48%)
Black or African American	127,229 (12.52%)	30,303 (11.03%)	19,733 (12.61%)	42,460 (13.88%)	4,238 (10.23%)	223,963 (12.48%)
Asian	24,565 (2.42%)	5,472 (1.99%)	2,740 (1.75%)	7,625 (2.49%)	974 (2.35%)	41,376 (2.31%)
Unknown	104,137 (10.25%)	26,575 (9.67%)	15,533 (9.93%)	38,818 (12.69%)	4,700 (11.35%)	189,763 (10.57%)
Native Hawaiian or Other Pacific Islander	3,388 (0.33%)	743 (0.27%)	619 (0.40%)	1,176 (0.38%)	157 (0.38%)	6,083 (0.34%)
American Indian or Alaska Native	6,984 (0.69%)	1,479 (0.54%)	1,528 (0.98%)	2,153 (0.70%)	350 (0.85%)	12,494 (0.70%)
Other Race	41,285 (4.06%)	11,096 (4.04%)	5,786 (3.70%)	14,448 (4.72%)	1,530 (3.69%)	74,145 (4.13%)
Ethnicity						

	Semaglutide	Tirzepatide	Liraglutide	Dulaglutide	Other	Overall
Hispanic or Latino	122,787 (12.08%)	32,953 (11.99%)	17,709 (11.32%)	42,131 (13.77%)	4,999 (12.07%)	220,579 (12.29%)
Not Hispanic or Latino	837,380 (82.40%)	226,511 (82.44%)	129,160 (82.55%)	244,030 (79.78%)	33,478 (80.83%)	1,470,559 (81.94%)
Unknown	56,053 (5.52%)	15,284 (5.56%)	9,601 (6.14%)	19,704 (6.44%)	2,939 (7.10%)	103,581 (5.77%)
Characteristics						
Type II Diabetes	451,627 (44.44%)	93,162 (33.91%)	73,494 (46.97%)	218,191 (71.34%)	29,054 (70.15%)	865,528 (48.23%)
Overweight or Obesity	723,519 (71.20%)	199,832 (72.73%)	98,509 (62.96%)	166,117 (54.31%)	21,189 (51.16%)	1,209,166 (67.37%)
Labeled Use						
ADM	488,317 (48.05%)	152,388 (55.46%)	59,486 (38.02%)	305,865 (100.00%)	41,416 (100.00%)	1,047,472 (58.36%)
AOM	188,782 (18.58%)	72,134 (26.25%)	47,816 (30.56%)	n/a	n/a	308,732 (17.20%)
Unknown	339,121 (33.37%)	50,226 (18.28%)	49,168 (31.42%)	n/a	n/a	438,515 (24.43%)
Year						
2018	3,673 (0.36%)	n/a	29,346 (18.76%)	22,584 (7.38%)	9,386 (22.66%)	64,989 (3.62%)
2019	18,181 (1.79%)	n/a	26,940 (17.22%)	29,229 (9.56%)	7,696 (18.58%)	82,046 (4.57%)
2020	31,872 (3.14%)	n/a	19,662 (12.57%)	31,664 (10.35%)	6,069 (14.65%)	89,267 (4.97%)
2021	71,149 (7.00%)	n/a	23,039 (14.72%)	47,636 (15.57%)	4,921 (11.88%)	146,745 (8.18%)
2022	163,372 (16.08%)	24,616 (8.96%)	25,420 (16.25%)	67,715 (22.14%)	4,978 (12.02%)	286,101 (15.94%)
2023	361,903 (35.61%)	68,911 (25.08%)	22,785 (14.56%)	69,386 (22.69%)	4,701 (11.35%)	527,686 (29.40%)
2024	366,070 (36.02%)	181,221 (65.96%)	9,278 (5.93%)	37,651 (12.31%)	3,665 (8.85%)	597,885 (33.31%)
Comorbidity						
Asthma	147,090 (14.47%)	39,728 (14.46%)	22,559 (14.42%)	35,763 (11.69%)	4,510 (10.89%)	249,650 (13.91%)
CKD	127,498 (12.55%)	23,765 (8.65%)	20,493 (13.10%)	55,675 (18.20%)	7,475 (18.05%)	234,906 (13.09%)
COPD	50,761 (5.00%)	10,075 (3.67%)	9,014 (5.76%)	21,473 (7.02%)	3,085 (7.45%)	94,408 (5.26%)
Hyperlipidemia	533,110 (52.46%)	129,525 (47.14%)	70,483 (45.05%)	168,549 (55.11%)	22,741 (54.91%)	924,408 (51.51%)
Hypertension	549,879 (54.11%)	131,431 (47.84%)	80,292 (51.31%)	182,017 (59.51%)	24,660 (59.54%)	968,279 (53.95%)
Major Depression	127,887 (12.58%)	31,772 (11.56%)	19,888 (12.71%)	29,392 (9.61%)	3,856 (9.31%)	212,795 (11.86%)

Prescribing over time

Overall

We describe the monthly rate of prescriptions by medication and FDA labeled use over time:

First-time

We describe the monthly rate of first-time prescriptions by medication and FDA labeled use over time.

Characteristics of patients by year of first GLP-1 RA use are as follows:

Table 2: Patient Characteristics by Year of First GLP-1 RA Prescription.

	2018	2019	2020	2021	2022	2023	2024
N	64,989 (100.00%)	82,046 (100.00%)	89,267 (100.00%)	146,745 (100.00%)	286,101 (100.00%)	527,686 (100.00%)	597,885 (100.00%)
Age Group							
18-44	10,725 (16.50%)	13,696 (16.69%)	15,448 (17.31%)	28,755 (19.60%)	69,398 (24.26%)	140,084 (26.55%)	166,063 (27.78%)
45-64	35,837 (55.14%)	43,685 (53.24%)	46,708 (52.32%)	74,694 (50.90%)	143,105 (50.02%)	261,484 (49.55%)	283,093 (47.35%)
65+	18,427 (28.35%)	24,665 (30.06%)	27,111 (30.37%)	43,296 (29.50%)	73,598 (25.72%)	126,118 (23.90%)	148,729 (24.88%)
Sex							
Female	36,448 (56.08%)	45,532 (55.50%)	49,067 (54.97%)	85,949 (58.57%)	182,181 (63.68%)	349,767 (66.28%)	395,398 (66.13%)

	2018	2019	2020	2021	2022	2023	2024
Male	28,541 (43.92%)	36,514 (44.50%)	40,200 (45.03%)	60,796 (41.43%)	103,920 (36.32%)	177,919 (33.72%)	202,487 (33.87%)
Race							
White	47,348 (72.86%)	59,078 (72.01%)	62,910 (70.47%)	102,679 (69.97%)	200,588 (70.11%)	363,739 (68.93%)	410,553 (68.67%)
Black or African American	7,558 (11.63%)	9,330 (11.37%)	10,529 (11.79%)	18,302 (12.47%)	36,614 (12.80%)	70,240 (13.31%)	71,390 (11.94%)
Asian	1,261 (1.94%)	1,866 (2.27%)	2,256 (2.53%)	4,089 (2.79%)	6,794 (2.37%)	11,630 (2.20%)	13,480 (2.25%)
Unknown	6,549 (10.08%)	8,282 (10.09%)	9,321 (10.44%)	14,043 (9.57%)	26,129 (9.13%)	53,905 (10.22%)	71,534 (11.96%)
Native Hawaiian or Other Pacific Islander	213 (0.33%)	282 (0.34%)	345 (0.39%)	594 (0.40%)	1,077 (0.38%)	1,664 (0.32%)	1,908 (0.32%)
American Indian or Alaska Native	470 (0.72%)	627 (0.76%)	662 (0.74%)	1,233 (0.84%)	2,210 (0.77%)	3,471 (0.66%)	3,821 (0.64%)
Other Race	1,590 (2.45%)	2,581 (3.15%)	3,244 (3.63%)	5,805 (3.96%)	12,689 (4.44%)	23,037 (4.37%)	25,199 (4.21%)
Ethnicity							
Hispanic or Latino	6,151 (9.46%)	8,554 (10.43%)	10,301 (11.54%)	18,271 (12.45%)	37,507 (13.11%)	65,456 (12.40%)	74,339 (12.43%)
Not Hispanic or Latino	54,722 (84.20%)	68,124 (83.03%)	73,349 (82.17%)	119,909 (81.71%)	232,767 (81.36%)	432,545 (81.97%)	489,143 (81.81%)
Unknown	4,116 (6.33%)	5,368 (6.54%)	5,617 (6.29%)	8,565 (5.84%)	15,827 (5.53%)	29,685 (5.63%)	34,403 (5.75%)
Characteristics							
Type II Diabetes	45,608 (70.18%)	60,511 (73.75%)	63,898 (71.58%)	95,224 (64.89%)	148,765 (52.00%)	230,186 (43.62%)	221,336 (37.02%)
Overweight or Obesity	42,313 (65.11%)	52,277 (63.72%)	54,288 (60.82%)	93,870 (63.97%)	183,990 (64.31%)	358,881 (68.01%)	423,547 (70.84%)
Medication							
Semaglutide	3,673 (5.65%)	18,181 (22.16%)	31,872 (35.70%)	71,149 (48.48%)	163,372 (57.10%)	361,903 (68.58%)	366,070 (61.23%)
Tirzepatide	n/a	n/a	n/a	n/a	24,616 (8.60%)	68,911 (13.06%)	181,221 (30.31%)
Liraglutide	29,346 (45.16%)	26,940 (32.84%)	19,662 (22.03%)	23,039 (15.70%)	25,420 (8.88%)	22,785 (4.32%)	9,278 (1.55%)
Dulaglutide	22,584 (34.75%)	29,229 (35.63%)	31,664 (35.47%)	47,636 (32.46%)	67,715 (23.67%)	69,386 (13.15%)	37,651 (6.30%)
Other	9,386 (14.44%)	7,696 (9.38%)	6,069 (6.80%)	4,921 (3.35%)	4,978 (1.74%)	4,701 (0.89%)	3,665 (0.61%)
Labeled Use							
ADM	41,667 (64.11%)	51,497 (62.77%)	63,789 (71.46%)	109,311 (74.49%)	203,214 (71.03%)	308,737 (58.51%)	269,257 (45.03%)
AOM	8,092 (12.45%)	7,162 (8.73%)	5,950 (6.67%)	10,365 (7.06%)	21,605 (7.55%)	88,267 (16.73%)	167,291 (27.98%)
Unknown	15,230 (23.43%)	23,387 (28.50%)	19,528 (21.88%)	27,069 (18.45%)	61,282 (21.42%)	130,682 (24.77%)	161,337 (26.98%)

	2018	2019	2020	2021	2022	2023	2024
Comorbidity							
Asthma	7,232 (11.13%)	10,148 (12.37%)	11,175 (12.52%)	19,604 (13.36%)	38,682 (13.52%)	74,757 (14.17%)	88,052 (14.73%)
CKD	10,292 (15.84%)	14,602 (17.80%)	16,787 (18.81%)	26,196 (17.85%)	39,456 (13.79%)	61,222 (11.60%)	66,351 (11.10%)
COPD	4,012 (6.17%)	5,500 (6.70%)	6,087 (6.82%)	9,055 (6.17%)	14,681 (5.13%)	25,764 (4.88%)	29,309 (4.90%)
Hyperlipidemia	37,106 (57.10%)	49,403 (60.21%)	52,054 (58.31%)	82,067 (55.92%)	145,974 (51.02%)	260,084 (49.29%)	297,720 (49.80%)
Hypertension	39,377 (60.59%)	52,714 (64.25%)	55,111 (61.74%)	87,185 (59.41%)	154,576 (54.03%)	273,554 (51.84%)	305,762 (51.14%)
Major Depression	6,521 (10.03%)	9,531 (11.62%)	10,624 (11.90%)	19,349 (13.19%)	36,388 (12.72%)	61,532 (11.66%)	68,850 (11.52%)

60-day dispensing over time

We describe the proportion of patients with any GLP-1 RA prescriptions (first-time and repeat) in a given month who filled any GLP-1 RA in the 60 days following their prescription.

60-day initiation over time

We describe the proportion of patients first-prescribed a GLP-1 RA who filled any GLP-1 RA in the subsequent 60 days over time.

Patient characteristics are given in Table 3:

Table 3: Patient Characteristics by Dispense Status. Dispense status refers to presence of any GLP-1 RA dispense within 60 days of first GLP-1 RA prescription. Table contains the subset of patients with observable 60-day dispense outcomes.

	Dispensed in 60 Days	Not Dispensed in 60 Days	Overall
N	1,003,585 (100.00%)	710,291 (100.00%)	1,713,876 (100.00%)
Age Group			
18-44	220,541 (21.98%)	200,978 (28.30%)	421,519 (24.59%)
45-64	510,194 (50.84%)	341,838 (48.13%)	852,032 (49.71%)
65+	272,850 (27.19%)	167,475 (23.58%)	440,325 (25.69%)
Sex			
Female	621,035 (61.88%)	472,322 (66.50%)	1,093,357 (63.79%)
Male	382,550 (38.12%)	237,969 (33.50%)	620,519 (36.21%)
Race			
White	685,335 (68.29%)	506,139 (71.26%)	1,191,474 (69.52%)
Black or African American	124,292 (12.38%)	90,778 (12.78%)	215,070 (12.55%)
Asian	25,133 (2.50%)	14,800 (2.08%)	39,933 (2.33%)
Unknown	114,281 (11.39%)	64,681 (9.11%)	178,962 (10.44%)
Native Hawaiian or Other Pacific Islander	3,661 (0.36%)	2,219 (0.31%)	5,880 (0.34%)
American Indian or Alaska Native	7,123 (0.71%)	4,814 (0.68%)	11,937 (0.70%)
Other Race	43,760 (4.36%)	26,860 (3.78%)	70,620 (4.12%)
Ethnicity			
Hispanic or Latino	125,367 (12.49%)	85,645 (12.06%)	211,012 (12.31%)
Not Hispanic or Latino	819,338 (81.64%)	586,940 (82.63%)	1,406,278 (82.05%)
Unknown	58,880 (5.87%)	37,706 (5.31%)	96,586 (5.64%)
Characteristics			
Type II Diabetes	560,020 (55.80%)	276,983 (39.00%)	837,003 (48.84%)
Overweight or Obesity	659,978 (65.76%)	518,699 (73.03%)	1,178,677 (68.77%)
Medication			
Semaglutide	544,988 (54.30%)	426,248 (60.01%)	971,236 (56.67%)
Tirzepatide	156,982 (15.64%)	101,362 (14.27%)	258,344 (15.07%)
Liraglutide	77,790 (7.75%)	73,124 (10.29%)	150,914 (8.81%)
Dulaglutide	201,319 (20.06%)	92,252 (12.99%)	293,571 (17.13%)
Other	22,506 (2.24%)	17,305 (2.44%)	39,811 (2.32%)
Labeled Use			
ADM	673,927 (67.15%)	327,921 (46.17%)	1,001,848 (58.46%)
AOM	128,412 (12.80%)	165,853 (23.35%)	294,265 (17.17%)
Unknown	201,246 (20.05%)	216,517 (30.48%)	417,763 (24.38%)
Year			
2018	30,223 (3.01%)	33,687 (4.74%)	63,910 (3.73%)
2019	43,756 (4.36%)	36,559 (5.15%)	80,315 (4.69%)
2020	48,351 (4.82%)	39,077 (5.50%)	87,428 (5.10%)
2021	86,380 (8.61%)	57,494 (8.09%)	143,874 (8.39%)

	Dispensed in 60 Days	Not Dispensed in 60 Days	Overall
2022	177,153 (17.65%)	100,996 (14.22%)	278,149 (16.23%)
2023	311,461 (31.03%)	202,116 (28.46%)	513,577 (29.97%)
2024	306,261 (30.52%)	240,362 (33.84%)	546,623 (31.89%)
Comorbidity			
Asthma	137,500 (13.70%)	104,904 (14.77%)	242,404 (14.14%)
CKD	143,185 (14.27%)	86,424 (12.17%)	229,609 (13.40%)
COPD	55,899 (5.57%)	35,649 (5.02%)	91,548 (5.34%)
Hyperlipidemia	532,933 (53.10%)	365,455 (51.45%)	898,388 (52.42%)
Hypertension	552,365 (55.04%)	382,972 (53.92%)	935,337 (54.57%)
Major Depression	113,680 (11.33%)	96,153 (13.54%)	209,833 (12.24%)

By Age

As a proxy for Medicare insurance, we segment initiation rate by populations 65+ versus 18-64.

Patient characteristics for patients aged 65+ are given in Table 4:

Table 4: Patient Characteristics by Dispense Status, Age 65+. Dispense status refers to presence of any GLP-1 RA dispense within 60 days of first GLP-1 RA prescription. Table contains the subset of patients with observable 60-day dispense outcomes.

	Dispensed in 60 Days	Not Dispensed in 60 Days	Overall
N	272,850 (100.00%)	167,475 (100.00%)	440,325 (100.00%)
Age Group			
65+	272,850 (100.00%)	167,475 (100.00%)	440,325 (100.00%)
Sex			

	Dispensed in 60 Days	Not Dispensed in 60 Days	Overall
Female	151,369 (55.48%)	94,027 (56.14%)	245,396 (55.73%)
Male	121,481 (44.52%)	73,448 (43.86%)	194,929 (44.27%)
Race			
White	202,910 (74.37%)	130,232 (77.76%)	333,142 (75.66%)
Black or African American	24,128 (8.84%)	14,593 (8.71%)	38,721 (8.79%)
Asian	5,901 (2.16%)	2,881 (1.72%)	8,782 (1.99%)
Unknown	29,040 (10.64%)	14,823 (8.85%)	43,863 (9.96%)
Native Hawaiian or Other Pacific Islander	556 (0.20%)	268 (0.16%)	824 (0.19%)
American Indian or Alaska Native	1,478 (0.54%)	803 (0.48%)	2,281 (0.52%)
Other Race	8,837 (3.24%)	3,875 (2.31%)	12,712 (2.89%)
Ethnicity			
Hispanic or Latino	22,554 (8.27%)	11,438 (6.83%)	33,992 (7.72%)
Not Hispanic or Latino	233,203 (85.47%)	145,972 (87.16%)	379,175 (86.11%)
Unknown	17,093 (6.26%)	10,065 (6.01%)	27,158 (6.17%)
Characteristics			
Type II Diabetes	187,923 (68.87%)	98,637 (58.90%)	286,560 (65.08%)
Overweight or Obesity	170,829 (62.61%)	108,536 (64.81%)	279,365 (63.45%)
Medication			
Semaglutide	148,450 (54.41%)	93,482 (55.82%)	241,932 (54.94%)
Tirzepatide	29,768 (10.91%)	15,471 (9.24%)	45,239 (10.27%)
Liraglutide	17,582 (6.44%)	16,929 (10.11%)	34,511 (7.84%)
Dulaglutide	67,878 (24.88%)	34,536 (20.62%)	102,414 (23.26%)
Other	9,172 (3.36%)	7,057 (4.21%)	16,229 (3.69%)
Labeled Use			
ADM	207,931 (76.21%)	99,798 (59.59%)	307,729 (69.89%)
AOM	10,779 (3.95%)	21,859 (13.05%)	32,638 (7.41%)
Unknown	54,140 (19.84%)	45,818 (27.36%)	99,958 (22.70%)
Year			
2018	8,284 (3.04%)	9,771 (5.83%)	18,055 (4.10%)
2019	12,570 (4.61%)	11,433 (6.83%)	24,003 (5.45%)
2020	14,382 (5.27%)	12,006 (7.17%)	26,388 (5.99%)
2021	25,076 (9.19%)	17,132 (10.23%)	42,208 (9.59%)
2022	45,440 (16.65%)	25,239 (15.07%)	70,679 (16.05%)
2023	83,434 (30.58%)	38,748 (23.14%)	122,182 (27.75%)
2024	83,664 (30.66%)	53,146 (31.73%)	136,810 (31.07%)
Comorbidity			
Asthma	32,894 (12.06%)	20,620 (12.31%)	53,514 (12.15%)
CKD	82,326 (30.17%)	48,975 (29.24%)	131,301 (29.82%)
COPD	27,859 (10.21%)	17,999 (10.75%)	45,858 (10.41%)
Hyperlipidemia	183,046 (67.09%)	118,137 (70.54%)	301,183 (68.40%)
Hypertension	191,656 (70.24%)	124,392 (74.27%)	316,048 (71.78%)
Major Depression	30,050 (11.01%)	20,010 (11.95%)	50,060 (11.37%)

Patient characteristics for patients aged 18-65 are given in Table 5:

Table 5: Patient Characteristics by Dispense Status, Age 18-64. Dispense status refers to presence of any GLP-1 RA dispense within 60 days of first GLP-1 RA prescription. Table contains the subset of patients with observable 60-day dispense outcomes.

	Dispensed in 60 Days	Not Dispensed in 60 Days	Overall
N	730,735 (100.00%)	542,816 (100.00%)	1,273,551 (100.00%)
Age Group			
18-44	220,541 (30.18%)	200,978 (37.03%)	421,519 (33.10%)
45-64	510,194 (69.82%)	341,838 (62.97%)	852,032 (66.90%)
Sex			
Female	469,666 (64.27%)	378,295 (69.69%)	847,961 (66.58%)
Male	261,069 (35.73%)	164,521 (30.31%)	425,590 (33.42%)
Race			
White	482,425 (66.02%)	375,907 (69.25%)	858,332 (67.40%)
Black or African American	100,164 (13.71%)	76,185 (14.04%)	176,349 (13.85%)
Asian	19,232 (2.63%)	11,919 (2.20%)	31,151 (2.45%)
Unknown	85,241 (11.67%)	49,858 (9.19%)	135,099 (10.61%)
Native Hawaiian or Other Pacific Islander	3,105 (0.42%)	1,951 (0.36%)	5,056 (0.40%)
American Indian or Alaska Native	5,645 (0.77%)	4,011 (0.74%)	9,656 (0.76%)
Other Race	34,923 (4.78%)	22,985 (4.23%)	57,908 (4.55%)
Ethnicity			
Hispanic or Latino	102,813 (14.07%)	74,207 (13.67%)	177,020 (13.90%)
Not Hispanic or Latino	586,135 (80.21%)	440,968 (81.24%)	1,027,103 (80.65%)
Unknown	41,787 (5.72%)	27,641 (5.09%)	69,428 (5.45%)
Characteristics			
Type II Diabetes	372,097 (50.92%)	178,346 (32.86%)	550,443 (43.22%)
Overweight or Obesity	489,149 (66.94%)	410,163 (75.56%)	899,312 (70.61%)
Medication			
Semaglutide	396,538 (54.27%)	332,766 (61.30%)	729,304 (57.27%)
Tirzepatide	127,214 (17.41%)	85,891 (15.82%)	213,105 (16.73%)
Liraglutide	60,208 (8.24%)	56,195 (10.35%)	116,403 (9.14%)
Dulaglutide	133,441 (18.26%)	57,716 (10.63%)	191,157 (15.01%)
Other	13,334 (1.82%)	10,248 (1.89%)	23,582 (1.85%)
Labeled Use			
ADM	465,996 (63.77%)	228,123 (42.03%)	694,119 (54.50%)
AOM	117,633 (16.10%)	143,994 (26.53%)	261,627 (20.54%)
Unknown	147,106 (20.13%)	170,699 (31.45%)	317,805 (24.95%)
Year			
2018	21,939 (3.00%)	23,916 (4.41%)	45,855 (3.60%)
2019	31,186 (4.27%)	25,126 (4.63%)	56,312 (4.42%)
2020	33,969 (4.65%)	27,071 (4.99%)	61,040 (4.79%)

	Dispensed in 60 Days	Not Dispensed in 60 Days	Overall
2021	61,304 (8.39%)	40,362 (7.44%)	101,666 (7.98%)
2022	131,713 (18.02%)	75,757 (13.96%)	207,470 (16.29%)
2023	228,027 (31.21%)	163,368 (30.10%)	391,395 (30.73%)
2024	222,597 (30.46%)	187,216 (34.49%)	409,813 (32.18%)
Comorbidity			
Asthma	104,606 (14.32%)	84,284 (15.53%)	188,890 (14.83%)
CKD	60,859 (8.33%)	37,449 (6.90%)	98,308 (7.72%)
COPD	28,040 (3.84%)	17,650 (3.25%)	45,690 (3.59%)
Hyperlipidemia	349,887 (47.88%)	247,318 (45.56%)	597,205 (46.89%)
Hypertension	360,709 (49.36%)	258,580 (47.64%)	619,289 (48.63%)
Major Depression	83,630 (11.44%)	76,143 (14.03%)	159,773 (12.55%)

Discussion and Conclusion

This analysis summarized GLP-1 RA prescribing trends and patient characteristics from January 2018 to December 2024. During this period 1,828,203 patients were prescribed a total of 8,412,584 total prescriptions. ADMs accounted for 81.0% of prescriptions, and AOMs accounted for 19.0%, among prescriptions where use was known. A larger proportion of ADMs, compared to AOMs, were dispensed within 60 days: 67.3% of first-time ADM prescriptions and 43.6% of first-time AOM prescriptions had a fill within 60 days.

Several limitations of our analysis should be noted. All data are preliminary and may change as additional data are obtained. These findings are consistent with data accessed March 03, 2025. Data presented in this analysis represent raw counts and/or rates, and post-stratification methods have not been conducted. In addition, cohorts with small counts may be suppressed during the de-identification process leading to the appearance of zero patients for a given time period. Given relatively large numbers of patients prescribed and dispensed GLP-1 RA, we do not expect substantial suppression in the most recent periods. The unknowns in this report either indicate the value was not included in the individual's electronic health record or that it was excluded from the data to protect an individual's identity as a part of Truveta's commitment to privacy (Truveta, 2024). Finally, this report is a description of prescribing trends and patient characteristics only. Results should not be interpreted as causal.

References

Truveta. (2024). Truveta's Approach to Patient Privacy.
https://resources.truveta.com/patient-privacy

Centers for disease control and prevention. "National diabetes statistics report". 2020.
https://www.cdc.gov/diabetes/data/statistics-report/index.html.

Davies, Melanie J., et al. "Efficacy of liraglutide for weight loss among patients with type 2 diabetes: the SCALE diabetes randomized clinical trial." JAMA 314.7 (2015): 687-699.

Davies, Melanie, et al. "Semaglutide 2· 4 mg once a week in adults with overweight or obesity, and type 2 diabetes (STEP 2): a randomised, double-blind, double-dummy, placebo-controlled, phase 3 trial." The Lancet 397.10278 (2021): 971-984.

Hales, Craig M., et al. "Prevalence of obesity and severe obesity among adults: United States, 2017–2018. NCHS Data Brief, no 360." National Center for Health Statistics 360 (2020): 1-7.

Han, Sabrina H., et al. "Public interest in the off-label use of glucagon-like peptide 1 agonists (Ozempic) for cosmetic weight loss: a google trends analysis." Aesthetic Surgery Journal 44.1 (2024): 60-67.

Garvey, W. Timothy, et al. "Tirzepatide once weekly for the treatment of obesity in people with type 2 diabetes (SURMOUNT-2): a double-blind, randomised, multicentre, placebo-controlled, phase 3 trial." The Lancet (2023).

Jastreboff, Ania M., et al. "Tirzepatide once weekly for the treatment of obesity." New England Journal of Medicine 387.3 (2022): 205-216.

Lincoff, A. Michael, et al. "Semaglutide and cardiovascular outcomes in obesity without diabetes." New England Journal of Medicine 389.24 (2023): 2221-2232.

Ussher, John R., and Daniel J. Drucker. "Glucagon-like peptide 1 receptor agonists: cardiovascular benefits and mechanisms of action." Nature Reviews Cardiology 20.7 (2023): 463-474.

Wilding, John PH, et al. "Once-weekly semaglutide in adults with overweight or obesity." New England Journal of Medicine 384.11 (2021): 989-1002.

Whitley, Heather P., Jennifer M. Trujillo, and Joshua J. Neumiller. "Potential Strategies for Addressing GLP-1 and Dual GLP-1/GIP Receptor Agonist Shortages." Clinical Diabetes (2023): cd230023.

Yeh, Tzu-Lin, et al. "Effect of glucagon-like peptide-1 receptor agonists on glycemic control, and weight reduction in adults: A multivariate meta-analysis." PLOS One 18.1 (2023): e0278685.

Acknowledgements

The authors thank Dr. Ty Gluckman for his contributions and feedback on describing GLP-1 RA prescribing trends.