Monitoring Report: GLP-1 RA Prescribing Trends – June 2025 Data

Samuel Gratzl, PhD*

Brianna M Goodwin Cartwright, MS*

Patricia J Rodriguez, PhD, MPH

Karen Gilbert, PhD

Duy Do, PhD

Nina Masters, MPH, PhD

Nicholas L Stucky, MD, PhD

*Denotes co-first author

Affiliations: All authors are affiliated with Truveta, Inc.

Corresponding author:
Dr Nicholas Stucky
Truveta, Inc
1745 114th Ave SE, Bellevue, WA 98004
Email: nicholass@truveta.com

Abstract

Background: Limited recent data exist on prescribing patterns and patient characteristics for glucagon-like peptide 1 receptor agonists (GLP-1 RAs), an important drug class used as anti-diabetic medication (ADM) for patients with type 2 diabetes mellitus (T2D) and/or anti-obesity medication (AOM) in patients with overweight or obesity.

For brevity, we use the term GLP-1 RA to refer to both GLP-1 RA and dual GLP-1 RA/GIP medications.

Objective: To describe recent trends in prescribing and dispensing of GLP-1-based medications in the US.

Methods: Using a subset of real-world electronic health record (EHR) data from Truveta, a growing collective of health systems that provide more than 18% of all daily clinical care in the US, we identified people who were prescribed a GLP-1-based medication between January 01, 2018 and June 30, 2025. We describe prescribing volumes and patient characteristics over time, by medication, and by FDA-labeled use. Among the subset of patients for whom post-prescription dispensing data is available, we describe the proportion and characteristics of patients who were and were not dispensed a GLP-1 RA following their prescription.

Results: 2,196,747 patients were prescribed a GLP-1 RA between January 2018 and June 2025, with 10,622,359 total prescriptions during this period. Among first-time prescriptions for which use could be established, ADMs accounted for 73.9% and AOMs accounted for 26.1%. The most common first-time medication was semaglutide (n = 1,175,286). Among those with available data, 69.5% of first-time ADM prescriptions overall and 72.9% in March 2025 had a fill within 60 days. By comparison, 46.9% of first-time AOM prescriptions overall and 49.2% in March 2025 had a fill within 60 days.

Trends in Prescribing

	GLP-1 RA	Semaglutide	Tirzepatide	Liraglutide	Dulaglutide	Other
Overall						
March 2025	5.79%	2.71%	2.68%	0.07%	0.43%	0.03%
June 2025	6.24%	2.62%	3.27%	0.07%	0.40%	0.02%
Change (percentage)	+7.71%	-3.33%	+21.84%	-10.90%	-6.74%	-28.99%
Difference (percentage points)	+0.45%	-0.09%	+0.59%	-0.01%	-0.03%	-0.01%
ADM						
March 2025	2.88%	1.27%	1.18%	0.01%	0.43%	0.03%
June 2025	2.94%	1.19%	1.36%	0.00%	0.40%	0.02%
Change (percentage)	+2.11%	-6.40%	+15.00%	-48.88%	-6.74%	-28.99%
Difference (percentage points)	+0.06%	-0.08%	+0.18%	-0.00%	-0.03%	-0.01%
AOM						
March 2025	1.61%	0.62%	1.00%	0.01%	n/a	n/a

	GLP-1 RA	Semaglutide	Tirzepatide	Liraglutide	Dulaglutide	Other
June 2025	1.97%	0.67%	1.32%	0.01%	n/a	n/a
Change (percentage)	+22.39%	+8.05%	+31.98%	-2.59%		
Difference (percentage points)	+0.36%	+0.05%	+0.32%	-0.00%		
Unknown						
March 2025	1.57%	0.91%	0.62%	0.06%	n/a	n/a
June 2025	1.61%	0.84%	0.73%	0.05%	n/a	n/a
Change (percentage)	+2.84%	-7.07%	+18.34%	-9.12%		
Difference (percentage points)	+0.04%	-0.06%	+0.11%	-0.01%		

- Overall prescribing rates (GLP-1 RA prescriptions per total prescriptions) in June 2025 increased slightly relative to March 2025 (+7.71%).
- Month-over-month ADM prescribing increased in April and then remained relatively stable through June 2025. ADM prescribing in June 2025 are about the same as March 2025 (+2.11%).
- Among ADMs, tirzepatide is now the most prescribed ADM medication.
- AOM prescribing increased each month between March and June 2025, which an overall increase of +22.39%.
- AOM tirzepatide and semaglutide both increased in rates between March and June 2025, however tirzepatide increased by 31.98% to reach an overall prescription rate of 1.32%, while AOM semaglutide increased by 8.05% to reach an overall prescription rate of 0.67%.

Trends in First-Time Prescribing

Changes in Rate of First-Time GLP-1 RA Prescribing over Time, by Medication and Labeled Use between February 2025 and June 2025

	GLP-1 RA	Semaglutide	Tirzepatide	Liraglutide	Dulaglutide	Other
Overall						
March 2025	1.34%	0.66%	0.61%	0.01%	0.06%	0.01%
June 2025	1.28%	0.55%	0.67%	0.01%	0.05%	0.00%
Change (percentage)	-4.84%	-15.76%	+9.87%	-33.98%	-23.59%	-52.99%
Difference (percentage points)	-0.06%	-0.10%	+0.06%	-0.00%	-0.01%	-0.00%
ADM						
March 2025	0.48%	0.23%	0.18%	0.00%	0.06%	0.01%
June 2025	0.41%	0.18%	0.18%	0.00%	0.05%	0.00%
Change (percentage)	-15.11%	-22.81%	-1.38%	-69.63%	-23.59%	-52.99%
Difference (percentage points)	-0.07%	-0.05%	-0.00%	-0.00%	-0.01%	-0.00%
AOM						
March 2025	0.43%	0.15%	0.27%	0.00%	n/a	n/a
June 2025	0.47%	0.14%	0.32%	0.00%	n/a	n/a
Change (percentage)	+8.54%	-5.35%	+16.39%	-8.68%		
Difference (percentage points)	+0.04%	-0.01%	+0.05%	-0.00%		
Unknown						
March 2025	0.44%	0.28%	0.15%	0.01%	n/a	n/a
June 2025	0.41%	0.23%	0.17%	0.01%	n/a	n/a
Change (percentage)	-6.75%	-15.71%	+11.57%	-33.74%		
Difference (percentage points)	-0.03%	-0.04%	+0.02%	-0.00%		

- First-time prescribing rates (first-time GLP-1 RA prescriptions per total prescriptions) in June 2025 are slightly decreased compared to March 2025 (-4.84%).
- First-time prescribing of ADMs decreased in June 2025, relative to March 2025 (-15.11%).
- First-time prescribing of AOMs increased in June 2025, relative to March 2025 (+8.54%).

- First-time prescribing of both ADM (-22.81%) and AOM (-5.35%) semaglutide decreased throughout the quarter. First-time prescribing of ADM tirzepatide slightly decreased (-1.38%), while first-time prescribing of AOM tirzepatide (16.39%) increased throughout the quarter.

Trends in Dispensing

	GLP-1 RA	Semaglutide	Tirzepatide	Liraglutide	Dulaglutide	Other
Overall						
December 2024	55.06%	52.63%	55.84%	63.26%	74.70%	61.78%
March 2025	54.88%	53.35%	54.09%	59.40%	77.97%	67.60%
Change (percentage)	-0.34%	+1.37%	-3.14%	-6.10%	+4.38%	+9.42%
Difference (percentage points)	-0.19%	+0.72%	-1.76%	-3.86%	+3.27%	+5.82%
ADM						
December 2024	69.94%	68.16%	71.04%	81.18%	74.70%	61.78%
March 2025	72.87%	71.71%	72.92%	61.82%	77.97%	67.60%
Change (percentage)	+4.19%	+5.20%	+2.65%	-23.85%	+4.38%	+9.42%
Difference (percentage points)	+2.93%	+3.54%	+1.88%	-19.36%	+3.27%	+5.82%
AOM						
December 2024	49.82%	48.32%	51.07%	60.94%	n/a	n/a
March 2025	49.18%	49.21%	49.18%	46.38%	n/a	n/a
Change (percentage)	-1.30%	+1.83%	-3.69%	-23.89%		
Difference (percentage points)	-0.65%	+0.89%	-1.89%	-14.56%		

	GLP-1 RA	Semaglutide	Tirzepatide	Liraglutide	Dulaglutide	Other
Unknown						
December 2024	40.47%	40.33%	39.38%	58.36%	n/a	n/a
March 2025	40.12%	40.08%	38.79%	61.65%	n/a	n/a
Change (percentage)	-0.88%	-0.61%	-1.51%	+5.64%		
Difference (percentage points)	-0.36%	-0.25%	-0.59%	+3.29%		

- During the full time period, 71.1% of GLP-1 RA prescriptions had a dispense within 60 days of their prescription (e.g., were initiated within 60 days).
- For patients first prescribed an ADM in March 2025, 72.9% filled a GLP-1 RA within 60 days. Initiation rates for ADM GLP-1 RA are about the same in March 2025, relative to December 2024 (+4.19%).
- For patients first prescribed an AOM in March 2025, 49.2% filled a GLP-1 RA within 60 days. Initiation rates for AOM GLP-1 RA are about the same in March 2025, relative to December 2024 (-1.30%).

About this report

This report provides recent trends in the prescribing and dispensing of glucagon-like peptide-1 receptor agonist (GLP-1 RAs)-based medications, including dual GLP-1 RA/gastric inhibitory polypeptide (GIP) over time, by medication, and by FDA-labeled use. We used near real-time, aggregated electronic health record (EHR) data from a collective of US healthcare systems to describe prescribing and dispensing of GLP-1 RA medications from January 2018 to June 2025.

Introduction

Obesity and type 2 diabetes mellitus (T2D) are interconnected epidemics with a high prevalence and individual and combined negative impacts on health (CDC, 2020, Hales, 2020). Glucagon-like peptide-1 receptor agonist (GLP-1 RAs) -based medications, including dual GLP-1 RA/gastric inhibitory polypeptide (GIP), have been shown to improve glycemic control in patients with T2D and reduce weight in patients with and without T2D (Davies, 2015, Davies, 2021, Garvey, 2023, Jastreboff, 2022, Wilding, 2023, Yeh, 2023). GLP-1 RA have also demonstrated cardiovascular benefit in patients with and without T2D (Lincoff, 2023, Ussher, 2023).

Interest in these medications has recently accelerated, largely for their weight-loss effects; however, limited empiric data exist on prescribing, dispensing, and patient

characteristics. Access to and use of GLP-1 RA may be impacted by high cost, limited insurance coverage for patients without T2D and medication shortages.

Methods

Data

We evaluated prescribing and dispensing of GLP-1 RAs between January 2018 and June 2025 using a subset of Truveta Data. Truveta provides access to continuously updated, linked, and de-identified electronic health record (EHR) from a collective of US health care systems that provide 18% of all daily clinical care in the US, including structured information on demographics, encounters, diagnoses, vital signs (e.g., weight, BMI, blood pressure), medication requests (prescriptions), medication administration, laboratory and diagnostic tests and results (e.g., HbA1c tests and values), and procedures. Updated EHR data are provided daily to Truveta by member health care systems. In addition to EHR data for care delivered within Truveta member health care systems, medication dispensing and social drivers of health (SDOH) information are made available through linked third-party data. Medication dispense (via e-prescribing data) includes fills for prescriptions written both within and outside Truveta member health care systems, providing greater observability into patients' medication history. Medication dispense histories are updated at the time of the encounter and include fill dates, NDC or RxNorm codes, quantity dispensed, and days of medication supplied.

Population

We identified GLP-1 RA prescriptions (medication requests) and dispenses among adults using RxNorm and NDC codes for dulaglutide, exenatide, liraglutide, lixisenatide, semaglutide, and tirzepatide. Exenatide and lixisenatide were grouped as 'other' given small relative sizes. Brands were identified when possible using RxNorm and NDC codes. Brands labeled for treatment of T2D were classified as ADMs and brands labeled for treatment of overweight and obesity were classified as AOMs. When brand could not be established, the labeled use was classified as unknown.

Prescribing Analysis

Characteristics of patients ever prescribed a GLP-1 RA were summarized, including demographics, social drivers of health (SDOH), and presence of T2D and/or overweight or obesity. Characteristics are provided by medication and by year of first prescription.

Overall and first-time prescribing rates of GLP-1 RA were summarized over time by medication and labeled use. The prescribing rate was calculated monthly as the

number of patients with a GLP-1 RA prescription divided by the number of patients with any prescription in the month. Absolute differences and percentage changes in the rate of GLP-1 RA prescribing over the most recent 3-month period were summarized overall and by medication and labeled use.

Dispensing Analysis

We describe the proportion of GLP-1 RA prescriptions (overall and first-time) with a dispense occurring within 60 days. Dispensing analyses were restricted to prescriptions for which complete 60-day dispensing outcomes could be observed. Dispensing outcomes were considered observable if (1) a dispense was recorded in the 60 days following the prescription, and/or (2) an encounter took place after the 60-day period ended (at which time any dispenses would be observed). To allow sufficient follow-up, we report dispensing with a 90-day delay. Given shortages and expected substitutions at the pharmacy, we considered the dispense of any GLP-1 RA , which could differ from the prescribed GLP-1 RA.

We plot the proportion of prescriptions (overall and first-time) dispensed within 60 days over time, by medication and use. Differences in dispensing were expected between ADMs and AOMs given ADMs are typically covered by insurance, whereas insurance coverage for AOMs is limited. As such, we provide dispensing rates stratified by FDA labeled use. Further, we stratify dispensing by age 65+ vs. 18-64 as a proxy for Medicare coverage. Medicare never provides coverage for AOMs, whereas commercial insurers may.

Results

Overall population

The demographics of patients by first-prescribed medication are as follows:

Table 1: Patient Characteristics by First-Prescribed GLP-1 RA Medication.

	Semaglutide	Tirzepatide	Liraglutide	Dulaglutide	Other	Overall
N	1,175,286 (100.00%)	436,794 (100.00%)	150,779 (100.00%)	321,747 (100.00%)	41,562 (100.00%)	2,126,168 (100.00%)
Age Group						
18-44	311,703 (26.52%)	139,856 (32.02%)	41,005 (27.20%)	46,110 (14.33%)	4,845 (11.66%)	543,519 (25.56%)
45-64	567,088 (48.25%)	215,145 (49.26%)	74,752 (49.58%)	161,619 (50.23%)	19,653 (47.29%)	1,038,257 (48.83%)

	Semaglutide	Tirzepatide	Liraglutide	Dulaglutide	Other	Overall
65+	296,495 (25.23%)	81,793 (18.73%)	35,022 (23.23%)	114,018 (35.44%)	17,064 (41.06%)	544,392 (25.60%)
Sex						
Female	766,409 (65.21%)	295,758 (67.71%)	102,109 (67.72%)	173,582 (53.95%)	22,155 (53.31%)	1,360,013 (63.97%)
Male	408,877 (34.79%)	141,036 (32.29%)	48,670 (32.28%)	148,165 (46.05%)	19,407 (46.69%)	766,155 (36.03%)
Race						
White	839,530 (71.43%)	320,382 (73.35%)	108,650 (72.06%)	216,760 (67.37%)	30,184 (72.62%)	1,515,506 (71.28%)
Black or African American	158,261 (13.47%)	50,510 (11.56%)	20,776 (13.78%)	50,746 (15.77%)	4,937 (11.88%)	285,230 (13.42%)
Asian	30,059 (2.56%)	9,480 (2.17%)	2,783 (1.85%)	8,619 (2.68%)	1,000 (2.41%)	51,941 (2.44%)
Unknown	84,725 (7.21%)	34,632 (7.93%)	10,144 (6.73%)	26,276 (8.17%)	3,240 (7.80%)	159,017 (7.48%)
Native Hawaiian or Other Pacific Islander	4,259 (0.36%)	1,242 (0.28%)	668 (0.44%)	1,383 (0.43%)	179 (0.43%)	7,731 (0.36%)
American Indian or Alaska Native	8,630 (0.73%)	2,463 (0.56%)	1,696 (1.12%)	2,417 (0.75%)	378 (0.91%)	15,584 (0.73%)
Other Race	49,822 (4.24%)	18,085 (4.14%)	6,062 (4.02%)	15,546 (4.83%)	1,644 (3.96%)	91,159 (4.29%)
Ethnicity						
Hispanic or Latino	145,241 (12.36%)	53,249 (12.19%)	17,555 (11.64%)	45,049 (14.00%)	5,192 (12.49%)	266,286 (12.52%)
Not Hispanic or Latino	965,332 (82.14%)	358,777 (82.14%)	124,565 (82.61%)	256,255 (79.64%)	33,585 (80.81%)	1,738,514 (81.77%)
Unknown	64,713 (5.51%)	24,768 (5.67%)	8,659 (5.74%)	20,443 (6.35%)	2,785 (6.70%)	121,368 (5.71%)
Characteristics						
Type II Diabetes	504,048 (42.89%)	137,369 (31.45%)	70,398 (46.69%)	228,689 (71.08%)	29,544 (71.08%)	970,048 (45.62%)
Overweight or Obesity	830,625 (70.67%)	323,057 (73.96%)	94,283 (62.53%)	172,774 (53.70%)	21,130 (50.84%)	1,441,869 (67.82%)
Labeled Use						
ADM	516,033 (43.91%)	201,758 (46.19%)	51,838 (34.38%)	321,747 (100.00%)	41,562 (100.00%)	1,132,938 (53.29%)

	Semaglutide	Tirzepatide	Liraglutide	Dulaglutide	Other	Overall
AOM	226,672 (19.29%)	144,290 (33.03%)	29,959 (19.87%)	n/a	n/a	400,921 (18.86%)
Unknown	432,581 (36.81%)	90,746 (20.78%)	68,982 (45.75%)	n/a	n/a	592,309 (27.86%)
Year						
2018	3,670 (0.31%)	n/a	25,711 (17.05%)	22,153 (6.89%)	9,255 (22.27%)	60,789 (2.86%)
2019	17,262 (1.47%)	n/a	24,160 (16.02%)	28,874 (8.97%)	7,631 (18.36%)	77,927 (3.67%)
2020	30,373 (2.58%)	n/a	18,423 (12.22%)	31,379 (9.75%)	5,911 (14.22%)	86,086 (4.05%)
2021	69,488 (5.91%)	n/a	22,072 (14.64%)	47,486 (14.76%)	4,846 (11.66%)	143,892 (6.77%)
2022	160,250 (13.63%)	24,724 (5.66%)	24,698 (16.38%)	67,566 (21.00%)	4,832 (11.63%)	282,070 (13.27%)
2023	361,573 (30.76%)	70,398 (16.12%)	22,942 (15.22%)	70,763 (21.99%)	4,560 (10.97%)	530,236 (24.94%)
2024	375,928 (31.99%)	189,299 (43.34%)	9,822 (6.51%)	39,876 (12.39%)	3,447 (8.29%)	618,372 (29.08%)
2025	156,742 (13.34%)	152,373 (34.88%)	2,951 (1.96%)	13,650 (4.24%)	1,080 (2.60%)	326,796 (15.37%)
Comorbidity						
Asthma	169,985 (14.46%)	64,915 (14.86%)	22,105 (14.66%)	37,481 (11.65%)	4,391 (10.56%)	298,877 (14.06%)
CKD	138,878 (11.82%)	36,159 (8.28%)	19,068 (12.65%)	56,968 (17.71%)	7,290 (17.54%)	258,363 (12.15%)
COPD	58,322 (4.96%)	16,504 (3.78%)	8,609 (5.71%)	22,592 (7.02%)	2,997 (7.21%)	109,024 (5.13%)
Hyperlipidemia	611,191 (52.00%)	209,683 (48.01%)	66,769 (44.28%)	175,906 (54.67%)	22,578 (54.32%)	1,086,127 (51.08%)
Hypertension	626,972 (53.35%)	208,893 (47.82%)	76,360 (50.64%)	190,720 (59.28%)	24,695 (59.42%)	1,127,640 (53.04%)
Major Depression	146,118 (12.43%)	51,398 (11.77%)	19,397 (12.86%)	30,425 (9.46%)	3,724 (8.96%)	251,062 (11.81%)

Prescribing over time

Overall

We describe the monthly rate of prescriptions by medication and FDA labeled use over time:

First-time

We describe the monthly rate of first-time prescriptions by medication and FDA labeled use over time.

Characteristics of patients by year of first GLP-1 RA use are as follows:

Table 2: Patient Characteristics by Year of First GLP-1 RA Prescription.

	2018	2019	2020	2021	2022	2023	2024
N	60,789 (100.00%)	77,927 (100.00%)	86,086 (100.00%)	143,892 (100.00%)	282,070 (100.00%)	530,236 (100.00%)	618,372 (100.00%)
Age Group							
18-44	10,004 (16.46%)	12,952 (16.62%)	14,907 (17.32%)	28,270 (19.65%)	68,516 (24.29%)	140,841 (26.56%)	172,106 (27.83%)
45-64	33,616 (55.30%)	41,527 (53.29%)	45,040 (52.32%)	73,177 (50.86%)	141,171 (50.05%)	262,849 (49.57%)	292,806 (47.35%)
65+	17,169 (28.24%)	23,448 (30.09%)	26,139 (30.36%)	42,445 (29.50%)	72,383 (25.66%)	126,546 (23.87%)	153,460 (24.82%)
Sex							
Female	33,930 (55.82%)	43,076 (55.28%)	47,359 (55.01%)	84,375 (58.64%)	179,848 (63.76%)	350,977 (66.19%)	408,119 (66.00%)
Male	26,859 (44.18%)	34,851 (44.72%)	38,727 (44.99%)	59,517 (41.36%)	102,222 (36.24%)	179,259 (33.81%)	210,253 (34.00%)
Race							

	2018	2019	2020	2021	2022	2023	2024
White	45,824 (75.38%)	57,805 (74.18%)	62,595 (72.71%)	103,286 (71.78%)	202,685 (71.86%)	375,325 (70.78%)	437,612 (70.77%)
Black or African American	7,816 (12.86%)	9,815 (12.60%)	11,287 (13.11%)	19,765 (13.74%)	39,266 (13.92%)	75,916 (14.32%)	80,187 (12.97%)
Asian	1,244 (2.05%)	1,831 (2.35%)	2,256 (2.62%)	4,141 (2.88%)	6,907 (2.45%)	12,334 (2.33%)	15,145 (2.45%)
Unknown	3,637 (5.98%)	5,003 (6.42%)	5,703 (6.62%)	9,056 (6.29%)	17,235 (6.11%)	37,252 (7.03%)	51,453 (8.32%)
Native Hawaiian or Other Pacific Islander	234 (0.38%)	304 (0.39%)	363 (0.42%)	637 (0.44%)	1,119 (0.40%)	1,802 (0.34%)	2,174 (0.35%)
American Indian or Alaska Native	468 (0.77%)	662 (0.85%)	690 (0.80%)	1,315 (0.91%)	2,300 (0.82%)	3,723 (0.70%)	4,355 (0.70%)
Other Race	1,566 (2.58%)	2,507 (3.22%)	3,192 (3.71%)	5,692 (3.96%)	12,558 (4.45%)	23,884 (4.50%)	27,446 (4.44%)
Ethnicity							
Hispanic or Latino	5,799 (9.54%)	8,179 (10.50%)	9,958 (11.57%)	18,050 (12.54%)	37,255 (13.21%)	66,681 (12.58%)	79,106 (12.79%)
Not Hispanic or Latino	51,568 (84.83%)	65,092 (83.53%)	70,987 (82.46%)	117,847 (81.90%)	229,956 (81.52%)	434,631 (81.97%)	503,384 (81.40%)
Unknown	3,422 (5.63%)	4,656 (5.97%)	5,141 (5.97%)	7,995 (5.56%)	14,859 (5.27%)	28,924 (5.45%)	35,882 (5.80%)
Characteristics							
Type II Diabetes	43,073 (70.86%)	58,241 (74.74%)	62,376 (72.46%)	94,042 (65.36%)	147,001 (52.12%)	230,157 (43.41%)	226,497 (36.63%)
Overweight or Obesity	39,860 (65.57%)	49,868 (63.99%)	52,663 (61.17%)	92,476 (64.27%)	181,963 (64.51%)	357,669 (67.45%)	430,453 (69.61%)
Medication							
Semaglutide	3,670 (6.04%)	17,262 (22.15%)	30,373 (35.28%)	69,488 (48.29%)	160,250 (56.81%)	361,573 (68.19%)	375,928 (60.79%)
Tirzepatide	n/a	n/a	n/a	n/a	24,724 (8.77%)	70,398 (13.28%)	189,299 (30.61%)
Liraglutide	25,711 (42.30%)	24,160 (31.00%)	18,423 (21.40%)	22,072 (15.34%)	24,698 (8.76%)	22,942 (4.33%)	9,822 (1.59%)
Dulaglutide	22,153 (36.44%)	28,874 (37.05%)	31,379 (36.45%)	47,486 (33.00%)	67,566 (23.95%)	70,763 (13.35%)	39,876 (6.45%)
Other	9,255 (15.22%)	7,631 (9.79%)	5,911 (6.87%)	4,846 (3.37%)	4,832 (1.71%)	4,560 (0.86%)	3,447 (0.56%)
Labeled Use							

	2018	2019	2020	2021	2022	2023	2024
ADM	37,404 (61.53%)	46,185 (59.27%)	59,510 (69.13%)	105,027 (72.99%)	197,283 (69.94%)	302,669 (57.08%)	271,003 (43.83%)
AOM	1,823 (3.00%)	2,282 (2.93%)	3,394 (3.94%)	8,806 (6.12%)	20,392 (7.23%)	86,791 (16.37%)	169,255 (27.37%)
Unknown	21,562 (35.47%)	29,460 (37.80%)	23,182 (26.93%)	30,059 (20.89%)	64,395 (22.83%)	140,776 (26.55%)	178,114 (28.80%)
Comorbidity							
Asthma	6,834 (11.24%)	9,699 (12.45%)	10,865 (12.62%)	19,341 (13.44%)	38,384 (13.61%)	74,583 (14.07%)	89,470 (14.47%)
CKD	9,666 (15.90%)	13,812 (17.72%)	15,945 (18.52%)	25,021 (17.39%)	37,750 (13.38%)	58,517 (11.04%)	63,585 (10.28%)
COPD	3,737 (6.15%)	5,191 (6.66%)	5,942 (6.90%)	8,897 (6.18%)	14,506 (5.14%)	25,510 (4.81%)	29,351 (4.75%)
Hyperlipidemia	34,891 (57.40%)	47,244 (60.63%)	50,599 (58.78%)	81,015 (56.30%)	144,329 (51.17%)	259,675 (48.97%)	302,502 (48.92%)
Hypertension	36,830 (60.59%)	50,250 (64.48%)	53,593 (62.26%)	86,010 (59.77%)	152,577 (54.09%)	272,656 (51.42%)	310,212 (50.17%)
Major Depression	6,263 (10.30%)	9,197 (11.80%)	10,458 (12.15%)	19,331 (13.43%)	36,406 (12.91%)	61,655 (11.63%)	69,466 (11.23%)

60-day dispensing over time

We describe the proportion of patients with any GLP-1 RA prescriptions (first-time and repeat) in a given month who filled any GLP-1 RA in the 60 days following their prescription.

60-day initiation over time

We describe the proportion of patients first-prescribed a GLP-1 RA who filled any GLP-1 RA in the subsequent 60 days over time.

GLP-1 RA Initiation: Proportion of First-Time GLP-1 RA Prescriptions Dispensed within 60 days

Patient characteristics are given in Table 3:

Table 3: Patient Characteristics by Dispense Status. Dispense status refers to presence of any GLP-1 RA dispense within 60 days of first GLP-1 RA prescription. Table contains the subset of patients with observable 60-day dispense outcomes.

	Dispensed in 60 Days	Not Dispensed in 60 Days	Overall
N	1,175,816 (100.00%)	808,790 (100.00%)	1,984,606 (100.00%)
Age Group			
18-44	265,362 (22.57%)	234,967 (29.05%)	500,329 (25.21%)
45-64	591,130 (50.27%)	383,920 (47.47%)	975,050 (49.13%)
65+	319,324 (27.16%)	189,903 (23.48%)	509,227 (25.66%)
Sex			
Female	727,406 (61.86%)	540,815 (66.87%)	1,268,221 (63.90%)
Male	448,410 (38.14%)	267,975 (33.13%)	716,385 (36.10%)
Race			
White	826,844 (70.32%)	589,935 (72.94%)	1,416,779 (71.39%)
Black or African American	160,825 (13.68%)	107,379 (13.28%)	268,204 (13.51%)
Asian	31,669 (2.69%)	17,425 (2.15%)	49,094 (2.47%)
Unknown	90,154 (7.67%)	53,659 (6.63%)	143,813 (7.25%)
Native Hawaiian or Other Pacific Islander	4,677 (0.40%)	2,651 (0.33%)	7,328 (0.37%)
American Indian or Alaska Native	8,869 (0.75%)	5,812 (0.72%)	14,681 (0.74%)
Other Race	52,778 (4.49%)	31,929 (3.95%)	84,707 (4.27%)
Ethnicity			
Hispanic or Latino	151,084 (12.85%)	97,536 (12.06%)	248,620 (12.53%)
Not Hispanic or Latino	956,265 (81.33%)	669,746 (82.81%)	1,626,011 (81.93%)

	Dispensed in 60 Days	Not Dispensed in 60 Days	Overall
Unknown	68,467 (5.82%)	41,508 (5.13%)	109,975 (5.54%)
Characteristics			
Type II Diabetes	632,627 (53.80%)	292,875 (36.21%)	925,502 (46.63%)
Overweight or Obesity	772,206 (65.67%)	599,192 (74.08%)	1,371,398 (69.10%)
Medication			
Semaglutide	625,064 (53.16%)	478,207 (59.13%)	1,103,271 (55.59%)
Tirzepatide	231,096 (19.65%)	157,766 (19.51%)	388,862 (19.59%)
Liraglutide	79,498 (6.76%)	65,666 (8.12%)	145,164 (7.31%)
Dulaglutide	217,321 (18.48%)	90,148 (11.15%)	307,469 (15.49%)
Other	22,837 (1.94%)	17,003 (2.10%)	39,840 (2.01%)
Labeled Use			
ADM	744,843 (63.35%)	327,557 (40.50%)	1,072,400 (54.04%)
AOM	172,353 (14.66%)	194,963 (24.11%)	367,316 (18.51%)
Unknown	258,620 (21.99%)	286,270 (35.39%)	544,890 (27.46%)
Year			
2018	30,167 (2.57%)	29,672 (3.67%)	59,839 (3.02%)
2019	43,145 (3.67%)	33,234 (4.11%)	76,379 (3.85%)
2020	48,088 (4.09%)	36,326 (4.49%)	84,414 (4.25%)
2021	86,917 (7.39%)	54,380 (6.72%)	141,297 (7.12%)
2022	177,830 (15.12%)	96,782 (11.97%)	274,612 (13.84%)
2023	320,183 (27.23%)	197,090 (24.37%)	517,273 (26.06%)
2024	332,077 (28.24%)	254,310 (31.44%)	586,387 (29.55%)
2025	137,409 (11.69%)	106,996 (13.23%)	244,405 (12.32%)
Comorbidity			
Asthma	159,614 (13.57%)	123,667 (15.29%)	283,281 (14.27%)
CKD	156,576 (13.32%)	91,951 (11.37%)	248,527 (12.52%)
COPD	63,918 (5.44%)	40,079 (4.96%)	103,997 (5.24%)
Hyperlipidemia	616,885 (52.46%)	416,084 (51.45%)	1,032,969 (52.05%)
Hypertension	637,035 (54.18%)	430,209 (53.19%)	1,067,244 (53.78%)
Major Depression	130,335 (11.08%)	111,716 (13.81%)	242,051 (12.20%)

By Age

As a proxy for Medicare insurance, we segment initiation rate by populations 65+ versus 18-64.

Patient characteristics for patients aged 65+ are given in Table 4:

Table 4: Patient Characteristics by Dispense Status, Age 65+. Dispense status refers to presence of any GLP-1 RA dispense within 60 days of first GLP-1 RA prescription. Table contains the subset of patients with observable 60-day dispense outcomes.

	Dispensed in 60 Days	Not Dispensed in 60 Days	Overall
N	319,324 (100.00%)	189,903 (100.00%)	509,227 (100.00%)
Age Group			
65+	319,324 (100.00%)	189,903 (100.00%)	509,227 (100.00%)
Sex			
Female	176,772 (55.36%)	107,916 (56.83%)	284,688 (55.91%)
Male	142,552 (44.64%)	81,987 (43.17%)	224,539 (44.09%)
Race			
White	244,239 (76.49%)	151,414 (79.73%)	395,653 (77.70%)
Black or African American	30,986 (9.70%)	17,399 (9.16%)	48,385 (9.50%)
Asian	7,327 (2.29%)	3,186 (1.68%)	10,513 (2.06%)
Unknown	23,516 (7.36%)	11,993 (6.32%)	35,509 (6.97%)

	Dispensed in 60 Days	Not Dispensed in 60 Days	Overall
Native Hawaiian or Other Pacific Islander	721 (0.23%)	335 (0.18%)	1,056 (0.21%)
American Indian or Alaska Native	1,816 (0.57%)	961 (0.51%)	2,777 (0.55%)
Other Race	10,719 (3.36%)	4,615 (2.43%)	15,334 (3.01%)
Ethnicity			
Hispanic or Latino	26,693 (8.36%)	12,651 (6.66%)	39,344 (7.73%)
Not Hispanic or Latino	272,469 (85.33%)	166,362 (87.60%)	438,831 (86.18%)
Unknown	20,162 (6.31%)	10,890 (5.73%)	31,052 (6.10%)
Characteristics			
Type II Diabetes	215,090 (67.36%)	105,406 (55.51%)	320,496 (62.94%)
Overweight or Obesity	199,435 (62.46%)	126,356 (66.54%)	325,791 (63.98%)
Medication			
Semaglutide	171,501 (53.71%)	107,240 (56.47%)	278,741 (54.74%)
Tirzepatide	46,499 (14.56%)	26,917 (14.17%)	73,416 (14.42%)
Liraglutide	18,171 (5.69%)	14,852 (7.82%)	33,023 (6.48%)
Dulaglutide	73,751 (23.10%)	34,153 (17.98%)	107,904 (21.19%)
Other	9,402 (2.94%)	6,741 (3.55%)	16,143 (3.17%)
Labeled Use			
ADM	234,357 (73.39%)	101,190 (53.29%)	335,547 (65.89%)
AOM	14,565 (4.56%)	26,796 (14.11%)	41,361 (8.12%)
Unknown	70,402 (22.05%)	61,917 (32.60%)	132,319 (25.98%)
Year			
2018	8,274 (2.59%)	8,550 (4.50%)	16,824 (3.30%)
2019	12,516 (3.92%)	10,340 (5.44%)	22,856 (4.49%)
2020	14,377 (4.50%)	11,085 (5.84%)	25,462 (5.00%)
2021	25,273 (7.91%)	16,172 (8.52%)	41,445 (8.14%)
2022	45,517 (14.25%)	24,091 (12.69%)	69,608 (13.67%)
2023	85,124 (26.66%)	37,708 (19.86%)	122,832 (24.12%)
2024	89,469 (28.02%)	55,958 (29.47%)	145,427 (28.56%)
2025	38,774 (12.14%)	25,999 (13.69%)	64,773 (12.72%)
Comorbidity			
Asthma	38,413 (12.03%)	24,499 (12.90%)	62,912 (12.35%)
CKD	92,286 (28.90%)	53,304 (28.07%)	145,590 (28.59%)
COPD	32,334 (10.13%)	20,400 (10.74%)	52,734 (10.36%)
Hyperlipidemia	213,026 (66.71%)	135,254 (71.22%)	348,280 (68.39%)

	Dispensed in 60 Days	Not Dispensed in 60 Days	Overall
Hypertension	222,874 (69.80%)	140,841 (74.16%)	363,715 (71.42%)
Major Depression	34,774 (10.89%)	23,560 (12.41%)	58,334 (11.46%)

Patient characteristics for patients aged 18-65 are given in Table 5:

Table 5: Patient Characteristics by Dispense Status, Age 18-64. Dispense status refers to presence of any GLP-1 RA dispense within 60 days of first GLP-1 RA prescription. Table contains the subset of patients with observable 60-day dispense outcomes.

	Dispensed in 60 Days	Not Dispensed in 60 Days	Overall
N	856,492 (100.00%)	618,887 (100.00%)	1,475,379 (100.00%)
Age Group			
18-44	265,362 (30.98%)	234,967 (37.97%)	500,329 (33.91%)
45-64	591,130 (69.02%)	383,920 (62.03%)	975,050 (66.09%)
Sex			
Female	550,634 (64.29%)	432,899 (69.95%)	983,533 (66.66%)
Male	305,858 (35.71%)	185,988 (30.05%)	491,846 (33.34%)
Race			
White	582,605 (68.02%)	438,521 (70.86%)	1,021,126 (69.21%)
Black or African American	129,839 (15.16%)	89,980 (14.54%)	219,819 (14.90%)
Asian	24,342 (2.84%)	14,239 (2.30%)	38,581 (2.61%)
Unknown	66,638 (7.78%)	41,666 (6.73%)	108,304 (7.34%)
Native Hawaiian or Other Pacific Islander	3,956 (0.46%)	2,316 (0.37%)	6,272 (0.43%)
American Indian or Alaska Native	7,053 (0.82%)	4,851 (0.78%)	11,904 (0.81%)
Other Race	42,059 (4.91%)	27,314 (4.41%)	69,373 (4.70%)
Ethnicity			
Hispanic or Latino	124,391 (14.52%)	84,885 (13.72%)	209,276 (14.18%)
Not Hispanic or Latino	683,796 (79.84%)	503,384 (81.34%)	1,187,180 (80.47%)
Unknown	48,305 (5.64%)	30,618 (4.95%)	78,923 (5.35%)
Characteristics			
Type II Diabetes	417,537 (48.75%)	187,469 (30.29%)	605,006 (41.01%)
Overweight or Obesity	572,771 (66.87%)	472,836 (76.40%)	1,045,607 (70.87%)

	Dispensed in 60 Days	Not Dispensed in 60 Days	Overall
Medication			
Semaglutide	453,563 (52.96%)	370,967 (59.94%)	824,530 (55.89%)
Tirzepatide	184,597 (21.55%)	130,849 (21.14%)	315,446 (21.38%)
Liraglutide	61,327 (7.16%)	50,814 (8.21%)	112,141 (7.60%)
Dulaglutide	143,570 (16.76%)	55,995 (9.05%)	199,565 (13.53%)
Other	13,435 (1.57%)	10,262 (1.66%)	23,697 (1.61%)
Labeled Use			
ADM	510,486 (59.60%)	226,367 (36.58%)	736,853 (49.94%)
AOM	157,788 (18.42%)	168,167 (27.17%)	325,955 (22.09%)
Unknown	188,218 (21.98%)	224,353 (36.25%)	412,571 (27.96%)
Year			
2018	21,893 (2.56%)	21,122 (3.41%)	43,015 (2.92%)
2019	30,629 (3.58%)	22,894 (3.70%)	53,523 (3.63%)
2020	33,711 (3.94%)	25,241 (4.08%)	58,952 (4.00%)
2021	61,644 (7.20%)	38,208 (6.17%)	99,852 (6.77%)
2022	132,313 (15.45%)	72,691 (11.75%)	205,004 (13.90%)
2023	235,059 (27.44%)	159,382 (25.75%)	394,441 (26.73%)
2024	242,608 (28.33%)	198,352 (32.05%)	440,960 (29.89%)
2025	98,635 (11.52%)	80,997 (13.09%)	179,632 (12.18%)
Comorbidity			
Asthma	121,201 (14.15%)	99,168 (16.02%)	220,369 (14.94%)
CKD	64,290 (7.51%)	38,647 (6.24%)	102,937 (6.98%)
COPD	31,584 (3.69%)	19,679 (3.18%)	51,263 (3.47%)
Hyperlipidemia	403,859 (47.15%)	280,830 (45.38%)	684,689 (46.41%)
Hypertension	414,161 (48.36%)	289,368 (46.76%)	703,529 (47.68%)
Major Depression	95,561 (11.16%)	88,156 (14.24%)	183,717 (12.45%)

Discussion and Conclusion

This analysis summarized GLP-1 RA prescribing trends and patient characteristics from January 2018 to June 2025. During this period 2,196,747 patients were prescribed a total of 10,622,363 total prescriptions. ADMs accounted for 77.6% of prescriptions, and AOMs accounted for 22.4%, among prescriptions where use was known. A larger proportion of ADMs, compared to AOMs, were dispensed within 60 days: 69.5% of

first-time ADM prescriptions and 46.9% of first-time AOM prescriptions had a fill within 60 days.

Several limitations of our analysis should be noted. All data are preliminary and may change as additional data are obtained. These findings are consistent with data accessed July 11, 2025. Data presented in this analysis represent raw counts and/or rates, and post-stratification methods have not been conducted. In addition, cohorts with small counts may be suppressed during the de-identification process leading to the appearance of zero patients for a given time period. Given relatively large numbers of patients prescribed and dispensed GLP-1 RA, we do not expect substantial suppression in the most recent periods. The unknowns in this report either indicate the value was not included in the individual's electronic health record or that it was excluded from the data to protect an individual's identity as a part of Truveta's commitment to privacy (Truveta, 2024). Finally, this report is a description of prescribing trends and patient characteristics only. Results should not be interpreted as causal.

References

Truveta. (2024). Truveta's Approach to Patient Privacy.
https://resources.truveta.com/patient-privacy

Centers for disease control and prevention. "National diabetes statistics report". 2020.
https://www.cdc.gov/diabetes/data/statistics-report/index.html.

Davies, Melanie J., et al. "Efficacy of liraglutide for weight loss among patients with type 2 diabetes: the SCALE diabetes randomized clinical trial." JAMA 314.7 (2015): 687-699.

Davies, Melanie, et al. "Semaglutide 2· 4 mg once a week in adults with overweight or obesity, and type 2 diabetes (STEP 2): a randomised, double-blind, double-dummy, placebo-controlled, phase 3 trial." The Lancet 397.10278 (2021): 971-984.

Hales, Craig M., et al. "Prevalence of obesity and severe obesity among adults: United States, 2017–2018. NCHS Data Brief, no 360." National Center for Health Statistics 360 (2020): 1-7.

Han, Sabrina H., et al. "Public interest in the off-label use of glucagon-like peptide 1 agonists (Ozempic) for cosmetic weight loss: a google trends analysis." Aesthetic Surgery Journal 44.1 (2024): 60-67.

Garvey, W. Timothy, et al. "Tirzepatide once weekly for the treatment of obesity in people with type 2 diabetes (SURMOUNT-2): a double-blind, randomised, multicentre, placebo-controlled, phase 3 trial." The Lancet (2023).

Jastreboff, Ania M., et al. "Tirzepatide once weekly for the treatment of obesity." New England Journal of Medicine 387.3 (2022): 205-216.

Lincoff, A. Michael, et al. "Semaglutide and cardiovascular outcomes in obesity without diabetes." New England Journal of Medicine 389.24 (2023): 2221-2232.

Ussher, John R., and Daniel J. Drucker. "Glucagon-like peptide 1 receptor agonists: cardiovascular benefits and mechanisms of action." Nature Reviews Cardiology 20.7 (2023): 463-474.

Wilding, John PH, et al. "Once-weekly semaglutide in adults with overweight or obesity." New England Journal of Medicine 384.11 (2021): 989-1002.

Whitley, Heather P., Jennifer M. Trujillo, and Joshua J. Neumiller. "Potential Strategies for Addressing GLP-1 and Dual GLP-1/GIP Receptor Agonist Shortages." Clinical Diabetes (2023): cd230023.

Yeh, Tzu-Lin, et al. "Effect of glucagon-like peptide-1 receptor agonists on glycemic control, and weight reduction in adults: A multivariate meta-analysis." PLOS One 18.1 (2023): e0278685.

Acknowledgements

The authors thank Dr. Ty Gluckman for his contributions and feedback on describing GLP-1 RA prescribing trends.