Running Title: Wildfire Smoke Linked to Lower Sperm Count

Title: Wildfire Smoke Exposure is Associated with Decreased Sperm Concentration and Total Motile Sperm Count

Authors: Lillian X. Lindell,[1] MD; Sarah K. Holt, PhD; [1] Erin Petersen, MS;[2] Navya Gunaje, MSPH;[1] Arash Amighi, MD; [1] Amanda Haack, PhD; [2] Anthony Bui, MD; [1] Ryan Nasseri, MD; [1] Theodore Crisostomo-Wynne, MD;[1] Catherine J. Karr, MD, PhD;[3,4] Charles H. Muller, PhD, HCLD; [1] Thomas J. Walsh, MD; [1] Tristan M. Nicholson, MD, PhD[1, 3]

[1] *Department of Urology, School of Medicine, University of Washington*
[2] *School of Medicine, University of Washington*
[3] *Department of Environmental & Occupational Health Sciences, School of Public Health, University of Washington*
[4] *Department of Pediatrics, School of Medicine, University of Washington*

Corresponding Author: Tristan M. Nicholson, MD, PhD
Academic Affiliations: Department of Urology, School of Medicine and Department of Environmental & Occupational Health Sciences, School of Public Health, University of Washington
Mailing Address: 1959 NE Pacific St, Box 356510, Seattle, WA 98195
Phone: (970)333-9796
Email: tristann@uw.edu

Article Type: cohort study

Funding Statement: Research reported in this publication was funded by the University of Washington Medicine Endowed Professorship for Men's Health (Dr. Thomas J. Walsh), the Urology Care Foundation Research Scholar Award (Dr. Tristan M. Nicholson), and the National Institute Of Environmental Health Sciences of the National Institutes of Health under Award Number K12ES033584 and P30ES007033. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health. The REDCap instance in this research is funded by the National Center for Advancing Translational Sciences of the National Institutes of Health under award number UL1TR002319. Special thanks to Dr. Charles H. Muller and the University of Washington Male Fertility Lab staff for making this work possible.

Disclosure Statement: We have no conflicts of interest to disclose. All funding for this study are reported above.

Attestation statements: Data regarding any of the subjects in the study has not been previously published. Data will be made available to the editors of the journal for review or query upon request.

Word Count: 264/350 words for abstract, 3599 words for manuscript

Capsule: In this retrospective cohort study we demonstrate that wildfire smoke exposure is associated with declines in sperm concentration, total sperm count, and total motile sperm count across three wildfire events.

Structured Abstract

Objective: As wildfires become more prevalent, their impact on fertility warrants evaluation. We aimed to examine the impact of smoke exposure on semen analysis parameters of intrauterine insemination patients in the greater Seattle, WA area. We hypothesized that wildfire smoke exposure was associated with a decline in total motile sperm count.

Design: Retrospective cohort study

Subjects: Patients undergoing fertility treatments at the University of Washington in 2018-2022

Exposure: Subjects were exposed to seasonal wildfire events in the fall of 2018, 2020, and 2022. Pre-exposure semen was a diagnostic fresh sample prior to each respective wildfire event while post-exposure semen was taken at time of intrauterine insemination (IUI) during the wildfire smoke exposure windows. All subjects acted as their own controls in a paired pre-post analysis.

Main Outcome Measures: Primary outcome measure was total motile sperm count; secondary outcomes measures are semen volume, sperm concentration, total sperm count, total progressively motile sperm count, percent motile sperm, percent progressively motile sperm.

Results: We identified 84 subjects who underwent IUI across the 2018 (n = 27), 2020 (n = 30), and 2022 (n = 27) wildfire smoke events. Median time between initial semen analysis and semen analysis for IUI was 4 months. We observed a decline in sperm concentration, total sperm count, total motile sperm count, and total progressively motile sperm count. We also observed an increase in percent progressively motile sperm. These trends did not differ across event year.

Conclusions: Our results are consistent with a prior small study demonstrating that wildfire smoke exposure is associated with declines in sperm quality. These findings highlight the need for further research on the effects of wildfire smoke exposure on human sperm and fertility treatments, especially as smoke exposures are expected to increase with climate change.

Keywords
Wildfires, semen analysis, male fertility, assisted reproductive technology

Introduction

Wildfires in the last decade have been increasingly devastating, burning millions of acres and resulting in high intensity of hazardous smoke exposure for days to weeks on end.[1] The western United States is disproportionately affected by wildfire smoke, with states like California and Oregon experiencing seasonal exacerbations regularly.[1,2] The Puget Sound region provides an excellent setting for the study of wildfire smoke exposure due to its close proximity to seasonal wildfire activity, and allows researchers to examine the effects of short-term air quality disruptions within a context of otherwise clean air.[3,4] As wildfire events become more commonplace due to climate change,[5] it is imperative to understand the effects of their resultant smoke exposure on human health. The impact of wildfire smoke on men's reproductive health and semen parameters are not well-described and warrant further study, especially as infertility increases in prevalence. This study leverages our institution's location in the Puget Sound region where wildfire smoke events create distinct pre- and post-exposure periods, allowing for a natural experiment to assess how sudden, temporary surges in wildfire smoke affect semen parameters.

The reproductive health impacts due to air pollution from various sources including fossil fuels, agriculture, and industrial activities are well-characterized. Air pollution of this kind is generally composed of coarse particulate matter (PM) with diameters larger than 2.5 microns (μm),[6] and is linked to reduced sperm concentration, motility, and morphology.[7,8] Air pollution has also been linked to declines in conception and adverse pregnancy outcomes, with exposure associated with declines in sperm quality, ovarian dysfunction, and inferior fertility treatments results.[7]

In contrast, $PM_{2.5}$, which are 2.5 μm in diameter or less, is a major component of

wildfire smoke.[6] The smallest particles, ultrafine particulates, defined as compounds less than

0.1 μm in diameter, have been found to make up 30 to 90 percent of wildfire smoke.[9] Due to

the small size of these particulates, they easily diffuse through buildings and affect both indoor

and outdoor air quality, making wildfire smoke exposure particularly difficult to avoid.[10]

Therefore, lengthier periods of wildfire smoke exposure can lead to high, sustained levels of

fine particulate $PM_{2.5}$ levels both outdoors and indoors.[11,12]

Exposure to $PM_{2.5}$ has been linked to respiratory problems, heart attack, stroke, lung

cancer, and cognitive impairment.[13] However, the impact of $PM_{2.5}$ and wildfire smoke exposure

on human male fertility has not been well-characterized in the scientific literature. In mouse

models, there is evidence that exposure to wildfire smoke is associated with decreased

spermatogenesis and increased sperm DNA methylation, which is a marker for overall male

reproductive health.[14,15] One small cohort study in humans observed that the resultant smoke

exposure from the September 2020 wildfire event was associated with decreases in total motile

sperm count and sperm concentration in men undergoing intrauterine insemination (IUI) in

Portland, OR.[16] In another retrospective analysis, exposure to high concentrations of $PM_{2.5}$ was

associated with an increased odds of azoospermia and lower total progressive motile sperm

count.[17]

As wildfire activity increases, it is critical to understand how exposure affects human

sperm quality. One in six couples in the world face infertility, with male-factor contributing to

approximately 50% of these cases.[18] Fertility treatments such as IUI and other assisted

reproductive technologies (ART) are responsible for a growing share of live births in the United

States.[19] The effect of environmental exposures on sperm viability and reproductive outcomes

could have significant impacts on patient counseling and timing of intervention.

We hypothesized that acute hazardous levels of smoke exposure would be associated

with a decline in total motile sperm (TMC) in our patient population. Our study seeks to

contribute to a growing body of scientific literature on the health impacts of wildfire smoke

exposure. Our location in the Puget Sound region offers a unique opportunity to study this

phenomenon through a natural experiment. Understanding how wildfire smoke exposure

affects semen parameters would impact how patients are counseled prior to fertility

interventions, and stimulate future studies on mitigating environmental toxicant exposures in

these settings.

Methods

This retrospective, Institutional Review Board–approved cohort study analyzed

diagnostic semen analysis (SA) and SA at time of IUI completed at our institution. We evaluated

the last seven years of publicly available air quality data collected by a network of over 60

monitoring stations sponsored by the United States Environmental Protection Agency (EPA),

the Washington Department of Ecology, and local Washington clean air agencies.[20] We defined

a smoke event as any period lasting three days above the Washington Air Quality Advisory

(WAQA) threshold for fine particulates ($PM_{2.5}$ > 20.4 μg/m³) at the EPA monitoring station

closest to our clinic.[21] We defined a smoke exposure window as a three-month period starting

on the first day of the smoke event. We chose to use a three-month window because the

process of spermatogenesis takes, on average, 74 days to complete with an additional 14 days

of epididymal storage and maturation time prior to emission.[22,23]

Subjects acted as their own controls. We included subjects who had at least one IUI

performed during a smoke exposure window, at least one SA performed prior to the first day of

the smoke event, and those who lived in the surrounding Puget Sound region during the smoke

events. We included individuals' first IUI that occurred the smoke exposure window, although

this was not necessarily their initial IUI. For subjects with multiple SAs before or multiple IUIs

during the smoke exposure window, we compared SA temporally closest to the smoke

exposure window with the first SA for IUI following the onset of the smoke event. Medical

records review was performed to obtain subject age, race, ethnicity, smoking status, marijuana

use, infertility history, and illness status at time of IUI. Average amount of smoke exposure and

number of smoke days, defined as any day with PM $_5$ > 20.4 μg/m³ per the WAQA threshold,

were tabulated for all patients at time of SA for IUI based on home zipcodes and the Childs

Model (Stanford, CA) of daily smoke $PM_{2.5}$ concentration estimates based on a combination of

surface air quality sensor measurements, satellite smoke coverage analysis, and machine

learning.[1,2] Female partner data including fertility history and relevant medical diagnoses were

also collected. Our primary outcome was total motile sperm count (TMC), assessed before and

after the onset of the smoke exposure. Our secondary outcomes included semen volume,

sperm concentration, percent motility, percent progressive motility, total motile sperm, and

total progressively motile sperm.

Semen samples for SA were collected and processed following the World Health

Organization (WHO) 5th edition criteria for 2018 and 2020 data and 6th edition criteria for 2022

data[24,25] with the following modifications: Sperm concentration and total sperm count were

obtained with use of volumetric measures of semen and hemacytometer counts of sperm in

semen diluted with purified water. Sperm motility was evaluated in semen on both cover-

slipped wet mounts and Leja Standard Count 20 micron 2 chamber fixed coverslip slides (Leja

Products BV, Netherlands). A motility category "Rapid & Linear," corresponding to the WHO

6th "Rapid Progressive" motility and motility category A of WHO editions prior to the 5th

edition was evaluated and recorded for all patients. The diagnostic SA and the SA for IUI were

identical, except that morphology and immature germ cell evaluations were only performed for

the diagnostic SA. Study data were maintained in a Research Electronic Data Capture (REDCap)

database.[26]

Statistical analyses were conducted using R Studio version 4.4.1 (Auckland, NZ) and

RStudio IDE (Boston, MA). Linear mixed-effects models were employed to evaluate the impact

of wildfire smoke exposure on semen parameters including sperm volume, concentration,

motility, and count. The number of days between peak wildfire-related $PM_{2.5}$ exposure and

semen analysis was also calculated. This variable was used as a covariate in separate time

sensitivity analyses to assess whether the timing of IUI relative to the exposure event

influenced outcomes. "lme4" package was used to fit models with semen parameters before

and after smoke exposure as dependent variables. Models included random intercepts both for

patient and wildfire event year to account for temporal variations in the severity of smoke

exposure. *p*-Values were determined under the assumption of a two-tailed distribution, with

<0.05 considered statistically significant. Results were reported as estimated marginal means.

109   We identified and included three wildfire events from 2018, 2020, and 2022 as they

110 met our definition of a smoke event as described previously (Figure 1). The 2018 smoke

111 exposure window was from August 14, 2018 to November 14, 2018 (Figure 1A). Peak $PM_{2.5}$

112 occurred on August 21, 2018 at 102.2 $\mu g/m^3$. The smoke event lasted for 10 days until August

113 25, 2018, before returning to baseline below 20.4 $\mu g/m^3$ (Figure 1A). The 2020 smoke exposure

114 window was from September 9, 2020 to December 9, 2020 (Figure 1B). Peak $PM_{2.5}$ occurred on

115 September 15, 2020 at 177.7 $\mu g/m^3$. The smoke event lasted for 9 days until September 18,

116 2020, before returning to baseline (Figure 1B). The 2022 smoke exposure window was from

117 October 7, 2022 to January 7, 2023. Peak $PM_{2.5}$ occurred on October 20, 2022 at 108.0 $\mu g/m^3$.

118 The smoke event lasted for 14 days before returning to baseline on October 21, 2022 (Figure

119 1C). Notably in 2022, there was a period of unhealthy air from September 10 to September 12,

120 where the average daily $PM_{2.5}$ ranged from 22.2 to 44.5 $\mu g/m^3$. Although this period does not

121 qualify as a smoke event by our definition, the air quality was not in the normal range;

122 therefore, individuals with SAs completed from September 10 to October 7, 2022 were

123 excluded from analysis.

125 **Results**

126   We identified 84 subjects who underwent IUI during the wildfire smoke exposure

127 windows in 2018 (27 subjects), 2020 (30 subjects), and 2022 (27 subjects). Average age at IUI

128 was 37 ± 8 years, with a median time between initial SA and SA for IUI of 4 (3-9) months. 96% of

129 men did not have changes in their fertility history between initial SA and SA for IUI; one

130 individual discontinued a clomid trial, one patient underwent right varicocelectomy, and one

131 patient was recommended over-the-counter CoEnzyme Q10 supplementation, although it is

132 not known if he began taking it in the interim. Median time from day of peak $PM_{2.5}$ to SA for IUI

133 was 17 (1 - 69) days in 2018, 18.5 (1 - 76) days in 2020, and 14.5 (1 - 67) days in 2022. Mean

134 daily $PM_{2.5}$ wildfire smoke exposure for patients in the 90 days preceding IUI was 6.4 μg/m^3 in

135 2018, 9.4 μg/m^3 in 2020, and 6.2 μg/m^3 in 2022. Mean number of smoke days experienced by

136 patients in the 90 days preceding IUI was 8 ± 2 days in 2018, 9 ± 3 days in 2020, and 7 ± 3 days

137 in 2022.

138   Most patients were non-Hispanic white individuals with average BMI of 28 ± 5.

139 Average Yost Score of socioeconomic status was 25, with 1 being the highest socioeconomic

140 status and 100 being the lowest.[27] History taken at time of IUI demonstrated that most men

141 were non-smokers, who had never used marijuana, and had intermittent alcohol intake (Table

142 1). 69% of men had primary infertility at the time of IUI, and 24% of men were evaluated by our

143 urology practice prior to IUI. The average age of those evaluated by a urologist was 40 ± 13

144 years-old, with infertility being the most common reason for evaluation (Table 1). The most

145 common infertility-related diagnosis was erectile dysfunction.

146   Smoke exposure was estimated to correlate with a decline of 133 million total sperm (*p*

147 = 2.3 x 10^{-5}) and 71 million total motile sperm (*p* = 2.4 x 10^{-5}, Table 2). Semen volume also

148 declined by an estimated 0.6mL (*p* = 1.6 x 10^{-6}) and sperm concentration by 22 million/mL (*p* =

149 2.2 x 10^{-3}, Table 2). In our linear mixed-effect models, event year and time from peak $PM_{2.5}$

150 exposure as covariates did not have statistically significant impacts on semen parameters. All *p*-

151 values for event year effect and time from peak exposure were not statistically significant.

Partner data and observed reproductive outcomes were also summarized (Table 3). Of

the 84 female partners, average age was 34 ± 5 years at time of IUI. The majority of these

women had never been pregnant. All partners were evaluated by reproductive

endocrinologists, and no female factor infertility was identified in 36% of cases following

workup. Two semen samples for IUI were ultimately not inseminated due to low sperm count,

and one sample was taken to an outside institution for insemination. Of the remaining 81 IUIs,

9 (11%) resulted in pregnancy and 7 (9%) resulted in live birth with average gestational age of

39 ± 3 weeks at birth (Table 3).

Discussion

The increasing frequency and intensity of wildfires highlight the need to understand

how poor air quality and fine particulate pollution affect fertility. In our study, patients

undergoing IUI treatment had significant declines in numerous semen parameters after wildfire

smoke exposure. We found that wildfire smoke exposure was associated with declines in

semen volume, sperm concentration, total sperm count, total motile sperm count, total

progressively motile sperm count, and total rapid and linear motile sperm count (Table 2).

There was also a statistically significant increase in percent progressively motile sperm post-

exposure. Our findings are similar to those in a smaller cohort study examining changes in

semen parameters after the 2020 wildfire event,[16] and corroborate a larger analysis that

associates high $PM_{2.5}$ levels with increased odds of lower total progressively motile sperm

count.[17] Together, these findings support our hypothesis that wildfire smoke exposure is

associated with declines in sperm parameters.

Spermatogenesis occurs in cyclical waves over an average of 74 days.[22] Additionally,

fully-formed sperm undergo maturation and storage in the epididymis for approximately two

weeks prior to ejaculation.[23] It is unclear at which points during this process spermatozoa and

sperm are vulnerable to environmental toxicant exposures such as wildfire smoke. Although

individuals in our cohort were completing their SAs for IUI throughout the smoke exposure

period, the median absolute time from exposure to SA each year was 14-19 days. One plausible

pathway for this rapid impact of wildfire smoke exposure on semen parameters is through

direct toxicant effect on mature sperm during epididymal storage or transit through the vas

deferens, as sperm are particularly susceptible to noxious insults due to the lack of DNA repair

mechanisms in the epididymis.[28] Oxidative stress induced by inhaled pollutants have been

shown to damage sperm membranes and DNA rapidly, even in the absence of effects on

spermatogenesis.[29] Additionally, toxicants may disrupt epididymal function, which may impair

sperm maturation and may lead to changes in semen quality within a few weeks.[30] It is also

possible that toxicity during a specific stage of spermatogenesis may underlie our findings.

However, addressing this latter hypothesis will likely require large-scale epidemiological studies

with greater sample sizes and temporal resolution, or targeted investigations using animal

models.

We also observed a statistically significant rise in the percentage of progressively motile

sperm after wildfire exposure (Table 2). One possible theory for this observation is that

abnormally motile sperm are more vulnerable to the oxidative stress imparted by inhaled fine

particulates. Sperm progressive motility has been found to correlate positively with DNA

integrity, whereas higher levels of DNA fragmentation are found in poorly motile sperm.[31]

Sperm with higher levels of baseline DNA fragmentation, as evidenced by poor motility, may be

more susceptible to culling during interaction with acute stressors like fine particulate wildfire

smoke. These findings highlight the need for additional research on the effects of wildfire

smoke exposure on vulnerable populations such as men with elevated levels of sperm DNA

fragmentation.

In 2020, the maximum fine particulate $PM_{2.5}$ level in the Seattle area during a regional

wildfire event was of 177.7 µg/m^3, which was less than half of the maximum fine particulate

level in the Portland, OR region of $PM_{2.5}$ 388 µg/m^3.[16] Importantly, we observed the same

magnitude of impairment of semen parameters with a lower $PM_{2.5}$ level when compared to the

Portland, OR study that same year. When comparing between events in our Seattle area study,

we note that the maximum $PM_{2.5}$ level in 2018 was slightly over half of the $PM_{2.5}$ level in 2020

(102.2 vs 177.7 µg/m^3, respectively); however, event year, and by association severity of

wildfire smoke, did not seem to correlate with worse declines in semen parameters. Whether

these findings suggest fine particulate exposure has a threshold effect on changes in semen

parameters warrants further research.

While this study was not specifically designed to evaluate the impact of wildfire smoke

exposure on reproductive outcomes, our observational partner data highlight areas that merit

further investigation. In our cohort of women, whose average age was 34 years at time of IUI,

pregnancy rate was 11% and live birth rate was 9%, which are similar to the 12-20% pregnancy

rate and 9-15% live birth rate per cycle for women in their mid-thirties reported in the

literature.[32-34] Data collected at our institution from 1995-2017 demonstrate a pregnancy rate

of 16% and live birth rate of 6% after IUI.[35] This underscores the need for future research, as

the observed impact on IUI outcomes has important implications for patient counseling and

ART procedural timing.

Our study had several strengths. We were able to take advantage of a natural

experiment in a region with good air quality outside of wildfire smoke season where subjects

could act as their own controls. The data analyzed only included couples undergoing their first

IUI during a smoke window, and all our analyses were performed at the same center and by the

same staff. We also collected detailed clinical information about our subjects taken at time of

IUI, and all collections were done in person. Although we could not measure total time spent

outdoors versus indoors for subjects, prolonged unhealthy air quality outdoors is associated

with parallel declines in indoor air quality[10,11] and all individuals in this study traveled to our

clinic for semen analysis and IUI during periods of wildfire smoke, inevitably resulting in some

level of exposure to poor outdoor air.

Limitations of our study include small sample size and unmeasured variables that may

mitigate or exacerbate the health effects of wildfire smoke exposure. These include factors that

influence individual exposure such as the estimated duration of outdoor smoke exposure,

wearing of high-quality masks, use of indoor air cleaners, occupation, change in behaviors due

to wildfire smoke exposure, and exercise habits. Additionally, the subjects in our study live in

relatively high socioeconomic regions, which is not representative of the general population. It

is also possible that baseline declines in semen analysis parameters occur around the time of

IUI due to the inherent stress of the procedure.

Another potential unmeasured influence on sperm quality is the COVID-19 pandemic for

2020 and 2022 cohorts, as COVID-19 infection has been shown to have transient impact on

sperm quality.[36,37] Even mild or asymptomatic COVID infection has been found to decrease

sperm concentration, total sperm count, motility, and morphology.[38] Our intake forms at time

of IUI asked patients about recent illness including fever and cough, to which the vast majority

of subjects answered "no" (Table 2). However, mandatory COVID testing prior to both

diagnostic SA and IUI appointments was not conducted. It is therefore unknown if

asymptomatic COVID infection before or after smoke exposure could have had an impact on

semen parameter results. However, the likelihood this impacted our results is low, as the

semen parameters were not statistically different in the 2018 wildfire event year. Future

prospective studies where data are collected on subject behavior and symptoms are required

to address these concerns.

Most of the smoke days in the Puget Sound region are also the result of remote wildfires

that began in northeast Washington and Oregon. It is well-documented that the chemical

composition of wildfire smoke changes over time and space as it travels from its location of

origin.[39] Oxidation of organic compounds and the formation of secondary pollutants such as

ozone and peroxyacetyl nitrates may have a distinct effect on semen parameters, which

warrant further investigation.

Lastly, the timeline for exposure was not studied and is a critical area for further

research. We used a three-month smoke exposure window as a biologically plausible

timeframe; however, we do not know if there are more time-resolved sensitive windows of

sperm vulnerability from acute smoke exposure, if the effects of exposure could persist beyond

three months, and if and when semen parameters eventually recover. Interestingly, when we

accounted for time from peak wildfire smoke $PM_{2.5}$ exposure to SA for IUI, we did not see

statistically significant effects. However, previous studies have shown that the timing of

exposure may influence certain semen parameters. Ramsay et al's analysis of the Subfertility,

Health and Assisted Reproduction (SHARE) data demonstrates that progressively motile sperm

count appears most affected by exposure 60-74 days prior to SA, whereas azoospermia was

more likely with exposure 30-59 days prior to SA.[17] These are compelling evidence that wildfire

smoke has acute deleterious effects during the process of spermatogenesis, storage, and

transport of sperm, and perhaps suggest that cumulative exposure may play a more critical role

than the timing relative to peak exposure. Ultimately, the window of exposure relevant for

human sperm and fertility is not fully understood and requires further study.

Conclusions

In this study, we examined the impact of wildfire smoke exposure on male fertility,

finding significant declines in several semen parameters, including sperm concentration, total

motile sperm count, progressively motile sperm count, and rapid and linear motile sperm count

after exposure to fine particulate pollution. These results emphasize the need for further

research on the short- and long-term effects of wildfire smoke exposure on reproductive

health, especially as wildfire smoke events become more frequent with climate change.

Sources

1. Childs ML, Li J, Wen J, et al. Daily Local-Level Estimates of Ambient Wildfire Smoke PM(2.5) for the Contiguous US. *Environ Sci Technol*. Oct 4 2022;56(19):13607-13621. doi:10.1021/acs.est.2c02934
2. Childs ML, Martins M, Wilson AJ, Heft-Neal S, Qiu M, Burke M. Growing wildfire-derived PM2.5across the contiguous U.S. and implicationsfor air quality regulation. *Earth ArXiv*. 2024;doi:10.31223/X57H95
3. 2020 Data Summary | Puget Sound Clean Air Agency, WA.
4. 2022 Data Summary | Puget Sound Clean Air Agency, WA.
5. Liu JC, Pereira G, Uhl SA, Bravo MA, Bell ML. A systematic review of the physical health impacts from non-occupational exposure to wildfire smoke. *Environ Res*. Jan 2015;136:120-32. doi:10.1016/j.envres.2014.10.015
6. Vicente A, Alves C, Calvo AI, et al. Emission factors and detailed chemical composition of smoke particles from the 2010 wildfire season. *Atmospheric Environment*. 2013;71:295-303. doi:https://doi.org/10.1016/j.atmosenv.2013.01.062
7. Checa Vizcaino MA, Gonzalez-Comadran M, Jacquemin B. Outdoor air pollution and human infertility: a systematic review. *Fertil Steril*. Sep 15 2016;106(4):897-904 e1. doi:10.1016/j.fertnstert.2016.07.1110
8. Ramsay JM, Fendereski K, Horns JJ, et al. Environmental exposure to industrial air pollution is associated with decreased male fertility. *Fertil Steril*. Sep 2023;120(3 Pt 2):637-647. doi:10.1016/j.fertnstert.2023.05.143
9. Hata M, Chomanee J, Thongyen T, et al. Characteristics of nanoparticles emitted from burning of biomass fuels. *J Environ Sci (China)*. Sep 1 2014;26(9):1913-20. doi:10.1016/j.jes.2014.07.005
10. Agency USEP. Why Wildfire Smoke is a Health Concern. Accessed Feb 18. 2025, 2025. https://www.epa.gov/wildfire-smoke-course/why-wildfire-smoke-health-concern
11. Liang Y, Sengupta D, Campmier MJ, Lunderberg DM, Apte JS, Goldstein AH. Wildfire smoke impacts on indoor air quality assessed using crowdsourced data in California. *Proc Natl Acad Sci U S A*. Sep 7 2021;118(36)doi:10.1073/pnas.2106478118
12. May NW, Dixon C, Jaffe DA. Impact of Wildfire Smoke Events on Indoor Air Quality and Evaluation of a Low-cost Filtration Method. *Aerosol and Air Quality Research*. 2021/05/12 2021;21(7):210046. doi:10.4209/aaqr.210046
13. Kim KH, Kabir E, Kabir S. A review on the human health impact of airborne particulate matter. *Environ Int*. Jan 2015;74:136-43. doi:10.1016/j.envint.2014.10.005
14. Sosedova LM, Vokina VA, Novikov MA, et al. Reproductive Function of Male Rats and Motor Activity of Their Offspring in Fire Emissions Modeling. *Bull Exp Biol Med*. Feb 2022;172(4):472-477. doi:10.1007/s10517-022-05416-3
15. Schuller A, Bellini C, Jenkins TG, et al. Simulated Wildfire Smoke Significantly Alters Sperm DNA Methylation Patterns in a Murine Model. *Toxics*. Aug 27 2021;9(9)doi:10.3390/toxics9090199
16. Rubin ES, Kornfield M, Parker P, et al. Poor air quality from wildfire smoke is associated with decrease in total motile sperm count at time of intrauterine insemination. *Fertil Steril*. May 2024;121(5):881-883. doi:10.1016/j.fertnstert.2024.02.014

17. Ramsay JM, Horns JJ, Emery BR, Aston KI, Hotaling JM. EXPOSURE TO WILDFIRE SMOKE IS
ASSOCIATED WITH DECREASED SEMEN QUALITY. *Fertil Steril*. 2024;122(4):e210.
doi:/10.1016/j.fertnstert.2024.07.690
18. Organization WH. *Infertility Prevalence Estimates, 1991-2021*. 2023.
19. Technologies SfAR. *Preliminary National Summary Report for 2022*. 2025. *National*
Summary Report. Jan 17 2025.
https://sartcorsonline.com/CSR/PublicSnapshotReport?ClinicPKID=0&reportingYear=2022
20. Ecology WSDo. About the air monitoring network. Website. Accessed Jan 14, 2024.
https://ecology.wa.gov/regulations-permits/guidance-technical-assistance/air-monitoring-
network
21. Washington Air Quality Advisory (WAQA) Guidance for Public Health Actions. Washington
State Department of Health; 2019.
https://resources.finalsite.net/images/v1622568385/ksdorg/p6b8vphzd5zvm0axtfmt/wildfi
resmokeandoutdoorpublicactivities.pdf
22. Mahoudeau JA, Messing B. [Basic elements of testicular physiology]. *Concours Med*. Oct 25
1975;97(37):5841-4, 5847-50, 5835-5. Elements de physiologie testiculaire.
23. Barrachina F, Battistone MA, Castillo J, et al. Sperm acquire epididymis-derived proteins
through epididymosomes. *Hum Reprod*. Apr 1 2022;37(4):651-668.
doi:10.1093/humrep/deac015
24. Organization WH. *WHO laboratory manual for the examination and processing of human*
semen, Sixth Edition. 2021:292.
25. Organization WH. *WHO laboratory manual for the examination and processing of human*
semen, Fifth edition. 2010:271.
26. Harris PA, Taylor R, Thielke R, Payne J, Gonzalez N, Conde JG. Research electronic data
capture (REDCap)--a metadata-driven methodology and workflow process for providing
translational research informatics support. *J Biomed Inform*. Apr 2009;42(2):377-81.
doi:10.1016/j.jbi.2008.08.010
27. Yost K, Perkins C, Cohen R, Morris C, Wright W. Socioeconomic status and breast cancer
incidence in California for different race/ethnic groups. *Cancer Causes Control*. Oct
2001;12(8):703-11. doi:10.1023/a:1011240019516
28. Sakkas D, Alvarez JG. Sperm DNA fragmentation: mechanisms of origin, impact on
reproductive outcome, and analysis. *Fertil Steril*. Mar 1 2010;93(4):1027-36.
doi:10.1016/j.fertnstert.2009.10.046
29. Agarwal A, Mulgund A, Hamada A, Chyatte MR. A unique view on male infertility around the
globe. *Reprod Biol Endocrinol*. Apr 26 2015;13:37. doi:10.1186/s12958-015-0032-1
30. Robaire B, Hinton B. The Epididymis. In: Plant T, Zeleznik A, eds. *Knobil and Neill's*
Physiology of Reproduction. 2015.
31. Xin Z, Han N, Jin H. Correlation analysis of age and MTHFR C677T polymorphism with sperm
motility and sperm DNA integrity. *Cell Mol Biol (Noisy-le-grand)*. Jun 30 2023;69(6):110-115.
doi:10.14715/cmb/2023.69.6.17
32. Goverde AJ, McDonnell J, Vermeiden JP, Schats R, Rutten FF, Schoemaker J. Intrauterine
insemination or in-vitro fertilisation in idiopathic subfertility and male subfertility: a
randomised trial and cost-effectiveness analysis. *Lancet*. Jan 1 2000;355(9197):13-8.
doi:10.1016/S0140-6736(99)04002-7

33. Merviel P, Heraud MH, Grenier N, Lourdel E, Sanguinet P, Copin H. Predictive factors for
pregnancy after intrauterine insemination (IUI): an analysis of 1038 cycles and a review of
the literature. *Fertil Steril*. Jan 2010;93(1):79-88. doi:10.1016/j.fertnstert.2008.09.058
34. Haebe J, Martin J, Tekepety F, Tummon I, Shepherd K. Success of intrauterine insemination
in women aged 40-42 years. *Fertil Steril*. Jul 2002;78(1):29-33. doi:10.1016/s0015-
0282(02)03168-0
35. Nayan M, Punjani N, Grober E, Lo K, Jarvi K. The use of assisted reproductive technology
before male factor infertility evaluation. *Transl Androl Urol*. Aug 2018;7(4):678-685.
doi:10.21037/tau.2018.06.08
36. Gacci M, Coppi M, Baldi E, et al. Semen impairment and occurrence of SARS-CoV-2 virus in
semen after recovery from COVID-19. *Hum Reprod*. May 17 2021;36(6):1520-1529.
doi:10.1093/humrep/deab026
37. Guo TH, Sang MY, Bai S, et al. Semen parameters in men recovered from COVID-19. *Asian J*
Androl. Sep-Oct 2021;23(5):479-483. doi:10.4103/aja.aja_31_21
38. Che BW, Chen P, Yu Y, et al. Effects of mild/asymptomatic COVID-19 on semen parameters
and sex-related hormone levels in men: a systematic review and meta-analysis. *Asian J*
Androl. May-Jun 2023;25(3):382-388. doi:10.4103/aja202250
39. National Academies of Sciences Engineer (WASHINGTON District of Columbia). *The*
chemistry of fires at the wildland-urban interface. The National Academies Press;
2022:pages cm.
40. Esri. (2025). Air Quality and Smoke Forecast [Screenshot]. Retrieved from https://www.arcgis.com/apps/dashboards/

Table 1. Demographics data of included subjects

Parameter	All Cohorts	2018	2020	2022
Total number of individuals	84	27	30	27
Average age at IUI (years)	37 ± 8	34 ± 6	36 ± 4	40 ± 12
Median time between SA and IUI (months)	4 (3 – 9)	3 (2.5 – 5)	4 (2 – 9)	5 (3 – 10)
Average BMI	28 ± 5	27 ± 4	29 ± 6	27 ± 6
Average Yost Score of socioeconomic status	25 ± 22	21 ± 21	20 ± 17	31 ± 24
Census Bureau race group				
White	41 (49%)	12 (44%)	17 (57%)	12 (44%)
Black or African American	6 (7%)	2 (7%)	2 (7%)	2 (7%)
Asian	26 (31%)	11 (41%)	5 (17%)	10 (37%)
Native American or Alaskan Native	1 (1%)	0 (0%)	1 (3%)	0 (0%)
Declined to answer or unknown	10 (12%)	2 (7%)	5 (17%)	3 (11%)
Ethnicity				
Not Hispanic, Latino, or Spanish	71 (85%)	23 (85%)	25 (83%)	23 (85%)
Hispanic, Latino, or Spanish	3 (4%)	1 (4%)	0 (0%)	2 (7%)
Declined to answer or unknown	10 (12%)	3 (11%)	5 (17%)	2 (7%)
Smoking History				
Never	65 (77%)	21 (8%)	24 (80%)	20 (74%)
Former	13 (15%)	3 (11%)	4 (13%)	6 (22%)
Current	4 (5%)	3 (11%)	0 (0%)	1 (4%)
Unknown	2 (2%)	0 (0%)	2 (7%)	0 (0%)
Marijuana History				
Never	50 (60%)	20 (74%)	15 (50%)	15 (56%)
Former	14 (17%)	4 (15%)	5 (17%)	5 (19%)
Current	17 (20%)	3 (11%)	7 (23%)	7 (26%)
Unknown	3 (4%)	0 (0%)	3 (10%)	0 (0%)
Alcohol History				
None	20 (24%)	10 (37%)	4 (13%)	6 (22%)
Social (intermittent)	50 (60%)	11 (41%)	20 (67%)	19 (70%)
Regular (daily)	3 (4%)	2 (7%)	1 (3%)	0 (0%)
Unknown	11 (13%)	4 (15%)	5 (17%)	2 (2%)
Male Infertility Type				
Primary Infertility	58 (69%)	19 (70%)	23 (77%)	16 (60%)
Secondary Infertility	26 (31%)	8 (30%)	7 (23%)	11 (40%)
Infertility-Related Diagnoses				
Solitary testicle	1 (1%)	1 (4%)	0 (0%)	0 (0%)
Erectile dysfunction	5 (6%)	1 (4%)	1 (3%)	3 (11%)
HIV	1 (1%)	0 (0%)	0 (0%)	1 (4%)
Pituitary micro/macroadenoma	1 (1%)	1 (4%)	0 (0%)	0 (0%)
Prior testosterone use	1 (1%)	0 (0%)	0 (0%)	1 (4%)
Radiation exposure	2 (2%)	1 (4%)	1 (3%)	0 (0%)
Elevated scrotal temperatures (regular hot tub use)	7 (8%)	2 (8%)	4 (13%)	1 (4%)
Pesticides or other organic compounds	2 (2%)	0 (0%)	0 (0%)	2 (8%)
Febrile Illness within prior 3 months of IUI				
Yes	3 (4%)	1 (4%)	1 (3%)	1 (4%)
No	77 (92%)	24 (89%)	29 (97%)	24 (89%)
Unknown	4 (5%)	2 (7%)	0 (0%)	2 (7%)
Number of men evaluated by urology prior to IUI	24 (29%)	7 (26%)	6 (20%)	11 (41%)
Average age of men evaluated by urology (years)	40 ± 13	34 ± 3	39 ± 4	45 ± 18
Reasons for initial urologic evaluation (multiple selection)				
Infertility evaluation	20 (83%)	6 (86%)	6 (100%)	8 (73%)
Fertility preservation	1 (4%)	0 (0%)	0 (0%)	1 (9%)
History of vasectomy, desire for restoration of fertility	1 (4%)	0 (0%)	0 (0%)	1 (9%)
Varicocele	1 (4%)	1 (14%)	0 (0%)	0 (0%)
Testicular or pelvic pain	1 (4%)	1 (14%)	0 (0%)	0 (0%)
Low testosterone	1 (4%)	0 (0%)	0 (0%)	1 (9%)
Erectile dysfunction	1 (4%)	0 (0%)	1 (17%)	0 (0%)
Ejaculatory Dysfunction	1 (4%)	0 (0%)	1 (17%)	0 (0%)

Table 2. Estimated effect of wildfire smoke on median semen parameters of 84 subjects before and after exposure

Semen Parameters	Before Exposure	After Exposure	Estimated Effect*	P value**
Semen volume (mL)	3.5 (2.3 – 4.3)	3.1 (2.3 – 3.7)	-0.6	**1.6 x 10^{-6}**
Sperm concentration (mil/mL)	76 (49 – 127)	74 (33 – 102)	-22	**2.2 x 10^{-3}**
Percent motile (%)	54 (47 – 60)	53 (44 – 62)	-0.4	0.800
Percent progressively motile (%)	39 (32 – 66)	46 (38 – 56)	7	**4.4 x 10^{-5}**
Percent rapid & linear motile (%)	13.5 (10 – 18)	12.5 (5.5 – 18)	-2	0.055
Total sperm count (x10^6)	286 (177 – 419)	184 (98 – 332)	-133	**2.3 x 10^{-5}**
Total motile sperm (x10^6)	157 (80 – 250)	90 (35 – 188)	-71	**2.4 x 10^{-5}**
Total progressively motile sperm (x10^6)	106 (46 – 175)	77 (38 – 158)	-37	**4.9 x 10^{-3}**
Total rapid & linear motile sperm (x10^6)	34 (15 – 79)	20 (7 – 60)	-22	**1.2 x 10^{-4}**

Linear mixed model estimate of the effect of wildfire smoke exposure when controlling for event year
**Linear mixed model evaluation with p < 0.05 considered statistically significant, event year nor time from peak wildfire exposure did not have statistically significant impact for any independent variables so p-values are not reported

Table 3. Partner data and reproductive outcomes

Parameter	All Cohorts
Average female partner age at IUI (years)	34 ± 5
Median number of pregnancies prior	
Gravida	0 (0 – 1)
Para	0 (0 – 0)
Abortus	0 (0 – 1)
Partner female factory infertility history	
No female factor noted after workup	30 (36%)
Tubal factors	5 (6%)
Uterine factors	10 (12%)
Cervical factors	1 (1%)
Diminished Ovarian Reserve	12 (14%)
Polycystic Ovarian Syndrome	20 (24%)
Endometriosis	4 (5%)
Other	17 (20%)
Number of non-inseminated samples due to low count	2
Number of pregnancies achieved from IUI	9 (11%)
<35 years-old at conception	5 (56%)
35-40 years-old at conception	4 (44%)
>40 years-old at conception	0 (0%)
Number of live births	7 (9%)
<35 years-old at birth	3 (43%)
35-40 years-old at birth	4 (57%)
>40 years-old at birth	0 (0%)
Average gestational age at birth (weeks)	39 ± 3

Figure 1. Daily Mean PM$_{2.5}$ in Seattle WA

Daily mean PM$_{2.5}$ concentration (blue line) as documented by the 10[th] & Weller Monitoring Station closest to our Men's Health Lab and Clinic in the years 2018 (**A**), 2020 (**B**), and 2022 (**C**). The red line indicates the Washington Air Quality Advisory (WAQA) threshold for unhealthy air of 20.4 μg/m^3, the area highlighted in yellow indicates the three-month wildfire smoke exposure window, and the blue dots represent intrauterine insemination (IUI) dates of included individuals.