Impact of PCSK9 inhibitors on bone disease: A comprehensive drug-target Mendelian randomization study

Wangyu Yang[1,2], Xiaohui Wang[2,4], Hao Yang[3]* and Dingjun Hao[1,2,4]*

[1]Shaanxi University of Chinese Medicine, Xianyang, 712046, China

[2]Department of Spine Surgery, Honghui Hospital, Xian Jiaotong University.Xi'an, Shaanxi 710054,China

[3] Translational Medicine Center, Hong Hui Hospital, Xi'an Jiaotong University, Xi'an, 710054, China

[4] Shaanxi Key Laboratory of Spine Bionic Treatment, Xi'an, Shaanxi, 710054, China

* Correspondence:

Hao Yang

yanghao71_99@yeah.net

Dingjun Hao

haodingjun@mail.xjtu.edu.cn

Abstract

Background:

While certain studies suggest a relationship between hyperlipidemia and bone metabolism, the exact nature of proprotein convertase subtilisin/kexin 9 inhibitors (PCSK9i, targeted by, eg, alirocumab),

which were originally developed for lowering LDL cholesterol inhibitors, with bone disease is still unclear. This research endeavors to uncover the potential causal relationship between PCSK9i and several most popular bone diseases (osteoporosis (OP), osteoarthritis (OA), rheumatoid osteoarthritis (RA)) using Mendelian randomization (MR).

Methods: This study employed a comprehensive approach involving the extraction of single-nucleotide polymorphisms (SNPs) from genome-wide association studies (GWAS), followed by rigorous quality checks. PCSK9i instrumental variables were utilized to evaluate the effect of cholesterol-lowering drugs on osteoporosis, osteoarthritis, and rheumatoid arthritis.

Results: PCSK9i instrumental variables were validated using familial combined hyperlipidemia summary data. Our analysis did not reveal a significant causal relationship between PCSK9i and OP. However, there was an observed an increased Lumbar-spine bone mineral density (LS-BMD) with PCSK9i intaking (OR = 1.157, 95% CI: 0.963 to 1.330, P = 0.044). PCKS9i significantly increased genetic risk of knee OA (OR = 1.136, 95% CI: 1.027 to 1.228, P = 0.013), but not for hip OA. Genetic risk of seropositive RA was strongly reduced while consuming PCSK9i (OR = 0.796, 95% CI: 0.580 to 0.964, P=0.020) and this effect is independent with LDL levels, while we don't observe causal relationships with PCSK9i on seronegative RA.

Conclusions:

This study elucidates the causal relationship between PCSK9i and genetic predisposition to OP, OA, and RA. PCSK9i would benefit LS-BMD and protect the genetic risk of seropositive RA. Meanwhile, PCSK9i might be a risk factor for knee OA.

Keywords: PCSK9 inhibitor, Osteoarthritis, Rheumatoid Osteoarthritis, Mendelian randomization, Bone mineral density

1   Introduction

Hyperlipidemia, a well-established risk factor for cardiovascular diseases(1), has been increasingly

implicated in the pathophysiology of bone disorders(2, 3). Osteoporosis (OP), osteoarthritis (OA),

and rheumatoid arthritis (RA) are among the most prevalent bone diseases, each characterized by

distinct pathological mechanisms yet sharing potential links with lipid metabolism(4, 5, 6).

Proprotein convertase subtilisin/kexin 9 inhibitors (PCSK9i), which could inhibit the PCSK9 protein

to increase clearance of low-density lipoprotein cholesterol (LDL-C) from the bloodstream(7), have

primarily been recognized for their role in hyperlipidemia treatment(8). However, emerging evidence

suggests that PCSK9i has implications beyond lipid metabolism(9). In addition, another well-known

cholesterol-lowering drug, statins have been reported to improve BMD, as well as reduce risk of

fractures(10). This study aims to explore the potential impact of PCSK9i on bone diseases, in

comparison to traditional statins, using Mendelian randomization (MR) to elucidate potential causal

relationships.

While studies have started to unveil the relationship between hyperlipidemia and bone mineral

density (BMD), the role of PCSK9i in this context remains less understood. The current research

employs a comprehensive approach, utilizing single-nucleotide polymorphisms (SNPs) extracted

from genome-wide association studies (GWAS) to investigate the influence of PCSK9i on OP, OA,

and RA. This approach provides a robust framework to assess the impact of PCSK9i on these bone

diseases, thereby contributing to a more nuanced understanding of the systemic effects of lipid-

lowering agents. This study not only aims to clarify the role of PCSK9i in bone disease management

but also seeks to offer insights into the broader implications of cholesterol metabolism in skeletal

health.

2   Materials and Methods

Data Acquisition and Processing

The BMD outcomes considered in this research pertained to BMD variation by specific skeletal sites.

OA outcomes were split by hip and knee, and RA was characterized by seropositive or seronegative.

Our foundational dataset incorporated GWAS summary outcomes: eBMD (N = 265,627), Knee-

OA(n= 403,124), seronegative RA (n= 174,771) and seronegative RA (n= 177,430) from MRC-IEU

and TB-BMD (N = 56,284), LS-BMD (N = 28,498), FN-BMD (N = 32,735), and FA-BMD (N =

8,143) from GEFOS(11), Hip-OA(n=14,275) from arcOGEN(12). Table 1 offers an overview of the

data sources used and their respective participant demographics. The majority of participants were

from European backgrounds. Given the study's dependence on existing GWAS summary datasets,

there was no need for institutional board approval, though consent was acquired from all

contributors.

Genetic instrument selection for PCSK9 inhibitors

Genetic associations with LDL cholesterol (LDL-C) was sourced from GWAS summary statistics

encompassing 173,082 individuals(13). By identifying instrumental variables targeting PCSK9 to

lower LDL-C, we aimed to mimic the effects of PCSK9 inhibitors(14). To circumvent linkage

disequilibrium issues, we established a linkage disequilibrium threshold ($r^2 \leq 0.3$) and excluded SNPs

that exhibited strong LD with each other. Following this process, 8 significant PCSK9i SNPs were

retained, deemed appropriate as instrumental variables. In a complementary approach, FCHL as the

positive control outcome, guided by the Consensus criteria (N = 39,961 cases; 309,261 controls)

derived from UK Biobank(15).

MR analysis

In our article, we highlight the importance of genetic variation as an instrumental variable (IV) in

Mendelian Randomization (MR) studies. This requires adherence to three key assumptions: First, a

significant correlation between the genetic variation and the exposure (e.g., protein levels); second,

independence from confounding factors affecting both exposure and outcome; and third, the

influence of the genetic variation on the outcome should occur solely through the exposure. Meeting

these criteria ensures the accuracy and validity of causal inferences in MR studies.

The overarching objective of our Mendelian Randomization (MR) examination was to validate

PCSK9i instrumental variables, as well as PCSK9i on BMD, OP, OA, and RA. Our primary modus

operandi was the inverse variance weighted (IVW) technique(16). In a quest for heightened

analytical rigor, we also engaged the MR-Egger regression and median-based estimator

techniques(17). Situations with discernible horizontal pleiotropy saw the deployment of the MR-

PRESSO outlier test(18). Furthermore, our analyses leveraged both IVW and MR Egger regression

to probe horizontal pleiotropic SNP effects, with Cochran's Q-test being instrumental(19) in

heterogeneity quantifications. To extend our results, we performed the same analytical procedure for

LDL cholesterol and bone disease.

Robust analysis

We employed the Cochran Q-test metrics to measure heterogeneities. To evaluate possible horizontal

pleiotropic impacts of the SNPs, we used IVW (random effect) and MR-Egger regression techniques.

Moreover, we executed a "leave-one-out" sensitivity assessment to pinpoint any SNPs that might

have a pronounced influence. In this approach, we sequentially removed each SNP to ascertain if it

drove the observed association.

Statistical Analysis

Bonferroni correction (corrected p = 0.05/X*Y, where X represents the number of exposures and Y

represents the number of outcomes) was used to adjust for multiple tests in this MR analysis. All

statistical analyses employed the Two-Sample MR (version 0.5.7) (20) R software (version 4.1.3).

3   Results

3.1   Positive control analysis for PCSK9 inhibitors instrumental variables

PCSK9 inhibitors (PCSK9i), known for their effectiveness in hyperlipidemia treatment. Using the

ieu-a300 dataset, we identified eight significant SNPs associated with PCSK9 inhibitors as

instrumental variables (Table 2). To validate identified PCSK9i instrumental variables, we chose

GWAS summary data of Familial Combined Hyperlipidemia (FCHL), which is characterized by high

blood cholesterols(21) as positive controls for PCSK9i instrumental variables. As expected, our

Mendelian Randomization (MR) analysis revealed that PCSK9i significantly reduced the risk of

FCHL (Odds Ratio: 0.3311, 95% CI: 0.286-0.383, P.adj = 3.2e-50; Figure 1).

3.2   PCSK9 inhibitor usage associated with higher lumbar spine BMD

We next conduct two sample Mendelian Randomization analysis for PCSK9i and osteoporosis (OP),

which is marked by reduced Bone Mineral Density (BMD) and impaired bone integrity(22). In

conclusion, our results have not shown a causal effect between PCSK9i on susceptibility of OP

neither postmenopausal osteoporosis (PMOP) with fracture. However, a positive association of

PCSK9 inhibitors with increased Lumbar Spine-BMD (LS-BMD; OR=1.157, 95% CI: 0.963 -1.330,

p.adj=0.0442) was identified (Figure 2). The relationship we did is similar with previous

research(23), as shown in Supplementary Figure 1.

3.3   PCSK9i intake leads to high genetic risk of knee OA

We utilized data from two Genome-Wide Association Studies (GWAS) cohorts diagnosed with hip

and knee osteoarthritis, respectively. This approach allowed us to explore the potential influence of

PCSK9i on these specific subtypes of osteoarthritis at a genetic level.

While we found no causal relationship between PCSK9i intake and hip osteoarthritis, we observed a

different scenario for knee osteoarthritis. Notably, our results indicated that the genetic risk of knee

osteoarthritis was significantly increased with PCSK9i intake (OR=1.136, 95% CI: 1.027-1.228,

p.adj=0.0126, Figure 3). MR analysis of LDL and genetic risk for knee OA suggests that this effect is

related to reduced LDL levels (OR=0.909, 95% CI: 0.859-0.951, p.adj=5.46e-5, Figure 3).

3.4　PCSK9i intake reduces the risk of seropositive RA and is independent of LDL levels

Rheumatoid arthritis (RA) is a chronic inflammatory disease that affects the joints. RA can be

classified into seropositive RA and seronegative RA based on the presence or absence of rheumatoid

factor (RF) and anti-citrullinated protein antibodies (ACPA) in the blood(24). Seropositive RA is

more common and tends to have a more severe course and worse prognosis than seronegative RA(25,

26). A notable outcome of our study is the lack of evidence supporting a significant risk association

between PCSK9i and seronegative RA. More interestingly, the analysis revealed that the genetic

susceptibility to seropositive RA could be significantly mitigated by the intake of PCSK9i

(OR=0.796, 95% CI: 0.580-0.964, p.adj=0.0202, Figure 4). The subsequent LDL MR analysis

corroborates this conclusion, showing that the beneficial role of PCSK9i in seropositive RA is not a

consequence of LDL reduction, since LDL levels do not correlate with genetic risk of seropositive

RA (OR=1.034, 95% CI: 0.932-1.135, p.adj=0.506, Figure 4). This observation is particularly

significant as it implies a potential therapeutic role of PCSK9i in the management of seropositive RA,

independent of its lipid-lowering effects.

3.5   Robustness

The MR-Egger regression method was utilized to investigate the potential existence of horizontal

pleiotropy among the SNPs and the result. Our results did not identify any evidence of such

pleiotropy (all p > 0.05) (Supplementary Tables 1). Similarly, the funnel plots did not display any

apparent horizontal pleiotropy for the evaluated outcomes (Supplementary Figure 2). Additionally,

the leave-one-out sensitivity charts revealed that no individual SNP significantly impacted the causal

association, emphasizing the strength of our conclusions (Supplementary Figure 3).

4   Discussion

Although previous studies indicating that a reduction in genetically predicted low-density lipoprotein

cholesterol (LDL-C) is linked to an increase in bone mineral density(23), furthermore the

osteoporosis(27). Our finding that PCSK9i does not significantly impact overall osteoporosis (OP)

risk contrasts with some literature suggesting statins may positively influence bone health(28). This

discrepancy underscores the complexity of lipid-lowering agents' effects on bone metabolism. While

statins are thought to exert osteoanabolic effects possibly via the mevalonate pathway(29), the

mechanism through which PCSK9i might affect bone density remains less clear. The observed

increase in lumbar-spine bone mineral density (LS-BMD) with PCSK9i use, demands a more refined

understanding, particularly when contrasted with the broader systemic effects of statins.

The study's revelation of an increased genetic risk for knee osteoarthritis (OA) but not hip OA. This

specificity suggests a potential biomechanical or local metabolic factor influenced by PCSK9i that

differentially affects joints. The mechanistic pathways linking cholesterol metabolism, PCSK9i

action, and joint-specific OA development need to be rigorously investigated. This could pave the

way for joint-specific therapeutic strategies and possibly uncover unforeseen side effects of PCSK9i

therapy in susceptible individuals. Statins have been hypothesized to exert anti-inflammatory and

chondroprotective effects(30), potentially beneficial in OA management(31). The joint-specific risk

increase associated with PCSK9i raises critical questions about the differential impact of cholesterol-

lowering drugs on joint health. It suggests that the systemic lipid-lowering effect might not uniformly

translate into protective benefits across all joints.

The discovery that PCSK9 inhibitors can reduce the genetic risk of seropositive rheumatoid arthritis

(RA) is a significant advancement in understanding the interplay between lipid metabolism and

autoimmune diseases. However, an intriguing aspect of this finding is the lack of a direct correlation

between low-density lipoprotein (LDL) cholesterol levels and the risk of RA. This suggests that the

beneficial effects of PCSK9 inhibitors in reducing the risk of seropositive RA might not be solely due

to their primary role in lowering LDL cholesterol levels.

One possible explanation for this phenomenon could be the pleiotropic effects of PCSK9 inhibitors.

Beyond their well-known impact on lipid levels, PCSK9 inhibitors may exert additional biological

effects that influence immune response and inflammation(32, 33), two key factors in the

pathogenesis of RA. Another hypothesis is that the mechanism by which PCSK9 inhibitors reduce

RA risk may involve modulation of lipid subfractions or other metabolic pathways(34), rather than a

straightforward reduction in LDL cholesterol. Alterations in these components can have significant

effects on inflammatory processes and immune responses, potentially contributing to the observed

reduction in RA risk.

Considering these findings, it is paramount to adopt a critically analytical lens towards the

application of PCSK9i in bone diseases. While the potential benefits in LS-BMD enhancement and

seropositive RA risk reduction are promising, the risk of exacerbating knee OA poses a significant

clinical dilemma. Future research should aim to dissect the molecular and cellular pathways through

which PCSK9i exerts these differential effects on bone and joint health. Investigating the role of

PCSK9i in bone remodeling, joint-specific cartilage metabolism, and immune modulation could

unravel the complex interplay between lipid metabolism and musculoskeletal diseases. Ultimately,

these efforts could aid in developing more nuanced and effective treatment strategies for bone

diseases.

5   Conclusion

This research clarifies the causal link between PCSK9 inhibitors (PCSK9i) and genetic susceptibility

to osteoporosis (OP), osteoarthritis (OA), and rheumatoid arthritis (RA). The study suggests that

PCSK9i positively impacts lumbar spine bone mineral density (LS-BMD) and may offer protective

effects against the genetic risk associated with seropositive RA. However, it also indicates that

PCSK9i could potentially be a risk factor for the development of knee OA.

6   Abbreviation

MR: Mendelian Randomization

IVs: Instrumental variables

FCHL: Familial combined hyperlipidemia

BMD: Bone mineral density

TB-BMD: Total body-bone mineral density

LS-BMD: Lumbar spine bone mineral density

FN-BMD: Femoral neck bone mineral density

FA-BMD: Forearm bone mineral density

eBMD: Heel bone mineral density

RA: Rheumatoid osteoarthritis

OP: Osteoporosis

PMOP: postmenopausal osteoporosis

OA: Osteoarthritis

IVW: Inverse variance weighted

SNPs: single nucleotide polymorphisms

OR: Odds ratio

7 Declarations

7.1 Ethics approval and consent to participate.

Not applicable.

7.2 Consent for publication

Not applicable.

7.3 Competing interests

The authors declare that the research was conducted in the absence of any commercial or financial

relationships that could be construed as a potential conflict of interest.

7.4 Availability of data and material

The datasets analyzed for this study can be found in the IEU OpenGWAS

https://gwas.mrcieu.ac.uk/datasets/.

7.5 Funding

This work was supported by Natural Science Basic Research Program of Shaanxi(Program

No.2025JC-YBQN-1050)

7.6 Authors' contributions

Conceptualization, Dingjun Hao; Data curation, Xiaohui Wang; Formal analysis, Wangyu Yang;

Funding acquisition, Xiaohui Wang; Supervision, Dingjun Hao; Visualization, Xiaohui Wang;

Writing – original draft, Wangyu Yang; Writing – review & editing, Xiaohui Wang, Hao Yang and

Dingjun Hao. This study was completed with teamwork. Each author had made corresponding

contribution to the study. All authors contributed to the article and approved the submitted version.

8 Reference

1. Goldstein JL, Schrott HG, Hazzard WR, Bierman EL, Motulsky AG. Hyperlipidemia in
coronary heart disease. II. Genetic analysis of lipid levels in 176 families and delineation of a new
inherited disorder, combined hyperlipidemia. J Clin Invest. 1973;52(7):1544-68.

2. Parhami F, Garfinkel A, Demer LL. Role of lipids in osteoporosis. Arterioscler Thromb Vasc
Biol. 2000;20(11):2346-8.

3. Graham LS, Tintut Y, Parhami F, Kitchen CM, Ivanov Y, Tetradis S, et al. Bone density and
hyperlipidemia: the T-lymphocyte connection. J Bone Miner Res. 2010;25(11):2460-9.

4. Brownbill RA, Ilich JZ. Lipid profile and bone paradox: higher serum lipids are associated
with higher bone mineral density in postmenopausal women. J Womens Health (Larchmt).
2006;15(3):261-70.

5. de Munter W, van den Bosch MH, Sloetjes AW, Croce KJ, Vogl T, Roth J, et al. High LDL
levels lead to increased synovial inflammation and accelerated ectopic bone formation during
experimental osteoarthritis. Osteoarthritis Cartilage. 2016;24(5):844-55.

6. de Munter W, Blom AB, Helsen MM, Walgreen B, van der Kraan PM, Joosten LA, et al.
Cholesterol accumulation caused by low density lipoprotein receptor deficiency or a cholesterol-rich
diet results in ectopic bone formation during experimental osteoarthritis. Arthritis Res Ther.
2013;15(6):R178.

7. Abifadel M, Varret M, Rabes JP, Allard D, Ouguerram K, Devillers M, et al. Mutations in
PCSK9 cause autosomal dominant hypercholesterolemia. Nat Genet. 2003;34(2):154-6.

8. Sabatine MS. PCSK9 inhibitors: clinical evidence and implementation. Nat Rev Cardiol.
2019;16(3):155-65.

9. Ragusa R, Basta G, Neglia D, De Caterina R, Del Turco S, Caselli C. PCSK9 and
atherosclerosis: Looking beyond LDL regulation. Eur J Clin Invest. 2021;51(4):e13459.

10.   Wang Z, Li Y, Zhou F, Piao Z, Hao J. Effects of Statins on Bone Mineral Density and
Fracture Risk: A PRISMA-compliant Systematic Review and Meta-Analysis. Medicine (Baltimore).
2016;95(22):e3042.

11.   Zheng HF, Forgetta V, Hsu YH, Estrada K, Rosello-Diez A, Leo PJ, et al. Whole-genome
sequencing identifies EN1 as a determinant of bone density and fracture. Nature.
2015;526(7571):112-7.

12.   arc OC, arc OC, Zeggini E, Panoutsopoulou K, Southam L, Rayner NW, et al. Identification
of new susceptibility loci for osteoarthritis (arcOGEN): a genome-wide association study. Lancet.
2012;380(9844):815-23.

13.   Elsworth B, Lyon M, Alexander T, Liu Y, Matthews P, Hallett J, et al. The MRC IEU
OpenGWAS data infrastructure. 2020.

14.   Richardson TG, Sanderson E, Palmer TM, Ala-Korpela M, Ference BA, Davey Smith G, et
al. Evaluating the relationship between circulating lipoprotein lipids and apolipoproteins with risk of
coronary heart disease: A multivariable Mendelian randomisation analysis. PLoS Med.
2020;17(3):e1003062.

15.   Trinder M, Vikulova D, Pimstone S, Mancini GBJ, Brunham LR. Polygenic architecture and
cardiovascular risk of familial combined hyperlipidemia. Atherosclerosis. 2022;340:35-43.

16.   Burgess S, Butterworth A, Thompson SG. Mendelian randomization analysis with multiple
genetic variants using summarized data. Genet Epidemiol. 2013;37(7):658-65.

17.   Bowden J, Davey Smith G, Burgess S. Mendelian randomization with invalid instruments:
effect estimation and bias detection through Egger regression. Int J Epidemiol. 2015;44(2):512-25.

18.   Verbanck M, Chen CY, Neale B, Do R. Detection of widespread horizontal pleiotropy in
causal relationships inferred from Mendelian randomization between complex traits and diseases. Nat
Genet. 2018;50(5):693-8.

19.   Bowden J, Del Greco MF, Minelli C, Zhao Q, Lawlor DA, Sheehan NA, et al. Improving the
accuracy of two-sample summary-data Mendelian randomization: moving beyond the NOME
assumption. Int J Epidemiol. 2019;48(3):728-42.

20.   Hemani G, Zheng J, Elsworth B, Wade KH, Haberland V, Baird D, et al. The MR-Base
platform supports systematic causal inference across the human phenome. Elife. 2018;7.

21.   Veerkamp MJ, de Graaf J, Bredie SJ, Hendriks JC, Demacker PN, Stalenhoef AF. Diagnosis
of familial combined hyperlipidemia based on lipid phenotype expression in 32 families: results of a
5-year follow-up study. Arterioscler Thromb Vasc Biol. 2002;22(2):274-82.

22.   Rachner TD, Khosla S, Hofbauer LC. Osteoporosis: now and the future. The Lancet.
2011;377(9773):1276-87.

23.   Li GH, Cheung CL, Au PC, Tan KC, Wong IC, Sham PC. Positive effects of low LDL-C and
statins on bone mineral density: an integrated epidemiological observation analysis and Mendelian
randomization study. Int J Epidemiol. 2020;49(4):1221-35.

24.   Aletaha D, Smolen JS. Diagnosis and Management of Rheumatoid Arthritis: A Review.
JAMA. 2018;320(13):1360-72.

25.   Rantapaa Dahlqvist S, Andrade F. Individuals at risk of seropositive rheumatoid arthritis: the
evolving story. J Intern Med. 2019;286(6):627-43.

26.   Malmstrom V, Catrina AI, Klareskog L. Author Correction: The immunopathogenesis of
seropositive rheumatoid arthritis: from triggering to targeting. Nat Rev Immunol. 2022;22(7):459.

27.   You L, Sheng ZY, Tang CL, Chen L, Pan L, Chen JY. High cholesterol diet increases
osteoporosis risk via inhibiting bone formation in rats. Acta Pharmacol Sin. 2011;32(12):1498-504.

28.   Bauer DC, Mundy GR, Jamal SA, Black DM, Cauley JA, Ensrud KE, et al. Use of statins and
fracture: results of 4 prospective studies and cumulative meta-analysis of observational studies and
controlled trials. Arch Intern Med. 2004;164(2):146-52.

29.   Deng L, Ding Y, Peng Y, Wu Y, Fan J, Li W, et al. gamma-Tocotrienol protects against
ovariectomy-induced bone loss via mevalonate pathway as HMG-CoA reductase inhibitor. Bone.
2014;67:200-7.

30.   Yudoh K, Karasawa R. Statin prevents chondrocyte aging and degeneration of articular
cartilage in osteoarthritis (OA). Aging (Albany NY). 2010;2(12):990-8.

31.   Kadam UT, Blagojevic M, Belcher J. Statin use and clinical osteoarthritis in the general
population: a longitudinal study. J Gen Intern Med. 2013;28(7):943-9.

32.   Hellberg S, Shavva VS, Metso J, Chen G, Li Q-Z, Manfè V, et al. Alternate models of acute
dyslipidemia reveal divergent pathways upon atherosclerosis initiation. 2020.

33.   Momtazi-Borojeni AA, Sabouri-Rad S, Gotto AM, Pirro M, Banach M, Awan Z, et al.
PCSK9 and inflammation: a review of experimental and clinical evidence. Eur Heart J Cardiovasc
Pharmacother. 2019;5(4):237-45.

34.   Waldmann E, Wu L, Busygina K, Altenhofer J, Henze K, Folwaczny A, et al. Effect of
PCSK9 inhibition with evolocumab on lipoprotein subfractions in familial dysbetalipoproteinemia
(type III hyperlipidemia). PLoS One. 2022;17(3):e0265838.

9   Figure legends

Figure 1. Causal impact of PCSK9i on FCHL. The Forest plot was drawn using IVW results. OR:

odds ratio; IVW: inverse variance weighting; CI: confidence interval.

Figure 2. Causal impact of PCSK9i on OP and BMD. The Forest plot was drawn using IVW results.

OR: odds ratio; IVW: inverse variance weighting; CI: confidence interval.

Figure 3. Causal impact of PCSK9i on hip OA and knee OA. The Forest plot was drawn using IVW

results. OR: odds ratio; IVW: inverse variance weighting; CI: confidence interval.

Figure 4. Causal impact of PCSK9i on seronegative RA and seropositive RA. The Forest plot was

drawn using IVW results. OR: odds ratio; IVW: inverse variance weighting; CI: confidence interval.

Exposure	Outcome	Method		OR(95% CI)	p.adj
PCSK9i	**FCHL**	MR Egger		0.277(0.230 to 0.333)	9.660432e-06
		Weighted median		0.318(0.274 to 0.368)	1.792810e-52
		Inverse variance weighted		0.331(0.286 to 0.383)	3.235587e-50
		Simple mode		0.355(0.285 to 0.443)	3.645771e-05
		Weighted mode		0.293(0.252 to 0.340)	8.932427e-07

0   0.5   1   1.5   2

← Lower Risk of FCHL   Higher Risk of FCHL →

Exposure	Outcome	Method		OR(95% CI)	p.adj
PCSK9i	**TB-BMD**	MR Egger		0.998(0.880 to 1.116)	0.97386007
		Weighted median		1.010(0.925 to 1.094)	0.82240245
		Inverse variance weighted		0.995(0.924 to 1.066)	0.89579775
		Simple mode		1.077(0.927 to 1.221)	0.35811190
		Weighted mode		1.023(0.915 to 1.130)	0.69037377
	FN-BMD	MR Egger		1.077(0.456 to 1.692)	0.32002922
		Weighted median		1.016(0.840 to 1.191)	0.32299897
		Inverse variance weighted		1.010(0.852 to 1.167)	0.33818871
		Simple mode		1.091(0.848 to 1.326)	0.20589835
		Weighted mode		1.089(0.887 to 1.283)	0.18309426
	FA-BMD	MR Egger		0.963(0.708 to 1.216)	0.49240345
		Weighted median		0.935(0.745 to 1.121)	0.30261068
		Inverse variance weighted		0.915(0.739 to 1.084)	0.19521950
		Simple mode		0.902(0.608 to 1.187)	0.32782483
		Weighted mode		0.957(0.754 to 1.157)	0.43027732
	eBMD	MR Egger		1.024(0.983 to 1.066)	0.29839417
		Weighted median		1.017(0.981 to 1.053)	0.36371687
		Inverse variance weighted		1.004(0.976 to 1.032)	0.77685467
		Simple mode		0.967(0.892 to 1.041)	0.41297285
		Weighted mode		1.022(0.983 to 1.059)	0.31119563
	LS-BMD	MR Egger		1.049(0.267 to 1.828)	0.34672970
		Weighted median		1.149(0.943 to 1.335)	0.06160699
		Inverse variance weighted		1.157(0.963 to 1.330)	0.04418783
		Simple mode		1.192(0.909 to 1.442)	0.12231069
		Weighted mode		1.147(0.897 to 1.377)	0.14240915
	OP	MR Egger		1.019(0.712 to 1.325)	0.90940786
		Weighted median		1.071(0.817 to 1.320)	0.59270056
		Inverse variance weighted		1.178(0.940 to 1.387)	0.15070944
		Simple mode		1.093(0.677 to 1.501)	0.68413474
		Weighted mode		1.085(0.814 to 1.349)	0.56917140
	PMOP with fracture	MR Egger		1.234(0.515 to 1.907)	0.57466825
		Weighted median		1.111(0.566 to 1.644)	0.70310221
		Inverse variance weighted		0.957(0.451 to 1.461)	0.86426739
		Simple mode		0.983(−0.081 to 2.047)	0.97565637
		Weighted mode		1.085(0.463 to 1.700)	0.80289329

Lower BMD   Higher BMD

Exposure	Outcome	Method		OR(95% CI)	p.adj
LDL cholesterol	**Hip OA**	MR Egger		0.945(0.705 to 1.181)	4.352457e-01
		Weighted median		0.972(0.755 to 1.188)	5.395996e-01
		Inverse variance weighted		1.028(0.865 to 1.190)	5.007404e-01
		Simple mode		0.949(0.534 to 1.361)	5.460691e-01
		Weighted mode		0.981(0.800 to 1.162)	5.679543e-01
	Knee OA	MR Egger		0.912(0.840 to 0.976)	9.590069e-03
		Weighted median		0.909(0.840 to 0.968)	3.238987e-03
		Inverse variance weighted		0.909(0.859 to 0.951)	5.461831e-05
		Simple mode		0.863(0.726 to 0.979)	2.513062e-02
		Weighted mode		0.906(0.851 to 0.953)	3.148892e-04
PCSK9i	**Hip OA**	MR Egger		0.972(-0.518 to 2.461)	8.498958e-01
		Weighted median		1.168(0.610 to 1.701)	5.046624e-01
		Inverse variance weighted		1.237(0.780 to 1.645)	2.934361e-01
		Simple mode		1.202(0.446 to 1.923)	5.619942e-01
		Weighted mode		1.132(0.546 to 1.701)	6.033842e-01
	Knee OA	MR Egger		1.192(1.025 to 1.326)	6.185851e-02
		Weighted median		1.166(1.028 to 1.279)	1.642244e-02
		Inverse variance weighted		1.136(1.027 to 1.228)	1.257783e-02
		Simple mode		1.188(0.979 to 1.366)	1.245932e-01
		Weighted mode		1.184(1.037 to 1.301)	3.997747e-02

0   0.5   1   1.5   2

← Lower Risk   Higher Risk →

Exposure	Outcome	Method		OR(95% CI)	p.adj
LDL cholesterol	**Seronegative RA**	MR Egger		0.916(0.698 to 1.127)	0.41541422
		Weighted median		0.925(0.685 to 1.159)	0.50495859
		Inverse variance weighted		1.017(0.866 to 1.167)	0.81040273
		Simple mode		1.136(0.700 to 1.555)	0.54665188
		Weighted mode		0.993(0.776 to 1.209)	0.92480538
	Seropositive RA	MR Egger		1.152(1.000 to 1.284)	0.05303717
		Weighted median		1.081(0.926 to 1.230)	0.30865110
		Inverse variance weighted		1.034(0.932 to 1.135)	0.50635511
		Simple mode		0.955(0.642 to 1.267)	0.75631733
		Weighted mode		1.117(0.981 to 1.241)	0.09645099
PCSK9i	**Seronegative RA**	MR Egger		1.018(0.624 to 1.411)	0.93217253
		Weighted median		1.183(0.843 to 1.494)	0.31096487
		Inverse variance weighted		1.275(0.924 to 1.563)	0.13574372
		Simple mode		1.223(0.735 to 1.667)	0.42569608
		Weighted mode		1.184(0.857 to 1.480)	0.32431464
	Seropositive RA	MR Egger		0.775(0.482 to 1.008)	0.10662528
		Weighted median		0.774(0.525 to 0.963)	0.02196391
		Inverse variance weighted		0.796(0.580 to 0.964)	0.02017828
		Simple mode		0.706(0.301 to 1.004)	0.09347236
		Weighted mode		0.772(0.529 to 0.952)	0.04745203

0   0.5   1   1.5   2

← Lower Risk   Higher Risk →