Title:

Effects of Upper Trunk Flexion on Ventilation Volume in Semi-Fowler's Position

Autohors:

Sayuki Miyashita (1), Satoshi Kubota (2)(3), Takuya Furudate (2)

(1) Medihos Kawaguchiko, Minamitsuru, Yamanashi, Japan

(2) School of Health Sciences at Odawara, International University of Health and Welfare (IUHW),

Odarawa, Kanagawa, Japan

(3) Graduate school, IUHW

Corresponding Author:

Satoshi Kubota, PhD

Email: satoshi@iuhw.ac.jp

ORCID: https://orcid.org/0000-0001-5930-6862

1-2-25, Shiroyama, Odawara, Kanagawa, 250-8588, Japan

Abstract:

**Introduction**: Semi-Fowler's position, a semi-sitting position with the trunk raised 30° from supine,

is commonly used to improve respiratory function and reduce cardiovascular stress. Previous studies

have shown that upper trunk flexion in this position reduces cardiovascular burden compared to

conventional whole trunk elevation. However, its effects on ventilation have not been investigated.

**Objective**: To compare ventilation volumes between semi-Fowler's position with upper trunk

flexion (UT30) and with whole trunk elevation (WT30) during resting conditions.

**Methods**: Fourteen healthy young women (age: 20.57±0.73 years; BMI: 20.19±1.46) participated in

this study. Ventilation parameters were measured in three positions: supine (SPIN), 30° upper trunk

flexion (UT30), and 30° whole trunk flexion (WT30). Tidal volume (TV), minute ventilation (MV),

respiratory rate (RespiR), and RR intervals (RRi) were recorded using a ventilation measurement system and ECG. Subjective breathing difficulty was assessed using a numerical rating scale. Data were analyzed using linear mixed models for comparison among the three positions.

**Results**: TV was significantly higher in both UT30 and WT30 compared to SPIN ($p < 0.05$), with differences exceeding 90 mL (approximately 30% increase). MV was significantly higher in UT30 compared to SPIN ($p < 0.05$). RespiR remained consistent across all postures. RRi was significantly lower in WT30 compared to SPIN ($p < 0.05$), while no significant difference was observed between UT30 and SPIN.

**Conclusion**: Upper trunk flexion in semi-Fowler's position improves ventilation volume similar to whole trunk elevation while maintaining cardiovascular stability. This simple postural modification may provide clinical benefits for patients requiring respiratory support with minimal hemodynamic compromise.

Keywords:

Semi-Fowler's position, Tidal Volume, Minute Ventilation, Posture, Respiratory Function

1. Introduction

Fowler's position and Semi-Fowler's position, which involve raising the trunk from supine, are

commonly used in clinical settings to improve respiratory function and reduce cardiovascular stress (Carol A Rauen , Mary Beth Flynn Makic 2009; Kubota et al. 2015, 2017; Mary Jo Grap 2005).

Fowler's position typically raises the trunk 45-60°, while Semi-Fowler's position involves a more moderate elevation of 30-45°. Kubota et al. demonstrated that flexing the upper trunk results in less reduction in stroke volume and less increase in heart rate compared to the conventional Fowler's position that raises the entire trunk(Kubota et al. 2015, 2017). On the other hand, a posture during the Fowler's position or Semi-Fowler's position that minimizes respiratory stress has not been proposed.

During quiet breathing, it has been reported that tidal volume and minute ventilation increase when the trunk is raised(Agostoni and Robert E Hyatt 2011; Romei et al. 2010). This occurs due to changes in diaphragm shape and movement associated with gravitational changes on the trunk. In the supine position, gravity causes abdominal organs to spread toward the thoracic cavity. Consequently, diaphragmatic movement is restricted, and thoracic volume decreases. In contrast, in the upright sitting position, gravity causes the abdominal organs to move downward, resulting in a lowered diaphragm and increased thoracic volume. Therefore, ventilation volume is greater in Fowler's position compared to the supine position. Additionally, abdominal wall tension may affect respiratory function through its influence on intra-abdominal pressure(Goldman, Silver, and Lehr 1986; Novak et al. 2021). This study aimed to investigate the effects of upper trunk flexion in a

Semi-Fowler's position on ventilation volume, hypothesizing that it would reduce abdominal wall

tension and thereby improve respiratory function.

## 2. Method

### 2.1 Participants

The study was conducted in accordance with the Declaration of Helsinki and was approved by the

Ethics Committee of International University of Health and Welfare (approval number: 21-lg-16).

All participants were provided with detailed information about the study procedures and potential

risks before giving their written informed consent. The study participants were healthy young

women. Exclusion criteria included individuals with internal medical conditions such as respiratory

or cardiovascular diseases, those with low back pain (including past history), and those with obesity.

While 15 participants were initially measured, one was excluded due to experimental equipment

malfunction, resulting in a final analysis of 14 participants(age, $20.57 \pm 0.73$ years; height, $162.07 \pm$

5.4 cm; weight, $53.13 \pm 5.57$ kg; BMI, $20.19 \pm 1.46$kg/m²).

### 2.2 Experimental Procedure

Participants were positioned in three postures: supine (SPIN) position and two types of semi-

Fowler's positions (Figure 1), during which ventilation volumes were measured. The semi-Fowler's

positions consisted of a 30° upper trunk flexion position (hereafter UT30) and a 30° whole trunk

flexion position (hereafter WT30). WT30 represented a position where the entire trunk was raised.

This study employed 30° semi-Fowler's positions rather than Fowler's positions exceeding 30°, as lower-angle semi-Fowler's positions (around 30°) are commonly used for frail patients(Carol A Rauen , Mary Beth Flynn Makic 2009; Grap et al. 2005; Mary Jo Grap 2005; Zhu et al. 2020).

Postures were achieved using a fully electric hospital bed (Paramount Co., Japan) and mattress (KE-553, Paramount Co., Japan). For WT30, participants were asked to sit as deeply as possible on the bed. For UT30, following Kubota et al.'s method, trunk flexion was adjusted at the tenth thoracic vertebra(Kubota et al. 2015). During both UT30 and WT30 positions, participants were instructed to maintain their knees in slight flexion.

A preliminary experiment was conducted on a separate day to allow participants to become accustomed to breathing while wearing the ventilation measurement mask. Participants were instructed to avoid strenuous exercise, caffeine, alcohol consumption, and food intake after dinner the day before the experiment, and to completely refrain from food and drink for 2 hours before the experiment. The laboratory temperature was maintained at 28°C, within the thermoneutral zone, based on previous research(Kubota et al. 2017, 2022). Participants were asked to wear sleeveless tops and shorts, with properly fitted, non-constrictive bras to avoid chest compression.

The experimental procedure began with a 15-minute rest period in the supine position, followed by measurements in the three postures. Each postural measurement lasted 5 minutes, with a 10-

minute rest period between measurements. The measurement order was randomized to eliminate order effects.

Ventilation volumes were measured using a ventilation measurement mask (Hans Rudolph, Inc., USA) and a pneumotachograph (TSD117, BIOPAC Systems, Inc., USA). Electrocardiogram (ECG) data were collected using an ECG100C module (BIOPAC Systems, Inc., USA). Data acquisition was performed with the MP150 system and AcqKnowledge software (BIOPAC Systems, Inc., USA), sampled at 1000 Hz.

2.3 Data Analysis and Statistical Analysis

Pulmonary volume fractions were calculated as Tidal Volume (TV), Minute Ventilation (MV), and Respiratory Rate (RespiR), while RR intervals (RRi) were derived from the ECG. RRi represents the time required for one cardiac cycle and corresponds to the reciprocal of heart rate. Subjective assessment of respiratory difficulty was obtained using a Numerical Rating Scale (NRS). NRS responses were collected with 0 indicating easy breathing and 10 indicating difficult breathing. A Linear Mixed Model (LMM) was used to analyze the effects of postures on these parameters, with postural differences as fixed effects and individual differences as random intercepts. Estimated marginal means and confidence intervals were calculated, and comparisons between postures were performed using Tukey's method. Cohen's d was calculated as the effect size. Statistical analyses were conducted using R version 4.3.3 (R Core Team). The significance level was set at less than 5%.

3. Results

Table 1 presents the estimated marginal means and 95% confidence intervals for lung volume

fractions and RRi in each posture, along with the p-values and effect sizes from between-group

comparisons.

TV was significantly higher in both UT30 and WT30 compared to SPIN ($p < 0.05$), with differences

exceeding 90 mL in both cases. TV values were comparable between UT30 and WT30, with no

significant difference. MV was significantly higher in UT30 compared to SPIN ($p<0.05$), with a

difference of more than 1.5 L/min. In contrast, RespiR remained consistent across all postures at

approximately 14 breaths/min, and no significant differences were observed among the postures.

RRi was significantly lower in WT30 compared to SPIN ($p<0.05$), while no significant differences

were found between other postures. NRS showed no significant differences between any of the

postures.

The effect sizes for significant between-group differences ranged from approximately 1 to 2.4.

4. Discussion

The major findings suggested that raising the upper trunk improves ventilation volume similar to

raising the entire trunk, as TV was larger in UT30 compared to SPIN.

The significantly higher TV in WT30 compared to SPIN was consistent with previous studies comparing sitting and supine positions(Romei et al. 2010; Agostoni and Robert E Hyatt 2011; Takahashi et al. 1998). This can be attributed to the gravitational effect causing abdominal organs to move downward when the trunk is raised, resulting in a lowered diaphragm and increased thoracic volume(Agostoni and Robert E. Hyatt 2011; Mezidi and Guérin 2018).

In contrast, while UT30 involves raising the thoracic region and head, the abdomen remains unraised. Therefore, the downward movement of abdominal organs that occurs in WT30 is less likely to occur. Nevertheless, UT30 showed increased ventilation volume compared to SPIN. This may be explained by the relaxation of abdominal wall tension due to upper trunk flexion, which affects intra-abdominal pressure reduction and diaphragm descent. A positive correlation between abdominal tension and intra-abdominal pressure has been suggested(Novak et al. 2021). Furthermore, decreased intra-abdominal pressure has been shown to increase abdominal compliance, facilitating diaphragm descent(Goldman et al. 1986). Thus, the increase in TV likely resulted from the relaxation of abdominal muscles accompanying upper trunk flexion, which lowered intra-abdominal pressure and facilitated diaphragm descent. UT30 showed approximately 90 mL higher TV than SPIN, representing about a 30% increase. UT30, which involves raising only the upper trunk, can be achieved through a simple intervention using positioning cushions to raise the patient's upper trunk from the supine position. This minor postural modification may potentially provide a simple way to improve

ventilation volume.

Regarding MV, which is the product of TV and RR, considering that RR remained consistent at approximately 14 breaths/min across all postures with no significant differences, we conclude that the postural differences observed in TV influenced MV, resulting in differences between SPIN and the two semi-Fowler's positions.

RRi was significantly lower in WT30 compared to SPIN, which may be attributed to a potential decrease in stroke volume when raising the entire trunk in WT30, possibly leading to a compensatory increase in heart rate. In contrast, UT30 showed no significant change in RRi compared to SPIN, indicating that stroke volume might be less affected in this position. These findings are consistent with the hemodynamic results demonstrated by Kubota et al(Kubota et al. 2015, 2017).

Although no significant differences were observed in NRS, this experiment was conducted under resting conditions and represented a low physical burden for healthy individuals. Therefore, we believe that participants had difficulty perceiving differences in breathing between the different postures.

While our findings cannot be directly extrapolated to patient populations, they provide preliminary evidence for a beneficial effect of upper trunk flexion on ventilation. Previous studies have shown that semi-Fowler's position can increase lung capacity and improve diaphragm range of motion in various populations(Grap et al. 2005; Zhu et al. 2020). Our finding of a substantial increase in tidal volume in healthy subjects provides a physiological basis for future clinical studies.

In clinical settings such as post-operative care, where patients often experience reduced lung volumes and atelectasis, the UT30 position might reduce respiratory complications. This is particularly relevant for frail or elderly patients who may not tolerate full Fowler's positioning due to hemodynamic instability. Additionally, for mechanically ventilated patients, this positioning strategy might allow for lower tidal volume settings while maintaining adequate minute ventilation, potentially reducing the risk of ventilator-induced lung injury(Jiang et al. 2024; Yamamoto et al. 2022).

This study has several limitations that should be acknowledged. First, our sample consisted only of young healthy women, which limits the generalizability of our findings to other populations such as elderly patients, males, or individuals with respiratory or cardiovascular conditions. Second, measurements were conducted under resting conditions for short durations (5 minutes per posture), which may not reflect ventilatory responses during prolonged positioning or physical exertion. Third, while we measured ventilation parameters, we did not assess gas exchange efficiency or arterial blood gases, which would provide more comprehensive information about respiratory function. Fourth, the laboratory setting may not fully replicate clinical environments where patients are typically positioned.

Despite these limitations, our findings provide valuable preliminary evidence for the potential benefits of upper trunk flexion on ventilation volume. Future studies should include diverse patient populations, longer measurement periods, additional respiratory parameters, and investigate whether this increase in ventilation volume translates to meaningful clinical outcomes such as reduced dyspnea,

improved arterial blood gases, decreased length of mechanical ventilation, or reduced incidence of respiratory complications in various patient populations.

5. Conclusion

This study demonstrated that semi-Fowler's position with upper trunk flexion improves ventilation volume. This posture may provide similar ventilation volume increases as raising the entire trunk while potentially causing less cardiovascular stress.

References

Agostoni, Emilio, and Robert E Hyatt. 2011. "Static Behavior of the Respiratory System." *Comprehensive Physiology* 113–30.

Agostoni, Emilio, and Robert E. Hyatt. 2011. "Static Behavior of the Respiratory System." Pp. 113–30 in *Comprehensive Physiology*.

Carol A Rauen , Mary Beth Flynn Makic, Elizabeth Bridges. 2009. "Evidence-Based Practice Habits: Transforming Research into Bedside Practice." *Crit Care Nurse* 29(2):46–59.

Goldman, J. M., J. R. Silver, and R. P. Lehr. 1986. "An Electromyographic Study of the Abdominal Muscles of Tetraplegic Patients." *Spinal Cord* 24(4):241–46. doi: 10.1038/sc.1986.33.

Grap, Mary Jo, Cindy L. Munro, Russell S. Hummel III, R. K. Elswick Jr, Jessica L. McKinney, and Curtis N. Sessler. 2005. "Effect of Backrest Elevation on the Development of Ventilator-Associated Pneumonia." *American Journal of Critical Care* 14(4):325–32. doi: 10.4037/ajcc2005.14.4.325.

Jiang, Jie, Feiping Xia, Zhonghua Lu, Yuying Tang, Haibo Qiu, Yi Yang, and Fengmei Guo. 2024. "Effects of Tidal Volume on Physiology and Clinical Outcomes in Patients with One-lung Ventilation Undergoing Surgery: A Meta-analysis of Randomized Controlled Trials."

*Biomedical Reports* 20(5):73. doi: 10.3892/br.2024.1761.

Kubota, Satoshi, Yutaka Endo, Mitsue Kubota, Yusuke Ishizuka, and Takuya Furudate. 2015. "Effects of Trunk Posture in Fowler's Position on Hemodynamics." *Autonomic Neuroscience: Basic and Clinical* 189:56–59. doi: 10.1016/j.autneu.2015.01.002.

Kubota, Satoshi, Yutaka Endo, Mitsue Kubota, Hiroko Miyazaki, and Tomohiko Shigemasa. 2022. *The Pressor Response to the Drinking of Cold Water and Cold Carbonated Water in Healthy Younger and Older Adults*. Vol. 12.

Kubota, Satoshi, Yutaka Endo, Mitsue Kubota, and Tomohiko Shigemasa. 2017. "Assessment of Effects of Differences in Trunk Posture during Fowler's Position on Hemodynamics and Cardiovascular Regulation in Older and Younger Subjects." *Clinical Interventions in Aging* 12:603–10. doi: 10.2147/CIA.S132399.

Mary Jo Grap, Cindy L. Munro. 2005. "Quality Improvement in Backrest Elevation: Improving Outcomes in Critical Care." *AACN Clin Issues* 16(2):133–39.

Mezidi, Mehdi, and Claude Guérin. 2018. "Effects of Patient Positioning on Respiratory Mechanics in Mechanically Ventilated ICU Patients." *Annals of Translational Medicine* 6(19):1–9.

Novak, Jakub, Jakub Jacisko, Andrew Busch, Pavel Cerny, Martin Stribrny, Martina Kovari, Patricie Podskalska, Pavel Kolar, and Alena Kobesova. 2021. "Intra-Abdominal Pressure Correlates with Abdominal Wall Tension during Clinical Evaluation Tests." *Clinical Biomechanics* 88:105426. doi: 10.1016/j.clinbiomech.2021.105426.

Romei, M., A. Lo Mauro, M. G. D'Angelo, A. C. Turconi, N. Bresolin, A. Pedotti, and A. Aliverti. 2010. "Effects of Gender and Posture on Thoraco-Abdominal Kinematics during Quiet Breathing in Healthy Adults." *Respiratory Physiology and Neurobiology* 172(3):184–91. doi: 10.1016/j.resp.2010.05.018.

Takahashi, Tetsuya, Sumio Yamada, Kazuhiko Tanabe, Masaru Nakayama, Naohiko Osada, Haruki Itoh, and Masahiro Murayama. 1998. "The Effects of Posture on the Ventilatory Responses During Exercise." *Journal of the Japanese Physical Therapy Association* 1(1):13–17. doi: 10.1298/jjpta.1.13.

Yamamoto, Ryohei, Satoru Robert Okazaki, Yoshihito Fujita, Nozomu Seki, Yoshufumi Kokei, Shusuke Sekine, Soichiro Wada, Yasuhiro Norisue, and Chihiro Narita. 2022. "Usefulness of Low Tidal Volume Ventilation Strategy for Patients with Acute Respiratory Distress

Syndrome: A Systematic Review and Meta-Analysis." *Scientific Reports* 12(1):9331. doi: 10.1038/s41598-022-13224-y.

Zhu, Qiongfang, Zheyan Huang, Qiaomei Ma, Zehui Wu, Yubo Kang, Miaoyin Zhang, Tiantian Gan, Minxue Wang, and Fei Huang. 2020. "Supine versus Semi-Fowler's Positions for Tracheal Extubation in Abdominal Surgery-a Randomized Clinical Trial." *BMC Anesthesiology* 20(1):185. doi: 10.1186/s12871-020-01108-5.

Figure and Table

Figure 1 Bed positions for each condition

SPIN: Supine.

UT30: Lower and upper trunk inclined at 0° and 30°. Segments were subdivided based on the spinous process of 10th thoracic vertebra.

WT30: Lower and upper trunk inclined at 30°.

Table 1 Respiratory Function values and RR Interval in all positions

|  | Estimated Marginal Means(95% Confidence Intervals) | | | p-value(effect size) | | |
|---|---|---|---|---|---|---|
|  | SPIN | TH30 | WT30 | SPIN vs TH30 | SPIN vs WT30 | TH30 vs WT30 |
| Tidal Volume | 318.34 | 397.06 | 412.46 | 0.0002 | <0.0001 |  |
| (mL) | (251.26 - 385.42) | (329.98 - 464.14) | (345.37 - 479.54) | (1.77) | (2.11) |  |
| Minuitu Ventilation | 4.61 | 5.5 | 5.79 | 0.0002 | <0.0001 |  |
| (L/min) | (3.69 - 5.53) | (4.58 - 6.42) | (4.87 - 6.71) | (1.81) | (2.39) |  |
| Respiratory Rate | 14.91 | 14.33 | 14.74 |  |  |  |
| (times/min) | (13.68 - 16.13) | (13.11 - 15.55) | (13.52 - 15.97) |  |  |  |
| RRinterval | 0.929 | 0.908 | 0.9 |  | 0.0384 |  |
| (sec) | (0.884 - 0.975) | (0.863 - 0.954) | (0.854 - 0.945) |  | (0.99) |  |

SPIN: Supine.

UT30: Lower and upper trunk inclined at 0° and 30°.

WT30: Lower and upper trunk inclined at 30°.